

An Introduction to Normal Modal Logics

End Session

Cosimo Perini Brogi

Logic Group Working Seminar

Sept. 29, 2020

Past Seminars

1. Normal Modal Logics

Axiomatic Calculi and Canonical Model Theorems

↪ S. POPKORN, *First Steps in Modal Logic*, CUP 1994

2. Gödel-Löb Logic

Modal Completeness and Solovay Theorem

↪ G. BOOLOS, *The Logic of Provability*, CUP 1995

3. Structural Proof Theory

Cut Elimination and Some Corollaries

↪ S. NEGRI, J. VON PLATO, *Structural Proof Theory*, CUP 2001

↪ A. TROELSTRA, H. SWICHTENBERG, *Basic Proof Theory* 2nd ed., CUP 2000

4. Labelled Sequent Calculi

Cut Elimination "for" Kripke Frames with First Order Properties

↪ S. NEGRI, J. VON PLATO, *Proof Analysis*, CUP 2011

↪ S. NEGRI, R. DYCKHOFF, *Geometrization of First Order Logic*, The Bulletin of Symbolic Logic, 21(2) 2015

Past Seminars

1. Normal Modal Logics

Axiomatic Calculi and Canonical Model Theorems

↪ S. POPKORN, *First Steps in Modal Logic*, CUP 1994

2. Gödel-Löb Logic

Modal Completeness and Solovay Theorem

↪ G. BOOLOS, *The Logic of Provability*, CUP 1995

3. Structural Proof Theory

Cut Elimination and Some Corollaries

↪ S. NEGRI, J. VON PLATO, *Structural Proof Theory*, CUP 2001

↪ A. TROELSTRA, H. SWICHTENBERG, *Basic Proof Theory* 2nd ed., CUP 2000

4. Labelled Sequent Calculi

Cut Elimination "for" Kripke Frames with First Order Properties

↪ S. NEGRI, J. VON PLATO, *Proof Analysis*, CUP 2011

↪ S. NEGRI, R. DYCKHOFF, *Geometrization of First Order Logic*, The Bulletin of Symbolic Logic, 21(2) 2015

Past Seminars

1. Normal Modal Logics

Axiomatic Calculi and Canonical Model Theorems

↪ S. POPKORN, *First Steps in Modal Logic*, CUP 1994

2. Gödel-Löb Logic

Modal Completeness and Solovay Theorem

↪ G. BOOLOS, *The Logic of Provability*, CUP 1995

3. Structural Proof Theory

Cut Elimination and Some Corollaries

↪ S. NEGRI, J. VON PLATO, *Structural Proof Theory*, CUP 2001

↪ A. TROELSTRA, H. SWICHTENBERG, *Basic Proof Theory* 2nd ed., CUP 2000

4. Labelled Sequent Calculi

Cut Elimination "for" Kripke Frames with First Order Properties

↪ S. NEGRI, J. VON PLATO, *Proof Analysis*, CUP 2011

↪ S. NEGRI, R. DYCKHOFF, *Geometrization of First Order Logic*, The Bulletin of Symbolic Logic, 21(2) 2015

Past Seminars

1. Normal Modal Logics

Axiomatic Calculi and Canonical Model Theorems

↪ S. POPKORN, *First Steps in Modal Logic*, CUP 1994

2. Gödel-Löb Logic

Modal Completeness and Solovay Theorem

↪ G. BOOLOS, *The Logic of Provability*, CUP 1995

3. Structural Proof Theory

Cut Elimination and Some Corollaries

↪ S. NEGRI, J. VON PLATO, *Structural Proof Theory*, CUP 2001

↪ A. TROELSTRA, H. SWICHTENBERG, *Basic Proof Theory* 2nd ed., CUP 2000

4. Labelled Sequent Calculi

Cut Elimination “for” Kripke Frames with First Order Properties

↪ S. NEGRI, J. VON PLATO, *Proof Analysis*, CUP 2011

↪ S. NEGRI, R. DYCKHOFF, *Geometrization of First Order Logic*, The Bulletin of Symbolic Logic, 21(2) 2015

Today Seminar

Labels, Trees, Hypersequents

Tree Hypersequents and Labelled Calculi

- ↪ S. NEGRI, J. VON PLATO, *Proof Analysis*, CUP 2011
- ↪ F. POGGIOLESI, *Gentzen Calculi for Modal Propositional Logic*, Springer 2011
- ↪ R. GORÉ, R. RAMANAYAKE, *Labelled tree sequents, Tree hypersequents and Nested (Deep) Sequents*, in *Advances in Modal Logic (AiML 2012)*, College Publications 2012

Outline

Overview

G3GL

THS

LTS

Translation

Application

Questions

Löb Rule

Remark

For any $\mathcal{M} \in ITF$ $x \models_{\mathcal{M}} \Box A$ iff for any y s.t. $Rxy, y \models_{\mathcal{M}} \Box A \rightarrow A$.

Proof.

If $x \models_{\mathcal{M}} \Box A$, then, for any y such that $Rxy, y \models_{\mathcal{M}} A$, therefore $y \models_{\mathcal{M}} \Box A \rightarrow A$. Conversely, if for any y such that $Rxy, y \models_{\mathcal{M}} \Box A \rightarrow A$, then assuming $x \not\models_{\mathcal{M}} \Box A$, there exists a y such that Rxy and $y \not\models_{\mathcal{M}} A$, so that $y \not\models_{\mathcal{M}} \Box A$, therefore there is a z such that Ryz and $z \not\models_{\mathcal{M}} A$, ..., contra $\mathcal{M} \in ITF$.



Löb Rule

Remark

For any $\mathcal{M} \in ITF$ $x \models_{\mathcal{M}} \Box A$ iff for any y s.t. $Rxy, y \models_{\mathcal{M}} \Box A \rightarrow A$.

Proof.

If $x \models_{\mathcal{M}} \Box A$, then, for any y such that $Rxy, y \models_{\mathcal{M}} A$, therefore $y \models_{\mathcal{M}} \Box A \rightarrow A$. Conversely, if for any y such that $Rxy, y \models_{\mathcal{M}} \Box A \rightarrow A$, then assuming $x \not\models_{\mathcal{M}} \Box A$, there exists a y such that Rxy and $y \not\models_{\mathcal{M}} A$, so that $y \not\models_{\mathcal{M}} \Box A$, therefore there is a z such that Ryz and $z \not\models_{\mathcal{M}} A$, ..., contra $\mathcal{M} \in ITF$.



G3GL

Let **G3GL** denote the following modification/extension of G3K:

Initial sequents: $+ \quad x : \Box A, \Gamma \Rightarrow \Delta, x : \Box A$

Rules: $+ \quad \frac{}{Rxx, \Gamma \Rightarrow \Delta} Irref \quad \& \quad \frac{Rxy, Ryz, Rxz, \Gamma \Rightarrow \Delta}{Rxy, Ryz, \Gamma \Rightarrow \Delta} Trans$

$$\frac{Rxy, y : \Box A, \Gamma \Rightarrow \Delta, y : A}{\Gamma \Rightarrow \Delta, x : \Box A} \Box_L - K$$

$$\frac{x : \Box A, Rxy, \Gamma \Rightarrow \Delta, y : \Box A \quad y : A, x : \Box A, Rxy, \Gamma \Rightarrow \Delta}{x : \Box A, Rxy, \Gamma \Rightarrow \Delta} \Box_L - A$$

$\Box_L - A$ and $\Box_L - K$ substitute the standard rules for \Box .

Moreover, in $\Box_L - K$, we require y to be a fresh variable

Structural Properties

Lemma

For G3GL the following hold:

- (i) Weakening and substitution rules are hp admissible;
- (ii) All rules are invertible;
- (iii) Let the **range** of x in a derivation δ be the finite set of worlds y such that Rxy or the relational terms $Rxx_1, x_1Rx_2, \dots, x_nRy$ appear in the antecedents of sequents in δ ; let also ranges be ordered by set inclusion. Then contraction rules are range-preserving admissible.

Proof.

To (i): By induction on the height of the derivation.

To (ii): The relevant case is $\Gamma \Rightarrow \Delta, x : \Box A$. If $x : \Box A$ is not principal, then $Rxy, y : \Box A; \Gamma \Rightarrow \Delta, y : A$ or we have *Irref*. But if $x : \Box A$ is principal, then $\Gamma \equiv x : \Box A, \Gamma'$ and we have $Rxy, y : \Box A, x : \Box A, \Gamma' \xRightarrow{m+1} \Delta, y : A$ by $\Box_L - A$ with $Rxy, y : \Box A, x : \Box A, \Gamma' \xRightarrow{0} \Delta, y : A, y : \Box A$ and $y : A, Rxy, y : \Box A, x : \Box A, \Gamma' \xRightarrow{m} \Delta, y : A$.

Structural Properties

Lemma

For G3GL the following hold:

- (i) Weakening and substitution rules are hp admissible;
- (ii) All rules are invertible;
- (iii) Let the **range** of x in a derivation δ be the finite set of worlds y such that Rxy or the relational terms $Rxx_1, x_1Rx_2, \dots, x_nRy$ appear in the antecedents of sequents in δ ; let also ranges be ordered by set inclusion. Then contraction rules are range-preserving admissible.

Proof.

To (i): By induction on the height of the derivation.

To (ii): The relevant case is $\Gamma \xRightarrow{0} \Delta, x : \Box A$. If $x : \Box A$ is not principal, then $Rxy, y : \Box A; \Gamma \xRightarrow{0} \Delta, y : A$ or we have *Irref*. But if $x : \Box A$ is principal, then $\Gamma \equiv x : \Box A, \Gamma'$ and we have $Rxy, y : \Box A, x : \Box A, \Gamma' \xRightarrow{m+1} \Delta, y : A$ by $\Box_L - A$ with $Rxy, y : \Box A, x : \Box A, \Gamma' \xRightarrow{0} \Delta, y : A, y : \Box A$ and $y : A, Rxy, y : \Box A, x : \Box A, \Gamma' \xRightarrow{m} \Delta, y : A$.

Structural Properties

Proof.

To (iii): By simultaneous induction for left and right contraction, with primary induction on the contraction formula, and secondary induction on the height of the derivation.

Relevant case: Let $x : \Box A$ to be contracted on the right. If it is not principal, use SIH and then the rule. Otherwise, the premise must be $Rxy, y : \Box A, \Gamma \Rightarrow \Delta, x : \Box A, y : A$, so we can invert $\Box_L - K$ obtaining $Rxy, y : \Box A, Rxy, y : \Box A, \Gamma \Rightarrow \Delta, y : A, y : A$ and by IH we have the result.



Cut Elimination for G3GL

Cut Elimination Theorem

The cut rule is admissible in G3GL.

Proof.

By induction on the **weight** of the cut, given by the cut formula complexity, the range of the cut label in the derivation and the cut height.

First note that if a loop occurs in the the conclusion of a *Cut*, then we can obtain that conclusion by *Irref* and *Trans*. Otherwise, there cannot be any loop in the premisses, and by the restrictions on variables we cannot introduce a loop by $\Box_L - K$.

Then if $y \in \text{rg}(x)$, then $\text{rg}(y) \subset \text{rg}(x)$.

Now, if the cut formula is not principal in the premisses of *Cut*, we can lift the cut. Otherwise, if $x : \Box A$ is principal in both premisses, then it must be introduced by \Box_L rules. Hence we can

1. eliminate the cut of $x : \Box A$ in the conclusion of $\Box_L - K$ and in the first premiss of $\Box_L - A$ by TIH, obtaining the sequent $Rxz, \Gamma, \Gamma' \Rightarrow \Delta, \Delta', z : \Box A$;
2. eliminate the cut of $x : \Box A$ in the conclusion of $\Box_L - K$ and in the second premiss of $\Box_L - A$ by TIH, obtaining the sequent $Rxz, z : A, \Gamma, \Gamma' \Rightarrow \Delta, \Delta'$;
3. substitute the eigenvariable in the premiss of $\Box_L - K$ to match that one of 1. and cut $z : \Box A$, obtaining by SIH $Rxz, Rxz, \Gamma, \Gamma', \Gamma \Rightarrow \Delta, \Delta', \Delta, z : A$ without cut;
4. eliminate the cut of $z : A$ between 3. and 2. by IH, obtaining by contraction $Rxz, \Gamma, \Gamma' \Rightarrow \Delta, \Delta'$, as desired.



Cut Elimination for G3GL

Cut Elimination Theorem

The cut rule is admissible in G3GL.

Proof.

By induction on the [weight](#) of the cut, given by the cut formula complexity, the range of the cut label in the derivation and the cut height.

First note that if a loop occurs in the the conclusion of a *Cut*, then we can obtain that conclusion by *Irref* and *Trans*. Otherwise, there cannot be any loop in the premisses, and by the restrictions on variables we cannot introduce a loop by $\Box_L - K$.

Then if $y \in \text{rg}(x)$, then $\text{rg}(y) \subset \text{rg}(x)$.

Now, if the cut formula is not principal in the premisses of *Cut*, we can lift the cut. Otherwise, if $x : \Box A$ is principal in both premisses, then it must be introduced by \Box_L rules. Hence we can

1. eliminate the cut of $x : \Box A$ in the conclusion of $\Box_L - K$ and in the first premiss of $\Box_L - A$ by TIH, obtaining the sequent $Rxz, \Gamma, \Gamma' \Rightarrow \Delta, \Delta', z : \Box A$;
2. eliminate the cut of $x : \Box A$ in the conclusion of $\Box_L - K$ and in the second premiss of $\Box_L - A$ by TIH, obtaining the sequent $Rxz, z : A, \Gamma, \Gamma' \Rightarrow \Delta, \Delta'$;
3. substitute the eigenvariable in the premiss of $\Box_L - K$ to match that one of 1. and cut $z : \Box A$, obtaining by SIH $Rxz, Rxz, \Gamma, \Gamma', \Gamma \Rightarrow \Delta, \Delta', \Delta, z : A$ without cut;
4. eliminate the cut of $z : A$ between 3. and 2. by IH, obtaining by contraction $Rxz, \Gamma, \Gamma' \Rightarrow \Delta, \Delta'$, as desired.



Tree Hypersequents

Recall that a sequent is a pair $X \Rightarrow Y$ where X and Y are finite *multisets* of formulas.

A **tree hypersequent** (THS) is defined as follows:

- Every sequent is a THS
- if \mathcal{S} is a sequent and G_1, \dots, G_n are THS, then $\mathcal{S}/G_1; \dots; G_n$ is a THS.

The intended interpretation I of a THS is

$$(X \Rightarrow Y)^I := \bigwedge X \rightarrow \bigvee Y$$

$$(\mathcal{S}/G_1; \dots; G_n)^I := \mathcal{S}^I \vee \Box G_1^I \vee \dots \vee \Box G_n^I.$$

Tree Hypersequents

Recall that a sequent is a pair $X \Rightarrow Y$ where X and Y are finite *multisets* of formulas.

A **tree hypersequent** (THS) is defined as follows:

- Every sequent is a THS
- if \mathcal{S} is a sequent and G_1, \dots, G_n are THS, then $\mathcal{S}/G_1; \dots; G_n$ is a THS.

The intended interpretation I of a THS is

$$(X \Rightarrow Y)^I := \bigwedge X \rightarrow \bigvee Y$$

$$(\mathcal{S}/G_1; \dots; G_n)^I := \mathcal{S}^I \vee \Box G_1^I \vee \dots \vee \Box G_n^I.$$

Tree Hypersequents

Two sequents $\mathcal{S}_1, \mathcal{S}_2$ have *equivalent position* in G and H (and we write $G\{\mathcal{S}_1\} \sim H\{\mathcal{S}_2\}$) if

- $G \equiv \mathcal{S}_1$ and $H \equiv \mathcal{S}_2$, or
- $G \equiv \mathcal{S}_1/\underline{X}_1$ and $H \equiv \mathcal{S}_2/\underline{X}_2$, or
- $G \equiv \mathcal{T}_1/(H_1\{\mathcal{S}_1\}; \underline{X}_1) \sim \mathcal{T}_2/(H_2\{\mathcal{S}_2\}; \underline{X}_2) \equiv H$ where $H_1\{\mathcal{S}_1\} \sim H_2\{\mathcal{S}_2\}$, $\mathcal{T}_1, \mathcal{T}_2$ are sequents and $\underline{X}_1, \underline{X}_2$ are sequences of THS.

Cut Rule

Let $X \Rightarrow Y \otimes U \Rightarrow V := X, U \Rightarrow Y, V$.

For $H\{\mathcal{S}\} \sim H'\{\mathcal{S}'\}$ define $H\{\mathcal{S}\} \star H'\{\mathcal{S}'\}$ as follows:

- $\mathcal{S} \star \mathcal{S}' := \mathcal{S} \otimes \mathcal{S}'$
- $(\mathcal{S}/\underline{X}) \star (\mathcal{S}'/\underline{Y}) := \mathcal{S} \otimes \mathcal{S}'/\underline{X}; \underline{Y}$
- $(\mathcal{T}/H\{\mathcal{S}\}; \underline{X}) \star (\mathcal{T}'/H'\{\mathcal{S}'\}; \underline{Y}) := \mathcal{T} \otimes \mathcal{T}'/H\{\mathcal{S}\} \star H'\{\mathcal{S}'\}; \underline{X}; \underline{Y}$
where $\mathcal{T}, \mathcal{T}'$ are sequents.

The Rule

For $G\{X \Rightarrow Y, A\}$ and $G'\{A, U \Rightarrow V\}$

s.t. $G\{X \Rightarrow Y, A\} \sim G'\{A, U \Rightarrow V\}$ define

$$\frac{G\{X \Rightarrow Y, A\} \quad G'\{A, U \Rightarrow V\}}{G\{X \Rightarrow Y\} \star G'\{U \Rightarrow V\}} \text{Cut}$$

Note that *Cut* merges trees, so that its conclusion is a THS indeed

Cut Rule

Let $X \Rightarrow Y \otimes U \Rightarrow V := X, U \Rightarrow Y, V$.

For $H\{\mathcal{S}\} \sim H'\{\mathcal{S}'\}$ define $H\{\mathcal{S}\} \star H'\{\mathcal{S}'\}$ as follows:

- $\mathcal{S} \star \mathcal{S}' := \mathcal{S} \otimes \mathcal{S}'$
- $(\mathcal{S}/\underline{X}) \star (\mathcal{S}'/\underline{Y}) := \mathcal{S} \otimes \mathcal{S}'/\underline{X}; \underline{Y}$
- $(\mathcal{T}/H\{\mathcal{S}\}; \underline{X}) \star (\mathcal{T}'/H'\{\mathcal{S}'\}; \underline{Y}) := \mathcal{T} \otimes \mathcal{T}'/H\{\mathcal{S}\} \star H'\{\mathcal{S}'\}; \underline{X}; \underline{Y}$
where $\mathcal{T}, \mathcal{T}'$ are sequents.

The Rule

For $G\{X \Rightarrow Y, A\}$ and $G'\{A, U \Rightarrow V\}$

s.t. $G\{X \Rightarrow Y, A\} \sim G'\{A, U \Rightarrow V\}$ define

$$\frac{G\{X \Rightarrow Y, A\} \quad G'\{A, U \Rightarrow V\}}{G\{X \Rightarrow Y\} \star G'\{U \Rightarrow V\}} \text{Cut}$$

Note that *Cut* merges trees, so that its conclusion is a THS indeed

Labelled Tree Sequents

Let \mathcal{R} denote a set of relational terms.

We say that \mathcal{R} is a tree when the frame it defines is a tree (possibly the empty one).

A **labelled tree sequent** (LTS) is a labelled sequent $\mathcal{R}, \Gamma \Rightarrow \Delta$ where

- \mathcal{R} is a tree, and
- if $\mathcal{R} = \emptyset$, then all formulas in the sequent have the same label;
- otherwise, each label occurring in Γ or Δ occurs in \mathcal{R} also.

Cut Rule

Note first that if \mathcal{R}_1 and \mathcal{R}_2 are tree, $\mathcal{R}_1 \cup \mathcal{R}_2$ might not be so.

The Rule

$$\frac{\mathcal{R}, \Gamma \Rightarrow \Delta, x : A \quad \mathcal{R}, x : A, \Gamma \Rightarrow \Delta}{\mathcal{R}, \Gamma \Rightarrow \Delta} \textit{Cut}$$

Translation

TL

For a label x define $\mathsf{TL}_x:\mathsf{THS}\rightarrow\mathsf{LTS}$ as follows:

- $\mathsf{TL}_x(X \Rightarrow Y) := x : X \Rightarrow x : Y$
- $\mathsf{TL}_x(X \Rightarrow Y/G_1; \cdots; G_n) :=$
 $(\bigotimes_{j=1}^n \mathsf{TL}_{x_j}(G_j)) \otimes (Rxx_1, Rxx_2, \cdots, Rxx_n, x : X \Rightarrow x : Y),$

where $(\mathcal{R}_1, X \Rightarrow Y) \otimes (\mathcal{R}_2, U \Rightarrow V) := \mathcal{R}_1 \cup \mathcal{R}_2, X, U \Rightarrow Y, V.$

LT

For any LTS $\mathcal{R}, \Gamma \Rightarrow \Delta$, define $\mathsf{LT}:\mathsf{LTS}\rightarrow\mathsf{THS}$ as follows:

- if $\mathcal{R} = \emptyset$, then $\mathsf{LT}(x : \Gamma \Rightarrow x : \Delta) := \Gamma \Rightarrow \Delta$
- otherwise, if x is the root of $\mathcal{R} \equiv \{Rxy_1, \cdots, Rxy_n\}$, let $v_i := \{z : Rxy_i z\}$; then $\mathsf{LT}(\mathcal{R}, \Gamma \Rightarrow \Delta) :=$

$$\Gamma_x \Rightarrow \Delta_x / \mathsf{LT}(\mathcal{R}_{v_1}, \Gamma_{v_1} \Rightarrow \Delta_{v_1}); \cdots; \mathsf{LT}(\mathcal{R}_{v_n}, \Gamma_{v_n} \Rightarrow \Delta_{v_n}).$$

Translation

TL

For a label x define $\mathbb{TL}_x : \text{THS} \rightarrow \text{LTS}$ as follows:

- $\mathbb{TL}_x(X \Rightarrow Y) := x : X \Rightarrow x : Y$
- $\mathbb{TL}_x(X \Rightarrow Y/G_1; \dots; G_n) :=$
 $(\bigotimes_{j=1}^n \mathbb{TL}_{x_j}(G_j)) \otimes (Rxx_1, Rxx_2, \dots, Rxx_n, x : X \Rightarrow x : Y),$

where $(\mathcal{R}_1, X \Rightarrow Y) \otimes (\mathcal{R}_2, U \Rightarrow V) := \mathcal{R}_1 \cup \mathcal{R}_2, X, U \Rightarrow Y, V.$

LT

For any LTS $\mathcal{R}, \Gamma \Rightarrow \Delta$, define $\mathbb{LT} : \text{LTS} \rightarrow \text{THS}$ as follows:

- if $\mathcal{R} = \emptyset$, then $\mathbb{LT}(x : \Gamma \Rightarrow x : \Delta) := \Gamma \Rightarrow \Delta$
- otherwise, if x is the root of $\mathcal{R} \equiv \{Rxy_1, \dots, Rxy_n\}$, let $v_i := \{z : R y_i z\}$; then $\mathbb{LT}(\mathcal{R}, \Gamma \Rightarrow \Delta) :=$

$$\Gamma_x \Rightarrow \Delta_x / \mathbb{LT}(\mathcal{R}_{v_1}, \Gamma_{v_1} \Rightarrow \Delta_{v_1}); \dots; \mathbb{LT}(\mathcal{R}_{v_n}, \Gamma_{v_n} \Rightarrow \Delta_{v_n}).$$

Substitution

Let Lbl be set of labels, and $Var(\mathcal{S}) \subset Lbl$ the finite set of labels occurring in the labelled sequent \mathcal{S} . A renaming of \mathcal{S} consists of a one-to-one function $f_{\mathcal{S}} : Var(\mathcal{S}) \rightarrow Lbl$.

For \mathcal{S}' and a renaming $f_{\mathcal{S}}$, let $\mathcal{S}'_{f_{\mathcal{S}}}$ the labelled sequent obtained by substituting in \mathcal{S}' x with $f_{\mathcal{S}}(x)$ for any $x \in Dom(f_{\mathcal{S}}) \cap Var(\mathcal{S}')$.

Note that if \mathcal{S}' is a LTS, $\mathcal{S}'_{f_{\mathcal{S}}}$ needs not to be a LTS, but it is so for \mathcal{S} LTS.

Lemma

Let G be a THS and \mathcal{S} a LTS. Then $LT(TL_x(G)) = G$, and $TL_x(LT(\mathcal{S})) = \mathcal{S}_{f_{\mathcal{S}}}$ for some $f_{\mathcal{S}}$.

Proof.

By definition of LTS, we cannot loose variables applying LT to \mathcal{S} ; however a renaming might be necessary, to obtain the desired equality, since TL does assign variables. The equality $LT(TL_x(G)) = G$ holds by definitions of the translations.

Substitution

Let Lbl be set of labels, and $Var(\mathcal{S}) \subset Lbl$ the finite set of labels occurring in the labelled sequent \mathcal{S} . A renaming of \mathcal{S} consists of a one-to-one function $f_{\mathcal{S}} : Var(\mathcal{S}) \rightarrow Lbl$.

For \mathcal{S}' and a renaming $f_{\mathcal{S}}$, let $\mathcal{S}'_{f_{\mathcal{S}}}$ the labelled sequent obtained by substituting in \mathcal{S}' x with $f_{\mathcal{S}}(x)$ for any $x \in Dom(f_{\mathcal{S}}) \cap Var(\mathcal{S}')$.

Note that if \mathcal{S}' is a LTS, $\mathcal{S}'_{f_{\mathcal{S}}}$ needs not to be a LTS, but it is so for \mathcal{S} LTS.

Lemma

Let G be a THS and \mathcal{S} a LTS. Then $LT(TL_x(G)) = G$, and $TL_x(LT(\mathcal{S})) = \mathcal{S}_{f_{\mathcal{S}}}$ for some $f_{\mathcal{S}}$.

Proof.

By definition of LTS, we cannot loose variables applying LT to \mathcal{S} ; however a renaming might be necessary, to obtain the desired equality, since TL does assign variables. The equality $LT(TL_x(G)) = G$ holds by definitions of the translations.

Substitution

Let Lbl be set of labels, and $Var(\mathcal{S}) \subset Lbl$ the finite set of labels occurring in the labelled sequent \mathcal{S} . A renaming of \mathcal{S} consists of a one-to-one function $f_{\mathcal{S}} : Var(\mathcal{S}) \rightarrow Lbl$.

For \mathcal{S}' and a renaming $f_{\mathcal{S}}$, let $\mathcal{S}'_{f_{\mathcal{S}}}$ the labelled sequent obtained by substituting in \mathcal{S}' x with $f_{\mathcal{S}}(x)$ for any $x \in Dom(f_{\mathcal{S}}) \cap Var(\mathcal{S}')$.

Note that if \mathcal{S}' is a LTS, $\mathcal{S}'_{f_{\mathcal{S}}}$ needs not to be a LTS, but it is so for \mathcal{S} LTS.

Lemma

Let G be a THS and \mathcal{S} a LTS. Then $\mathbb{LT}(\mathbb{TL}_x(G)) = G$, and $\mathbb{TL}_x(\mathbb{LT}(\mathcal{S})) = \mathcal{S}_{f_{\mathcal{S}}}$ for some $f_{\mathcal{S}}$.

Proof.

By definition of LTS, we cannot loose variables applying \mathbb{LT} to \mathcal{S} ; however a renaming might be necessary, to obtain the desired equality, since \mathbb{TL} does assign variables. The equality $\mathbb{LT}(\mathbb{TL}_x(G)) = G$ holds by definitions of the translations.

Substitution Lemma

Let C be a LTS calculus, and \mathcal{S} a LTS. Let $f_{\mathcal{S}}$ be a renaming of \mathcal{S} . If \mathcal{S} is derivable in C with height n , then so is $\mathcal{S}_{f_{\mathcal{S}}}$.

Proof.

Let δ be the derivation of \mathcal{S} with height n . We proceed by induction on n :

- if $n = 0$, then \mathcal{S} is initial, so that $\mathcal{S}_{f_{\mathcal{S}}}$ is initial too.
- if $n > 0$, and ρ is the last rule of δ (from $\mathcal{S}_1, \dots, \mathcal{S}_n$ to \mathcal{S}) then $(\mathcal{S}_i)_{f_{\mathcal{S}}}$ might not be a LTS, since $\bigcup_i \text{Var}(\mathcal{S}_i)$ may contain labels that do not occur in \mathcal{S} . Hence define $g : \bigcup_i \text{Var}(\mathcal{S}_i) \setminus \text{Dom}(f_{\mathcal{S}}) \rightarrow \text{Lbl} \setminus \text{Im}(f_{\mathcal{S}})$. Then the composition $f_{\mathcal{S}} \circ g$ is a renaming of \mathcal{S}_i 's. By applying the IH to the \mathcal{S}_i 's and then applying ρ , we have a derivation of $((\mathcal{S})_g)_{f_{\mathcal{S}}} = \mathcal{S}_{f_{\mathcal{S}}}$.



Substitution Lemma

Let C be a LTS calculus, and \mathcal{S} a LTS. Let $f_{\mathcal{S}}$ be a renaming of \mathcal{S} . If \mathcal{S} is derivable in C with height n , then so is $\mathcal{S}_{f_{\mathcal{S}}}$.

Proof.

Let δ be the derivation of \mathcal{S} with height n . We proceed by induction on n :

- if $n = 0$, then \mathcal{S} is initial, so that $\mathcal{S}_{f_{\mathcal{S}}}$ is initial too.
- if $n > 0$, and ρ is the last rule of δ (from $\mathcal{S}_1, \dots, \mathcal{S}_n$ to \mathcal{S}) then $(\mathcal{S}_i)_{f_{\mathcal{S}}}$ might not be a LTS, since $\bigcup_i \text{Var}(\mathcal{S}_i)$ may contain labels that do not occur in \mathcal{S} . Hence define $g : \bigcup_i \text{Var}(\mathcal{S}_i) \setminus \text{Dom}(f_{\mathcal{S}}) \mapsto \text{Lbl} \setminus \text{Im}(f_{\mathcal{S}})$. Then the composition $f_{\mathcal{S}} \circ g$ is a renaming of \mathcal{S}_i 's. By applying the IH to the \mathcal{S}_i 's and then applying ρ , we have a derivation of $((\mathcal{S})_g)_{f_{\mathcal{S}}} = \mathcal{S}_{f_{\mathcal{S}}}$.



Induced Rules

$$\frac{\overbrace{\mathcal{R}, Rxy, y : \Box A, \Gamma \Rightarrow \Delta, y : A}^{\text{principal}}}{\mathcal{R}, \Gamma \Rightarrow \Delta, \underbrace{x : \Box A}_{\text{principal}}} \quad \xrightarrow{\text{LT}} \quad \frac{G\{X \Rightarrow Y / \Box A \Rightarrow A\}}{G\{X \Rightarrow Y, \Box A\}}$$

$$\frac{G\{\overbrace{X \Rightarrow Y}^x / \overbrace{\Box A \Rightarrow A}^y\}}{G\{\overbrace{X \Rightarrow Y, \Box A}^x / \underbrace{\emptyset}_y\}} \quad \xrightarrow{\text{TL}} \quad \frac{\mathcal{R}, Rxy, y : \Box A, \Gamma \Rightarrow \Delta, y : A}{\mathcal{R}, \Gamma \Rightarrow \Delta, x : \Box A}$$

Induced Calculus

Translation Lemma

Let C be a THS calculus. Then the following hold:

- (i) $C \vdash^n G$ iff $\text{TLC} \vdash^n \text{TL}G$
- (ii) $\text{TLC} \vdash^n S$ iff $C \vdash^n \text{LTS}$.

Proof.

To (i): Let δ be a derivation of G in C with height n . Then by substituting each THS H in δ with $\text{TL}H$ and any rule ρ in δ with $\text{TL}(\rho)$ we obtain a derivation of $\text{TL}G$ whose height is n by construction.

To (ii): We can reason analogously to (i).



Corollary

For any THS calculus C and any formula A , $C \vdash \Rightarrow A$ iff $\text{TLC} \vdash \Rightarrow x : A$.

Induced Calculus

Translation Lemma

Let C be a THS calculus. Then the following hold:

- (i) $C \vdash^n G$ iff $\text{TLC} \vdash^n \text{TL}G$
- (ii) $\text{TLC} \vdash^n S$ iff $C \vdash^n \text{LTS}$.

Proof.

To (i): Let δ be a derivation of G in C with height n . Then by substituting each THS H in δ with $\text{TL}H$ and any rule ρ in δ with $\text{TL}(\rho)$ we obtain a derivation of $\text{TL}G$ whose height is n by construction.

To (ii): We can reason analogously to (i).



Corollary

For any THS calculus C and any formula A , $C \vdash \Rightarrow A$ iff $\text{TLC} \vdash \Rightarrow x : A$.

Induced Calculus

Translation Lemma

Let C be a THS calculus. Then the following hold:

- (i) $C \vdash^n G$ iff $\text{TLC} \vdash^n \text{TL}G$
- (ii) $\text{TLC} \vdash^n S$ iff $C \vdash^n \text{LTS}$.

Proof.

To (i): Let δ be a derivation of G in C with height n . Then by substituting each THS H in δ with $\text{TL}H$ and any rule ρ in δ with $\text{TL}(\rho)$ we obtain a derivation of $\text{TL}G$ whose height is n by construction.

To (ii): We can reason analogously to (i).



Corollary

For any THS calculus C and any formula A , $C \vdash \Rightarrow A$ iff $\text{TLC} \vdash \Rightarrow x : A$.

THSGL

Let **THSGL** denote the following THS calculus for \mathbb{GL} :

Initial THS: $G\{p, X \Rightarrow Y, p\} \quad G\{\Box A, X \Rightarrow Y, \Box A\}$

Propositional rules:

$$\frac{G\{X \Rightarrow Y, A\}}{G\{\neg A, X \Rightarrow Y\}} \neg A \quad \frac{G\{A, X \Rightarrow Y\}}{G\{X \Rightarrow Y, \neg A\}} \neg K$$

$$\frac{G\{A, B, X \Rightarrow Y\}}{G\{A \wedge B, X \Rightarrow Y\}} \wedge A \quad \frac{G\{X \Rightarrow Y, A\} \quad G\{X \Rightarrow Y, B\}}{G\{X \Rightarrow Y, A \wedge B\}} \wedge K$$

Modal rules:

$$\frac{G\{\Box A, X \Rightarrow Y/(U \Rightarrow V, \Box A/\underline{X})\} \quad G\{\Box A, X \Rightarrow Y/(A, U \Rightarrow V/\underline{X})\}}{G\{\Box A, X \Rightarrow Y/(U \Rightarrow V/\underline{X})\}} \Box A$$

$$\frac{G\{X \Rightarrow Y/\Box A \Rightarrow A\}}{G\{X \Rightarrow Y, \Box A\}} \Box K$$

Special logical rule:

$$\frac{G\{\Box A, X \Rightarrow Y/(\Box A, U \Rightarrow V/\underline{X})\}}{G\{\Box A, X \Rightarrow Y/(U \Rightarrow V/\underline{X})\}} 4$$

Structural Properties

Lemma

For THSGL the following hold:

- (i) Weakening rules are hp admissible;
- (ii) All rules are invertible;
- (iii) Contraction rules are admissible.

Proof.

To (i): By induction on the height of the derivation.

To (ii): By induction on the height of the derivation.

To (iii): By simultaneous induction for left and right contraction, with primary induction on the contraction formula, and secondary induction on the height of the derivation.



Cut Elimination

Cut Elimination Theorem

The cut rule is admissible in THSGL.

Proof.

By induction on the cut formula, with secondary induction on the **position** of the cut, and ternary induction on the height of the cut.

The position of a sequent S in a THS G in a derivation δ is given by the difference between the length of the longest THS occurring in δ and the number of sequents that precede S .



Cut Elimination

Cut Elimination Theorem

The cut rule is admissible in THSGL.

Proof.

By induction on the cut formula, with secondary induction on the **position** of the cut, and ternary induction on the height of the cut.

The position of a sequent S in a THS G in a derivation δ is given by the difference between the length of the longest THS occurring in δ and the number of sequents that precede S .



TLTHSGL

By applying the map \mathbb{TL} to THSGL we obtain the following LTS calculus:

Initial LTS: $\mathcal{R}, x : p, \Gamma \Rightarrow \Delta, x : p \quad \mathcal{R}, x : \Box A, \Gamma \Rightarrow \Delta, x : \Box A$

Propositional rules:

$$\frac{\mathcal{R}, \Gamma \Rightarrow \Delta, x : A}{\mathcal{R}, x : \neg A, \Gamma \Rightarrow \Delta} \neg A \quad \frac{\mathcal{R}, x : A, \Gamma \Rightarrow \Delta}{\mathcal{R}, \Gamma \Rightarrow \Delta, x : \neg A} \neg K$$

$$\frac{\mathcal{R}, x : A, x : B, \Gamma \Rightarrow \Delta}{\mathcal{R}, x : A \wedge B, \Gamma \Rightarrow \Delta} \wedge A \quad \frac{\mathcal{R}, \Gamma \Rightarrow \Delta, x : A \quad \mathcal{R}, \Gamma \Rightarrow \Delta, x : B}{\mathcal{R}, \Gamma \Rightarrow \Delta, x : A \wedge B} \wedge K$$

Modal rules:

$$\frac{\mathcal{R}, Rxy, x : \Box A, \Gamma \Rightarrow \Delta, y : \Box A \quad \mathcal{R}, Rxy, x : \Box A, y : A, \Gamma \Rightarrow \Delta}{\mathcal{R}, Rxy, x : \Box A, \Gamma \Rightarrow \Delta} \mathbb{TL}\Box A$$

$$\frac{\mathcal{R}, Rxy, y : \Box A, \Gamma \Rightarrow \Delta, y : A}{\mathcal{R}, \Gamma \Rightarrow \Delta, x : \Box A} \mathbb{TL}\Box K$$

Special logical rule:

$$\frac{\mathcal{R}, Rxy, x : \Box A, y : \Box A, \Gamma \Rightarrow \Delta}{\mathcal{R}, Rxy, x : \Box A, \Gamma \Rightarrow \Delta} \mathbb{TL}4$$

Note three main **differences** from G3GL:

- \mathcal{R} is a tree;
- $\mathbb{TL}4$ does not appear in G3GL;
- *Irref* and *Trans* do not appear in $\mathbb{TLTHSGL}$.

Structural Properties

Lemma

Both weakening and contraction rules are admissible in $\mathsf{TLTHSGL}$.

Proof.

It follows by the definition TL and the translation lemma since the corresponding results hold for THSGL .



Lemma

Let δ be a derivation in $\mathsf{G3GL} + \mathsf{TL4}$ and assume that δ does not contain any occurrence of Irref . Then Trans can be eliminated in δ .

Proof.

Assume Trans occurs $s + 1$ -times in δ , and let ρ be the topmost occurrence of Trans :

$$\frac{Rxy, Ryz, Rxz, \Gamma \Rightarrow \Delta}{Rxy, Ryz, \Gamma' \Rightarrow \Delta'} \rho$$

Let $w(\rho)$ denote the number of rules above ρ that make Rxz principal. We proceed by induction on $w(\rho)$.

Structural Properties

Lemma

Both weakening and contraction rules are admissible in $\mathbb{T}\mathbb{L}\mathbb{T}\mathbb{H}\mathbb{S}\mathbb{G}\mathbb{L}$.

Proof.

It follows by the definition $\mathbb{T}\mathbb{L}$ and the translation lemma since the corresponding results hold for $\mathbb{T}\mathbb{H}\mathbb{S}\mathbb{G}\mathbb{L}$.



Lemma

Let δ be a derivation in $\mathbb{G}\mathbb{3}\mathbb{G}\mathbb{L} + \mathbb{T}\mathbb{L}4$ and assume that δ does not contain any occurrence of *Irref*. Then *Trans* can be eliminated in δ .

Proof.

Assume *Trans* occurs $s + 1$ -times in δ , and let ρ be the topmost occurrence of *Trans*:

$$\frac{Rxy, Ryz, Rxz, \Gamma \Rightarrow \Delta}{Rxy, Ryz, \Gamma' \Rightarrow \Delta'} \rho$$

Let $w(\rho)$ denote the number of rules above ρ that make *Rxz* principal. We proceed by induction on $w(\rho)$.

Structural Properties

Proof.

- if $w(\rho) = 0$, then Rxz is not principal in any rule above ρ , then it can be eliminated and we obtain a derivation with s occurrences of *Trans*;
- if $w(\rho) = k + 1$, then since the derivation above ρ does not contain *Irref* nor *Trans*, Rxz can be principal just by TL4 or $\text{TL}\Box A$:
 - if Rxz is principal by $\text{TL}\Box A$, we have

$$\begin{array}{c}
 Rxy, Ryz, Rxz, \mathcal{R}', x : \Box B, \Gamma' \Rightarrow \Delta', z : \Box B \\
 Rxy, Ryz, Rxz, \mathcal{R}', x : \Box B, z : B, \Gamma' \Rightarrow \Delta' \\
 \hline
 Rxy, Ryz, Rxz, \mathcal{R}', x : \Box B, \Gamma' \Rightarrow \Delta' \quad \text{TL}\Box A \\
 \\
 \vdots \\
 \hline
 \frac{Rxy, Ryz, Rxz, \mathcal{R}, \Gamma \Rightarrow \Delta}{Rxy, Ryz, \mathcal{R}, \Gamma \Rightarrow \Delta} \rho
 \end{array}$$

Then we can weaken the premises with $y : \Box B$, then apply $\text{TL}\Box A$ with Ryz principal formula and TL4 with Rxy principal. This way $w(\rho) = k$, and we apply IH;

Structural Properties

Proof.

- if Rxz is principal by $\mathbb{TL4}$, we have

$$\frac{Rxy, Ryz, Rxz, \mathcal{R}', x : \Box B, z : \Box B, \Gamma' \Rightarrow \Delta'}{Rxy, Ryz, Rxz, \mathcal{R}', x : \Box B, \Gamma' \Rightarrow \Delta'} \mathbb{TL4}$$

$$\vdots$$

$$\frac{Rxy, Ryz, Rxz, \mathcal{R}, \Gamma \Rightarrow \Delta}{Rxy, Ryz, \mathcal{R}, \Gamma \Rightarrow \Delta} \rho$$

Then we can weaken with $y : \Box B$, then apply $\mathbb{TL4}$ with Ryz principal formula, and again $\mathbb{TL4}$ with Rxy principal, so that $w(\rho) = k$ and we can use IH.

The result now follows by induction on s .



Cut Elimination

Adequacy Theorem

For any formula A , $\mathbb{T}\mathbb{L}\text{THSGL} \vdash \Rightarrow x : A$ iff $\text{G3GL} \vdash \Rightarrow x : A$.

Cut Elimination Theorem

The cut rule is admissible for THSGL.

Proof.

It follows by the corresponding result for G3GL along with the translation lemma. □

Cut Elimination

Adequacy Theorem

For any formula A , $\mathbb{T}\mathbb{L}\text{THSGL} \vdash \Rightarrow x : A$ iff $\text{G3GL} \vdash \Rightarrow x : A$.

Cut Elimination Theorem

The cut rule is admissible for THSGL.

Proof.

It follows by the corresponding result for G3GL along with the translation lemma. □

Cut Elimination

Adequacy Theorem

For any formula A , $\mathbb{TL}4\text{HSGL} \vdash \Rightarrow x : A$ iff $\text{G3GL} \vdash \Rightarrow x : A$.

Proof.

To \Rightarrow : It suffices to prove that $\mathbb{TL}4$ is admissible in G3GL. Assume then that $\mathcal{R}, Rxy, y : \Box A, x : \Box A, \Gamma \Rightarrow \Delta$ is derived. Consider now

$$\begin{array}{c}
 \frac{z : \Box A \Rightarrow z : \Box A}{Rxx, x : \Box A, z : \Box A \Rightarrow z : A, z : \Box A} \quad \frac{z : A \Rightarrow z : A}{Rxx, z : A, x : \Box A, z : \Box A \Rightarrow z : A} \\
 \hline
 \frac{Rxx, x : \Box A, z : \Box A \Rightarrow z : A}{Rxy, Ryz, Rxx, x : \Box A, z : \Box A \Rightarrow z : A} \text{TL}\Box A \\
 \hline
 \frac{Rxy, Ryz, Rxx, x : \Box A, z : \Box A \Rightarrow z : A}{Rxy, Ryz, x : \Box A, z : \Box A \Rightarrow z : A} \text{LW} \\
 \hline
 \frac{Rxy, Ryz, x : \Box A, z : \Box A \Rightarrow z : A}{Rxy, x : \Box A \Rightarrow y : \Box A} (\text{Trans}) \\
 \hline
 Rxy, x : \Box A \Rightarrow y : \Box A \quad \text{TL}\Box K
 \end{array}$$

By applying contractions and *Cut* we obtain $Rxy, \mathcal{R}, x : \Box A, \Gamma \Rightarrow \Delta$ which is turned into a contraction- and cut-free derivation by contraction admissibility and cut elimination for G3GL.

Cut Elimination

Adequacy Theorem

For any formula A , $\mathbb{T}\mathbb{L}\mathbb{T}\mathbb{H}\mathbb{S}\mathbb{G}\mathbb{L} \vdash \Rightarrow x : A$ iff $\mathbb{G}\mathbb{3}\mathbb{G}\mathbb{L} \vdash \Rightarrow x : A$.

Proof.

To \Leftarrow : Note that in any derivation in G3GL, a label can disappear only by $\Box_L - K$ rule when it occurs in a relational term Rxy with $x \neq y$. Now if in the derivation δ of $\Rightarrow x : A$ *Irref* occurs, we could not have empty antecedent in the final sequent, therefore δ does not contain *Irref*. By the previous lemma, $\Rightarrow x : A$ is derivable in $\mathbb{G}\mathbb{3}\mathbb{G}\mathbb{L} + \mathbb{T}\mathbb{L}\mathbb{4} - \text{Trans} - \text{Irref}$. Moreover such a derivation involves only LTS, for, by inspection of the rules in $\mathbb{G}\mathbb{3}\mathbb{G}\mathbb{L} + \mathbb{T}\mathbb{L}\mathbb{4} - \text{Trans} - \text{Irref}$, we see that each rule, read upwards, preserves LTS: since $\Rightarrow x : A$ is LTS, we have the result.



Questions

By means of Negri's and Poggiolesi's calculi we have uncontroversial **syntactic** cut elimination for provability logic.

(Do you recall its long history, until the mechanized proof in HOL for multiset version of the original sequent calculus for \mathbb{GL} ?)

As Blamey and Humberstone state:

[T]he move from truth-functional to modal logic is not one best made simply by adding a new primitive connective with new rules governing it, but rather by extending one's conception of the objects to be manipulated by such rules.

By applying Goré and Ramanayake's translation lemma to THSGL we see that the construction of tree-labelled and \neg -hypersequent calculi has a "common" structure.

(Kind of) Psychoanalytic Question

What mathematical structure is behind these calculi?

Could we extend this construction to different non-classical operators?

Questions

We have strongly well-behaved calculi for Gödel-Löb provability logic, and it seems easy to construct intuitionistic versions of them. Are there results concerning interpretability logics, and Solovay logic?

Questions about Interpretability

Is it possible to extend these calculi to capture ILL and other provability & interpretability logics?

Should we use *standard* Veltman semantics or *generalized* Veltman semantics?

- ⇒ Mixing labels for worlds and neighbourhoods?
- ⇒ How to “index” THS?

Questions

Recall that Solovay theorem can be proved (as far as *everybody* knows) *only* by encoding a Kripke countermodel in our arithmetical language.

Main Question

Is it possible to develop a **syntactic proof of Solovay theorem**, for instance by encoding THS, or LTS?

Related to previous questions: Could we prove *syntactically* Berarducci arithmetical completeness theorem for ILM?

Questions

Recall that Solovay theorem can be proved (as far as *everybody* knows) *only* by encoding a Kripke countermodel in our arithmetical language.

Main Question

Is it possible to develop a **syntactic proof of Solovay theorem**, for instance by encoding THS, or LTS?

Related to previous questions: Could we prove *syntactically* Berarducci arithmetical completeness theorem for ILLM ?