Where the Ladder Must Stand

A test for forced mathematical objects, three verdicts, and one measurement that failed

Yuanjie Liu

Independent researcher

yuanjieliu65@gmail.com

Abstract

Mathematicians often say that an object *had to* exist: you cannot subtract in the natural numbers, so the integers appear; you cannot solve $x^2 + 1 = 0$ in the reals, so the complex numbers appear. The same objects are also called inventions, and both descriptions are usually offered without a test. This paper proposes one, in the form of questions that can be answered yes or no, and reports what happened when the test was turned on the author's own work. Question 1: can the demand be met on the same elements? Question 2: among all extensions, is there a smallest one, determined up to a unique relabelling? Question 3: is the result new, judged against a catalogue of known objects? If Question 2 is answered yes, the object is pinned down: any two smallest extensions are the same up to a relabelling, and the relabelling is unique. Three steps of the number tower pass Questions 1 and 2; the step from the rationals to the reals does not, and is recorded as a named obligation instead; a counting invariant proposed inside the same programme is refused, because it is already definable on the same elements. Question 3 broke: the implementation compared the sizes of carriers, and the integers, the rationals and the complex numbers all have the same size, so the test refused everything. The paper argues that the first two questions are worth keeping, that the third needs to compare structures rather than carriers, and that a test which refuses its author's candidates is worth more than one which admits them all.

Keywords: mathematical discovery; minimal extension; initial object; rigidity; number systems; formal verification; negative result.

> *"Whatever can be defined may be questioned."*
>
> — Galactic Empire *(Isaac Asimov)*

1 The puzzle

1.1 A sentence with no test

When mathematicians introduce a number system they usually say something causal: *we could not subtract, so we made the integers.* The sentence is not a description of the

construction. It is a claim about necessity, and it is the claim that does the philosophical work. It is also the claim that is easiest to abuse.

Consider a program that announces a new number α with $\alpha^2 = 7$. Nothing in the announcement is false. The rationals contain no square root of seven, so α does extend them; but the extension is the field everyone already knows, with α written in place of $\sqrt{7}$. A test that asks only 'was there a solution before?' will certify this as new, and will certify a great deal else besides.

The same weakness runs through the standard justifications. 'The natural numbers cannot solve $x + 1 = 0$' is true. It is also true of the one-element set, of a free group, and of the field of rational functions. Each of them contains a solution; none of them is the integers. So the sentence *there was no solution, therefore we added one* does not pick out an object. It picks out a crowd.

1.2 The literature settles what, not why

The tower itself is settled mathematics. The integers can be built as pairs of naturals modulo a difference, or as the group completion of the additive monoid; the rationals as ordered pairs of integers, or as the fraction field; the reals as Dedekind cuts (Dedekind, 1872) or as equivalence classes of Cauchy sequences; the complex numbers as pairs of reals (Ebbinghaus et al., 1991, following Hamilton, 1837) or as the quotient ring $\mathbb{R}[x]/(x^2+1)$. Any two constructions give isomorphic objects; Landau (1930) builds the tower from Peano's axioms with the standard uniqueness statements, and Ebbinghaus et al. (1991) survey several constructions of each system together with their equivalence.

That literature settles *what* the objects are, and it leaves the causal sentence untouched. Mathematicians do have a name for 'had to exist' — the object is *initial*, or *free*, or *universal*, in the relevant category — and the standard fact that an initial object is determined up to a unique isomorphism is in Mac Lane (1971). Model theory supplies a parallel notion: when a model completion exists it is unique up to equivalence, and the comparison embeddings are elementary and unique over the base model (Eklof and Sabbagh, 1971; Hodges, 1993). So the ingredients are old, and this paper claims no new mathematics for them.

Programs that propose mathematics have a long history, from Lenat's AM (Lenat, 1976) to recent systems that use search or learning to generate conjectures and constructions (Davies et al., 2021; Raayoni et al., 2021; Romera-Paredes et al., 2024). Their pattern is consistent: generate candidates, then decide which ones are worth keeping. The decision is made by a human, by a heuristic of interestingness, or by a checker that tests truth or performance rather than novelty. This paper is about the decision step, and should be read as complementary to those systems: a generator supplies candidates, a criterion decides which of them may be called forced.

What is missing, as far as we can tell from the literature cited here, is a *test*: a short list of questions that a reader can put to a claim of the form 'this object had to exist' — including claims made by a program — together with an honest account of which questions can be settled mechanically and which cannot. The textbook statement is about a fixed category. A test has to survive a change of category, because that is where the abuses live.

1.3 What this paper does

The paper builds such a test, states what it can and cannot settle, and then runs it on the programme that produced it. Three results are negative and are reported as such: one step of the tower is not certified, one candidate is refused, and the third question is not implemented. The positive content is small and is stated in the same breath: two questions work, they are construction-invariant, and they are enough to refuse a candidate that a weaker test would have published.

The order of the paper follows the argument. §2 distinguishes four things that 'forced' can mean and says which one is tested. §3 states the test. §4 explains why the second question settles the object. §5 reports the three verdicts. §6 reports the measurement that failed. §7 answers six objections. §8 concludes.

2 Four senses of 'forced'

The word carries at least four different claims, and a paper that does not say which one it means can be attacked from any direction.

(i) *Metaphysical*: the object could not have failed to exist.

(ii) *Conceptual*: the object is already contained in the concept we started from.

(iii) *Practical*: no working alternative exists, given the state of the field.

(iv) *Determined*: every acceptable answer is the same object up to a unique relabelling.

This paper tests (iv) and nothing else. (i) is not touched: we have no argument that the integers could not have failed to exist, and we doubt that a formal test could supply one. (ii) is not touched either; our demands are data, not concepts, and a demand can be met in a way that tells us nothing about the concept that motivated it. (iii) is a claim about a community, and it changes when the community changes; it is not the kind of thing a proof can settle.

The fourth sense is weaker than the first three, and it is the only one that can be settled by reading a proof. It is also the one that Benacerraf's argument leaves standing. Benacerraf (1965) observed that if numbers are sets, then the natural number 4 could be the von Neumann 4 or the Zermelo 4, and nothing in arithmetic tells us which; so either the identification is arbitrary or numbers are not objects at all. Structuralism replies that numbers are positions in a structure rather than objects with an inner nature (Shapiro, 1997). If that is right, then 'the same position' is exactly determination up to relabelling, and nothing stronger is on offer.

We therefore do not claim to have shown that the integers had to exist. We claim to have shown that, given the demand and the company it keeps, nothing else could have answered it. That is not a hedge; it changes what counts as a refutation. A reader who wants metaphysical necessity will find the test silent, and should. A reader who wants to know whether a program's 'discovery' is a renaming will find it decisive.

3 The test

3.1 Three pieces of data

Before any question can be asked, three things must be fixed.

(a) An *ambient theory*: what the new object is allowed to carry.

(b) A *reading* of the old structure inside the new one.

(c) A *demand*: a constant to add, a property to impose, or both.

'Minimal extension' is not an absolute notion, and the reason is easy to see. Given only a system and a missing solution, one may always take a trivial extension (throw in a formal symbol and nothing else) or a chaotic one (throw in a proper class of symbols), and there is no order between them that would single out a winner. A minimum becomes definable only once (a)–(c) are fixed.

The first step of the tower shows why (a) matters, and it is worth stating slowly. The demand is '$x + 1 = 0$ must become solvable'. If we are allowed to stay among structures where addition is only a semigroup operation, we may add a formal ε with $\varepsilon + 1 = 0$ and stop. The result solves the equation and is not the integers: its addition is not a group operation. If instead we first agree to work among commutative rings, the same demand produces the integers, and produces nothing else. The difference is not in the demand; it is in the company the demand is allowed to keep. Changing the ambient theory changes the object, and the change of theory is the real content of the step.

3.2 Question 1: can the demand be met on the same elements?

The question is about the elements, not about the solution. The wording is doing work. Both the natural numbers and the integers are countable, so any ring structure on the integers can be carried over to the natural numbers by a bijection. The statement 'the natural numbers admit no ring structure' is therefore *false*, and an early version of our own formalisation asserted it. The true statement keeps the old structure fixed: no ring structure on $\mathbb{N}$ whose addition and multiplication extend the usual ones solves $x + 1 = 0$. The episode is worth recording because the test caught its author.

A gap, then, is not 'this carrier admits no such structure'. It is 'this carrier, with the structure it already has, cannot support the required one'. The distinction is the formal counterpart of a platitude: you cannot solve an equation by changing what the symbols mean.

3.3 Question 2: is the extension the smallest one?

Among all extensions that meet the demand, is there one that maps into every other, and does exactly one such map exist? When the answer is yes we call the extension *minimal*. The name is traditional — 'the minimal field containing a root' — but the property is not smallness; it is universality, and universality is what makes the object determined. The next section states what follows from it.

Two clarifications, because 'smallest' invites misreading. First, the comparison is between *extensions*, not between carriers: a smaller carrier is neither necessary nor sufficient. Second, 'smallest' is relative to the triple fixed above; the same system and the same demand can have different smallest extensions under different ambient theories, and that is a finding rather than a defect.

3.4 Question 3: is it new?

New relative to what is already known, not relative to the system's own state, which can always be arranged to report novelty. In practice this means an explicit catalogue of known objects, supplied from outside. The requirement is not stylistic. Whether an object is new is a fact about what is already known, and a system that computes novelty from its own state will always find itself original.

§6 reports what happened when this question was implemented. The short version is that the implementation was wrong in a way that only the examples revealed, and that the test as a whole is therefore a two-question test with a third question specified and unimplemented. We prefer to say that at the outset rather than let a reader discover it.

4 Why the second question settles the object

4.1 The statement

If an extension is smallest in the sense above, then:

(1) it maps into every other extension, and the map is unique;

(2) any two smallest extensions are the same up to a relabelling of their elements, and that relabelling is itself unique;

(3) the smallest extension has no symmetries of its own beyond the identity.

None of this mentions rings, order, or completeness. The three statements use four properties of the surrounding category — an identity map, a composition, an underlying-function map, and the fact that maps agreeing on every value are equal — and nothing else. That is why one argument covers the integers, the rationals and the complex numbers at once, and why it can be checked once rather than three times.

4.2 What the theorem is worth

We should be plain about this, because the alternative is to let a reader discover it. Once you *define* 'smallest' as 'maps into everything, uniquely', the first and third statements are restatements of the definition, and the proof is four lines. The content is not the proof. The content is that this is the right definition to have chosen: it is the only one on the table that does not depend on how the object is constructed. Two people who build the complex numbers differently — as pairs of reals, as a quotient ring, as a matrix algebra — get the same answer to Question 2, and their constructions are related by a

relabelling that is itself unique. That is what it means for the object, rather than the construction, to be determined.

A philosophical reader may object that this is precisely the structuralist commonplace, and that a four-line proof of a commonplace is not worth a paper. The objection is right about the proof and wrong about the use. What the four lines buy is *transfer*: because the statement is parameterised by the ambient theory and the reading, the same receipt covers every step of the tower, and the receipt can be checked by machine. The interest lies in what the receipt refuses, not in how hard it was to obtain.

4.3 Pinning the witness

For the step from the reals to the complex numbers, the demand is 'add a constant j with $j \cdot j = -1$'. The extension $(\mathbb{C}, i)$ is smallest, and it has no symmetry beyond the identity. Now drop the constant from the demand and ask only that *some* square root of -1 exists. The same field is still an extension, but it is no longer smallest: the comparison is no longer unique. Identity and complex conjugation are two distinct maps over $\mathbb{R}$, and both are allowed; pinning i is what removes the second one.

The moral is general. Finding a solution and pinning it down are different achievements, and only the second produces an object. A test that accepts the unpinned version will accept a field with a symmetry group, which is exactly the ambiguity it was supposed to remove.

4.4 What the theorem does not settle

The demand 'the order is complete' behaves differently from '$j^2 = -1$', and the difference is worth stating in words rather than symbols. An equation like $j^2 = -1$ is preserved by every ring map: if j satisfies it, so does the image of j. A condition like 'every bounded non-empty set has a least upper bound' is not preserved by ring maps, and it has no term model: there is no expression you can write down, quotient by a relation, and be done. What follows is that the smallest extension cannot be *computed* the way the complex numbers can. It does not follow that no smallest extension exists.

An earlier draft of this paper drew the stronger conclusion. It was wrong, and the correction matters for how the fourth step of the tower is reported: the asymmetry is about computation and about proof debt, not about a metaphysical difference between adjoining a root and completing an order.

5 Three verdicts

We ran the test on four steps of the number tower and on two further pieces of the same programme. Table 1 gives the summary; the text explains the interesting rows.

5.1 Three steps that pass

The integers are the smallest extension of the naturals once we agree to work among commutative rings; the rationals are the smallest extension of the integers once we agree

Case	Same elements	Smallest	New
$\mathbb{N} \to \mathbb{Z}$	proved	proved	not used
$\mathbb{Z} \to \mathbb{Q}$	proved	proved	not used
$\mathbb{Q} \to \mathbb{R}$	proved	*asserted*	not used
$\mathbb{R} \to \mathbb{C}$	proved	proved	not used
counting invariant	*fails* (argued)	—	—

Table 1: What passed, what did not, and what could not be judged. 'Proved' means machine-checked modulo the standard-maps condition discussed in §7; 'asserted' means written down as an obligation and not proved; 'argued' means a prose argument, not a formalised verdict. No entry in the 'New' column was produced by an implementation.

to work among fields; and $(\mathbb{C}, i)$ is the smallest extension of the reals once we pin the square root of -1. The underlying facts are classical (Ebbinghaus et al., 1991); the formulation as a smallest extension, with the old structure pinned, is our packaging. What the test adds is that the three are instances of one statement, and that in each case the answer does not depend on the construction. For the integers, the side condition that the extension really contains the old system also gives characteristic zero for free, which is a pleasant sign that the definition is carrying its weight.

5.2 One step we could not certify

For the step from the rationals to the reals we prove the failure — the rationals are not complete — and we prove that every real is generated by a cut of rationals below it. We do not prove that the reals are the smallest complete ordered field over the rationals. That statement is written down as a named obligation in the formal development.

We think the obligation is true and provable; a page of standard argument about complete ordered fields should do it. We did not do it. Nothing in the development consumes the obligation, so nothing collapses if it is later proved or refuted; what would change is the status of the fourth row of Table 1, which is exactly what should change.

5.3 One candidate we refused

The programme also proposed an invariant of integers, meant to capture the 'tension' between two numbers: count the primes dividing one but not the other, subtract the count in the opposite direction, and call the result $\tau(a, b)$. The proposal was that the surface $\tau = 0$ organises arithmetic, and that τ is a new object in its own right.

The axiomatic part worked out cleanly. Under four natural assumptions the invariant is unique up to sign and equals a difference of counts,

$$\tau(a, b) = \omega(a) - \omega(b), \tag{1}$$

where $\omega(n)$ is the number of distinct prime factors of n. That is the whole content: τ knows how many prime factors each side has and never which ones. Since ω is a century-old function and the formula is definable on the same elements, the test returns *refused* at Question 1. There is no new object here; there is a theorem about ω, and a clear statement of why the construction cannot say more.

Two qualifications. The refusal is argued by hand: the counting part of the development contains no demand and no catalogue, so 'Question 1 returns refused' is a mathematical observation, not a machine-checked verdict. And the refusal does not go through Question 3: the first draft said the novelty score was zero there, which neither follows from Question 1 nor is available, since no catalogue exists.

The episode is still the most useful thing in the paper. It is the case where the test said no to us, and it came with a reason: the invariant collapses to a count, and counts are already on the same elements. A weaker test would have produced a paper claiming a discovery.

6 The measurement that failed

Question 3 is the one we could not deliver, and the failure is instructive enough to describe in detail.

6.1 Two scores

The design had two scores. One was a weighted average of five qualities — coverage, relatedness, closure gain, minimality, novelty — meant to *rank* candidates. The other was a product of four — novelty, consistency, generativity, compression — meant to *admit* them. The distinction is real: a weighted average gives a high mark to a candidate whose novelty is zero but whose other qualities are excellent, which is precisely the $\sqrt{7}$ story; a product gives it zero.

The claim that the two scores are not interchangeable is machine-checked in the only form available: there is a score with novelty zero and generativity $4/5$, and a snapshot with the same entries whose product is zero. We stress what this does and does not show. It shows that the two formulas are not interchangeable. It does not model renaming at all: no catalogue, no candidate, and no map between objects appears in those two statements. The $\sqrt{7}$ story remains an argument, not a theorem.

6.2 Why the implementation failed

Modelling renaming is where the implementation failed. To decide whether a candidate was new, the code compared the *sizes* of carriers: if a candidate's carrier had the same cardinality as a known object's carrier, it was called a renaming. The integers, the rationals and the complex numbers are all countably infinite. One countable object in the catalogue therefore makes all three renamings of it, and the gate assigns all three a novelty of zero. The gate refuses exactly the objects the rest of the paper certifies.

We did not patch this. Comparing sizes can only produce false refusals, never false admissions, so the risk runs in the opposite direction from the one our first draft described. A correct comparison is between structures, not carriers, which requires the catalogue to carry its own maps; that is a specification, and we have not implemented it.

6.3 What follows

Three consequences, stated plainly. First, no admission decision in this paper uses Question 3, and the trace distributed with the code prints novelty values that were entered by hand. Second, the product structure and the zero-factor lemmas survive, because they are statements about a product; what fails is the *measurement* of one factor. Third, the failure is evidence for the test rather than against it: the example that was supposed to demonstrate the gate demolished it, and the demolition was visible only because the gate had been applied to objects whose status we already knew.

7 Objections and replies

7.1 'Your verdicts are relative to chosen data, so they are arbitrary'

Two parts of the objection should be separated. The first is true and is stated in the paper: every verdict is relative to the ambient theory, the reading, and (for Question 3) the catalogue. Change any of them and the verdict can change. The second part — that relativity makes the verdict arbitrary — does not follow. The data are published, so a reader can recompute the verdict with different data and see whether it changes. A relative verdict that anyone can recheck is a different thing from an arbitrary one.

There is also a point of substance behind the relativity. Changing the ambient theory changes the object: the same demand for subtraction produces the integers among rings and a semiring monster among semirings. That is not a defect of the test; it is the phenomenon the test was built to expose. A test that gave the same answer under every ambient theory would be insensitive to precisely the choice that does the work.

7.2 'A degenerate framework makes every object smallest'

This is the most serious objection, and the paper concedes it. The four properties that the main argument uses do not say that the maps between structures are the structure-preserving ones. A degenerate choice — one map between any two structures — satisfies all four, and in that setting every object is both smallest and largest, so the test says nothing about anything.

The paper therefore states a precondition, called C0: the framework is the standard one of the theory at hand. It is not provable inside the framework, because 'being the standard maps' is a schema rather than a sentence; it is checkable in each instance, and in our instances it holds by construction, since the maps are defined to be the structure-preserving ones. Every statement that needs C0 is marked as such.

A test with a stated precondition is better than a test with an unstated one. But the objection does have a residue: the general theorem is less general than its statement suggests, because its content lives in the instances rather than in the abstraction. We accept that description.

7.3 'Benacerraf already showed that determination is all you get'

Correct, and the paper says so. If numbers are positions in a structure, then determination up to a unique relabelling is the most that could be on offer, and the test is not adding a metaphysical result. What it adds is a procedure: given a candidate and a catalogue, a reader can decide whether the word 'discovery' is warranted, and can be shown the step where the decision is made.

Whether that is philosophy is a fair question. Our answer is that it is the part of the philosophy that can be done with a proof rather than a intuition, and that the two refusals reported here — one by proof, one by argument — are the kind of result the procedure produces. A reader who wants the metaphysics will not find it, and we have tried not to pretend otherwise.

7.4 'Your main theorem is a definition'

Yes. The paper says so in the section that states it, and repeats it in the conclusion. The theorem is worth stating because it is the receipt that the definition is well posed, and because the receipt transfers across theories. It is not worth stating as a discovery, and we do not.

The objection would bite if the paper's contribution were the theorem. It is not: the contribution is the triple of questions, the two refusals, and the report of the failure. A reader who judges the paper by the length of its main proof will judge it correctly and will not want to read it.

7.5 'One refusal of your own candidate is not evidence'

It is not much evidence, and the paper says that too. A single refusal, argued by hand, of a candidate the author was already prepared to abandon, is close to the weakest form of evidence a test can produce. What would count as stronger: a candidate proposed by someone else and believed by them, refused by the test; an independent user applying the test to a case neither of us chose; or a repaired, structure-comparing gate that still refuses an object the test ought to admit.

None of these exists yet. We list them because a reader is entitled to know what would change our mind, and because the list is short enough to be a research programme rather than a wish.

7.6 'Machine-checking here is decoration'

Some of it is, and the paper tries to say which parts. The development is in Lean 4 (de Moura et al., 2015) with Mathlib (The Mathlib Community, 2020). The main theorem's proof is short enough that checking it adds little; the four failures are textbook; the two-score separation is arithmetic. Three things were not decoration. The proof assistant forced the definition of 'gap' to be stated in a form that is not false, which is how the ring-structure error was found. It made the four-line nature of the main argument visible, so that the paper could not pretend otherwise. And it turned a claim about two

scoring formulas into a statement a reader can check in a minute. A tool whose main contribution is to make the author's mistakes visible is worth its cost in a paper of this kind.

8 Conclusion

The practical lesson is about the order in which to explain an object. The usual order is: here is the object, here are the properties that make it necessary. The order that survived the test is the reverse. First the failure that cannot be repaired in place. Then the company the repair is allowed to keep, because the same failure repaired in different company gives different objects. Then the smallest repair, which is determined once the company is fixed. Whatever is left over after those three steps is not forced, and should be called a construction.

The second lesson is that a negative result is cheap and worth publishing. The counting invariant was, for a while, the centrepiece of a different paper. Running it through Question 1 took an afternoon and produced a theorem about ω instead. That is a better outcome than a paper claiming a discovery, and it cost less.

The third lesson is about what a test can be. This one is two questions and a failure. It does not generate candidates, it does not decide its own questions, and it cannot certify the fourth step of the tower. What it can do is stop a particular kind of sentence — 'there was no solution, therefore I discovered an object' — from being uttered without a receipt. On the evidence of this paper, that is enough to refuse one candidate and to expose one measurement, and not enough to claim more.

A What is checked, argued, and assumed

Availability. The Lean development, the seven-round self-audit, and a build pinned to a fixed Mathlib revision are in the repository at tag `paper-forced-objects-v1.1` (commit `720ae3b`). The artifact contains only the modules used in this paper; the repository's simulation code is outside its boundary. No theorem in the development consumes the single named axiom.

References

Paul Benacerraf. What numbers could not be. *The Philosophical Review*, 74(1):47–73, 1965.

Alex Davies, Petar Veličković, Lars Buesing, et al. Advancing mathematics by guiding human intuition with AI. *Nature*, 600:70–74, 2021.

Leonardo de Moura, Soonho Kong, Jeremy Avigad, Floris van Doorn, and Jakob von Raumer. The Lean theorem prover (system description). In *Automated Deduction – CADE-25*, volume 9195 of *Lecture Notes in Computer Science*, pages 378–388, 2015. doi: 10.1007/978-3-319-21401-6_26.

Richard Dedekind. *Stetigkeit und irrationale Zahlen*. Vieweg, Braunschweig, 1872.

Claim	Status	Note
Uniqueness and rigidity of minimal extensions	checked	modulo C0
Four failures of the tower	checked	natural, integer, rational, real
$\mathbb{Z}$, $\mathbb{Q}$, $(\mathbb{C}, i)$ are minimal	checked	modulo C0
Characteristic zero of the field extension	checked	follows from injectivity
Two-score separation	checked	arithmetic, not a model of renaming
Counting collapse to $\omega(a) - \omega(b)$	checked	four axioms
Refusal of the counting invariant	argued	no demand, no catalogue instance
Real is the smallest complete ordered field over $\mathbb{Q}$	asserted	obligation, consumed by nothing
Preservation of equations vs closure conditions	argued	about the language, not inside it
Novelty gate	refuted	carrier counting; repair specified, not built

Table 2: The status of every substantive claim. 'Checked' means a Lean 4 declaration with no `sorry` and no named axiom beyond the obligation above; 'argued' means prose; 'asserted' means written down and not proved; 'refuted' means the implementation was shown wrong by the paper's own examples.

H.-D. Ebbinghaus, H. Hermes, F. Hirzebruch, M. Koecher, K. Mainzer, J. Neukirch, A. Prestel, and R. Remmert. *Numbers*, volume 123 of *Graduate Texts in Mathematics*. Springer, New York, 1991. doi: 10.1007/978-1-4612-1005-4.

Paul C. Eklof and Gabriel Sabbagh. Model-completions and modules. *Annals of Mathematical Logic*, 2:251–295, 1971. doi: 10.1016/0003-4843(71)90016-7.

Wilfrid Hodges. *Model Theory*, volume 42 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1993.

Edmund Landau. *Grundlagen der Analysis*. Akademische Verlagsgesellschaft, Leipzig, 1930.

Douglas B. Lenat. *AM: An artificial intelligence approach to discovery in mathematics as heuristic search*. PhD thesis, Stanford University, 1976.

Saunders Mac Lane. *Categories for the Working Mathematician*, volume 5 of *Graduate Texts in Mathematics*. Springer, New York, 1971.

Gal Raayoni, Shahar Gottlieb, Yahel Manor, et al. Generating conjectures on fundamental constants with the Ramanujan Machine. *Nature*, 590:67–73, 2021.

Bernardino Romera-Paredes, Mohammadamin Barekatain, Alexander Novikov, et al. Mathematical discoveries from program search with large language models. *Nature*, 625:468–475, 2024.

Stewart Shapiro. *Philosophy of Mathematics: Structure and Ontology*. Oxford University Press, New York, 1997.

The Mathlib Community. The Lean mathematical library. In *Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020)*, pages 367–381, 2020. doi: 10.1145/3372885.3373824.