

# Week 2 Lab

---

Yuanchao Zhang

January 23, 2018

# Plan for today

1. Changes of lab session administration
2. Recap
3. Install packages
4. The normal distribution
5. Check normality
6. One-sample tests of proportion

# Changes of lab session administration

---

# Changes of lab session administration

- Lab handout:
  - Merged to homework assignment.
  - Written mostly by instructors.
- Optional sections of lab handout:
  - Separated from the handout & homework assignment.
  - Will show up in the "Pages" section of Canvas.

# Recap

---

# Recap

- R environment
  - Associated with each R interpreter session
- R current working directory
  - Do **NOT** change during analysis
- Path
  - Absolute path
  - Relative path
  - Path delimiter:
    - \ in Windows
    - / in Linux and MacOS
    - / in R works for Windows, Linux, and MacOS

# Recap

- Tutorial of R data analysis
  - Input:
    - `read.csv()`
  - Analyze:
    - Plotting: `ggplot2`
    - Data manipulation: indexing, `subset()`, and `dplyr`
    - Statistical testing: `t.test()`
  - Output: `write.csv()` and `pdf()`

# Install packages

---



# Install packages

---

*# For normality test*

```
install.packages('nortest')
```

*# For power analysis of one sample proportion test*

```
install.packages('pwr')
```

*# You do not need to run this if you already*

*# installed `ggplot2`*

```
install.packages('ggplot2')
```

---

# The normal distribution

---

# The normal distribution

When a random variable  $X$  is distributed normally with mean  $\mu$  and variance  $\sigma^2$ , we write:

$$X \sim \mathcal{N}(\mu, \sigma^2)$$


Probability density distribution (PDF) of the normal distribution:

$$f(x|\mu, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$


Source: Normal Distribution at [en.wikipedia.org](https://en.wikipedia.org)

# PDF of the normal distribution

$$f(x|\mu, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$



# The normal distribution





- Single Peaked
- Symmetric
- $E[x] = \mu$
- $Var[x] = \sigma^2$
- $SD[x] = \sigma$

Source: Paul E. Johnson. CLT. 2011

# Generate normal random numbers with **rnorm**

```
opar <- par(no.readonly = TRUE)
par(mfrow=c(2, 2))
normdata <- rnorm(10, mean=0, sd=1)
hist(normdata)
hist(normdata)
par(opar)
```




Blank Graph 3


Blank Graph 4

```
par(mfrow=c(2, 2))
```

Histogram of normdata



Histogram of normdata




Blank Graph 3

Blank Graph 4

# par(opar)

```
# Plotting parameters have been set to old ones  
# in `opar`  
hist(normdata)
```






# Generate normal random numbers with **rnorm**

```
normdata <- rnorm(10, mean=0, sd=1)
mean(normdata)
## [1] -0.1397156
sd(normdata)
## [1] 0.5693181
```

# Generate 1 million normal random numbers

```
normdata <- rnorm(1000000, mean=0, sd=1)
mean(normdata)
## [1] -9.904492e-05
sd(normdata)
## [1] 1.000252
hist(normdata)
```




# Calculate probability density of normal distribution with `dnorm`

```
z_score_range <- seq(-4, 4, by = 0.2)
den_zscores <- dnorm(z_score_range)
plot(den_zscores, type = "l",
     main = "PDF on the Standard Normal Distribution",
     xlab = "Z-score", ylab = "Density", xaxt = "n")

den_zscore_sigmas <- c(dnorm(4), dnorm(3), dnorm(2),
                      dnorm(1), dnorm(0), dnorm(1),
                      dnorm(2), dnorm(3), dnorm(4))

den_score_labels <- c(-4, -3, -2, -1, 0, 1, 2, 3, 4)
axis(1, at = which(den_zscores %in% den_zscore_sigmas),
     labels = den_score_labels)
```

# Calculate probability density of normal distribution with `dnorm`




# Calculate cumulative probability of normal distribution with `pnorm`

```
pnorm_scores <- pnorm(z_score_range)
plot(pnorm_scores, type = "l",
     main = "CDF of the Normal Distribution",
     xlab = "quantile", ylab = "density", xaxt = "n")

pnorm_values <- c(pnorm(-4), pnorm(-3), pnorm(-2),
                  pnorm(-1), pnorm(0), pnorm(1),
                  pnorm(2), pnorm(3), pnorm(4))

axis(1, at = which(pnorm_scores %in% pnorm_values),
     labels = round(pnorm_values, 4), las = 3)
```

# Calculate probability density of normal distribution with `dnorm`



# Calculate quantiles of normal distribution with `qnorm`

```
qnorm(0.99)
## [1] 2.326348

qnorm(0.9999)
## [1] 3.719016

pnorm(qnorm(0.9999))
## [1] 0.9999

# lower.tail: if TRUE (default), probabilities are  $P[X \leq x]$ 
# otherwise,  $P[X > x]$ .
qnorm(0.9999, lower.tail = FALSE)
## [1] -3.719016

qnorm(1e-04)
## [1] -3.719016

pnorm(qnorm(0.9999, lower.tail = FALSE))
## [1] 1e-04
```

# Central limit theorem

## Theorem

Let  $X_1, X_2, \dots, X_n$  be a sequence of identically distributed (i.i.d.) random variables with mean  $E[X_i] = \mu$  and finite variance  $\text{Var}(X_i) = \sigma^2$ . Define  $S_n = \frac{1}{n} \sum_i X_i$ . Then, as  $n \rightarrow \infty$ ,  $S_n \xrightarrow{\mathcal{D}} \mathcal{N}(\mu, \sigma^2/n)$ .

Central limit theorem is difficult to prove algebraically, but it is quite easy to demonstrate with simulation.



# Demonstrate central limit theorem with simulation

## Theorem

Let  $X_1, X_2, \dots, X_n$  be a sequence of i.i.d. random variables with mean  $E[X_i] = \mu$  and finite variance  $\text{Var}(X_i) = \sigma^2$ .

Define  $S_n = \frac{1}{n} \sum_i X_i$ . Then, as  $n \rightarrow \infty$ ,  $S_n \xrightarrow{\mathcal{D}} \mathcal{N}(\mu, \sigma^2/n)$ .

- Simulation procedure:
  1. Let  $X_1, X_2, \dots, X_n$  be a sequence of i.i.d. uniformly distributed random variables.
  2. Calculate sample average  $S_n$ .
  3. Repeat  $m$  times.
  4. Plot the empirical PDF of  $S_n$  and normal PDF stated by the theorem.

# Demonstrate central limit theorem with simulation

```
# Let X1, X2, ..., X10 be a sequence of i.i.d.  
# random variables uniformly distributed  
# from 0 to 1.  
runif(n = 10, min = 0, max = 1)  
## [1] 0.89804708 0.85956656 0.63841285  
## [4] 0.59958342 0.68575572 0.28295208  
## [7] 0.42904749 0.15577247 0.26559273  
## [10] 0.01854827
```

# Demonstrate central limit theorem with simulation

```
number_of_samples <- 5000
size_of_each_sample <- 5000
unif_min <- 0
unif_max <- 100

unif_mean <- (unif_max - unif_min) / 2
unif_sd <- (((unif_max - unif_min) ^ 2) / 12) ^ 0.5


sample_average_vector <- replicate(number_of_samples, {
  mean(runif(n = size_of_each_sample,
            min = unif_min,
            max = unif_max))
})
```

# Demonstrate central limit theorem with simulation

```
clt_norm_mean <- unif_mean
clt_norm_sd <- unif_sd / (size_of_each_sample ^ 0.5)

ggplot(data = data.frame(x = sample_average_vector),
       mapping = aes(x = x)) +
  geom_histogram(mapping = aes(y = ..density..),
                alpha = 0.4) +
  geom_density(mapping = aes(color = 'Sample Ave. Dens.')) +
  stat_function(fun = dnorm,
               args = list(mean=clt_norm_mean,
                           sd=clt_norm_sd),
               aes(colour = 'CLT Norm. Dens.')) +
  labs(x = 'Sample Average', y = 'Density') +
  ggtitle("Central Limit Theorem Simulation")
```

# Demonstrate central limit theorem with simulation



# Check normality

---


# Methods to check normality

- Explorative data analysis (EDA)
  - Histogram
  - Density plot
  - Quantile-Quantile plot
- Statistical tests
  - Shapiro-Wilk normality test
  - Anderson-Darling normality test

# Use EDA to check normality

Histogram:

```
num_points <- 30  
normdata2 <- rnorm(num_points)  
  
qplot(normdata2) +  
  geom_histogram() +  
  theme_bw() +  
  ggtitle(paste("rnorm (", num_points, ")", sep=""))
```






# Use EDA to check normality

Quantile-Quantile (Q-Q) plot:


```
qqnorm(normdata2, pch=16, cex=0.5,  
       main=paste("Generated using rnorm(",  
                  num_points, ")", sep=""))  
qqline(normdata2, col=3)
```



# Q-Q plot

**Quantiles** divide probability distribution or sample observations into even intervals.

Example, 4-quantiles (quartiles) and 100-quantiles (percentiles):




# Q-Q plot

Q-Q plot compares the quantiles of one sample or distribution against the other.

```
quantile(0:10, probs=seq(0, 1, 0.1))  
##      0%    10%    20%    30%    40%    50%    60%    70%    80%    90%   100%  
##      0      1      2      3      4      5      6      7      8      9     10  
qqnorm(0:10, ylim = c(0,10))  
qqline(0:10, ylim = c(0,10))
```

Normal Q-Q Plot




# Histogram of TCGA-LUAD cigarettes per day

```
luad_cpd_sd <- sd(tcga_luad$cigarettes_per_day,
                  na.rm=TRUE)
luad_cpd_mean <- mean(tcga_luad$cigarettes_per_day,
                      na.rm=TRUE)


qplot(x = cigarettes_per_day, xlim=c(-5, 10),
      data = tcga_luad, geom = "blank") +
  geom_histogram(aes(y = ..density..),
                 alpha = 0.4) +
  geom_line(aes(y = ..density.., colour = 'Empirical'),
            stat = 'density') +
  stat_function(fun = dnorm,
                args = list(mean=luad_cpd_mean,
                             sd=luad_cpd_sd),
                aes(colour = 'Normal Approx')) +
  scale_colour_manual(name = 'Density',
                      values = c('red', 'blue')) +
  theme(legend.position = c(0.85, 0.85)) +
  ggtitle("TCGA LUAD: Cigarettes Per Day")
```

# Histogram of TCGA-LUAD cigarettes per day




# Q-Q plot of TCGA-LUAD cigarettes per day

```
qqnorm(tcga_luad$cigarettes_per_day,  
       pch=16, cex=0.5, main="TCGA-LUAD: Cigarettes per Day")  
qqline(tcga_luad$cigarettes_per_day, col=3)
```




# Histogram of TCGA-LUAD years smoked



# Q-Q plot of TCGA-LUAD years smoked

TCGA-LUAD: Years smoked





# Use statistical tests to check normality

Shapiro-Wilk normality test:

```
nortest_points <- 100
normdata_nortest <- rnorm(nortest_points)
shapiro.test(normdata_nortest)
##
##  Shapiro-Wilk normality test
##
## data:  normdata_nortest
## W = 0.97583, p-value = 0.06271
shapiro.test(1:1000)
##
##  Shapiro-Wilk normality test
##
## data:  1:1000
## W = 0.95481, p-value < 2.2e-16

# Smaller p-value indicates significant deviation from normal
# distribution.
```

# Use statistical tests to check normality

## Anderson-Darling normality test

```
ad.test(normdata_nortest)
##
##  Anderson-Darling normality test
##
## data:  normdata_nortest
## A = 0.38284, p-value = 0.3912
ad.test(1:1000)
##
##  Anderson-Darling normality test
##
## data:  1:1000
## A = 11.085, p-value < 2.2e-16

# Smaller p-value indicates significant deviation from normal
# distribution.
```

# Shapiro-Wilk test of TCGA-LUAD cigarettes per day

```
shapiro.test(tcga_luad$cigarettes_per_day)
```

```
##
```


```
##  Shapiro-Wilk normality test
```

```
##
```

```
## data:  tcga_luad$cigarettes_per_day
```


```
## W = 0.90998, p-value = 9.873e-14
```

TCGA-LUAD: Cigarettes per Day



# Shapiro-Wilk test of TCGA-LUAD years smoked

```
shapiro.test(tcga_luad$years_smoked)
##
##  Shapiro-Wilk normality test
##
## data:  tcga_luad$years_smoked
## W = 0.98572, p-value = 0.04686
```



# One-sample tests of proportion

---

# One-sample tests of proportion

- z-test
- $\chi^2$ -test using `prop.test`
- One-sided
- Two-sided
- Power analysis

# Overview of one-sample tests of proportion

- Proportion:

$$\frac{\text{The number of observations with certain condition}}{\text{The number of total observations}}$$

- Question: How likely the observed proportion ( $prop_{obs}$ ) is different from another proportion ( $prop_{H_0}$ )?
  - $prop_{obs}$  is determined by observations.
  - $prop_{H_0}$  is determined by us when specifying the null hypothesis ( $H_0$ ).

# z-test

$$z = \frac{\hat{p} - p_0}{\sqrt{\frac{p_0(1-p_0)}{n}}}$$

Intuition: apply central limit theorem on  $n$  random variables *i.i.d.* Bernoulli distribution.



## z-test

Example: A new study is performed where 22 out of 200 patients smoked more than 5 cigarettes per day. We would like to check if this group of patients has a statistically greater proportion of heavy smokers compared to the TCGA study. In TCGA, the proportion of heavy smokers 0.1.

Null hypothesis: The proportion of heavy smokers in this new study is no more than the proportion of heavy smokers in the TCGA study.

# z-test

Null hypothesis: The porportion of heavy smokers in this new study is no more than 0.1.

```
alpha <- 0.05
z0 <- qnorm(1 - alpha)
print(z0)
## [1] 1.644854

prop2 <- 22/200
p0 <- 0.1
n <- 200

z <- (prop2 - p0)/sqrt(p0 * (1 - p0)/n)
print(z)
## [1] 0.4714045

z >= z0
## [1] FALSE
```

# $\chi^2$ -test

Intuition: z-test with multiple categories. For each category, apply central limit theorem on  $n$  random variables *i.i.d.* Bernoulli distribution.

$$\chi^2 = \sum_{i=1}^k \frac{(x_i - np_i)^2}{np_i}$$

There are  $k$  total categories and  $n$  total observations.  $x_i$  is the number of observations of category  $i$ .  $p_i$  is the hypothesized probability of observing category  $i$ .

$\chi^2$  followed the  $\chi^2$  distribution with  $(k - 1)$  degrees of freedom.

# $\chi^2$ -test

Null hypothesis: The porportion of heavy smokers in the new study is no more than 0.1.

```
alpha = 0.05
z0 = qnorm(1 - alpha)
print(z0)
## [1] 1.644854
hs = 22
p0 = 0.1
n2 = 200

propresult <- prop.test(hs, n2, p = p0, correct = FALSE,
                        alternative = "greater")
```

# $\chi^2$ -test

```
print(propresult)
##
## 1-sample proportions test without continuity correction
##
## data:  hs out of n2, null probability p0
## X-squared = 0.22222, df = 1, p-value = 0.3187
## alternative hypothesis: true p is greater than 0.1
## 95 percent confidence interval:
##  0.0786844 1.0000000
## sample estimates:
##      p
## 0.11
```

# $\chi^2$ -test

```
# Comparing prop.test to z-score method.
```

```
Xstat <- propresult$statistic[[1]]
```

```
print(sqrt(Xstat))
```


```
## [1] 0.4714045
```

```
z = (hs/n2 - p0)/sqrt(p0 * (1 - p0)/n2)
```

```
print(abs(z))
```

```
## [1] 0.4714045
```

# One-sided versus two-sided test





A **p-value** (shaded green area) is the probability of an observed (or more extreme) result arising by chance


Source: [https://en.wikipedia.org/wiki/One-\\_and\\_two-tailed\\_tests](https://en.wikipedia.org/wiki/One-_and_two-tailed_tests)

# One-sided versus two-sided test

One-sided:



Two-sided:






# Two-sided one sample test of proportion

```
hs = 30 # second study patient proportion
p0 = 0.1 # TCGA LUAD proportion
n2 = 200 # second study sample size
z_twotail = (hs/n2 - p0)/sqrt(p0 * (1 - p0)/n)
pval_z_twotail = 2 * pnorm(z_twotail, lower.tail = FALSE)
pval_z_twotail
## [1] 0.01842213

proptest_twotail <- prop.test(hs, n2, p = p0, correct = FALSE,
                             alternative = "two.sided")

proptest_twotail$p.value
## [1] 0.01842213
```




# One-sided one sample test of proportion

```
hs = 22 # second study heavy smokers
p0 = 0.1 # TCGA LUAD proportion
n2 = 200 #Number of tests

propresult <- prop.test(hs, n2, p = p0, correct = FALSE,
                        alternative = "greater")

propresult$p.value
## [1] 0.3186759

z = (hs/n2 - p0)/sqrt(p0 * (1 - p0)/n2)
print(abs(z))
## [1] 0.4714045
pvalue_z_onesided <- pnorm(-abs(z))
pvalue_z_onesided
## [1] 0.3186759
```



# Power analysis for one-sample test of proportion

- The **power** of a binary hypothesis test is the probability that the test correctly rejects the null hypothesis ( $H_0$ ) when a specific alternative hypothesis ( $H_1$ ) is true.

$$\text{power} = \Pr(\text{reject } H_0 \mid H_1 \text{ is true})$$

- Main factors in calculating the power of a test:
  - Effect size, or the difference in means between two groups
  - Sample size:  $n$
  - Significance threshold,  $\alpha$  (often 0.05)

Source: [https://en.wikipedia.org/wiki/Statistical\\_power](https://en.wikipedia.org/wiki/Statistical_power)

# Power analysis for one-sample test of proportion

Calculate power given effect size, sample size, and significance threshold.

```
# h is effect size
pwr.p.test(h = 0.5, n = 55, sig.level = 0.05)
##
##      proportion power calculation for binomial distribution
##      (arcsine transformation)
##
##              h = 0.5
##              n = 55
##      sig.level = 0.05
##      power = 0.9597797
##      alternative = two.sided
```

# Power analysis for one-sample test of proportion

Calculate sample size given power, effect size, and significance threshold.

```
pwr.p.test(h = 0.5, n = NULL, sig.level = 0.05, power = 0.9)
##
##      proportion power calculation for binomial distribution
##      (arcsine transformation)
##
##              h = 0.5
##              n = 42.02968
##      sig.level = 0.05
##      power = 0.9
##      alternative = two.sided
```

# Power analysis for one-sample test of proportion

Calculate effect size given power, sample size, and significance threshold.

```
pwr.p.test(h = NULL, n = 100, sig.level = 0.05, power = 0.9)
##
##      proportion power calculation for binomial distribution
##      (arcsine transformation)
##
##              h = 0.3241514
##              n = 100
##      sig.level = 0.05
##      power    = 0.9
##      alternative = two.sided
```

# Power analysis for one-sample test of proportion

Estimate effect size `h` using `ES.h`.

```
pwr.p.test(h = ES.h(p1 = 0.75, p2 = 0.5), sig.level = 0.05,  
           n = NULL, power = 0.8,  
           alternative = "greater")  
  
##  
##      proportion power calculation for binomial distribution  
##      (arcsine transformation)  
##  
##              h = 0.5235988  
##              n = 22.55126  
##      sig.level = 0.05  
##      power = 0.8  
##      alternative = greater
```

Questions?



test

Just a test