

Visualizations for the Quantum Bouncer

Yen Lee Loh and Chee Kwan Gan, 2021-5-30

1 General-purpose setup

2 One-Sided Bouncer (1SB)

2.1 Eigenvalues and eigenfunctions

2.2 Classical paths (interactive)

2.3 Feynman path integral in semiclassical approximation: infographic

2.4 Wavepacket evolution (animation)

2.5 Propagator visualized as heatmap

2.6 Phase along path: detailed study

2.7 Phase diagram as function of (x_f, T) for fixed x_i : obtaining caustics as loci of foci

Plotting many paths by brute force reveals that the paths overlap strongly along caustic curves:

```

In[273]:= xi = 1; (* initial position *)
g = 1; (* gravitational field strength *)
viList = Range[-3, 3, .05]; (* list of initial velocities for plotting *)
Tmax = 12; (* maximum time for plotting *)
xmax = 2; (* maximum position for plotting *)
plotPathVi[xi_, vi_, T_, g_,
  style_ : Directive[AbsoluteThickness@0, Opacity[0.25, Black]] ] :=
Module[{vm, xt, tb, t0, amax, color},
  vm =  $\sqrt{v_i^2 + 2 g x_i}$ ; t0 =  $\frac{v_i}{g}$ ; tb =  $\frac{2 v_m}{g}$ ; (* vm and tb are lists *)


  xt = Function[{t},  $\frac{v_m^2}{2 g} - \frac{g}{2} \left( t - t_0 - t_b \text{Floor}\left[\frac{t - t_0}{t_b} + \frac{1}{2}\right] \right)^2$ ];

  Plot[Evaluate[xt[t]], {t, 0, T}, PlotStyle → style, Exclusions → None];

gr = Show[
  Table[ plotPathVi[xi, vi, Tmax, g,
    Directive[AbsoluteThickness@0.1, Black]] , {vi, viList}],
  Frame → True, FrameLabel → {"Time t", "Position x"}, RotateLabel → True,
  PlotRange → {{0, Tmax}, {0, xmax}}, PlotRangePadding → 0,
  ImageSize → {648, 216}, AspectRatio → Full,
  ImagePadding → {{Automatic, Automatic}, {Automatic, 8}}]

```

Out[275]=



From our analysis of the symmetric bouncer, we find that for a path with initial position x_i , initial velocity v_i , and k bounces, there is a focus at time

$$t_k^F = \frac{2k}{g} \frac{v_i^2 + 2kv_i v_m + v_m^2}{2kv_i + v_m} = \frac{2k}{g} \frac{2v_i^2 + 2gx_i + 2kv_i \sqrt{v_i^2 + 2gx_i}}{2kv_i + \sqrt{v_i^2 + 2gx_i}}.$$

The position at this time is

$$x_k^F = X(t_k^F) \quad \text{where } X(t) = x_i + \frac{v_i^2}{2g} - \frac{g}{2} \left(t - \frac{v_i}{g} - \frac{2\sqrt{v_i^2 + 2gx_i}}{g} k \right)^2.$$

If we loop over each value of $k = 1, 2, 3, \dots$, loop over many values of v_i , and make a Parametric-Plot of the locus of (t_k^F, x_k^F) , we obtain the caustics. The caustics are the loci of the foci.


```

In[587]:= Clear[n, vi, xi, g];
xi = 1; g = 1;
ParametricPlot[
  Table[{
    
$$\frac{2n}{g} * \frac{2vi^2 + 2gxi + 2nvi \sqrt{vi^2 + 2gxi}}{\sqrt{vi^2 + 2gxi} + 2nvi},$$

    
$$xi + \frac{vi^2}{2g} - \frac{g}{2} \left( \frac{2n}{g} * \frac{2vi^2 + 2gxi + 2nvi \sqrt{vi^2 + 2gxi}}{\sqrt{vi^2 + 2gxi} + 2nvi} - \frac{vi}{g} - \frac{2 \sqrt{vi^2 + 2gxi}}{g} n \right)^2$$

  }, {n, 1, 6}]
  , {vi, -5, 5}, PlotRange -> {{0, 15}, {0, 2}},
  ImageSize -> {648, 216}, AspectRatio -> Full,
  FrameLabel -> {"T", "xf"}]

```



The pattern of caustics above is universal. Changing x_f merely results in rescaling.

2.8 Phase diagram as function of (x_f, T) for fixed x_i : discriminant method

2.9 Phase diagram as function of (x_i, x_f) for fixed T – a bit old

2.10 Van Vleck determinant D_α – details of derivation

3 Symmetric Bouncer

3.1 Eigenvalues and eigenfunctions

3.2 Feynman path integral in semiclassical approximation: infographic

3.3 Phase diagram as function of (x_f, T) for fixed x_i : discriminant method

3.4 Path divergence function $f_\alpha(t)$ – slightly messy

3.5 Focal times t_k^F – slightly messy

3.6 Morse index m as a function of (T, v_i) – slightly messy

3.7 Propagator visualized as heatmap

3.8 Wavepacket evolution (animation)

3.9 Phase diagram as function of (x_f, T) for fixed x_i : obtaining caustics as loci of foci

The expression for the k th focal time is surprisingly messy. (See section on focal times.) Careful consideration allows us to plot the caustics using ParametricPlot:

```

In[500]:= Clear[n, vi, xi, g];
xi = 1; g = 1;
gr2 = Show[{
  (*----- Plot one branch -----*)
  Table[
    ParametricPlot[{
      
$$\frac{2n}{g} * \frac{2vi^2 + 2gxi + 2nvi \sqrt{vi^2 + 2gxi}}{\sqrt{vi^2 + 2gxi} + 2nvi},$$


      
$$\left( xi + \frac{vi^2}{2g} - \frac{g}{2} \left( \frac{2n}{g} * \frac{2vi^2 + 2gxi + 2nvi \sqrt{vi^2 + 2gxi}}{\sqrt{vi^2 + 2gxi} + 2nvi} - \frac{vi}{g} - \frac{2 \sqrt{vi^2 + 2gxi}}{g} n \right)^2 \right) (-1)^n$$

    ], {vi, - $\frac{1}{\sqrt{2} \sqrt{n^2 + n}}$ , 2}]
    , {n, 1, 6}],
  (*----- Plot the other branch -----*)
  Table[
    ParametricPlot[{
      
$$\frac{2n}{g} * \frac{2vi^2 + 2gxi + 2nvi \sqrt{vi^2 + 2gxi}}{\sqrt{vi^2 + 2gxi} + 2nvi},$$

      
$$\left( xi + \frac{vi^2}{2g} - \frac{g}{2} \left( \frac{2n}{g} * \frac{2vi^2 + 2gxi + 2nvi \sqrt{vi^2 + 2gxi}}{\sqrt{vi^2 + 2gxi} + 2nvi} - \frac{vi}{g} - \frac{2 \sqrt{vi^2 + 2gxi}}{g} n \right)^2 \right) (-1)^n$$

    ], {vi, -2, - $\frac{1}{\sqrt{2} \sqrt{n^2 - n}}$ }]
    , {n, 2, 6}]
  },
  PlotRange -> {{0, 15}, {-2, 2}}, PlotRangePadding -> 0,
  ImageSize -> {648, 126 * 2}, AspectRatio -> Full, Axes -> False,
  FrameLabel -> {"T", "x_f"}]

```




Comparison of phase diagrams of one-sided bouncer and symmetric bouncer

```
In[579]:= Clear[n, vi, xi, g];
xi = 1; g = 1;
gr1 = Show[
  Table[
    ParametricPlot[ {
      
$$\frac{2n}{g} * \frac{2vi^2 + 2gxi + 2nvi \sqrt{vi^2 + 2gxi}}{\sqrt{vi^2 + 2gxi} + 2nvi},$$

      
$$xi + \frac{vi^2}{2g} - \frac{g}{2} \left( \frac{2n}{g} * \frac{2vi^2 + 2gxi + 2nvi \sqrt{vi^2 + 2gxi}}{\sqrt{vi^2 + 2gxi} + 2nvi} - \frac{vi}{g} - \frac{2\sqrt{vi^2 + 2gxi}}{g} n \right)^2$$

    }, {vi, - $\frac{1}{\sqrt{2} \sqrt{n^2 + n}}$ , 2}, PlotStyle ->
      {{Dashed, CapForm@"Butt", AbsoluteThickness@3, ColorData[97][2]}}]
    , {n, 1, 6}],
  ImageSize -> {648, 162}, AspectRatio -> Full,
  PlotRange -> {{0, 15}, {0, 2}}, FrameLabel -> {"T", "x_f"}];
```

```
In[586]:= Show[gr2, gr1, PlotRange -> {{0, 15}, {-2, 2}}, ImageSize -> {648, 162 * 2},
Axes -> True, AxesOrigin -> {0, 0}, AxesStyle -> LightGray]
```



In the figure above, the dashed orange curves are the caustics (phase boundaries) of the one-sided bouncer. The solid blue curves are the caustics of the two-sided bouncer. We see that these boundaries are related to each other by reflection.