



Associate editor: H. Bonisch

Novel therapeutics for type 2 diabetes: Incretin hormone mimetics (glucagon-like peptide-1 receptor agonists) and dipeptidyl peptidase-4 inhibitors

E.J. Verspohl

Department of Pharmacology, Institute of Medicinal Chemistry, University of Münster, Germany

ARTICLE INFO

Keywords:

Diabetes

GLP-1

Incretin mimetics

DPP-4

Incretin enhancers

ABSTRACT

Known treatments of type 2 diabetes mellitus have limitations such as weight gain, and hypoglycaemias. A new perspective is the use of incretin hormones and incretin enhancers. Incretins are defined as being responsible for the higher insulin release after an oral glucose load compared to an intravenous glucose load. The delicate balance of glucose homeostasis, in which incretin hormones are involved, is disturbed in type 2 diabetes mellitus. The incretin GLP-1 helps to maintain glucose homeostasis through stimulation of insulin secretion and inhibition of glucagon release in a glucose-dependent manner. This is associated with reductions in body weight, and no risk of hypoglycaemias. When classical oral agents have failed to maintain adequate glycaemic control, incretin mimetics may be of particular value for obese patients and those who have little control over meal sizes. Exenatide was marketed as a GLP-1 analogue and longer acting incretin mimetics such as liraglutide, albiglutide and others have the same pharmacological profile.

In addition to incretin mimetics incretin enhancers which inhibit/delay degradation of incretins were developed: so-called DPP-4 inhibitors such as sitagliptin and vildagliptin are approved in Europe. Their differences from incretin mimetics include: oral bioavailability, less side effects with overdose, no direct CNS effects (nausea and vomiting) and no effect on weight. In rodent models of diabetes, but not yet in humans, GLP-1 receptor agonists and DPP-4 inhibitors increase islet mass and preserve β -cell function.

Incretin mimetics and enhancers expand type 2 diabetes treatment, are still not first line therapy and it is discussed if they are to be prophylactically used.

© 2009 Elsevier Inc. All rights reserved.

Contents

1. History	113
2. The incretin hormones (GLP-1 and GIP)	114
3. Physiology of GLP-1 and GIP (release kinetics, effects, similarities and dissimilarities, kinetics including degradation, receptors and 2nd messengers involved)	115
4. Pathophysiological impact of the incretin concept (similarities and differences between the incretins GLP-1 and GIP in pathophysiological terms)	119
5. GLP-1 receptor agonists (exenatide)	120
6. Longer acting GLP-1 analogues	123
7. DPP-4 (dipeptidyl-4) inhibition as a drug target.	125
References	131

Abbreviations: GLP-1, glucagon-like peptide 1; GIP, glucose-dependent insulinotropic peptide (formerly gastrin inhibiting peptide); DPP-4, (dipeptidyl peptidase 4); WNT, cell-secreted glycoprotein ligand; PYY, peptide YY; s.c., subcutaneous; min, minute; TNF- α , tumor necrosis factor alpha; PAI-1, plasminogen activator inhibitor-1; VCAM-1, vascular cell adhesion molecule; ICAM-1, intercellular adhesion molecule; DAPD, dual-acting peptide for diabetes; PACAP, pituitary adenyl cyclase-activating peptide; GH-RF, growth hormone releasing factor, somatoliberin; SP, substance P; RANTES, regulated on activation normal T cells expressed and secreted; GRH, growth hormone-releasing hormone; IGF-1, insulin-like growth factor 1; HCG, human chorionic gonadotropin.

E-mail address: verspohl@uni-muenster.de.

1. History

By 2030, the World Health Organization predicts more than 300 million people to be diagnosed with type 2 diabetes (Zimmet et al., 2001). Current treatment modalities of those patients include exercise, diet and a variety of therapeutics. B-cell dysfunction in type 2 diabetes is characterized by reduced B-cell sensitivity to glucose and a delay and—in addition (see below)—a reduction in the meal-induced insulin secretion, a loss of regular oscillatory insulin




Fig. 1. Incretin effect (schematically shown differences in curves after i.v. and per os application of glucose).

secretion and of first phase insulin response to a glucose challenge, by an elevated proinsulin:insulin ratio, and by an abnormal secretion of islet amyloid polypeptide.

Known treatments have limitations and side effects, e.g. weight gains. The main classes of antidiabetic drugs either suppress hepatic glucose production (biguanides), stimulate insulin secretion (sulfonylureas and glinides), delay digestion and absorption of intestinal carbohydrates (α -glucosidase inhibitors) or improve insulin sensitivity and peripheral glucose uptake (thiazolidinediones and metformin) or insulin (Krentz & Bailey, 2005). Major pathophysiological issues are not addressed by current therapies (Baron et al., 1987; NN, 1995, 1998; Kahn et al., 2006) especially the progressive B-cell failure in type 2 diabetes. Since obesity is the engine that is driving the epidemic of diabetes, it is disappointing that most antidiabetic treatments are associated with weight gain.

New innovative concepts of treatment had to be launched: one new possibility is based on incretins. The incretin concept (the name did not exist yet) was indirectly initiated 100 years ago by showing that

duodenal mucosa extracts are effective in inhibiting glycosuria (Moore et al., 1906). An incretin is a compound which is responsible for the higher insulin release in response to an oral glucose load compared to an equal intravenous glucose load (reaching the same glucose level) as shown in Fig. 1. The term incretin, first used by La Barre in 1932, refers to gut derived hormones that stimulate insulin secretion with nutrient ingestion (Elrick et al., 1964; McIntyre et al., 1964; Perley & Kipnis, 1967; Creutzfeldt, 1979; Nauck et al., 1986; Creutzfeldt, 2005; Leon et al., 2006). E.g. glucose was given intravenously and compared with an intrajeunal application (McIntyre et al., 1965). The incretin effect is the most important component of endocrine signalling from the gut affecting pancreatic islet physiology and function.

The incretin effect is believed to be mediated by mainly two incretin hormones: glucose dependent insulinotropic polypeptide (GIP, originally referred to as gastric inhibitory peptide) and GLP-1 (glucagon-like peptide-1) (Holst, 2004). Both hormones enhance insulin secretion to an extent that can fully account for the incretin effect (Kreymann et al., 1987; Nauck et al., 1989; Nauck et al., 1993). It is difficult to state which of both incretins is more important: in normal healthy subjects GLP-1 is several times more potent than GIP on a molar basis at equivalent glycemic conditions (Elahi et al., 1994). Nevertheless GIP may be more effective than GLP-1; this can be indirectly derived from the fact that postprandial concentrations of GIP reach 250–500 pmol/l which is much higher than the minimum concentration affecting insulin release compared to GLP-1 for which postprandial concentrations of 30–50 pmol/l hardly overcome the minimum concentration for insulin release (reviewed by Creutzfeldt and Nauck, 1992). The fact that the physiologic increase after an oral glucose load is much more pronounced for GIP than for GLP-1 was described (Nauck et al., 1993).

More than 40 years ago an enzyme was purified not knowing that it would later be called DPP-4 (DPP-IV, dipeptidyl-peptidase IV; EC 3.4.14.5) (Hopsu-Havu & Glenner, 1966). The enzyme cleaves N-terminal dipeptides with a proline or alanine residue (Fig. 2), an area which is safe with respect to degradation by nonspecific proteases (Yaron & Naider, 1993). DPP-4 cleaves both the incretins GLP-1 and GIP which are substrates (Mentlein et al., 1993) (Fig. 2).

2. The incretin hormones (GLP-1 and GIP)

Eating provokes the secretion of multiple gastrointestinal hormones which have pleiotropic effects such as the regulation of gut motility, secretion of gastric acid and of pancreatic enzymes, induction of gall bladder contraction, and effects on nutrient absorption. Some



Fig. 2. Amino acid sequence of GLP-1 (two types), GIP, exenatide and liraglutide (deviations from GLP-1(7-36)amide are marked by underlining).

gut hormones also facilitate the disposal of absorbed glucose through the stimulation of insulin secretion from the endocrine pancreas: two out of them are incretins, GLP-1 and GIP.

The first incretin to be identified, glucose-dependent insulinotropic polypeptide (GIP), was purified from porcine intestinal extracts; it has weak effects on gastric acid secretion but potent insulinotropic actions (Dupré et al., 1973). The GIP gene is expressed mainly in K cells, enterochromaffin cells of the proximal small intestine (enteroendocrine duodenal and jejunal mucosa). GIP, a 42-amino acid hormone, is stimulated by enteral glucose, lipids and products of meal digestion in a concentration dependent manner (Schirra et al., 1996). Elevated GIP plasma levels augment glucose-stimulated insulin secretion (Nauck et al., 1989). Consistent with the incretin concept, GIP acts as a feed-forward mechanism to signal the endocrine pancreas of impending substrate fluxes from the gut (Ebert & Creutzfeldt, 1987).

The second incretin hormone, called glucagon-like peptide-1 (GLP-1), was identified after the cloning of the cDNAs and genes encoding the human proglucagon (Fig. 3). The human proglucagon gene located on the long arm of chromosome 2 consists of six exons and five introns (Hansotia & Drucker, 2005). The proglucagon gene is expressed in a type of enterochromaffin cells, the L-cells of the intestine ileum and colon. Processing of the expressed peptide by proconvertase 1/3 leads to GLP-1 (Fehmann et al., 1995); another processing of the proglucagon gene product by proconvertase 2 in the A-cells of pancreatic islets leads to glucagon (Fig. 3).

GLP-1(1–37) is produced as an inactive 37-amino acid peptide (Fig. 3). The active form is produced by post-translational cleavage of six amino acids from the N-terminal end. GLP-1 exists in two circulating forms, GLP-1(7–37) and GLP-1(7–36) amide (amidation at the glycine end of the C-terminal) (Fig. 2), from which GLP-1(7–36) amide is more abundant in the circulation after being stimulated by eating (Kieffer & Habener, 1999). Both GLP-1s are equipotent with respect to their receptor interaction as well as their insulinotropic effect.

Cholecystokinin is important for the incretin that did not get clinical importance because its incretin effect is mainly observed in animals (Verspohl et al., 1986a, 1986b; Rushakoff et al., 1987; Rushakoff et al., 1993; Liddle, 1995).

3. Physiology of GLP-1 and GIP (release kinetics, effects, similarities and dissimilarities, kinetics including degradation, receptors and 2nd messengers involved)

3.1. Regulation of incretin secretion (stimuli for incretin release)

Ingestion of nutrients such as carbohydrate, protein and fat leads to release of GLP-1 as well as GIP; of these, carbohydrate ingestion is the best stimulus for GLP-1 secretion. The very rapid increase in plasma levels of GLP-1 after eating implies a combination of endocrine and neural signals well before digested food transits through the gut is able to directly activate the release from L cells; stimulation of GLP-1 secretion may involve vagal pathways (Rocca & Brubaker, 1999). The Rho GTPase Cdc42 is involved in the secretion process of GLP-1 (Lim & Brubaker, 2008).

A rapid gastric emptying, e.g. evidenced by post-gastrectomy dumping syndromes, causes marked increase in GIP and GLP-1 secretion (Ranganath et al., 1998). The impact of circulating free fatty acids on GLP-1 secretion is a matter of controversy: while an inhibition was observed (Ranganath et al., 1999b, 1999c) the opposite was found when the long-chain fatty acid-derivate oleoylethanolamide was investigated which interacts with novel fatty acid receptor (GPR 119) (Lauffer et al., 2009). The effects of α -glycosidase inhibitors are controversial: they increase indirectly GLP-1 secretion (Göke et al., 1994; Qualmann et al., 1995; Seifarth et al., 1998; Ranganath & Morgan, 2000); this was also shown for voglibose (Göke et al. 1995a)




Fig. 3. Proglucagon gene and processing of glucagon and GLP-1.

and miglitol (Lee et al., 2002). However, others did not confirm this effect of acarbose (Hücking et al., 2005). GIP may be reduced by acarbose (Groop et al., 1986).

Plasma levels of GLP-1 are low in the fasted state (5–10 pmol/l), and increase rapidly 3–5 times after eating. Circulating GIP and GLP-1 concentrations rise within 15 min and peak concentrations of GIP and GLP-1 of approx. 200 and 50 pmol/l, respectively, are obtained after 30–45 min, returning to basal values after 2–3 h (Vilsbøll et al., 2001). The concentration of intact GLP-1 after a meal rises proportionally with the size of the meal (Vilsbøll et al., 2003b). Since circulating GLP-1 levels are low, it cannot be excluded that GLP-1 receptors in the portal vein (reached directly after secretion of GLP-1) mediate some effects as alternative to the endocrine signaling (insulin release) (Perabo et al., 2008).

Interestingly insulin and exendin-4 (a GLP-1 analogue, see below) appear to stimulate the proliferation of GLP-1 secreting cells (Miller et al., 2008).

3.2. Effects in vitro (pancreatic islets)

GLP-1 increases insulin secretion and the biosynthesis of important B-cell products besides insulin: glucokinase and GLUT 2 glucose transporters (Wang et al. 1997).

GLP-1 inhibits the secretion of glucagon in a glucose-dependent way (Ørskov et al., 1988; Schirra et al., 2006), but the mechanism is not fully understood and may be a consequence of several factors: since a controversy exists whether receptors for GLP-1 on A-cells exist, intra-islet insulin released from B-cells in response to GLP-1 may locally inhibit glucagon secretion. The resulting improvement in insulin/glucagon ratio may account at least partly for the reduction of insulin resistance in both liver and peripheral tissues such as the skeletal muscle (Ritzel et al., 1995). Surprisingly, GLP-1 does not inhibit, but stimulates glucagon secretion when isolated rat pancreatic A-cells are investigated (Ding et al., 1997), making a direct inhibitory effect on the A-cells unlikely. An important notion is that in vivo, the inhibitory effect of GLP-1 on glucagon secretion fades away at glucose levels just below fasting levels, so that the normal counter-regulatory glucagon secretion at hypoglycaemic levels is preserved even at high GLP-1 concentration (Nauck et al., 2002). Administration of GLP-1 is thus unlikely to be associated with an increased risk of hypoglycaemia (see below).

GLP-1 has been shown to stimulate somatostatin secretion (Ørskov et al., 1988), which in turn may inhibit insulin and glucagon secretion mediated by direct receptor activation of A-cells (de Heer et al., 2005). This indicates a complex interplay between GLP-1 and all the major endocrine cells of the islets possibly being important for the fine-tuning of homeostatic control of glucose metabolism.

3.3. Effects on non-endocrine cells (stomach, brain, liver) (Fig. 4)

GLP-1 slows gastric emptying (Wettergren et al., 1993; Nauck et al., 1997; Naslund et al., 1999b) using cholinergic pathways (Schirra et al., 2009), and induces gastric relaxation via nitric oxide (Andrews et al., 2007) and satiety (Flint et al., 1998). By delaying gastric emptying postprandial glycaemia is minimized; also the distension of the stomach and peripheral satiety signals are influenced.

The effects of GLP-1 and GIP on gastric acid secretion *in vivo* at physiological concentrations are disputed. Receptors for GIP and GLP-1 are present in the stomach. The mechanisms involved are not clear, but endocrine or neural mechanisms, or a combination, may be involved. Experiments with vagal deafferentiation of rats and humans show the involvement of vagus (Wettergren et al., 1994; Imeryuz et al., 1997; Wettergren et al., 1997b). Peptide YY (PYY) may participate in inhibitory effects on gastric acid release which is released in parallel with GLP-1 from the L cells (Wettergren et al., 1997a). Both hormones show an additive effect and are able to block, for example, gastrin-stimulated gastric acid secretion (Holst, 1997).

Effects of incretins (and incretin mimetics) on body weight may be mediated through effects on the adipocytes and brain centres controlling food intake and energy expenditure (Kastin et al., 2002). Administration of GLP-1 into cerebral ventricles of rats sharply decreased their energy intake (Turton et al., 1996). GLP-1 effects may be mediated by interactions of central GLP-1 (released from noncatecholaminergic neurons in the solitary tract nucleus) (Larsen et al., 1997) with hypothalamic and extrahypothalamic nuclei in the brain, as well as via peripheral GLP-1, which can reach the area postrema and subfornical organs with access to hypothalamic centres controlling energy intake (Ørskov et al., 1996). Thus, although GLP-1 synthesis was demonstrated in the brain, peripheral GLP-1 may bind to regions of the brain.

GIP receptor expression has also been found in the brain; GIP administration in healthy subjects, however, appears to increase food intake and to decrease energy expenditure, in contrast to GLP-1 (Lui et al., 1990; Ranganath et al., 2007). Beyond acting as an incretin (endocrine axis) other prominent actions are obvious: GIP seems to increase glucose uptake and triglyceride storage in adipocytes (Miyawaki et al., 2002; Gault et al., 2005), and to stimulate possibly bone mass and bone turnover (Tsukiyama et al., 2006; Xie et al., 2007).

In Table 1 the physiologic characteristics of GIP and GLP-1 are summarized and compared. These incretins share a number of similarities: both peptides are released in response to meal consumption with a dose dependent relationship to meal size. Both are inactivated

by DPP-4. Early intracellular signaling events following ligand binding to different receptors are comparable. They cause enhanced insulin secretion from B-cells in a glucose-dependent manner. The general conformities shared by GIP and GLP-1 suggest the evolution of redundant systems to mediate the incretin effect. In fact, there is evidence that one system compensates in the absence of normal functioning of the other incretin (Pamir et al., 2003). GIP and GLP-1 have physiologic roles that are complementary and in some aspect distinct.

Despite the many similarities between the incretins there are several dissimilarities. Key points of distinction are site of synthesis and the manner in which nutrient stimuli is coupled to incretin secretion, mechanism of action, and extra-incretin effects. GIP and GLP-1 have only modest degrees of sequence homology as do their receptors. GIP is produced primarily in the upper intestine and there is convincing data connecting K-cell secretion of GIP with the digestion and absorption of lipid and carbohydrate (Falko et al., 1975; Ross & Dupré, 1978; Schirra et al., 1996).

GLP-1 is produced throughout the small intestine but the highest concentrations of producing L-cells are in the distal gut (Eissele et al., 1992). The GLP-1 receptor is expressed in cells of the endocrine pancreas and in many other tissues, mainly those of the central and peripheral nervous systems, heart, kidney, lung, and gastrointestinal tract and GLP-1, therefore, mediates a broader spectrum of effects than GIP. This is why GLP-1 also inhibits gastric emptying, food ingestion, and promotes enhanced glucose disposal through neural mechanisms (Burcelin et al., 2001) which all contribute to glucoregulation. A direct effect on muscle and adipose tissue (Fig. 4) as originally propagated is questionable.

3.4. Receptors and signalling

The actions of both GIP and GLP-1 are mediated by specific and structurally distinct G-protein-coupled receptors (Amiranoff et al., 1984; Wheeler et al., 1995; Thorens & Widmann, 1996). GIP interacts with two subtypes of receptors, GIP receptor 1 and 2, which are predominantly expressed on islet A and B cells, and to a lesser extent in adipose tissue, the upper gastrointestinal tract, adrenal cortex, bone, pituitary and a variety of brain regions (Bollag et al., 2000; Drucker, 2006). The GIP receptor has substantial homology with the secretin-VIP receptor family.

The GLP-1 receptor is expressed in pancreatic A-cells (controversial, see above), B-cells and D-cells. The inhibition of glucagon release by GLP-1 may be a direct effect or an indirect one via somatostatin release (Sinclair and Drucker, 2005b; Leon et al., 2006) (as discussed

Table 1
Summary of the physiological characteristics of glucose-dependent insulinotropic polypeptide (GIP) and glucagon-like peptide-1 (GLP-1): Similarities and differences (modified from Ranganath, 2008; Green and Flatt, 2007).

	GIP	GLP-1
Number of amino acids	42	30/31
Cell of origin in the intestine	K cells/proximal	L-cells/distal intestine/colon
Degradation by	DPP-4	DPP-4
Insulin secretion (glucose-dependent)	Increased	Increased
B-cell glucose responsiveness	Yes	Yes
Lowering of blood glucose	Yes	Yes
Extrapankreatic glucose lowering effect	Yes	Yes
Inhibition of hepatic insulin extraction	Yes	Yes
Gastric emptying	Inconclusive, eventually no effect	Slowed
B-cell proliferation	Stimulated	Stimulated
Glucagon secretion	Unaffected or increased (but not suppressed)	Suppressed
Somatostatin secretion	Unchanged	Increased
Food intake	Increased	Decreased
Reduce body weight	No	Yes
Gastric acid secretion	Decreased	Decreased
Insulin sensitivity	Unchanged	Probably improved
Secretion in Type 2 diabetes	Reduced (questionable)	Reduced (questionable)
Insulin secretion to administration in type 2 diabetes	Impaired	Relatively well preserved




Fig. 4. Physiology of GLP-1 secretion and action on various tissues (modified from Kieffer and Habener, 1999; Meier et al., 2002; Ranganath, 2008). The pleiotropic effects of GLP-1 include the brain (satiety), peripheral organs (anabolic effects), endocrine pancreas (increase of insulin and somatostatin release and inhibition of glucagon release). An effect of GIP on GLP-1 release is not obvious in humans.

above). GLP-1 receptors are also present on mucosal cells in the gastric and small intestinal mucosa, on cardiac myocytes, neurons in the hypothalamus, hindbrain, the vagus nerve and several other brain regions.

Activation of GIP and GLP-1 receptors on B-cells leads to rapid increases in levels of cAMP (Fig. 5) which is glucose-dependent

(Drucker et al., 1987) and which is linked to both acute and long-term effects. Acutely a subsequent activation of either protein kinase A (PKA) (Wang et al. 2001) or cAMP-regulated guanine nucleotide exchange factor II (Epac 2) (Holz et al., 2006) alters ion channel activity, handling of intracellular calcium and enhances exocytosis of insulin containing granules (Holz, 2004). The glucose dependency of




Fig. 5. Mechanisms of action of GLP-1.

insulin release may depend on the K_{ATP} channels of the B-cell which close when activated by a rise in the intracellular ATP concentration, hence making them sensitive to the intracellular glucose metabolism and therefore the plasma glucose concentration. The effect of GLP-1 appears to involve further channel closure and membrane depolarization (through accumulation of cAMP and activation of protein kinase A), but apparently, a parallel glucose mediated closure is required for the effect of GLP-1.

GIP receptor signaling in the B-cells shares many similarities with GLP-1 (production of cAMP and activation of PKA, increased intracellular Ca²⁺, and closure of K_{ATP} channels, causing insulin exocytosis). Like GLP-1, a significant component of GIP signaling is independent of PKA and acts through Epac 2 pathway (Kashima et al., 2001).

3.5. Kinetics

After being secreted both GIP and GLP-1 have a very short circulating half-life of 3–5 min due to the action of proteases such as dipeptidyl peptidase IV (DPP-4) and other endopeptidases, resulting in inactivation (Ørskov et al., 1993). An important organ for GLP-1 clearance is the kidney (Deacon, 2005). Both GIP and GLP-1 contain alanine at position 2 (Fig. 2), where DPP-4 is active (see above and later). The degradation of GLP-1 and GIP occurs through N-terminal removal of the two amino acids histidine and alanine (GLP-1) and tyrosine and alanine (GIP) (Fig. 2), yielding the inactive (with regards to stimulating insulin secretion) truncated form of e.g. GLP-1 (9–36) amide (Mentlein, 1999). As a result of DPP-4 activity, intact, biologically active GLP-1 represents only 10–20% of total plasma GLP-1 (Deacon et al., 1995b). Truncation of GLP-1(7–36)amide to GLP-1(9–36) amide reduces receptor affinity by 1000 and rather completely eliminates its insulin-releasing activity (Knudsen & Pridal, 1996; Deacon et al., 2002; Green et al., 2004). The major significance of DPP-4 for incretin inactivation was shown in DPP-4 gene knock out mice which have elevated levels of plasma GIP and GLP-1, increased insulin secretion, and reduced glucose increases after glycaemic challenge (Marguet et al., 2000). It is not clear to which degree additional proteases, such as a human neutral endopeptidase, are also essential for GLP-1 inactivation. Neutral endopeptidase 24.11 (NEP24.11, also known as neprilysin) is a widespread membrane-bound zinc metallopeptidase (Plamboeck et al., 2005); the molecule is cleaved at the C-terminal region.

The quick degradation by DPP-4 and the short half-life raises the question of whether significant amounts of intact, bioactive, peptide really reach the arterial circulation and target tissues (Hansen et al., 1999) (see above: receptors in the portal vein). In general some uncertainty exists (with this respect) over how GLP-1 mediates its actions at this low concentration.

3.6. Clinical aspects

A short-term intravenous infusion of GLP-1 (1–1.2 pmol kg⁻¹ min⁻¹) leading to plasma concentrations of 70–150 pmol/l (total GLP-1) or of 10–20 pmol/l (intact biologically active GLP-1), lowers blood glucose in patients with type 2 diabetes through a transient stimulation of insulin secretion and suppression of glucagon secretion and gastric emptying (Nauck et al., 1993b; Willms et al., 1996; Toft-Nielsen et al., 2001b). A long-term 6-week s.c. infusion of GLP-1 in patients with type 2 diabetes, achieving plasma levels of 60–70 pmol/L, led to substantial improvements in insulin secretory capacity, insulin sensitivity, a reduction in HbA_{1c} by 1.2% and a modest weight loss (1.9 kg) (Zander et al., 2002).

GLP-1 suppresses hepatic glucose production, and promotes insulin independent glucose disappearance (D'Alessio et al., 1994; Egan et al., 2002b). The ability of GLP-1 to lower plasma glucose is preserved even in type 1 diabetic patients (no residual B-cell function)

(Creutzfeldt et al., 1996), indicating that a paracrine effect of insulin or other B-cell products is not involved.

The impact of endogenous GIP and GLP-1 on glucose homeostasis has been investigated indirectly both by using receptor antagonists and by looking at mice with receptor defects. The GLP-1 receptor antagonist exendin (9–39) provokes a defective glucose-stimulated insulin secretion, reduces glucose clearance, increases levels of glucagon and induces a quicker gastric emptying (Schirra et al., 1998). Mutated mice with abnormal GIP or GLP-1 receptors have impaired glucose-stimulated insulin secretion and glucose tolerance and become diabetic (Scrocchi et al., 1996; Miyawaki et al., 1999).

3.6.1. Effects on glucose homeostasis and HbA_{1c}

Many of the already mentioned effects of GLP-1 are important for glucose homeostasis during therapy: stimulation of insulin, restoration of first phase of insulin release, suppression of glucagon secretion, inhibition of gastric emptying, and reduction of appetite and food intake. When glucose levels reach normal fasting levels, insulin and glucagon levels return to baseline concentrations despite continued GLP-1 infusion, indicating the glucose-dependent nature of the stimulation of insulin and suppression of glucagon by GLP-1 (Nauck et al., 1993b). The fact that the effects are glucose dependent is a major advantage over sulfonylureas. The question in general is whether therapeutic approaches should be targeted towards relieving the demand on the B-cell to secrete insulin; this option should at least be delayed as long as possible (Aston-Mourney et al., 2008); the primary focus may be to overcome insulin resistance when starting a therapy (see below the relevance of sulfonylureas and incretins for the therapy in the future).

3.6.2. Effects on B-cell mass

Some remodelling of size and function of islet B-cells occurring throughout life may play a role in the maintenance of normal glucose tolerance (Bonner-Weir, 2000). In addition to functional abnormalities destabilizing the islet cell mass primarily through changes in numbers and size of B-cells, a disturbed interplay of division and growth of these cells (neogenesis, rates of apoptosis and even loss) is very important (Finegood et al., 1995; Butler et al., 2003). An inhibition of B-cell apoptosis by GLP-1 is observed in human B cells (Farilla et al. 2003; Buteau et al. 2004). Therefore, any agent that normalizes this balance may delay or prevent the decline in insulin secretory capacity. GLP-1 (including its analogues) possibly has this disease-modifying capability in type 2 diabetes (Xu et al., 1999; Stoffers et al., 2000; Buteau et al., 2001; Baggio & Drucker, 2006; Aaboe et al., 2009).

GLP-1 seems to stimulate the proliferation of developed B-cells. In addition GLP-1 stimulates the differentiation and maturation of foetal islet cells to insulin-producing islet cells (Abraham et al., 2002). In animals a GLP-1 analogue stimulates islet cell differentiation from ductal progenitor cells or exocrine cells (Xu et al., 1999; Zhou et al., 1999). Activation of GLP-1 receptor signaling promotes conversion of human or rat pancreatic ductal cell lines to islet-like cells that produce insulin in a glucose-dependent fashion (Hui et al., 2001; Miyawaki et al., 2002; Bulotta et al., 2002; Drucker, 2003).

The effect of GLP-1 on B-cell differentiation involves the transcription factors PDX-1 and its upstream regulator Foxa2 (formerly HNF3β) (Hui et al., 2001; Miyawaki et al., 2002; Bulotta et al., 2002), and appears to be mediated by MAP kinases (Miyawaki et al., 2002), including extra-cellular signal related kinase (ERK) (Gomez et al., 2002), and an isoform of PKC (Buteau et al., 2001). Besides CREB (Fig. 5), the key intermediary compounds of GLP-1 signaling that mediate more chronic B-cell effects include insulin receptor substrate 2 (IRS-2) and the transcription factor PDX-1 (Drucker et al., 1987; Wang et al., 1997; Skoglund et al., 2000; Wang et al., 2005; Bregenholt et al., 2005). Activation of PDX-1 appears to be involved in all cytoprotective effects of GLP-1. Enhanced PI-3K activity as a result of IRS-2

phosphorylation or EGFR transactivation leads to the upregulation of PDX-1 via nuclear exclusion of FoxO1 (Buteau et al., 2006). Transactivation of epidermal growth factor receptor (EGFR) is involved in inducing PI-3K activity (proliferation) (Buteau et al., 2003). Support for involvement of the transcription factor NFAT (nuclear factor of activated T-cells) has been presented (Lawrence et al., 2002; Heit et al. 2006). The data are summarized in Fig. 5.

The role of neogenesis of endocrine precursors from pancreatic duct cells has been questioned as a significant mechanism for changes in B-cell mass in vivo (Dor et al., 2004; Dor, 2006; Teta et al., 2007).

GLP-1 and similar receptor agonists reduce apoptosis in primary rodent islets or islet cell lines exposed to cytotoxic agents, such as cytokines, free fatty acids, hydrogen peroxide, or streptozotocin (Perfetti et al., 2000; Pospisilik et al., 2002; Buteau et al., 2003; Hui et al., 2003; Bregenholt et al., 2005).

The data raise the interesting possibility that activation of the GLP-1 receptor could increase B-cell mass, a novel and potentially effective approach to inhibit the progression of diabetes. It has to be differentiated between in vitro/animal studies and the extrapolated clinical outcome. It is important to note that despite the accumulation of an impressive body of evidence the GLP-1 receptor signaling promotes the expansion or protection of B-cell mass; this information is limited to studies of cultured cells or rodent models. There are important differences in rates and capacity for islet cell turnover and growth between rodents and humans (Butler et al., 2007), such that extension of findings from animal studies cannot be assumed. The mechanisms of prolonged effects are summarized in Fig. 6 (Brubaker & Drucker, 2004; Li et al., 2005). Some of these effects may have a therapeutic implication (see details later). Thus basic research in this area has proceeded over the past years, albeit translation of this work to human application has not yet been accomplished. Because it is not currently possible to evaluate B-cell mass non-invasively in humans the question of GLP-1 effects on B-cell proliferation and apoptosis cannot be reliably addressed in longitudinal or intervention studies.

Chronic effects of GLP-1, such as gene transcription through the actions of CREB (Fig. 5), activation of cellular growth and inhibition of

apoptosis, can be attributed to PI-3K downstream pathways possibly including PI3-K, protein kinase B (PKB) and protein kinase C (PKC) (Buteau et al., 2001; Wang & Brubaker, 2002; Buteau et al., 2003) enhancement of insulin biosynthesis (Drucker, 2006), increased resistance to apoptosis and enhanced B-cell survival (Li et al., 2003; Farilla et al., 2003).

WNT signalling (WNT = cell-secreted glycoprotein ligand) is a key pathway for B-cell growth and differentiation. There are some hints that GLP-1 is linked as a direct activator of the WNT pathway (Gustafson & Smith, 2008).

The other incretin GIP has similar proliferative and anti-apoptotic effects on B-cells using rather the same intracellular mechanisms (PKA activation, PI-3K/PKB activity and p38 MAPK) (Trumper et al., 2001; Wideman & Kieffer, 2004). Administration of a GIP-antagonist for 11 days to *ob/ob* mice partially reversed islet hypertrophy with a concurrent improvement of hyperinsulinemia and insulin resistance (Gault et al., 2005). There are yet no studies of chronic effects of GIP receptor agonists in humans; thus the intriguing results of these animal studies have not been extended to possible clinical applications as it holds for GLP-1 receptor agonists.

3.6.3. Effects on the gastrointestinal tract and exocrine pancreas

GLP-1 inhibits gastrointestinal secretion and motility (Wettergren et al. 1993; Nauck et al. 1997). It has further been established that GLP-1 inhibits pancreatic exocrine secretion independently of its inhibition of gastric emptying (Groger et al. 1997).

3.6.4. Central effects with respect to appetite and food intake

GLP-1 receptors in the brain are expressed particularly in the regions responsible for the regulation of food intake. Intracerebroventricular administration of low GLP-1 doses results in inhibition of food intake (Göke et al., 1995b; Tang-Christensen et al. 1996; Turton et al., 1996). Interestingly also peripheral administration of GLP-1 enhances satiety and reduces food intake in normal subjects (Flint et al., 1998; Verdich et al., 2001), as well as obese subjects (Naslund et al., 1999a) and type 2 diabetic patients (Gutzwiller et al., 1999a; Zander et al., 2002). The local mechanisms of these effects are unclear. As the enhancement of satiety and decrease of hunger is preserved in fasting subjects (Gutzwiller et al., 1999b), it is unlikely that the mechanisms are direct, i.e. related to the effect on gastric emptying and motility. This is corroborated by experiments when exendin 9-39 is given intracerebroventricularly.

3.6.5. Cardiovascular effects (GLP-1)

It is evident from in vitro and in vivo experiments that GLP-1 has beneficial effects on myocardial function: in rat models of myocardial ischemia a reduction in infarction size has been demonstrated (Bose et al., 2005). GLP-1 improves myocardial contractility and glucose uptake in normal and postischemic rat hearts (Zhao et al., 2006). In humans with type 2 diabetes and congestive heart failure, GLP-1 may improve myocardial function (Thrainsdottir et al., 2004; Nikolaidis et al., 2004).

GLP-1 has been shown to induce an endothelial-dependent reduction in vascular tone of rat lungs (Golpon et al., 2001) and improves endothelial function in a salt-sensitive rat model (Yu et al. 2003). In type 2 diabetic patients with established coronary artery disease an expression of GLP-1 receptors in coronary artery endothelial cells was observed and a beneficial effect of GLP-1 infusion on endothelial function was demonstrated (Nystrom et al., 2004).

4. Pathophysiological impact of the incretin concept (similarities and differences between the incretins GLP-1 and GIP in pathophysiological terms)

Failure of incretin effect leads to a relative insulin deficiency and hyperglycaemia. The incretin effect (oral glucose elicits a considerably




Fig. 6. Mechanisms involved in apoptosis and proliferation with respect to GLP-1 (modified from Brubaker & Drucker, 2004). (Bcl-2 and Bcl-xL are antiapoptotic proteins; PDX-1 is pancreatic and duodenal homeobox 1, a transcription factor; c-src is a protein with tyrosine kinase activity).

Table 2
Pathophysiological state of type 2 diabetes and related biological effects of GIP and GLP-1.

Feature of type 2 diabetes	GIP	GLP-1
↓ Insulin secretion	Ineffective in type 2 diabetics	Effectiveness preserved in type 2 diabetics; ↑ Insulin secretion
↓ Insulin synthesis		↑ Insulin synthesis
Hyperglucagonaemia (↑ Glucagon secretion)		↓ Glucagon secretion (possibly indirectly)
↑ B cell apoptosis	↓ B cell apoptosis	↓ B cell apoptosis
↓ B cell mass	↑ B cell replication (humans?)	↑ B cell replication (humans?)
Obesity		↓ Food intake, ↓ body weight
↑ ± gastric emptying	No effect	↓ gastric emptying
Hyperlipidemia	No effects	↓ triglycerides, ↓ free fatty acids
Insulin resistance	No effects	No immediate effect (↑ Insulin sensitivity (probably))

(↑ increase, ↓ decrease, ± no effect) (adapted from Van Gaal et al., 2008).

higher insulin secretory response compared to the same amount of intravenous glucose; see above) was believed to be substantially reduced in patients with type 2 diabetes (Nauck et al., 1986; Nauck et al., 2004b). Studies have reported that plasma GLP-1 levels are increased (Ørskov et al., 1991; Fukase et al., 1995), decreased (Vaag et al., 1996; Vilsbøll et al., 2001; Toft-Nielsen et al., 2001a; Vilsbøll et al., 2003b), and not changed (O'Donovan et al., 2004; Ryskjaer et al., 2006; Knop et al., 2007) in this situation. These differences are probably due to technical discrepancies using different assays and due to the relatively small sample sizes studied. Regardless, in studies where GLP-1 secretion has been reported to be reduced in diabetic subjects the relative decrease was modest, suggesting that GLP-1 deficiency probably does not contribute significantly to B-cell dysfunction in type 2 diabetes. The GLP-1 release, but not that much the GLP-1 effect is hampered in type 2 diabetics (Højberg et al., 2007) which, however, can be improved by a 4-week normalization of blood glucose (Højberg et al., 2007, 2008). Supplementing the incretin GLP-1 can overcome this type of defect in type 2 diabetics.

There is no evidence for reduced GIP secretion as an explanation for a reduced incretin effect in diabetes. GIP and fortunately not GLP-1 shows attenuated insulinotropic action in patients with type 2 diabetes (Nauck et al., 1993a). GIP, by contrast, has proved to be disappointing in terms of therapeutic use since its intravenous administration is not associated with unequivocally improved insulin secretion and glucose tolerance (Elahi et al., 1994). GIP is an insulin secretagogue in healthy subjects but is relatively inactive in patients with type 2 diabetes. The defect in GIP is acquired rather than representing a primary feature of type 2 diabetes (Meier et al., 2003b). The defect is reversible, and GIP responsiveness can be restored by sulfonyleureas or by improving blood glucose control (Menelly et al., 1993; Aaboe et al., 2007; Højberg et al., 2007). Studies in type 2 diabetes report decreased, normal and increased GIP secretion (Ebert & Creutzfeldt, 1987; Krarup, 1988; Nauck et al., 1993a; Fehmann et al., 1995; Toft-Nielsen et al., 2001a; Vilsbøll et al., 2003a).

GLP-1 secretion is reduced in non-diabetic obese subjects, suggesting that incretin secretion may be altered early in the natural history of diabetes (Ranganath et al., 1996). Since GLP-1 actions remain relatively preserved in diabetic patients, the pharmaceutical efforts have mainly focused on GLP-1 receptor agonists since they are of higher therapeutic relevance than GIP.

It now has to be concluded that the impaired incretin effect is a result of the diabetic state rather than a primary pathogenic factor (Vaag et al., 1996; Meier et al., 2005b). The paradigm has changed: it is now clear that diabetes affects the incretin hormones.

In Table 2 the pathophysiological situation in type 2 diabetes including the influences of GIP and GLP-1 is summarized.

Increased circulating glucagon levels are not suppressed in type 2 diabetics after a carbohydrate-rich meal and occurs early in prediabetes (Muller et al., 1970; Unger et al., 1970; Åhren and Larson,

2001). The resultant hyperglucagonaemia contributes both to an inability to suppress hepatic glucose production postprandially and to excessive plasma glucose excursions. Therapies currently available do not influence the hyperglucagonaemia; an approach that targets glucagon suppression (Table 2) would offer a valuable mechanism to improve glycaemic control.

Gastric emptying may be more rapid in type 2 diabetes and may be a key determinant of postprandial glucose excursions (Phillips et al., 1992; Horowitz et al., 1994). A slowing in gastric emptying occurs to improve the match between rise in glucose due to nutrient absorption and glucose disappearance from the circulation, despite which, the gastric half-emptying time is significantly shorter in patients with type 2 diabetes. GLP-1 inhibits the gastric-emptying rate (Table 2).

Effects as a result of an incretin lack and therapeutic effect of incretins are summarized in Table 3.

One characteristic of diabetic islet function is the classic absence of first-phase insulin release (Table 4), an apparent loss of B-cell sensitivity to glucose (Byrne et al., 1996), and delayed insulin secretion in response to oral glucose (Table 4). This may be partly due to a decrease in the actions of GI hormones and neural signals, engaged after meals, to potentiate the insulin response (Quddusi et al., 2003). Additionally there may be a disruption of normal pulsatile insulin secretion (Polonsky et al., 1988), abnormal potentiation of non-glucose secretagogues, and a decreased maximal secretory capacity (Ward et al., 1984).

Two different approaches can be used: one would be to administer incretin(s) exogenously (incretin mimetics). An alternate strategy would be to enhance endogenous incretins by inhibiting their degradation (= incretin enhancers).

There are some aspects (pathophysiological factors) in type 2 diabetes addressed now (Table 4), but not fully addressed by other antidiabetic treatments (Table 5) except incretin mimetics (avoiding weight gain and metformin (hyperglucagonaemia).

5. GLP-1 receptor agonists (exenatide)

GLP-1 itself is rapidly degraded and could only be used as a continuous infusion (Nielsen & Baron, 2003). Chemically modified degradation-resistant peptides with an acceptable pharmacokinetic profile have been developed. In the search for those peptides GLP-1 was compared with known structures from databanks. Exendin-4 (36 amino acids) was discovered in a lizard (salivary gland venom of the Gila monster) and found to be more stable and less rapidly degraded than GLP-1 (Eng et al., 1992; Göke et al., 1993; Raufman, 1996). Although exendin-4 (synthetic = exenatide) is transcribed from a distinct gene, it has a 53% overlap of the amino acid sequence with mammalian GLP-1 (Chen & Drucker, 1997) (Fig. 2). Exendin-4 has an in vivo potency up to 5–10 times higher than GLP-1 itself.

As a GLP-1 agonist, exenatide binds to the pancreatic GLP-1 receptor and improves glucose homeostasis by mimicking the actions of naturally occurring GLP-1: it stimulates insulin release in a glucose-dependent manner (Egan et al., 2002a; Kolterman et al., 2003; Degn et al., 2004a), improves first phase insulin release (Fineman et al.,

Table 3
Consequences of incretin deficiency and substitution (modified from Ranganath, 2008; Holst et al., 2008).

Presumed phenotype	Incretin actions correcting this
Weight gain	Decrease in food intake
Postprandial hyperglycaemia	Delay in gastric emptying, increase in glucose-induced insulin secretion
Hyperglucagonaemia	Increase in glucose-mediated suppression of glucagons
Increase in loss of B cells	Increase in B cell neogenesis, proliferation snf decrease in apoptosis
Delayed insulin response (lacking first phase insulin release) (partly mediated)	Reversed

Table 4

Comparison of defects in type 2 diabetes and biological actions of GLP-1, incretin mimetics, and DPP-4 inhibitors (partly modified from several authors: Drucker & Nauck, 2006; Åhrén, 2007; Handelsman, 2008).

Defect in type-2 diabetes	Biological action of compounds	GLP-1, Exenatide, Liraglutide	Sitagliptin, Vildagliptin (DPP-4 inhibitors)
↓ insulin secretion	↑ insulin secretion (glucose-dependent)	Yes	Yes
Lack of first phase insulin secretion	Restoration of first phase insulin secretion	yes ^a	Not known
↓ incretin effect	Replacement of incretin effect	yes	Yes
↑ glucagon secretion	↓ glucagon secretion	yes	Yes
Hypoglycemia counterregulation	↑ Glucagon secretion at low glucose	yes	Not known
↓ βcell insulin content	↑ proinsulin biosynthesis	yes	Yes
↓ βcell mass	↑ proliferation of βcells	yes	Not known
↑ βcell apoptosis	↓ βcell apoptosis	yes	Not known
No pathophysiological equivalent	↓ velocity of gastric emptying	yes	No (nearly not detectable)
↑ energy/food uptake	↓ energy/food uptake	Yes	No
↑ weight gain	↓ weight gain or even weight loss	Yes	No

Note: the outcome of data presented in the table is in part from animal studies. The first phase insulin release is a measure of clinical diagnosis albeit artificial because not seen after an oral glucose load (slow glucose increase).

^a Not known for liraglutide.

2003; Quddusi et al., 2003; Fehse et al., 2005), induces a delay of gastric emptying (Kolterman et al., 2003, 2005; Blase et al., 2005) and decreases food intake. It improves the GLUT-4 expression in target cells of STZ-diabetic rats (Tornerio-Exteban et al., 2007). On a long run B-cell function is improved for which a permanently ongoing treatment is necessary (Bunck et al., 2007).

5.1. Clinical aspects

The incretin mimetic exenatide has been developed for the treatment of type 2 diabetes (Polonsky et al., 1988; Gerich, 1989; Rachman et al., 1997; Quddusi et al., 2003; Degn et al., 2004a; Fehse et al., 2005; Kolterman et al., 2005). The efficacy of adding exenatide to ongoing therapy in patients suboptimally controlled by oral anti-diabetic agents has been investigated: metformin (DeFronzo et al., 2005), sulfonylureas (Buse et al., 2004) including a combination of both (Kendall et al., 2005) or thiazolidinediones (Zinman et al., 2006). Exenatide is indicated as an adjunctive therapy to improve glycaemic control in patients with type 2 diabetes mellitus who are already receiving metformin, a sulfonylurea, or both but continue to have suboptimal glycaemic control. Exenatide is approved only for the use in combination with metformin, a sulfonylurea, a thiazolidinedione or the combination of metformin with either a sulfonylurea or with a thiazolidinedione. Exenatide nevertheless is effective as monotherapy, but this is not an immediately attractive option for most patients with respect to the type of administration. Not surprisingly, there also is some interest in using GLP-1 mimetics for weight loss in patients without diabetes, but this has not yet been approved.

Appropriate doses were derived from various studies (Buse et al., 2004; DeFronzo et al., 2005; Kendall et al., 2005; Heine et al., 2005; Zinman et al., 2006). 5 or 10 µg exenatide by two subcutaneous injections per day shows reductions in fasting and postprandial glucose concentrations and in weight loss (2–5 kg). Given chronically it enhances hepatic insulin sensitivity and splanchnic glucose uptake

(Cersosimo, 2007). Exenatide reduces HbA_{1c} concentrations by 0.8–1.0% (Buse et al., 2004; DeFronzo et al., 2005; Heine et al., 2005; Kendall et al., 2005; Kim et al., 2006; Zinman et al., 2006) for a long time (over 30 weeks), with prevention of weight gain or even induction of a modest weight loss of 1.5–3 kg which may be higher (4–5 kg) after 80 weeks (Riddle et al., 2006).

Sixteen-week treatment with exenatide added to existing thiazolidinedione treatment in type 2 diabetic patients suboptimally controlled with or without metformin reduced HbA_{1c} by 0.98% from a mean baseline value of 7.8–8.0%, with a weight loss of 1.51 kg (Zinman et al., 2007). The addition of either insulin or exenatide in patients not sufficiently controlled by sulfonylurea and metformin resulted in a better reduction of HbA_{1c} in the exenatide-group and in a weight reduction only in this group (weight increase in the insulin-group) (Nauck et al., 2007a) although a more stringent insulin regime might have equalized the differences. Among patients with type 2 diabetes requiring insulin therapy, exenatide exhibits less potent HbA_{1c} reduction compared with its combination with oral agents (Davis et al., 2007); this observation was expected because such individuals tend to be more insulin deficient with less available insulin secretory reserve.

The main metabolic benefits demonstrated from exenatide therapy have been on indexes of glycaemic control.

On a long term exenatide has additional effects: it also improves markers of hepatic functions such as ALT and AST (Okerson et al., 2007); nonalcoholic fatty liver disease often coexists with diabetes. The observed weight reduction over time may have an indirect benefit on cardiovascular risk, including blood pressure, cholesterol levels, inflammatory markers, and insulin resistance. One clinical report on exenatide suggests a modest benefit on certain cardiovascular risk factors (Buse et al., 2007). GLP-1 receptors have been demonstrated in cardiac myocytes and in those regions of the brain that regulate autonomic function (Bullock et al., 1996). The observed effects on blood pressure are controversial: GLP-1 may decrease (Avogaro, 2008) or increase (Yamamoto et al., 2002) blood pressure in humans. Rat

Table 5

Drugs used in type 2 diabetes and effects on factors with pathophysiological relevance (+ = improved or increased; - = reduced; 0 = not changed; ? = questionable; -/? = reduced or questionable effect) (adapted from Ranganath, 2008).

	Bigua-nide (Met-formin)	Thiazoli-dindione	α-Glucosi-dase inhibitor	Sulfo-nylurea	Glinides	Insulin	Amylin	Glucagon-like peptide-1 mimetics	Dipeptidyl peptidase-4 inhibitors
Insulin deficiency	0	0	0	+	+	+	0	+	+
Insulin resistance	+	+	0	0	0	0	0	+	0
Hepatic glucose output	+	+	0	0	0	0	0	-?	0
Excess glucagon	0	0	0	0	0	0	-	-	-?
Gastric emptying	0	0	-	0	0	0	-	-	-/?
Body weight	0	+	0	+	0	+	-/?	-	0
B-cell mass	0	+	0	0	0	0	0	+	+
B-cell function	0	+	0	0	0	0	0	+	+

experiments (antihypertensive or hypertensive effect of GLP-1 (Barra-gán et al., 1996, 1999; Yu et al., 2003; Okerson et al., 2008)) especially those using the intracerebroventricular application (increase in blood pressure by GLP-1) may not be helpful (Isbil-Buyukcoskun and Gulec, 2004). The heart rate may be increased indirectly (Yamamoto et al., 2002) as well as a positive inotropic effect may be induced. Liraglutide decreases blood pressure. It is still a matter of speculation whether GLP-1-based drugs might be useful as adjunctive therapy in patients with decompensated heart failure. Improvements in high density lipoprotein cholesterol and triglycerides were reported (Blonde et al., 2006) but need further verification.

Exendin-4 has also extrapancreatic effects: increase in glucose transport in muscle (Moreno et al., 2007), glucose transporter expression and glycogen synthesis in liver (Arnés et al., 2008).

5.2. Pharmacokinetics (note: some data information were only available from a company)

The s.c. absorption of exenatide is quick. Its relative bioavailability is 93–97% and similar when exenatide is injected s.c. into the upper arm, abdomen or thigh (Calara et al., 2005). The starting dose of exenatide is 5 µg twice daily for 4 weeks, followed eventually by an increase to 10 µg (Fineman et al., 2002, 2004). Since exenatide is DPP-4 resistant, its pharmacological action lasts for 4–6 h after a single subcutaneous injection (Nielsen & Baron, 2003; Nielsen et al., 2004). It reaches peak plasma concentrations after approximately 2 h, its circulating half-life is 2.4 h, and it is predominantly eliminated by glomerular filtration (N.N., 2005; Kolterman et al., 2005). The pharmacokinetics of exenatide is linear with respect to AUC (dose-dependency), albeit not with respect to C_{max} (reviewed by Iltz et al., 2006). The V_d is 28.3 l (after a single s.c. dose).

Since exenatide will be degraded enzymatically in the kidneys, it is not recommended for patients with severe renal insufficiency (creatinine clearance < 30 ml/min); no dose adjustment, however, is necessary in patients with mild to moderate renal insufficiency. Hepatic dysfunction is not expected to influence exenatide pharmacokinetics (see elimination pathways above). Obesity does not appear to modify the pharmacokinetic profile of exenatide. The fetal/maternal exenatide concentrations are < 0.017 (*ex vivo* study) (Hiles et al., 2003).

5.3. Side effects

Nausea is the most commonly (45–51%; placebo 7–23%) (Mikhail, 2008) reported adverse effect of exenatide (see also other GLP-1 receptor agonists), which lessens over time. It is more obvious in fasting subjects and is probably a direct central effect. Other adverse events occurring in >10% of patients are diarrhoea (12.8%), and vomiting (12.8%) (Buse et al., 2004; DeFronzo et al., 2005; Kendall et al., 2005). The etiology of nausea is not fully clear, but may be partly related to the delay in gastric emptying. Nausea is not the major reason for weight loss induced by exenatide, since there was no correlation between change in body weight and the duration of nausea (DeFronzo et al., 2005; Kendall et al., 2005).

Hypoglycaemic events are mainly observed when combinations are used: in 20–30% of the patients when being treated together with a sulfonylurea and in 13% of the patients treated in combination with a thiazolidinedione (Buse et al., 2004; Kendall et al., 2005; Zinman et al., 2007). When sulfonylurea doses were reduced, the incidence of hypoglycemia decreased from 26.9 to 6.1 events/patient-year (Nauck et al., 2007a). When exenatide produces hypoglycaemia (especially in combination therapy) the counter-regulatory responses to hypoglycaemia are unimpaired (Degen et al., 2004a). Hypoglycaemias were not observed when it was given together with metformin (DeFronzo et al., 2005).

Other side effects of exenatide are feeling jittery (12–15%), dizziness (9–15%), constipation (9%), sweating (8%), and backache (6%) (Mikhail, 2008).

Exenatide was rarely (5–10%) discontinued because of side effects (Buse et al., 2004; DeFronzo et al., 2005; Kendall et al., 2005; Heine et al., 2005; Mikhail, 2008).

40–50% of patients develop anti-exenatide antibodies with low binding affinity and low titres (Buse et al., 2004; DeFronzo et al., 2005; Kendall et al., 2005) which, however, did not affect patients' clinical glycaemic control in most cases. About 6% of patients had high titres of antibodies; in about one half of these the glycaemic response to exenatide appeared to be diminished (N.N., 2005).

In 2006 the first case of acute pancreatitis was reported in a patient, who had been treated with exenatide and neutral protamine Hagedorn insulin (Denker & Dimarco, 2006). Recently, the US Food and Drug Administration reviewed 30 post-marketing reports of acute pancreatitis in patients: an association between exenatide and acute pancreatitis has been suspected in some of these cases (N.N. Food and Drug Administration Information for Healthcare Professionals Exenatide, 2008); it is, however, not clear whether the incidence of acute pancreatitis in patients receiving exenatide is higher than expected in diabetic patients for which it is known to be increased generally. Most of the patients had an additional risk factor for pancreatitis such as gall stones, severe hypertriglyceridemia or alcoholism. A causal association with exenatide is not clear. Nevertheless US manufacturer labelling was updated in this respect giving the advice: "Patients should be informed that persistent severe abdominal pain, which may be accompanied by vomiting, is the hallmark symptom of acute pancreatitis. If pancreatitis is suspected exenatide and other potentially suspect drugs should be discontinued, confirmatory tests performed and appropriate treatment initiated." The problem of pancreatitis is unsolved and still to be discussed (Ahmad & Swann, 2008).

5.4. Drug and food interactions (absorption problems)

Exenatide should be administered 60 min before a meal (in accordance with its mechanism of action); it should not be administered after a meal.

Due to the delay of gastric emptying induced by exenatide, a modified absorption of oral compounds simultaneously given can be expected. A decrease of e.g. paracetamol (acetaminophen) plasma concentrations can be seen (Kolterman et al., 2003; Blase et al., 2005; Iltz et al., 2006). Compounds of which the effectiveness relies on a minimal drug concentration (e.g. oral contraceptives, antibiotics) should be given 1 h before an exenatide injection. Acid resistant preparations (e.g. proton pump inhibitors) should not be given simultaneously with exenatide. When e.g. digoxin (Kothare et al., 2005), lisinopril (Iltz et al., 2006) or warfarin are given 30 min after exenatide, the t_{max} values are delayed by 2 h; in the case of lisinopril, however, no clinical effect was observed. Exenatide decreases the AUC and C_{max} of lovastatin severely.

Exenatide does not appear to influence the pharmacokinetics of sulfonylureas or of metformin, used for a combination therapy (see above).

Erythromycin reverses the delay of stomach emptying induced by GLP-1 (Meier et al., 2005a).

5.5. Contraindications

Contraindications are ketoacidosis and kidney failure (see above).

Exenatide should not be used in type 1 diabetics although it was very recently claimed to have promising effects (interesting because of weight loss, see information above) (reviewed by Aaboe et al., 2009).

5.6. Comparison with other therapy regimes (insulin glargine) or replacement of insulin by exenatide

Exenatide has been compared with insulin glargine when either compound was added to metformin and a sulfonylurea because an

effective glucose control was not reached (Heine et al., 2005): fasting glucose concentrations were reduced more effectively by insulin glargine, albeit postprandial glucose reduction was better with exenatide. Patients receiving insulin glargine gained an average of 1.8 kg compared with a 2.3 kg weight loss observed in exenatide-treated patients; no significant differences were seen in levels of HbA_{1c} (reduction by 1.1% over 26 weeks) or in hypoglycaemias as a side effect (7.3 events/patient-year) compared with insulin glargine patients (6.3 events/patient-year) (Heine et al., 2005).

Replacing insulin treatment by exenatide therapy does not seem to be advisable (Davis et al., 2007). In a crossover non-inferiority study, the overall incidences of hypoglycaemia were <15% in patients on exenatide compared with 25% on insulin glargine treatment (Barnett et al., 2007). This was rather the same with respect to hypoglycaemias when it was compared with insulin aspart.

5.7. Aspects for the future

Several therapeutic implications need clinical confirmation: the durability of the weight loss, the ability to preserve functional B-cell mass and the applicability to other patients than type 2 diabetics. Long-term studies especially with cardiovascular end-points are needed to confirm the probable benefits of this new class of anti-diabetic drugs.

6. Longer acting GLP-1 analogues

Although intravenous or subcutaneous GLP-1 application could be useful for the short-term control of hyperglycaemia (Meier et al., 2004; Nauck et al., 2004c), long-term treatment of type 2 diabetes is necessary and application therefore, should be less than twice a day as for exenatide.

6.1. Liraglutide

Liraglutide (NN 2211) is a long-acting GLP-1 analogue with the following chemical modification (Arg34Lys substitution, and a glutamic acid and 16-C free-fatty-acid addition to Lys26) (Fig. 2) (Knudsen et al., 2000; Juhl et al., 2002). The fatty-acyl-GLP-1 structure binds to interstitial albumin at the injection site and the GLP-1 resembling structure is released slowly from the albumin complex to be absorbed into the circulation.

Liraglutide inhibits appetite (Hansen et al., 2001). The counter-regulation by glucagon induced by a hypoglycaemia is not impaired by liraglutide (Nauck et al., 2003).

Liraglutide inhibits apoptosis (Bregenholt et al., 2001) and increases B-cell mass (Rolin et al., 2002).

6.1.1. Clinics

The clinical effects of liraglutide are similar to those of GLP-1 (Saad et al., 2002; Juhl et al., 2002; Degn et al., 2004b). In phase II studies 14 weeks of treatment as monotherapy with 2 mg (the highest dose) reduced fasting plasma glucose and HbA_{1c} was lowered up to 1.75% from a baseline value of 8.1–8.5%. Liraglutide reduces HbA_{1c} by 0.7% (Madsbad et al., 2004; Mikhail, 2008). Patients experienced a weight loss of approximately 3 kg (1.2 kg corrected for placebo) (Viltsbøll et al., 2007) or at least weight gain was prevented (Nauck et al., 2004a, 2006). Overall liraglutide has similar effects like the incretin mimetic exenatide. Recently several studies called LEAD (Liraglutide Effect and Action in Diabetes) were presented on congresses and/or were published (Marre et al., 2009; Nauck et al., 2009); higher doses (1.2 and 1.8 mg) were much more effective than 0.6 mg with respect to various parameters such as HbA_{1c} in absolute terms and as a percentage of improvements with respect to the envisaged goal. Nausea did not increase in a dose-dependent manner. Liraglutide was superior to rosiglitazone (either compound was combined with

glimepiride), liraglutide was superior to glimepiride (when both were combined with metformin) with respect to reduction of body weight and the occurrence of hypoglycaemias whereas the glycemic control was the same.

Furthermore, liraglutide has been shown to significantly lower systolic blood pressure by 7–9 mm Hg (Viltsbøll et al., 2007). Atherosclerosis is a major contributor to morbidity and mortality in diabetes. Liraglutide improves parameters which are associated with atherosclerosis: it inhibits TNF α -induced PAI-1, VCAM-1, ICAM-1, and E-Selectin mRNA and protein secretion in an in vitro model of vascular endothelial cell dysfunction (Dear et al., 2007): E-selectin is involved in inflammation, recruiting leucocytes, VCAM is important for the immune system, recruiting monocytes to atherosclerotic sites, ICAM is an intercellular adhesion molecule and is involved in subarachnoid hemorrhage, and PAI-1 has an impact on thrombosis and cell migration.

6.1.2. Pharmacokinetics

Liraglutide (modified GLP-1-albumin complex) results in a favourable pharmacokinetic profile because it resists DPP-4 metabolism, is slowly absorbed from injection site with a maximum after 9–12 h (Jonker et al., 2008), and its renal clearance is reduced. It is first absorbed in zero-order kinetics and later within first-order kinetics, the absolute bioavailability is 51% (Jonker et al., 2008), the V_d is very low indicating that the compound is concentrated mainly in the central compartment, and the clearance is in the range of 6 ml/h \times kg and not dose-dependent (Jonker et al., 2008). After s.c. injection, liraglutide has a half-life of 10–14 h allowing once-daily injection (Agero et al., 2002; Elbrond et al., 2002; Jonker et al., 2008). The acyl moiety promotes non-covalent binding to albumin with 1–2% of liraglutide circulating as the non-albumin bound free peptide (Knudsen et al., 2003). The variations of plasma levels are lower when liraglutide is given once compared to when exenatide is given twice a day (Jonker et al., 2008). The liraglutide dose is between 0.45 and 0.75 mg daily (Madsbad et al., 2004; Nauck et al., 2006; Viltsbøll et al., 2006).

Liraglutide kinetics is not significantly modified in patients with renal kidney insufficiency (Jacobsen et al., 2007, 2008) or liver impairment (Flint et al., 2008).

6.1.3. Side effects

As expected from exenatide, GI symptoms (e.g., nausea, vomiting) are prominent adverse effects of liraglutide and led to discontinuation in 3% of patients (Viltsbøll et al., 2007). Side effects are generally mild and transient (Nauck et al., 2006; Viltsbøll et al., 2006). No patient experienced any hypoglycaemic episodes, and there was no anti-liraglutide antibody induction (Viltsbøll et al., 2007) as it was observed for exenatide (see above).

In Table 6 liraglutide is compared with exenatide. Long term data are needed to fully assess the potential of liraglutide in treatment of type 2 diabetes.

The impact of liraglutide on the absorption of simultaneously administered drugs is summarized and appears to be low (Zdravkovic et al., 2008).

Table 6
Comparison of effects with respect to exenatide and liraglutide (adapted from Ranganath, 2008).

Exenatide	Liraglutide
Twice daily injection	Once daily injection
Plasma levels: rise to a peak and then a fall	More or less peakless plasma levels
Significant weight loss	Weight loss
Good effect in HbA _{1c}	Good effect on HbA _{1c} and fasting glucose
Antibodies to exenatide	No antibodies to liraglutide
Injection site reactions occur	Negligible injection site reaction

6.2. Exenatide long-acting release (exenatideLAR)

The action of one subcutaneous injection of exenatide lasts for only 6–8 h. Exenatide long-acting release (LAR) is a special preparation containing a polylactide-glycolide microsphere suspension with biodegradable microparticles (polymers: Medisorb®) with 3% exenatide peptide. This leads to a sustained dose-dependent glycaemic control in diabetic rats, for up to 28 days after only one s. c. injection (Gedulin et al., 2005). It was planned to be given once weekly to patients (Kim et al., 2006).

Experience with this dose regime of exenatideLAR indicates a much greater reduction in fasting glucose concentrations and HbA_{1c} compared with normal exenatide administered twice daily (Kim et al., 2006). In a small, trial with patients suboptimally controlled by metformin and/or diet, 15-week treatment with 2.0 mg LAR once a week showed a reduction in HbA_{1c} by 1.7% from a baseline value of 8.5% and a weight loss of roughly 4 kg (Kim et al., 2007).

Altogether the efficacy with respect to fasting blood glucose and postprandial glucose appears to be similar and the side effects are not different compared to normal exenatide. The pharmacokinetics will be different (Trautmann et al., 2008) although many details are not yet available. It has to be mentioned that 14 injections of normal exenatide is now reduced to once each week, that the exenatide levels in the periphery are kept a little bit lower (to avoid peak values mainly responsible for above mentioned side effects), that there is no tachyphylaxis (vanishing of effects) as often recorded after continuous application of hormones, and that the time point of injection is not important at all.

6.3. Albiglutide (Albugon)

Recombinant albumin-GLP-1 fusion proteins have been developed that mimic the full range of GLP-1 actions (Baggio et al., 2004). Albiglutide is one example: it stimulates GLP-1 receptor-dependent pathways coupled with glucose homeostasis and gastrointestinal motility, improves insulin secretion, and reduces blood glucose (Baggio et al., 2004). Little clinical information is currently available regarding this drugs safety in humans (Kim et al., 2003; Baggio et al., 2004). Albiglutide has less potent anorectic effects in animal studies compared to exenatide and liraglutide, and it is not clear whether this disparity is secondary to the impaired blood–brain barrier's permeability due to the larger molecule and not to albiglutide itself (Baggio et al., 2004).

6.4. Taspoglutide (Ro 1583)

Taspoglutide is a matrix free sustained release formulation for GLP-1 and is just going to be converted to phase III studies.

6.5. Other compounds (Table 7)

The following strategies are used to circumvent rapid renal filtration: modification of the molecule so that the incretin binding to circulating plasma proteins is increased; this is performed by acylation, i.e. attachment of a fatty acid side-chain; see liraglutide, or by PEGylation, i.e. attachment of polyethylene glycol chains. An N-terminal modification will result in prevention of degradation by DPP-4; this is done by attaching chemical groups or by substituting amino acids with others. Another way is C-terminal modification which will reduce renal clearance; this is done by either attaching fatty acids or PEG moieties to facilitate binding to blood proteins or a direct fusion with a blood protein like albumin or transferrin.

CJC-1131 (Conjuchem) is a synthetic GLP-1 analogue produced by a single amino acid substitution and by attachment of a chemical reactant at the carboxyterminal end; it allows for covalent binding to endogenous serum albumin and is protected against DPP-4 degradation (Guivarch et al., 2004). Its half-life is similar to that of circulating

albumin, approximately 10–15 days (Sinclair & Drucker, 2005a; Leon et al., 2006). It does not seem to be of further clinical interest.

CJC-1134 is a similar conjugate of albumin, albeit with exendin-4 (Baggio & Ducker, 2006; Baggio et al., 2006; Kim et al., 2003).

Because of the naturally limited intestinal absorption of GLP-1 a series of GLP-1 analogues via site-specific conjugation of biotin-NHS and/or of biotin-poly(ethylene glycol)-NHS at Lys26 and Lys34 of GLP-1 (7–36), respectively, was developed. The resultant GLP-1 analogues, Lys26,34-DiBiotin-GLP-1 (DB-GLP-1) and Lys26-Biotin-Lys34-(Biotin-PEG)-GLP-1 (DBP-GLP-1), were investigated. DB-GLP-1 and DBP-GLP-1 show enhanced GLP-1 intestinal bioavailability. DBP-GLP-1 concentration in plasma rapidly increased 30 min after oral administration in rats. Both DB-GLP-1 and more DBP-GLP-1 had markedly better proteolytic stabilities than native GLP-1 and preserved their biological and pharmacological activities (Chae et al., 2008).

It was reported that site-specific PEGylated GLP-1 at the Lys34 position profoundly improves its enzymatic stability while retaining its biological activities, and thus enhances its therapeutic potential in type 2 diabetic *db/db* mice (Lee et al., 2005, 2006; Youn et al., 2007). Both show potent in vitro insulinotropic effects which resemble that of native GLP-1. Altogether these orally active bioconjugated GLP-1 drugs might be considered as potential oral antidiabetic agents for type 2 diabetes mellitus.

In general site-specific bioconjugation offers a convenient means of developing also orally active peptide drugs. In general many different types of peptide bioconjugation, with vitamins, fatty acids, bile acids, and PEG, have been used to develop orally active peptide agents (reviewed by Chae et al., 2008). For an oral application of GLP-1 Eligen™ technology is under investigation (Arbit et al., 2007). Eligen is a drug delivery agent due to a conformational complex with the peptide which can protect against degradation and helps the peptide to be absorbed. In HTS (high throughput screening) for an oral incretin mimetic Boc5 (a substituted cyclobutane) was identified (Chen et al., 2007).

Even intranasal administration of exenatide and pulmonary administration of GLP-1 as a Technosphere powder are under investigation (Blase et al., 2008; Cassidy et al., 2008).

For the GLP-1 receptor agonist ZP10 a prolonged-release formulation exists (Thorkildsen et al., 2003). For the novel, rationally designed peptide ZP10A (H-HGEGTFTSDLSKQMEEEAVRLFIEWLKNGGPSS-GAPPSK KKKKK-NH₂) the binding affinity for the human GLP-1 receptor is 4-fold higher than that of GLP-1 (7–36) amide (Thorkildsen et al., 2003). An antiapoptotic effect was demonstrated (Thews et al., 2008).

Albiglutide is generated by genetic fusion of a DPP-4 resistant GLP-1 dimer to human serum albumin (Stewart et al., 2008).

More modifications/substitutions of GLP-1 are summarized in a review (Green & Flatt, 2007) (Table 7).

Table 7
Other compounds.

Compound	Chemistry
CJC-1131	GLP-1 plus Gly ³⁷ replaced with Lys ³⁷ , Lys ³⁷ contains a reactive chemical linker, Ala ⁸ replaced with D-Ala ⁸
CJC-1134-PC	same molecule as CJC-1134 except that it originates from exendin-4(1-39) instead of GLP-1
ZP 10 (AVE0010), ZP 10A	going to be tested positively (Rosenstock et al., 2008a) exendin-4(1-39) plus C-terminally extended with 6 Lys residues
BIM51077	structure not disclosed
LY307161	31 amino acid analog of GLP-1; long acting
LY315902	GLP-1 plus C8 fatty acid chain linked to Lys ³⁴ , Lys ²⁶ replaced with Arg ²⁶ , His ⁷ replaced with des-His ⁷ → half-life of 3-6 hours
PEG-DAPD	PEGylated DAPD; DAPD stands for dual acting peptide for diabetes = GLP-1/glucagon hybrid peptide containing a maleimide-polyethylene glycol polymer
Taspoglutide (Ro 1583)	Matrix free sustained release formulation for GLP-1

Additionally a GLP-1/glucagon hybrid peptide was engineered that acts both as a GLP-1 receptor agonist and as a glucagon receptor antagonist; it is named “dual-acting peptide for diabetes” or DAPD.

6.6. Gene therapy with respect to GLP-1

A single administration of rAd-GLP-1 (recombinant adenoviral vector expressing GLP-1) via the tail vein into diabetic (streptozotocin) nonobese mice with severe combined immunodeficiency (NOD/SCID) resulted in remission of diabetes within 10 days, and normoglycaemia remained until the experiment was terminated (Liu et al., 2007). Intramuscular gene transfer of the plasmid coding for the GLP-1/Fc peptide in *db/db* mice demonstrated that its expression normalized glucose tolerance by enhancing insulin secretion and suppressing glucagon release (Kumar et al., 2007). Hence gene therapy leading to overexpression of GLP-1 related peptides may have a potential in the future.

7. DPP-4 (dipeptidyl-4) inhibition as a drug target

GLP-1 as an endogenous peptide is characterized by a very short half-life of 2–3 min due to the hydrolysis by DPP-4 (Fig. 2). Thus DPP-4 came into the focus as a relevant drug target (Holst & Deacon, 1998) since specific inhibitors would prevent the rapid fall of GLP-1 in circulating plasma. Two inhibitors of DPP-4 enzyme activity are in the market and many more are going to be developed.

7.1. DPP-4 enzyme characteristics

DPP-4 was already described in 1966 (Hopsu-Havu & Glenner, 1966); it is also known as CD26 regarding its activity in the immune system (see below). It is a 110 kDa plasma membrane-spanning cell-surface glycoprotein ectopeptidase ubiquitously expressed in tissues such as liver, lung, kidney, intestinal brush-border membranes, lymphocytes, and endothelial cells (Barth & Schulz, 1974; Elovson, 1980; Kreisel et al., 1984; Mrazkova et al., 1986; Imai et al., 1992; Darmoul et al., 1994; Kozakova et al., 1998; Mentlein, 1999; De Meester et al., 2000, 2003). The enzyme is also present in the circulation since the extracellular domain of DPP-4 can be cleaved from its membrane-anchored form and protrudes from endothelial cells into plasma, thereby retaining its full enzymatic activity. DPP-4 is also present in other fluids besides serum (Iwaki Egawa et al., 1998): in urine, seminal (Wilson et al., 1998) and amnion liquid (reviewed by Hildebrandt, 2004). Serum DPP-4 can also bind plasminogen and streptokinase (Gonzalez Gronow & Weber, 1998).

The enzyme DPP-4 preferentially cleaves peptides with a proline or alanine residue in the second last aminoterminal position (see above, Fig. 2) and removes X-Pro and X-Ala dipeptides from the N-terminal end of peptides and proteins which may have relevance in inhibiting degradation by unspecific proteases (Yaron & Naider, 1993; Mentlein et al., 1993; Kieffer et al., 1995; Deacon et al., 1995a, 1995b; Holst & Deacon, 1998; Kieffer & Habener, 1999; Deacon et al., 2000; Deacon, 2004). DPP-4 inhibitors act by binding to the enzyme pocket, leaving the incretins GLP-1 and GIP circulating in intact (active) form. The modification of a protein in the proline region (DPP-4 enzyme) has a general impact since protein influences the secondary structure of a protein and, therefore, its biological activity (reviewed by Hildebrandt, 2004).

Many substrates of DPP-4 have been identified (Table 8): gastrointestinal hormones, neuropeptides, cytokines, and chemokines possessing a proline in the right position are in vitro substrates of DPP-4 (Mentlein, 1999; De Meester et al., 2000, 2003).

The expression of DPP-4 may be increased by a proline diet (Suzuki et al., 1993) and by interferon- γ (Dang et al., 1991; Schmitz et al., 1996).

DPP-4 is not only responsible for inhibition of peptide degradation, but has also an effect on immunomodulation, cell adhesion and cell

Table 8

Substrates of DPP-4 (Mentlein, 1999; De Meester et al., 2000, 2003).

Substrates which may have clinical relevance	GLP-1, GLP-2, GIP, NPY, PYY, PACAP (pituitary acenylate cyclase-activating polypeptide), Substance P, RANTES (regulated on activation normal T cells expressed and secreted).
Substrates with questionable clinical relevance	GRH (growth hormone-releasing hormone), IGF-1, Prolactin, HCG (human chorionic gonadotropin), Bradykinin, Interleukin-18, Interleukin-2

movement. These effects are mediated by the rather identical functional protein, also called T-cell antigen CD26, but not by its enzymatic activity (Bednarczyk et al., 1991). It has an impact on the immune system with respect to T-lymphocyte memory cells activity and movement (Masuyama et al., 1992, 1999), differentiation of lymphocytes as a thymic maturation marker in differentiation of lymphocytes, and the anchoring of the ecto-adenosin deaminase in the lymphocyte membrane (involved in lowering extracellular adenosine) (Blanco et al., 1996; Iwaki Egawa et al., 1998).

The enzyme may be released from activated T-lymphocytes (Uematsu et al., 1996) and hepatomas of rats (Hanski et al., 1988). Cell adhesion (Hanski et al., 1985), and cell movement across the extracellular matrix (reviewed by Hildebrandt, 2004) were observed. This means that DPP-4 plays an important role in immune system, being a T-cell co-stimulator (Barnett, 2006), activating T-lymphocytes and cleaving numerous physiological peptides (Gorrell, 2005; Green et al., 2006). Note: all these additional effects may be relevant for unselectivity of DPP-4 inhibitory drugs as discussed later.

Clinically DPP-4 may also be linked to some immune diseases: low serum DPP-4 in patients with systemic lupus erythematoses (Stancikova et al., 1992), in synovial fluid of patients with rheumatoid arthritis (Kamori et al., 1991), and in Morbus Crohn patients (Th-1 cytokine profile) (Willheim et al., 1997). The clinical data are not conclusive since DPP-4 has contradictory effects, increasing and inhibiting immune responses (reviewed by Hildebrandt, 2004).

Interestingly, mice lacking the DPP-4 gene appear to be protected against obesity and insulin resistance (Conarello et al., 2003).

7.2. DPP-4 inhibitors

DPP-4 inhibitors increase the effects of above mentioned enzymes. The therapeutic focus is on increasing the activity of GLP-1 for type 2 diabetics. They are not incretin mimetics such as GLP-1, but incretin enhancers. DPP-4 inhibitors mainly act indirectly via incretin hormones which is evident from animal studies: their beneficial effect is lost in mice lacking the GLP-1 and GIP receptors (Hansotia et al., 2004) or lacking the DPP-4 enzyme (Mitani et al., 2002).

Small-molecule DPP-4 inhibitors have been developed that can be used orally. Typically, the inhibitors reduce serum DPP-4 activity to a high degree with some inhibition maintained for 24 h after one dose which allows once daily treatment (Åhren et al., 2004a, 2004b; Herman et al., 2005). There is evidence that plasma DPP-4 must be inhibited by rather 80% for an effectively high and sustained GLP-1 level increase since it has to be kept in mind that DPP-4 is also present in tissues with still potentially degrading activity for GLP-1.

Many effects different from incretin mimetics exist (Table 4).

7.2.1. Clinics

The DPP-4 inhibition is accompanied by a rise in postprandial levels of intact GLP-1 (Åhren et al., 2004a, 2004b; Herman et al., 2005; Nauck & El-Quaghli, 2005) and GIP. Though the total pool of the peptides is decreased during DPP-4 inhibition (El-Quaghli et al., 2007) (probably a negative feedback mechanism of GLP-1 on its secretion), the levels of active hormones are increased (Mari et al., 2005). DPP-4 inhibitors mimic the therapeutic effects of the incretin mimetics including stimulation of insulin secretion, inhibition of

glucagon secretion and possibly preservation of B-cell mass (Deacon, 2004) and inhibition of apoptosis (see above Fig. 5) (Drucker et al., 1987; Deacon 2004). Two compounds are in the market (Fig. 6). As shown for sitagliptin DPP-4 inhibitors improve HOMA- β , a marker of fasting insulin secretion, the proinsulin-to-insulin ratio, and the insulinogenic index. The moderate increase in GLP-1 by DPP-4 inhibition is probably not sufficient for a direct central effect on satiety. In contrast, DPP-4 inhibitors are so far not believed to alter gastric emptying (Vella et al., 2007) and are generally not associated with weight loss (reviewed by Aaboe et al., 2009) (Table 9). However, compared with sulfonylureas, thiazolidinediones, and insulin, DPP-4 inhibitors have the advantage that patients at least do not gain weight. Sitagliptin reduces weight by approx. 1.5 kg or is weight neutral whereas the sulfonylurea glipizide increases weight by approx. 1.1 kg.

Sitagliptin rather doubles the GLP-1 levels after a meal for a long time (Herman et al., 2005; Åhren et al., 2004a; Pratley et al., 2006; Rosenstock et al., 2006a; Garber et al., 2006).

Incretin enhancers do not pass the blood-brain barrier which may be the reason that there is rather no effect on satiety compared to GLP-1. The in vivo effect of exenatide is much greater than that of sitagliptin in rodents with respect to inhibition of food intake, amelioration of weight gain, and better inhibition of gastric emptying (Gedulin et al., 2007).

Sitagliptin and vildagliptin therapy in comparison with placebo resulted in an HbA_{1c} reduction of approximately 0.7%. There is no major difference in decreasing HbA_{1c} by either exenatide or the DPP-4 inhibitors sitagliptin and vildagliptin being 0.8 and 0.7% (Mikhail, 2008).

DPP-4 inhibitors are not associated with weight gain or oedema (except for combination with glitazones).

Clinical studies with both vildagliptin (Åhren et al., 2004a, 2004b; Åhren, 2006; Rosenstock et al., 2008b; Dejager et al., 2006; 2007; Pi-Sunyer et al., 2007) and sitagliptin (Scott et al., 2007; Aschner et al., 2006a; Charbonnel et al., 2006) have shown their efficacy in the treatment of type 2 diabetes. Compared with metformin (52 weeks) (Dejager et al., 2006;) and rosiglitazone (24 weeks) (Rosenstock et al., 2007b) in drug-naïve type 2 diabetic patients, treatment with vildagliptin as monotherapy showed slight inferiority and non-inferiority, respectively, with regards to lowering HbA_{1c}. DPP-4 inhibitors can be used as monotherapy (patients with type 2 diabetes uncontrolled by diet) and in combination (Mikhail, 2008). A combina-

tion of either sitagliptin or vildagliptin with metformin was recently approved; the HbA_{1c} is decreased by an extra of 0.7% compared to metformin alone. The addition of sitagliptin to glimepiride reduced the HbA_{1c} more effectively (0.89% versus 0.57%). As add-on therapy, both DPP-4 inhibitors induce additional lowering of HbA_{1c} with approximately 0.8–1.0% (Charbonnel et al., 2006; Bosi et al., 2007; Fonseca et al. 2007; Garber et al. 2007).

In general DPP-4 inhibitors seem to have positive effects on B-cell mass (Pospisilik et al., 2003; Mu et al., 2006).

DPP-4 inhibition seems to be well tolerated. Few adverse effects in humans have been demonstrated thus far in clinical trials (Raz et al., 2006; Rosenstock et al., 2006c; Åhren et al. 2004a). No characteristic pattern of adverse events has been associated with the use of vildagliptin (Åhren et al., 2004a; Åhren et al., 2004b) or other DPP-4 inhibitors (Scott et al., 2005; Hanefeld et al., 2005; Brazg et al., 2005).

There is practically no risk for hypoglycaemias (Nauck et al., 2007b; Aschner et al., 2006a; Charbonnel et al., 2006) which is due to the fact that the therapeutic effect of indirectly elevated GLP-1 is dependent on the glucose concentration (Nauck et al., 2007b) (for comparison e.g. glipizide (32%)). The risk of hypoglycaemic episodes during monotherapy with a DPP-4 inhibitor or a DPP-4 inhibitor in combination with metformin is negligible.

7.2.2. Sitagliptin (MK-0431)

Sitagliptin is the first compound of its class introduced in the market as a DPP-4 inhibitor. Its chemical structure is (2R)-4-Oxo-4[3-(trifluoromethyl)-5,6-dihydro[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl] 1-(2,4,5-trifluorophenyl)butan-2-amide (Fig. 6). It has an IC₅₀ of 18 nM.

It is approved as a monotherapy in the U.S. and as an add-on therapy to metformin or glitazone, when metformin plus diet are not sufficient (Aschner et al., 2006c; Charbonnel et al., 2006). In Europe a monotherapy with sitagliptin primarily for newly diagnosed diabetes or during therapeutic failure of other oral antidiabetics is not allowed; approved and effective are combinations of sitagliptin with metformin, insulin sensitizers (glitazones), and a sulfonylurea or in a triple combination with both metformin and a sulfonylurea, but not with a glinide (Karasik et al., 2006; Raz et al., 2006; Rosenstock et al., 2006b). The combination with metformin shows an improvement of the B-cell function (Xu & Williams-Herman, 2007; Migoya et al., 2007); probably different mechanisms are involved.

Table 9
Comparison of GLP-1 and DPP-4 inhibitors.

GLP-1 receptor agonists (GLP-1 mimetics)	DPP-4 inhibitor
Parenteral administration (s.c.), administration 60 min before morning and evening meal	Oral administration
Short acting*	Long acting
Strong effect	Effect less strong compared with incretin mimetics (except decrease in HbA _{1c})
Selective (one single target = GLP-1 receptor)	Nonselective (multiple targets: other enzymes, many DPP-4 substrates)
	Non-specific: GLP-2, NPY, SP, PACAP and others
Add-on therapy (metformin, sulfonylurea or to a combination of metformin and a thiazolidinedione)	Monotherapy or adjunctive therapy (combinations as shown left)
Increase in levels of exogenous GLP-1 (supraphysiologic, sustained)	Increase in endogenous GLP-1 concentrations (close to physiologic, retained diurnal pattern) and GIP
Stimulated insulin secretion	Stimulated insulin secretion
Inhibited glucagon secretion	Inhibited glucagon secretion
No change in fasting glucose	No change in fasting glucose
Marked reduction prandial glucose	Slight reduction in prandial glucose
Reduction in HbA _{1c} (~ -0.8%)	Reduction in HbA _{1c} (~ -0.7%)
Inhibition of gastric emptying	No effect on gastric emptying
May induce early satiety	No clear effect on satiety
Drug overdosage problematic	Drug overdose is non-toxic (except liver toxicity and QT prolongation for vildagliptin)
Potential of CNS side effects	No CNS side effects (no passing the blood-brain barrier)
Weight loss	Weight neutral (no effect)
Side effects well known (e.g. probably pancreatitis)	Side effects still to be evaluated (completely)
Increased hypoglycaemias, especially in combination with sulfonylureas	Low/no incidence of hypoglycaemias
Dose-related nausea	No nausea or vomiting
Not recommended in severe renal failure	Dose adjustment in severe renal failure (for only some compounds)

* Effect may be extended by pharmaceutical technology manipulations (once weekly injections).

The normal dose is 100 mg once a day (Raz et al., 2006; Aschner et al., 2006a, 2006b, 2006c; Kipnes, 2007; Karasik et al., 2006; Rosenstock et al., 2006b).

7.2.2.1. Pharmacokinetics (note: some data were obtained from a company). Sitagliptin is well absorbed and its oral bioavailability is 87%. The maximum plasma concentration after intake of 100 mg sitagliptin is reached after 1 to 4 hours. Binding to plasma proteins is low (38%), the V_d is 198 litres. 70–80% of a dose are renally excreted unchanged, and only a small amount is metabolized in which mainly CYP3A4 and CYP2C8 are involved. Its plasma half-life is between 8 and 14 hours.

The dose should not be changed when it is combined with other antidiabetic drugs (e.g. metformin or glitazones). In kidney malfunction or even renal failure the dose of sitagliptine has to be reduced one half or one fourth or sitagliptin should even no longer be used.

7.2.2.2. Side effects and contraindications. Medication with DPP-4 inhibitors appears to be reasonably safe. Side effects of sitagliptin are cold (running nose), stuffy nose and diarrhoea, also sore throat, headache and arthralgias were observed. Recently, postmarketing reports of anaphylaxis, angioedema, and rashes, including Stevens-Johnson syndrome, in sitagliptin-treated patients have emerged. A causal link to the drug, however, is not known in some cases.

Contraindications are rarely known except severe kidney malfunction.

7.2.2.3. Drug interactions. The combination of sitagliptin with pioglitazone results in peripheral oedema (4%). Glitazones probably posses a cardiotoxic effect (increased heart attack) and worsening of heart failure (Slørdal & Spigset, 2006). It is not sure whether the increase of cardiotoxic effects is only a result of this combination therapy.

Though CYP3A4 and CYP2C8 are partly involved in the degradation of sitagliptin, no drug interactions are expected except when this metabolizing way is extremely because of a kidney failure. In this case ketoconazole, Itraconazole, ritonavir, clarithromycin and possibly other CYP3A4 inhibitors may be relevant with respect to drug interactions.

Plasma digoxin is slightly increased by sitagliptin; an adaptation of doses of either drug is not necessary.

An interaction with any antidiabetic drugs has not been observed (Mistry et al., 2008).

7.2.3. Vildagliptin (LAF237)

Vildagliptin is the second DPP-4 inhibitor being approved in Europe. Sitagliptin and Vildagliptin are different in molecular structure (Fig. 7) and in pharmacokinetics. Sitagliptin is a competitive antagonist at the enzyme, and vildagliptin (as saxagliptin, see below) is a substrate for DPP-4 and inhibits the target (receptor or enzyme) molecule. Vildagliptin has a high affinity (Åhren et al., 2007; Villhauer et al., 2003; El-Ouaghli et al., 2003).




Fig. 7. Chemical structures of sitagliptin and vildagliptin.

7.2.3.1. Clinical aspects. Note: 100 mg doses of vildagliptin have been used in many studies, but only 50 mg are approved. Some above mentioned clinical results as well as some side effects (see below) have to be recognized under this aspect.

The in vitro actions have translated into significant effects in improving glycaemic control in type 2 diabetic patients, as illustrated by the clinically meaningful reductions in HbA_{1c} (by 0.8%) observed over 24 weeks (Rosenstock et al., 2007a) and 1 year (Schweizer et al., 2007) when a monotherapy of 100 mg/day is used. This was also observed using over 52 weeks a combination of vildagliptin 50 mg/day with metformin (Åhren et al., 2004a). Vildagliptin is used as a monotherapy or in combination with metformin (50 mg together with 850 mg metformin). Vildagliptin (as sitagliptin) may be used as monotherapy in patients who cannot tolerate metformin or sulfonylureas. Sitagliptin may be used as alternative to metformin in modest renal insufficiency. 100 mg vildagliptin can be combined with metformin or glitazone, but only 50 mg should be combined with a sulfonylurea (no higher effect when 100 mg are used).

Mechanistic studies in type 2 diabetics indicate that vildagliptin increases fasting and postprandial GLP-1 levels, improves B-cell sensitivity to glucose, A-cell functions, and insulin sensitivity, reduces postprandial lipaemia but exerts no clinically important effect on gastric emptying (Garber et al., 2006; Pratley et al., 2006; Rosenstock et al., 2006a). These effects lead to improved glucose tolerance and a reduction in fasting plasma glucose levels. At a dose of 100 mg once daily, fasting and postprandial glucose concentrations were reduced after 4 weeks of vildagliptin treatment and plasma glucagon concentrations were suppressed and the ratio of insulin to glucose was increased (Åhren et al., 2004b). An increased insulin secretory rate was observed with vildagliptin (Balas et al., 2007): calculation of the insulin secretory rate area under the curve (ISR) divided by the glucose AUC showed significantly improved insulin response at 0–30 min, 0–120 min and 0–840 min. Vildagliptin improves glucose-dependent pancreatic A-cell function in type 2 diabetes mellitus, as shown by reduced glucagon levels in the postmeal period (Åhren et al., 2004b; Azuma et al., 2008). The reduction in glucagon/insulin ratio is associated with reduced endogenous glucose production during both the postprandial period and the postabsorptive period (Balas et al., 2007). These results illustrate the dual effects of vildagliptin to improve glucose tolerance during meals and to directly reduce fasting hepatic glucose production. Clinical trials indicate that these mechanisms of action are associated with clinically significant improvements in glycaemic control as assessed by HbA_{1c} when vildagliptin is used alone or in combination, in patients with type 2 diabetes. Insulin sensitivity is significantly improved following meal ingestion for up to 52 weeks (Åhren et al., 2005). Such improvement could be an indirect effect of improved glycaemia, or related to glucagon reduction, but it may also be a direct effect shown in a hyperinsulinaemic euglycaemic clamp study in which vildagliptin improved insulin sensitivity under conditions where neither glucagon nor glucose toxicity could explain the improved insulin-mediated glucose disposal rate.

Vildagliptin has no effect on gastric emptying or rate of entry of ingested glucose into the systemic circulation (Vella et al., 2007) which is in contrast to GLP-1 (Naslund et al., 1999b; Kolterman et al., 2000; Meier et al., 2003a; Kolterman et al., 2005). DPP-4 inhibitors are generally not associated with a deceleration of gastric emptying, perhaps due to the modest increase of postprandial plasma levels of intact (biologically active) GLP-1 (doubled to 15–25 pmol/l). For vildagliptin there exists one opposing outcome demonstrating an inhibition of gastric emptying (Dardik et al., 2003).

Like sitagliptin vildagliptin therapy does not result in weight gain.

B-cell function is improved by vildagliptin treatment over 1 year in patients with type 2 diabetes, indicated by measuring C-peptide secretion after meal ingestion, (Åhren et al., 2005). Levels of proinsulin during meal ingestion are reduced by vildagliptin (Åhren et al., 2007); proinsulin is an additional marker of abnormal B-cell function.

Fasting lipolysis is decreased by vildagliptin (Azuma et al., 2008). Postprandial hypertriglyceridaemia is an important metabolic abnormality in many patients with type 2 diabetes mellitus; animal studies suggest that incretin enhancement may reduce intestinal triglyceride absorption (D'Alessio & Tso 2005). Postprandial triglyceride-rich lipoprotein levels were reduced (Matikainen et al., 2006); this happens with respect to total serum triglycerides and chylomicron triglycerides, reflecting reductions in chylomicron apolipoprotein B-48 and chylomicron cholesterol.

As already shown for GLP-1 alone a combination of GLP-1 with a DPP-4 inhibitor has positive effects on B-cell viability and proliferation in a model of B-cell regeneration (partially pancreatectomized rats) (Joanny et al., 2007).

Vildagliptin monotherapy, either 50 mg twice daily or 100 mg once daily, showed in preliminary reports of longer phase III sustained efficacy but non-inferiority compared with metformin after 1 year of therapy, (Dejager et al., 2006); it has to be emphasized that vildagliptin was better tolerated than metformin. Similarly, vildagliptin was as effective as rosiglitazone in direct comparison monotherapy study (Rosenstock et al., 2006a) and also produced significant reductions in HbA_{1c} when used in combination with metformin (Garber et al., 2006). Since metformin increases GLP-1 being a GLP-1 secretagogue and not a DPP-4 inhibitor (Sinha Roy et al., 2007), the clinical observation to combine metformin with a DPP-4 inhibitor is reasonable.

The DPP-4 inhibitor vildagliptin enhances the HbA_{1c} reduction when already induced by insulin in type 2 diabetic patients (Fonesca et al., 2007).

7.2.4. Side effects and contraindications

Hypoglycaemias are rare in response to vildagliptin combined with pioglitazone (0.3%) but were higher with pioglitazone being used alone (1.9%). Pioglitazone increased body weight (1.4 kg) which is further increased using a combination with 100 mg vildagliptin (2.7 kg).

Adverse effects of sitagliptin and vildagliptin were nasopharyngitis (6.4%), urinary tract infection (3.2%), and headache (5.1%), although placebo effects are close to these values (Mikhail, 2008). There are some hints on skin lesions.

The combination with pioglitazone increased the incidence of peripheral oedema.

Vildagliptin should not be used by patients with impaired kidney function or being dialysed or having a severe liver disease.

7.2.5. Drug interactions / combinations

A vildagliptin dose of 100 mg (approved are 50 mg in Germany) daily should be reduced to 50 mg when combined with other drugs such as sulfonylurea and can later be increased again but not to more than 100 mg. There is no interaction with food.

The metabolism of vildagliptin is not influenced by P450 enzymes so that drug interactions by this mechanism can not be expected.

7.3. Nonselectivity as a problem

In view of the widespread expression of DPP-4 on many cell types (see above), unselective effects have to be expected. Taking into account the large number of potential substrates for DPP-4 (Table 8) (Mentlein, 1999; De Meester et al., 2000) the aspect of nonselectivity is a major issue with respect to long-term safety. Since DPP-4 enzymes metabolise a wide variety of peptides, they potentially affect other regulatory systems and have pleiotropic additional effects (Hildebrandt et al., 2000; Durinx et al., 2000; Gorrell, 2005).

PACAP (pituitary adenyl cyclase-activating peptide) is one of the DPP-4 substrates (Mentlein et al., 1993), but the inhibition of PACAP degradation in porcine pancreas is not associated with an acute effect on endocrine secretion (Hjellund et al., 2007).

Other substrates of DPP-4 are trypsinogen (Erlanson Albertsson & Larsson, 1988; Heymann et al., 1986) and procolipase (which has

impact on enterostatin by which the lipid uptake is regulated) (Bouras et al., 1996). The inactivation of GH-RF (somatoliberin) and of other mitogens may be inhibited by DPP-4 inhibitors.

GLP-2 degradation will be inhibited by DPP-4 inhibitors which will increase its activity (Mentlein et al., 1993); it has effects on intestinal trophic activity (proliferation), suppresses parathormone, decreases bone resorption and increases gut motility.

The inhibition of NPY (neuropeptide Y) degradation has an impact on its orexigenic effect since intact NPY stimulates food intake and feeding motivation (Karl et al., 2003b). The not degraded NPY is one of the most orexigenic peptides (Bray, 1993; Chance & Fischer, 1993; Karydis & Tolis, 1998), and by truncating this peptide its receptor preference is modified (Grandt et al., 1993). It would be expected that diminishing NPY interaction with NPY1 receptors will not only modify hunger and satiety, but also the GI motility (Chen et al., 1997). It is also involved in modifications of behavioural tests; DPP-4 deficient rats exhibit increased pain indicative of a reduced stress-induced analgesia (Karl et al., 2003a). Psychoneuroendocrine effects may perhaps be underestimated by not concerning the less degradation of NPY (also substance P) (Myers, 1994; Hildebrandt, 2004). Rats lacking DPP-4 like activity have a reduced behavioural stress response which may be linked to NPY processing. Cleavage of NPY by DPP-4 results in NPY (3-36) which lacks affinity for the NPY-1 receptor mediating the anxiolytic-like effects of NPY. The nonspecific effect of DPP-4 inhibitors on NPY degradation (truncated from NPY(1-36) (original) to NPY(3-36)) may influence the antilipolytic effect (Kos et al., 2007).

Possibly the impact of an inhibition of above mentioned compounds degradation is underestimated.

Long-term safety based on CD26 effects remains unknown, but to date no significant alterations of immune function have been observed (Richter et al., 2008a) except a nonspecific increase of infections after sitagliptin (Richter et al., 2008b).

Non-selectivity with respect to actions on the related enzymes DPP-8, DPP-9 or both, could be of particular importance (Lankas et al., 2005). The relevance of DPP-8/9 inhibition is not clear yet, although skin lesions observed in primates have to be reevaluated, the FDA says. DPP-8 and DPP-9 are active in hematologic and immune cells (Gorrell, 2005). DPP-4 inhibitors do not substantially inhibit cell proliferation in experiments with human lymphocytes in vitro (Lankas et al., 2005). The toxicity of DPP-8/DPP-9 inhibitor that were reported as 100% mortality in mice, alopecia, thrombocytopenia, reticulocytopenia, enlarged lymph nodes, splenomegaly and 20% mortality in rats was not confirmed by others (Burkey et al., 2008).

7.4. Comparison of properties of GLP-1 receptor agonists and DPP-4 inhibitors (Table 9)

Exenatide (GLP-1 receptor agonist) is an incretin mimetic, while sitagliptin and vildagliptin (DPP-4 inhibitors) are incretin enhancers. No difference is observed with respect to HbA_{1c} values: exenatide is associated with reduction by approximately 0.8% that with either sitagliptin or vildagliptin is 0.7%; no difference was observed with respect to hypoglycaemias.

Exenatide treatment leads to a mild weight loss of approximately 2 kg after 30 weeks, whereas sitagliptin and vildagliptin have generally a neutral effect on weight. Weight loss is a common outcome of therapy with native GLP-1 (Zander et al., 2002), exenatide (DeFronzo et al., 2005; Buse et al., 2004; Kendall et al., 2005), and liraglutide (Nauck et al., 2006; Hansen et al., 2001), whereas treatment with DPP-4 inhibitors is associated with prevention of weight gain (Ahren et al., 2004a; 2004b; Scott et al., 2005; Hanefeld et al., 2005).

GLP-1 mimetics markedly inhibit gastric emptying whereas DPP-4 inhibitors have no such effect. This is of major importance for the glycaemic effect of the two strategies because a main mechanism of GLP-1 mimetics is to reduce prandial glucose, which is achieved mainly by reduction in gastric emptying, whereas DPP-IV inhibitors

reduced the overall 24-hour glucose with no marked difference in reduction in fasting versus prandial glycaemia.

GLP-1 mimetics heavily activate GLP-1 receptors whereas DPP-4 inhibitors moderately increase endogenous GLP-1 levels, which will retain the diurnal pattern and the primary portal action (Vahl et al., 2007) of the incretin. The difference of effects of incretin mimetics and DPP-4 inhibitors might be partly because of the relative low increase in GLP-1 levels by the enzyme inhibitors. This is explained in part by the relatively modest stabilisation of postprandial GLP-1 seen after DPP-4 inhibition, compared with the high increases in circulating levels of GLP-1 receptor agonists exemplified by exenatide. Gastro-intestinal side-effects, predominantly nausea, are often reported after treatment with injectable GLP-1 receptor agonists but have not been described for DPP-4 inhibitors (Åhren et al., 2004a, 2004b; Demuth et al., 2005; Scott et al., 2005; Hanefeld et al., 2005; Brazg et al., 2005). Although nausea is a common side-effect of in exenatide therapy, many patients lose weight independently of nausea. The lack of a weight loss by DPP-4 inhibitors obese patients compared to lean subjects probably due the low activity of DPP-4 in fat cells of obese patients (Kos et al., 2007).

When exenatide is compared with sitagliptin, the effectiveness of exenatide is higher with respect to inhibition of food uptake and in inhibiting gastric emptying (Herrmann et al., 2008) which may be linked to weight reduction (mentioned above). Twice daily exenatide through subcutaneous injection is indicated for the treatment of patients with type 2 diabetes mellitus in whom one or more oral agents do not work, often as an alternative to insulin treatment. By contrast, once daily DPP-4 inhibitors could be used alone or as add-on therapy to patients failing one or more oral agents.

In contrast to incretin mimetics, DPP-4 inhibitors do not exclusively act via pharmacological concentrations of GLP-1 like activity, but raise the concentration of other peptide hormones possibly involved in metabolic control.

Differences between GLP-1 agonists and DPP-4 inhibitors are summarized in Table 9.

7.5. DPP-4 inhibitors under investigation

In addition to the mentioned two drugs being in the market, more are going to be developed as DPP-4 inhibitors. Concerning the chemical differences of distinct DPP-4 inhibitors it seems likely that one or more of these agents could be associated with unexpected adverse effects. Some characteristics of new compounds are summarized in Table 10.

The dose of saxagliptin (BMS-477118; active metabolite; developed by Bristol and Astra Zeneca) has to be changed due to kidney functioning (Augeri et al., 2005; List et al., 2007). Denagliptin (developed by Glaxo Smith Kline, albeit no longer clinically investigated for being approved) (Demuth et al., 2005) show significant differences in their pharmacokinetic profiles, side-effects, or clinical activity which are not investigated in detail.

Table 10
Comparison of some DPP-4 inhibitors.

	Sitagliptin	Vildagliptin	Alogliptin	Saxagliptin	BI 1356 ^b
Enzyme inhibition in vivo after 24 hours (ED ₅₀ as mg/kg)	>30	14	10	2.7	0.9
IC ₅₀ (nM)	19	62	24	50	1
Dissociation velocity from enzyme		$2.1 \times 10^{-4} \text{ sec}^{-1}$			$3.0 \times 10^{-5} \text{ sec}^{-1}$
Selectivity versus DPP-8 and DPP-9		Approx. 300 and 30 fold, respectively ^a			10000 fold

^a Burkey et al., 2008.

^b Thomas et al., 2007, 2008b.

Alogliptin (SYR-322; developed by Takeda) is a novel quinazolinone-based dipeptidyl peptidase-4 (DPP-4) inhibitor (IC₅₀ ~ 6.9 nM) and exhibited a >10,000-fold selectivity for DPP-4 over the closely related serine proteases DPP-2, DPP-8, DPP-9, fibroblast activation protein/seprase, prolyl endopeptidase, and trypsin (IC₅₀>100,000 nM); the absolute oral bioavailability of alogliptin is between 45% and 88% (Lee et al., 2008). Alogliptin has a kinetic profile which may allow a once-a-day dosing regime (Takeuchi et al., 2007; Christopher et al., 2007). Alogliptin improves the glycemic control (Fleck et al., 2008).

P32/98 (Isoleucine thiazolidide) increased B-cell mass in animal studies (Pospisilik et al., 2003).

BI 1356 ((R)-8-(3-amino-piperidin-1-yl)-7-but-2-ynyl-3-methyl-1-(4-methyl-quinazolin-2-ylmethyl)-3,7-dihydro-purine-2,6-dione; developed by Boehringer-Ingelheim, proposed tradename Oндero®) differs from those in the market (Thomas et al., 2008a, 2008b) since it is a xanthine derivative, has higher in vitro potency (IC₅₀) combined with a stronger in vivo effect, has a lower dissociation velocity from the enzyme DPP-4, which means it binds merely to the target at rather low plasma levels in humans. BI 1356 has a longer duration of action in vivo so that a once a day regimen is possible (Thomas et al., 2007). BI 1356 increases the basal GLP-1 level, but not the plasma insulin levels in rats after 24 hours. Under specific conditions it may (yet not proved) improve a regeneration of B-cells (see above for GLP-1). BI 1356 induces a more than 80% enzyme inhibition at plasma concentration of 5.3 nM (Dugi et al., 2007). It is not renally excreted and, therefore, a modification of the dose in kidney failure may not be necessary.

BI 1356 competitively inhibits DPP-4 activity in vitro with an IC₅₀ of approximately 1 nM (Thomas et al., 2008a, 2008b), compared with sitagliptin (19 nM), alogliptin (24 nM), saxagliptin (50 nM), and vildagliptin (62 nM). The calculated k(off) rate for BI 1356 was $3.0 \times 10^{-5}/\text{s}$ (versus $2.1 \times 10^{-4}/\text{s}$ for vildagliptin). BI 1356 is $\geq 10,000$ -fold more selective for DPP-4 than DPP-8, DPP-9, aminopeptidases N and P, prolyloligopeptidase, trypsin, plasmin, and thrombin and is 90-fold more selective than for fibroblast activation protein in vitro (Thomas et al., 2008a, 2008b).

PHX1149 (developed by Phenomix Corporation) is water-soluble, not metabolized, excreted renally, and has a half-life of 10 to 13 hours (Guler, 2007). The SK-0405 characteristics are an IC₅₀ of 3.3 nM, high selectivity (17000 fold selectivity over DPP8, DPP9, FAP = fibroblast activation protein), and a longer lasting inhibition compared to vildagliptin (Yasuda et al., 2007).

There are even many more DPP-4 inhibitors including P93/01, NVP-DPP728, 815541A, GSK23A and valine-pyrrolidide.

Up to 50% of GLP-1 entering the circulation may be degraded by NEP-24.11; therefore combined inhibition of DPP-4 and NEP-24.11 is superior to DPP-4 inhibition alone in preserving intact GLP-1, which potential therapeutic effect (Plamboeck et al., 2005).

7.6. Additional remarks and general comment with respect to DPP-4 inhibitors

Some antidiabetic compounds (e.g. pioglitazone), which are in the market for a long time, partly may have DPP-4 inhibitory effects (Lenhard et al., 2004), also atorvastatin (Taldone et al., 2006). In addition to the well known direct insulinotropic effect of nateglinide via KATP channels, also an inhibition of GLP-1 degradation may have some minor relevance (McKillop et al., 2007). Metformin may inhibit DPP-4 activity (Lindsay et al., 2005) which could be valuable for a combination with incretin hormones. A confirmation of these data by other laboratories is awaited.

Generally speaking, the DPP-4 inhibitors are modestly effective glucose-lowering drugs although HbA_{1c} values lowering effect is sufficient and it has to be admitted that mainly patients with not extremely high HbA_{1c} values were included in some trials and could not show up with higher effects. This new class of antidiabetic agents

(incretin enhancers) expand B-cell mass in preclinical studies. However, long-term clinical studies are needed to determine the benefits of targeting the incretin axis for the treatment of type 2 diabetes.

A flat dose–response curve for DPP-4 inhibition was observed in trials of vildagliptin and sitagliptin and is sufficient. On the other hand an inhibition of at least 70–80% is necessary.

So far, no study reported on patient-oriented parameters like mortality, diabetic complications, costs of treatment and health-related quality of life. When compared to placebo treatment sitagliptin and vildagliptin improved metabolic control. Comparison with other already established blood-glucose lowering drugs did not reveal major advantages of DPP-4 treatment. DPP-4 inhibitors like sitagliptin and vildagliptin probably have advantages over existing therapies with oral antidiabetic compounds (Table 5), e.g. with respect to absence of hypoglycaemia, weight gain etc. Long-term data on cardiovascular outcomes and safety are urgently needed before widespread use of these new agents. More information on the benefit-risk ratio of DPP-4 inhibitor treatment is necessary especially analysing adverse effects on parameters of immune function although not the enzymatic component of the enzyme (DPP-4, CD 26) is responsible for this effect. Also, long-term data are needed investigating patient-oriented parameters like health-related quality of life, diabetic complications and all-cause mortality.

7.7. Developments to be expected in general with respect to incretins and future perspectives

Much effort continues to be directed towards improvement of the pharmacokinetic profile of GLP-1 receptor agonists, to minimise peak levels of the drug and thus reduce the extent of nausea. Longer-acting GLP-1 receptor agonists should ideally provide more uniform and sustained GLP-1 receptor activation over a 24-h period while requiring less frequent administration (as mentioned before). Furthermore, there is great interest in determining whether chronic therapy with GLP-1 receptor agonists will be associated with sustained long-term control of HbA_{1c} and improvement in B-cell function beyond that achievable with existing agents. Similar questions pertain to the DPP-4 inhibitors, which also (indirectly) target B-cells.

Several issues remain to be elucidated. The durability of the weight loss induced by GLP-1 receptor agonists needs to be established, and even though both GLP-1 receptor agonists and DPP-4 inhibitors improve B cell function, the ability of either treatment to preserve functional B cell mass in humans needs clinical confirmation. The optimal role on B-cell mass has to be established. However, it is important to note that new drugs that act by blocking the action of DPP-4 increase levels of GIP as well as GLP-1. Therefore, manipulations that increase islet mass in rodents by reducing DPP-4 action are potentially due to actions of both incretins (Pospisilik et al., 2002; Conarello et al., 2003).

This leaves no direct means for testing direct effects of incretin based drugs on B-cell mass in diabetic patients. Given this limitation, the best alternative currently available to gain insights into the effects of GLP-1 receptor agonists and DPP-4 inhibitors on the course of B-cell function in diabetes may be careful natural history studies. An excellent example of this is the recently published ADOPT study (Kahn et al., 2006). In this trial, patients were followed for 4 years to the relative rate of decline in glycemic control. The reflected data worsened glycemic control on stable therapy, which is thought to be at least partially the result of loss of islet mass. Therefore the first step in ascertaining whether incretin based drugs can affect islet mass in humans is likely to come from clinical trials such as ADOPT. Comparison of the course of diabetes in patients treated with GLP-1 receptor agonists or DPP-4 inhibitors, using durability of effect on glycemic control as the primary outcome, would provide some insight into whether these agents have chronic effects on B-cell health. While such studies cannot prove that the effects of incretin-based stimula-

tion of islet mass in animals also pertain in humans, they would go a long way towards establishing the optimal role for these drugs in clinical medicine.

In patients with newly diagnosed type 1 diabetes in remission phase, treatment with intravenous infusions of GLP-1 without concomitant insulin injection displays beneficial effects on postprandial glucose excursions and glucagon suppression, and in patients with long lasting type 1 diabetes and no residual insulin secretory capacity, administration of exenatide together with insulin has similar effects (Behme et al., 2003; Dupré et al., 1995, 2004). But, further investigations are needed to fully evaluate the ability of GLP-1 to prolong the remission period, to reduce the insulin requirements and to evaluate the overall efficacy of incretin based treatments in the field of type 1 diabetes.

The observation that GLP-1 receptor agonists improve myocardial function in human patients after myocardial infarction and reduce blood pressure highlights the need for studies with cardiovascular end-points, especially as short-term studies with GLP-1 receptor agonists show promising results. Because patients with type 2 diabetes have increased risks of cardiovascular morbidity and mortality, the observation that GLP-1 receptor agonists improve myocardial function in human patients after myocardial infarction (Nikolaidis et al., 2004) even more strengthens the need for studies that assess cardiovascular endpoints in patients treated with DPP-4 inhibitors or GLP-1 receptor agonists, which will help to get a greater understanding of the true benefits and role of these drugs for the treatment of diabetes mellitus.

Interestingly a non-peptide GLP-1 receptor agonist was developed which is also an allosteric modulator of GLP-1 binding (6,7-dichloro-2-methylsulfonyl-3-N-tert-butylaminoquinoxaline).

Besides using GLP-1 analogues and inhibitors of GLP-1 degradation, a third strategy aims at enhancing the endogenous GLP-1 secretion by using α -glucosidase inhibitors (controversy about this possible effect as mentioned above under 3.1. These drugs induce a lasting improvement in the GLP-1 response to oral sucrose in both patients with type 2 diabetes (Seifarth et al., 1998) and in healthy volunteers. They result in a more than 90% increase in the GLP-1 concentration after a meal (Göke et al., 1995a). However, a clinical relevance (similar effect on GLP-1 secretion) in patients with type 2 diabetes has not been demonstrated yet (Hücking et al., 2005). Inhibition of intestinal α -glucosidase has been shown to augment secretion of both GIP and GLP-1. The biological consequences of GLP-1 enhancement by α -glucosidase inhibition are also not characterised beyond their ability to minimise postprandial glucose fluctuations and delay in gastric emptying.

Suppression of circulating NEFA by inhibition of adipose tissue hormone sensitive lipase may augment circulating GLP-1 concentrations (Ranganath et al., 1999a).

7.8. Overall clinical aspects

It is currently not clear at which point in the natural history of type 2 diabetes incretin therapy should be initiated and more research is needed to address this issue. Logic dictates that since incretin failure may occur early and can address many pathophysiological features of type 2 diabetes that lead to progression despite use of existing treatments, incretin therapy should be used early. However, most current data on incretin therapy are to be found in combination with existing treatments often when type 2 diabetes is well advanced. The optimal agent(s) that may mimic and replace the endogenous incretin effect is not fully known and awaits the outcome of ongoing clinical trials. Important issues with regard to practicalities such as oral or parenteral route of administration, long term safety and efficacy may determine which of the proposed options to increase the incretin effect is preferred in clinical practice.

In view of the above limitations, metformin will be the drug of choice for initial treatment of type 2 diabetes.

Incretin mimetics and enhancers may be established as first-line treatment; however, the exact place in therapy remains to be explored.

DPP-4 inhibitors could eventually be used in prediabetic stages and in early stages of diabetes in order to prevent the progression of type 2 diabetes (prophylactic use).

There is no conflict of interest with respect to any comment or interpretation of data in this review.

The help of critically reading the manuscript by Dr. M. Mark, Biberach, Germany, is greatly acknowledged.

References

- Aaboe, K., Knop, F. K., Vilsbøl, T., et al. (2007). Acute administration of a sulphonylurea compound restores the insulinotropic effect of glucose-dependent insulinotropic polypeptide in patients with type-2 diabetes. *Diabetologia* 50, S248–S249.
- Aaboe, K., Krarup, T., Madsbad, S., & Holst, J. J. (2009). GLP-1: physiological effects and potential therapeutic applications. *Diabetes Obes Metab* 9, 603–608.
- Abraham, E. J., Leech, C. A., Lin, J. C., Zulewski, H., & Habener, J. F. (2002). Insulinotropic hormone glucagon-like peptide-1 differentiation of human pancreatic islet-derived progenitor cells into insulin-producing cells. *Endocrinology* 143, 3152–3161.
- Agerso, H., Jensen, L. B., Elbrønd, B., et al. (2002). The pharmacokinetics, pharmacodynamics, safety and tolerability of NN2211, a new long-acting GLP-1 derivative, in healthy men. *Diabetologia* 45, 195–202.
- Ahmad, S. R., & Swann, J. (2008). Exenatide and rare adverse events. *New Engl J Med*, 358, 1970–1971.
- Åhren, B. (2006). Vildagliptin: an inhibitor of dipeptidyl peptidase-4 with antidiabetic properties. *Expert Opin Investig Drugs* 15, 431–442.
- Åhren, B. (2007). GLP-1 – based therapy of type 2 diabetes: GLP-1 mimetics and DPP-IV inhibitors. *Current Diabetes Reports* 7, 340–347.
- Åhren, B., & Larsson, H. (2001). Impaired glucose tolerance (IGT) is associated with reduced insulin-induced suppression of glucagon concentrations. *Diabetologia* 44, 1998–2003.
- Åhren, B., Gomis, R., Standl, E., Mills, D., & Schweizer, A. (2004a). Twelve- and 52-week efficacy of the dipeptidyl peptidase IV inhibitor LAF237 in metformin-treated patients with type 2 diabetes. *Diabetes Care* 27, 2874–2880.
- Åhren, B., Landin-Olsson, M., Jansson, P. A., Svensson, M., Holmes, D., & Schweizer, A. (2004b). Inhibition of dipeptidyl peptidase-4 reduces glycemia, sustains insulin levels, and reduces glucagon levels in type 2 diabetes. *J Clin Endocrinol Metab* 89, 2078–2084.
- Åhren, B., Pacini, G., Foley, J. E., & Schweizer, A. (2005). Improved meal-related b-cell function and insulin sensitivity by the dipeptidyl peptidase-IV inhibitor vildagliptin in metformin-treated patients with type 2 diabetes over 1 year. *Diabetes Care* 28, 1936–1940.
- Åhren, B., Pacini, G., Tura, A., Foley, J. E., & Schweizer, A. (2007). Improved meal related insulin processing contributes to the enhancement of B cell function by the DPP 4 inhibitor vildagliptin in patients with type 2 diabetes. *Horm Metab Res* 39, 826–829.
- Amiranoff, B., Vauclin-Jacques, N., & Laburthe, M. (1984). Functional GIP receptors in a hamster pancreatic cell line, In 11: specific binding and biological effects. *Biochem Biophys Res Commun* 17, 671–676.
- Andrews, C. N., Bharucha, A. E., Camilleri, M., Low, P. A., Seide, B., Burton, D., et al. (2007). Nitrogenic contribution to gastric relaxation induced by glucagon-like peptide-1 (GLP-1) in healthy adults. *Am J Physiol Gastrointest Liver Physiol* 292, G1359–G1365.
- Arbit, E., Poller, B., Ganzoni, C., Gass, S., Gomez-Orellana, I., Drewe, J., et al. (2007). Pharmacokinetics and pharmacodynamic effects of oral GLP-1 and PYY3-36: a proof of concept study in healthy subjects. *Diabetologia* 50, Abstract # 112.
- Arnés, L., Sacho, V., Valverde, I., & Villanueva-Penarcarillo, M. L. (2008). Action of exendin-4 treatment upon glucose transport in liver and muscle, in normal and type 2 diabetic state. *Diabetologia* 51(Suppl.), S361.
- Aschner, P., et al. (2006a). Effect of the dipeptidyl peptidase-4 inhibitor sitagliptin as monotherapy on glycemic control in patients with type 2 diabetes. *Diabetes Care* 29, 2632–2637.
- Aschner, P., Kipnes, M., Lunceford, J., Mickel, C., Davies, M., & Williams-Herman, D. (2006b). Sitagliptin monotherapy improved glycemic control in the fasting and postprandial states and beta-cell function after 24 weeks in patients with type 2 diabetes (T2DM). *Diabetes* 55(suppl 1), 462 (abstr 1995-PO).
- Aschner, P., Kipnes, M. S., Lunceford, J. K., Sanchez, M., Mickel, C., & Williams-Herman, D. E. (2006c). Effect of the dipeptidyl peptidase-4 inhibitor sitagliptin as monotherapy on glycemic control in patients with type 2 diabetes. *Diabetes Care* 29, 2632–2637.
- Aston-Mourme, K., Proietto, J., & Morahan, G. (2008). Too much of a good thing: why it is bad to stimulate the beta cell to secrete insulin. *Diabetologia* 51, 540–545.
- Augeri, D. J., Robl, J. A., Betebeuner, D. A., et al. (2005). Discovery and preclinical profile of saxagliptin (BMS-477118): a highly potent, long-acting, orally active dipeptidyl peptidase IV inhibitor for the treatment of type 2 diabetes. *J Med Chem* 48, 5025–5037.
- Avogaro, A. (2008). Glucagon-like peptide-1: pleiotropic effects on the cardiovascular system. *G Ital Cardiol (Rome)* 9, 753–758.
- Azuma, K., Rádiková, Z., Mancino, J., et al. (2008). Measurements of islet function and glucose metabolism with the DPP-4 inhibitor vildagliptin in patients with type 2 diabetes. *J Clin Endocrinol Metab* 93, 459–464.
- Baggio, L. L., & Drucker, D. J. (2006). Therapeutic approaches to preserve islet mass in type 2 diabetes. *Annual Rev Med* 57, 265–281.
- Baggio, L. L., Huang, Q. L., Brown, T. J., & Drucker, D. J. (2004). A recombinant human glucagon-like peptide (GLP)-1-albumin protein (albugon) mimics peptidergic activation of GLP-1 receptor-dependent pathways coupled with satiety, gastrointestinal motility, and glucose homeostasis. *Diabetes* 53, 2492–2500.
- Baggio, L. L., Huang, Q. L., & Drucker, D. J. (2006). The long-acting albumin-exendin-4 GLP-1R agonist CJC-1134 engages central and peripheral mechanisms regulating glucose homeostasis. *Diabetes* 55, A85.
- Balas, B., Baig, M. R., Watson, C., et al. (2007). The dipeptidyl peptidase IV inhibitor vildagliptin suppresses endogenous glucose production and enhances islet function after single-dose administration in type 2 diabetic patients. *J Clin Endocrinol Metab* 92, 1249–1255.
- Barnett, A. (2006). DPP-4 inhibitors and their potential role in the management of type 2 diabetes. *Int J Clin Pract* 60, 1454–1470.
- Barnett, A. H., Burger, J., Johns, D., Brodows, R., Kendall, D. M., Roberts, A., et al. (2007). Tolerability and efficacy of exenatide and titrated insulin glargine in adult patients with type 2 diabetes previously uncontrolled with metformin or a sulphonylurea: a multinational, randomized, open-label, two-period crossover noninferiority trial. *Clin Ther* 29, 2333–2348.
- Baron, A. D., Schaeffer, L., Shragg, P., & Kolterman, O. G. (1987). Role of hyperglucagonemia in maintenance of increased rates of hepatic glucose output in type II diabetics. *Diabetes* 36, 274–283.
- Barragán, J. M., Eng, J., Rodríguez, R., & Blázquez, E. (1999). Neural contribution to the effect of glucagon-like peptide-1-(7–36) amide on arterial blood pressure in rats. *Am J Physiol* 277, E784–E791.
- Barragán, J. M., Rodríguez, R. E., Eng, J., & Blázquez, E. (1996). Interactions of exendin-(9–39) with the effects of glucagon-like peptide-1-(7–36) amide and of exendin-4 on arterial blood pressure and heart rate in rats. *Regul Pept* 67, 63–68.
- Barth, A., & Schulz, H. (1974). Untersuchungen und Charakterisierung der Dipeptidylpeptidase IV. *Acta biologica et medica germanica* 32, 157–174.
- Bednarczyk, J. L., Carroll, S. M., Marin, C., & McIntyre, B. W. (1991). Triggering of the proteinase dipeptidyl peptidase IV (CD26) amplifies human T lymphocyte proliferation. *J Cell Biochem* 46, 206–218.
- Behme, M. T., Dupre, J., & McDonald, T. J. (2003). Glucagon-like peptide 1 improved glycemic control in type 1 diabetes. *BMC Endocr Disord* 3, 3.
- Blanco, J., Marie, I., Callebaut, C., Jacotot, E., Krust, B., & Hovanessian, A. G. (1996). Specific binding of adenosine deaminase but not HIV-1 transactivator protein Tat to human CD26. *Exp. Cell Res* 225, 102–111.
- Blase, E., Deng, W., Walsh, B., Fineman, M., & Rhodes, C. (2008). Intranasal administration of exenatide in patients with type 2 diabetes: pharmacokinetics, pharmacodynamics, safety and tolerability. *Diabetologia* 51(Suppl.), S346.
- Blase, E., Taylor, K., Gao, H., et al. (2005). Pharmacokinetics of an oral drug (acetaminophen) administered at various times in relation to subcutaneous injection of exenatide (exendin-4) in healthy subjects. *J Clin Pharmacol* 45, 570–577.
- Blonde, L., Klein, E. J., Han, J., Zhang, B., Mac, S. M., Poon, T. H., et al. (2006). Interim analysis of the effects of exenatide treatment on A1C, weight and cardiovascular risk factors over 82 weeks in 314 overweight patients with type 2 diabetes. *Diabetes Obes Metab* 8, 436–447.
- Bollag, R. J., Zhong, Q., Phillips, P., Min, L., Zhong, L., Cameron, R., et al. (2000). Osteoblast-derived cells express functional glucocorticoid-dependent insulinotropic peptide receptors. *Endocrinology* 141, 1228–1235.
- Bonner-Weir, S. (2000). Islet growth and development in the adult. *J Mol Endocrinol* 24, 297–302.
- Bose, A. K., Mocanu, M. M., Carr, R. D., Brand, C. L., & Yellon, D. M. (2005). Glucagon-like peptide 1 can directly protect the heart against ischemia/reperfusion injury. *Diabetes* 54, 146–151.
- Bosi, E., Camisasca, R. P., Colloher, C., Rochotte, E., & Garber, A. J. (2007). Effects of vildagliptin on glucose control over 24 weeks in patients with type 2 diabetes inadequately controlled with metformin. *Diabetes Care* 30, 890–895.
- Bouras, M., Huneau, J. F., & Tome, D. (1996). The inhibition of intestinal dipeptidylaminopeptidase-IV promotes the absorption of enterostatin and des-arginine-enterostatin across rat jejunum in vitro. *Life Sci* 59, 2147–2155.
- Bray, G. A. (1993). The nutrient balance hypothesis: peptides, sympathetic activity, and food intake. *Ann N Y Acad Sci* 676, 223–241.
- Brazg, R., Thomas, K., Zhao, P., Xu, L., Chen, X., & Stein, P. (2005). Effect of adding MK-0431 to ongoing metformin therapy in type 2 diabetic patients who have inadequate glycemic control on metformin. *Diabetes* 54(suppl 1), 3 (Abstract).
- Bregenholt, S., Moldrup, A., Blume, N., Karlens, A. E., Nissen Friedrichsen, B., Tornhave, D., et al. (2005). The long-acting glucagon-like peptide-1 analogue, liraglutide, inhibits beta-cell apoptosis in vitro. *Biochem Biophys Res Commun* 330, 577–584.
- Bregenholt, S., Moldrup, A., Knudsen, L. B., & Petersen, J. S. (2001). The GLP-1 derivative NN2211 inhibits cytokine-induced apoptosis in primary rat β -cells. *Diabetes* 50(suppl 1), 31 (Abstract).
- Brubaker, P. L., & Drucker, D. J. (2004). Minireview: Glucagon-like peptides regulate cell proliferation and apoptosis in the pancreas, gut and central nervous system. *Endocrinology* 145, 2653–2659.
- Bullock, B. P., Heller, R. S., & Habener, J. F. (1996). Tissue distribution of messenger ribonucleic acid encoding the rat glucagon-like peptide-1 receptor. *Endocrinology* 137, 2968–2978.
- Bulotta, A., Hui, H., Anastasi, E., Bertolotto, C., Boros, L. G., Di Mario, U., et al. (2002). Cultured pancreatic ductal cells undergo cell cycle re-distribution and beta-cell-like differentiation in response to glucagon-like peptide-1. *J Mol Endocrinol* 29, 347–360.
- Bunck, M. C., Diamant, M., Corner, A., Eliasson, B., Malloy, J. L., Shaginian, R. M., et al. (2007). One year treatment with exenatide improves β cell function and glycaemic control in metformin treated type 2 diabetes patients. *Diabetologia* 50, (Suppl.) Abstract # 251.

- Burcelin, R., Da Costa, A., Drucker, D., & Thorens, B. (2001). Glucose competence of the hepatoportal vein sensor requires the presence of an activated glucagon-like peptide-1 receptor. *Diabetes* 50, 1720–1728.
- Burkey, B. F., Hoffmann, P. K., Hassiepen, U., Trappe, J., Juedes, M., & Foley, J. E. (2008). Adverse effects of dipeptidyl peptidases 8 and 9 inhibition in rodents revisited. *Diabetes Obes Metab* 10, 1057–1061.
- Buse, J. B., Henry, R. R., Han, J., Kim, D. D., Fineman, M. S., & Baron, A. D. (2004). Effects of exenatide (exendin-4) on glycemic control over 30 weeks in sulfonylurea-treated patients with type 2 diabetes. *Diabetes Care* 27, 2628–2635.
- Buse, J. B., Klonoff, D. C., Nielsen, L. L., Guan, X., Bowlus, C. L., Holcombe, J. H., et al. (2007). Metabolic effects of two years of exenatide treatment on diabetes, obesity, and hepatic biomarkers in patients with type 2 diabetes. *Clin Ther* 29, 139–153.
- Buteau, J., El-Assaad, W., Rhodes, C. J., Rosenberg, L., Joly, E., & Prentki, M. (2004). Glucagon-like peptide-1 prevents beta cell glucolipotoxicity. *Diabetologia* 47, 806–815.
- Buteau, J., Foisy, S., Joly, E., & Prentki, M. (2003). Glucagon-like peptide 1 induces pancreatic betacell proliferation via transactivation of the epidermal growth factor receptor. *Diabetes* 52, 124–132.
- Buteau, J., Foisy, S., Rhodes, C. J., Carpenter, L., Biden, T. J., & Prentki, M. (2001). Protein kinase C ζ activation mediates glucagon-like peptide-1-induced pancreatic betacell proliferation. *Diabetes* 50, 2237–2243.
- Buteau, J., Spatz, M. L., & Accili, D. (2006). Transcription factor FoxO1 mediates glucagon-like peptide-1 effects on pancreatic beta-cell mass. *Diabetes* 55, 1190–1196.
- Butler, A. E., Janson, J., Bonner-Weir, S., Ritzel, R., Rizza, R. A., & Butler, P. C. (2003). Beta-cell deficit and increased beta-cell apoptosis in humans with type 2 diabetes. *Diabetes* 52, 102–110.
- Butler, P. C., Meier, J. J., Butler, A. E., & Bhushan, A. (2007). The replication of beta cells in normal physiology, in disease and for therapy. *Nat Clin Pract* 3, 758–768.
- Byrne, M. M., Sturis, J., Sobel, R. J., & Polonsky, K. S. (1996). Elevated plasma glucose 2 h postchallenge predicts defects in beta-cell function. *Am J Physiol* 270, E572–579.
- Calara, F., Taylor, K., Han, J., et al. (2005). A randomized, open-label, crossover study examining the effect of injection site on bioavailability of exenatide (synthetic exendin-4). *Clin Ther* 27, 210–215.
- Cassidy, J., Baughman, R., Costello, D., Levy, B., va Vliet, A., Diaz, M., et al. (2008). Pulmonary administration of GLP-1 (GLP-1 Technosphere powder): Kinetics. *Diabetologia* 51(Suppl.), S346.
- Cersosimo, E. (2007). Mechanisms of exenatide attenuation of postprandial hyperglycaemia in T2DM subjects. *Diabetologia* 50, (Suppl.) Abstract # 250.
- Chae, S. Y., Cheng-Hao, J., Han, J. S., Yu, S. Y., Seulki, L., & Kang, C. L. (2008). Preparation, Characterization, and Application of Biotinylated and Biotin-PEGylated Glucagon-Like Peptide-1 Analogues for Enhanced Oral Delivery. *Bioconjugate Chem* 19, 334–341.
- Chance, W. T., & Fischer, J. E. (1993). Neurotransmitters and food intake. *Nutrition* 9, 470–471.
- Charbonnel, B., Karasik, A., Liu, J., Wu, M., & Meininger, G. (2006). Efficacy and safety of the dipeptidyl peptidase-4 inhibitor sitagliptin added to ongoing metformin therapy in patients with type 2 diabetes inadequately controlled with metformin alone. *Diabetes Care* 29, 2638–2643.
- Chen, Y. E., & Drucker, D. J. (1997). Tissue-specific expression of unique mRNAs that encode proglucagon-derived peptides or exendin 4 in the lizard. *J Biol Chem* 272, 4108–4115.
- Chen, D., Lao, J., Li, N., et al. (2007). A nonpeptidic agonist of glucagon-like peptide I receptors with efficacy in diabetic db/db mice. *Proc Natl Acad Sci (USA)* 104, 943–948.
- Chen, C. H., Stephens, R. L., Jr., & Rogers, R. C. (1997). PYY and NPY: control of gastric motility via action on Y1 and Y2 receptors in the DVC. *Neurogastroenterol Motil* 9, 109–116.
- Christopher, R., Covington, P., Davenport, M., Fleck, P., Mekki, Q., Wann, E., et al. (2007). Single-rising-dose pharmacodynamics, and tolerability of alogliptin benzoate (SYR-322), a novel, dipeptidyl peptidase-IV inhibitor, in healthy male subjects. *Diabetologia* 50, (Suppl.) Abstract # 891.
- Conarello, S. L., Li, Z., Ronan, J., Roy, R. S., Zhu, L., Jiang, G., et al. (2003). Mice lacking dipeptidyl peptidase IV are protected against obesity and insulin resistance. *PNAS (USA)* 100, 6825–6830.
- Creutzfeldt, W. (1979). The incretin concept today. *Diabetologia* 16, 75–85.
- Creutzfeldt, W. (2005). The [pre-] history of the incretin concept. *Regul Pept* 128, 87–91.
- Creutzfeldt, W., & Nauck, M. (1992). Gut hormones and diabetes mellitus. *Diabetes/Metabol Rev* 8, 149–177.
- Creutzfeldt, W. O., Kleine, N., Willms, B., Orskov, C., Holst, J. J., & Nauck, M. A. (1996). Glucagonostatic actions and reduction of fasting hyperglycemia by exogenous glucagon-like-peptide I (7–36) amide in type I diabetic patients. *Diabetes Care* 19, 580–586.
- D'Alessio, D. A., & Tso, P. (2005). GLP-1 reduces intestinal lymph flow, triglyceride absorption and lipoprotein production in rats. *Am J Physiol* 288, G943–G949.
- D'Alessio, D. A., Kahn, S. E., Leusner, C. R., & Ensink, J. W. (1994). Glucagon-like peptide 1 enhances glucose tolerance both by stimulation of insulin release and by increasing insulin-independent glucose disposal. *J Clin Invest* 93, 2263–2266.
- Dang, N. H., Torimoto, Y., Shimamura, K., Tanaka, T., Daley, J. F., Schlossman, S. F., et al. (1991). IF17 (CD26): a marker of thymic maturation involved in the differential regulation of the CD3 and CD2 pathways of human thymocyte activation. *J Immunol* 147, 2825–2832.
- Dardik, B., Valentin, M., Schwartzkopf, C., et al. (2003). NVP-LAF237, a dipeptidyl peptidase IV inhibitor, improves glucose tolerance and delays gastric emptying in obese insulin resistant cynomolgus monkeys. *Diabetes* 52(suppl 1), 322 (Abstr 1292-P).
- Darmoul, D., Voisin, T., Couvineau, A., Rouyer Fessard, C., Salomon, R., Wang, Y., et al. (1994). Regional expression of epithelial dipeptidyl peptidase IV in the human intestines. *Biochem Biophys Res Commun* 203, 1224–1229.
- Davis, S. N., Johns, D., Maggs, D., Xu, H., Northrup, J. H., & Brodows, R. G. (2007). Exploring the substitution of exenatide for insulin in patients with type 2 diabetes treated with insulin in combination with oral antidiabetic agents. *Diabetes Care* 30, 2767–2772.
- de Heer, J., Hoy, M. K., & Holst, J. J. (2005). GLP-1, but not GIP, inhibits glucagon secretion via somatostatin in the perfused rat pancreas. *Diabetologia* (Abstract 164). *European Association for the Study of Diabetes, 41st Annual Meeting, Athens, Greece*.
- De Meester, I., Durinx, C., Bal, G., et al. (2000). Natural substrates of dipeptidyl peptidase IV. *Adv Exp Med Biol* 477, 67–87.
- De Meester, I., Lambeir, A. M., Proost, P., & Scharpe, S. (2003). Dipeptidyl peptidase IV substrates: an update on in vitro peptide hydrolysis by human DPP-IV. *Adv Exp Med Biol* 524, 3–17.
- Deacon, C. F., Plamboeck, A., Moller, S., et al. (2002). GLP-1-(9–36) amide reduces blood glucose in anesthetized pigs by a mechanism that does not involve insulin secretion. *Am J Physiol (Endocrinol Metab)* 282, E873–E879.
- Deacon, C. F. (2004). Therapeutic strategies based on glucagonlike peptide 1. *Diabetes* 53, 2181–2189.
- Deacon, C. F. (2005). What do we know about the secretion and degradation of incretin hormones? *Regul Pept* 128, 117–124.
- Deacon, C. F., Johnsen, A. H., & Holst, J. J. (1995b). Degradation of glucagon-like peptide-1 by human plasma in vitro yields an N-terminally truncated peptide that is a major endogenous metabolite in vivo. *J Clin Endocrinol Metab* 80, 952–957.
- Deacon, C. F., Nauck, M. A., Meier, J., Hücking, K., & Holst, J. J. (2000). Degradation of endogenous and exogenous gastric inhibitory polypeptide in healthy and in type 2 diabetic subjects as revealed using a new assay for the intact peptide. *J Clin Endocrinol Metab* 85, 3575–3581.
- Deacon, C. F., Nauck, M. A., Toft-Nielsen, M., Pridal, L., Willms, B., & Holst, J. J. (1995a). Both subcutaneously and intravenously administered glucagonlike peptide 1 are rapidly degraded from NH₂-terminus in type II diabetic patients and in healthy subjects. *Diabetes* 44, 1126–1131.
- Dear, A. E., Liu, H., Knudsen, L. B., & Simpson, R. W. (2007). Liraglutide modulates expression of vascular adhesion molecules and PAI-1 in an in vitro model of endothelial cell dysfunction. *Diabetologia* 50, (Suppl.) Abstract # 594.
- DeFronzo, R. A., Ratner, R. E., Han, J., Kim, D. D., Fineman, M. S., & Baron, A. D. (2005). Effects of exenatide (exendin-4) on glycemic control and weight over 30 weeks in metformin-treated patients with type 2 diabetes. *Diabetes Care* 28, 1092–1100.
- Degn, K. B., Brock, B., Juhl, C. B., et al. (2004a). Effect of intravenous infusion of exenatide (synthetic exendin-4) on glucose-dependent insulin secretion and counterregulation during hypoglycemia. *Diabetes* 53, 2397–2403.
- Degn, K. B., Juhl, C. B., Sturis, J., et al. (2004b). One week's treatment with the long-acting glucagon-like peptide 1 derivative liraglutide (NN2211) markedly improves 24-h glycaemia and alpha- and beta-cell function and reduces endogenous glucose release in patients with type 2 diabetes. *Diabetes* 53, 1187–1194.
- Dejager, S., Lebeaut, A., Couturier, A., & Schweizer, A. (2006). Sustained reduction in HbA_{1c} during one-year treatment with vildagliptin in patients with type 2 diabetes (T2DM). *Diabetes* 55(Suppl 1), 29 (Abstr 120-OR).
- Dejager, S., Razac, S., Foley, J. E., & Schweizer, A. (2007). Vildagliptin in drug-naïve patients with type 2 diabetes: a 24-week, double-blind, randomized, placebo-controlled, multiple-dose study. *Horm Metab Res* 39, 218–223.
- Demuth, H. U., McIntosh, C. H., & Pederson, R. A. (2005). Type 2 diabetes—therapy with dipeptidyl peptidase IV inhibitors. *Biochim. Biophys Acta* 1751, 33–44.
- Denker, P. S., & Dimarco, P. E. (2006). Exenatide (exendin-4)-induced pancreatitis: a case report. *Diabetes Care* 29, 471.
- Ding, W. G., Renstrom, E., Rorsman, P., Buschard, K., & Gromada, J. (1997). Glucagon-like peptide I and glucosedependent insulinotropic polypeptide stimulate Ca²⁺-induced secretion in rat alpha-cells by a protein kinase A-mediated mechanism. *Diabetes* 46, 792–800.
- Dor, Y. (2006). Beta-cell proliferation is the major source of new pancreatic beta cells. *Nat Clin Pract* 2, 242–243.
- Dor, Y., Brown, J., Martinez, O. I., & Melton, D. A. (2004). Adult pancreatic beta-cells are formed by self-duplication rather than stem-cell differentiation. *Nature* 429, 41–46.
- Drucker, D. J. (2003). Glucagon-like peptides: regulators of cell proliferation, differentiation, and apoptosis. *Mol Endocrinol* 17, 161–171.
- Drucker, D. J. (2006). The biology of incretin hormones. *Cell Metab* 3, 153–165.
- Drucker, D. J., & Nauck, M. A. (2006). The incretin system: glucagon-like peptide-1 receptor agonists and dipeptidyl peptidase-4 inhibitors in type 2 diabetes. *Lancet* 368, 1696–1705.
- Drucker, D. J., Philippe, J., Mojsov, S., Chick, W. L., & Habener, J. F. (1987). Glucagon-like peptide I stimulates insulin gene expression and increases cyclic AMP levels in a rat islet cell line. *Proc Natl Acad Sci (USA)* 84, 3434–3438.
- Dugi, K. A., Heise, T., Ring, A., Graef-Mody, U., Ritzhaupt, A., & Huettner, S. (2007). BI 1356, a novel xanthine-based DPP-IV inhibitor, exhibits high potency with a wide therapeutic window and significantly reduces postprandial glucose excursions after an OGTT. *Diabetologia* 50, (Suppl.) Abstract # 890.
- Dupré, J., Behme, M. T., Hramiak, I. M., et al. (1995). Glucagon-like peptide I reduces postprandial glycemic excursions in IDDM. *Diabetes* 44, 626–630.
- Dupré, J., Behme, M. T., & McDonald, T. J. (2004). Exendin-4 normalized postcibal glycemic excursions in type 1 diabetes. *J Clin Endocrinol Metab* 89, 3469–3473.
- Dupré, J., Ross, S. A., Watson, D., & Brown, J. C. (1973). Stimulation of insulin secretion by gastric inhibitory polypeptide in man. *J Clin Endocrinol Metab* 37, 826–828.
- Durinx, C., Lambeir, A. M., Bosmans, E., et al. (2000). Molecular characterization of dipeptidyl peptidase activity in serum: soluble CD26/dipeptidyl peptidase IV is responsible for the release of X-Pro dipeptides. *Eur J Biochem* 267, 5608–5613.
- Ebert, R., & Creutzfeldt, W. (1987). Gastrointestinal peptides and insulin secretion. *Diabetes Metab Rev* 3, 1–26.

- Eissele, R., Göke, R., Willemer, S., Harthus, H. P., Vermeer, H., Arnold, R., et al. (1992). Glucagon-like peptide-1 cells in the gastrointestinal tract and pancreas of rat, pig and man. *Eur J Clin Invest* 22, 283–291.
- Egan, J. M., Cloquet, A. R., & Elahi, D. (2002a). The insulinotropic effect of acute exendin-4 administered to humans: Comparison of nondiabetic state to type 2 diabetes. *J Clin Endocrinol Metab* 87, 1282–1290.
- Egan, J. M., Meneilly, G. S., Habener, J. F., & Elahi, D. (2002b). Glucagon-like peptide-1 augments insulin-mediated glucose uptake in the obese state. *J Clin Endocrinol Metab* 87, 3768–3773.
- Elahi, D., McAloon-Dyke, M., Fukagawa, N. K., Meneilly, G. S., Sclater, A. L., Minaker, K. L., et al. (1994). The insulinotropic actions of glucose-dependent insulinotropic polypeptide (GIP) and glucagon-like peptide-1 (7–37) in normal and diabetic subjects. *Regul Pept* 51, 63–74.
- Elbrond, B., Jakobsen, G., Larsen, S., et al. (2002). Pharmacokinetics, pharmacodynamics, safety, and tolerability of a single-dose of NN2211, a long-acting glucagon-like peptide 1 derivative, in healthy male subjects. *Diabetes Care* 25, 1398–1404.
- El-Ouaghli, A., Rehling, E., Holst, J. J., et al. (2007). The dipeptidyl peptidase 4 inhibitor vildagliptin does not accentuate glibenclamide-induced hypoglycemia, but reduces glucagon-like peptide 1 and gastric inhibitory polypeptide secretion in response to glucose. *J Clin Endocrinol Metab* 92, 4165–4171.
- El-Ouaghli, A., Rehling, E., Schweizer, A., Holmes, D., & Nauck, M. A. (2003). The dipeptidyl peptidase IV inhibitor LAF237 does not accentuate reactive hypoglycemia caused by the sulfonylurea glibenclamide administered before an oral glucose load in healthy subjects. *Diabetes* 52 (Suppl 1), 118 (Abstr 507-P).
- Elovson, J. (1980). Biogenesis of plasma membrane glycoproteins. Tracer kinetic study of two rat liver plasma membrane glycoproteins in vivo. *J Biol Chem* 255, 5816–5825.
- Elrick, H., Stimmiller, L., Hlad, C. J., & Arai, Y. (1964). Plasma insulin response to oral and intravenous glucose administration. *J Clin Endocrinol Metab* 24, 1076–1082.
- Eng, J., Kleinman, W. A., Singh, L., Singh, G., & Raufman, J. P. (1992). Isolation and characterization of exendin-4, an exendin-3 analogue, from *Heloderma suspectum* venom. Further evidence for an exendin receptor on dispersed acini from guinea pig pancreas. *J Biol Chem* 267, 7402–7405.
- Erlanson Albertsson, C., & Larsson, A. (1988). The activation peptide of pancreatic procolipase decreases food intake in rats. *Regul Pept* 22, 325–331.
- Falko, G. M., Crockett, S. E., Cataland, S., & Mazzaferri, E. L. (1975). Gastric inhibitory polypeptide (GIP) stimulated by fat ingestion by man. *J Clin Endocrinol Metab* 41, 260–265.
- Farilla, L., Bulotta, A., Hirshberg, B., et al. (2003). Glucagon-like peptide 1 inhibits cell apoptosis and improves glucose responsiveness of freshly isolated human islets. *Endocrinology* 144, 5149–5158.
- Fehmann, H. C., Göke, R., & Göke, B. (1995). Cell and molecular biology of the incretin hormones glucagon-like peptide-1 and glucose-dependent insulin releasing polypeptide. *Endocr Rev* 16, 390–410.
- Fehse, F. C., Trautmann, M., Holst, J. J., et al. (2005). Exenatide augments first- and second-phase insulin secretion in response to intravenous glucose in subjects with type 2 diabetes. *J Clin Endocrinol Metab* 90, 5991–5997.
- Finegood, D. T., Scaglia, L., & Bonner-Weir, S. (1995). Dynamics of beta-cell mass in the growing rat pancreas. Estimation with a simple mathematical model. *Diabetes* 44, 249–256.
- Fineman, M. S., Bicsak, T. A., Shen, L. Z., et al. (2003). Effect on glycemic control of exenatide (synthetic exendin-4) additive to existing metformin and/or sulfonylurea treatment in patients with type 2 diabetes. *Diabetes Care* 26, 2370–2377.
- Fineman, M. S., Shen, L. Z., Smitzi, J., et al. (2002). Unique study design: Evaluation of the effect of dose titration on dose-limiting nausea. *Clin Pharmacol Ther* 71, 19 (Abstr).
- Fineman, M. S., Shen, L. Z., Taylor, K., Kim, D. D., & Baron, A. D. (2004). Effectiveness of progressive dose-escalation of exenatide (exendin-4) in reducing dose-limiting side effects in subjects with type 2 diabetes. *Diabetes Metab Res Rev* 20, 411–417.
- Fleck, P., Mekki, Q., Kipnes, M., Wilson, C., & Pratley, R. (2008). Efficacy and safety of alogliptin and glyburide combination therapy in patients with type 2 diabetes. *Diabetologia* 51 (Suppl.), S37.
- Flint, A., Nazzari, K., Jagielski, P., Hindsberger, C., & Zdravkovic, M. (2008). Einfluss von Leberfunktionsstörungen auf die Pharmakokinetik des langwirksamen humanen GLP-1 Analogs Liraglutid. *Diabetologie und Stoffwechsel* 3, S70.
- Flint, A., Raben, A., Astrup, A., & Holst, J. J. (1998). Glucagon-like peptide 1 promotes satiety and suppresses energy intake in humans. *J Clin Invest* 101, 515–520.
- Fonseca, V., Schweizer, A., Albrecht, D., Baron, M. A., Chang, L., & DeJager, S. (2007). Addition of vildagliptin to insulin improves glycaemic control in type 2 diabetes. *Diabetologia* 50, 1148–1155.
- Fukase, N., Manaka, H., Sugiyama, K., Takahashi, H., Igarashi, M., Daimon, M., et al. (1995). Response of truncated glucagon-like peptide-1 and gastric inhibitory polypeptide to glucose ingestion in non-insulin dependent diabetes mellitus. Effect of sulfonylurea therapy. *Acta Diabetol* 32, 165–169.
- Garber, A., Camisasca, R. P., Ehrsam, E., Colloher-Maugeais, C., Rochotte, E., & Lebeaut, A. (2006). Vildagliptin added to metformin improves glycemic control and may mitigate metformin-induced GI side effects in patients with type 2 diabetes (T2DM). *Diabetes* 55 (Suppl 1), 29 (Abstr 121-OR).
- Garber, A. J., Schweizer, A., Baron, M. A., Rochotte, E., & DeJager, S. (2007). Vildagliptin in combination with pioglitazone improves glycaemic control in patients with type 2 diabetes failing thiazolidinedione monotherapy: a randomized, placebo-controlled study. *Diabetes Obes Metab* 9, 166–174.
- Gault, V. A., Irwin, N., Green, B. D., McCluskey, J. T., Greer, B., Bailey, C. J., et al. (2005). Chemical ablation of gastric inhibitory polypeptide receptor action by daily (Pro3)GIP administration improves glucose tolerance and ameliorates insulin resistance and abnormalities of islet structure in obesity-related diabetes. *Diabetes* 54, 2436–2446.
- Gedulin, B. R., Smith, P., Prickett, K. S., et al. (2005). Dose-response for glycaemic and metabolic changes 28 days after single injection of long-acting release exenatide in diabetic fatty Zucker rats. *Diabetologia* 48, 1380–1385.
- Gedulin, B., Tatarkiewicz, K., Smith, P., Wilson, J., Bhole, D., Kendall, D., et al. (2007). Contrasting actions of sitagliptin and exenatide on food intake, body weight, glucose stimulated insulin secretion and gastric emptying in rodents. *Diabetologia* 50, (Suppl.) Abstract # 576.
- Gerich, J. E. (1989). Abnormal glucagon secretion in type 2 (noninsulin-dependent) diabetes mellitus: causes and consequences. In W. Creutzfeldt & P. Lefèbvre (Eds.), *Diabetes mellitus: pathophysiology and therapy* (pp. 127–133). Berlin, Heidelberg: Springer Verlag; Risk Publications.
- Göke, R., Fehmann, H. C., Linn, T., et al. (1993). Exendin-4 is a high potency agonist and truncated exendin-(9–39) amide and antagonist at the glucagon-like peptide 1-(7–36)-amide receptor of insulin-secreting β -cells. *J Biol Chem* 268, 19650–19655.
- Göke, B., Fuder, H., Wieckhorst, G., et al. (1995a). Voglibose (AO-128) is an efficient alpha-glucosidase inhibitor and mobilizes the endogenous GLP-1 reserve. *Digestion* 56, 493–501.
- Göke, B., Herrmann, C., Göke, R., Fehmann, H. C., Berghöfer, P., Richter, G., et al. (1994). Intestinal effects of alpha-glucosidase inhibitors: absorption of nutrients and enterohormonal changes. *Eur J Clin Invest Suppl* 3, 25–30.
- Göke, R., Larsen, P. J., Mikkelsen, J. D., & Sheikh, S. P. (1995b). Distribution of GLP-1 binding sites in the rat brain: evidence that exendin-4 is a ligand of brain GLP-1 binding sites. *Eur J Neurosci* 7, 2294–2300.
- Golpon, H. A., Puechner, A., Welte, T., Wichert, P. V., & Feddersen, C. O. (2001). Vasorelaxant effect of glucagon-like peptide-(7–36)amide and amylin on the pulmonary circulation of the rat. *Regul Pept* 102, 81–86.
- Gomez, E., Pritchard, C., & Herbert, T. P. (2002). cAMP-dependent protein kinase and Ca²⁺ influx through L-type voltage-gated calcium channels mediate Raf-independent activation of extracellular regulated kinase in response to glucagon-like peptide-1 in pancreatic beta-cells. *J Biol Chem* 277, 48146–48151.
- Gonzalez Gronow, M., & Weber, M. R. (1998). Dipeptidyl peptidase IV (CD26) is a receptor for streptokinase on rheumatoid synovial fibroblasts. *Fibrinolysis Proteolysis* 12, 129–135.
- Grøll, M. D. (2005). Dipeptidyl peptidase IV and related enzymes in cell biology and liver disorders. *Clin Sci (London)* 108, 277–292.
- Grandt, D., Dahms, P., Schimiczek, M., Eysselein, V. E., Reeve, J. R., Jr., & Mentlein, R. (1993). Proteolytic processing by dipeptidyl aminopeptidase IV generates receptor selectivity for peptide YY (PYY). *Med Klin* 88, 143–145.
- Green, B. D., & Flatt, P. R. (2007). Incretin hormone mimetics and analogues in diabetes therapeutics. *Best Practice Res Clin Endocrinol Metab* 21, 497–516.
- Green, B. D., Flatt, P. R., & Bailey, C. J. (2006). Inhibition of dipeptidyl IV activity as a therapy of type-2 diabetes. *Expert Opinion on Emerging Drugs* 11, 525–539.
- Green, B. D., Mooney, M. H., Gault, V. A., et al. (2004). Lys9 for Glu9 substitution in glucagon-like peptide-1(7–36)amide confers dipeptidylpeptidase IV Resistance with cellular and metabolic actions similar to those of established antagonists glucagon-like peptide-1(9–36)amide and exendin (9–39). *Metabolism* 53, 252–259.
- Groger, G., Unger, A., Holst, J. J., Goebell, H., & Layer, P. (1997). Ileal carbohydrates inhibit cholinergically stimulated exocrine pancreatic secretion in humans. *Int J Pancreatol* 22, 23–29.
- Groop, P. H., Groop, L., Töttermann, K. J., & Fyhrquist, F. (1986). Effects of acarbose on the relationship between changes in GIP and insulin responses to meals in normal subjects. *Acta Endocrinol (Copenh)* 112, 361–366.
- Guivarch, P. H., Castaigne, J. P., Gagnon, C., Peshherbe, L., Dreyfus, J. H., & Drucker, D. J. (2004). CJC-1131, a long acting GLP-1 analog safely normalizes post-prandial glucose excursion and fasting glycemia in type 2 diabetes mellitus. *Diabetes* 53, A127.
- Guler, H. P. (2007). PHX1149, a selective DPP4 inhibitor, improves postprandial blood glucose control in patients with type 2 diabetes. *Diabetologia* 50, (Suppl.) Abstract #114.
- Gustafson, B., & Smith, U. (2008). WNT signalling is both an inducer and effector of glucagon-like peptide-1. *Diabetologia* 51, 1768–1770.
- Gutzwiller, J. P., Drewe, J., Göke, B., et al. (1999a). Glucagon-like peptide-1 promotes satiety and reduces food intake in patients with diabetes mellitus type 2. *Am J Physiol* 276, R1541–R1544.
- Gutzwiller, J. P., Göke, B., Drewe, J., et al. (1999b). Glucagon-like peptide-1: a potent regulator of food intake in humans. *Gut* 44, 81–86.
- Handelsman, Y. (2008). Treating diabetes with incretin hormones - clinical experience. *Pediatr Endocrinol Rev* 5, 897–903.
- Hanefeld, M., Herman, G., Mickel, C., et al. (2005). Effect of MK-0431, a dipeptidyl peptidase IV (DPP-IV) inhibitor, on glycemic control after 12 weeks in patients with type 2 diabetes. *Diabetologia* 48 (Suppl 1) (Abstr #791).
- Hansen, B. C., Bjerring, C., & Bjerrre Knudsen, L. (2001). Sustained appetite suppression and weight loss in obese rhesus monkeys treated with a long-acting GLP-1 derivative, NN2211. *Diabetologia* 44 (Suppl 1) (Abstr #196).
- Hansen, L., Deacon, C. F., Orskov, C., & Holst, J. J. (1999). Glucagon-like peptide-1-(7–36) amide is transformed to glucagon-like peptide-1-(9–36)amide by dipeptidyl peptidase IV in the capillaries supplying the L cells of the porcine intestine. *Endocrinology* 140, 5356–5363.
- Hanski, C., Huhle, T., & Reutter, W. (1985). Involvement of plasma membrane dipeptidyl peptidase IV in fibronectin-mediated adhesion of cells on collagen. *Biol Chem Hoppe Seyler* 366, 1169–1176.
- Hanski, C., Huhle, T., Gossrau, R., & Reutter, W. (1988). Direct evidence for the binding of rat liver DPP IV to collagen in vitro. *Exp Cell Res* 178, 64–72.
- Hansotia, T., & Drucker, D. J. (2005). GIP and GLP-1 as incretin hormones: lessons from single and double incretin receptor knockout mice. *Regul Pept* 128, 125–134.
- Hansotia, T., Baggio, L. L., Delmeire, D., et al. (2004). Double incretin receptor knockout (DIRKO) mice reveal an essential role for the enteroinsular axis in transducing the glucoregulatory actions of DPP-IV inhibitors. *Diabetes* 53, 1326–1335.

- Heine, R. J., Van Gaal, L. F., Johns, D., Mihm, M. J., Widell, M. H., & Brodows, R. G. (2005). Exenatide versus insulin glargine in patients with suboptimally controlled type 2 diabetes: a randomized trial. *Ann Int Med* 143, 559–569.
- Heit, J. J., Apelqvist, A. A., Gu, X., Winslow, M. M., Neilson, J. R., Crabtree, G. R., et al. (2006). Calcineurin/NFAT signalling regulates pancreatic beta-cell growth and function. *Nature* 443, 345–349.
- Herman, G. A., Stevens, C., Van Dyck, K., et al. (2005). Pharmacokinetics and pharmacodynamics of sitagliptin, an inhibitor of dipeptidyl peptidase IV, in healthy subjects: results from two randomized, double-blind, placebo-controlled studies with single oral doses. *Clin Pharmacol Ther* 78, 675–688.
- Herrmann, K., Gedulin, B., Tatarkiewicz, K., Bhole, D., Kendall, D., Hargrove, D., et al. (2008). Wirkungsunterschiede von Sitagliptin und Exenatide auf Nahrungsaufnahme, Körpergewicht, Glucosestimulierte Insulinsekretion und Magenentleerung in Tierexperimenten. *Diabetologie und Stoffwechsel* 3, S29.
- Heymann, E., Mentlein, R., Nausch, J., Erlanson-Albertson, C., Yoshimoto, T., & Feller, A. C. (1986). Processing of procolipase and trypsinogen by pancreatic dipeptidyl peptidase IV. *Biomed Biochim Acta* 45, 575–584.
- Hildebrandt, M. (2004). Vom Modell zur Therapie: Tumorzelldetektion und Immunmodulation. III Dipeptidylpeptidase IV. Habilitationsschrift Berlin.
- Hildebrandt, M., Reutter, W., Arck, P., et al. (2000). A guardian angel: the involvement of the dipeptidyl peptidase IV in psychoneuroendocrine function, nutrition and immune defence. *Clin Sci* 99, 93–104.
- Hiles, R. A., Bawdon, R. E., & Petrella, E. M. (2003). Ex vivo human placental transfer of the peptides pramlintide and exenatide (synthetic exendin-4). *Hum Exp Toxicol* 22, 623–628.
- Hjellund, K. R., Hughes, T. E., & Holst, J. J. (2007). The dipeptidyl peptidase 4 inhibitor vildagliptin has no acute effect on the endocrine secretion in the isolated perfused porcine pancreas. *Diabetologia* 50, (Suppl.) Abstract # 691.
- Højberg, P. V., Vilsbøll, T., Bache, M., Knop, F. K., Rabøl, R., Krarup, T., et al. (2007). Four weeks of near-normalization of blood glucose restores the insulin response to GIP and improves the insulin response to GLP-1 in patients with type 2 diabetes. *Diabetologia* 50, (Suppl.) Abstract #249.
- Højberg, P. V., Zander, M., Vilsbøll, T., Knop, F. K., Krarup, T., Vølund, A., et al. (2008). Near normalisation of blood glucose improves the potentiating effect of GLP-1 on glucose-induced insulin secretion in patients with type 2 diabetes. *Diabetologia* 51, 632–640.
- Holst, J. J. (1997). Enteroglucagon. *Ann Rev Physiol* 59, 257–271.
- Holst, J. J. (2004). On the physiology of GIP and GLP-1. *Horm Metab Res* 36, 747–754.
- Holst, J. J., & Deacon, C. F. (1998). Perspectives in diabetes. Inhibition of the activity of dipeptidyl-peptidase IV as a treatment for type 2 diabetes. *Diabetes* 47, 1663–1670.
- Holst, J. J., Deacon, C. F., Vilsbøll, T., Krarup, T., & Madsbad, S. (2008). Glucagon-like peptide-1, glucose homeostasis and diabetes. *Trends Mol Med* 14, 161–168.
- Holz, G. G. (2004). Epac: a new cAMP-binding protein in support of glucagon-like peptide-1 receptor-mediated signal transduction in the pancreatic beta-cell. *Diabetes* 53, 5–13.
- Holz, G. G., Kang, G., Harbeck, M., Roe, M. W., & Chepurny, O. G. (2006). Cell physiology of cAMP sensor Epac. *J Physiol* 577, 5–15.
- Hopsu-Havu, V. K., & Glenner, G. G. (1966). A new dipeptide naphthylamidase hydrolyzing glycyl-prolyl-beta-naphthylamide. *Histochemie* 7, 197–201.
- Horowitz, M., Dent, J., Fraser, R., et al. (1994). Role and integration of mechanisms controlling gastric emptying. *Dig Dis Sci* 39 (12 Suppl), 7S–13S.
- Hücking, K., Kostic, Z., Pox, C., et al. (2005). Alpha-Glucosidase inhibition (acarbose) fails to enhance secretion of glucagon-like peptide 1 (7–36 amide) and to delay gastric emptying in type 2 diabetic patients. *Diabet Med* 22, 470–476.
- Hui, H., Nourparvar, A., Zhao, X., & Perfetti, R. (2003). Glucagon-like peptide-1 inhibits apoptosis of insulin-secreting cells via a cyclic 5'-adenosine monophosphate-dependent protein kinase A and a phosphatidylinositol 3-kinase-dependent pathway. *Endocrinology* 144, 1444–1455.
- Hui, H., Wright, C., & Perfetti, R. (2001). Glucagon-like peptide 1 induces differentiation of islet duodenal homeobox-1-positive pancreatic ductal cells into insulin-secreting cells. *Diabetes* 50, 785–796.
- Ilitz, J. L., Baker, D. E., Setter, S. M., & Keith Campbell, R. (2006). Exenatide: an incretin mimetic for the treatment of type 2 diabetes mellitus. *Clin Ther* 28, 652–665.
- Imai, K., Maeda, M., Fujiwara, H., Kariya, M., Takakura, K., Kanzaki, H., et al. (1992). Dipeptidyl peptidase IV as a differentiation marker of the human endometrial glandular cells. *Hum Reprod* 7, 1189–1194.
- Imeryuz, N., Yegen, B. C., Bozkurt, A., Coskun, T., Villanueva-Penacarrillo, M. L., & Ulusoy, N. B. (1997). Glucagon-like peptide-1 inhibits gastric emptying via vagal afferent-mediated central mechanisms. *Am J Physiol* 273, G920–G927 1997.
- Isbil-Buyukcoskun, N., & Gulec, G. (2004). Effects of intracerebroventricularly injected glucagon-like peptide-1 on cardiovascular parameters; role of central cholinergic system and vasopressin. *Regul Pept* 118, 33–38.
- Iwaki Egawa, S., Watanabe, Y., Kikuya, Y., & Fujimoto, Y. (1998). Dipeptidyl peptidase IV from human serum: Purification, characterization, and N-terminal amino acid sequence. *J Biochem* 124, 428–433.
- Jacobsen, L. V., Hindsberger, C., Robson, R., & Zdravkovic, M. (2008). Pharmakokinetik des langwirksamen GLP-1 Analogons Liraglutid zur einmal täglichen Gabe bei Patienten mit Nierenfunktionsstörungen. *Diabetologie und Stoffwechsel* 3, S70.
- Jacobsen, L. V., Hindsberger, C., Robson, R., & Zdravkovic, M. (2007). Pharmacokinetics of the long-acting human GLP-1 analogue liraglutide in subjects with renal impairment. *Diabetologia* 50, (Suppl.) Abstract # 852.
- Joanny, G., Téllez, N., Estilles, E., Nacher, M., Rodriguez-Mulero, S., & Montanya, E. (2007). Increased beta cell replication and mass in 95% pancreatized rats treated with a DPP-IV stabilized GLP-1 analogue. *Diabetologia* 50, (Suppl.) Abstract # 690.
- Jonker, D., Watson, E., Toft, A. D., Kristensen, P., Knudsen, L., & Ingwersen, S. H. (2008). Pharmakokinetisches Modell von Liraglutid, einem humanen GLP-1 Analogon zur einmal täglichen Gabe bei Gesunden und im Vergleich zu Exenatide. *Diabetologie und Stoffwechsel* 3, S78.
- Juhl, C. B., Hollingdal, M., Sturis, J., et al. (2002). Bedtime administration of NN2211, a long-acting GLP-1 derivative, substantially reduces fasting and postprandial glycemia in type 2 diabetes. *Diabetes* 51, 424–429.
- Kahn, S. E., Haffner, S. M., Heise, M. A., Herman, W. H., Holman, R. R., Jones, N. P., et al. (2006). Glycemic durability of rosiglitazone, metformin, or glyburide monotherapy. *N Engl J Med* 355, 2427–2443.
- Kamori, M., Magihara, M., Nagatsu, T., Iwata, H., & Miura, T. (1991). Activities of dipeptidyl peptidase II, dipeptidyl peptidase IV, prolyl endopeptidase, and collagenase-like peptidase in synovial membrane from patients with rheumatoid arthritis and osteoarthritis. *Biochem Med Metab Biol* 45, 154–160.
- Karask, A., Charbonnell, B., Liu, J., Wu, M., Meehan, A., & Meininger, G. (2006). Sitagliptin added to ongoing metformin therapy enhanced glycemic control and beta-cell function in patients with type 2 diabetes. *Diabetes* 55(suppl 1), 119–120 (abstr 501-P).
- Karl, T., Hoffmann, T., Pabst, R., & van Hörsten, S. (2003a). Extreme reduction of dipeptidyl peptidase IV activity in F344 rat substrains is associated with various behavioral differences. *Physiol Behav* 80, 123–134.
- Karl, T., Hoffmann, T., Pabst, R., & van Hörsten, S. (2003b). Behavioral effects of neuropeptide Y in F344 rat substrains with a reduced dipeptidyl-peptidase IV activity. *Pharmacol Biochem Behav* 75, 869–879.
- Karydis, I., & Tolis, G. (1998). Orexis, anorexia, and thyrotropin-releasing hormone. *Thyroid* 8, 947–950.
- Kashima, Y., Miki, T., Shibasaki, T., Ozaki, N., Miyazaki, M., Yano, H., et al. (2001). Critical role of cAMP-GEFII-Rim2 complex in incretin-potentiated insulin secretion. *J Biol Chem* 276, 46046–46053.
- Kastin, A. J., Akerstrom, V., & Pan, W. (2002). Interactions of glucagon-like peptide-1 (GLP-1) with the blood-brain barrier. *J Mol Neurosci* 18, 7–14.
- Kendall, D. M., Riddle, M. C., Rosenstock, J., et al. (2005). Effects of exenatide (exendin-4) on glycemic control over 30 weeks in patients with type 2 diabetes treated with metformin and a sulfonylurea. *Diabetes Care* 28, 1083–1091.
- Kieffer, T. J., McIntosh, C. H., & Pederson, R. A. (1995). Degradation of glucose-dependent insulinotropic polypeptide and truncated glucagon-like peptide 1 in vitro and in vivo by dipeptidyl peptidase IV. *Endocrinology* 136, 3585–3596.
- Kieffer, T. J., & Habener, J. F. (1999). The glucagon-like peptides. *Endocr Rev* 20, 876–913.
- Kim, J. G., Baggio, L. L., Bridon, D. P., et al. (2003). Development and characterization of a glucagon-like peptide 1-albumin conjugate: the ability to activate the glucagon-like peptide 1 receptor in vivo. *Diabetes* 52, 751–759.
- Kim, D., MacConnell, L., Zhuang, D., et al. (2006). Safety and efficacy of a once-weekly, long-acting release formulation of exenatide over 15 weeks in patients with type 2 diabetes. *Diabetes* 55(suppl 1), 116 (Abstr 487-P).
- Kim, D., MacConnell, L., Zhuang, D., Kothare, P. A., Trautmann, M., Fineman, M., et al. (2007). Effects of once weekly dosing of a long-acting release formulation of exenatide on glucose control and body weight in subjects with type 2 diabetes. *Diabetes Care* 30, 1987–1995.
- Kipnes, M. S. (2007). Sitagliptin: a viewpoint by Mark S. Kipnes. *Drugs* 67, 598–599.
- Knop, F. K., Vilsbøll, T., Højberg, P. V., Larsen, S., Madsbad, S., Vølund, A., et al. (2007). Reduced incretin effect in type 2 diabetes: cause or consequence of the diabetic state? *Diabetes* 56, 1951–1959.
- Knudsen, L. B., & Pridal, L. (1996). Glucagon-like peptide-1-(9–36) amide is a major metabolite of glucagon-like peptide-1-(7–36) amide after in vivo administration to dogs, and it acts as an antagonist on the pancreatic receptor. *Europ J Pharmacol* 318, 429–435.
- Knudsen, L. B., Knudsen, S. M., Wilken, M., et al. (2003). Plasma protein binding of NN2211, a long-acting derivative of GLP-1, is important for its efficacy. *Diabetes* 52, A321–A322.
- Knudsen, L. B., Nielsen, P. F., Huusfeldt, P. O., et al. (2000). Potent derivatives of glucagon-like peptide-1 with pharmacokinetic properties suitable for once daily administration. *J Med Chem* 43, 1664–1669.
- Kolterman, O. G., Buse, J. B., Fineman, M. S., et al. (2003). Synthetic exendin-4 (exenatide) significantly reduces postprandial and fasting plasma glucose in subjects with type 2 diabetes. *J Clin Endocrinol Metab* 88, 3082–3089.
- Kolterman, O., Gottlieb, A., Prickett, K., Gaines, E., & Young, A. (2000). Dose-response for inhibition of glucagon secretion and gastric emptying by synthetic exendin-4 (AC2993) in subjects with type 2 diabetes. *Diabetes* 49(Suppl 1), 114 (Abstr).
- Kolterman, O. G., Kim, D. D., Shen, L., et al. (2005). Pharmacokinetics, pharmacodynamics, and safety of exenatide in patients with type 2 diabetes mellitus. *Am Health Syst Pharm* 62, 173–181.
- Kos, K., Baker, A. R., Harte, A. L., O'Hare, J. P., McTernan, P. G., & Kumar, S. (2007). DPP-IV inhibitors influence fat metabolism through modulation of NPY's paracrine antilipolytic action in adipose tissue. *Diabetologia* 50(Suppl.) Abstract # 109.
- Kothare, P. A., Soon, D. K., Linnebjerg, H., et al. (2005). Effect of exenatide on the steady-state pharmacokinetics of digoxin. *J Clin Pharmacol* 45, 1032–1037.
- Kozakova, H., Stepankova, R., Kolinska, J., Farre, M. A., Funda, D. P., Tuckova, L., et al. (1998). Brush border enzyme activities in the small intestine after long-term gliadin feeding in animal models of human coeliac disease. *Folia Microbiol Praha* 43, 497–500.
- Krarup, T. (1988). Immunoreactive gastric inhibitory polypeptide. *Endocr Rev* 9, 122–134.
- Kreisel, W., Reutter, W., & Gerok, W. (1984). Modification of the intramolecular turnover of terminal carbohydrates of dipeptidylaminopeptidase IV isolated from rat-liver plasma membrane during liver regeneration. *Eur J Biochem* 138, 435–438.
- Krentz, A. J., & Bailey, C. J. (2005). Oral antidiabetic agents: current role in type 2 diabetes mellitus. *Drugs* 65, 385–411.

- Kreymann, B., Williams, G., Gbatei, M. A., & Bloom, S. R. (1987). Glucagon-like peptide-1 7-36: a physiological incretin in man. *Lancet* 2(8571), 1300–1304.
- Kumar, M., Hunag, Y., Glinka, Y., Prud'Homme, G. J., & Wang, Q. (2007). Gene therapy of diabetes using a novel GLP-1/IgG1-Fc fusion construct normalizes glucose levels in db/db mice. *Gene Ther* 14, 162–172.
- Lankas, G. R., Leiting, B., Roy, R. S., et al. (2005). Dipeptidyl peptidase IV inhibition for the treatment of type 2 diabetes: potential importance of selectivity over dipeptidyl peptidases 8 and 9. *Diabetes* 54, 2988–2994.
- Larsen, P. J., Tang-Christensen, M., Holst, J. J., & Orskov, C. (1997). Distribution of glucagon-like peptide-1 and other preproglucagon-derived peptides in the rat hypothalamus and brainstem. *Neuroscience* 77, 257–270.
- Lauffer, L. M., Iakubov, R., & Brubaker, P. L. (2009). GPR119 is essential for oleylethanolamide-induced glucagon-like peptide-1 secretion from the intestinal enteroendocrine L-cell. *Diabetes* 58, 1059–1066.
- Lawrence, M. C., Bhatt, H. S., & Easom, R. A. (2002). NFAT regulates insulin gene promoter activity in response to synergistic pathways induced by glucose and glucagon-like peptide-1. *Diabetes* 51, 691–698.
- Lee, S. H., Lee, S., Youn, Y. S., Na, D. H., Chae, S. Y., Byun, Y., et al. (2005). Synthesis, characterization, and pharmacokinetic studies of PEGylated glucagon-like peptide-1. *Bioconjugate Chem* 16, 377–382.
- Lee, A., Patrick, P., Wishart, J., Horowitz, M., & Morley, J. E. (2002). The effects of miglitol on glucagon-like peptide-1 secretion and appetite sensations in obese type 2 diabetics. *Diabetes Obes Metab* 4, 329–335.
- Lee, B., Shi, L., Kassel, D. B., Asakawa, T., Takeuchi, K., & Christopher, R. J. (2008). Pharmacokinetic, pharmacodynamic, and efficacy profiles of alogliptin, a novel inhibitor of dipeptidyl peptidase-4, in rats, dogs, and monkeys. *Eur J Pharmacol* 28, 306–314.
- Lee, S., Youn, Y. S., Lee, S. H., Byun, Y., & Lee, K. C. (2006). PEGylated glucagon-like peptide-1 displays preserved effects on insulin release in isolated pancreatic islets and improved biological activity in db/db mice. *Diabetologia* 49, 1608–1611.
- Lenhard, J. M., Croom, D. K., & Minnick, D. T. (2004). Reduced serum dipeptidyl-IV after metformin and pioglitazone treatments. *Biochem Biophys Res Commun* 324, 92–97.
- Leon, D. D., Crutchlow, M. F., Ham, J. Y., & Stoffers, D. A. (2006). Role of glucagon-like peptide-1 in the pathogenesis and treatment of diabetes mellitus. *Int J Biochem Cell Biol* 38, 845–859.
- Li, Y., Cao, X., Li, L. X., Brubaker, P. L., Edlund, H., & Drucker, D. J. (2005). β -cell PDX1 expression is essential for the glucoregulatory, proliferative, and cytoprotective actions of glucagon-like peptide-1. *Diabetes* 54, 482–491.
- Li, Y., Hansotia, T., Yusta, B., Ris, F., Halban, P. A., & Drucker, D. J. (2003). Glucagon-like peptide-1 receptor signaling modulates beta cell apoptosis. *J Biol Chem* 278, 471–478.
- Liddle, R. A. (1995). Regulation of cholecystokinin secretion by intraluminal releasing factors. *Am J Physiol* 269, G319–327.
- Lim, G. E., & Brubaker, P. L. (2008). The RhoGTPase Cdc42 regulates glucagon-like peptide-1 secretion from the enterendocrine L cell. *Diabetologia* 51(Suppl.), S58.
- Lindsay, J. R., Duffy, N. A., McKillop, A. M., Ardill, J., O'Harte, F. P., Platt, R. R., et al. (2005). Inhibition of dipeptidyl peptidase IV activity by oral metformin in type 2 diabetes. *Diabet Med* 22, 654–657.
- List, J., Rosenstock, J., Sankoh, S., & Chen, R. (2007). Saxagliptin improves glycaemic control as monotherapy in type 2 diabetes. *Diabetologia* 50, (Suppl.) Abstract # 889.
- Lui, E. Y., Asa, S. L., Drucker, D. J., et al. (1990). Glucagon and related peptides in fetal rat hypothalamus in vivo and in vitro. *Endocrinology* 126, 110–117.
- Liu, M. J., Shin, S., Li, N., Shigihara, T., Lee, Y. S., Yoon, J. W., et al. (2007). Prolonged remission of diabetes by regeneration of beta cells in diabetic mice treated with recombinant adenoviral vector expressing glucagon-like peptide-1. *Mol Ther* 15, 86–93.
- Madsbad, S., Schmitz, O., Rasmussen, J., Jakobsen, G., & Matthews, D. R. (2004). Improved glycaemic control with no weight increase in patients with type 2 diabetes after once-daily treatment with the long-acting glucagon-like peptide 1 analog liraglutide (NN2211): A 12-week, double-blind, randomized, controlled trial. *Diabetes Care* 27, 1335–1342.
- Marguet, D., Baggio, L., Kobayashi, T., et al. (2000). Enhanced insulin secretion and improved glucose tolerance in mice lacking CD26. *Proc Natl Acad Sci (USA)* 97, 6874–6879.
- Mari, A., Sallaz, W. M., He, Y. L., et al. (2005). Vildagliptin, a dipeptidyl peptidase-IV inhibitor, improves model-assessed beta-cell function in patients with type 2 diabetes. *J Clin Endocrinol Metab* 90, 4888–4894.
- Marre, M., Shaw, J., Brändle, M., Bebakar, W. M., Kamaruddin, N. A., Strand, J., et al. LEAD-1 SU study group (2009). Liraglutide, a once-daily human GLP-1 analogue, added to a sulphonylurea over 26 weeks produces greater improvements in glycaemic and weight control compared with adding rosiglitazone or placebo in subjects with type 2 diabetes (LEAD-1). *Diabet Med* 26, 268–278.
- Masuyama, J., Berman, J. S., Cruikshank, W. W., Morimoto, C., & Center, D. M. (1992). Evidence for recent as well as long term activation of T cells migrating through endothelial cell monolayers in vitro. *J Immunol* 148, 1367–1374.
- Masuyama, J., Yoshio, T., Suzuki, K., Kitagawa, S., Iwamoto, M., Kamimura, T., et al. (1999). Characterization of the 4C8 antigen involved in transendothelial migration of CD26(hi) T cells after tight adhesion to human umbilical vein endothelial cell monolayers. *J Exp Med* 189, 979–990.
- Matikainen, N., Manttari, S., Schweizer, A., et al. (2006). Vildagliptin therapy reduces postprandial intestinal triglyceride-rich lipoprotein particles in patients with type 2 diabetes. *Diabetologia* 49, 2049–2057.
- McIntyre, N., Holdsworth, C. D., & Turner, D. S. (1964). New interpretation of oral glucose tolerance. *Lancet* 41, 20–21.
- McIntyre, N., Holdsworth, C. D., & Turner, D. S. (1965). Intestinal factors in the control of insulin secretion. *J Clin Endocrinol* 25, 1317–1324.
- McKillop, A. M., Duffy, N. A., Green, B. D., Irwin, N., Gault, V. A., O'Harte, F. P. M., et al. (2007). Antidiabetic action of nateglinide includes augmentation of glucagon-like peptide-1 activity mediated partly through inhibition of DPP IV mediated degradation. *Diabetologia* 50, (Suppl.) Abstract # 882.
- Meier, J. J., Gallwitz, B., Askenas, M., et al. (2005b). Secretion of incretin hormones and the insulinotropic effect of gastric inhibitory polypeptide in women with a history of gestational diabetes. *Diabetologia* 48, 1872–1881.
- Meier, J. J., Gallwitz, B., Salmen, S., et al. (2003a). Normalization of glucose concentrations and deceleration of gastric emptying after solid meals during intravenous glucagon-like peptide 1 in patients with type 2 diabetes. *J Clin Endocrinol Metab* 88, 2719–2725.
- Meier, J. J., Kemmeries, G., Holst, J. J., & Nauck, M. A. (2005a). Erythromycin antagonizes the deceleration of gastric emptying by glucagon-like peptide 1 and unmasks its insulinotropic effect in healthy subjects. *Diabetes* 54, 2212–2218.
- Meier, J. J., Nauck, M. A., Schmidt, W. E., & Gallwitz, B. (2002). Gastric inhibitory polypeptide: the neglected incretin revisited. *Regul Peptides* 107, 1–13.
- Meier, J., Nauck, M. A., Siepmann, N., et al. (2003b). Similar insulin secretory response to a gastric inhibitory polypeptide bolus injection to euglycemia in first-degree relatives of patients with type-2 diabetes and control subjects. *Metabolism* 52, 1579–1585.
- Meier, J. J., Weyhe, D., Michael, M., et al. (2004). Intravenous glucagon-like peptide 1 normalizes blood glucose after major surgery in patients with type 2 diabetes. *Crit Care Med* 32, 848–851.
- Menelly, G. S., Bryer-Ash, M., & Elahi, D. (1993). The effect of glyburide on beta-cell sensitivity to glucose-dependent insulinotropic polypeptide. *Diabetes Care* 16, 110–114.
- Mentlein, R. (1999). Dipeptidyl-peptidase IV (CD26)—role in the inactivation of regulatory peptides. *Regul Pept* 85, 9–24.
- Mentlein, R., Gallwitz, B., & Schmidt, W. E. (1993). Dipeptidyl-peptidase IV hydrolyses gastric inhibitory polypeptide, glucagon-like peptide-1(7–36)amide, peptide histidine methionine and is responsible for their degradation in human serum. *Eur J Biochem* 214, 829–835.
- Migoya, E. M., Miller, J. L., Larson, P. J., Tanen, M. R., Hilliard, D., Deacon, C. F., et al. (2007). Sitagliptin, a selective DPP-4 inhibitor, and metformin have complementary effects to increase active GLP-1 concentrations. *Diabetologia* 50, (Suppl.) Abstract # 111.
- Mikhail, N. (2008). Incretin mimetics and dipeptidyl peptidase 4 inhibitors in clinical trials for the treatment of type 2 diabetes. *Expert Opin Investig Drugs* 17, 845–853.
- Miller, C. B., Zhang, Q., & Sjöholm, A. (2008). Insulin and exendin-4 stimulate proliferation of GLP-1-secreting cells. *Diabetologia* 51(Suppl.), S256.
- Mistry, G. C., Bergman, A. J., Zheng, W., Hreniuk, D., Zinny, M. A., Gottesdiener, K. M., et al. (2008). Sitagliptin, a dipeptidyl peptidase-4 inhibitor, does not alter the pharmacokinetics of the sulphonylurea, glyburide, in healthy subjects. *Br J Clin Pharmacol* 66, 36–42.
- Mitani, H., Takimoto, M., Hughes, T. E., & Kimura, M. (2002). Dipeptidyl peptidase IV inhibition improves impaired glucose tolerance in high fat diet-fed rats; study using a Fischer 344 rat substrain deficient in its enzyme activity. *Jpn J Pharmacol* 88, 442–450.
- Miyawaki, K., Yamada, Y., Ban, N., Ihara, Y., Tsukiyama, K., Zhou, H., et al. (2002). Inhibition of gastric inhibitory polypeptide signaling prevents obesity. *Nat Med* 8, 738–742.
- Miyawaki, K., Yamada, Y., Yano, H., et al. (1999). Glucose intolerance caused by a defect in the entero-insular axis: A study in gastric inhibitory polypeptide receptor knockout mice. *Proc Natl Acad Sci (USA)* 96, 14843–14847.
- Moore, B., Edia, E. S., & Abram, J. H. (1906). On the treatment of diabetes mellitus by an extract of duodenal mucous membrane. *Biochem J* 1, 28–38.
- Moreno, P., Sancho, V., Cancelas, J., Nuche-Berenguer, B., Valverde, I., & Villanueva-Peñacarrillo, M. L. (2007). GLP-1 and Exendin-4 normalise D-glucose uptake in fat and muscle of insulin-resistant rats. *Diabetologia* 50, (Suppl.) Abstract # 583.
- Mrazkova, O., Grim, M., & Carlson, B. M. (1986). Enzymatic heterogeneity of the capillary bed of rat skeletal muscles. *Am J Anat* 177, 141–148.
- Mu, J., Woods, J., Zhou, Y. P., et al. (2006). Chronic inhibition of dipeptidyl peptidase-4 with a sitagliptin analog preserves pancreatic (beta)-cell mass and function in a rodent model of type 2 diabetes. *Diabetes* 55, 1695–1704.
- Muller, W. A., Faloona, G. R., Aguilar-Parada, E., et al. (1970). Abnormal alpha-cell function in diabetes: response to carbohydrate and protein ingestion. *N Engl J Med* 283, 109–115.
- Myers, R. D. (1994). Neuroactive peptides: unique phases in research on mammalian brain over three decades. *Peptides* 15, 367–381.
- Naslund, E., Barkeling, B., King, N., et al. (1999a). Energy intake and appetite are suppressed by glucagon-like peptide-1 (GLP-1) in obese men. *Int J Obes Relat Metab Disord* 23, 304–311.
- Naslund, E., Bogefors, J., Skogar, S., et al. (1999b). GLP-1 slows solid gastric emptying and inhibits insulin, glucagon, and PYY release in humans. *Am J Physiol* 277, R910–916.
- Nauck, M. A., Baller, B., & Meier, J. J. (2004b). Gastric inhibitory polypeptide and glucagon-like peptide-1 in the pathogenesis of type 2 diabetes. *Diabetes* 53, S190–196.
- Nauck, M. A., Bartels, E., Orskov, C., Ebert, R., & Creutzfeldt, W. (1993). Additive insulinotropic effects of exogenous synthetic human gastric inhibitory polypeptide and glucagon-like peptide 1 (7–36) amide infused at near-physiological insulinotropic hormone and glucose concentrations. *J Clin Endocrinol Metab* 76, 912–917.
- Nauck, M. A., Duran, S., Kim, D., Johns, D., Northrup, J., Festa, A., et al. (2007a). A comparison of twice-daily exenatide and biphasic insulin aspart in patients with type 2 diabetes who were suboptimally controlled with sulfonylurea and metformin: a non-inferiority study. *Diabetologia* 50, 259–267.
- Nauck, M. A., & El-Ouaghli, A. (2005). The therapeutic actions of DPP-IV inhibition are not mediated by glucagon-like peptide-1. *Diabetologia* 48, 608–611.

- Nauck, M. A., El-Ouaghli, A., Hompesch, M., et al. (2003). No impairment of hypoglycemia counterregulation via glucagon with NN2211, a GLP-1 derivative, in subjects with type 2 diabetes. *Diabetes* 52(Suppl 1), A128 (Abstr 550-P).
- Nauck, M. A., Frid, A., Hermansen, K., Shah, N. S., Tankova, T., Mitha, I. H., et al. LEAD-2 Study Group(2009). Efficacy and safety comparison of liraglutide, glimepiride, and placebo, all in combination with metformin, in type 2 diabetes: the LEAD (liraglutide effect and action in diabetes)-2 study. *Diabetes Care* 32, 84–90.
- Nauck, M. A., Heimesaat, M. M., Behle, K., et al. (2002). Effects of glucagon-like peptide 1 on counterregulatory hormone responses, cognitive functions, and insulin secretion during hyperinsulinemic, stepped hypoglycemic clamp experiments in healthy volunteers. *J Clin Endocrinol Metab* 87, 1239–1246.
- Nauck, M. A., Heimesaat, M. M., Ørskov, C., Holst, J. J., Ebert, R., & Creutzfeldt, W. (1993a). Preserved incretin activity of glucagon-like peptide 1 [7–36 amide] but not of synthetic human gastric inhibitory polypeptide in patients with type-2 diabetes mellitus. *J Clin Invest* 91, 301–307.
- Nauck, M., Hompesch, M., Filipczak, R., et al. (2004a). Liraglutide significantly improves glycemic control and reduces body weight compared with glimepiride as add-on to metformin in type 2 diabetes. *Diabetes* 53(Suppl 2), A83.
- Nauck, M. A., Homberger, E., Siegel, E. G., et al. (1986). Incretin effects of increasing glucose loads in man calculated from venous insulin and C-peptide responses. *J Clin Endocrinol Metab* 63, 492–498.
- Nauck, M. A., Hompesch, M., Filipczak, R., Le, T. D., Zdravkovic, M., & Glumprecht, J. (2006). Five weeks of treatment with the GLP-1 analogue liraglutide improves glycaemic control and lowers body weight in subjects with type 2 diabetes. *Exp Clin Endocrinol Diabetes* 114, 417–423.
- Nauck, M. A., Kleine, N., Ørskov, C., Holst, J. J., Willms, B., & Creutzfeldt, W. (1993b). Normalization of fasting hyperglycaemia by exogenous glucagon-like peptide 1 (7–36 amide) in type 2 (non-insulin-dependent) diabetic patients. *Diabetologia* 36, 741–744.
- Nauck, M., Meininger, G., Sheng, D., Terranella, L., & Stein, P. P. Sitagliptin Study 024 Group. (2007b). Efficacy and safety of the dipeptidyl peptidase-4 inhibitor, sitagliptin, compared with the sulfonylurea, glipizide, in patients with type 2 diabetes inadequately controlled on metformin alone: a randomized, double-blind, non-inferiority trial. *Diabetes Obes, Metab*, 9, 194–205.
- Nauck, M. A., Niedereichholz, U., Ettler, R., et al. (1997). Glucagonlike peptide 1 inhibition of gastric emptying outweighs its insulinotropic effects in healthy humans. *Am J Physiol* 273, E981–E988.
- Nauck, M., Schmidt, W. E., Ebert, R., et al. (1989). Insulinotropic properties of synthetic human gastric inhibitory polypeptide in man: interactions with glucose, phenylalanine, and cholecystokinin-8. *J Clin Endocrinol Metab* 69, 654–662.
- Nauck, M., Stöckmann, F., Ebert, R., & Creutzfeldt, W. (1986). Reduced incretin effect in type 2 (non-insulin-dependent) diabetes. *Diabetologia* 29, 46–52.
- Nauck, M. A., Walberg, J., Vethacke, A., et al. (2004c). Blood glucose control in healthy subject and patients receiving intravenous glucose infusion or total parenteral nutrition using glucagon-like peptide 1. *Regul Pept* 118, 89–97.
- Nielsen, L. L., & Baron, A. D. (2003). Pharmacology of exenatide (synthetic exendin-4) for the treatment of type 2 diabetes. *Curr Opin Investig Drugs* 4, 401–405.
- Nielsen, L. L., Young, A. A., & Parkes, D. G. (2004). Pharmacology of exenatide (synthetic exendin-4): a potential therapeutic for improved glycemic control of type 2 diabetes. *Regul Pept* 117, 77–88.
- N.N. Amylin Pharmaceuticals, Inc (2005). *Byetta (exenatide) package insert*. San Diego, CA: Amylin (www.byetta.com).
- N.N. Food and Drug Administration Information for Healthcare Professionals Exenatide (marketed as Byetta). Available at www.fda.gov/cder/drug/InfoSheets/HCP/exenatideHCP.htm. Accessed January 17, 2008.
- N.N. UK Prospective Diabetes Study (UKPDS) Group. (1998). Effect of intensive blood-glucose control with metformin on complications in overweight patients with type 2 diabetes (UKPDS 34). *Lancet* 352, 854–865.
- N.N. UK Prospective Diabetes Study 16 (1995). Overview of 6 years' therapy of type II diabetes: a progressive disease. U.K. Prospective Diabetes Study Group. *Diabetes* 44, 1249–1258.
- Nikolaïdis, L. A., Mankad, S., Sokos, G. G., et al. (2004). Effects of glucagon-like peptide-1 in patients with acute myocardial infarction and left ventricular dysfunction after successful reperfusion. *Circulation* 109, 962–965.
- Nystrom, T., Gutniak, M. K., Zhang, Q., et al. (2004). Effects of glucagon-like peptide-1 on endothelial function in type 2 diabetes patients with stable coronary artery disease. *Am J Physiol (Endocrinol Metab)* 287, E1209–E1215.
- O'Donovan, D. G., Doran, S., Feinle-Bisset, C., Jones, K. L., Meyer, J. H., Wishart, J. M., et al. (2004). Effect of variations in small intestinal glucose delivery on plasma glucose, insulin, and incretin hormones in healthy subjects and type 2 diabetes. *J Clin Endocrinol Metab* 89, 3431–3435.
- Okerson, T., Maggs, D., Guan, X., Bowlus, C., Holcombe, J., Wintle, M., et al. (2007). Exenatide improved markers of hepatic function over 3 years in patients with type 2 diabetes. *Diabetologia* 50, (Suppl.) Abstract # 110.
- Okerson, T., Yan, P., Stonehouse, A., Brodows, R., & Bhole, D. (2008). Exenatide improved systolic blood pressure compared to insulin or placebo in patients with type 2 diabetes. *Diabetologia* 51(Suppl.), S350.
- Ørskov, C., Holst, J. J., & Nielsen, O. V. (1988). Effect of truncated glucagon-like peptide-1 [proglucagon-(78–107) amide] on endocrine secretion from pig pancreas, antrum, and nonantral stomach. *Endocrinology* 123, 2009–2013.
- Ørskov, C., Jeppesen, J., Madsbad, S., & Holst, J. J. (1991). Proglucagon products in plasma of noninsulin-dependent diabetes and nondiabetic controls in the fasting state and after oral glucose and intravenous arginine. *J Clin Invest* 87, 415–423.
- Ørskov, C., Poulsen, S. S., Møller, M., & Holst, J. J. (1996). Glucagon-like peptide I receptors in the subfornical organ and the area postrema are accessible to circulating glucagon-like peptide I. *Diabetes* 45, 832–835.
- Ørskov, C., Wettergren, A., & Holst, J. J. (1993). Biological effects and metabolic rates of glucagonlike peptide-1 7–36 amide and glucagonlike peptide-1 7–37 in healthy subjects are indistinguishable. *Diabetes* 42, 658–661.
- Pamir, N., Lynn, F. C., Buchan, A. M., Ehse, J., Hinke, S. A., Pospisilik, J. A., et al. (2003). Glucose-dependent insulinotropic polypeptide receptor null mice exhibit compensatory changes in the enteroinsular axis. *Am J Physiol Endocrinol Metab* 284, E931–939.
- Perabo, M., Auling, B. A., Elfers, E. E., Heyl, J., Ellis, K., & Seeley, R. J. (2008). The effects of jugular vs. portal GLP-1 during hyperglycaemic clamps in rats. *Diabetologia* 51 (Suppl.), S56.
- Perfetti, R., Zhou, J., Doyle, M. E., et al. (2000). Glucagon-like peptide-1 induces cell proliferation and pancreatic-duodenum homeobox-1 expression and increases endocrine cell mass in the pancreas of old glucose-intolerant rats. *Endocrinology* 141, 4600–4605.
- Perley, M. J., & Kipnis, D. M. (1967). Plasma insulin responses to oral and intravenous glucose: studies in normal and diabetic subjects. *J Clin Invest* 46, 1954–1962.
- Phillips, W. T., Schwartz, J. G., & McMahon, C. A. (1992). Rapid gastric emptying of an oral glucose solution in type 2 diabetic patients. *J Nucl Med* 33, 1496–1500.
- Pi-Sunyer, F. X., Schweizer, A., Mills, D., & DeJager, S. (2007). Efficacy and tolerability of vildagliptin monotherapy in drug-naïve patients with type 2 diabetes. *Diabetes Res Clin Pract* 76, 132–138.
- Plamboeck, A., Holst, J. J., Carr, R. D., & Deacon, C. F. (2005). Neutral endopeptidase 24.11 and dipeptidyl peptidase IV are both mediators of the degradation of glucagon-like peptide 1 in the anaesthetized pig. *Diabetologia* 48, 1882–1890.
- Polonsky, K. S., Given, B. D., Hirsch, L. J., Tillil, H., Shapiro, E. T., Beebe, C., et al. (1988). Abnormal patterns of insulin secretion in non-insulin-dependent diabetes mellitus. *N Engl J Med* 318, 1231–1239.
- Pospisilik, J. A., Martin, J., Doty, T., et al. (2003). Dipeptidyl peptidase IV inhibitor treatment stimulates beta-cell survival and islet neogenesis in streptozotocin-induced diabetic rats. *Diabetes* 52, 741–750.
- Pospisilik, J. A., Stafford, S. G., Demuth, H. U., Brownsey, R., Parkhouse, W., Finegood, D. T., et al. (2002). Long-term treatment with the dipeptidyl peptidase IV inhibitor P32/98 causes sustained improvements in glucose tolerance, insulin sensitivity, hyperinsulinemia, and beta-cell glucose responsiveness in VDF (fa/fa) Zucker rats. *Diabetes* 51, 943–950.
- Pratley, R. E., Jauffret-Kamel, S., Galbreath, E., & Holmes, D. (2006). Twelve-week monotherapy with the DPP-4 inhibitor vildagliptin improves glycemic control in subjects with type 2 diabetes. *Horm Metab Res* 38, 423–428.
- Qualmann, C., Nauck, M. A., Holst, J. J., Ørskov, C., & Creutzfeldt, W. (1995). Glucagon-like peptide 1 (7–36 amide) secretion in response to luminal sucrose from the upper and lower gut. A study using alpha-glucosidase inhibition (acarbose). *Scand J Gastroenterol* 30, 892–896.
- Quddusi, S., Vahl, T. P., Hanson, K., Prigeon, R. L., & D'Alessio, D. A. (2003). Differential effects of acute and extended infusions of glucagon-like peptide-1 on first- and second-phase insulin secretion in diabetic and nondiabetic humans. *Diabetes Care* 26, 791–798.
- Rachman, J., Gribble, F. M., Levy, J. C., & Turner, R. C. (1997). Near-normalization of diurnal glucose concentrations by continuous administration of glucagon-like peptide 1 (GLP-1) in subjects with NIDDM. *Diabetologia* 40, 205–211.
- Ranganath, L. R. (2008). Incretins: pathophysiological and therapeutic implications of glucose-dependent insulinotropic polypeptide and glucagon-like peptide-1. *J Clin Pathol* 61, 401–409.
- Ranganath, L., & Morgan, L. (2000). An osmotic stimulus-mediating glucagon-like peptide-1 (7–36 amide) (GLP-1) secretion in acarbose-induced sucrose malabsorption? *Nutrition* 16, 64–65.
- Ranganath, L. R., Beety, J. M., Morgan, L. M., et al. (1996). Attenuated GLP-1 secretion in obesity: cause or consequence. *Gut* 38, 916–919.
- Ranganath, L., Daousi, C., Cleator, J., et al. (2007). Effects of GLP-1 and GIP on energy expenditure and appetite in type 2 diabetes. *National Meeting of the Association of Clinical Biochemistry, Poster T25*.
- Ranganath, L., Norris, F., Morgan, L., et al. (1998). Delayed gastric emptying occurs following acarbose administration and is a further mechanism for its anti-hyperglycaemic effect. *Diabetes Med* 15, 120–124.
- Ranganath, L., Norris, F., Morgan, L., et al. (1999a). Inhibition of carbohydrate-mediated glucagon-like peptide-1 (7–36) amide secretion by circulating non-esterified fatty acids. *Clin Sci* 96, 335–342.
- Ranganath, L., Norris, F., Morgan, L., et al. (1999b). The effect of circulating non-esterified fatty acids on the entero-insular axis. *Eur J Clin Invest* 29, 27–32.
- Ranganath, L., Schaper, F., Gama, R., et al. (1999c). Effect of glucagon on carbohydrate-mediated secretion of glucose-dependent insulinotropic polypeptide (GIP) and glucagon-like peptide-1(7–36 amide) (GLP-1). *Diabetes Metab Res Rev* 15, 390–394.
- Raufman, J. P. (1996). Bioactive peptides from lizard venoms. *Regul Pept* 61, 1–18.
- Raz, I., Hanefeld, M., Xu, L., Caria, C., Williams-Herman, D., & Khatami, H. for the Sitagliptin Study 023 Group. (2006). Efficacy and safety of the dipeptidyl peptidase-4 inhibitor sitagliptin as monotherapy in patients with type 2 diabetes mellitus. *Diabetologia* 49, 2564–2571.
- Richter, B., Bandeira-Echtler, E., Bergerhoff, K., & Lerch, C. (2008a). Emerging role of dipeptidyl peptidase-4 inhibitors in the management of type 2 diabetes. *Vasc Health Risk Manag* 4, 753–768.
- Richter, B., Bandeira-Echtler, E., Bergerhoff, K., & Lerch, C. L. (2008b). Dipeptidyl peptidase-4 (DPP-4) inhibitors for type 2 diabetes mellitus. *Cochrane Database Syst Rev* 2, CD006739.
- Riddle, M. C., Henry, R. R., Poon, T. H., et al. (2006). Exenatide elicits sustained glycaemic control and progressive reduction of body weight in patients with type 2 diabetes inadequately controlled by sulphonylureas with or without metformin. *Diabetes Metab Res Rev* 22, 483–491.

- Ritzel, R., Ørskov, C., Holst, J. J., et al. (1995). Pharmacokinetic, insulinotropic, and glucagonostatic properties of GLP-1 (7–36 amide) after subcutaneous injection in healthy volunteers. Dose-response-relationships. *Diabetologia* 38, 720–725.
- Rocca, A. S., & Brubaker, P. L. (1999). Role of the vagus nerve in mediating proximal nutrient-induced glucagon-like peptide-1 secretion. *Endocrinology* 140, 1687–1694.
- Rolin, B., Larsen, M. O., Gotfredsen, C. F., et al. (2002). The long-acting GLP-1 derivative NN2211 ameliorates glycemia and increases beta-cell mass in diabetic mice. *Am J Physiol (Endocrinol Metab)* 283, 745–752.
- Rosenstock, J., Baron, M. A., Camisasca, R. P., Cressier, F., Couturier, A., & Dejager, S. (2007a). Efficacy and tolerability of initial combination therapy with vildagliptin and pioglitazone compared with component monotherapy in patients with type 2 diabetes. *Diabetes Obes Metab* 9, 175–185.
- Rosenstock, J., Baron, M. A., Dejager, S., Mills, D., & Schweizer, A. (2007b). Comparison of vildagliptin and rosiglitazone monotherapy in patients with type 2 diabetes: a 24-week, double-blind, randomized trial. *Diabetes Care* 30, 217–223.
- Rosenstock, J., Baron, M. A., Schweizer, A., Mills, D., & Dejager, S. (2006a). Vildagliptin is as effective as rosiglitazone in lowering HbA1c but without weight gain in drug-naïve patients with type 2 diabetes (T2DM). *Diabetes* 55(Suppl 1), 133 (abstr 557–P).
- Rosenstock, J., Brazg, R., Andryuk, P. J., Lu, K., & Stein, P. for the Sitagliptin Study 019 Group. (2006c). Efficacy and safety of the dipeptidyl peptidase-4 inhibitor sitagliptin added to ongoing pioglitazone therapy in patients with type 2 diabetes: a 24-week, multicenter, randomized, double-blind, placebo-controlled, parallel-group study. *Clin Ther* 28, 1556–1568.
- Rosenstock, J., Brazg, R., Andryuk, P. J., McCrary Sisk, C., Lu, K., & Stein, P. (2006b). Addition of sitagliptin to pioglitazone improved glycemic control with neutral weight effect over 24 weeks in inadequately controlled type 2 diabetes (T2DM). *Diabetes* 55(Suppl 1), 132–133 (Abstr 556–P).
- Rosenstock, J., Ratner, R. E., & Boka, G. (2008a). Dose range effects of the new once daily GLP-1 receptor agonist AVE0010 added to metformin in type 2 diabetes. *Diabetologia* 51(Suppl.), S66.
- Rosenstock, J., Rendell, M. S., Landin-Olsson, M., Foley, J. E., Rochotte, E., & Baron, M. A. (2008b). Effects of vildagliptin in subjects with IGT. *Diabetes Care* 31, 30–35.
- Ross, S. A., & Dupré, J. (1978). Effects of ingestion of triglyceride or galactose on secretion of gastric inhibitory polypeptide and on responses to intravenous glucose in normal and diabetic subjects. *Diabetes* 27, 327–333.
- Rushakoff, R. J., Goldfine, I. D., Beccaria, L. J., Mathur, A., Brand, R. J., & Liddle, R. A. (1993). Reduced postprandial cholecystokinin (CCK) secretion in patients with noninsulin-dependent diabetes mellitus: evidence for a role for CCK in regulating postprandial hyperglycemia. *J Clin Endocrinol Metab* 76, 489–493.
- Rushakoff, R. J., Goldfine, I. D., Carter, J. D., & Liddle, R. A. (1987). Physiological concentrations of cholecystokinin stimulate amino acid-induced insulin release in humans. *J Clin Endocrinol Metab* 65, 395–401.
- Ryskjaer, J., Deacon, C. F., Carr, R. D., Krarup, T., Madsbad, S., Holst, J., et al. (2006). Plasma dipeptidyl peptidase-IV activity in patients with type-2 diabetes mellitus correlates positively with HbA1c levels, but is not acutely affected by food intake. *Eur J Endocrinol/ Eur Fed Endocr Soc* 155, 485–493.
- Saad, M., An, B., & Santiago, O., and on behalf of the 2072 study group. (2002). The effect of NN2211, a long-acting GLP-1 derivative, on glycemic control and body weight in obese patients with type 2 diabetes. *Diabetologia* 45(Suppl 2), 44 (Abstr).
- Schirra, J., Katschinski, M., Weidmann, C., Schafer, T., Wank, U., Arnold, R., et al. (1996). Gastric emptying and release of incretin hormones after glucose ingestion in humans. *J Clin Invest* 97, 92–103.
- Schirra, J., Nicolaus, M., Roggel, R., Katschinski, M., Storr, M., Woerle, H. J., et al. (2006). Endogenous glucagon-like peptide 1 controls endocrine pancreatic secretion and antro-pyloroduodenal motility in humans. *Gut* 55, 243–251.
- Schirra, J., Nicolaus, M., Woerle, H. J., Struckmeier, C., Katschinski, M., & Göke, B. (2009). GLP-1 regulates gastroduodenal motility involving cholinergic pathways. *Neurogastroenterol Motil* Feb 6. [Electronic publication ahead of print].
- Schirra, J., Sturm, K., Leicht, P., Arnold, R., Göke, B., & Katschinski, M. (1998). Exendin (9–39)amide is an antagonist of glucagon-like peptide-1 (7–36) amide in humans. *J Clin Invest* 101, 1421–1430.
- Schmitz, T., Underwood, R., Khiroua, R., Bachovchin, W. W., & Huber, B. T. (1996). Potentiation of the immune response in HIV-1+ individuals. *J Clin Invest* 97, 1545–1549.
- Schweizer, A., Couturier, A., Foley, J. E., & Dejager, S. (2007). Comparison between vildagliptin and metformin to sustain reductions in HbA1c over 1 year in drug-naïve patients with type 2 diabetes. *Diabet Med* 24, 955–961.
- Scott, R., Herman, G., Zhao, P., Chen, X., Wu, M., & Stein, P. (2005). Twelve-week efficacy and tolerability of MK-0431, a dipeptidyl peptidase IV (DPP-IV) inhibitor, in the treatment of type 2 diabetes (T2D). *Diabetes* 54(Suppl 1), 10–11 (Abstr 41–OR).
- Scott, R., Wu, M., Sanchez, M., & Stein, P. (2007). Efficacy and tolerability of the dipeptidyl peptidase-4 inhibitor sitagliptin as monotherapy over 12 weeks in patients with type 2 diabetes. *Int J Clin Pract* 61, 171–180.
- Scrocchi, L. A., Brown, T. J., McCluskey, N., et al. (1996). Glucose intolerance but normal satiety in mice with a null mutation in the glucagon-like peptide 1 receptor gene. *Nat Med* 2, 1254–1258.
- Seifarth, C., Bergmann, J., Holst, J. J., Ritzel, R., Schmiegel, W., & Nauck, M. A. (1998). Prolonged and enhanced secretion of glucagon-like peptide 1 (7–36 amide) after oral sucrose due to alpha-glucosidase inhibition (acarbose) in type 2 diabetic patients. *Diabet Med* 15, 485–491.
- Sinclair, E. M., & Drucker, D. J. (2005a). Glucagon-like peptide 1 receptor agonists and dipeptidyl peptidase IV inhibitors: new therapeutic agents for the treatment of type 2 diabetes. *Curr Opin Endocrinol Diabetes* 12, 146–151.
- Sinclair, E. M., & Drucker, D. J. (2005b). Proglucagon-derived peptides: mechanisms of action and therapeutic potential. *Physiology (Bethesda)* 20, 357–365.
- Sinha Roy, R., Bergeron, R., Zhu, L., He, H., Jiang, G., Liu, F., et al. (2007). Metformin is a GLP-1 secretagogue, not a dipeptidyl peptidase-4 inhibitor. *Diabetologia* 50, (Suppl.) Abstract # 688.
- Skoglund, G., Hussain, M. A., & Holz, G. G. (2000). Glucagon-like peptide 1 stimulates insulin gene promoter activity by protein kinase A-independent activation of the rat insulin I gene cAMP response element. *Diabetes* 49, 1156–1164.
- Slørdal, L., & Spigset, O. (2006). Heart failure induced by noncardiac drugs. *Drug Saf* 29, 567–589.
- Stancikova, M., Lajda, Z., Lukac, J., & Ruzickova, M. (1992). Dipeptidyl peptidase IV in patients with systemic lupus erythematosus. *Clin Exp Rheumatol* 10, 381–385.
- Stewart, M. W., Matthews, J., De Boever, E. J., Dobbins, R. L., Hodge, R. J., Walker, S. E., et al. (2008). Albiglutide, a long acting GLP-1 mimetic; pharmacodynamics, pharmacokinetics, safety and tolerability in subjects with type 2 diabetes. *Diabetologia* 51 (Suppl.), S347.
- Stoffers, D. A., Kieffer, T. J., Hussain, M. A., Drucker, D. J., Bonner-Weir, S., Habener, J. F., et al. (2000). Insulinotropic glucagon-like peptide 1 agonists stimulate expression of homeodomain protein IDX-1 and increase islet size in mouse pancreas. *Diabetes* 49, 741–748.
- Suzuki, Y., Erickson, R. H., Sedlmayer, A., Chang, S. K., Ikehara, Y., & Kim, Y. S. (1993). Dietary regulation of rat intestinal angiotensin-converting enzyme and dipeptidyl peptidase IV. *Am J Physiol* 264, G1153–1159.
- Takeuchi, K., Moritoh, Y., Asakawa, T., Kataoka, O., Shi, L., & Odaka, H. (2007). Alogliptin (SYR-322) is a novel, potent, highly selective dipeptidyl peptidase-IV inhibitor that improves glucose tolerance and β cell function in ob/ob mice. *Diabetologia* 50, (Suppl.) Abstract # 878.
- Taldone, T., Zito, S. W., & Talele, T. T. (2006). Inhibition of dipeptidyl peptidase-IV (DPP-IV) by atorvastatin. *Bioorg Med Chem Lett* 18, 479–484.
- Tang-Christensen, M., Larsen, P. J., Göke, R., et al. (1996). Central administration of GLP-1 (7–36) amide inhibits food and water intake in rats. *Am J Physiol* 271, R848–R856.
- Teta, M., Rankin, M. M., Long, S. Y., Stein, G. M., & Kushner, J. A. (2007). Growth and regeneration of adult beta cells does not involve specialized progenitors. *Developmental Cell* 12, 817–826.
- Thews, D., Werner, U., & Eckel, J. (2008). Enhanced protection against cytokine- and fatty acid-induced apoptosis in pancreatic beta cells by combined treatment with glucagon-like peptide-1 receptor agonists and insulin analogues. *Horm Metab Res* 40, 172–180.
- Thomas, L., Eckhardt, M., Langkopf, F. E., Tadayyon, M., Himmelsbach, F., & Mark, M. (2008b). (R)-8-(3-amino-piperidin-1-yl)-7-but-2-ynyl-3-methyl-1-(4-methyl-quinazolin-2-ylmethyl)-3,7-dihydro-purine-2,6-dione (BI 1356), a novel xanthine-based dipeptidyl peptidase 4 inhibitor, has a superior potency and longer duration of action compared with other dipeptidyl peptidase-4 inhibitors. *J Pharmacol Exp Ther* 325, 175–182.
- Thomas, L., Himmelsbach, F., Eckhardt, M., Langkopf, E., & Mark, M. (2007). BI 1356, a novel and selective xanthine based DPP-IV inhibitor, exhibits a superior profile when compared to sitagliptin and vildagliptin. *Diabetologia* 50, (Suppl.) Abstract # 879.
- Thomas, L., Himmelsbach, F., Eckhardt, M., Langkopf, E., & Mark, M. (2008a). BI 1356 (geplanter Handelsname ONDERO), ein neuer DPP-4-Inhibitor aus der Strukturklasse der Xanthine, zeigt ein herausragendes präklinisches Profil. *Diabetologie und Stoffwechsel* 3, S20.
- Thorens, B., & Widmann, C. (1996). Signal transduction and desensitization of the glucagon-like peptide-1 receptor. *Acta Physiol Scand* 157, 317–319.
- Thorkildsen, C., Neve, S., Larsen, B. D., Meier, E., & Petersen, J. S. (2003). Glucagon-like peptide 1 receptor agonist ZP10A increases insulin mRNA expression and prevents diabetic progression in db/db mice. *J Pharmacol Exp Ther* 307, 490–496.
- Thrainsdottir, I., Malmberg, K., Olsson, A., Gutniak, M., & Ruydén, L. (2004). Initial experience with GLP-1 treatment on metabolic control and myocardial function in patients with type 2 diabetes mellitus and heart failure. *Diab Vasc Dis Res* 1, 40–43.
- Toft-Nielsen, M. B., Damholt, M. B., Madsbad, S., Hilsted, L. M., Hughes, T. E., Michelsen, B. K., et al. (2001a). Determinants of the impaired secretion of glucagon-like peptide-1 in type 2 diabetic patients. *J Clin Endocrinol Metabolism* 86, 3717–3723.
- Toft-Nielsen, M. B., Madsbad, S., & Holst, J. J. (2001b). Determinants of the effectiveness of glucagon-like peptide-1 in type 2 diabetes. *J Clin Endocrinol Metab* 86, 3853–3860.
- Tornero-Esteban, P., Arnés, L., Sancho, V., Moreno, P., Garcia-Arevalo, M., Valverde, I., et al. (2007). Effect of Exendin-4 upon glucotransporter GLUT-4 expression in the adipose tissue of normal and type 2 diabetic states. *Diabetologia* 50, (Suppl.) Abstract # 591.
- Trautmann, M., MacConell, L., Taylor, K., Zhuang, D., Kothare, P. A., Li, W. I., et al. (2008). Pharmakokinetik und Pharmakodynamik von Exenatide in langwirksamer Formulierung (LAR) als Einzeldosis und nach Mehrfachgabe. *Diabetologie und Stoffwechsel* 3, S72.
- Trumper, A., Trumper, K., Trusheim, H., Arnold, R., Göke, B., & Horsch, D. (2001). Glucosedepeendent insulinotropic polypeptide is a growth factor for beta (INS-1) cells by pleiotropic signaling. *Mol Endocrinol (Baltimore)* 15, 1559–1570.
- Tsukiyama, K., Yamada, Y., Yamada, C., Harada, N., Kawasak, I. Y., Ogura, M., et al. (2006). Gastric inhibitory polypeptide as an endogenous factor promoting new bone formation after food ingestion. *Mol Endocrinol* 20, 1644–1651.
- Turton, M. D., O'Shea, D., Gunn, I., Beak, S. A., Edwards, C. M., Meeran, K., et al. (1996). A role for glucagon-like peptide-1 in the central regulation of feeding. *Nature* 379, 69–72.
- Uematsu, T., Urade, M., Yamaoka, M., & Yoshioka, W. (1996). Reduced expression of dipeptidyl peptidase (DPP) IV in peripheral blood T lymphocytes of oral cancer patients. *J Oral Pathol Med* 25, 507–512.
- Unger, R. H., Aguilar-Parada, E., Muller, W. A., et al. (1970). Studies of pancreatic alpha cell function in normal and diabetic subjects. *J Clin Invest* 49, 837–848.
- Vaag, A. A., Holst, J. J., Volund, A., & Beck-Nielsen, H. B. (1996). Gut incretin hormones in identical twins discordant for non-insulin-dependent diabetes mellitus (NIDDM)–

- evidence for decreased glucagon-like peptide 1 secretion during oral glucose ingestion in NIDDM twins. *Eur J Endocrinol / Eur Fed Endocr Soc* 135, 425–432.
- Vahl, T. P., Tauchi, M., Durler, T. S., Elfers, E. E., Fernandes, T. M., Bitner, R. D., et al. (2007). GLP-1 receptors expressed on nerve terminals in the portal vein mediate the effects of endogenous GLP-1 on glucose tolerance in rats. *Endocrinology* 148, 4965–4973.
- Van Gaal, L. F., Gutkin, S. W., & Nauck, M. A. (2008). Exploiting the antidiabetic properties of incretins to treat type 2 diabetes mellitus: glucagon-like peptide 1 receptor agonists or insulin for patients with inadequate glycemic control? *Eur J Endocrinol* 158, 773–784.
- Vella, A., Bock, G., Giesler, P. D., et al. (2007). Effects of dipeptidyl peptidase-4 inhibition on gastrointestinal function, meal appearance, and glucose metabolism in type 2 diabetes. *Diabetes* 56, 1475–1480.
- Verdich, C., Flint, A., Gutzwiller, J. P., et al. (2001). A meta-analysis of the effect of glucagon-like peptide-1 (7-36) amide on ad libitum energy intake in humans. *J Clin Endocrinol Metab* 86, 4382–4389.
- Verspohl, E. J., Ammon, H. P. T., Williams, J. A., & Goldfine, I. D. (1986a). Evidence that cholecystokinin interacts with specific receptors and regulates insulin release in isolated rat islets of Langerhans. *Diabetes* 35, 38–43.
- Verspohl, E. J., Wunderle, G., Ammon, H. P. T., Williams, J. A., & Goldfine, I. D. (1986b). Proglumide (gastric and cholecystokinin receptor antagonist) inhibits insulin secretion in vitro. *N-S Arch Pharmacol* 332, 284–287.
- Villhauer, E. B., Brinkman, J. A., Naderi, G. B., et al. (2003). 1-[(3-hydroxy-1-adamantyl)amino]acetyl-2-cyano-(S)-pyrrolidine: a potent, selective, and orally bioavailable dipeptidyl peptidase IV inhibitor with antihyperglycemic properties. *J Med Chem* 46, 2774–2789.
- Viltsbøll, T., Knop, F. K., Krarup, T., et al. (2003a). The pathophysiology of diabetes involves a defective amplification of the late-phase insulin response to glucose by glucose dependent insulinotropic polypeptide – regardless of etiology and phenotype. *J Clin Endocrinol Metab* 88, 4897–4903.
- Viltsbøll, T., Krarup, T., Deacon, C. F., Madsbad, S., & Holst, J. J. (2001). Reduced postprandial concentrations of intact biologically active glucagon-like peptide 1 in type 2 diabetic patients. *Diabetes* 50, 609–613.
- Viltsbøll, T., Krarup, T., Sonne, J., Madsbad, S., Volund, A., Juul, A. G., et al. (2003b). Incretin secretion in relation to meal size and body weight in healthy subjects and people with type 1 and type 2 diabetes mellitus. *J Clin Endocrinol Metab* 88, 2706–2713.
- Viltsbøll, T., Zdravkovic, M., Le Thi, T., Krarup, T., Schmitz, O., Courrèges, J. P., et al. (2007). Liraglutide, a long-acting human glucagon-like peptide-1 analog, given as monotherapy significantly improves glycemic control and lowers body weight without risk of hypoglycemia in patients with type 2 diabetes. *Diabetes Care* 30, 1608–1610.
- Viltsbøll, T., Zdravkovic, M., Le-Thi, T., et al. (2006). Liraglutide significantly improves glycemic control, and lowers body weight without risk of either major or minor hypoglycemic episodes in subjects with type 2 diabetes. *Diabetes* 55(Suppl 1), 27–28 (Abstr. 115–OR).
- Wang, Q., & Brubaker, P. L. (2002). Glucagon-like peptide-1 treatment delays the onset of diabetes in 8 week-old db/db mice. *Diabetologia* 45, 1263–1273.
- Wang, H., Lezzi, M., Theander, S., Antinozzi, P. A., Gauthier, B. R., Halban, P. A., et al. (2005). Suppression of Pdx-1 perturbs proinsulin processing, insulin secretion and GLP-1 signalling in INS-1 cells. *Diabetologia* 48, 720–731.
- Wang, Y., Perfetti, R., Greig, N. H., Holloway, H. W., DeOre, K. A., Montrose-Rafizadeh, C., et al. (1997). Glucagon-like peptide-1 can reverse the age-related decline in glucose tolerance in rats. *J Clin Invest* 99, 2883–2889.
- Wang, X., Zhou, J., Doyle, M. E., & Egan, J. M. (2001). Glucagon-like peptide-1 causes pancreatic duodenal homeobox-1 protein translocation from the cytoplasm to the nucleus of pancreatic beta-cells by a cyclic adenosine monophosphate/protein kinase A-dependent mechanism. *Endocrinology* 142, 1820–1827.
- Ward, W. K., Bolgiano, D. C., McKnight, B., Halter, J. B., & Porte, D., Jr. (1984). Diminished B cell secretory capacity in patients with noninsulin-dependent diabetes mellitus. *J Clin Invest* 74, 1318–1328.
- Wettergren, A., Maina, P., Boesby, S., & Holst, J. J. (1997a). Glucagonlike peptide-1 7-36 amide and peptide YY have additive inhibitory effect on gastric acid secretion in man. *Scand J Gastroenterol* 32, 552–555.
- Wettergren, A., Petersen, H., Orskov, C., Christiansen, J., Sheikh, S. P., & Holst, J. J. (1994). Glucagon-like peptide-1 7-36 amide and peptide YY from the L-cell of the ileal mucosa are potent inhibitors of vagally induced gastric acid secretion in man. *Scand J Gastroenterol* 29, 501–505.
- Wettergren, A., Schjoldager, B., Mortensen, P. E., Myhre, J., Christiansen, J., & Holst, J. J. (1993). Truncated GLP-1 (proglucagon 78–107-amide) inhibits gastric and pancreatic functions in man. *Dig Dis Sci* 38, 665–673.
- Wettergren, A., Wojdemann, M., Meisner, S., Stadil, F., & Holst, J. J. (1997b). The inhibitory effect of glucagon-like peptide-1 (GLP-1) 7-36 amide on gastric acid secretion in humans depends on an intact vagal innervation. *Gut* 40, 597–601.
- Wheeler, M. B., Gelling, R. W., McIntosh, C. H., Georgiou, J., Brown, J. C., & Pederson, R. A. (1995). Functional expression of the rat pancreatic islet glucose-dependent insulinotropic polypeptide receptor: ligand binding and intracellular signaling properties. *Endocrinology* 136, 4629–4639.
- Wideman, R. D., & Kieffer, T. J. (2004). Glucose-dependent insulinotropic polypeptide as a regulator of beta cell function and fate. *Horm. Metab Res* 36, 782–786.
- Willheim, M., Ebner, C., Baier, K., Kern, W., Schratlbauer, K., Thien, R., et al. (1997). Cell surface characterization of T lymphocytes and allergen-specific T cell clones: correlation of CD26 expression with T(H1) subsets. *J Allergy Clin Immunol* 100, 348–355.
- Willms, B., Werner, J., Holst, J. J., Ørskov, C., Creutzfeldt, W., & Nauck, M. A. (1996). Gastric emptying, glucose responses, and insulin secretion after a liquid test meal: effects of exogenous glucagon-like peptide-1 (GLP-1)-(7-36) amide in type 2 (noninsulin-dependent) diabetic patients. *J Clin Endocrinol Metab* 81, 327–332.
- Wilson, M. J., Ruhland, A. R., Pryor, J. L., Ercole, C., Sinha, A. A., Hensleigh, H., et al. (1998). Prostate specific origin of dipeptidylpeptidase IV (CD-26) in human seminal plasma. *J Urol* 160, 1905–1909.
- Xie, D., Zhong, Q., Ding, K. H., Cheng, H., Williams, S., Correa, D., et al. (2007). Glucose-dependent insulinotropic peptide-overexpressing transgenic mice have increased bone mass. *Bone* 40, 1352–1360.
- Xu, G., Stoffers, D. A., Habener, J. F., & Bonner-Weir, S. (1999). Exendin-4 stimulates both beta-cell replication and neogenesis, resulting in increased beta-cell mass and improved glucose tolerance in diabetic rats. *Diabetes* 48, 2270–2276.
- Xu, L., & Williams-Herman, D. E. (2007). Initial combination therapy with sitagliptin, a selective DPP-4 inhibitor, and metformin leads to marked improvement in β cell function in patients with type 2 diabetes. *Diabetologia* 50, (Suppl.) Abstract # 884.
- Yamamoto, H., Lee, C. E., Marcus, J. N., Williams, T. D., Overton, J. M., Lopez, M. E., et al. (2002). Glucagon-like peptide-1 receptor stimulation increases blood pressure and heart rate and activates autonomic regulatory neurons. *J Clin Invest* 110, 43–52.
- Yaron, A., & Naider, F. (1993). Proline-dependent structural and biological properties of peptides and proteins. *Crit Rev Biochem Mol Biol* 28, 31–81.
- Yasuda, Y., Goto, M., Nakaya, K., Kawamura, S., Furuta, S., Tamura, M., et al. (2007). Pharmacodynamic characteristics of SK-403, a novel highly selective dipeptidyl peptidase IV inhibitor, in animal models. *Diabetologia* 50, (Suppl.) Abstract # 689.
- Youn, Y. S., Chae, S. Y., Lee, S., Jeon, J. E., Shin, H. G., & Lee, K. C. (2007). Evaluation of therapeutic potentials of site-specific PEGylated glucagon-like peptide-1 isomers as a type 2 anti-diabetic treatment: Insulinotropic activity, glucose-stabilizing capability, and proteolytic stability. *Biochem Pharmacol* 73, 84–93.
- Yu, M., Moreno, C., Hoagland, K. M., et al. (2003). Antihypertensive effect of glucagon-like peptide 1 in Dahl salt-sensitive rats. *J Hypertens* 21, 1125–1135.
- Zander, M., Madsbad, S., Madsen, J. L., & Holst, J. J. (2002). Effect of 6-week course of glucagon-like peptide 1 on glycaemic control, insulin sensitivity, and beta-cell function in type 2 diabetes: a parallel-group study. *Lancet* 359, 824–830.
- Zdravkovic, M., Ekblom, M., Bronsted, L., Vouis, J., Lennernäs, H., & Malm-Erfjelt, M. (2008). The effect of liraglutide on the absorption pharmacokinetics of concomitant oral drugs with different solubility and permeability properties in healthy subjects. *Diabetologia* 51(Suppl.), S355.
- Zhao, T., Parikh, P., Bhashyam, S., et al. (2006). Direct effects of glucagon-like peptide-1 on myocardial contractility and glucose uptake in normal and posts ischemic isolated rat hearts. *J Pharmacol Exp Ther* 317, 1106–1113.
- Zhou, J., Wang, X., Pineyro, M. A., & Egan, J. M. (1999). Glucagon-like peptide 1 and exendin-4 convert pancreatic AR42J cells into glucagon- and insulin-producing cells. *Diabetes* 48, 2358–2366.
- Zimmet, P., Alberti, K. G. M. M., & Shaw, J. (2001). Global and societal implications of the diabetes epidemic. *Nature* 414, 782–787.
- Zinman, B., Hoogwerf, B., Garcia, S. G., et al. (2006). Safety and efficacy of exenatide in patients with type 2 diabetes mellitus using thiazolidinediones with or without metformin. *Diabetes* 55(Suppl. 1), 28 (Abstract 117-OR).
- Zinman, B., Hoogwerf, B. J., Duran, G. S., et al. (2007). The effect of adding exenatide to a thiazolidinedione in suboptimally controlled type 2 diabetes: a randomized trial. *Ann Intern Med* 146, 477–485.