

Icta Oncol. Author manuscript; available in PMC 2013 February 13.

Published in final edited form as:

Acta Oncol. 2011 June; 50(Suppl 1): 61-75. doi:10.3109/0284186X.2010.542174.

Tumor markers in prostate cancer I: blood-based markers

Shahrokh F. Shariat¹, Axel Semjonow², Hans Lilja³, Caroline Savage⁴, Andrew J. Vickers⁴, and Anders Bjartell⁵

¹Department of Urology and Medical Oncology, Weill Cornell Medical Center, New York, NY, USA

²Department of Urology, Prostate Center, University Hospital Muenster, Muenster, Germany

³Department of Surgery (Urology Service), Clinical Laboratories, and Medicine (Genito-Urinary Oncology Service), Memorial Sloan-Kettering Cancer Center, New York, NY, USA

⁴Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center, New York, New York, USA

⁵Department of Urology Malmö-Lund, Skåne University Hospital, Lund University, Sweden

Abstract

INTRODUCTION—The introduction of total prostate specific antigen (total PSA) testing in blood has revolutionized the detection and management of men with prostate cancer (PCa). The objective of this review was to discuss the challenges of PCa biomarker research, definition of the type of PCa biomarkers, the statistical considerations for biomarker discovery and validation, and to review the literature regarding total PSA velocity and novel blood-based biomarkers.

METHODS—An English-language literature review of the Medline database (1990 to August 2010) of published data on blood-based biomarkers and PCa was undertaken.

RESULTS—The inherent biological variability of total PSA levels affects the interpretation of any single result. Men who will eventually develop PCa have increased total PSA levels years or decades before the cancer is diagnosed. Total PSA velocity improves predictiveness of total PSA only marginally, limiting its value for PCa screening and prognostication. The combination of PSA molecular forms and other biomarkers improve PCa detection substantially. Several novel blood-based biomarkers such as human glandular kallikrein 2 (hK2), urokinase plasminogen activator (uPA) and its receptor (uPAR), transforming growth factor-beta 1 (TGF- β 1); interleukin-6 (IL-6) and its receptor (IL-6R) may help PCa diagnosis, staging, prognostication, and monitoring. Panels of biomarkers that capture the biologic potential of PCa are in the process of being validated for PCa prognostication.

CONCLUSIONS—PSA is a strong prognostic marker for long-term risk of clinically relevant cancer. However, there is a need for novel biomarkers that aid clinical decision making about biopsy and initial treatment. There is no doubt that progress will continue based on the integrated collaboration of researchers, clinicians and biomedical firms.

Keywords

Prostate neoplasms; mole	ecular markers; prostate specif	nc antigen

INTRODUCTION

In Western societies, prostate cancer (PCa) is the most common solid malignancy and the second leading cause of cancer death in men [1]. The wide availability of total prostatespecific antigen (PSA, formal name human kallikrein 3, hK3) revolutionized PCa screening and ushered in the PSA era resulting in a decrease of PCa metastasis and death. However, the ubiquitous application of PSA screening has also led to over-detection and overtreatment. The lifetime risk of PCa diagnosis is estimated at ~18%, whereas that for death from PCa is ~3%. In addition, PSA is neither cancer specific nor a surrogate for the biologic behavior of PCa. An elevated PSA level can reflect the presence of cancer but can also be caused by benign prostatic hyperplasia (BPH), infection, and/or chronic inflammation. Virtually all prostate epithelial cells, whether normal, hyperplastic or cancerous, synthesize PSA. Therefore, there has been a concerted effort to discover and validate novel PCa biomarkers. In this review article, we first discuss the challenge of PCa biomarker research, types of PCa biomarkers, and the statistical considerations for biomarker discovery and validation. Then, we discuss the limitations of measuring total PSA and its derivatives such as total PSA velocity (total PSAV) and different molecular forms (i.e., free PSA, BPSA, pro-PSA, and intact PSA). Moreover, we briefly discuss several promising novel blood-based biomarkers for PCa diagnosis, staging, prognostication, and monitoring i.e., human glandular kallikrein 2 (hK2), urokinase plasminogen activator (uPA) and its receptor (uPAR), transforming growth factor-beta 1 (TGF-β1); interleukin-6 (IL-6) and its receptor (IL-6R).

BIOMARKER CHALLENGE IN PROSTATE CANCER

A PubMed Search on "prostate cancer" AND ("biomarker" OR "marker") in English language yielded 3016 hits (accessed 8/8/2010). The number of articles published on PCa biomarkers has increased steadily over the years. Despite this plethora of biomarkers reported to be "promising", only one biomarker, (i.e. total PSA in blood) is routinely used by urologists. Why are PCa biomarkers not living up to their promise? For one, there are remarkable analytical and regulatory barriers to the application of biomarkers in PCa care. [2, 3] These include but are not limited to the status of intellectual property protection, availability of standard reference materials for the assay, complexity of assay format, implementation of quality control to assure reproducibility and accuracy, sufficient market testing size to assess methods of commercialization, lack of clear guidelines for good manufacturing/laboratory practice and quality control requirements for all phases of biomarker development, cost and effort required to accumulate clinical data under appropriately designed and Institutional Review Board-approved prospective trials, and the interval required for resolution of patent issues, assay standardization, validation, testing, and regulatory approval.

Besides analytical and regulatory barriers, the lack of PCa biomarker use in daily clinical practice is also a result of poor application of statistics and study design. There has been a lack of effective and efficient strategies to determine which biomarker candidates justify the great investment of time and money required for assay development, optimization and demonstration of analytical robustness. There are now guidelines intended to ensure that biomarker studies conform to some basic standards of design and reporting.[2, 3] For a new biomarker to be clinically useful, it has to answer a clinically relevant question and provide information that is not available in a more simple and cost-effective way. Any new biomarker needs to provide a benefit over these standard criteria or at least improve their accuracy. Before a biomarker assay can be implemented in the community setting, it needs to address four concepts: "better, easier, faster and cheaper".[2]

Conceptually, the development of new biomarkers should be a process that is similar to therapeutic drug evaluation. In 2002, the National Cancer Institute's Early Detection Research Network developed a five phase approach to systematic discovery and validation of biomarkers (Figure 1).[2-6] This schema is not only an intellectual process but also provides a clear scale by which researchers, patients, and investors can evaluate the status of the biomarker in the development process. The expected failure rate of biomarkers in development can be expected to be similar to the one of drugs. A large concerted effort is required to advance the field of PCa biomarker through systematic discovery, verification, and validation- each step coupled with adequate statistical analysis.

REFINING THE DEFINITION OF PROSTATE CANCER BIOMARKER

According to the National Institute of Health (NIH) in the US, a biomarker is a characteristic that is objectively measured and evaluated as an indicator of normal biological processes, pathogenic processes or pharmaceutical responses to a therapeutic intervention [7]. Cancer biomarkers are either produced by the tumor or by the body in response to the tumor. Six different types of biomarker can be differentiated in PCa:

- **1.** Detection/screening: this biomarker is used for evaluating patients with either risk factors for or symptoms of PCa.
- **2.** Diagnostic: this biomarker can help classical histopathological characteristics in assessing presence or absence of cancer.
- **3.** Prognostic: this biomarker is used to predict the outcome of patients based on different risk of recurrence or progression thereby allowing individualized management.
- **4.** Predictive: this biomarker is used to predict whether the treatment (drug or other therapy) will be effective, and/or to monitor the effectiveness of the treatment. It can help identify the best treatment modality.
- **5.** Therapeutic target: this biomarker can help identify the patients who will benefit from a particular treatment regimen. It identifies the molecular targets of novel therapies and is affected by therapy. As of now there is no such blood-based biomarker in clinical use for prostate PCa.
- 6. Surrogate endpoint: this biomarker is used to substitute for a clinical endpoint and/ or to measure clinical benefit, harm or lack of benefit or harm. Surrogates could replace traditional endpoints, such as mortality due to disease or the recurrence or relapse of disease. Biomarkers can reduce time factors and costs for Phase I and II clinical trials by replacing clinical endpoints.

For the purposes of the current review, we will focus on blood-based biomarkers that are the end result of a bioassay, for processing biological material from humans, expressed quantitatively or categorically. Tissue-based and urinebased PCa biomarkers are discussed in separate reports.

STATISTICAL CONSIDERATIONS FOR BIOMARKER DISCOVERY AND VALIDATION

An issue that has received less attention is the degree to which research on biomarkers has made sufficient use of clinically relevant statistics, such as the assessment of predictive accuracy, decision analysis, and/or experimental methodology. Most biomarkers do not provide sufficient information to be used independent of other information. The optimal use of biomarkers lies in incorporating it in a model that also includes standard clinical data.[3,

8-12] To determine the value of a new biomarker, it is not sufficient to show that it is significantly related to the outcome, statistically significant in a multivariable model including the standard clinical and pathologic factors, or more significant than the standard clinical and pathologic factors. A variable that is statistically significant in a multivariable model might not improve the model's predictive accuracy. P-value and odds/hazard ratio do not meaningfully describe a biomarkers' ability to classify patients. For a biomarker to be potentially clinically useful, it is necessary to show that adding the biomarker to an existing model based on the most important clinical and pathologic factors improves the predictive accuracy (discrimination and calibration) of the model.[3, 11, 13-16]

One major issue with model development is the need for appropriate validation. There are two general types of validation: internal validation on original dataset and external validation on independent dataset.[2, 3] External validation on a different data set allows for evaluation of the generalizability of the risk prediction tool to wider populations than originally reported. Finally, methods that incorporate clinical consequences such as decision curve analysis are crucial to the evaluation of biomarkers. Several methods are available including decision curve analysis which combines simplicity with efficient computations. [17-20]

TOTAL PSA AND ITS LIMITATIONS

Neoplastic cells produce somewhat lower and varying tissue levels of PSA compared to benign epithelial cells although both conditions cause total PSA elevation in the blood.[3] Therefore, it has been suggested that total PSA should be considered as a marker of BPH-related prostate volume, growth, and outcome rather than a reliable marker of PCa.[3] Moreover, some aggressive PCa 's do not produce PSA.

PSA levels are inherently variable thereby affecting the interpretation of any single result. [21] Variation in total PSA includes both analytical (i.e., pre-analytical sample handling, laboratory processing, assay performance, and standardization) and biological variation (i.e., metabolism, renal elimination, medication, physical and sexual activity, size and integrity of the prostate). Oscillations up to 20-30% in the total PSA range 0.1-20 ng/mL may be due to biologic variation.[22, 23] Furthermore, the use of different detection assays may be another important cause of variation. Differences in assay standardization can give an artifactually high or low estimate of total PSA and total PSAV.[24-26] Assays are not interchangeable and caution should be exercised when comparing results from different commercial total PSA assays. Patients and physicians should be aware of which assay was used each time a total PSA measurement is performed, and an effort should be made to use the same assay at the next visit. In addition, studies of total PSA kinetics over time using different assays should be interpreted with caution. Of note, there is an expected 20% lower value when the World Health Organization standard is adopted and laboratories should be obliged to mention the name of the PSA assay used on the lab report as well as stating the assay specific reference range and the type of master-calibration (i.e. WHO standardized or traditional calibration).[27-29]

The effect of previous BPH treatment on total PSA remains mostly unpredictable. For example, the effect of commonly used $5-\alpha$ -reductase inhibitors on the predictive value of total PSA kinetics for tumor progression is uncertain. Because $5-\alpha$ -reductase inhibitors are known to decrease the PSA level with ~50% and mostly suppress the benign components of PSA secretion, they may enhance the utility of total PSA.[30] In addition, by shrinking the prostate gland, finasteride has been suggested to increase the likelihood of detecting a small cancer on needle biopsy.[30]

This large normal variability of total PSA requires larger changes between two consecutive measurements to distinguish pathological changes from changes resulting from analytical and biological variations. Nixon et al. calculated the coefficient of variation over 2 weeks and demonstrated that a change between two total PSA measurements of approximately 25% indicated a significant change. [31, 32] Bunting et al. reported a critical difference, defined as the minimum percent change between two consecutive measurements that suggests a significant change beyond the normal variation, close to 60% over a time period of 1 year. [33] Bruun et al. recently assessed the long-term variability of the different forms of PSA at several different total PSA levels in a randomly selected population of asymptomatic and apparently healthy men whose total PSA levels were <2.0 ng/mL at the end of the 8-year observation period.[34] They found that the total intra-individual variation of total PSA was much less than that reported by Bunting et al.[33] and somewhat higher than the intra-individual variation for either free PSA or percent free PSA. This suggests that free PSA concentration in blood may vary less than complexed PSA concentration, which is the major contributor to total PSA. One explanation is that free PSA and complexed PSA may have different elimination pathways, and hence different elimination rates.[35-38]

OPTIMAL TOTAL PSA CUT-OFF VALUES

No single total PSA cut-off separates men at high risk for PCa from men at low risk, nor men affected with high-grade disease from those with low-grade disease. At a total PSA cutoff of 4 ng/mL, a significant number of PCas remain undetected.[39-42] In addition, intervention at lower total PSA levels has been proposed to improve patient outcomes.[43, 44] Catalona et al. found that 22% of men with a normal digital rectal examination (DRE) and a serum total PSA level between 2.6 and 4.0 ng/mL have PCa, and 81% of them have organ-confined disease.[45] Data from the Prostate Cancer Prevention Trial (PCPT) revealed that as many as 15% of men with normal DRE and a serum total PSA less than 4.0 ng/mL have PCa.[39] Among men with total PSA levels 0.5, 0.6–1.0, 1.1–2.0, 2.1–3.0, and 3.1–4.0 ng/mL, PCa was detected in 6.6%, 10.1%, 17.0%, 23.9%, and 26.9%, respectively. Moreover, approximately 25% of these men had a tumor with Gleason score of 7 or higher. These and other investigators demonstrated that increasing levels of total PSA are associated with increasing probability of PCa risk within the 0-4.0 ng/mL interval.[39, 41, 46] There is no total PSA threshold at age 62-91 below which PCa can be ruled out with high specificty. [39] No single total PSA cut-off separates men with "significant" (high grade, high volume) cancer from those with low-grade, possibly insignificant cancer. Similar to PCa presence, high-grade cancer can be found in men with low total PSA levels.

On the other hand, as of now, there is no evidence that lowering the total PSA threshold below 4 ng/mL improves the long-term survival in men with PCa Lowering the total PSA threshold combined with decreasing the age of total PSA screening may be beneficial for men who are at an increased risk for PCa (i.e., strong family history of PCa and/or African-American race). However, consideration must be given to the possibility that lowering the total PSA threshold could result in unnecessary biopsies and an increased detection of insignificant cancers. Finally, determination of the optimal, institution-specific, and management-guiding threshold involves not only clinical and epidemiologic features but should also consider the social and psychological implications of prostate biopsy and possible PCa detection.

The difficulty in selecting a cut-off to define what constitutes an abnormal total PSA suggests that total PSA is most useful as a continuous variable, providing a spectrum of prostate cancer risk. Therefore, we prefer to include serum total PSA levels in an overall estimate of the risk of cancer, inform the patient of his particular risk, then make a shared decision about a biopsy.[39-41, 47-51] Nam et al., for example, developed a model that

predicts an individual's risk for PCa in a cohort of 3,108 men who underwent a prostate biopsy for the first time.[51] This model comprises factors that can be easily determined at the time of screening such as age, ethnicity, family history of PCa, the presence of urinary symptoms, total PSA, percent free PSA, and DRE. Addition of all these risk factors improved the predictive accuracy of a base model from 0.62 to 0.74. The main advantage of this and other predictive tools[47] is that clinicians can assess PCa risk on an individual basis and make management decisions. However, despite the reasonable accuracy, similar to all predictive tools, the exact probability cut-off for undergoing or foregoing a biopsy is left with the patient and his treating physician and should be individualized.

LONG-TERM PREDICTION OF THE FUTURE RISK OF PROSTATE CANCER USING TOTAL PSA

Several studies have suggested that total PSA levels are associated with the risk of PCa years, or even decades, before its diagnosis. The first long-term prediction study, which reported that total PSA levels >2.5 ng/mL predicted diagnosis of PCa over the subsequent decade was limited by the small number of cancer cases (n=44) and by the degradation of total PSA in archived serum samples.[52] In a prospective study involving a large number of cases, the lead time between total PSA levels 4 ng/mL and the subsequent clinical diagnosis of PCa was estimated at 5.5 years.[53] Similarly, Fang *et al.* studied the risk of PCa diagnosis in a cohort of 549 men following a baseline total PSA measurement at age 40–60 while providing a median follow-up of ~13 years.[54] They concluded a total PSA value above the age-adjusted median carried a relative risk of subsequent cancer diagnosis of ~3.6.

Two larger studies extended prediction models to lower total PSA ranges and longer followup intervals. Loeb et al. examined 1,178 men in their 40s who had risk factors for PCa.[55] The risk of subsequent PCa diagnosis was 14.6-fold higher for men with a baseline total PSA level between 0.7 and 2.5 ng/mL compared to men with total PSA <0.7 ng/mL. Lilja et al. assessed PCa risk among 21,277 men younger than 50 years when they attended the Malmö Preventive Medicine study (MPM), a cardiovascular risk assessment study. [56] The investigators measured total PSA levels in archived plasma obtained from 462 participants diagnosed with PCa within a median of 18 years from start of the study and from 1,222 matched controls. Total PSA level at age 44-50 was very strongly associated with the likelihood of developing PCa up to 25 years later. The odds ratio for a PCa diagnosis at a total PSA value of 0.51-1.0 ng/mL was 2.51 compared to total PSA 0.50 ng/mL, which roughly corresponded to the population average. The odds ratio increased to 7.02 for a total PSA of 1.0-1.5 ng/mL, and further up to 19.01 for a total PSA of 2.01-3.0 ng/ml compared to a total PSA 0.50 ng/mL. In a follow-up study, the authors have further shown that total PSA level at age 44–50 predicts the likelihood of developing advanced PCa, defined as either locally advanced (clinical T3 or higher) or metastatic disease at the time of diagnosis. [57] In another analysis of the MPM-study cohort, the prognostic accuracy of PSA (both total PSA and complexed PSA, described below) decreased with age. [58] The authors hypothesized that these findings result from a greater prevalence of BPH (and therefore of non-cancer-related total PSA increase) among older men.

An analysis of the same cohort demonstrated that PSA at 60 is an extremely strong predictor of the risk of prostate cancer metastasis (AUC 0.86) and death (AUC 0.90) by age 85. Almost all deaths (90%) occurred in men in the top quartile of PSA levels (> 2 ng / ml); men with PSA below the median (< 1 ng / ml), had an extremely low risk of clinically relevant prostate cancer by age of 85 (0.5% risk of metastasis, 0.2% risk of death from prostate cancer). This suggests that at least half of men can be exempted from prostate cancer

screening at age 60, with early detection efforts focusing on a sub-group of men at elevated risk.[59]

In summary, these studies indicate that men who will eventually develop PCa have increased total PSA levels years or decades before the cancer is diagnosed. These total PSA levels may reflect the long duration of prostate carcinogenesis or could reflect a causal role of total PSA in PCa development and/or progression. A total PSA measurement before age 50 could help risk-stratify men for frequency and/or type of later PCa screening; a PSA at 60 could determine which men need to continue with screening.

PROSTATE-SPECIFIC ANTIGEN DERIVATIVES

Enhancing the diagnostic accuracy of total PSA, particularly specificity, is critical, since higher specificity would reduce the number of biopsies performed in men not affected by PCa. Several different strategies have been investigated, including the use of age-specific total PSA cut-offs, total PSA density, total PSA density of the transition zone, total PSA velocity (total PSAV), and the measurement of various molecular forms of PSA.[47, 60-62] In this section, we will focus on total PSAV and the measurement of various molecular forms of PSA.

Total PSA Velocity

Total PSAV refers to the serial evaluation of serum total PSA concentration over time. [63, 64] Different methods of calculating total PSAV are available (eg, based on the first and the last measured values only or on a regression line through all available measurements, based on normal or logarithmic values), but only small differences in predictive value have been found among these derivatives. Connolly et al found that using all available total PSA measurements in a linear regression analysis should be the method of choice for calculating total PSAV. [65] When using the first and last measurements only, these should at least be separated by a sufficiently long time period.

Carter et al. showed that patients with BPH demonstrated a linear increase in total PSA levels over time, whereas patients with PCa had an initial linear increase with a subsequent exponential rise that occurred approximately 5 years before cancer detection.[63] In men with an initial total PSA level between 4 and 10 ng/mL, a total PSAV cut-off value of 0.75 ng/mL per year provided a sensitivity and specificity for PCa of 79% and >90%, respectively. If the initial total PSA concentration was less than 4 ng/mL, the specificity of total PSAV remained >90%, but the sensitivity dropped to 11%. These results were questioned using relatively short total PSA intervals of 1 and 2 years.[66] Subsequently, Carter et al. showed that total PSAV values are only useful if a minimum of three consecutive measurements are taken over a two year period.[67] While the specificity of total PSAV is high, its sensitivity is too low to advise against prostate biopsy in a patient with an elevated total PSA level who is otherwise healthy and a good candidate for curative therapy. Other limitations of total PSAV include imprecision due to biological and analytical intra-individual variability (see section entitled "Total PSA and its limitations") and total PSA stability.

Prospective screening studies have reported that total PSAV does not appear to add diagnostic value for PCa detection beyond that of a single total PSA level. In an analysis of PCPT data, Thompson et al. found that when total PSAV was used alone, it was an independent predictor of PCa presence and aggressiveness.[40] However, when total PSAV was adjusted for the effect of total PSA and other standard variables, it lost independent predictive value. Similarly, the first two screening rounds of the Rotterdam section of the ERSPC found that total PSAV did not improve accuracy when combined with total PSA in

the prospective setting.[68, 69] A recent analysis from the Prostate, Lung, Colon, and Ovarian (PLCO) cancer screening trial showed that although total PSAV was an independent predictor of high-grade disease, addition of total PSAV to total PSA only slightly increased its performance for prediction of high-grade tumors.[70] Using a large population-based cohort of men in early middle age who were likely to have a low incidence of BPH, Ulmert et al. found no benefit to calculate total PSAV or the velocity of any other PSA form over total PSA for long-term PCa prediction.[71] Of note, the predictive value of total PSAV alone was 71.2%, while the predictive value of a single total PSA was higher (77.1%) and the combined model including both total PSAV and total PSA did not alter the predictive accuracy. The observed lack of additional predictive value for total PSAV indicates that total PSA levels do not increase sharply before PCa diagnosis but rise gradually and slowly over many years, also in those men who later present with advanced cancer.

Several studies have shown that that a high pre-treatment total PSAV is strongly associated with a poor disease-specific survival following diagnosis and could help identify men with low total PSA values who are at increased risk of harboring a potentially lethal tumor. [72-75] Carter et al. found a strong association between survival and higher total PSAV as early as 10-15 years before diagnosis in the Baltimore Longitudinal Study of Aging project. [75] Based on these findings, they proposed that a total PSAV threshold of 0.35 ng/mL per year be used in screening men with low total PSA levels to increase the detection of potentially lethal tumors still in the window of curability. These data have prompted debate as to whether this would suffice as evidence to warrant the National Comprehensive Cancer Network to recommend a prostate biopsy if the total PSAV is greater than 0.5 ng/mL per year. [76] However, in analysis of the PCPT data, Vickers and colleagues showed that biopsying men with low PSA but elevated PSAV, led to a large increase in unnecessary biopsies without detecting an important number of clinically significant cancers. [77]

D'Amico et al. reported that men with a pre-operative total PSAV greater than 2.0 ng/mL per year had a 9.8-fold increased relative risk of death from PCa than men with a lower total PSAV.[72] This analysis is compromised by a low number of events; accordingly, it is impossible to tell whether PSAV adds predictive value to standard predictors such as stage, grade and absolute level of PSA. In a more recent study, these investigators reported that total PSAV was also significantly associated with the risk of cancer-specific mortality following external beam radiation therapy.[73] Conversely, using data from 267 Scandinavian men with localized PCa and baseline total PSA levels <50 ng/mL, Fall and colleagues found that, although prognostically relevant, baseline total PSA levels and relative total PSAV in the first 2 years following diagnosis were not able to predict accurately which patients would have a lethal PCa outcome.[78] Several other studies have found that PSAV does not aid prediction in men treated by radical prostatectomy [79, 80]or conservatively[81].

It may be that the observation period necessary for obtaining a valid calculation of total PSAV that is not disturbed by considerable short-term fluctuations is be too long, or that number of total PSA measurements is too high for use in clinical practice. In addition, total PSAV may not correlate with early tumor progression, but could be a mere indicator of aggressive disease for which the window of curability has already closed. Furthermore, a quickly rising total PSA is more common in men with a high starting total PSA level.[82] This proportion of men is expected to be much smaller in a screened cohort than in a clinically diagnosed cohort. PSAV is a practical parameter after treatment, when PSA is a sensitive measure of cancer; its value in men with a prostate, in whom rises in PSA may be cause by benign disease, remains to be established.

PSA molecular forms

Improvements in measuring PSA isoforms have allowed the measurement of free PSA and its ratio to total PSA.[83-85] Other forms include complexed PSA, which is a measure of how much PSA in serum is bound to $\alpha 2$ -macroglobulin, $\alpha 1$ -protease inhibitor, or $\alpha 1$ -antichymotrypsin. Currently there is no assay commercially available which specifically measures the complex of $\alpha 2$ -macroglobulin with PSA.

The FDA has approved the use of percent free PSA testing [i.e., (fPSA/tPSA) × 100] as an adjunct to total PSA in men with a serum total PSA concentration between 4 and 10 ng/mL. A higher percent free PSA value indicates a lower probability of finding PCa on biopsy and raises the likelihood that the elevation in total PSA is due to the presence of BPH.[86, 87] In a multicenter, prospective trial, Catalona et al. reported that when a percent free PSA of <25% is used for triggering a sextant prostate biopsy, it yielded a 95% sensitivity for PCa detection and increased the specificity by 20% over PSA alone.[86] The area under the curve for percent free PSA was significantly higher than that for total PSA (AUC=0.72 versus AUC=0.53). However, in response to the realization that sextant biopsies misclassify up to one third of patients who have PCa as without cancer, a more recent evaluation of the utility of percent free PSA in patients undergoing extended 10- or 12-core biopsy has suggested a lower diagnostic efficiency of percent free PSA.[88] While most investigators agree that percent free PSA can improve the diagnostic performance of total PSA between 4 and 10 ng/mL, the most appropriate percent free PSA cutoff value remains debatable. Catalona et al. determined that with a percent free PSA cutoff of less or equal to 27%, they were able to obtain a sensitivity of 90% and avoid 18% of unnecessary biopsies in men over the age of 50 with a total PSA of 2.6 to 4.0 ng/mL.[45] In addition, 83% of these cancers were clinically significant. Finally, data on the utility of percent free PSA for the prediction of pathologic grade and stage of PCa is inconclusive. Catalona et al. determined that with a percent free PSA cutoff of less or equal to 27%, they were able to obtain a sensitivity of 90% and avoid 18% of unnecessary biopsies in men over the age of 50 with a total PSA of 2.6 to 4.0 ng/mL.[45] In addition, 83% of these cancers were clinically significant.

In response to the realization that sextant biopsies misclassify up to one third of patients who have CaP as being without cancer, a more recent evaluation of the usefulness of free PSA values in patients undergoing extended 10- or 12-core biopsy has suggested a lower diagnostic efficiency of free PSA.[88] While most investigators agree that free PSA can improve the diagnostic performance of total PSA between 4 and 10 ng/mL, the most appropriate free PSA cutoff value remains debated.

Data on the usefulness of free PSA to predict clinical outcomes is inconclusive. Graefen and coworkers[89] failed to detect an independent association of preoperative free PSA with biochemical failure in 581 unscreened patients who underwent radical prostatectomy for clinically localized CaP.[90] In contrast, Shariat and colleagues found that lower preoperative serum free PSA is an independent predictor of advanced pathologic features, biochemical progression, and patterns of aggressive disease progression in 402 consecutive men treated with radical prostatectomy for clinically localized CaP who had total PSA levels less than 10 ng/mL.[84]

There are three distinct cleavage isoforms of free PSA in the serum: pro-PSA, BPH-associated PSA (BPSA), and intact free PSA.[91] The precursor of PSA is a 261 amino acid pre-pro-protein. Subsequent processing by human glandular kallikrein 2 (hK2) and other proteases produces the active 237 amino acid mature PSA.[91] Studies have shown that higher levels of pro-PSA are associated with PCa. In men with PSA levels between 6.0-24.0 ng/ml, the [-2]proPSA fraction was found to be significantly higher in men with PCa. [91, 92] Moreover, authors demonstrated the utility of the pro-PSA to free PSA ratio for

screening patients with PSA levels between 2.5-4.0 ng/ml and between 4.0-10.0 ng/ml.[93] Elevated pro-PSA to free PSA ratios have also been associated with aggressive pathological features and decreased biochemical disease free survival after radical prostatectomy[94, 95]. A new automated tool using the [-2]proPSA assay with a percent free PSA based artificial neural network was capable of detecting PCa and more aggressive disease with higher accuracy than total PSA or percent free PSA alone.[96] In a recent prospective cohort of men enrolled into active surveillance for PCa, serum and tissue levels of pro-PSA at diagnosis were associated with need for subsequent treatment.[97] The authors hypothesized that the increase in the ratio of serum pro-PSA to percent free PSA might be driven by increased pro-PSA production from "premalignant" cells.

Molecular forms of PSA may differ in their in-vitro stability properties and information about the preanalytical conditions is therefore essential for proper clinical interpretation. For proper measurement of [-2]proPSA, blood samples should be centrifuged within 3 hours of blood draw. Serum may be stored at room temperature or refrigerated (+4°C) for a maximum of 48 hours and should be frozen if stored for a longer period. Two freeze-thaw cycles have no effect on [-2]proPSA stability.[98]

BPH-associated PSA (BPSA) is formed by the internal cleavage of free PSA between Lys182 and Ser183. BPSA is expressed in nodular hyperplasia limited to the transitional zone of men with BPH. BPSA can be detected in semen, blood and prostate, and its levels correlate with transitional zone volume and obstructive voiding symptoms.[91, 99] BPSA seems to be a promising marker of BPH since it has been shown a direct association between its secretion and the volume of the transition zone.[99] As such, BPSA is a better predictor of prostate enlargement than total and free PSA.[100] In addition, BPSA is not affected by age and is significantly higher in the presence of BPH symptoms. Adjusting the level of free PSA for BPSA resulted in 13-17% improvement in specificity compared to free PSA alone while maintaining a sensitivity of 90-95%.[101]

Combination of PSA molecular forms for improved cancer detection

It is most unlikely that a single biomarker will have the single decision as to a diagnosis and/or a prognosis of a particular pathology. The future of cancer profiling relies on the combination of a panel of complimentary biomarkers that can give accurate molecular staging and indicate the likelihood of aggressive behavior.[10, 11, 102, 103]

The group of Vickers and Lilja has developed a statistical model that predicts prostate biopsy outcomes based on age, DRE and a panel of four kallikrein markers - total PSA, free PSA, intact PSA and hK2. Using data from the randomized prostate cancer screening trial in Göteborg, Sweden [one centre of the European Randomized Study of Prostate Cancer screening (ERSPC)], they estimated that for every 1000 previously unscreened men with elevated total PSA, use of the model to determine biopsy would reduce biopsy rates by 573, while missing only a small number of cancers (31 out of 152 low-grade cancers and 3 out of 40 high-grade cancers).[15] These findings were subsequently replicated in an independent cohort (reduction in biopsy by 513 per 1000 men with elevated total PSA, missing 54 out of 177 low-grade cancer and 12 out of 100 high-grade cancers).[104] These findings have also been verified in men who recently have undergone previous screening, with resultant improvements in predictive accuracy.[105, 106] Recently, Gupta et al. demonstrated that the panel of four kallikrein markers can predict the outcome of prostate biopsy in men who had previously undergone prostate biopsy during previous screening.[107] This model in addition to age and DRE substantially improved the predictive accuracy of a base model (comprising of total PSA, age and DRE), for both low- and high-grade cancers.

EMERGING BLOOD-BASED BIOMARKERS FOR BOTH PROSTATE CANCER DETECTION AND PROGNOSTICATION

Human Kallikrein 2 (hK2)

Human kallikrein related peptidase 2 is a secreted serine protease from the same gene family as PSA.[108] They share an 80% sequence homology and are both primarily expressed in the prostate gland.[108] Despite these structural similarities, hK2 and total PSA differ in their enzymatic activity. The levels of hK2 in prostate tissue, plasma, semen, and serum are less than 2% that of total PSA, although hK2 mRNA transcript expression represents half that of total PSA. Similar to total PSA, serum hK2 is present in two forms in the blood: one bound to various protease inhibitors, and the other (preponderant) free in the circulation. Several studies have shown that, when used in conjunction with free and total PSA, serum hK2 could improve the discrimination of men with PCa from men without cancer.[109, 110] [111] It has also been suggested that hK2 could predict poor differentiation, extra-capsular extension and biochemical recurrence in patients treated with radical prostatectomy. [112-114] However, this finding has not been validated by other authors.[115] The usefulness of hK2 for the preoperative staging of localized PCa therefore remains controversial. As mentioned above (see "Combination of PSA molecular forms for improved cancer detection" section), the addition of hK2 to three other kallikreins (total, free, and intact PSA) improved the prediction of prostate biopsy results in men with elevated total PSA (increase of predictive accuracy from 68-72% to ~83%).[15] Considering the risk of PCa at 20%, the number of biopsies would have reduced by half, missing only 3 out of 40 high-grade tumors.[15]

EMERGING BLOOD-BASED BIOMARKERS FOR PROSTATE CANCER PROGNOSTICATION

Urokinase Plasminogen Activation (uPA)

The urokinase plasminogen activation (uPA) axis represents a potential target for PCa markers by being involved in various phases of tumor development and progression though degradation of the extra-cellular matrix. The serum protease uPA may play a role in cancer progression by binding to the uPA receptor (uPAR) and consequently converting plasminogen to plasmin, which activates proteases related to the degradation of extracellular matrix proteins.[116] Immunohistochemical staining of radical prostatectomy specimens revealed that overexpression of both uPA and its inhibitor (PAI-1) were associated with aggressive PCa recurrence.[117] In patients with a total PSA level above 2 ng/mL, soluble uPAR and free PSA measured in serum before prostate biopsy improved the regression model accuracy for prediction of PCa.[118] Steuber et al recently showed that uPAR fragments were significant predictors of PCa on biopsy specimens of patients with an elevated PSA.[118]

Both uPA and uPAR might also have a prognostic value. Elevated circulating levels of uPA and uPAR have been linked to PCa stage and bone metastases.[117, 119-122] In a study of 429 patients treated with radical prostatectomy, preoperative plasma uPA was a strong predictor of biochemical recurrence after surgery. Both preoperative uPA and uPAR were associated with features of aggressive biochemical recurrence such as development of distant metastasis suggesting an association with occult metastatic disease at time of local therapy. Moreover, elevation of plasma uPA and uPAR levels in PCa patients seemed to be partly caused by local release from the prostate. Larger multi-institutional studies are under way to validate the potential role of uPA and uPAR as markers of metastatic PCa.

Transforming Growth Factor-Beta 1 (TGF-β1) and Interleukin-6 (IL-6)

TGF-β1 is a growth factor involved in the regulation of several cellular mechanisms including proliferation, immune response, differentiation and angiogenesis.[123] TGF-β1 has been shown to promote cell progression in PCa models and its local expression has been associated with higher tumor grade, tumor invasion and metastasis in PCa patients.[124, 125] Several studies have shown that increased levels of circulating TGF-β1 were associated with cancer progression, occult and documented metastasis and biochemical progression in PCa patients.[125-127]

IL-6 is a cytokine with variable effects on immune and hematopoietic mechanisms. In vitro and in vivo studies have shown that both IL-6 and its receptors (IL-6R) were expressed in PCa.[128, 129] Several authors reported that elevated serum levels of IL-6 and IL-6R were associated with metastatic and hormone refractory disease, and suggested that IL-6 could predict progression and survival of PCa patients.[130, 131]

Based on these findings, Kattan and associates developed and internally validated a prognostic model that incorporates plasma TGF- $\beta1$ and IL-6R into a standard nomogram for prediction of biochemical recurrence following radical prostatectomy.[14] This combination of serum markers and classical clinical parameters improved the predictive accuracy by a statistically and prognostically substantial margin (increase in predictive accuracy from 75 to 84%). However, before a biomarker can become useful in daily clinical management, it needs to be externally validated in an independent cohort of patients (figure 1).[2, 3] Therefore, in a multi-institutional dataset of 423 patients treated with radical prostatectomy, Shariat et al. confirmed that plasma levels of TGF-beta1 and IL6-SR considerably enhance the accuracy of the standard preoperative nomogram for the prediction of biochemical recurrence (accuracy of clinical features plus biomarkers 87.9% versus 71.1% for clinical features alone; p<0.001). Such prognostic models refine our ability to identify patients at a high risk of biochemical recurrence after radical prostatectomy who may benefit from inclusion into perioperative clinical trials and/other intensified follow-up protocols.

Endoglin

Endoglin, or CD 105, is a transmembrane glycoprotein that is typically expressed by human vascular endothelial cells. Functionally, it is a cell-surface coreceptor for transforming growth factor $\beta1$ (TGF- $\beta1$) and $\beta3$ (TGF- $\beta3$) [132] that modulates cellular responses to TGF-β in the early steps of endothelial cell proliferation. Its critical role in angiogenesis has prompted investigators to evaluate the role of Endoglin in cancer progression and metastasis. In PCa, Endoglin is preferentially found on new, immature blood vessels and immunohistochemical analysis supports an association between Endoglin expression and disease progression [133]. Urine levels of Endoglin may distinguish patients with PCa and may help in the staging of the disease. [134] In addition, pre-operative plasma Endoglin were found to be associated with metastasis to regional lymph nodes,[135] established features of biologically aggressive PCa such as higher pathologic Gleason sum, and biochemical recurrence following radical prostatectomy. [136] Use of pre-operative plasma Endoglin could help decide whether and how extensively to perform a lymphadenectomy as well as preoperative identification of patients at risk for disease progression. This would help select patients for neo-adjuvant and/or adjuvant therapy or enrollment into clinical trials. Moreover, Endoglin may be valuable as a surrogate biomarker for occult metastatic disease in patients with presumed organ-confined disease. Further investigation is needed to validate Endoglin as a useful biomarker in men with PCa and to elucidate the mechanistic role of this biomarker in the progression of PCa.

Combination of blood-based biomarkers for prostate cancer prognostication

A biomarker may reflect disruption of a biochemical pathway by a particular mechanism. Given the complexity of the molecular abnormalities associated with PCa, it is improbable that a single marker can accurately segregate tumors of similar clinico-pathologic phenotypes into distinct prognostic categories. Therefore, combinations of independent, yet complementary markers, may provide a more accurate prediction of outcome compared to a single marker.[62]

Shariat et al. found that the addition of a panel comprising pre-operative plasma levels of TGF-β1, soluble IL-6R, IL-6, Endoglin, vascular endothelial growth factor (VEGF), and vascular cell adhesion molecule-1 (VCAM-1 or CD 106)[121, 123, 125, 135-140] improved the predictive accuracy the Kattan pre-operative nomogram[9] by 15.0% (i.e., 71.6 to 86.6%).[11, 103] This increase substantially exceeds accuracy gains obtained from the consideration of detailed pathologic descriptors of prostate cancer at radical prostatectomy. Svatek et al. confirmed the strong predictive value of *pre*-operative levels of the candidate biomarkers after adjusting for the effect of postoperative features.[102] Addition of preoperative levels of the candidate biomarkers improved the accuracy of the base model (i.e., total PSA, surgical margin status, extracapsular extension, seminal vesicles invasion, lymph node involvement, and pathologic Gleason sum) for prediction of biochemical recurrence by a statistically and prognostically significant margin (79% to 86%, p<0.001). Predictive tools integrating biomarker levels could constitute the new standard for counseling patients regarding their risk of recurrence following curative therapy and for designing clinical trials to test neo-adjuvant and/or adjuvant treatment strategies in high-risk patients. However, while prediction of biochemical recurrence is important, prediction of response to therapy as well as metastasis and survival is more important for the management of PCa patients.[141]

BLOOD-BASED BIOMARKERS FOR MONITORING OF PROSTATE CANCER TREATMENT

TOTAL PSA

Treatment monitoring is the most accepted clinical application for total PSA. Total PSA is used for monitoring response to local treatments such as radical prostatectomy, various methods of radiation therapy, and other local therapies including cryosurgery, as well as systemic treatments such as androgen deprivation therapy and chemotherapy. Post-treatment total PSA levels can provide invaluable information about the effectiveness of the therapy given and the existence of residual cancer in men treated with local therapy with curative intent. In such patients, rising total PSA levels can signal cancer activity well before any clinical signs of recurrence appear. This lead-time can be further increased by months and even years.

In patients treated with systemic therapy, post-therapy total PSA changes are a seemingly perfect outcome measure because they are easily assessable, quantitative, reproducible, and inexpensive; this is true regardless of whether this outcome measure is applied to evaluate drugs in the clinical trial setting or in clinical practice. Critical to the successful application of total PSA measurement as an endpoint are the therapeutic objectives of the trial and the mechanisms of action of the treatment administered. Criteria were proposed to screen for treatment effects in prostate cancer clinical trials on the basis of the hypothesis that total PSA declines reflect significant cell kill in response to agents that cause reduction in overall tumor burden. With the recognition that short-term declines in total PSA levels might simply reflect the effect of the drug on the marker, and not on cell growth or survival, it is recommended that the declines be documented on more than one occasion and, equally importantly, over time. Indeed, different total PSA-based outcomes would be required for

different classes of drug. For example, drugs that are not anticipated to kill cells would not be expected to produce declines in total PSA. Similarly, 'differentiating effect' might produce an initial rise in total PSA that might be an indication that the drug is actually working. A single set of outcomes would not only be inapplicable to agents that act via diverse mechanisms, but could also be misleading.

Unfortunately, the significance of post-therapy changes in total PSA have been misinterpreted as indicators of tumor response, and/or as a measure of clinical benefit. An additional misconception is that the demonstration of a particular degree of decline in total PSA, in a proportion of patients, is an indication that the treatment prolongs life, and as such, should be used as an endpoint for accelerated drug approval.

The use of a post-therapy decline in total PSA as an outcome measure has been justified in part by statistical analyses exploring the relationship between the defined outcome measure and survival.[142] Most of these reports include multivariate techniques, and in some, the resultant models were validated on independent data sets. These early reports did not assess randomized comparative trials showing a survival benefit, a necessary but not sufficient condition in which to explore measures of surrogacy.[142, 143]

There is no clear demonstration that a post-therapy total PSA change can account for all of the treatment effects seen is not yet available. Large-scale prospective studies incorporating different post-therapy total PSA change definitions, as well as other potential biomarkers, are ongoing. At this point, one can conclude that a cytotoxic drug that does not produce any total PSA declines is unlikely to be effective, but that the corollary is not always true. In addition, it is important to recognize that there is a range of clinical benefits to patients that can improve the quality, and more than likely, the duration of survival, which are independent of total PSA levels. This includes differentiating agents that might produce a rise in total PSA before a decline is observed, or act through immunization strategies where the effect on total PSA levels might be delayed or not occur at all, and angiogenesis and growth factor inhibitors. It is also important to note that there remains significant variation in the methodologies and assays used by different laboratories, which makes it difficult to compare results between groups. All of these factors must be considered carefully to ensure that a drug is not discontinued prematurely on the basis of a total PSA rise that is not relevant to the drug under study.

CONCLUSIONS

The introduction of total PSA in clinical practice has resulted in early detection and reduced mortality from PCa.[144] However, PCa screening remains controversial, because of the risk of overdiagnosis reduced mortality and overtreatment and the inability to detect a significant proportion of dangerous tumors. A large concerted effort has been made to improve and/or monitoring the activity of PCa and to guide molecular targeted therapy and/or assess therapeutic response. An integrated approach with blood-based measurement of different molecular forms of PSA in combination with genetic and urine biomarkers hold the promise of improving screening for and diagnosis of PCa. Panels of blood-based biomarkers will allow a fingerprinting of the tumors biologic behavior resulting in individualized therapy and monitoring. In addition, the emergence of new therapeutic approaches for PCa cannot flourish without a set of markers to serve as prognosticators, predictors, therapeutic targets, and/or surrogate end points.

REFERENCES

[1]. Jemal A, Siegel R, Ward E, Hao Y, Xu J, Thun MJ. Cancer statistics, 2009. CA Cancer J Clin. Jul-Aug;2009 59(4):225–49. [PubMed: 19474385]

[2]. Bensalah K, Montorsi F, Shariat SF. Challenges of cancer biomarker profiling. Eur Urol. Dec; 2007 52(6):1601–9. [PubMed: 17919807]

- [3]. Shariat SF, Canto EI, Kattan MW, Slawin KM. Beyond prostate-specific antigen: new serologic biomarkers for improved diagnosis and management of prostate cancer. Rev Urol. Spring;2004 6(2):58–72. [PubMed: 16985579]
- [4]. Bensalah K, Lotan Y, Karam JA, Shariat SF. New circulating biomarkers for prostate cancer. Prostate Cancer Prostatic Dis. Nov 13.2007
- [5]. Verma M, Srivastava S. New cancer biomarkers deriving from NCI early detection research. Recent Results Cancer Res. 2003; 163:72–84. discussion 264-6. [PubMed: 12903844]
- [6]. Winget MD, Baron JA, Spitz MR, Brenner DE, Warzel D, Kincaid H, et al. Development of common data elements: the experience of and recommendations from the early detection research network. Int J Med Inform. Apr; 2003 70(1):41–8. [PubMed: 12706181]
- [7]. Ilyin SE, Belkowski SM, Plata-Salaman CR. Biomarker discovery and validation: technologies and integrative approaches. Trends in biotechnology. Aug; 2004 22(8):411–6. [PubMed: 15283986]
- [8]. Shariat SF, Karakiewicz PI. Perspectives on prostate cancer biomarkers. Eur Urol. Jul; 2008 54(1): 8–10. [PubMed: 18262336]
- [9]. Shariat SF, Karakiewicz PI, Roehrborn CG, Kattan MW. An updated catalog of prostate cancer predictive tools. Cancer. Dec 1; 2008 113(11):3075–99. [PubMed: 18823041]
- [10]. Shariat SF, Karam JA, Margulis V, Karakiewicz PI. New blood-based biomarkers for the diagnosis, staging and prognosis of prostate cancer. BJU Int. Mar; 2008 101(6):675–83. [PubMed: 17941930]
- [11]. Shariat SF, Karam JA, Walz J, Roehrborn CG, Montorsi F, Margulis V, et al. Improved prediction of disease relapse after radical prostatectomy through a panel of preoperative blood-based biomarkers. Clin Cancer Res. Jun 15; 2008 14(12):3785–91. [PubMed: 18559597]
- [12]. Shariat SF, Park S, Trinh QD, Roehrborn CG, Slawin KM, Karakiewicz PI. Plasminogen activation inhibitor-1 improves the predictive accuracy of prostate cancer nomograms. J Urol. Oct; 2007 178(4 Pt 1):1229–36. discussion 36-7. [PubMed: 17698116]
- [13]. Kattan MW. Judging new markers by their ability to improve predictive accuracy. J Natl Cancer Inst. May 7; 2003 95(9):634–5. [PubMed: 12734304]
- [14]. Kattan MW, Shariat SF, Andrews B, Zhu K, Canto E, Matsumoto K, et al. The addition of interleukin-6 soluble receptor and transforming growth factor beta1 improves a preoperative nomogram for predicting biochemical progression in patients with clinically localized prostate cancer. J Clin Oncol. Oct 1; 2003 21(19):3573–9. [PubMed: 12913106]
- [15]. Vickers AJ, Cronin AM, Aus G, Pihl CG, Becker C, Pettersson K, et al. A panel of kallikrein markers can reduce unnecessary biopsy for prostate cancer: data from the European Randomized Study of Prostate Cancer Screening in Goteborg, Sweden. BMC medicine. 2008; 6:19. [PubMed: 18611265]
- [16]. Vickers AJ, Jang K, Sargent D, Lilja H, Kattan MW. Systematic review of statistical methods used in molecular marker studies in cancer. Cancer. Apr 15; 2008 112(8):1862–8. [PubMed: 18320601]
- [17]. Vickers AJ, Elkin EB. Decision curve analysis: a novel method for evaluating prediction models. Med Decis Making. Nov-Dec;2006 26(6):565–74. [PubMed: 17099194]
- [18]. Vickers AJ, Cronin AM, Elkin EB, Gonen M. Extensions to decision curve analysis, a novel method for evaluating diagnostic tests, prediction models and molecular markers. BMC Med Inform Decis Mak. 2008; 8:53. [PubMed: 19036144]
- [19]. Vickers AJ. Decision analysis for the evaluation of diagnostic tests, prediction models and molecular markers. Am Stat. 2008; 62(4):314–20. [PubMed: 19132141]
- [20]. Elkin EB, Vickers AJ, Kattan MW. Primer: using decision analysis to improve clinical decision making in urology. Nat Clin Pract Urol. Aug; 2006 3(8):439–48. [PubMed: 16902520]
- [21]. Ankerst DP, Miyamoto R, Nair PV, Pollock BH, Thompson IM, Parekh DJ. Yearly prostate specific antigen and digital rectal examination fluctuations in a screened population. J Urol. May; 2009 181(5):2071–5. discussion 6. [PubMed: 19286205]

[22]. Soletormos G, Semjonow A, Sibley PE, Lamerz R, Petersen PH, Albrecht W, et al. Biological variation of total prostate-specific antigen: a survey of published estimates and consequences for clinical practice. Clin Chem. Aug; 2005 51(8):1342–51. [PubMed: 15961552]

- [23]. Roehrborn CG, Pickens GJ, Carmody T 3rd. Variability of repeated serum prostate-specific antigen (PSA) measurements within less than 90 days in a well-defined patient population. Urology. Jan; 1996 47(1):59–66. [PubMed: 8560664]
- [24]. Link RE, Shariat SF, Nguyen CV, Farr A, Weinberg AD, Morton RA, et al. Variation in prostate specific antigen results from 2 different assay platforms: clinical impact on 2304 patients undergoing prostate cancer screening. J Urol. Jun; 2004 171(6 Pt 1):2234–8. [PubMed: 15126793]
- [25]. Sotelo RJ, Mora KE, Perez LH, Novoa J, Carmona O, De Andrade R, et al. Assay standardization bias: different prostate cancer detection rates and clinical outcomes resulting from different assays for free and total prostate-specific antigen. Urology. Jun; 2007 69(6):1143–6. [PubMed: 17572203]
- [26]. Stephan C, Klaas M, Muller C, Schnorr D, Loening SA, Jung K. Interchangeability of measurements of total and free prostate-specific antigen in serum with 5 frequently used assay combinations: an update. Clin Chem. Jan; 2006 52(1):59–64. [PubMed: 16391327]
- [27]. Slev PR, La'ulu SL, Roberts WL. Intermethod differences in results for total PSA, free PSA, and percentage of free PSA. American journal of clinical pathology. Jun; 2008 129(6):952–8. [PubMed: 18480013]
- [28]. Stephan C, Bangma C, Vignati G, Bartsch G, Lein M, Jung K, et al. 20-25% lower concentrations of total and free prostate-specific antigen (PSA) after calibration of PSA assays to the WHO reference materials--analysis of 1098 patients in four centers. The International journal of biological markers. Apr-Jun;2009 24(2):65–9. [PubMed: 19634108]
- [29]. Semjonow A, Brandt B, Oberpenning F, Roth S, Hertle L. Discordance of assay methods creates pitfalls for the interpretation of prostate-specific antigen values. The Prostate. 1996; 7:3–16.
 [PubMed: 8950358]
- [30]. Thompson IM, Chi C, Ankerst DP, Goodman PJ, Tangen CM, Lippman SM, et al. Effect of finasteride on the sensitivity of PSA for detecting prostate cancer. Journal of the National Cancer Institute. Aug 16; 2006 98(16):1128–33. [PubMed: 16912265]
- [31]. Nixon RG, Wener MH, Smith KM, Parson RE, Blase AB, Brawer MK. Day to day changes in free and total PSA: significance of biological variation. Prostate cancer and prostatic diseases. Dec; 1997 1(2):90–6. [PubMed: 12496922]
- [32]. Nixon RG, Wener MH, Smith KM, Parson RE, Strobel SA, Brawer MK. Biological variation of prostate specific antigen levels in serum: an evaluation of day-to-day physiological fluctuations in a well-defined cohort of 24 patients. J Urol. Jun; 1997 157(6):2183–90. [PubMed: 9146611]
- [33]. Bunting PS, DeBoer G, Choo R, Danjoux C, Klotz L, Fleshner N. Intra-individual variation of PSA, free PSA and complexed PSA in a cohort of patients with prostate cancer managed with watchful observation. Clin Biochem. Sep; 2002 35(6):471–5. [PubMed: 12413608]
- [34]. Bruun L, Becker C, Hugosson J, Lilja H, Christensson A. Assessment of intra-individual variation in prostate-specific antigen levels in a biennial randomized prostate cancer screening program in Sweden. Prostate. Nov 1; 2005 65(3):216–21. [PubMed: 15948137]
- [35]. Birkenmeier G, Struck F, Gebhardt R. Clearance mechanism of prostate specific antigen and its complexes with alpha2-macroglobulin and alpha1-antichymotrypsin. J Urol. Sep; 1999 162(3 Pt 1):897–901. [PubMed: 10458404]
- [36]. Bjork T, Ljungberg B, Piironen T, Abrahamsson PA, Pettersson K, Cockett AT, et al. Rapid exponential elimination of free prostate-specific antigen contrasts the slow, capacity-limited elimination of PSA complexed to alpha 1-antichymotrypsin from serum. Urology. Jan; 1998 51(1):57–62. [PubMed: 9457289]
- [37]. Lilja H, Haese A, Bjork T, Friedrich MG, Piironen T, Pettersson K, et al. Significance and metabolism of complexed and noncomplexed prostate specific antigen forms, and human glandular kallikrein 2 in clinically localized prostate cancer before and after radical prostatectomy. J Urol. Dec; 1999 162(6):2029–34. discussion 34-5. [PubMed: 10569562]

[38]. Djavan B, Shariat S, Ghawidel K, Guven-Marberger K, Remzi M, Kovarik J, et al. Impact of chronic dialysis on serum PSA, free PSA, and free/total PSA ratio: is prostate cancer detection compromised in patients receiving long-term dialysis?PG - 1169-74. Urology. Jun.1999 53(6)

- [39]. Thompson IM, Pauler DK, Goodman PJ, Tangen CM, Lucia MS, Parnes HL, et al. Prevalence of prostate cancer among men with a prostate-specific antigen level < or =4.0 ng per milliliter. The New England journal of medicine. May 27; 2004 350(22):2239–46. [PubMed: 15163773]
- [40]. Thompson IM, Ankerst DP, Chi C, Goodman PJ, Tangen CM, Lucia MS, et al. Assessing prostate cancer risk: results from the Prostate Cancer Prevention Trial. J Natl Cancer Inst. Apr 19; 2006 98(8):529–34. [PubMed: 16622122]
- [41]. Thompson IM, Ankerst DP, Chi C, Lucia MS, Goodman PJ, Crowley JJ, et al. Operating characteristics of prostate-specific antigen in men with an initial PSA level of 3.0 ng/ml or lower. Jama. Jul 6; 2005 294(1):66–70. [PubMed: 15998892]
- [42]. Thompson IM, Pauler Ankerst D, Chi C, Goodman PJ, Tangen CM, Lippman SM, et al. Prediction of prostate cancer for patients receiving finasteride: results from the Prostate Cancer Prevention Trial. J Clin Oncol. Jul 20; 2007 25(21):3076–81. [PubMed: 17634486]
- [43]. Berger AP, Volgger H, Rogatsch H, Strohmeyer D, Steiner H, Klocker H, et al. Screening with low PSA cutoff values results in low rates of positive surgical margins in radical prostatectomy specimens. Prostate. Nov 1; 2002 53(3):241–5. [PubMed: 12386925]
- [44]. Catalona WJ, Ramos CG, Carvalhal GF, Yan Y. Lowering PSA cutoffs to enhance detection of curable prostate cancer. Urology. Jun; 2000 55(6):791–5. [PubMed: 10840078]
- [45]. Catalona WJ, Smith DS, Ornstein DK. Prostate cancer detection in men with serum PSA concentrations of 2.6 to 4.0 ng/mL and benign prostate examination. Enhancement of specificity with free PSA measurements. Jama. May 14; 1997 277(18):1452–5. [PubMed: 9145717]
- [46]. Efstathiou JA, Chen MH, Catalona WJ, McLeod DG, Carroll PR, Moul JW, et al. Prostate-specific antigen-based serial screening may decrease prostate cancer-specific mortality. Urology. Aug; 2006 68(2):342–7. [PubMed: 16904449]
- [47]. Shariat SF, Karakiewicz PI, Margulis V, Kattan MW. Inventory of prostate cancer predictive tools. Current opinion in urology. May; 2008 18(3):279–96. [PubMed: 18382238]
- [48]. Kattan MW. When and how to use informatics tools in caring for urologic patients. Nature clinical practice. Apr; 2005 2(4):183–90.
- [49]. Kattan MW, Eastham JA, Wheeler TM, Maru N, Scardino PT, Erbersdobler A, et al. Counseling men with prostate cancer: a nomogram for predicting the presence of small, moderately differentiated, confined tumors. J Urol. Nov; 2003 170(5):1792–7. [PubMed: 14532778]
- [50]. Steyerberg EW, Roobol MJ, Kattan MW, van der Kwast TH, de Koning HJ, Schroder FH. Prediction of indolent prostate cancer: validation and updating of a prognostic nomogram. J Urol. Jan; 2007 177(1):107–12. discussion 12. [PubMed: 17162015]
- [51]. Nam RK, Toi A, Klotz LH, Trachtenberg J, Jewett MA, Appu S, et al. Assessing individual risk for prostate cancer. J Clin Oncol. Aug 20; 2007 25(24):3582–8. [PubMed: 17704405]
- [52]. Stenman UH, Hakama M, Knekt P, Aromaa A, Teppo L, Leinonen J. Serum concentrations of prostate specific antigen and its complex with alpha 1-antichymotrypsin before diagnosis of prostate cancer. Lancet. 1994; 344(8937):1594–8. 1994. [PubMed: 7527116]
- [53]. Gann PH, Hennekens CH, Stampfer MJ. A prospective evaluation of plasma prostate-specific antigen for detection of prostatic cancer [see comments]. Jama. 1995; 273(4):289–94. [PubMed: 7529341]
- [54]. Fang J, Metter EJ, Landis P, Chan DW, Morrell CH, Carter HB. Low levels of prostate-specific antigen predict long-term risk of prostate cancer: results from the Baltimore Longitudinal Study of Aging. Urology. Sep; 2001 58(3):411–6. [PubMed: 11549490]
- [55]. Loeb S, Roehl KA, Antenor JA, Catalona WJ, Suarez BK, Nadler RB. Baseline prostate-specific antigen compared with median prostate-specific antigen for age group as predictor of prostate cancer risk in men younger than 60 years old. Urology. Feb; 2006 67(2):316–20. [PubMed: 16442597]
- [56]. Lilja H, Ulmert D, Bjork T, Becker C, Serio AM, Nilsson JA, et al. Long-term prediction of prostate cancer up to 25 years before diagnosis of prostate cancer using prostate kallikreins measured at age 44 to 50 years. J Clin Oncol. Feb 1; 2007 25(4):431–6. [PubMed: 17264339]

[57]. Ulmert D, Cronin AM, Bjork T, O'Brien MF, Scardino PT, Eastham JA, et al. Prostate-specific antigen at or before age 50 as a predictor of advanced prostate cancer diagnosed up to 25 years later: a case-control study. BMC medicine. 2008; 6:6. [PubMed: 18279502]

- [58]. Vickers AJ, Ulmert D, Serio AM, Bjork T, Scardino PT, Eastham JA, et al. The predictive value of prostate cancer biomarkers depends on age and time to diagnosis: towards a biologically-based screening strategy. Int J Cancer. Nov 15; 2007 121(10):2212–7. [PubMed: 17657743]
- [59]. Vickers AJ, Cronin AM, Bjork T, Manjer J, Nilsson PM, Dahlin A, et al. Prostate specific antigen concentration at age 60 and death or metastasis from prostate cancer: case-control study. Bmj. 341:c4521.
- [60]. Lilja H, Ulmert D, Vickers AJ. Prostate-specific antigen and prostate cancer: prediction, detection and monitoring. Nat Rev Cancer. Apr; 2008 8(4):268–78. [PubMed: 18337732]
- [61]. Schroder FH, Carter HB, Wolters T, van den Bergh RC, Gosselaar C, Bangma CH, et al. Early detection of prostate cancer in 2007. Part 1: PSA and PSA kinetics. Eur Urol. Mar; 2008 53(3): 468–77. [PubMed: 17997011]
- [62]. Shariat SF, Karam JA, Roehrborn CG. Blood biomarkers for prostate cancer detection and prognosis. Future Oncol. Aug; 2007 3(4):449–61. [PubMed: 17661720]
- [63]. Carter HB, Morrell CH, Pearson JD, Brant LJ, Plato CC, Metter EJ, et al. Estimation of prostatic growth using serial prostate-specific antigen measurements in men with and without prostate disease. Cancer research. Jun 15; 1992 52(12):3323–8. [PubMed: 1375867]
- [64]. Carter HB, Pearson JD, Metter EJ, Brant LJ, Chan DW, Andres R, et al. Longitudinal evaluation of prostate-specific antigen levels in men with and without prostate disease [see comments]. Journal of the American Medical Association. 1992; 267(16):2215–20. 1992. [PubMed: 1372942]
- [65]. Connolly D, Black A, Murray LJ, Napolitano G, Gavin A, Keane PF. Methods of calculating prostate-specific antigen velocity. Eur Urol. Oct; 2007 52(4):1044–50. [PubMed: 17197071]
- [66]. Smith DS, Catalona WJ. Rate of change in serum prostate specific antigen levels as a method for prostate cancer detection. J Urol. Oct; 1994 152(4):1163–7. [PubMed: 7520949]
- [67]. Carter HB, Pearson JD, Waclawiw Z, Metter EJ, Chan DW, Guess HA, et al. Prostate-specific antigen variability in men without prostate cancer: effect of sampling interval on prostate-specific antigen velocity. Urology. Apr; 1995 45(4):591–6. [PubMed: 7536366]
- [68]. Roobol MJ, Kranse R, de Koning HJ, Schroder FH. Prostate-specific antigen velocity at low prostate-specific antigen levels as screening tool for prostate cancer: results of second screening round of ERSPC (ROTTERDAM). Urology. Feb; 2004 63(2):309–13. discussion 13-5. [PubMed: 14972478]
- [69]. Schroder FH, Roobol MJ, van der Kwast TH, Kranse R, Bangma CH. Does PSA velocity predict prostate cancer in pre-screened populations? Eur Urol. Mar; 2006 49(3):460–5. discussion 5. [PubMed: 16442212]
- [70]. Pinsky PF, Andriole G, Crawford ED, Chia D, Kramer BS, Grubb R, et al. Prostate-specific antigen velocity and prostate cancer gleason grade and stage. Cancer. Apr 15; 2007 109(8):1689–95. [PubMed: 17330856]
- [71]. Ulmert D, Serio AM, O'Brien MF, Becker C, Eastham JA, Scardino PT, et al. Long-term prediction of prostate cancer: prostate-specific antigen (PSA) velocity is predictive but does not improve the predictive accuracy of a single PSA measurement 15 years or more before cancer diagnosis in a large, representative, unscreened population. J Clin Oncol. Feb 20; 2008 26(6): 835–41. [PubMed: 18281654]
- [72]. D'Amico AV, Chen MH, Roehl KA, Catalona WJ. Preoperative PSA velocity and the risk of death from prostate cancer after radical prostatectomy. The New England journal of medicine. Jul 8; 2004 351(2):125–35. [PubMed: 15247353]
- [73]. D'Amico AV, Renshaw AA, Sussman B, Chen MH. Pretreatment PSA velocity and risk of death from prostate cancer following external beam radiation therapy. Jama. Jul 27; 2005 294(4):440– 7. [PubMed: 16046650]
- [74]. Patel DA, Presti JC Jr. McNeal JE, Gill H, Brooks JD, King CR. Preoperative PSA velocity is an independent prognostic factor for relapse after radical prostatectomy. J Clin Oncol. Sep 1; 2005 23(25):6157–62. [PubMed: 16135482]

[75]. Carter HB, Ferrucci L, Kettermann A, Landis P, Wright EJ, Epstein JI, et al. Detection of lifethreatening prostate cancer with prostate-specific antigen velocity during a window of curability. Journal of the National Cancer Institute. Nov 1; 2006 98(21):1521–7. [PubMed: 17077354]

- [76]. National Comprehensive Cancer Network. National Comprehensive, 2004. 2004. http://www.nccn.org/professionals/physician_gls/PDF/prostate_detection.pdf
- [77]. Vickers A, Tangen C, Till C, Lilja H, Thompson I. An empirical evaluation of the national cancer center network guidelines on prostate specific antigen velocity in prostate cancer detection. J Urol. 2010; 183(4):e771.
- [78]. Fall K, Garmo H, Andren O, Bill-Axelson A, Adolfsson J, Adami HO, et al. Prostate-specific antigen levels as a predictor of lethal prostate cancer. Journal of the National Cancer Institute. Apr 4; 2007 99(7):526–32. [PubMed: 17405997]
- [79]. Stephenson AJ, Kattan MW, Eastham JA, Bianco FJ Jr. Yossepowitch O, Vickers AJ, et al. Prostate cancer-specific mortality after radical prostatectomy for patients treated in the prostate-specific antigen era. J Clin Oncol. Sep 10; 2009 27(26):4300–5. [PubMed: 19636023]
- [80]. O'Brien MF, Cronin AM, Fearn PA, Smith B, Stasi J, Guillonneau B, et al. Pretreatment prostate-specific antigen (PSA) velocity and doubling time are associated with outcome but neither improves prediction of outcome beyond pretreatment PSA alone in patients treated with radical prostatectomy. J Clin Oncol. Aug 1; 2009 27(22):3591–7. [PubMed: 19506163]
- [81]. Frank O'Brien M, Cronin AM, Fearn PA, Savage CJ, Smith B, Stasi J, et al. Evaluation of prediagnostic prostate-specific antigen dynamics as predictors of death from prostate cancer in patients treated conservatively. Int J Cancer. Jul 23.
- [82]. Yu X, Loeb S, Roehl KA, Han M, Catalona WJ. The association between total prostate specific antigen concentration and prostate specific antigen velocity. J Urol. Apr; 2007 177(4):1298–302. discussion 301-2. [PubMed: 17382716]
- [83]. Lilja H. Significance of different molecular forms of serum PSA. The free, noncomplexed form of PSA versus that complexed to alpha 1-antichymotrypsin. Urologic Clinics of North America. 1993; 20(4):681–6. [Review]. 1993. [PubMed: 7505976]
- [84]. Shariat SF, Abdel-Aziz KF, Roehrborn CG, Lotan Y. Pre-operative percent free PSA predicts clinical outcomes in patients treated with radical prostatectomy with total PSA levels below 10 ng/ml. Eur Urol. Feb; 2006 49(2):293–302. [PubMed: 16387412]
- [85]. Catalona WJ, Southwick PC, Slawin KM, Partin AW, Brawer MK, Flanigan RC, et al. Comparison of percent free PSA, PSA density, and age-specific PSA cutoffs for prostate cancer detection and staging. Urology. Aug 1; 2000 56(2):255–60. [PubMed: 10925089]
- [86]. Catalona WJ, Partin AW, Slawin KM, Brawer MK, Flanigan RC, Patel A, et al. Use of the percentage of free prostate-specific antigen to enhance differentiation of prostate cancer from benign prostatic disease: a prospective multicenter clinical trial. Jama. May 20; 1998 279(19): 1542–7. [PubMed: 9605898]
- [87]. Woodrum DL, Brawer MK, Partin AW, Catalona WJ, Southwick PC. Interpretation of free prostate specific antigen clinical research studies for the detection of prostate cancer. J Urol. Jan; 1998 159(1):5–12. [PubMed: 9400426]
- [88]. Canto EI, Singh H, Shariat SF, Kadmon D, Miles BJ, Wheeler TM, et al. Effects of systematic 12-core biopsy on the performance of percent free prostate specific antigen for prostate cancer detection. J Urol. Sep; 2004 172(3):900–4. [PubMed: 15310993]
- [89]. Graefen M, Karakiewicz PI, Cagiannos I, Hammerer PG, Haese A, Palisaar J, et al. Percent free prostate specific antigen is not an independent predictor of organ confinement or prostate specific antigen recurrence in unscreened patients with localized prostate cancer treated with radical prostatectomy. J Urol. Mar; 2002 167(3):1306–9. [PubMed: 11832719]
- [90]. Graefen M, Karakiewicz PI, Cagiannos I, Hammerer PG, Haese A, Palisaar J, et al. Percent free prostate specific antigen is not an independent predictor of organ confinement or prostate specific antigen recurrence in unscreened patients with localized prostate cancer treated with radical prostatectomy. J Urol. 2002; 167(3):1306–9. [PubMed: 11832719]
- [91]. Mikolajczyk SD, Marks LS, Partin AW, Rittenhouse HG. Free prostate-specific antigen in serum is becoming more complex. Urology. Jun; 2002 59(6):797–802. [PubMed: 12031356]

[92]. Mikolajczyk SD, Marker KM, Millar LS, Kumar A, Saedi MS, Payne JK, et al. A truncated precursor form of prostate-specific antigen is a more specific serum marker of prostate cancer. Cancer Res. 2001; 61(18):6958–63. [PubMed: 11559576]

- [93]. Sokoll LJ, Chan DW, Mikolajczyk SD, Rittenhouse HG, Evans CL, Linton HJ, et al. Proenzyme psa for the early detection of prostate cancer in the 2.5-4.0 ng/ml total psa range: preliminary analysis. Urology. Feb; 2003 61(2):274–6. [PubMed: 12597929]
- [94]. Catalona WJ, Bartsch G, Rittenhouse HG, Evans CL, Linton HJ, Horninger W, et al. Serum proprostate specific antigen preferentially detects aggressive prostate cancers in men with 2 to 4 ng/ml prostate specific antigen. J Urol. Jun; 2004 171(6 Pt 1):2239–44. [PubMed: 15126794]
- [95]. Catalona WJ, Bartsch G, Rittenhouse HG, Evans CL, Linton HJ, Amirkhan A, et al. Serum pro prostate specific antigen improves cancer detection compared to free and complexed prostate specific antigen in men with prostate specific antigen 2 to 4 ng/ml. J Urol. Dec; 2003 170(6 Pt 1): 2181–5. [PubMed: 14634374]
- [96]. Stephan C, Kahrs AM, Cammann H, Lein M, Schrader M, Deger S, et al. A [–2]proPSA-based artificial neural network significantly improves differentiation between prostate cancer and benign prostatic diseases. Prostate. Feb 1; 2009 69(2):198–207. [PubMed: 18942119]
- [97]. Makarov DV, Isharwal S, Sokoll LJ, Landis P, Marlow C, Epstein JI, et al. Pro-prostate-specific antigen measurements in serum and tissue are associated with treatment necessity among men enrolled in expectant management for prostate cancer. Clin Cancer Res. Dec 1; 2009 15(23): 7316–21. [PubMed: 19934305]
- [98]. Semjonow A, Kopke T, Eltze E, Pepping-Schefers B, Burgel H, Darte C. Pre-analytical in-vitro stability of [–2]proPSA in blood and serum. Clinical biochemistry. Jul; 43(10-11):926–8. [PubMed: 20450900]
- [99]. Mikolajczyk SD, Millar LS, Wang TJ, Rittenhouse HG, Wolfert RL, Marks LS, et al. "BPSA," a specific molecular form of free prostate-specific antigen, is found predominantly in the transition zone of patients with nodular benign prostatic hyperplasia. Urology. Jan; 2000 55(1):41–5. [PubMed: 10654892]
- [100]. Canto EI, Singh H, Shariat SF, Lamb DJ, Mikolajczyk SD, Linton HJ, et al. Serum BPSA outperforms both total PSA and free PSA as a predictor of prostatic enlargement in men without prostate cancer. Urology. May; 2004 63(5):905–10. discussion 10-1. [PubMed: 15134977]
- [101]. Stephan C, Cammann H, Deger S, Schrader M, Meyer HA, Miller K, et al. Benign prostatic hyperplasia-associated free prostate-specific antigen improves detection of prostate cancer in an artificial neural network. Urology. Oct; 2009 74(4):873–7. [PubMed: 19476981]
- [102]. Svatek RS, Jeldres C, Karakiewicz PI, Suardi N, Walz J, Roehrborn CG, et al. Pre-treatment biomarker levels improve the accuracy of post-prostatectomy nomogram for prediction of biochemical recurrence. Prostate. Jun 1; 2009 69(8):886–94. [PubMed: 19229851]
- [103]. Shariat SF, Karakiewicz PI, Ashfaq R, Lerner SP, Palapattu GS, Cote RJ, et al. Multiple biomarkers improve prediction of bladder cancer recurrence and mortality in patients undergoing cystectomy. Cancer. Jan 15; 2008 112(2):315–25. [PubMed: 18008359]
- [104]. Vickers A, Cronin A, Roobol M, Savage C, Peltola M, Pettersson K, et al. Reducing unnecessary biopsy during prostate cancer screening using a four-kallikrein panel: an independent replication. J Clin Oncol. May 20; 28(15):2493–8. [PubMed: 20421547]
- [105]. Vickers AJ, Cronin AM, Aus G, Pihl CG, Becker C, Pettersson K, et al. Impact of recent screening on predicting the outcome of prostate cancer biopsy in men with elevated prostate-specific antigen: data from the European Randomized Study of Prostate Cancer Screening in Gothenburg, Sweden. Cancer. Jun 1; 116(11):2612–20. [PubMed: 20336781]
- [106]. Vickers AJ, Cronin AM, Roobol MJ, Savage CJ, Peltola M, Pettersson K, et al. A four-kallikrein panel predicts prostate cancer in men with recent screening: data from the European Randomized Study of Screening for Prostate Cancer, Rotterdam. Clin Cancer Res. Jun 15; 16(12):3232–9. [PubMed: 20400522]
- [107]. Gupta A, Roobol MJ, Savage CJ, Peltola M, Pettersson K, Scardino PT, et al. A four-kallikrein panel for the prediction of repeat prostate biopsy: data from the European Randomized Study of Prostate Cancer Screening in Rotterdam, Netherlands. British journal of cancer. Jul 27.

[108]. Yousef GM, Diamandis EP. The new human tissue kallikrein gene family: structure, function, and association to disease. Endocr Rev. Apr; 2001 22(2):184–204. [PubMed: 11294823]

- [109]. Nam RK, Diamandis EP, Toi A, Trachtenberg J, Magklara A, Scorilas A, et al. Serum human glandular kallikrein-2 protease levels predict the presence of prostate cancer among men with elevated prostate-specific antigen. J Clin Oncol. Mar; 2000 18(5):1036–42. [PubMed: 10694554]
- [110]. Becker C, Piironen T, Pettersson K, Hugosson J, Lilja H. Clinical value of human glandular kallikrein 2 and free and total prostate-specific antigen in serum from a population of men with prostate-specific antigen levels 3.0 ng/mL or greater. Urology. May; 2000 55(5):694–9. [PubMed: 10792083]
- [111]. Becker C, Piironen T, Pettersson K, Bjork T, Wojno KJ, Oesterling JE, et al. Discrimination of men with prostate cancer from those with benign disease by measurements of human glandular kallikrein 2 (HK2) in serum. J Urol. Jan; 2000 163(1):311–6. [PubMed: 10604382]
- [112]. Haese A, Graefen M, Steuber T, Becker C, Pettersson K, Piironen T, et al. Human glandular kallikrein 2 levels in serum for discrimination of pathologically organ-confined from locally-advanced prostate cancer in total PSA-levels below 10 ng/ml. Prostate. Oct 1; 2001 49(2):101–9. [PubMed: 11582588]
- [113]. Kwiatkowski MK, Recker F, Piironen T, Pettersson K, Otto T, Wernli M, et al. In prostatism patients the ratio of human glandular kallikrein to free PSA improves the discrimination between prostate cancer and benign hyperplasia within the diagnostic "gray zone" of total PSA 4 to 10 ng/mL. Urology. Sep; 1998 52(3):360–5. [PubMed: 9730444]
- [114]. Recker F, Kwiatkowski MK, Piironen T, Pettersson K, Lummen G, Wernli M, et al. The importance of human glandular kallikrein and its correlation with different prostate specific antigen serum forms in the detection of prostate carcinoma. Cancer. Dec 15; 1998 83(12):2540– 7. [PubMed: 9874461]
- [115]. Kurek R, Nunez G, Tselis N, Konrad L, Martin T, Roeddiger S, et al. Prognostic value of combined "triple"-reverse transcription-PCR analysis for prostate-specific antigen, human kallikrein 2, and prostate-specific membrane antigen mRNA in peripheral blood and lymph nodes of prostate cancer patients. Clin Cancer Res. Sep 1; 2004 10(17):5808–14. [PubMed: 15355910]
- [116]. Duffy MJ. Urokinase-type plasminogen activator: a potent marker of metastatic potential in human cancers. Biochem Soc Trans. Apr; 2002 30(2):207–10. [PubMed: 12023852]
- [117]. Gupta A, Lotan Y, Ashfaq R, Roehrborn CG, Raj GV, Aragaki CC, et al. Predictive value of the differential expression of the urokinase plasminogen activation axis in radical prostatectomy patients. Eur Urol. May; 2009 55(5):1124–33. [PubMed: 18585843]
- [118]. Steuber T, Vickers A, Haese A, Kattan MW, Eastham JA, Scardino PT, et al. Free PSA isoforms and intact and cleaved forms of urokinase plasminogen activator receptor in serum improve selection of patients for prostate cancer biopsy. Int J Cancer. Apr 1; 2007 120(7):1499–504. [PubMed: 17205511]
- [119]. Hienert G, Kirchheimer JC, Pfluger H, Binder BR. Urokinase-type plasminogen activator as a marker for the formation of distant metastases in prostatic carcinomas. J Urol. Dec; 1988 140(6): 1466–9. [PubMed: 3193516]
- [120]. Miyake H, Hara I, Yamanaka K, Gohji K, Arakawa S, Kamidono S. Elevation of serum levels of urokinase-type plasminogen activator and its receptor is associated with disease progression and prognosis in patients with prostate cancer. Prostate. May; 1999 39(2):123–9. [PubMed: 10221568]
- [121]. Shariat SF, Roehrborn CG, McConnell JD, Park S, Alam N, Wheeler TM, et al. Association of the circulating levels of the urokinase system of plasminogen activation with the presence of prostate cancer and invasion, progression, and metastasis. J Clin Oncol. Feb 1; 2007 25(4):349– 55. [PubMed: 17264329]
- [122]. Miyake H, Hara I, Yamanaka K, Gohji K, Arakawa S, Kamidono S. Elevation of serum levels of urokinase-type plasminogen activator and its receptor is associated with disease progression and prognosis in patients with prostate cancer. Prostate. 1999; 39(2):123–9. [PubMed: 10221568]
- [123]. Shariat SF, Shalev M, Menesses-Diaz A, Kim IY, Kattan MW, Wheeler TM, et al. Preoperative plasma levels of transforming growth factor beta(1) (TGF-beta(1)) strongly predict progression

- in patients undergoing radical prostatectomy. J Clin Oncol. Jun 1; 2001 19(11):2856–64. [PubMed: 11387358]
- [124]. Truong LD, Kadmon D, McCune BK, Flanders KC, Scardino PT, Thompson TC. Association of transforming growth factor-beta 1 with prostate cancer: an immunohistochemical study. Hum Pathol. Jan; 1993 24(1):4–9. [PubMed: 7678092]
- [125]. Shariat SF, Kattan MW, Traxel E, Andrews B, Zhu K, Wheeler TM, et al. Association of preand postoperative plasma levels of transforming growth factor beta(1) and interleukin 6 and its soluble receptor with prostate cancer progression. Clin Cancer Res. Mar 15; 2004 10(6):1992–9. [PubMed: 15041717]
- [126]. Shariat SF, Kim JH, Andrews B, Kattan MW, Wheeler TM, Kim IY, et al. Preoperative plasma levels of transforming growth factor beta(1) strongly predict clinical outcome in patients with bladder carcinoma. Cancer. Dec 15; 2001 92(12):2985–92. [PubMed: 11753975]
- [127]. Ivanovic V, Melman A, Davis-Joseph B, Valcic M, Geliebter J. Elevated plasma levels of TGF-beta 1 in patients with invasive prostate cancer. Nat Med. Apr; 1995 1(4):282–4. [PubMed: 7585049]
- [128]. Hobisch A, Eder IE, Putz T, Horninger W, Bartsch G, Klocker H, et al. Interleukin-6 regulates prostate-specific protein expression in prostate carcinoma cells by activation of the androgen receptor. Cancer Res. Oct 15; 1998 58(20):4640–5. [PubMed: 9788616]
- [129]. Giri D, Ozen M, Ittmann M. Interleukin-6 is an autocrine growth factor in human prostate cancer. Am J Pathol. Dec; 2001 159(6):2159–65. [PubMed: 11733366]
- [130]. Michalaki V, Syrigos K, Charles P, Waxman J. Serum levels of IL-6 and TNF-alpha correlate with clinicopathological features and patient survival in patients with prostate cancer. Br J Cancer. Jun 14; 2004 90(12):2312–6. [PubMed: 15150588]
- [131]. Nakashima J, Tachibana M, Horiguchi Y, Oya M, Ohigashi T, Asakura H, et al. Serum interleukin 6 as a prognostic factor in patients with prostate cancer. Clin Cancer Res. Jul; 2000 6(7):2702–6. [PubMed: 10914713]
- [132]. Cheifetz S, Bellon T, Cales C, Vera S, Bernabeu C, Massague J, et al. Endoglin is a component of the transforming growth factor-beta receptor system in human endothelial cells. J Biol Chem. Sep 25; 1992 267(27):19027–30. [PubMed: 1326540]
- [133]. Wikstrom P, Lissbrant IF, Stattin P, Egevad L, Bergh A. Endoglin (CD105) is expressed on immature blood vessels and is a marker for survival in prostate cancer. Prostate. Jun 1; 2002 51(4):268–75. [PubMed: 11987155]
- [134]. Fujita K, Ewing CM, Chan DY, Mangold LA, Partin AW, Isaacs WB, et al. Endoglin (CD105) as a urinary and serum marker of prostate cancer. Int J Cancer. Feb 1; 2009 124(3):664–9.
 [PubMed: 19004009]
- [135]. Karam JA, Svatek RS, Karakiewicz PI, Gallina A, Roehrborn CG, Slawin KM, et al. Use of preoperative plasma endoglin for prediction of lymph node metastasis in patients with clinically localized prostate cancer. Clin Cancer Res. Mar 1; 2008 14(5):1418–22. [PubMed: 18316564]
- [136]. Svatek RS, Karam JA, Roehrborn CG, Karakiewicz PI, Slawin KM, Shariat SF. Preoperative plasma endoglin levels predict biochemical progression after radical prostatectomy. Clin Cancer Res. Jun 1; 2008 14(11):3362–6. [PubMed: 18519764]
- [137]. Shariat SF, Andrews B, Kattan MW, Kim J, Wheeler TM, Slawin KM. Plasma levels of interleukin-6 and its soluble receptor are associated with prostate cancer progression and metastasis. Urology. Dec; 2001 58(6):1008–15. [PubMed: 11744478]
- [138]. Shariat SF, Anwuri VA, Lamb DJ, Shah NV, Wheeler TM, Slawin KM. Association of preoperative plasma levels of vascular endothelial growth factor and soluble vascular cell adhesion molecule-1 with lymph node status and biochemical progression after radical prostatectomy. J Clin Oncol. May 1; 2004 22(9):1655–63. [PubMed: 15117988]
- [139]. Shariat SF, Shalev M, Menesses-Diaz A, Kim IY, Kattan MW, Wheeler TM, et al. Preoperative plasma levels of transforming growth factor beta(1) (tgf-beta(1)) strongly predict progression in patients undergoing radical prostatectomy. J Clin Oncol. 2001; 19(11):2856–64. [PubMed: 11387358]

[140]. Shariat SF, Walz J, Roehrborn CG, Zlotta AR, Perrotte P, Suardi N, et al. External validation of a biomarker-based preoperative nomogram predicts biochemical recurrence after radical prostatectomy. J Clin Oncol. Mar 20; 2008 26(9):1526–31. [PubMed: 18349404]

- [141]. Shariat SF, Kattan MW, Erdamar S, Nguyen C, Scardino PT, Spencer DM, et al. Detection of clinically significant, occult prostate cancer metastases in lymph nodes using a splice variant-specific rt-PCR assay for human glandular kallikrein. J Clin Oncol. Apr 1; 2003 21(7):1223–31. [PubMed: 12663708]
- [142]. Gignac GA, Morris MJ, Heller G, Schwartz LH, Scher HI. Assessing outcomes in prostate cancer clinical trials: a twenty-first century tower of Babel. Cancer. Sep 1; 2008 113(5):966–74.
 [PubMed: 18661513]
- [143]. Fleming MT, Morris MJ, Heller G, Scher HI. Post-therapy changes in PSA as an outcome measure in prostate cancer clinical trials. Nat Clin Pract Oncol. Dec; 2006 3(12):658–67. [PubMed: 17139317]
- [144]. Schroder FH, Hugosson J, Roobol MJ, Tammela TL, Ciatto S, Nelen V, et al. Screening and prostate-cancer mortality in a randomized European study. N Engl J Med. Mar 26; 2009 360(13): 1320–8. [PubMed: 19297566]

	Phase	Goals/aims	Experimentation	Sample details
Ц	Preclinical	Exploratory; nominate	Preclinical study	Possible bias: small size
	Testing	and rank candidate	for hypothesis generation	and
Ī		biomarker profiles		convenience sampling
	0	Develop an assay with	Reproducibility and robustness	
		clinically reproducible results	of assay; No assessment of benefit	
	1	Test on small sample to	Marker optimization, establish	Sample population assay
		determine benefit	prediction rules, determine cut-	developed from candidate
			offs	biomarker profile
	II	Determine operating	Retrospective design be the	Sample population should
		characteristics & internal validation	target population	be the target population
	III	External validation	Retrospective or prospective,	Multi-institutional, large
			Generaliziability, Impact on	study
			clinical decision-making	
	IV	Assess whether biomarker	Post-approval reporting and	
		reduces the burden of disease	testing for other disease	
Į			processes or disease stages	

Figure 1. Modification of the structured phase-approach to the systematic discovery, evaluation, and validation of biomarkers[2, 3]