



AMITY UNIVERSITY PUNJAB, MOHALI

B.SC. MATHEMATICS (5TH SEMESTER)
SET THEORY AND METRIC SPACES (MAT304)

Zorn's Lemma, the Axiom of Choice, and the Foundations of Set Theory

Dr. Lokesh Singh

`lsingh@pb.amity.edu`

August 11, 2026

Contents

1	Introduction: What is an Axiom?	3
1.1	The Foundation of Mathematics	3
1.2	The Game of Chess Analogy	3
1.3	Historical Origins: Euclid's Geometry	3
1.3.1	Euclid's Axioms (Postulates)	3
2	Why Do We Need Axioms in Set Theory?	4
2.1	Russell's Paradox (1901)	4
2.2	The Barber Paradox (The Russell Paradox in Disguise)	4
2.3	The Connection to Russell's Paradox	5
3	The ZFC Axioms: The Constitution of Mathematics	5
3.1	The 9 Axioms of ZF (Zermelo-Fraenkel)	5
3.2	The 10th "Optional" Member: The Axiom of Choice (AC)	7
3.3	Summary Table of the ZFC Axioms	8
4	The Axiom of Choice (AC): The Most Controversial Axiom	8
4.1	Formal Statement	8
4.2	Equivalent Formulations	8
4.3	Why Do We Need the Axiom of Choice?	8
4.4	The Famous "Socks vs. Shoes" Analogy (Bertrand Russell)	9
4.5	The "Independence" of the Axiom of Choice	9
5	Zorn's Lemma (ZL)	9
5.1	Formal Statement	10
5.2	Terminology Breakdown	10
5.3	The Mountain Climbing Analogy	10
5.4	Maximal vs. Maximum: A Critical Distinction	10
6	The Equivalence: AC, ZL, and the Well-Ordering Theorem	11
6.1	Proof: $AC \Rightarrow ZL$ (Detailed)	11
6.2	Proof: $ZL \Rightarrow WO$ (Detailed)	12
6.3	Proof: $WO \Rightarrow AC$ (Trivial)	12
7	Applications of Zorn's Lemma	12
7.1	Application 1: Every Vector Space has a Basis	12
7.2	Application 2: Existence of Maximal Ideals in Rings	13
7.3	Application 3: Comparability of Cardinals (Total Order of Cardinals)	13
8	Consequences of Accepting the Axiom of Choice	13
8.1	The "Good" Consequences (Why Mathematicians Love AC)	13
8.2	The "Weird" Consequences (Why Some Mathematicians Reject AC)	14
8.3	The Price of Rejecting AC	14
8.4	Countable Choice vs. Full Choice	14
9	Axioms Are Choices, Not Absolute Truths	14
9.1	The Role of Intuition	15

10 Summary: The Big Picture	15
Exercises	16
References	16

1 Introduction: What is an Axiom?

1.1 The Foundation of Mathematics

Imagine you are building a house. You cannot build walls without a foundation. You cannot construct the second floor without the first floor. Mathematics is exactly like a giant building—every theorem rests on previous theorems, which rest on definitions, which ultimately must rest on something we just *accept* as true without proof.

Those foundational "truths" are called **axioms**.

Definition 1.1 (Axiom). An **axiom** (or **postulate**) is a statement that is taken to be true without proof. It serves as a starting point for logical reasoning and deduction.

1.2 The Game of Chess Analogy

Think of mathematics as a game of chess.

- The **axioms** are the rules of the game (e.g., "A bishop moves diagonally," "A pawn moves forward one square").
- The **theorems** are the strategies and winning moves you can derive from those rules.

You cannot "prove" that a bishop moves diagonally—it is simply a rule you agree to follow. Similarly, in mathematics, we agree to follow certain fundamental rules, and from them we derive everything else. If you change the rules (axioms), you are playing a different game (a different mathematical system).

1.3 Historical Origins: Euclid's Geometry

The concept of axioms is ancient. The Greek mathematician **Euclid** (c. 300 BCE) wrote his famous work *The Elements*, which organized all known geometry into a logical system based on just a handful of self-evident truths.

1.3.1 Euclid's Axioms (Postulates)

Euclid began with five postulates (axioms). Here are the first four:

1. A straight line can be drawn from any point to any point.
2. A finite straight line can be extended continuously in a straight line.
3. A circle can be drawn with any center and any radius.
4. All right angles are equal to each other.

The **Fifth Postulate** (the "Parallel Postulate") was more complicated and controversial:

"If a straight line falling on two straight lines makes the interior angles on the same side less than two right angles, the two straight lines, if produced indefinitely, meet on that side on which the angles are less than two right angles."

For centuries, mathematicians tried to prove this fifth postulate from the other four. They failed. Eventually, in the 19th century, mathematicians discovered that by **replacing** this axiom, they could create entirely new, consistent geometries—now called **non-Euclidean geometries**.

Remark 1.1 (The Big Lesson). This historical episode teaches us a vital lesson: **Axioms are not "absolute truths" about the universe. They are assumptions we choose to make.** By choosing different axioms, we create different mathematical worlds. All are equally valid internally; they just describe different structures.

2 Why Do We Need Axioms in Set Theory?

In the late 19th century, Georg Cantor developed set theory in a *naive* way. He assumed that given any property $P(x)$, we could form the set of all objects satisfying $P(x)$. This is called the **Comprehension Principle**:

$$\{x : P(x) \text{ is true}\}.$$

This seems utterly intuitive. But it leads to disaster.

2.1 Russell's Paradox (1901)

Define the set of all sets that do not contain themselves:

$$R = \{X : X \notin X\}.$$

Now ask: Does $R \in R$?

- If $R \in R$, then by definition $R \notin R$, contradiction.
- If $R \notin R$, then by definition $R \in R$, contradiction.

This was a **devastating blow** to naive set theory. It showed that you cannot simply assume that every collection of objects forms a set. You need **rules**—**axioms**—to restrict what sets can be formed.

2.2 The Barber Paradox (The Russell Paradox in Disguise)

The Barber Paradox is the perfect "layman's version" of Russell's paradox. It goes like this:

In a certain village, there is a barber. This barber shaves **all and only** those men who do **not** shave themselves.

Now ask: **Who shaves the barber?**

Let us analyze both possibilities:

1. **If the barber shaves himself:** Since the barber shaves only those men who do not shave themselves, and he has shaved himself, this means he is among those who *do* shave themselves. But his rule says he shaves only those who *do not* shave themselves. Contradiction.

2. **If the barber does not shave himself:** Since the barber shaves all men who do not shave themselves, and he has not shaved himself, this means he falls into the category of men who do not shave themselves. Therefore, he *must* shave himself. Contradiction.

Either way, we get a logical contradiction. The only resolution is that **no such barber can exist** in that village. The "set" of all men who do not shave themselves cannot be formed in a way that includes the barber.

2.3 The Connection to Russell's Paradox

The Barber Paradox is structurally identical to Russell's paradox:

- **Barber** = The set R .
- **Men who shave themselves** = Sets that contain themselves ($X \in X$).
- **Men who do not shave themselves** = Sets that do not contain themselves ($X \notin X$).

Just as no such barber can exist, no such set R can exist in a consistent set theory. The barber paradox shows that the problem is not just a mathematical technicality—it is a fundamental logical issue about self-reference.

Example 2.1 (Other Verbal Paradoxes). • **The Liar Paradox:** "This statement is false." If it is true, then it is false; if it is false, then it is true.

- **Grelling's Paradox:** An adjective is *heterological* if it does not describe itself (e.g., "long" is not long). Is "heterological" heterological?

3 The ZFC Axioms: The Constitution of Mathematics

To avoid paradoxes, mathematicians developed the **Zermelo-Fraenkel axioms** (with the Axiom of Choice), collectively called **ZFC**.

These axioms act like a **strict constitution** for mathematics. They tell us what we can build and strictly forbid us from building "too big" or "self-referential" monstrosities. Here is the complete list of all 10 axioms of ZFC:

3.1 The 9 Axioms of ZF (Zermelo-Fraenkel)

1. Axiom of Extensionality (The "Equality" Rule)

Two sets are equal if and only if they have exactly the same elements.

Why we need it: This defines what a "set" fundamentally is. It tells us that the order of elements doesn't matter, and duplicates don't matter. A set is entirely determined by its members.

Example: $\{1, 2, 3\} = \{3, 2, 1\}$ and $\{1, 1, 2\} = \{1, 2\}$.

$$\forall x \forall y (\forall z (z \in x \leftrightarrow z \in y) \rightarrow x = y)$$

2. Axiom of Empty Set (Existence of "Nothing")

There exists a set that contains no elements.

Why we need it: In mathematics, we need a starting point—a "zero" for sets. Without it, we couldn't prove that *any* set exists at all.

Notation: This unique set is denoted by $\emptyset$ or $\{\}$.

3. Axiom of Pairing (Making Two into One)

For any two sets a and b , there exists a set whose only members are exactly a and b .

Why we need it: This allows us to build ordered pairs and, consequently, relations and functions.

Example: Given $a = \{1\}$ and $b = \{2\}$, this axiom guarantees the existence of the set $\{\{1\}, \{2\}\}$. From pairing, we can define the ordered pair (a, b) using Kuratowski's definition: $(a, b) = \{\{a\}, \{a, b\}\}$.

4. Axiom of Union (Gathering Elements Together)

For any set A whose elements are themselves sets, there exists a set U that contains all the elements of all the members of A .

Why we need it: This allows us to "flatten" a collection of sets. If you have a box full of boxes, this axiom lets you dump all the contents of the inner boxes into one big box.

Example: If $A = \{\{1, 2\}, \{3, 4\}, \{5\}\}$, then $\bigcup A = \{1, 2, 3, 4, 5\}$. This is essential for defining the union of two sets: $X \cup Y = \bigcup \{X, Y\}$.

5. Axiom of Power Set (Creating All Subsets)

For any set A , there exists a set $P(A)$ whose members are exactly all the subsets of A .

Why we need it: This is the axiom that generates larger and larger infinities. It allows us to form the set of all possible combinations of elements of A .

Example: If $A = \{1, 2\}$, then $P(A) = \{\emptyset, \{1\}, \{2\}, \{1, 2\}\}$.

Significance: This is what powers Cantor's Theorem, proving that $|A| < |P(A)|$ and giving us the infinite hierarchy of cardinalities.

6. Axiom of Infinity (The "First Infinity")

There exists a set that contains $\emptyset$ and is "closed" under the successor operation $S(x) = x \cup \{x\}$.

Why we need it: Without this axiom, we could only build finite sets. This axiom explicitly declares the existence of an infinite set—specifically, the set of natural numbers $\mathbb{N}$.

How it builds $\mathbb{N}$:

- $0 = \emptyset$
- $1 = \{\emptyset\}$
- $2 = \{\emptyset, \{\emptyset\}\}$

This axiom gives us a place to start doing calculus, analysis, and counting infinite sets.

7. Axiom of Separation (Subset / Specification) – [The Paradox-Solver]

If we already have a set A and a property $P(x)$, we can form a new set consisting of exactly those elements of A that satisfy $P(x)$.

Why we need it: This is the critical axiom that destroys Russell's Paradox. In naïve set theory, you could ask for $\{x : x \notin x\}$ directly, which led to disaster. This axiom forces you to write $\{x \in A : x \notin x\}$. Since you cannot have a universal set A containing everything, Russell's set cannot be formed.

Example: From the set of integers $\mathbb{Z}$, we can separate out the even numbers using the property $P(x) = "x \text{ is divisible by } 2"$.

8. Axiom of Replacement (The Image Axiom)

If we have a set A and a function (or rule) F that assigns a unique set to each element of A , then the collection of all those images $\{F(x) : x \in A\}$ is also a set.

Why we need it: This is a subtle but powerful axiom. It allows us to construct sets that are "large" relative to other sets. For example, if you have a set of indices and each index corresponds to a set, Replacement lets you collect all those sets together into a new set. It is essential for defining ordinals and cardinals and for proving that the hierarchy of infinities continues past ω .

9. Axiom of Foundation (Regularity) – [The "Anti-Circularity" Rule]

Every non-empty set A contains an element x such that $x \cap A = \emptyset$ (i.e., x is disjoint from A).

Why we need it: This axiom bans infinite descending chains of sets like $A_1 \ni A_2 \ni A_3 \ni \dots$ and also bans a set from containing itself ($A \in A$). It ensures that sets are built up in a well-founded, step-by-step hierarchy starting from the empty set. This guarantees that set theory is free from weird, circular self-reference.

3.2 The 10th "Optional" Member: The Axiom of Choice (AC)

For any collection of non-empty sets, there exists a function (called a **choice function**) that picks exactly one element from each set.

Why we need it: As we will discuss extensively in the next sections, AC is incredibly useful. It is equivalent to Zorn's Lemma and the Well-Ordering Theorem. It allows us to prove that every vector space has a basis, every ring has a maximal ideal, and that cardinalities are comparable.

Why it is separate: AC is **independent** of the other 8 axioms. You cannot prove or disprove it using the others (thanks to Gödel and Cohen). Most mathematicians accept it (forming ZFC) because it makes math much more powerful, despite leading to strange consequences like the Banach-Tarski paradox.

3.3 Summary Table of the ZFC Axioms

#	Axiom Name	Core Purpose
1	Extensionality	Defines set equality by members.
2	Empty Set	Declares $\emptyset$ exists.
3	Pairing	Creates $\{a, b\}$.
4	Union	Flattens a set of sets into one set.
5	Power Set	Creates all subsets (higher infinities).
6	Infinity	Declares an infinite set exists (creates $\mathbb{N}$).
7	Separation	Filters existing sets by properties. (Defeats Russell's Paradox).
8	Replacement	Substitutes images for elements (defines ordinals/cardinals).
9	Foundation	Bans self-membership and infinite loops.
10	Choice (AC)	Picks one element from each set.

4 The Axiom of Choice (AC): The Most Controversial Axiom

4.1 Formal Statement

Definition 4.1 (Axiom of Choice). For any collection $\{A_i\}_{i \in I}$ of non-empty sets, there exists a **choice function**

$$f : I \rightarrow \bigcup_{i \in I} A_i$$

such that $f(i) \in A_i$ for every $i \in I$.

4.2 Equivalent Formulations

1. **Product Formulation:** The Cartesian product of any non-empty family of non-empty sets is non-empty:

$$\prod_{i \in I} A_i \neq \emptyset.$$

2. **Partition Formulation (AC for disjoint sets):** For any partition of a set X into non-empty subsets, there exists a set Y that contains exactly one element from each member of the partition.

4.3 Why Do We Need the Axiom of Choice?

Consider the following situations:

- **Finite Collection:** If you have 100 boxes, each containing apples, you can pick one apple from each. This is provable by induction. No axiom needed.
- **Countably Infinite Collection with a Rule:** If you have infinitely many pairs of shoes, you can define the choice function: "Pick the left shoe." This is a rule. No AC needed.
- **Arbitrary Infinite Collection without a Rule:** If you have infinitely many pairs of identical socks, there is no natural way to distinguish a "left" sock from a "right" sock. You cannot write down a rule for picking one from each pair. Yet, AC says you can *assume* such a choice exists.

4.4 The Famous "Socks vs. Shoes" Analogy (Bertrand Russell)

This analogy beautifully illustrates the core issue:

Example 4.1 (Shoes vs. Socks). Suppose you have infinitely many pairs of shoes. You can define a choice function without any controversy: "From each pair, choose the left shoe." There is a definite rule.

Now suppose you have infinitely many pairs of **socks**. Socks are identical! There is no "left sock" or "right sock." For finitely many pairs, you can just reach into each and pull one out. But for **infinitely many** pairs, you cannot physically do this one by one. You need a mathematical guarantee that such a simultaneous selection exists, even though no explicit rule can be written down.

The **Axiom of Choice** is exactly that guarantee.

Key Insight: AC is not about finite choices; it is about **making infinitely many arbitrary choices simultaneously without a definable rule**.

4.5 The "Independence" of the Axiom of Choice

In 1963, the American mathematician **Paul Cohen** proved a landmark result:

Theorem 4.1 (Cohen, 1963). The Axiom of Choice is **independent** of the other ZF axioms. This means:

- You cannot prove AC from the other axioms.
- You cannot disprove AC from the other axioms.

So you are free to *choose* whether to accept it or reject it.

This is exactly analogous to Euclid's parallel postulate. You can accept it (Euclidean geometry) or reject it (non-Euclidean geometry). Both are valid systems.

5 Zorn's Lemma (ZL)

Zorn's Lemma, formulated by Max Zorn in 1935, is **equivalent** to the Axiom of Choice. It provides a different—and often more convenient—way to apply the same underlying idea.

5.1 Formal Statement

Definition 5.1 (Zorn's Lemma). Let P be a non-empty partially ordered set (poset). Suppose that **every chain** (i.e., every totally ordered subset) in P has an **upper bound** in P . Then P contains at least one **maximal element**.

5.2 Terminology Breakdown

Before we proceed, let us clearly define the key terms:

- **Chain:** A subset $C \subset P$ such that for any $x, y \in C$, either $x \leq y$ or $y \leq x$. Think of it as a "path" where every element is comparable.
- **Upper Bound:** An element $u \in P$ such that $c \leq u$ for all $c \in C$. Note: u does **not** have to be in the chain itself.
- **Maximal Element:** An element $m \in P$ such that there is **no** $x \in P$ with $m < x$. A maximal element is not necessarily a "largest" element (maximum); it is simply an element that cannot be exceeded within the poset.

5.3 The Mountain Climbing Analogy

Imagine you are exploring an infinite mountain range.

- **Poset (P):** The set of all possible points on the mountains.
- **Chain:** A climbing route where you only go upwards. Every point on the route is higher than the previous.
- **Upper Bound:** If your route goes on forever, Zorn's Lemma's hypothesis says that there is **some point** (maybe a plateau or a nearby peak) that is at least as high as every point on your route.
- **Maximal Element:** A peak. It might not be the highest mountain in the entire world, but it is a point from which you cannot climb *strictly* higher.

Zorn's Lemma essentially says: "If every infinite upward path has a ceiling above it, then there must be at least one peak that you cannot climb above."

5.4 Maximal vs. Maximum: A Critical Distinction

- **Maximum (Top element):** An element t such that $x \leq t$ for *all* $x \in P$. There can be at most one maximum.
- **Maximal:** An element m such that there is no x with $m < x$. There can be many maximal elements.

Example 5.1. Consider the poset of proper ideals of a ring (ordered by inclusion).

- A **maximal ideal** is one that cannot be enlarged to a strictly larger proper ideal.
- There may be many maximal ideals, and none of them is a **maximum** (since no single proper ideal contains all others).

Zorn's Lemma gives maximal elements, not maxima.

6 The Equivalence: AC, ZL, and the Well-Ordering Theorem

One of the most beautiful results in set theory is that these three seemingly different statements are **logically equivalent** (assuming the ZF axioms).

Principle	Intuition	Equivalent?
Axiom of Choice	Pick one element from each set in a family.	Yes
Zorn's Lemma	If every chain has an upper bound, there is a maximal element.	Yes
Well-Ordering Theorem	Every set can be arranged so every subset has a least element.	Yes

Theorem 6.1 (Zermelo, 1904; Zorn, 1935). $AC \iff ZL \iff WO$ over ZF.

6.1 Proof: $AC \Rightarrow ZL$ (Detailed)

This is the most technically involved direction. We will present the proof using **transfinite induction with ordinals**, which is conceptually clear.

Proof. Let P be a non-empty poset where every chain has an upper bound. Suppose, for contradiction, that P has no maximal element.

Define a **choice function** $c : P \rightarrow P$ using AC as follows: For each $x \in P$, the set

$$S_x = \{y \in P : y > x\}$$

is non-empty (since x is not maximal). Let $c(x) \in S_x$. Thus $c(x) > x$ for all $x \in P$.

We now build a strictly increasing sequence indexed by all ordinals using transfinite recursion:

- **Step 0:** Pick any $x_0 \in P$.
- **Successor Step:** For any ordinal α , define $x_{\alpha+1} = c(x_\alpha)$. Then $x_\alpha < x_{\alpha+1}$.
- **Limit Step:** For a limit ordinal λ (e.g., $\omega, \omega + \omega, \dots$), the set $\{x_\alpha : \alpha < \lambda\}$ is a chain in P (since the sequence is strictly increasing). By the hypothesis of Zorn's Lemma, this chain has an upper bound. Let x_λ be any such upper bound.

This recursive definition produces a strictly increasing sequence

$$x_0 < x_1 < x_2 < \dots < x_\alpha < \dots$$

for *all* ordinals α .

But this is impossible! The ordinals form a proper class (they are "too numerous" to be a set), while P is a set. The map $\alpha \mapsto x_\alpha$ would inject the class of all ordinals into the set P , which contradicts the axiom of replacement (or simply the fact that a set cannot have a proper class of elements).

Therefore, our assumption was false, and P must have a maximal element. □

6.2 Proof: ZL $\Rightarrow$ WO (Detailed)

Proof. Let A be any set. We need to prove that A can be well-ordered.

Consider the poset W whose elements are all **well-ordered subsets** of A . That is, an element of W is a pair $(S, \leq_S)$ where $S \subset A$ and $\leq_S$ is a well-order on S .

Order W by: $(S, \leq_S) \leq (T, \leq_T)$ if $S \subset T$, the order on S agrees with the order on T restricted to S , and S is an **initial segment** of T (meaning if $x \in S$ and $y \in T$ with $y \leq_T x$, then $y \in S$).

Chain Check: Let $\{(S_i, \leq_i)\}$ be a chain in W . Let $S = \bigcup S_i$. We can define a well-order on S : for $x, y \in S$, choose i such that both $x, y \in S_i$ (since they form a chain, one contains the other), and compare them using $\leq_i$. This is well-defined. The union S is a well-ordered subset of A . Thus $(S, \leq)$ is an upper bound for the chain.

By Zorn's Lemma, W has a maximal element $(M, \leq_M)$. We claim $M = A$. If not, pick $a \in A \setminus M$. We can extend the well-order on M by declaring a to be greater than every element of M . This gives a well-ordered set $M \cup \{a\}$ properly extending M , contradicting maximality. Hence $M = A$. So A is well-ordered. $\square$

6.3 Proof: WO $\Rightarrow$ AC (Trivial)

Proof. Given a family of non-empty sets $\{A_i\}_{i \in I}$, well-order the union $U = \bigcup_{i \in I} A_i$. Define the choice function $f(i)$ to be the **least element** of A_i with respect to this well-order. This gives a choice function without invoking AC separately. $\square$

7 Applications of Zorn's Lemma

Zorn's Lemma is the workhorse of modern algebra and analysis. Here are three of its most important applications.

7.1 Application 1: Every Vector Space has a Basis

This is perhaps the most famous application.

Theorem 7.1. Every vector space V over a field K has a basis (a maximal linearly independent set that spans V).

Proof. Let P be the set of all linearly independent subsets of V , ordered by inclusion.

Chain Check: Let $\{B_i\}_{i \in I}$ be a chain in P . Let $B = \bigcup_{i \in I} B_i$. We must prove B is linearly independent.

Take any finite subset $\{v_1, \dots, v_n\} \subset B$. Each v_j belongs to some B_{i_j} . Since the B_i 's form a chain, one of them (say B_{i_k}) contains all the v_j 's. Because B_{i_k} is linearly independent, any linear combination $\sum c_j v_j = 0$ forces all $c_j = 0$. Hence B is linearly independent. So $B \in P$ and is an upper bound for the chain.

By Zorn's Lemma, P has a maximal element B_0 . We claim B_0 spans V . If not, there exists $v \in V$ not in the span of B_0 . Then $B_0 \cup \{v\}$ is linearly independent (otherwise v would be in the span of B_0). This contradicts the maximality of B_0 . Therefore B_0 spans V , so it is a basis. $\square$

7.2 Application 2: Existence of Maximal Ideals in Rings

Theorem 7.2. Every commutative ring with unity R has a maximal ideal (assuming $R \neq \{0\}$).

Proof. Let P be the set of all **proper ideals** of R (ideals not containing the identity 1), ordered by inclusion.

Chain Check: Let $\{I_i\}_{i \in I}$ be a chain of proper ideals. Let $I = \bigcup I_i$. We must show I is a proper ideal.

- I is an ideal: The union of a chain of ideals is an ideal (check closure under addition and multiplication by ring elements).
- I is proper: If $1 \in I$, then $1 \in I_i$ for some i , contradicting that I_i is proper.

Thus $I \in P$ and is an upper bound.

By Zorn's Lemma, P has a maximal element M . This M is a maximal ideal. $\square$

7.3 Application 3: Comparability of Cardinals (Total Order of Cardinals)

Theorem 7.3. For any two sets A and B , either $|A| \leq |B|$ or $|B| \leq |A|$. In other words, cardinal numbers are totally ordered.

Proof. Consider the poset P of all triples (A', B', f) where $A' \subset A$, $B' \subset B$, and $f : A' \rightarrow B'$ is a bijection. Order by extension: $(A_1, B_1, f_1) \leq (A_2, B_2, f_2)$ if $A_1 \subset A_2$, $B_1 \subset B_2$, and $f_2|_{A_1} = f_1$.

Chain Check: The union of a chain of such bijections gives a bijection from the union of their domains to the union of their codomains. This is an upper bound.

By Zorn's Lemma, we get a maximal bijection (A_0, B_0, f_0) . If $A_0 \neq A$ and $B_0 \neq B$, we can pick $a \in A \setminus A_0$ and $b \in B \setminus B_0$, and extend f_0 by setting $f(a) = b$, contradicting maximality. Hence either $A_0 = A$ (so $|A| \leq |B|$) or $B_0 = B$ (so $|B| \leq |A|$). $\square$

8 Consequences of Accepting the Axiom of Choice

8.1 The "Good" Consequences (Why Mathematicians Love AC)

If you accept AC, you get:

1. **Zorn's Lemma:** Every vector space has a basis.
2. **Well-Ordering Theorem:** Every set can be well-ordered.
3. **Comparability of Cardinals:** Any two sets have comparable cardinalities.
4. **Existence of Maximal Ideals:** In every commutative ring with unity.
5. **Tychonoff's Theorem:** The product of compact topological spaces is compact.

8.2 The "Weird" Consequences (Why Some Mathematicians Reject AC)

If you accept AC, you also get:

1. **Banach-Tarski Paradox:** A solid ball can be cut into finitely many pieces and reassembled into two identical balls. This defies physical intuition (but the pieces are non-measurable).
2. **Existence of Non-Measurable Sets:** There are subsets of $\mathbb{R}$ that cannot be assigned a length (Vitali sets).

Remark 8.1. The Banach-Tarski paradox is not a logical contradiction; it is just highly counterintuitive. It relies on the fact that AC allows the existence of "wild" sets that cannot be physically constructed.

8.3 The Price of Rejecting AC

If we reject AC, many standard theorems fail:

- There may exist vector spaces without a basis.
- The Cartesian product of non-empty sets may be empty.
- There may be infinite sets that are not Dedekind-infinite (i.e., they cannot be put into bijection with a proper subset of themselves).
- The comparability of cardinals fails: there might be two sets whose cardinalities are incomparable.

8.4 Countable Choice vs. Full Choice

Most mathematicians are comfortable with the **Axiom of Countable Choice** (AC for countable families), which is needed for standard results in analysis (e.g., the equivalence of sequential continuity and continuity for metric spaces). The controversy mainly surrounds the **Full Axiom of Choice**, which applies to uncountable families.

Note 8.1. In this course, we will use Zorn's Lemma freely. It is standard practice in modern mathematics. However, we will occasionally note when a theorem explicitly relies on AC.

9 Axioms Are Choices, Not Absolute Truths

This is perhaps the most important takeaway for students:

Mathematics is not about discovering absolute truths about the universe.

Mathematics is about exploring the logical consequences of carefully chosen assumptions (axioms).

- Euclid chose his axioms $\rightarrow$ Euclidean geometry.

- Lobachevsky and Bolyai chose a different parallel postulate → Hyperbolic geometry.
- Zermelo and Fraenkel chose their axioms → Standard set theory (ZFC).
- If you reject AC, you get a different set theory (ZF without Choice).

All of these are valid mathematical systems. They are internally consistent. They just describe different "mathematical universes."

9.1 The Role of Intuition

Why do we choose one set of axioms over another?

- **Intuition:** They seem "self-evidently true" (e.g., Euclid's first four postulates).
- **Utility:** They allow us to prove useful theorems (e.g., AC).
- **Consistency:** They do not lead to contradictions (as far as we know).

The Axiom of Choice is accepted by the majority of mathematicians because it is *intuitive* (it seems obvious that you can pick an element from each set) and *useful* (it underpins vast swathes of modern mathematics). The weird consequences are tolerated as a price for this utility.

10 Summary: The Big Picture

- **Axiom** = A statement accepted without proof. It is the foundation of a mathematical system.
- **Why we need them:** To avoid contradictions (like Russell's paradox) and to provide a solid starting point for deduction.
- **The ZFC axioms** are the standard foundation. They consist of 9 ZF axioms (Extensionality, Empty Set, Pairing, Union, Power Set, Infinity, Separation, Replacement, Foundation) plus the Axiom of Choice (AC).
- **The Axiom of Choice** is a specific axiom that guarantees the existence of choice functions for arbitrary families of sets.
- AC is **independent** of the other axioms—you can take it or leave it.
- Accepting AC gives us powerful tools (Zorn's Lemma, bases for vector spaces) but also leads to bizarre results (Banach-Tarski).
- Ultimately, axioms are **choices**. We choose the mathematical system that best suits the problems we want to solve.

In this course, we will adopt ZFC as our foundation.
 We will use Zorn's Lemma freely because it is the standard tool in modern algebra and analysis.
 However, we will remain aware that we are *choosing* to use it.

Exercises

1. Explain in your own words why an axiom is not a theorem.
2. What is the difference between Euclid's parallel postulate and the Axiom of Choice? How are they historically similar?
3. Explain Russell's paradox in a way that a non-mathematician could understand. (Use the barber example.)
4. Why does the Axiom of Separation prevent Russell's paradox?
5. List three theorems that would fail if we rejected the Axiom of Choice.
6. What is the Banach-Tarski paradox? Does it disprove AC? Why or why not?
7. If you were a mathematician, would you accept AC? Justify your answer with reasons.
8. Use Zorn's lemma to prove that every chain in a partially ordered set is contained in a maximal chain.
9. Prove that the Axiom of Choice implies the Well-Ordering Theorem. (Hint: Use transfinite induction to build a well-ordering.)
10. Let P be a poset. Suppose every chain in P has a least upper bound. Prove that P has a maximal element. (This is a variant of Zorn's lemma.)
11. Use Zorn's lemma to prove that every ideal in a ring (not necessarily with unity) is contained in a maximal ideal, provided the ideal is proper. (Note: The union argument needs modification because 1 may not exist.)
12. Explain why the "Socks vs. Shoes" analogy illustrates the need for the Axiom of Choice. Why do shoes not require AC but socks do?
13. Prove that AC is equivalent to the statement: "For any non-empty set X , the Cartesian product of the non-empty subsets of X is non-empty."
14. Show that the Banach-Tarski paradox requires the Axiom of Choice. What would happen if we rejected AC?

References

1. Kaplansky, I. *Set Theory and Metric Spaces*. Allyn and Bacon, 1972. (Chapter 2-3)
2. Andersson, S., Björn, A., & Wiman, D. *An Introduction to Metric Spaces*. (Chapter 1)
3. Halmos, P. *Naive Set Theory*. Springer, 1960.
4. Jech, T. *Set Theory*. Springer, 2003. (For advanced equivalence proofs)
5. Euclid. *The Elements* (c. 300 BCE).
6. Russell, B. *Principles of Mathematics*. 1903. (For the paradox)