

---

## On the robustness of the relation $\beta_{tr} = 2\beta$

### Definitions:

1.

$$C_i(t) = \text{Tr} \left\{ \rho_\infty \left( \hat{n}_i(t) - \frac{1}{2} \right) \left( \hat{n}_{i_0}(t=0) - \frac{1}{2} \right) \right\} \quad R^2(L, t) = \sum_{i \in \Lambda} |i - i_0|^2 C_i(t)$$

$$\hat{N}_{\mathcal{D}} = \sum_{i \in \mathcal{D}} \hat{n}_i \quad w^2(L, t) = \left\langle \hat{N}_{\mathcal{D}}^2 \right\rangle - \left\langle \hat{N}_{\mathcal{D}} \right\rangle^2$$

2. Defining exponents  $\{\alpha_{tr}, \beta_{tr}, z_{tr} = \alpha_{tr}/\beta_{tr}\}$  for the FV scaling of MSD  $R^2(L, t)$  and exponents  $\{\alpha, \beta, z = \alpha/\beta\}$  for the FV scaling of subsystem number variance  $w^2(L, t)$ , i.e.,:

$$R^2(L, t) = \begin{cases} t^{\beta_{tr}} & \text{for } t \ll L^{z_{tr}} \\ L^{\alpha_{tr}} & \text{for } t \gg L^{z_{tr}} \end{cases}$$

$$w^2(L, t) = \begin{cases} t^{\beta} & \text{for } t \ll L^z \\ L^{\alpha} & \text{for } t \gg L^z \end{cases}$$

### Questions:

1. Do MSD  $R^2(L, t)$  and number variance  $w^2(L, t)$  saturate (to a system size dependent value) on the same time scale, i.e., is  $z = z_{tr}$ ?

We know that for a single particle subspace:

$$4C_i(t) = |\psi(i, t)|^2 = \langle \hat{n}_i \rangle_t$$

$R^2(L, t)$  saturating in the time scale  $L^{z_{tr}}$ , implies that 2-point correlators  $\langle \hat{n}_i \rangle_t$  saturate in the same time scale.

Whilst  $w^2(L, t)$  is made up of summation of 4-point and 2-point correlators-  $\langle \hat{n}_i \hat{n}_j \rangle_t$  and  $\langle \hat{n}_i \rangle_t$ . **Hence,  $z = z_{tr}$  implies that the 4-point correlators saturate in a time scale which is at the most equal to  $L^{z_{tr}}$ .**