


Cost analysis of the NIRUDAK & DHAKA clinical diagnostic model for volume deficit in patients with acute diarrhea



BROWN
Alpert Medical School

Anagha Lokhande BA¹, Monique Gainey MS MPH², Stephanie C. Garbern MD MPH¹, Sabiha Nasrin MBBS MPH³,
Nur H. Alam MD MBBS PhD³, Adam C. Levine MD MPH¹

¹ Alpert Medical School, Brown University, Providence, RI, USA

² Rhode Island Hospital, Providence, RI, USA

³ International Centre for Diarrhoeal Disease Research, Bangladesh (icddr,b), Dhaka, Bangladesh



Background

- With over 6.5 billion cases and 1.4 million deaths in 2019, diarrheal diseases are a major cause of morbidity and mortality and place a heavy burden on healthcare systems worldwide
- While the World Health Organization's (WHO) guidelines¹ on acute diarrhea management is the standard practice of care worldwide, the NIRUDAK² and DHAKA³ models, which assesses dehydration severity in patients over and under 5 years of age respectively, have been more recently developed
- Gaps remain in the literature with regards to cost of care for acute diarrhea management especially in LMICs
- This study aims to compare simulated treatment costs of acute diarrhea management of the WHO guidelines to the NIRUDAK and DHAKA models

Methods

- Cost of care for each patient presenting to the International Centre for Diarrhoeal Research, Bangladesh's (icddr,b) Dhaka Hospital between March 2019 – March 2020 represents the summed costs of:
 - fluid administered
 - hospital fees
 - equipment (e.g. IV tubing and needles)
- NIRUDAK model: total costs of resuscitation & cost of fluid required for initial resuscitation (within first 6 hours of admission) were calculated and reported as median and interquartile range (IQR) in USD
- DHAKA model: due to data limitations, total costs of fluid were calculated and reported as median and IQR in USD

Results




Figure 1. Summary of cost comparisons – NIRUDAK model




Figure 2. Total cost of fluid – DHAKA model

- NIRUDAK model, median total cost per patient: \$5.18 (IQR: 0 – 25.56)
 - WHO: \$5.23 (IQR: 5.09 – 22.17)
- NIRUDAK model, median initial fluid resuscitation cost per patient: \$3.27 (IQR: 0 – 4.27)
 - WHO: \$4.55 (IQR: 0 – 5.76)
- DHAKA model, total fluid cost: \$0.02 (IQR: 0 – 0.97)
 - WHO: \$0.04 (IQR: 0.03 – 1.24)

Conclusion

- Implementing the most cost-effective approach to diarrhea management will help optimize allocation of resources, which is especially critical in low resource settings
- Measuring the societal cost savings (e.g., of hospital beds, physician and nurse labor, transportation to healthcare facilities, etc.) is beyond the scope of this analysis
- However, more accurately diagnosing patient dehydration severity can free up hospital beds and resources for the most severely ill patients, saving moderately ill patients the costs of in-patient care
- Thus, the NIRUDAK and DHAKA models may provide positive externalities unable to be captured here

References & Disclosures

1. World Health Organization (WHO). IMAI District Clinician Manual: Hospital Care for Adolescents and Adults. World Heal Organ [Internet]. 2011; 2:780. Available from: http://apps.who.int/iris/bitstream/10665/77751/3/9789241548290_Vol2_eng.pdf?ua=1
2. Levine AC, Barry MA, Gainey M, Nasrin S, Qu K, Schmid CH, et al. (2021) Derivation of the first clinical diagnostic models for dehydration severity in patients over five years with acute diarrhea. PLoS Negl Trop Dis 15(3): e0009266. <https://doi.org/10.1371/journal.pntd.0009266>
3. Levine, A. C., Glavis-Bloom, J., Modi, P., Nasrin, S., Rege, S., Chu, C., Schmid, C. H., & Alam, N. H. (2015). Empirically Derived Dehydration Scoring and Decision Tree Models for Children With Diarrhea: Assessment and Internal Validation in a Prospective Cohort Study in Dhaka, Bangladesh. Global health, science and practice, 3(3), 405–418. <https://doi.org/10.9745/GHSP-D-15-00097>

The authors have no conflict of interests to declare.

Funding was provided by the National Institutes of Health (ID: 5R01DK116163-03).