

Solutions of HW9

Chapter 7

31. Let X_1, X_2, \dots, X_n be a random sample of size n from an exponential random variable with parameter β .

Let's find the moments based estimator:

$$E[X] = \beta$$

Replace both sides by its estimators, we have,

$$M_1 = \hat{\beta}$$

i.e., $\hat{\beta} = \bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$

Let's find the maximum likelihood estimator:

The likelihood function is:

$$L(\beta) = \prod_{i=1}^n f(X_i) = \prod_{i=1}^n \frac{1}{\beta} e^{-X_i/\beta} = \frac{1}{\beta^n} e^{-\frac{1}{\beta} \sum_{i=1}^n X_i}$$

The log likelihood function is:

$$\ln L(\beta) = -n \ln \beta - \frac{1}{\beta} \sum_{i=1}^n X_i$$

Take the derivative of the log likelihood function:

$$\frac{d}{d\beta} \ln L(\beta) = -\frac{n}{\beta} + \frac{1}{\beta^2} \sum_{i=1}^n X_i$$

Let the derivative equal to zero. We have,

$$\hat{\beta} = \frac{1}{n} \sum_{i=1}^n X_i = \bar{X}$$

The maximum likelihood estimator and the moments-based estimator has the same form.

32. (a) $\hat{\lambda} = \bar{x} = 10.3$ calls

- (b) The time of arrival of the first call of the day is exponential with parameter

$$\beta = \frac{1}{\lambda}.$$

$$\hat{\mu} = \hat{\beta} = \frac{1}{\hat{\lambda}} = \frac{1}{10.3} = .0971 \text{ hours}$$

Since .0971 hours = 5.83 minutes and the center opens at 9 a.m., the average time of arrival of the first call is estimated to be 9:058.

33. (a) Let X_1, X_2, \dots, X_n be a random sample of size n from a gamma random variable with parameter $\alpha = 2$ and β unknown.

We have, $f(x) = \frac{1}{\Gamma(2)\beta^2} x e^{-x/\beta}$

The likelihood function is:

$$L(\beta) = \prod_{i=1}^n f(X_i) = \prod_{i=1}^n \frac{1}{\Gamma(2)\beta^2} X_i e^{-X_i/\beta} = \frac{1}{\Gamma(2)\beta^{2n}} \prod_{i=1}^n X_i e^{-\frac{1}{\beta} \sum_{i=1}^n X_i}$$

The log likelihood function is:

$$\ln L(\beta) = -n(\ln \Gamma(2) + 2 \ln \beta) + \ln \prod_{i=1}^n X_i - \frac{1}{\beta} \sum_{i=1}^n X_i$$

Take the derivative of the log likelihood function, we have,

$$\frac{d}{d\beta} \ln L(\beta) = -\frac{2n}{\beta} + \frac{1}{\beta^2} \sum_{i=1}^n X_i$$

Let this derivative equal to zero. We have,

$$\hat{\beta} = \frac{1}{2n} \sum_{i=1}^n X_i = \frac{1}{2} \bar{X}$$

- (b) Since $\mu = E[X] = \alpha\beta = 2\beta$

We take the estimator of μ as $\hat{\mu} = 2\hat{\beta}$

$$E[\hat{\mu}] = E[2\hat{\beta}] = 2E[\hat{\beta}] = 2E\left[\frac{1}{2}\bar{X}\right] = E[\bar{X}] = \mu$$

Hence, this is an unbiased estimator.

- (c) we calculated that $\bar{X} = 70.5$. Therefore, $\hat{\beta} = 35.25$

- (d) $\bar{\mu} = 2\hat{\beta} = 70.5$

37. (a) Normal distribution: $\mu = 2, \sigma^2 = 9$
 (b) Normal distribution: $\mu = 0, \sigma^2 = 4$
 (c) Geometric distribution: $p = 0.25$
 (d) Binomial distribution: $n = 5, p = 0.5$
 (e) Poisson distribution: $k = 6$
 (f) Gamma distribution: $\alpha = 5, \beta = 3$
 (g) Chi-squared distribution: $\gamma = 16$
 (h) Exponential distribution: $\beta = 0.5$

$$39. m_Y(t) = E[e^{Yt}] = E[e^{(a_0 + a_1 X_1 + a_2 X_2 + \dots + a_n X_n)t}] = E[e^{a_0 t} e^{a_1 X_1 t} e^{a_2 X_2 t} \dots e^{a_n X_n t}]$$

Since X_1, X_2, \dots, X_n are independent. $e^{a_1 X_1 t}, e^{a_2 X_2 t}, \dots, e^{a_n X_n t}$ are also independent. Therefore, we have,

$$m_Y(t) = E[e^{a_0 t} e^{a_1 X_1 t} e^{a_2 X_2 t} \dots e^{a_n X_n t}] = E[e^{a_0 t}] E[e^{a_1 X_1 t}] \dots E[e^{a_n X_n t}]$$

$$= e^{a_0 t} m_{X_1}(a_1 t) m_{X_2}(a_2 t) \dots m_{X_n}(a_n t) = e^{a_0 t} \prod_{i=1}^n m_{X_i}(a_i t)$$

41. We have that $m_{X_i}(t) = e^{\mu_i t + \sigma_i^2 t^2 / 2}, i = 1, 2, \dots, n$. Then, applying the result from question 39 gives:

$$m_Y(t) = e^{a_0 t} \prod_{i=1}^n m_{X_i}(a_i t) = e^{a_0 t} e^{\mu_1 a_1 t + \sigma_1^2 a_1^2 t^2 / 2} \dots e^{\mu_n a_n t + \sigma_n^2 a_n^2 t^2 / 2}$$

$$= e^{(a_0 + \mu_1 a_1 + \dots + \mu_n a_n)t + (\sigma_1^2 a_1^2 + \dots + \sigma_n^2 a_n^2)t^2 / 2}$$

Hence, Y is following a normal distribution with

$$\mu = a_0 + \sum_{i=1}^n a_i \mu_i, \sigma^2 = \sum_{i=1}^n a_i^2 \sigma_i^2$$

$$42. \bar{X} = \frac{1}{n} X_1 + \frac{1}{n} X_2 + \dots + \frac{1}{n} X_n$$

In reference to Exercise 41, $a_0 = 0$ and $a_i = \frac{1}{n}$ for $i = 1, 2, \dots, n$.

$$\text{Thus, } \mu_{\bar{X}} = \sum_{i=1}^n \frac{1}{n} \mu = \frac{1}{n} \cdot n \mu = \mu \text{ and } \sigma_{\bar{X}}^2 = \sum_{i=1}^n \frac{1}{n^2} \sigma^2 = \frac{1}{n^2} \cdot n \sigma^2 = \frac{\sigma^2}{n}$$

Since X_1, X_2, \dots, X_n constitute a random sample from a normal population, and thus are independent normal random variables, \bar{X} is normal.

$$43. \text{ We have, } m_{X_1}(t) = (1 - 2t)^{-5/2}, m_{X_2}(t) = (1 - 2t)^{-10/2}$$

Since X_1 and X_2 are independent, we have

$$m_{X_1 + X_2}(t) = m_{X_1}(t) m_{X_2}(t) = (1 - 2t)^{-15/2}$$

Hence, $X_1 + X_2$ is a chi-squared R.V. with degree of freedom of 15

44. Since X_i is a chi-squared random variable with parameter γ_i ,

$$m_{X_i}(t) = (1 - 2t)^{\gamma_i / 2}$$

$$\text{Using Exercise 39, } m_Y(t) = \prod_{i=1}^n m_{X_i}(t)$$

$$\text{Substitution then yields } m_Y(t) = \prod_{i=1}^n (1 - 2t)^{\gamma_i / 2} = (1 - 2t)^{\sum \gamma_i / 2}$$

which is the moment generating function of a chi-squared random variable with

$$\text{parameter } \gamma = \sum_{i=1}^n \gamma_i.$$

45. We have, $X_i \sim N(\mu, \sigma^2)$, $i = 1, 2, \dots, n$. Then, $\frac{X_i - \mu}{\sigma} \sim N(0, 1)$. i.e., $\frac{X_i - \mu}{\sigma}$ is a standard normal R.V.

Hence, $\left(\frac{X_i - \mu}{\sigma}\right)^2 \sim \chi_1^2$

Applying the results from Q44 gives,

$$\sum_{i=1}^n \left(\frac{X_i - \mu}{\sigma}\right)^2 \sim \chi_n^2$$

i.e., $\sum_{i=1}^n \left(\frac{X_i - \mu}{\sigma}\right)^2$ has a chi-squared distribution with n degree of freedom.