

ESE 326 Practice Exam for the Final F2017

1. Consider the random variables (X, Y) with following joint density

$$f_{XY}(x, y) = cxy, \quad c \text{ is a constant}, \quad 0 < y < x < 1$$

- (1) Find the value of c so that f_{XY} is a valid density function
- (2) Find the marginal density for X . Find $E[X]$ and $E[X^2]$
- (3) Find the marginal density for Y . Find $E[Y]$ and $E[Y^2]$
- (4) Are X and Y independent? Why?
- (5) Find the curve of regression of X on Y . Is the regression linear?

2. let \bar{X}_1 and \bar{X}_2 be the sample means based on independent samples of size n_1 and n_2 drawn from normal distributions X_1 and X_2 with means μ_1 and μ_2 and variances σ_1^2 and σ_2^2 respectively.

- (1) Show that $\bar{X}_1 - \bar{X}_2$ is an unbiased estimator of $\mu_1 - \mu_2$
- (2) Show that $\bar{X}_1 - \bar{X}_2$ is a normal random variable with mean $\mu_1 - \mu_2$ and variance $\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}$

Let's assume that $\sigma_1^2 = \sigma_2^2 = \sigma^2$ for the following questions.

- (3) The pooled variance is defined as: $S_p^2 = \frac{(n_1-1)S_1^2 + (n_2-1)S_2^2}{n_1+n_2-2}$

Show that $\frac{(n_1+n_2-2)S_p^2}{\sigma^2} \sim \chi_{n_1+n_2-2}^2$

- (4) Show that $\frac{(\bar{X}_1 - \bar{X}_2) - (\mu_1 - \mu_2)}{\sqrt{S_p^2(\frac{1}{n_1} + \frac{1}{n_2})}} \sim T_{n_1+n_2-2}$


3. The following are two random samples of sizes n_1 and n_2 from two normal populations with means μ_1 and μ_2 and variances σ_1^2 and σ_2^2 respectively.

Population 1	Population 2
$n_1 = 11$	$n_2 = 9$
$\bar{x}_1 = 18.55$	$\bar{x}_2 = 14.67$
$S_1^2 = 10.27$	$S_2^2 = 7.75$

- (1) Test the equality of the variances at the significance level of 0.2
- (2) Find the 95% confidence interval on $\mu_1 - \mu_2$. Based on this interval, is there evidence that a real difference exists?
- (3) Test the hypothesis $H_0: \mu_1 = \mu_2$ $H_1: \mu_1 \neq \mu_2$ by using p test with significant level of $\alpha = 0.1$.
- (4) Verify the result of (3) by finding the rejection region of the test statistics.

The tables of T distribution and F distribution are attached.

TABLE VI
T distribution




Column heading = cumulative probability

Row heading = degrees of freedom

Row ∞ = standard normal values

γ	$P[T_\gamma \leq t]$								
	.6	.75	.9	.95	.975	.99	.995	.999	.9995
1	0.325	1.000	3.078	6.314	12.706	31.821	63.657	318.317	636.607
2	0.289	0.816	1.886	2.920	4.303	6.965	9.925	22.327	31.598
3	0.277	0.765	1.638	2.353	3.182	4.541	5.841	10.215	12.924
4	0.271	0.741	1.533	2.132	2.776	3.747	4.604	7.173	8.610
5	0.267	0.727	1.476	2.015	2.571	3.365	4.032	5.893	6.869
6	0.265	0.718	1.440	1.943	2.447	3.143	3.707	5.208	5.959
7	0.263	0.711	1.415	1.895	2.365	2.998	3.499	4.785	5.408
8	0.262	0.706	1.397	1.860	2.306	2.896	3.355	4.501	5.041
9	0.261	0.703	1.383	1.833	2.262	2.821	3.250	4.297	4.781
10	0.260	0.700	1.372	1.812	2.228	2.764	3.169	4.144	4.587
11	0.260	0.697	1.363	1.796	2.201	2.718	3.106	4.025	4.437
12	0.259	0.695	1.356	1.782	2.179	2.681	3.055	3.930	4.318
13	0.259	0.694	1.350	1.771	2.160	2.650	3.012	3.852	4.221
14	0.258	0.692	1.345	1.761	2.145	2.624	2.977	3.787	4.140
15	0.258	0.691	1.341	1.753	2.131	2.602	2.947	3.733	4.073
16	0.258	0.690	1.337	1.746	2.120	2.583	2.921	3.686	4.015
17	0.257	0.689	1.333	1.740	2.110	2.567	2.898	3.646	3.965
18	0.257	0.688	1.330	1.734	2.101	2.552	2.878	3.611	3.922
19	0.257	0.688	1.328	1.729	2.093	2.539	2.861	3.579	3.883
20	0.257	0.687	1.325	1.725	2.086	2.528	2.845	3.552	3.850
21	0.257	0.686	1.323	1.721	2.080	2.518	2.831	3.527	3.819
22	0.256	0.686	1.321	1.717	2.074	2.508	2.819	3.505	3.792
23	0.256	0.685	1.319	1.714	2.069	2.500	2.807	3.485	3.768
24	0.256	0.685	1.318	1.711	2.064	2.492	2.797	3.467	3.745
25	0.256	0.684	1.316	1.708	2.060	2.485	2.787	3.450	3.725
26	0.256	0.684	1.315	1.706	2.056	2.479	2.779	3.435	3.707
27	0.256	0.684	1.314	1.703	2.052	2.473	2.771	3.421	3.690
28	0.256	0.683	1.313	1.701	2.048	2.467	2.763	3.408	3.674
29	0.256	0.683	1.311	1.699	2.045	2.462	2.756	3.396	3.659
30	0.256	0.683	1.310	1.697	2.042	2.457	2.750	3.385	3.646
31	0.256	0.682	1.309	1.696	2.040	2.453	2.744	3.375	3.633
32	0.255	0.682	1.309	1.694	2.037	2.449	2.738	3.365	3.622
33	0.255	0.682	1.308	1.692	2.035	2.445	2.733	3.356	3.611
34	0.255	0.682	1.307	1.691	2.032	2.441	2.728	3.348	3.601
35	0.255	0.682	1.306	1.690	2.030	2.438	2.724	3.340	3.591
36	0.255	0.681	1.306	1.688	2.028	2.434	2.719	3.333	3.582
37	0.255	0.681	1.305	1.687	2.026	2.431	2.715	3.326	3.574
38	0.255	0.681	1.304	1.686	2.024	2.429	2.712	3.319	3.566
39	0.255	0.681	1.304	1.685	2.023	2.426	2.708	3.313	3.558
40	0.255	0.681	1.303	1.684	2.021	2.423	2.704	3.307	3.551
41	0.255	0.681	1.303	1.683	2.020	2.421	2.701	3.301	3.544
42	0.255	0.680	1.302	1.682	2.018	2.418	2.698	3.296	3.538
43	0.255	0.680	1.302	1.681	2.017	2.416	2.695	3.291	3.532
44	0.255	0.680	1.301	1.680	2.015	2.414	2.692	3.286	3.526

TABLE IX
F distributionColumn heading = numerator
degrees of freedomRow heading = denominator
degrees of freedomPoints given are $f_{.10}$ pointsFor degrees of freedom > 120,
use row or column 120

$$P[F_{\gamma_1, \gamma_2} \leq f] = .90$$

$\gamma_2 \backslash \gamma_1$	1	2	3	4	5	6	7	8	9	10
1	39.862	49.500	53.593	55.833	57.240	58.204	58.906	59.439	59.857	60.195
2	8.526	9.000	9.162	9.243	9.293	9.326	9.349	9.367	9.381	9.392
3	5.538	5.462	5.391	5.343	5.309	5.285	5.266	5.252	5.240	5.231
4	4.545	4.325	4.191	4.107	4.051	4.010	3.979	3.955	3.936	3.920
5	4.060	3.780	3.619	3.520	3.453	3.405	3.368	3.339	3.316	3.297
6	3.776	3.463	3.289	3.181	3.108	3.055	3.014	2.983	2.958	2.937
7	3.589	3.257	3.074	2.961	2.883	2.827	2.785	2.752	2.725	2.703
8	3.458	3.113	2.924	2.806	2.726	2.668	2.624	2.589	2.561	2.538
9	3.360	3.006	2.813	2.693	2.611	2.551	2.505	2.469	2.440	2.416
10	3.285	2.924	2.728	2.605	2.522	2.461	2.414	2.377	2.347	2.323
11	3.225	2.860	2.660	2.536	2.451	2.389	2.342	2.304	2.274	2.248
12	3.177	2.807	2.606	2.480	2.394	2.331	2.283	2.245	2.214	2.188
13	3.136	2.763	2.560	2.434	2.347	2.283	2.234	2.195	2.164	2.138
14	3.102	2.726	2.522	2.395	2.307	2.243	2.193	2.154	2.122	2.095
15	3.073	2.695	2.490	2.361	2.273	2.208	2.158	2.119	2.086	2.059
16	3.048	2.668	2.462	2.333	2.244	2.178	2.128	2.088	2.055	2.028
17	3.026	2.645	2.437	2.308	2.218	2.152	2.102	2.061	2.028	2.001
18	3.007	2.624	2.416	2.286	2.196	2.130	2.079	2.038	2.005	1.977
19	2.990	2.606	2.397	2.266	2.176	2.109	2.058	2.017	1.984	1.956
20	2.975	2.589	2.380	2.249	2.158	2.091	2.040	1.999	1.965	1.937
21	2.961	2.575	2.365	2.233	2.142	2.075	2.023	1.982	1.948	1.920
22	2.949	2.561	2.351	2.219	2.128	2.061	2.008	1.967	1.933	1.904
23	2.937	2.549	2.339	2.207	2.115	2.047	1.995	1.953	1.919	1.890
24	2.927	2.538	2.327	2.195	2.103	2.035	1.983	1.941	1.906	1.877
25	2.918	2.528	2.317	2.184	2.092	2.024	1.971	1.929	1.895	1.866
26	2.909	2.519	2.307	2.174	2.082	2.014	1.961	1.919	1.884	1.855
27	2.901	2.511	2.299	2.165	2.073	2.005	1.952	1.909	1.874	1.845
28	2.894	2.503	2.291	2.157	2.064	1.996	1.943	1.900	1.865	1.836
29	2.887	2.495	2.283	2.149	2.057	1.988	1.935	1.892	1.857	1.827
30	2.881	2.489	2.276	2.142	2.049	1.980	1.927	1.884	1.849	1.819
31	2.875	2.482	2.270	2.136	2.042	1.973	1.920	1.877	1.842	1.812
32	2.869	2.477	2.263	2.129	2.036	1.967	1.913	1.870	1.835	1.805
33	2.864	2.471	2.258	2.123	2.030	1.961	1.907	1.864	1.828	1.799
34	2.859	2.466	2.252	2.118	2.024	1.955	1.901	1.858	1.822	1.793
35	2.855	2.461	2.247	2.113	2.019	1.950	1.896	1.852	1.817	1.787
36	2.850	2.456	2.243	2.108	2.014	1.945	1.891	1.847	1.811	1.781
37	2.846	2.452	2.238	2.103	2.009	1.940	1.886	1.842	1.806	1.776
38	2.842	2.448	2.234	2.099	2.005	1.935	1.881	1.838	1.802	1.772
39	2.839	2.444	2.230	2.095	2.001	1.931	1.877	1.833	1.797	1.767
40	2.835	2.440	2.226	2.091	1.997	1.927	1.873	1.829	1.793	1.763
50	2.809	2.412	2.197	2.061	1.966	1.895	1.840	1.796	1.760	1.730
60	2.791	2.393	2.177	2.041	1.946	1.875	1.819	1.775	1.738	1.707
120	2.748	2.347	2.130	1.992	1.896	1.824	1.767	1.722	1.684	1.652