

Distribution of a linear combination of independent random variables:

Let X_1, X_2, \dots, X_n be a collection of independent random variables with mgfs $m_{X_i}(t), i = 1, 2, 3, \dots, n$.

Let $Y = a_0 + a_1X_1 + a_2X_2 + \dots + a_nX_n$, a_0, a_1, \dots, a_n are real numbers. Then the moment generating function for Y is given by:

$$m_Y(t) = e^{a_0t} \prod_{i=1}^n m_{X_i}(a_it)$$

Practice Example: Let X_1 and X_2 be independent normal random variables with means 2 and 5 and variances 9 and 1 respectively. Let $Y = 3X_1 + 6X_2 - 8$. What is the distribution of Y ?

$$m_{X_1}(t) = e^{2t + \frac{9}{2}t^2}; m_{X_2}(t) = e^{5t + \frac{1}{2}t^2}$$

$$m_Y(t) = e^{-8t} m_{X_1}(3t) m_{X_2}(6t) = e^{-8t} e^{6t + \frac{81}{2}t^2} e^{30t + 18t^2} = e^{28t + \frac{117}{2}t^2}$$

Hence, Y is a normal random variable with $\mu_Y = 28$ and $\sigma_Y^2 = 117$, i.e., $Y \sim N(28, 117)$

To verify the mean and variance of Y , we have,

$$E[Y] = E[3X_1 + 6X_2 - 8] = 3E[X_1] + 6E[X_2] - 8 = 3 \times 2 + 6 \times 5 - 8 = 28$$

$$VarY = Var(3X_1 + 6X_2 - 8) = 9VarX_1 + 36VarX_2 = 9 \times 9 + 36 = 117$$

Theorem 7.3.4: Let X_1, X_2, \dots, X_n be a random sample of size n from a normal distribution with mean μ and variance σ^2 . Then \bar{X} is normally distributed with mean μ and variance $\frac{\sigma^2}{n}$.

Proof:

Since X_1, X_2, \dots, X_n is a random sample of size n from a normal distribution with mean μ and variance σ^2 , we have, $m_{X_i}(t) = e^{\mu t + \frac{1}{2}\sigma^2 t^2}$, $i = 1, 2, \dots, n$

$$\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i = \frac{X_1}{n} + \frac{X_2}{n} + \dots + \frac{X_n}{n}$$

The moment generating function of \bar{X} can be found as:

$$\begin{aligned} m_{\bar{X}}(t) &= m_{X_1} \left(\frac{t}{n} \right) m_{X_2} \left(\frac{t}{n} \right) \dots m_{X_n} \left(\frac{t}{n} \right) \\ &= e^{\frac{\mu t}{n} + \frac{1}{2}\sigma^2 \left(\frac{t}{n} \right)^2} e^{\frac{\mu t}{n} + \frac{1}{2}\sigma^2 \left(\frac{t}{n} \right)^2} \dots e^{\frac{\mu t}{n} + \frac{1}{2}\sigma^2 \left(\frac{t}{n} \right)^2} = e^{\mu t + \frac{1}{2} \frac{\sigma^2}{n} t^2} \end{aligned}$$

Hence, we can claim that $\bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right)$

Theorem $Z \sim N(0,1)$. Let $X = Z^2$. Then $X \sim X_1^2$ (chi-squared distribution with degree of 1)

Proof:

Recall a Gamma random variable X with parameters α and β has the density function:

$$f(x) = \frac{1}{\Gamma(\alpha)\beta^\alpha} x^{\alpha-1} e^{-\frac{x}{\beta}}, \quad x > 0, \alpha > 0, \beta > 0$$

When $\beta = 2$ and $\alpha = \frac{\gamma}{2}$, we have a chi-squared distribution with γ degree of freedom, X_γ^2

$$f(x) = \frac{x^{\frac{\gamma}{2}-1} e^{-\frac{x}{2}}}{\Gamma(\frac{\gamma}{2}) 2^{\frac{\gamma}{2}}}$$

When $\gamma = 1$, the density function of X_1^2 is:

$$f(x) = \frac{x^{-1/2} e^{-\frac{x}{2}}}{\Gamma(\frac{1}{2}) 2^{1/2}} = \frac{x^{-1/2} e^{-\frac{x}{2}}}{\sqrt{2\pi}}, \quad \Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$$

Now, we have to prove that $X = Z^2$ has the same density function of X_1^2

Proof (continued):

First, let's find the cumulative distribution function of X .

$$\begin{aligned} F_X(x) &= P[X \leq x] = P[Z^2 \leq x] = P[-\sqrt{x} \leq Z \leq \sqrt{x}] \\ &= P[Z \leq \sqrt{x}] - P[Z \leq -\sqrt{x}] = F_Z(\sqrt{x}) - F_Z(-\sqrt{x}) \end{aligned}$$

Then, to find the density function $f_X(x)$, we take the derivative of $F_X(x)$ w.r.t. x

$$f_X(x) = \frac{1}{2}x^{-\frac{1}{2}}f_Z(\sqrt{x}) + \frac{1}{2}x^{-\frac{1}{2}}f_Z(-\sqrt{x})$$

Recall that the density function of a standard normal random variable is $f_Z(x) = \frac{1}{\sqrt{2\pi}}e^{-\frac{1}{2}x^2}$. Then,

$$f_X(x) = \frac{1}{2}x^{-\frac{1}{2}}\frac{1}{\sqrt{2\pi}}e^{-\frac{1}{2}x} + \frac{1}{2}x^{-\frac{1}{2}}\frac{1}{\sqrt{2\pi}}e^{-\frac{1}{2}x} = \frac{x^{-\frac{1}{2}}e^{-\frac{1}{2}x}}{\sqrt{2\pi}}$$

Therefore, $X \sim X_1^2$, i.e., $Z^2 \sim X_1^2$

Theorem *Distribution of a sum of independent chi-squared random variables*

Let X_1, X_2, \dots, X_n be independent chi-squared random variables with $\gamma_1, \gamma_2, \dots, \gamma_n$ degree of freedom respectively. Let $Y = X_1 + X_2 + \dots + X_n$. Then $Y \sim X_\gamma^2$, $\gamma = \sum_{i=1}^n \gamma_i$

Proof: Recall the mgf of a Gamma R.V. with parameters α and β is $(1 - \beta t)^{-\alpha}$. A chi-squared R.V. is a special case of Gamma R.V. with $\beta = 2$ and $\alpha = \frac{\gamma}{2}$. Hence the mgf of a chi-squared R.V. is $(1 - 2t)^{-\gamma/2}$

We have,

$$m_{X_i}(t) = (1 - 2t)^{-\gamma_i/2}, \quad i = 1, 2, \dots, n$$

Then,

$$m_Y(t) = \prod_{i=1}^n m_{X_i}(t) = \prod_{i=1}^n (1 - 2t)^{-\gamma_i/2} = (1 - 2t)^{-\sum_{i=1}^n \gamma_i/2}$$

Therefore,

$$Y \sim X_\gamma^2, \quad \gamma = \sum_{i=1}^n \gamma_i$$

Theorem Let X_1, X_2, \dots, X_n be a random sample of size n from a normal distribution with mean μ and variance σ^2 . Then

$$\sum_{i=1}^n \frac{(X_i - \mu)^2}{\sigma^2} \sim \chi_n^2$$

Proof: Since X_1, X_2, \dots, X_n is a random sample of size n from a normal distribution with mean μ and variance σ^2 . We have,

$$X_i \sim N(\mu, \sigma^2), \quad \frac{X_i - \mu}{\sigma} \sim N(0,1), \quad i = 1, 2, \dots, n$$

Then,

$$\left(\frac{X_i - \mu}{\sigma} \right)^2 \sim \chi_1^2$$




Therefore,

$$\sum_{i=1}^n \left(\frac{X_i - \mu}{\sigma} \right)^2 = \sum_{i=1}^n \frac{(X_i - \mu)^2}{\sigma^2} = \frac{\sum_{i=1}^n (X_i - \mu)^2}{\sigma^2} \sim \chi_n^2$$

The Central Limit Theorem




Theorem 7.4.2: Let X_1, X_2, \dots, X_n be a random sample of size n from a distribution with mean μ and variance σ^2 . Then for large n , \bar{X} is approximately normal with mean μ and variance $\frac{\sigma^2}{n}$. The random variable $\frac{\bar{X} - \mu}{\sigma/\sqrt{n}}$ is approximately standard normal.

- ***This applies to any population (not only normal population!!!)***
- For most population, if the sample size is greater than 30, the central limit theorem approximation is good.
- If the sample is drawn from a nearly symmetric distribution, the normal approximation can be good for a fairly small value of n ;
- If the population is heavily skewed, a fairly large n may be necessary.






Value of sample mean

Value of sample mean



Value of sample mean

Value of sample mean



Value of sample mean

Value of sample mean

The Central Limit Theorem

Example: Let X_1, X_2, \dots, X_{100} be a random sample of size 100 from a Gamma distribution with $\alpha = 5$ and $\beta = 3$. Find the distribution of \bar{X} using the moment generating function method and the central limit theorem.

Solution: (the moment generating function method)

Since $X_i, i = 1, 2, \dots, 100$ are the random samples from a Gamma distribution with $\alpha = 5$ and $\beta = 3$, The moment generating function of X_i is $(1 - 3t)^{-5}$. Then the mgf of $\bar{X} = \frac{1}{100} \sum_{i=1}^{100} X_i$ is:

$$m_{\bar{X}}(t) = \prod_{i=1}^{100} m_{X_i}\left(\frac{t}{100}\right) = (1 - 0.03t)^{-500}$$

Hence, \bar{X} is a Gamma R.V. with $\alpha = 500$ and $\beta = 0.03$. i.e.,

$$f_{\bar{X}}(x) = \frac{1}{\Gamma(500) \times 0.03^{500}} x^{499} e^{-\frac{x}{0.03}}$$

This is very hard to evaluate!

The Central Limit Theorem

Example: Let X_1, X_2, \dots, X_{100} be a random sample of size 100 from a Gamma distribution with $\alpha = 5$ and $\beta = 3$. Find the distribution of \bar{X} using the moment generating function method and the central limit theorem. Find the probability that \bar{X} is at most 14.

Solution: (the central limit theorem method)

According to the central limit theorem, \bar{X} is approximately following a normal distribution with mean μ and variance $\frac{\sigma^2}{100}$, where μ and σ^2 are the mean and variance of the population variable.

Since the population variable follows a Gamma distribution with $\alpha = 5$ and $\beta = 3$, we have,

$$\mu = \alpha\beta = 15; \sigma^2 = \alpha\beta^2 = 45$$

$$\bar{X} \sim N(15, 0.45)$$

$$P[\bar{X} \leq 14] = P[\bar{X} - 15 \leq -1] = P\left[\frac{\bar{X} - 15}{\sqrt{0.45}} \leq -\frac{1}{\sqrt{0.45}}\right] = P[Z \leq -1.49] = 0.0681$$