

UNIVERSITY OF HAMBURG

MASTER THESIS

Masterthesis_doc

Author:

Lona Frießner

Supervisor:

Prof. Dr. Simon Grund

*A thesis submitted in fulfillment of the requirements
for the degree of Master of Science*

in the

Institute of Psychology
Department of Psychology with focus on Quantitative Methods



Universität Hamburg
DER FORSCHUNG | DER LEHRE | DER BILDUNG

May 05, 2026

UNIVERSITY OF HAMBURG

Abstract

Institute of Psychology

Department of Psychology with focus on Quantitative Methods

Master of Science

Masterthesis__doc

by Lona Frießner

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Aliquam quis accumsan ante. Quisque lorem metus, varius id urna eget, lacinia dapibus sem. Etiam laoreet, quam ac mollis congue, arcu leo dictum neque, nec euismod sem enim luctus odio. Donec condimentum tortor sit amet mollis volutpat. Donec ornare libero vel velit malesuada consectetur. Vestibulum in sem non justo dignissim congue at quis erat. Integer quis erat vitae mi maximus fringilla tristique nec odio. Morbi non ipsum sapien. Vestibulum tortor est, ultricies in eros et, bibendum iaculis justo. Cras pellentesque enim quam, id pretium lacus lacinia non. Integer velit neque, ultrices a malesuada vel, imperdiet quis enim. Quisque eu facilisis urna, ut faucibus lorem. Donec mollis turpis sed arcu venenatis interdum. Nulla facilisis tortor ac scelerisque consequat.

Table of contents

Abstract	iii
1 Introduction	1
2 Theory	3
2.1 Multilevel models	3
2.2 Small sample sizes in multilevel models	6
2.3 Missing data in multilevel models	6

List of Figures

List of Tables

Chapter 1

Introduction

Chapter 2

Theory

2.1 Multilevel models

Example

Imagine you are interested in the musical self-concept of young adults and whether it is related to how many hours they practice their instrument each week.

To investigate this question, you collect data from several youth ensembles. You recruit 15 ensembles, each consisting of approximately 10 players. Every player completes a questionnaire assessing their musical self-concept as well as their weekly practice time.

At first glance, this seems like a simple dataset of 150 individuals. But is it really that simple?

Players from the same ensemble rehearse together, share teachers, and experience similar social and musical environments. Their responses may therefore be more similar within ensembles than between ensembles.

2.1.1 The idea behind multilevel models

Populations like this, where units of interest (e.g., music players, students, employees) are grouped into higher-level units (e.g., ensembles, classes, firms), are common in the human social life and therefore in psychological studies.

The resulting data is called *nested* or *clustered* (Snijders & Bosker, 2012, p. 7).

From a sampling perspective, a so-called *multistage sampling process* (sample groups first, then individuals inside these groups¹) is not independent: After selecting a group, you will try to sample all of the members of this group, making it more likely to select one person in this group over others in non-selected groups (Snijders & Bosker, 2012, p. 7). More importantly, individuals inside the same group typically share contextual influences (e.g., the same ensemble teacher, the same class room) and thus will give more similar responses in surveys and behavior. As a consequence, the basic assumption of independence in traditional linear modeling is violated, because error terms will be correlated (Raudenbush & Bryk, 2010, p. xx).

One well-suited and popular possibility to account for this fact is the use of *multilevel models* (also called *hierarchical linear models* or *mixed models*).

¹In this thesis, the term ‘groups’ refers to the higher-level units of the data and level-1 units are referred to as ‘persons’. However, it is important to note that these models can also be used for longitudinal research, where measurement points are on level 1 and individuals are the level-2 units. Furthermore, more than two levels are also possible, but are not discussed here.

2.1.2 The concept of multilevel regression models

The most basic aim of regression analysis is the explanation of variance in an outcome of interest (Snijders & Bosker, 2012, p. 80). Measurements of other associated variables are included in the model in an additive way to predict changes in the outcome, in other words: they explain variance in Y . The assumption of the classical linear model is that after inclusion of all *systematic* influences on Y , the deviations of the measured Y from the model-predicted Y are *random*. Specifically, the residuals are assumed to have expectation zero ($E(e_i) = 0$), constant variance across observations (homoscedasticity; $Var(e_i) = \sigma^2$), and to be independent of one another ($Cov(e_i, e_{i'}) = 0$ for $i \neq i'$) (Fahrmeir et al., 2021, p. ~87).

In single-level regression, the residual/error term e_{ij} captures the *unexplained variance* in Y , that is differences in the outcome from the prediction that the model made (Leonhart, 2013, p. 319):

$$Y_i = \mu_i + e_i, \quad \mu_i = \underbrace{\beta_0}_{\text{Intercept}} + \underbrace{\beta_1}_{\text{Slope}} X_i$$

Because in nested data there are populations on more than one level, multilevel models (MLM) extend regular linear regression by adding level-2 *random effects* to explain the variability between groups that cannot be accounted for by predictors (Snijders & Bosker, 2012, p. 80). The level-1 regression equation remains similar, but the regression coefficients are now group-dependent:

$$Y_{ij} = \mu_{ij} + e_{ij}, \quad \mu_{ij} = \beta_{0j} + \beta_{1j} X_{ij}$$

These group-dependent coefficients can be separated into fixed and random parts:

$$\beta_{0j} = \underbrace{\gamma_{00}}_{\text{Fixed part}} + \underbrace{u_{0j}}_{\text{Random part}}, \quad \beta_{1j} = \underbrace{\gamma_{10}}_{\text{Fixed part}} + \underbrace{u_{1j}}_{\text{Random part}}$$

The random parts reflect differences between groups, in their average Y -value (u_{0j}) and in the effect of covariates on Y (u_{1j}).

Each group has its own intercept and slope, respectively. Substitution of all single components leads to the combined model

$$Y_{ij} = \underbrace{\gamma_{00} + \gamma_{10} X_{ij}}_{\text{Fixed part}} + \underbrace{u_{0j} + u_{1j} X_{ij} + e_{ij}}_{\text{Random part}}$$

Y_{ij} : Outcome variable for individual i in group j

X_{ij} : Level-1 (within-group) predictor

γ_{00} : Fixed intercept; average value of Y_{ij} across all groups when $X_{ij} = 0$

γ_{10} : Fixed slope; average effect of X_{ij} on Y_{ij} across all groups

u_{0j} : Random intercept for group j ; group-specific deviation from the overall intercept

γ_{00}

u_{1j} : Random slope for group j ; group-specific deviation from the average effect γ_{10}

e_{ij} : Level-1 residual; individual deviation from the group-specific predicted value

Similar distributional assumptions are made for the level-2 random effects as for the residuals in single-level regression: they have mean zero, constant variances across groups, and are independent across clusters and independent of the level-1 residuals. However, random effects within the same cluster may be correlated and are often

expected to do so (Snijders & Bosker, 2012, p. 76). Additionally, residuals and random effects are often assumed to follow a normal distribution for convenience, as it facilitates the likelihood-based estimation of the unknown parameters (Fahrmeir et al., 2021, p. 388). In summary:

$$\begin{aligned} \begin{pmatrix} u_{0j} \\ u_{1j} \end{pmatrix} &\sim \mathcal{N} \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \mathbf{T} \right), & \mathbf{T} &= \begin{pmatrix} \tau_0^2 & \tau_{01} \\ \tau_{01} & \tau_1^2 \end{pmatrix} \\ e_{ij} &\sim \mathcal{N}(0, \sigma^2) \\ \text{Cov}(u_{0j}, e_{ij}) &= 0, & \text{Cov}(u_{1j}, e_{ij}) &= 0 \end{aligned}$$

2.1.3 The random intercept model with context effects

A simpler, but often used version of the random coefficients model is the *random intercept model*. This model allows for baseline differences in the outcome between groups (random intercept), but the strength of the relationship between predictors and the outcome is assumed to be constant (i.e., $u_{1j} = 0$). The strength of the effects of predictors is assumed to be the same between groups.

In such models, it is of special interest how the group context influences the outcome. The random intercept's variance indicates whether and how much groups vary in their baseline level, but to explain these differences, a certain kind of variables can be introduced: contextual effects. Adding a predictor X 's mean as an explanatory context variable, the within-group regression coefficient may vary from the between-group regression coefficient, making it more flexible and realistic (Snijders & Bosker, 2012, p. 57).

In these cases, *group-mean-centering* of the level-1 predictor is often applied. Instead of using X_{ij} directly, the centered predictor $(X_{ij} - \bar{X}_{.j})$ is entered into the model. This decomposes within- and between- group effects and facilitates their interpretation (Snijders & Bosker, 2012, p. 58). Such a random intercept model can look like this:

$$Y_{ij} = \gamma_{00} + \gamma_{10}(X_{ij} - \bar{X}_{.j}) + \gamma_{01}\bar{X}_{.j} + u_{0j} + e_{ij}$$

2.1.4 The intraclass correlation coefficient (ICC) in the random intercept model

In the RI model, the variance components are given by

$$u_{0j} \sim \mathcal{N}(0, \tau_0^2), \quad e_{ij} \sim \mathcal{N}(0, \sigma^2), \quad \text{Cov}(e_{ij}, u_{0j}) = 0.$$

From these variance components, the proportion of variability that is attributable to between-group differences can be calculated. This quantity is referred to as the *intraclass correlation coefficient* (ICC). The ICC is equal to the correlation between the outcomes of two randomly chosen individuals from the same group (Snijders & Bosker, 2012, p. 18). In models including predictors, the ICC is interpreted conditionally on the predictors, that is, as the correlation between two individuals from the same group who share identical predictor values.

Formally, the ICC is defined as

$$ICC = \frac{\tau_0^2}{\tau_0^2 + \sigma^2}$$

The ICC is an important diagnostic metric because it indicates whether the use of a multilevel model is advantageous. If $ICC = 0$, no between-group variance is present and individuals within the same group are not more similar to each other than to individuals from different groups (conditional on the predictors included in the model). In this case, the independence assumption of classical regression is not violated and a conventional single-level regression model is sufficient.

Conversely, larger ICC values indicate increasing similarity within groups and thus statistical dependency of observations. Ignoring this dependency and applying a classical linear model leads to biased standard errors, confidence intervals, and hypothesis tests (Fahrmeir et al., 2021, p. 370).

2.2 Small sample sizes in multilevel models

In the preceding chapter, the theoretical foundation and parameters of the multilevel model were introduced. In practice, however, these parameters are unknown and only exist in the theoretical realm as ‘true’ values. The fixed effects γ and variance components σ^2 and τ^2 must therefore be estimated from observed data. Based on these estimates, inferences about the underlying population parameters are made. In frequentist approaches, estimation procedures of multilevel models are based on ‘asymptotic theory’ and assume a sufficiently large sample size. In practice, this assumption is often difficult to meet, especially because estimation depends not only on the number of individual observations (Level 1), but also on the number of clusters (Level 2), which are often difficult or costly to obtain (McNeish, 2017). Fortunately, statistics has solutions at hand for those cases. The main challenges of small-sample multilevel models and possible solutions are presented in the following section.

2.2.1 Main problems

In multilevel modeling, statistical inference is primarily determined by three key components. If these are estimated accurately and appropriately specified, the inferential machinery of the model generally performs reliably (Elff et al., 2021):

1. Estimation of fixed-effect coefficients (regression parameters)
2. Estimation of variance–covariance components of the random effects (u) and residual errors (e)
3. The choice of an appropriate approximation of the sampling distribution of the test statistic.

Auch Bayesian, Raudenbush S. 399

2.3 Missing data in multilevel models

2.3.1 Methods for handling missing data

2.3.1.1 Listwise Deletion

2.3.1.2 Full Information Maximum Likelihood

2.3.1.3 Bayesian Model Estimation

Multiple Imputation

Elff, M., Heisig, J. P., Schaeffer, M., & Shikano, S. (2021). Multilevel Analysis with Few Clusters: Improving Likelihood-Based Methods to Provide Unbiased

- Estimates and Accurate Inference. *British Journal of Political Science*, 51(1), 412–426. <https://doi.org/10.1017/S0007123419000097>
- Fahrmeir, L., Kneib, T., Lang, S., & Marx, B. D. (2021). *Regression: Models, Methods and Applications*. Springer Berlin Heidelberg. <https://doi.org/10.1007/978-3-662-63882-8>
- Leonhart, R. (2013). *Lehrbuch Statistik: Einstieg und Vertiefung* (3., überarbeitete und erweiterte Auflage). Verlag Hans Huber.
- McNeish, D. (2017). Small Sample Methods for Multilevel Modeling: A Colloquial Elucidation of REML and the Kenward-Roger Correction. *Multivariate Behavioral Research*, 52(5), 661–670. <https://doi.org/10.1080/00273171.2017.1344538>
- Raudenbush, S., & Bryk, A. S. (2010). *Hierarchical linear models: Applications and data analysis methods* (2. ed., [Nachdr.]). Sage Publ.
- Snijders, T. A. B., & Bosker, R. J. (2012). *Multilevel analysis: An introduction to basic and advanced multilevel modeling* (2nd ed). Sage.