

UNIVERSITY OF HAMBURG

MASTER THESIS

Masterthesis_doc

Author:

Lona Frießner

Supervisor:

Prof. Dr. Simon Grund

*A thesis submitted in fulfillment of the requirements
for the degree of Master of Science*

in the

Institute of Psychology
Department of Psychology with focus on Quantitative Methods



Universität Hamburg
DER FORSCHUNG | DER LEHRE | DER BILDUNG

May 05, 2026

UNIVERSITY OF HAMBURG

Abstract

Institute of Psychology

Department of Psychology with focus on Quantitative Methods

Master of Science

Masterthesis__doc

by Lona Frießner

Hier kommt der Abstract hin

Table of contents

Abstract	iii
1 Introduction	1
2 Theory	3
2.1 Theoretical foundation of multilevel models	3
2.2 Estimation of multilevel models	8
2.3 Small sample sizes in multilevel models	8
2.4 Missing data in multilevel models	12
3 Methods	21
3.1 Data generation mechanism	21
3.2 Estimands of the focal analysis model	23
3.3 Methods of Missing Data Handling	23
3.4 Performance measures	24
4 Results	25
4.1 Bias of point estimation	25
4.2 Coverage of confidence/credible intervals	27
4.3 Standard deviation of point estimates	27
4.4 Width of confidence / credible intervals	28
4.5 Summary of results	29
5 Discussion	31
5.1 Interpretation of results	31
5.2 Limitations and further research	32
References	35
Appendices	41
A Derivations of equations of data-generating model	41
A.1 Calculation of σ_w^2	41
A.2 Calculation of the slope to generate W_j	42
A.3 Variances of random components of Y	43
B Convergence analysis for MI and fully Bayesian MCMC	45
B.1 Diagnostics of multiple imputation	45
B.2 Diagnostics of fully Bayesian analysis	45
C Complete simulation results for all performance measures	49

D	Default priors and sensitivity analysis	63
D.1	Default priors in <code>mdmb</code>	63
D.2	Default priors in <code>jomo</code>	63
D.3	Sensitivity analysis with weakly informative priors in <code>jomo</code>	63

List of Figures

2.1	Illustration of common multilevel model specifications.	4
2.2	Frequentist inferential process in a random intercept model	9
4.1	Bias of γ_{01} across simulation conditions under MAR.	26
4.2	Coverage performance for $N_2 = 15$	27
4.3	Empirical standard deviations for γ_{01} under MAR	28
4.4	Width of confidence/credible intervals for $\gamma_{01} = .4$	28
B.1	Trace and autocorrelation plot in an exemplary simulation run for $\beta_{Y,X}$ (multiple imputation).	46
B.2	Trace and autocorrelation plot in an exemplary simulation run for the level-2 variance of Y τ_{y0}^2 (multiple imputation).	46
B.3	Trace and autocorrelation plot in an exemplary simulation run for the level-2 variance of X τ_{x0}^2 (multiple imputation).	46
B.4	Trace and autocorrelation plot for the parameter $\beta_{wx}^{(imp)}$, representing the effect of the group mean of x on w in the imputation model (fully Bayesian estimation).	47
B.5	Trace and autocorrelation plot in an exemplary simulation run for the level-2 variance of X τ_{x0}^2 (fully Bayesian analysis)	47
B.6	Trace and autocorrelation plot in an exemplary simulation run for the level-2 variance of Y τ_{y0}^2 (fully Bayesian analysis)	48

List of Tables

3.1	Simulation conditions for the Data-Generating Model and the Generation of Missing Values	22
C.1	Simulation results ($N_2 = 15, \gamma_{01} = 0.0$)	50
C.2	Simulation results ($N_2 = 15, \gamma_{01} = 0.4$)	51
C.3	Simulation results ($N_2 = 30, \gamma_{01} = 0.0$)	52
C.4	Simulation results ($N_2 = 30, \gamma_{01} = 0.4$)	53
C.5	Simulation results ($N_2 = 60, \gamma_{01} = 0.0$)	54
C.6	Simulation results ($N_2 = 60, \gamma_{01} = 0.4$)	55
C.7	Monte Carlo standard errors of performance measures ($N_2 = 15, \gamma_{01} = 0.0$)	56
C.8	Monte Carlo standard errors of performance measures ($N_2 = 15, \gamma_{01} = 0.4$)	57
C.9	Monte Carlo standard errors of performance measures ($N_2 = 30, \gamma_{01} = 0.0$)	58
C.10	Monte Carlo standard errors of performance measures ($N_2 = 30, \gamma_{01} = 0.4$)	59
C.11	Monte Carlo standard errors of performance measures ($N_2 = 60, \gamma_{01} = 0.0$)	60
C.12	Monte Carlo standard errors of performance measures ($N_2 = 60, \gamma_{01} = 0.4$)	61
D.1	Bias reduction with alternative priors	64

Chapter 1

Introduction

Multilevel models are a standard statistical tool for analysing clustered data, which frequently occur in the social sciences (e.g., patients nested within therapists, students within classes, or employees within firms). In practice, such data sets are often characterized by a relatively small numbers of groups and by the presence of missing values.

Methodological research has proposed a range of approaches to address each of these challenges separately. Small-sample corrections for multilevel models have been developed to improve statistical inference when the number of clusters is limited (Elff et al., 2021; Holtmann et al., 2016; D. M. McNeish & Stapleton, 2016; D. McNeish & Stapleton, 2016). At the same time, modern missing-data methods have been extended to multilevel data structures (Audigier et al., 2018; Enders et al., 2016; see e.g. Enders, 2025; Grund et al., 2019; Kahan et al., 2016; Yucel, 2008).

Some research has investigated how these two challenges interact, both for single-level regressions (D. McNeish, 2017; see e.g. Von Hippel, 2013) and in multilevel contexts (e.g., D. McNeish, 2018; Shin et al., 2025; Shin & Shin, 2025; Zitzmann et al., 2015). However, the existing literature remains limited, and many relevant scenarios have not yet been systematically investigated. Among these, situations in which predictor variables contain missing values and are simultaneously used to construct group-mean contextual effects represent one particularly important but understudied case.

In applied research, effects at both the individual and the group level are often of interest and can be modeled within multilevel frameworks. Level-2 characteristics are frequently operationalized by aggregating Level-1 variables, resulting in so-called contextual effects (Croon & Van Veldhoven, 2007; Lüdtke et al., 2008; Snijders & Bosker, 2012, p. 102). This approach is widely used in psychological research (e.g., Anderman, 2002; Papaioannou et al., 2004). At the same time, missing values in predictor variables pose additional challenges for statistical inference, as they complicate missing data handling compared to situations in which only the outcome variable is incomplete. Standard likelihood-based approaches such as full information maximum likelihood are not easily applicable in this setting or require strong distributional assumptions.

This thesis therefore investigates the suitability of modern approaches to handling missing data in small-sample multilevel models in which predictor variables are partially missing and simultaneously included as group-mean contextual predictors. Specifically, the performance of multiple imputation and fully Bayesian estimation is evaluated under different data conditions.

The structure of this thesis is as follows: First, the theoretical foundations of multilevel models are introduced, and current methodological approaches for handling small samples and missing data are reviewed. Subsequently, a simulation study is conducted to investigate the performance of these methods under different data conditions.

Chapter 2

Theory

2.1 Theoretical foundation of multilevel models

Example

Imagine you are interested in the musical self-concept of young adults and whether it is related to how many hours they practice their instrument each week.

To investigate this question, you collect data from several youth ensembles. You recruit 15 ensembles, each consisting of approximately 10 players. Every player completes a questionnaire assessing their musical self-concept as well as their weekly practice time.

At first glance, this seems like a simple dataset of 150 individuals. But is it really that simple?

Players from the same ensemble rehearse together, share teachers, and experience similar social and musical environments. Their responses may therefore be more similar within ensembles than between ensembles.

Populations like this are common in the social sciences. Units of interest (e.g., students, employees, musicians) are grouped into higher-level units such as classes, firms, or ensembles. The resulting data is called *nested* or *clustered* (Snijders & Bosker, 2012, p. 7).

From a sampling perspective, a so-called *multistage sampling process* (sample groups first, then individuals inside these groups¹) is not independent: After selecting a group, you will try to sample all of the members of this group, making it more likely to select one person in this group over others in non-selected groups (Snijders & Bosker, 2012, p. 7). More importantly, individuals inside the same group typically share contextual influences (e.g., the same ensemble teacher, the same class room) and thus will give more similar responses in surveys and behavior. As a consequence, the basic assumption of independence in traditional linear modeling is violated, because error terms will be correlated (Raudenbush & Bryk, 2010, p. xx).

One well-suited and popular possibility to account for this fact is the use of *multilevel models* (also called *hierarchical linear models* or *mixed models*).

¹In this thesis, the term ‘groups’ refers to the higher-level units of the data and level-1 units are referred to as ‘persons’. However, it is important to note that these models can also be used for longitudinal research, where measurement points are on level 1 and individuals are the level-2 units. Furthermore, more than two levels are also possible, but are not discussed here.

2.1.1 The concept of multilevel regression models

The aim of regression analysis, whether single- or multilevel, is to explain variance in an outcome variable by including associated variables in the model as predictors (Snijders & Bosker, 2012, p. 80). In single-level regression, the residual e_i captures the *unexplained variance* in Y , that is differences in the outcome from the prediction that the model made (Leonhart, 2013, p. 319). The assumption of the classical linear model is that after inclusion of all *systematic* influences on Y , the residuals are *random*. Specifically, they are assumed to have expectation zero ($E(e_i) = 0$), constant variance across observations (homoscedasticity; $Var(e_i) = \sigma^2$), and to be independent across observations ($Cov(e_i, e_{i'}) = 0$ for $i \neq i'$) (Fahrmeir et al., 2021, p. 87).

Multilevel models (MLM) extend regular linear regression by adding level-2 *random effects* to explain the variability between groups that cannot be accounted for by predictors (Snijders & Bosker, 2012, p. 80). This partition of variance can be expressed in the multilevel regression model without predictors, the *empty model*:

$$Y_{ij} = \gamma_{00} + u_{0j} + e_{ij}$$

Here, the outcome value of person i in group j is modeled as the sum of the overall mean γ_{00} , a group level random effect u_{0j} and a random effect on the individual level, e_{ij} (Snijders & Bosker, 2012, p. 49). The random intercept u_{0j} captures the dependency of individuals within one group statistically, as all individuals in group j have the same u_{0j} value and their outcome values are therefore correlated. Conditional on this group component, however, the remaining level-1 residuals e_{ij} are assumed to be independent, addressing the issue of dependency that would arise if a single-level regression model were applied to nested data.

This and the models explained below are illustrated in Figure 2.1.




FIGURE 2.1: Illustration of common multilevel model specifications.

Note. Top row: empty model, random intercept model, and random slope model. Bottom row: group-level relationships between group means of the predictor $\bar{X}_{.j}$ and the group-specific intercepts β_{0j} .

2.1.2 Within- and between- group regressions and inclusion of predictors

Predictors can then be included to explain variability within groups (level 1) and between groups (level 2).

At level 1, the outcome of a person i in group j is modeled as a function of individual-level explanatory variables X_{ij} to obtain the within-group regression model. For simplicity, the model is illustrated with a single level-1 predictor:

$$Y_{ij} = \beta_{0j} + \beta_{1j}X_{ij} + e_{ij} \quad (2.1)$$

At level 2, the variation between groups in the group-dependent coefficients β_{0j} and β_{1j} is modeled. Each group may have its own intercept and slope. These coefficients can be separated into a fixed part (the average coefficient across groups) and a random deviation from this average (Snijders & Bosker, 2012, p. 75). Without level-2 predictors:

$$\begin{aligned} \beta_{0j} &= \underbrace{\gamma_{00}}_{\text{Fixed part}} + \underbrace{u_{0j}}_{\text{Random part}} \\ \beta_{1j} &= \underbrace{\gamma_{10}}_{\text{Fixed part}} + \underbrace{u_{1j}}_{\text{Random part}} \end{aligned}$$

On this level, in order to explain between-group variability in the intercepts and slopes, level-2 predictors Z_j may be included (Snijders & Bosker, 2012, p. 80). Including such predictors may reduce the unexplained variance of the random effects u_{0j} and u_{1j} . With a single level-2 predictor, the model becomes:

$$\beta_{0j} = \gamma_{00} + \gamma_{01}Z_j + u_{0j} \quad (2.2)$$

$$\beta_{1j} = \gamma_{10} + \gamma_{11}Z_j + u_{1j} \quad (2.3)$$

Substituting the level-2 equations Equation 2.2 and Equation 2.3 into the level-1 model Equation 2.1 yields the combined multilevel model:

$$Y_{ij} = \underbrace{\gamma_{00} + \gamma_{10}X_{ij} + \gamma_{01}Z_j + \gamma_{11}X_{ij}Z_j}_{\text{Fixed part}} + \underbrace{u_{0j} + u_{1j}X_{ij} + e_{ij}}_{\text{Random part}}$$

Y_{ij} : Outcome for individual i in group j

X_{ij} : Level-1 (within-group) predictor

Z_j : Level-2 (between-group) predictor

γ_{00} : Fixed intercept (overall mean when $X_{ij} = 0$, $Z_j = 0$)

γ_{10} : Average within-group effect of X_{ij} on Y_{ij}

γ_{01} : Effect of Z_j on the intercept (between-group effect)

γ_{11} : Cross-level interaction; change in the effect of X_{ij} depending on Z_j

u_{0j} : Random intercept (group-specific deviation from γ_{00})

u_{1j} : Random slope (group-specific deviation from γ_{10})

e_{ij} : Level-1 residual

If no level-2 predictors are included ($Z_j = 0$), the model reduces to the simpler random-coefficients model without contextual predictors.

2.1.3 Distributional assumptions of the multilevel regression model

Similar distributional assumptions are made for the level-2 random effects as for the residuals in single-level regression: they have mean zero, constant variances across groups, and are independent across clusters and independent of the level-1 residuals. However, random effects within the same cluster may be correlated and are often expected to do so (Snijders & Bosker, 2012, p. 76). Additionally, residuals and random effects are commonly assumed to follow a normal distribution for convenience, as it facilitates the likelihood-based estimation of the unknown parameters (Fahrmeir et al., 2021, p. 388). In summary:

$$\begin{aligned} \begin{pmatrix} u_{0j} \\ u_{1j} \end{pmatrix} &\sim \mathcal{N} \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \mathbf{T} \right), & \mathbf{T} &= \begin{pmatrix} \tau_0^2 & \tau_{01} \\ \tau_{01} & \tau_1^2 \end{pmatrix} \\ e_{ij} &\sim \mathcal{N}(0, \sigma^2) \\ \text{Cov}(u_{0j}, e_{ij}) &= 0, & \text{Cov}(u_{1j}, e_{ij}) &= 0 \end{aligned}$$

2.1.4 The random intercept model

In many empirical applications of multilevel models the slope β_{1j} is assumed to be constant across groups ($u_{1j} = 0$), leading to the *random intercept model* (Snijders & Bosker, 2012, p. 45). This model allows for baseline differences in the outcome between groups, whereas the strength of the relationship between predictors and the outcome is assumed to be constant:

$$Y_{ij} = \gamma_{00} + \gamma_{10}X_{ij} + \gamma_{01}Z_j + u_{0j} + e_{ij}$$

The random intercept model also provides a convenient framework for examining the use of *group means* of a given level-1 predictor as additional level-2 predictors. Including the group mean as a contextual variable allows the within-group regression coefficient to vary from the between-group regression coefficient (Snijders & Bosker, 2012, p. 57). This makes the model more flexible and often more realistic, as so-called *contextual effects* are frequently observed in empirical settings (Snijders & Bosker, 2012, p. 27). In such cases, the effect of an individual-level predictor differs from the effect of the average value of this predictor. The following example illustrates how processes operating within ensembles may differ from those operating between ensembles.

Example

Students often compare their own practice behavior with that of other ensemble members.

A student who practices more than their peers may feel more competent and report a higher musical self-concept.

At the same time, students in ensembles where everyone practices a lot may feel less competent because upward social comparisons become more likely.

This leads to two possible hypotheses:

- Students who practice more than others in their ensemble tend to report a higher musical self-concept.

- Students in ensembles with a higher average practice time may report a lower musical self-concept.

To separate within- and between-group effects, the level-1 predictor is then often *group-mean-centered*: Instead of using X_{ij} directly, the centered predictor $(X_{ij} - \bar{X}_j)$ is entered into the model. This decomposes within- and between- group effects and facilitates their interpretation (Snijders & Bosker, 2012, p. 58). A random intercept model including such a decomposition can be written like this:

$$Y_{ij} = \gamma_{00} + \gamma_{10}(X_{ij} - \bar{X}_j) + \gamma_{01}\bar{X}_j + u_{0j} + e_{ij} \quad (2.4)$$

In this random intercept model with a contextual effect, the musical self-concept of student i in ensemble j is predicted by the sum of several components:

- γ_{00} : the average musical self-concept across all ensembles
- $\gamma_{10}(X_{ij} - \bar{X}_j)$: the effect of the student's practice time relative to the average practice time in their ensemble
- $\gamma_{01}\bar{X}_j$: the effect of the ensemble's average practice time
- u_{0j} : a random deviation of ensemble j 's average musical self-concept from the overall mean (unexplained differences between ensembles)
- e_{ij} : a random deviation of the student's musical self-concept from the predicted value within their ensemble

The regression coefficients represent two conceptually different effects:

- γ_{10} : the *within-group effect*, indicating how a student's own practice time relates to their musical self-concept while controlling for the ensemble context
- γ_{01} : the *contextual effect*, indicating how the ensemble's average practice time relates to students' musical self-concept

For the remainder of this thesis, the focus lies on this particular random intercept model.

2.1.5 The intraclass correlation coefficient (ICC)

In the random intercept model, the variance components are given by

$$u_{0j} \sim \mathcal{N}(0, \tau_0^2), \quad e_{ij} \sim \mathcal{N}(0, \sigma^2), \quad \text{Cov}(e_{ij}, u_{0j}) = 0.$$

From these variance components, the proportion of variability that is attributable to between-group differences can be calculated. This quantity is referred to as the *intraclass correlation coefficient* (ICC). In the empty model, the ICC is equal to the correlation between the outcomes of two randomly chosen individuals from the same group (Snijders & Bosker, 2012, p. 18). In models including predictors, the ICC is interpreted conditionally on the predictors, that is, as the correlation between two individuals from the same group who share identical predictor values (also referred to as the *residual ICC*) (Snijders & Bosker, 2012, p. 52).

Formally, the ICC is defined as

$$ICC = \frac{\tau_0^2}{\tau_0^2 + \sigma^2}$$

The ICC is an important diagnostic metric because it indicates whether the use of a multilevel model is advantageous. If $ICC = 0$, no between-group variance is present and a conventional single-level regression model would typically be sufficient. Conversely, larger ICC values indicate increasing similarity within groups and thus statistical dependency of observations. Ignoring this dependency and applying a classical linear model leads to biased standard errors, confidence intervals, and hypothesis tests (Fahrmeir et al., 2021, p. 370).

2.2 Estimation of multilevel models

In the preceding chapter, the theoretical foundations and parameters of the multilevel model were introduced. In practice, however, these parameters are unknown and only exist in the theoretical realm as ‘true’ values. The fixed effects γ and variance components σ^2 and τ^2 must therefore be estimated from observed data. Based on these estimates, inferences about the underlying population parameters are drawn. This process is illustrated for the random intercept model in Figure 2.2. Provided that the model specification corresponds to the true data-generating process well enough, the reliable performance of this ‘inferential machinery’ depends on three main elements (Elff et al., 2021):


1. Estimation of fixed-effect coefficients (regression parameters)
2. Estimation of variance–covariance components of the random effects (u) and residual errors (e)
3. Selection of an appropriate approximation of the sampling distribution of the test statistic.

Aspects of the observed data may pose challenges to this estimation process. Moreover, the inferential machinery itself (i.e., its formulas and underlying assumptions) may be inapt for the empirical data reality. Two common challenging aspects, namely small sample sizes and missing data, will be discussed in the following chapters.

2.3 Small sample sizes in multilevel models

Frequentist estimation procedures for linear multilevel models typically rely on large-sample (asymptotic) theory. In multilevel models, the number of groups is particularly important. Simulation studies suggest that approximately 25–30 clusters are often required for reliable estimation, depending on the model specification and the parameters of interest (D. McNeish, 2017; D. M. McNeish & Stapleton, 2016). In practice, however, such numbers are often difficult to achieve because recruiting a large number of groups can be challenging and costly.

When the number of groups is small, several components of the inferential procedure may perform poorly, particularly the estimation of variance components, the calculation of standard errors, and the approximation of sampling distributions. The main issues associated with small-sample multilevel models and available corrections and alternatives are presented below.



Illustrative example. For a hypothetical data set with four persons nested in two groups:

The corresponding model matrices are

$$X = \begin{pmatrix} 1 & 2 \\ 1 & 1 \\ 1 & 3 \\ 1 & 5 \end{pmatrix} \quad Z = \begin{pmatrix} 1 & 0 \\ 1 & 0 \\ 0 & 1 \\ 0 & 1 \end{pmatrix} \quad Y = \begin{pmatrix} 5 \\ 4 \\ 2 \\ 5 \end{pmatrix}$$

Illustrative observed values

id	j	X_{ij}	Y_{ij}
1	A	2	5
2	A	1	4
3	B	3	2
4	B	5	5

and the implied covariance matrix is

$$\hat{V} = \begin{pmatrix} \hat{\tau}_0^2 + \hat{\sigma}^2 & \hat{\tau}_0^2 & 0 & 0 \\ \hat{\tau}_0^2 & \hat{\tau}_0^2 + \hat{\sigma}^2 & 0 & 0 \\ 0 & 0 & \hat{\tau}_0^2 + \hat{\sigma}^2 & \hat{\tau}_0^2 \\ 0 & 0 & \hat{\tau}_0^2 & \hat{\tau}_0^2 + \hat{\sigma}^2 \end{pmatrix}$$

Simplified frequentist inferential process in a random intercept model, based on Snijders & Bosker (2012) and Raudenbush & Bryk (2010).

Notes on notation

- X : design matrix for the fixed effects. Each row corresponds to an observation (e.g., a person) and each column to a predictor included in the fixed part of the model.
- Z : design matrix for the random effects. In the random intercept model it encodes group membership (1 = observation belongs to the group, 0 = otherwise), so observations from the same group share the same random intercept.
- Y : outcome vector containing the observed values of the dependent variable.
- $\hat{V}$: estimated variance–covariance matrix of the outcome vector Y . In the random intercept model it is determined by the estimated variance components of the random intercept ($\hat{\tau}_0^2$) and the residual variance ($\hat{\sigma}^2$), reflecting that observations within the same group are correlated.
- $\hat{\gamma}_k$: the k -th element of the estimated fixed-effect vector $\hat{\gamma}$.

FIGURE 2.2

2.3.0.1 Estimation of regression parameters

The regression parameters are estimates of the effects of interest. For example, how much does one additional hour of weekly practice influence a student’s musical self-concept? How does the musical self-concept differ between ensembles that practice on average three hours per week and those that practice thirteen hours per week?

Kackar & Harville (1981) showed that both ML and REML estimators of variance components are even and translation-invariant, implying that fixed effect point estimates are unbiased even in small samples.

2.3.0.2 Estimation of variance components

The estimation of the variance components is more delicate. In ML estimation, fixed effects and variance components are estimated simultaneously, typically via iterative procedures like the Expectation Maximization (EM) mechanism (Daniel McNeish, 2017). In this process fixed effects are effectively being treated as known when estimating variance components. Consequently, the sampling variability of the fixed-effect estimates is ignored and the loss of degrees of freedom due to their estimation is not accounted for. While these issues are not troublesome in large samples, they lead to systematic underestimation of variance components in small samples (Daniel McNeish, 2017).

REML addresses this issue by separating the estimation conceptually. The fixed effects are eliminated from the likelihood, and variance components are estimated solely based on residual information. Fixed effects are then estimated via generalized least squares (GLS), using the previously estimated variance components (Raudenbush & Bryk, 2010, p. 410). This leads to less downward-biased variance component estimates, while ML and REML become asymptotically equivalent as sample size increases.

Nevertheless, the the estimated variance of the fixed-effect estimators may still be too small in small samples for two reasons. First, the variance formula treats the estimated variance components $\hat{\theta}$ as known and therefore ignores their sampling variability (Kackar & Harville, 1984). Second, the REML variance expression itself relies on asymptotic approximations (Daniel McNeish, 2017). Both issues can lead to underestimated standard errors. Several small-sample corrections have been proposed to address this problem, among which the Kenward–Roger procedure is one of the most widely used (Daniel McNeish, 2017).

2.3.0.3 Approximation of the sampling distribution

The standardized test statistic used for inference (see Figure 2.2) can be regarded as ‘small-sample-adjusted’ at this point. The final step concerns the choice of a sampling distribution for p-values and confidence intervals. In small samples, a *t*-distribution is typically recommended instead of a standard normal distribution (Snijders & Bosker, 2012, p. 95). Compared to the standard normal distribution, the *t*-distribution has heavier tails, particularly when the degrees of freedom are small. As a result, it yields wider confidence intervals and larger p-values, making statistical inference more conservative in small samples.

The shape of the *t*-distribution is determined by the number of degrees of freedom. With small degrees of freedom, even small changes noticeably affect the shape of the sampling distribution and therefore the resulting p-values and confidence intervals. The concept of degrees of freedom is difficult to define concisely, but it can be described

as the number of independent pieces of information in the data that remain after accounting for restrictions such as estimated parameters (Pandey & Bright, 2008).

In multilevel models, the degrees of freedom generally cannot be calculated exactly and must therefore be approximated because of the nested data structure (Snijders & Bosker, 2012, p. 95). The Kenward–Roger correction mentioned earlier also includes a complex small-sample-adjusted approximation of the degrees of freedom, building on the Satterthwaite method (Daniel McNeish, 2017).

A more straightforward approximation is provided in textbooks by Raudenbush & Bryk (Raudenbush & Bryk, 2010, p. 58) and Snijders & Bosker (Snijders & Bosker, 2012, p. 95), and has also been proposed in methodological articles (Elff et al., 2021). This approach distinguishes between level-1 and level-2 predictors and reflects the concept of degrees of freedom as information minus estimated parameters:

$$df_{Lv1} = N_1 - r - 1, \quad df_{Lv2} = N_2 - q - 1$$

N_1 : Level-1 sample size (total number of individuals)

r : Total number of predictors

N_2 : Level-2 sample size (number of groups)

q : Number of predictors at level two

2.3.1 Fully Bayesian analysis

As alternative to the frequentist framework predominantly presented here, a Bayesian approach is often recommended for small sample sizes (Daniel McNeish, 2016a). In this framework, parameters are viewed as random variables with probability distributions. Prior knowledge about the parameters of interest in the form of prior distributions is combined with the information in the observed data in the form of a likelihood function. Bayes’ theorem yields the posterior distribution, representing the updated knowledge about the parameters (Van De Schoot et al., 2014). More general details on Bayesian analyses will not be discussed in this thesis.

Two aspects of Bayesian analyses are particularly helpful in small samples.

First, Bayesian inference does not rely on large-sample asymptotic approximations. Instead, the posterior distribution, the end result of a Bayesian analysis, is typically approximated using Markov Chain Monte Carlo (MCMC) sampling. Consequently, Bayesian inference does not require small-sample corrections for standard errors or degrees of freedom.

Second, Bayesian methods allow the incorporation of prior information about parameters. Informative or weakly informative priors may improve estimation accuracy by adding information when the likelihood is relatively flat in small samples (Gelman et al., 2025, p. 355). At the same time, prior specification becomes particularly influential under such conditions. Vague or software default priors may therefore have unintended effects and can perform poorly in small samples (Daniel McNeish, 2016a, 2016b; Gelman, 2006). Nevertheless, Bayesian estimation has in some settings been found to outperform maximum likelihood estimation in terms of RMSE, even when noninformative priors are used (Zitzmann et al., 2015). In addition, prior distributions may prevent inadmissible parameter estimates that could otherwise lead to convergence problems, such as negative variance estimates (Van De Schoot et al., 2014). Thus, the inclusion of prior distributions may be regarded either as a methodological advantage or as “a price to be paid” (Raudenbush & Bryk, 2010, p. 411) for Bayesian inference.

2.4 Missing data in multilevel models

Example

In the study about musical self-concept, you find after data collection that some students in several ensembles did not report how many hours they practice per week. They completed the questionnaire on musical self-concept but left the practice question blank.

The reason for this is unknown: they might have forgotten to answer the question, skipped it intentionally, or misunderstood it. Regardless of the reason, the data set now contains missing values. How should these missing observations be handled in the analysis?

Missing data (i.e., incomplete data sets) are a widespread issue in empirical research and of course also applies to multilevel data, e.g., because individuals drop out of studies or forget to fill out certain items of a questionnaire. Three major concerns commonly arise from data that contain missing values (Lüdtke et al., 2007):

1. The effective sample size is reduced, leading to lower efficiency in estimation (this is especially troublesome if the would-be sample size is already small)
2. Typical statistical methods are difficult to conduct, because they were designed for complete data.
3. Parameter estimation may be biased, if the missing and the observed data differ from each other systematically

Therefore, it is crucial to treat missing data appropriately. In the last three decades, this issue has received attention in psychological research, and modern methods were developed (Enders, 2022; Schafer & Graham, 2002), more recently also for the more complex case of multilevel data (Enders, 2025; Lüdtke et al., 2017). This chapter discusses these methods, after giving a short introduction into different missing data mechanisms and patterns.

2.4.1 Missing data mechanisms

Why data are missing can have multiple reasons, and different mechanisms may lead to different consequences in statistical inference. Rubin and Little (Little & Rubin, 1987; Rubin, 1976) introduced a typology and theoretical framework of missing data mechanisms that describes the relationship between the observed and the missing data. The missing data mechanism is generally not empirically testable from the observed data alone and assumptions about the mechanism must therefore be theoretically justified (Enders, 2022, p. 14ff.).

Missing completely at random (MCAR). Considering the hypothetical complete data matrix Y^2 , Y_{com} is partitioned into observed, Y_{obs} , and missing parts, Y_{mis} , and R denotes the missingness³. Missing values are considered to be MCAR if the distribution of missingness does not depend on the data (Schafer & Graham, 2002):

$$P(R | Y_{com}) = P(R)$$

²Here, Y represents the complete data set, including both outcome variables and predictors.

³ R is treated as a set of random variables that indicate for each value if it is missing or not (Schafer & Graham, 2002).

In other words, no variable (observed or unobserved) can predict missingness and every individual in every group has the same chance of missing values (Enders, 2022, p. 6).

Missing at random (MAR). The MAR mechanism describes the case when the probability of missingness does depend on observed data, but not on missing values (Enders, 2022, p. 8):

$$P(R | Y_{com}) = P(R | Y_{obs})$$

Missingness is therefore not random, as the name would suggest, but *conditionally random*: After controlling for the observed data, the missingness is purely random (Enders, 2022, p. 8).

For example, students who practice less might be more likely to skip the question about weekly practice duration. If this behavior can be predicted from other observed variables (musical self-concept or ensemble membership), the MAR assumption could be plausible.

Missing not at random (MNAR). If missingness depends on unobserved values, this is called a missing not at random process⁴:

$$P(R | Y_{com}) = P(R | Y_{obs}, Y_{mis})$$

Two individuals in the same group with identical scores in all observed variables therefore no longer have the same chance of missing values, because this chance depends on the unavailable, missing information itself (Enders, 2022, p. 11).

While these mechanisms also apply to multilevel data, the situation becomes more complex because missing values can occur in the outcome, in level-1 predictors, in level-2 predictors, or even in the grouping variable (or combinations of these; Buuren (2021)).

2.4.2 Methods for handling missing data

Many estimation procedures require complete data. Therefore, missing data must be handled explicitly before model estimation. If researchers do not actively specify a method, statistical software often defaults to using only complete observations (Lüdtke et al., 2007). This simple ad hoc procedure is called *listwise deletion* (LD) or *complete case analysis* and is a widely used approach to handling missing data. Cases with at least one missing value are discarded, and the model is estimated using only complete observations (Graham, 2009). In multilevel settings, missingness on the group level implies that all individuals in this group are excluded as well.

Listwise deletion has two major disadvantages (Enders, 2022, p. 24):

- Because partial data is unused, statistical power is reduced.
- If the missing data mechanism is not MCAR, parameter estimates may be highly biased.

LD can still be adequate under some conditions (Enders, 2022, p. 25; Graham, 2009), but in already small samples, additional data deletion should be avoided as power is already limited (D. McNeish, 2017).

⁴The special case $P(R | Y_{com}) = P(R | Y_{mis})$ is referred to as **focused MNAR** (Enders, 2022, p. 11).

More modern and better-performing methods are available and will be introduced in the following sections. These approaches generally rely on the assumption that data are missing at random (MAR). Methods for dealing with MNAR processes are not discussed in this thesis.

Modern approaches to missing data can be broadly specified into two classes:

1. *Model-based approaches* to missing data specify a statistical model for the complete data and base inference on the likelihood or posterior distribution implied by that model. Missing values are handled directly within the estimation procedure (Little & Rubin, 2020, p. 25).
2. *Imputation-based approaches* are two-fold: First, substitutions for the missing values are estimated, leading to complete data sets, which are then analyzed using standard methods.

2.4.2.1 Multiple Imputation (MI)

Multiple imputation is an imputation-based method, as the name suggests. The basic idea is to replace missing values with plausible draws ('imputations') from their predictive distribution. The substantive analysis model is then fitted to the imputed data. In order to include the uncertainty due to missing values, in MI several imputations for each missing observation are generated. By incorporating the variance between these imputations, the additional uncertainty can be captured (Enders, 2022, p. 283). The analysis of incomplete data is separated into three steps: imputation, analysis, and pooling.

2.4.2.1.1 Creating imputations

Imputations can be understood as "predicted values plus noise" (Enders, 2022, p. 266). Multiple imputation is closely related to Bayesian data augmentation techniques, in which missing values are treated as additional unknown quantities (Tanner & Wong, n.d.). From the joint distribution of regression parameters θ and the missing values $P(\theta, Y_{mis} | Y_{obs})$, parameters and imputations are simulated iteratively by an MCMC algorithm (Enders, 2022, p. 190). While during this process many thousand posterior draws are generated, only a small number (usually between $m = 20$ and $m = 100$ is recommended, Graham et al. (2007)) of imputations is saved, leading to m complete data sets. The posterior draws of the parameters are discarded (Gelman et al., 2025, p. 451).

The distribution of missing values (i.e., the imputation model) is either conceptualized jointly for all variables (*joint imputation*) or individually (*fully conditional specification*). Both approaches are also suitable for multilevel data structures (Enders et al., 2016). In joint modeling, a single multivariate distribution for all variables with missing values is specified, often the multivariate normal distribution in case of continuous variables (Enders, 2022, p. 263). *Fully Conditional Specification* approximates the joint distribution by cycling through a sequence of univariate models for each variable with missings, conditional on the others. This approach is suitable if many ordinal or nominal variables are included in the data, as for every variable the researcher specifies a separate imputation model (Lüdtke et al., 2007).

Compatibility of imputation and analysis model A crucial point in the choice of a valid imputation model is compatibility with the focal analysis model that is used later on. That is, the imputation model must reflect the key structural features of this analysis model (Grund et al., 2018). If it fails to do so, parameter estimates

may become biased because the imputed values no longer preserve the structure of the data. Consider, for example, the random intercept model with group means (Equation 2.4). In this setting, the imputation model must account for the nested data structure (i.e., random effects at the group level) and allow for different effects at the individual and the group level (Grund et al., 2018). **Auxiliary variables** Importantly, the imputation model is allowed to be ‘richer’ than the analysis model. Including additional effects or variables that are not part of the analysis model as *auxiliary variables*⁵ is considered beneficial for parameter estimation (Collins et al., 2001). These variables could either be a cause of missingness, thereby making the assumption of missing at random (MAR) more plausible. Furthermore, they may correlate with variables that contain missing values, which increases the precision of the imputations and reduces bias in parameter estimates (Collins et al., 2001).

Tip

In the musical self-concept study, the years of teaching experience of the ensemble teachers might also be available. Although this variable is not part of the analysis model, it may correlate with practice behavior or musical self-concept and can therefore serve as an auxiliary level-2 variable.

In this thesis only the joint imputation will be considered. The *multivariate empty model* represents a suitable joint imputation model for the random intercept analysis model (Grund et al., 2016). Revisiting the example from the previous chapter, an imputation model with one auxiliary variable can be written as

$$\begin{pmatrix} Y_{ij} \\ X_{ij} \\ W_j \end{pmatrix} = \begin{pmatrix} \beta_{0Y} \\ \beta_{0X} \\ \beta_{0W} \end{pmatrix} + \begin{pmatrix} u_{Yj} \\ u_{Xj} \\ u_{Wj} \end{pmatrix} + \begin{pmatrix} e_{Yij} \\ e_{Xij} \\ 0 \end{pmatrix} \quad (2.5)$$

Here, W_j represents a level-2 auxiliary variable and therefore has no level-1 residual term. The model preserves the clustered data structure by allowing random intercepts at the group level.

The random effects and residuals are assumed to follow multivariate normal distributions

$$\begin{pmatrix} u_{Yj} \\ u_{Xj} \\ u_{Wj} \end{pmatrix} \sim N(0, \Psi), \quad \begin{pmatrix} e_{Yij} \\ e_{Xij} \end{pmatrix} \sim N(0, \Sigma)$$

where the covariance matrix Ψ captures the between-group variability and covariation of the random intercepts and Σ represents the within-group covariance structure of the level-1 residuals:

$$\Psi = \begin{bmatrix} \psi_y^2 & \psi_{yx} & \psi_{yw} \\ \psi_{yx} & \psi_x^2 & \psi_{xw} \\ \psi_{yw} & \psi_{xw} & \psi_w^2 \end{bmatrix}, \quad \Sigma = \begin{bmatrix} \sigma_y^2 & \sigma_{yx} \\ \sigma_{yx} & \sigma_x^2 \end{bmatrix}$$

⁵Auxiliary variables are referred to as variables whose only purpose it is to improve the missing data imputation (Collins et al., 2001).

2.4.2.1.2 Analyzing the completed data sets

After the imputation step, researchers are in the comfortable position to have m completed data sets that can be analyzed using standard frequentist methods. To each of the data sets, the substantive analysis model of interest (e.g., the random intercept model) is now fitted. Each analysis proceeds according to the standard estimation procedure described in Section 2.2, up to the computation of parameter estimates and their standard errors. Hypothesis testing is not conducted at this stage. The result of this step is a set of m parameter estimates and corresponding standard errors.

2.4.2.1.3 Pooling the results

These m estimates and standard errors are aggregated into a single set of results following Rubin's pooling rules (Rubin, 1987). He defined the point estimates of effects as the arithmetic average of the m estimates:

$$\hat{\theta} = \frac{1}{m} \sum_{k=1}^m \hat{\theta}_k$$

$\hat{\theta}$ is the pooled point estimate, and $\hat{\theta}_k$ is the estimate from data set k .

For the pooled standard errors, both the within-imputation variance (that is based on a complete data set) and the between-imputation variance (that adds the uncertainty due to missing values) are combined. The within-imputation variance is the arithmetic average of the imputation's variances (standard errors squared):

$$\bar{V}_w = \frac{1}{m} \sum_{k=1}^m SE_k^2$$

The between-imputation variance is defined as the variance of point estimates between imputations that is due to uncertainty caused by the missing values:

$$V_b = \frac{1}{m-1} \sum_{k=1}^m (\hat{\theta}_k - \hat{\theta})^2$$

The total pooled standard error combines the within-imputation variance $\bar{V}_w$ and the between-imputation variance V_b :

$$SE = \sqrt{\bar{V}_w + V_b + \frac{V_b}{m}}$$

The third term in this expression acts as a correction factor for using a finite number of imputations in the Monte Carlo simulation (Enders, 2022, p. 284). As the number of imputations m increases, this correction term approaches zero. As the between-imputation variance is a measure of increase in uncertainty due to missing values, the influence of missing data on the variance of estimates can be expressed proportionally, e.g. as the relative increase of variance (RIV) due to missing values:

$$RIV = \frac{V_b + \frac{V_b}{m}}{\bar{V}_w}$$

With an appropriate point estimate and standard error, the remaining step (see Figure 2.2) is the choice of the sampling distribution. Rubin proposed a calculation method for degrees of freedom of the t-distribution, that is based on the *RIV* and the number of imputations m :

$$df_R = (m - 1) \left(1 + \frac{1}{RIV^2} \right)$$

This formula assumes that the degrees of freedom of the complete data are effectively infinite (Barnard & Rubin, 1999). In smaller samples, the classic Rubin formula therefore tends to produce degrees of freedom that are too large, sometimes even larger than the degrees of freedom of the complete data. Barnard and Rubin [Barnard] thus proposed a small-sample-adjusted degrees-of-freedom calculation for pooled parameters. Their formula downweights the complete-data degrees of freedom to account for the missing values. As the complete-data degrees of freedom cannot easily be obtained for multilevel analyses, a suitable degrees-of-freedom approximation must be chosen for the analysis. Possible approaches include the methods described in Section 2.3.0.3, such as the Kenward–Roger correction or the simple $m - l - 1$ rule.

2.4.2.2 Full Information Maximum Likelihood (FIML)

Full information maximum likelihood (FIML) is a model-based approach in which parameter estimates are obtained directly by maximizing the likelihood of the observed data (Enders, 2022, p. 98). Under the assumption that data are missing at random (MAR), FIML yields unbiased and efficient parameter estimates (D. McNeish, 2017).

Incorporating auxiliary variables is less straightforward in FIML than in multiple imputation. Nevertheless, several approaches have been proposed to include auxiliary information in FIML estimation, such as the saturated correlates model or the extra dependent variable approach (Graham, 2003), as well as the factored regression framework described below for the Bayesian missing-data approaches.

For the random intercept model with group-mean effects considered in this thesis, FIML is not always well suited when missingness occurs in predictor variables. In such cases, maximum likelihood estimation requires additional distributional assumptions about the distribution of the predictors in order to incorporate them into the likelihood function (Grund et al., 2018). In multilevel models with manifest group means, these assumptions may lead to biased estimates of between-group effects (Grund et al., 2019).

2.4.2.3 Fully Bayesian Estimation

The Bayesian framework introduced in Section 2.3.1 can naturally be applied to missing-data problems. In contrast to the two-step procedure of multiple imputation, Bayesian missing-data analysis directly uses the posterior distribution of the model parameters conditional on the observed data for statistical inference.

As described in Section 2.4.2.1.1, missing values are treated as additional unknown quantities. To simulate these values during estimation, a model for the variables with missing data must be specified. A convenient way to specify the joint distribution of the outcome and the variables with missing values is to apply the probability chain rule and express the joint distribution as a product of conditional distributions, which is referred to as *sequential specification* or *factored regression specification* (Lüdtke et al., 2020):

$$g(y, x; \gamma) = g_y(y|x; \gamma_y) \times g_x(x; \gamma_x)$$

where $\gamma = (\gamma_y, \gamma_x)$ denotes the vector of all model parameters. Here, the joint density is decomposed into two components. The first term $g_y(y | x; \gamma_y)$ corresponds to the analysis model, describing the conditional distribution of the outcome given the covariate X . The second term $g_x(x; \gamma_x)$ specifies a model for the distribution of the covariate X .

Each conditional distribution in a sequential modeling framework is typically specified as a regression model assuming normally distributed residuals. Consequently, the implied joint distribution is constructed from a sequence of conditional models, each imposing its own distributional assumptions.

An advantage of this formulation is its flexibility: it allows the inclusion of variables with different measurement scales and facilitates the use of auxiliary variables that are not part of the substantive analysis model (Daniels et al., 2014; Erler et al., 2016). For example, auxiliary variables such as W may be included to improve the imputation of missing values while the substantive model of interest is still estimated without W . For this purpose, auxiliary variables can be placed in the first position of the factored regression, so that the analysis variables predict the auxiliary variables, but not vice versa (Enders, 2022, p. 215):

$$g(y, x, w; \gamma) = g_w(w|y, x; \gamma_w) \times g_y(y|x; \gamma_y) \times g_x(x; \gamma_x) \quad (2.6)$$

Because all missing values are resampled in each iteration, the uncertainty due to missing data is automatically included in the posterior distribution (Erler et al., 2016).

2.4.3 Present study

Both challenges considered here — small sample sizes and missing data — often co-occur in empirical research, but missing-data methods have rarely been investigated in small-sample settings. For multiple imputation, some studies suggest that small-sample adjustments applied during pooling may improve inferential performance in both single-level and multilevel contexts (D. McNeish, 2017, 2018). However, these adjustments have not been compared to standard inferential approaches based on unadjusted degrees of freedom, and the models considered did not incorporate contextual predictors.

Fully Bayesian estimation may offer a promising model-based alternative, as it naturally accommodates both missing predictor values and small sample sizes. It is often recommended in more complex multilevel models, for example in the presence of nonlinearities or interaction effects (Depaoli & Clifton, 2015; Enders et al., 2020; Muthén & Asparouhov, 2012). However, it may also represent a suitable alternative to likelihood-based approaches such as FIML in simpler models with contextual effects, where the handling of missing predictor variables and the construction of group means may lead to biased estimates (Grund et al., 2019).

To evaluate this, the present study uses a simulation design to compare methods for handling missing data in a predictor variable in a small-sample multilevel random intercept model. In particular, it examines whether applying small-sample-adjusted degrees-of-freedom approximations after multiple imputation improves inferential performance, and whether fully Bayesian estimation provides a suitable alternative to imputation-based or likelihood-based approaches.

Simulation studies can be understood as computer experiments (Morris et al., 2019), in which data with known characteristics are generated and then analysed using the methods of interest. Because the true parameter values are known, the resulting estimates can be compared with the truth, allowing the performance of the methods to be evaluated (Siepe et al., 2024). The simulation design and evaluation criteria are described in the following section.

Chapter 3

Methods

Planning of the simulation study followed the ADEMP structure proposed by Siepe et al. (2024) and will be reported based on their guidelines.

3.1 Data generation mechanism

3.1.1 Data-generating model

The data-generating model was a two-level random intercept model:

$$Y_{ij} = \gamma_{10}(X_{ij} - \bar{X}_{\cdot j}) + \gamma_{01}\bar{X}_{\cdot j} + \gamma_{02}W_j + u_{0j} + e_{ij} \quad (3.1)$$

Y_{ij} , X_{ij} and W_j were parameterized to have a mean of zero and a variance of 1 on the population level.

First, between- and within-components of X_{ij} were simulated, independent of each other and with variances chosen to achieve a prespecified intraclass coefficient (see Section 3.1.3):

$$\begin{aligned} X_{ij} &= u_{Xj} + e_{Xij}, \\ u_{Xj} &\sim N(0, \tau_x^2), \quad e_{Xij} \sim N(0, \sigma_x^2) \end{aligned}$$

with $\tau_x^2 = ICC_x$ and $\sigma_x^2 = 1 - ICC_x$, respectively.

X_{ij} was then group-mean-centered.

The group-level covariate W_j was generated to have a prespecified correlation with the group mean of X ($\rho_{\bar{X}W}$) using a linear model¹:

$$W_j = a\bar{X}_{\cdot j} + e_{Wj} \quad e_{Wj} \sim N(0, \sigma_w^2)$$

Finally, the random effects on the group and the individual level were generated as

$$u_{0j} \sim N(0, \psi^2), \quad e_{ij} \sim N(0, \sigma^2)$$

and independent of each other.

Individual and group residual variances of Y_{ij} were set to achieve a prespecified residual intraclass correlation and $Var(Y_{ij}) = 1$, see Appendix A.

¹Under the constraint $Var(W_j) = 1$, it follows that $\sigma_w^2 = 1 - \rho_{\bar{X}W}^2$ and $a = \frac{\rho_{\bar{X}W}}{\sqrt{\tau_x^2 + \frac{\sigma_x^2}{n}}}$, see Appendix A for details.

TABLE 3.1: Simulation conditions for the Data-Generating Model and the Generation of Missing Values

Data conditions	
N_1	10
N_2	15 / 30 / 60
ICC of X and Y	.1 / .3
Correlation $\bar{X}W$	.3
Model parameters	
Effect of $(X_{ij} - \bar{X}_{.j})$	.2
Effect of $\bar{X}_{.j}$	0 / .4
Effect of W_j	0
Missing values	
Pattern	$X \sim W$
Mechanism	MCAR / MAR
Proportion	.3

3.1.2 Missing data generation

After simulation of the complete dataset, values of the X -variable have been deleted to simulate missingness. A ‘missingness tendency’ r was simulated for each X -value according to

$$r_{ij} = \alpha + \beta W_j + e_{ij}, \quad e_{ij} \sim N(0, 1 - \beta^2)$$

r_{ij} is normally distributed. Observations with $r_{ij} > 0$ were set to missing.

The intercept α controls the proportion of missingness in X , whereas the slope β determines the strength of the missing-at-random mechanism. When $\beta = 0$, missingness is completely at random (MCAR). With increasing β , the missingness mechanism becomes more dependent on the W -values; with $\beta = 1$, missingness would be completely deterministic.

For the present simulation, $\beta = 0.3$ was used for MAR conditions, corresponding to approximately 9% explained variance.

3.1.3 Factors of the data-generating mechanism

A summary of simulation conditions is provided in Table 3.1.

The number of Level-2 units was manipulated across conditions ($N_2 = 15, 30, 60$), while the number of Level-1 observations was kept constant ($N_1 = 10$). The between-cluster effect of $\bar{X}_{.j}$ was varied ($\gamma_{01} = 0, .40$), whereas the within-cluster effect of $(X_{ij} - \bar{X}_{.j})$ was held constant at .20 and the effect of W_j was set to zero. This results in conditions with either a negative or a positive context effect (i.e., $\gamma_{10} - \gamma_{01}$) of X in the population model. The ICC of X and the residual ICC of Y were varied ($\rho_X = \rho_{Y|X} = .10, .30$). Missing data were generated on X with a varying missingness mechanism (MCAR or MAR with a strength of $\beta = .30$), with a constant missingness rate of 30%. The factors were combined fully factorially, creating $3 \times 2 \times 2 \times 2 = 24$ conditions.

Taken together, the simulation design targets conditions that are particularly challenging for multilevel estimation (small Level-2 sample sizes and low ICCs), while also covering less challenging scenarios for comparison. The between-group effect of $\bar{X}_{.j}$ is varied to evaluate method performance under a null versus a substantial effect.

3.2 Estimands of the focal analysis model

The focal analysis model is

$$Y_{ij} = \gamma_{10}(X_{ij} - \bar{X}_{\cdot j}) + \gamma_{01}\bar{X}_{\cdot j} + u_{0j} + e_{ij} \quad (3.2)$$

The primary target is the Level-2-effect, γ_{01} . Additionally, the Level-1-effect γ_{10} was evaluated.

3.3 Methods of Missing Data Handling

Four methods will be compared, alongside the benchmark analysis of Complete Data (CD).

3.3.1 Listwise Deletion (LD)

Listwise Deletion was used as a standard reference method for missing data handling. Cases with missing X -values were excluded prior to analysis. The group means $\bar{X}_{\cdot j}$ were then calculated only from the remaining observations. Then, group-mean centering was applied.

3.3.2 Multiple Imputation (MI) with standard (MI-R) and adjusted (MI-a) degrees of freedom

For both multiple imputation methods, a joint modeling approach was employed. The imputation model was the multivariate empty model specified before (Equation 2.5), with W_j as an auxiliary variable on level 2.

Imputation was executed with the R package `mitml` (Grund et al., 2015) which provides an interface to the `jomo` package (Quartagno & Carpenter, 2023). The algorithm was run with 2000 burn-in-iterations and 3000 post-burn-in iterations, saving every 300th imputation, resulting in 10 imputations.

For MI-R, degrees of freedom for calculating the test statistic were calculated with the software standard (Rubin's rules, assuming infinite degrees of freedom in complete data).

For MI-a, complete data degrees of freedom were calculated with $df_{\gamma_{10}} = M - r - 1$ (M : total number of Level-1 units; r : total number of predictors, in the present case 2) and $df_{\gamma_{01}} = N_2 - q - 1$ (N_2 : number of Level-2 units; q : number of Level-2 predictors, here: 1).

3.3.3 Fully Bayesian Analysis

A factored regression specification with the auxiliary variable W was used within the Bayesian framework that corresponds to the model described in Equation 2.6.

The burn-in period was set to 1000 iterations, followed by 3000 post-burn-in iterations. The model was estimated using the package `mdmb` (Robitzsch & Luedtke, 2024) with its default noninformative priors.

3.3.4 Extracted quantities

For complete data, listwise deletion and both multiple imputation methods, point estimates of γ_{10} and γ_{01} as well as their standard errors and 95% - confidence intervals

were extracted.

For the fully Bayesian method, the posterior means of γ_{10} and γ_{01} were used as point estimates along with posterior standard deviations and 95% credible intervals. For comparability with the frequentist approaches and in line with common simulation practice, these posterior summaries were treated analogously to frequentist estimates and intervals (Morris et al., 2019).

3.4 Performance measures

Performance was evaluated with standard performance measures for simulation studies (Morris et al., 2019; Siepe et al., 2024); namely bias, coverage, standard deviation of the estimates across simulation replications and the average width of confidence/credible intervals.

Monte Carlo uncertainty of the estimated performance measures was quantified using Monte Carlo Standard Errors (MCSE), using the formulas provided in Siepe et al. (2024).

For each condition, 1000 simulation replications were conducted, in line with comparable simulation studies (Enders et al., 2016; Grund et al., 2018) and resulting in sufficiently small MCSEs for coverage. All simulations were conducted in R (version 4.5.2, R Core Team (2025)) using a reproducible workflow (e.g., fixed random seeds and controlled package environments). The full code and all relevant materials to reproduce the results are available in the digital version of this thesis (*LINK*).

Chapter 4

Results

Convergence diagnostics for the MCMC algorithm were evaluated under an exemplary critical condition and indicated satisfactory convergence with no problematic autocorrelation (see Appendix B). Monte Carlo uncertainty was quantified using intervals of ± 2 Monte Carlo standard errors (MCSEs), corresponding approximately to 95% confidence intervals for the estimated performance measures.

For the individual-level effect γ_{10} , bias was consistently negligible relative to their Monte Carlo standard errors, and coverage, empirical standard deviations and confidence/credible interval widths were comparable across methods and closely aligned with the corresponding complete-data estimates. Results are therefore reported here only for the group-level effect γ_{01} . Full results for all parameters of interest and performance measures are provided in Appendix C.

4.1 Bias of point estimation

The findings for γ_{01} are depicted in Figure 4.1 for all conditions with missingness at random (MAR). Results under MCAR followed very similar patterns and are therefore not shown. A bias of 10% of the true parameter value is considered substantial. For a true null effect, bias is considered substantial if it lies outside the interval of ± 2 Monte Carlo Standard Errors (MCSE).

Listwise deletion resulted in substantial bias, especially in conditions with low ICC. For a true null effect, estimates were positively biased, with a maximum of 0.042 (MCAR, $ICC = .1$, $N_2 = 30$). For a positive between-group effect, estimates were negatively biased, with a maximum of -0.05 . The magnitude of bias was similar across sample sizes and did not differ markedly between MAR and MCAR conditions. Multiple imputation resulted in larger bias than listwise deletion in conditions with low ICC and a positive between-group effect. Bias ranged from -0.044 to -0.078 , corresponding to approximately 10-20% of the true effect¹. In all other conditions, bias estimates for multiple imputation were small relative to their Monte Carlo standard errors (all $|\text{bias}| \leq 0.02$). In contrast, the Bayesian approach showed only small bias across all simulation conditions except the most challenging conditions ($N_2 = 15$, $ICC = .1$, $\gamma_{01} = .4$), where bias was -0.039 (MAR) and -0.036 (MCAR). These values remain below 10% of the true effect. All other biases were small ($|\text{bias}| \leq 0.015$).

¹However, in a sensitivity analysis, the use of alternative weakly informative priors in these conditions consistently reduced bias by approximately 30–60% (see Appendix D for details).




FIGURE 4.1: Bias of the between-group effect (γ_{01}) across simulation conditions under missing at random (MAR). Error bars represent ± 2 Monte Carlo standard errors. ES = Effect size.

4.2 Coverage of confidence/credible intervals

Coverage of the confidence and credible intervals for γ_{01} is shown for $N_2 = 15$ in Figure 4.2, as coverage for larger sample sizes showed similar but attenuated patterns. Overall, coverage rates were close to nominal (95%) across methods for $N_2 = 30$ and $N_2 = 60$.

Listwise deletion yielded coverage rates close to the nominal level across all conditions.

For multiple imputation, the small-sample adjusted degrees of freedom resulted in increased overcoverage compared to the standard degrees of freedom, particularly when $N_2 = 15$, with a similar but weaker tendency at larger sample sizes.

The Bayesian approach showed overcoverage for $N_2 = 15$, which was generally slightly higher than for multiple imputation with adjusted degrees of freedom. With increasing Level-2 sample size, this pattern attenuated. For $N_2 = 30$, coverage estimates were largely close to the Monte Carlo error bounds, and for $N_2 = 60$, they were fully within these bounds across conditions.




FIGURE 4.2: Deviation from nominal (95%) coverage of confidence/credible intervals across methods and simulation conditions for $N_2 = 15$. Error bars represent ± 2 Monte Carlo standard errors. MI = Multiple Imputation; ES = Effect size.

4.3 Standard deviation of point estimates

Standard deviations of point estimates across conditions and methods are illustrated in Figure 4.3 for missingness at random (MAR), as differences between missingness mechanisms were small.

Listwise deletion produced standard deviations very close to those obtained under complete-data analysis, ranging from 0.124 ($N_2 = 60$, MCAR, $\gamma_{01} = 0.4$) to 0.300 ($N_2 = 15$, MCAR, $\gamma_{01} = 0.0$). For multiple imputation, standard deviations were comparable to complete-data analysis across conditions, with a tendency toward smaller values in some low-ICC conditions. For example, for $N_2 = 15$, $\gamma_{01} = 0.0$, and MAR missingness, multiple imputation yielded a standard deviation of 0.247, compared to 0.283 under complete-data analysis. The Bayesian approach yielded standard deviations that were generally slightly larger than those from complete-data analysis, typically by approximately 0.006–0.02. Standard deviations were generally

smaller under high ICC conditions than under low ICC conditions in the Bayesian framework, except in the conditions with $\gamma_{01} = 0.4$ and $N_2 = 15$. In contrast, multiple imputation consistently yielded slightly larger standard deviations under high ICC conditions across all examined conditions.




FIGURE 4.3: Empirical standard deviation of point estimates for γ_{01} for conditions with missingness at random (MAR). Error bars represent ± 2 Monte Carlo standard errors.

4.4 Width of confidence / credible intervals

Results are shown in an exemplary manner in Figure 4.4, as this pattern was visible across conditions in a similar way.

Bayesian credible intervals were generally the widest, followed by the multiple imputation with adjusted degrees of freedom approach. Confidence intervals from listwise deletion and from the standard multiple imputation with Rubin's degrees of freedom were similar to complete data analysis.

With increasing Level-2 sample size, differences between methods became less pronounced, although interval widths for the Bayesian approach remained slightly larger. A slight tendency for larger intervals in low-ICC conditions was observed for the Bayesian approach, whereas this pattern was minimal for multiple imputation.




FIGURE 4.4: Width of confidence/credible intervals for $\gamma_{01} = .4$ under missingness at random (MAR) and $ICC = .1$. Error bars represent ± 2 Monte Carlo standard errors.

4.5 Summary of results

Listwise deletion showed increased bias while all other performance measures were closely aligned to the complete data analysis. Multiple imputation produced substantial downward bias in the estimation of the level-2 effect under conditions of low ICC and MAR missingness. The standard variant, but close to nominal coverage for all conditions, whereas the fully Bayesian approach resulted in overall wider confidence intervals and a tendency toward overcoverage, but small to no bias. Multiple imputation with adjusted degrees of freedom showed increased overcoverage and wider confidence intervals compared to the standard variant.

In the present simulation, the performance of confidence intervals based on uncorrected degrees of freedom was unexpectedly good. Coverage estimates were largely consistent with nominal levels within Monte Carlo uncertainty, suggesting that uncorrected degrees of freedom did not lead to substantial undercoverage in the present setting. Applying a small-sample correction to the degrees of freedom, which results in heavier tails of the sampling t -distribution, led to higher coverage, exceeding the nominal level.

Chapter 5

Discussion

This thesis was concerned with the treatment of missing data in predictor variables in small-sample multilevel models with group-mean predictors. It focused on a reduced setting with a single continuous predictor which was included at level 1 as an individual-level predictor and at level 2 as a manifest group-mean predictor, within a random intercept model. In addition, one auxiliary variable at level 2 was included to explain missingness under a MAR mechanism, without being directly related to the outcome. While applied settings are often more complex, this simplified design allowed for a more controlled investigation of the methods of interest. Furthermore, there is still limited literature addressing the combined challenges of multilevel data, small sample sizes, and missing data. Against this background, focusing on a simplified setting appears to be a reasonable starting point for examining the behavior of the methods before extending to more complex models. The analytical setup combined approaches that explicitly account for small-sample uncertainty with commonly used default settings (e.g., prior specifications), thereby approximating conditions that may also occur in applied research.

5.1 Interpretation of results

The results for listwise deletion are consistent with theoretical expectations regarding systematic bias. However, contrary to expectations, listwise deletion did not perform worse under MAR than under MCAR conditions. The MAR mechanism in the present study was relatively weak, suggesting that it was not sufficiently strong to induce additional bias beyond that already observed under MCAR.

A comparison of multiple imputation and the fully Bayesian approach suggests a trade-off between bias and interval width. In settings with low ICC, where unbiased estimation of level-2 effects is critical and default prior specifications are used, the fully Bayesian approach may be preferable to multiple imputation. Given the narrow scope of the simulation design, this should be interpreted as a cautious indication rather than a general recommendation. It is important to note that differences between Bayesian estimation and multiple imputation are often small in magnitude when based on comparable assumptions, and multiple imputation can be understood as an approximate Bayesian procedure, as it relies on draws from the posterior predictive distribution (e.g., Enders (2022), p. 299; Schafer & Graham (2002)). Against this background, the present results should not be interpreted as representing a general pattern between the two approaches. Prior specification seems to play an important role in this context, as the sensitivity analysis revealed. When reasonable prior information is available for level-2 variance components, these may reduce bias produced with multiple imputation while holding coverage at a nominal level.

One aspect of multiple imputation that was specifically investigated in this thesis was the behavior of inference based on uncorrected versus small-sample-adjusted degrees of freedom. The findings suggest that, in this setting, standard Rubin’s rules already yield approximately valid inference, and that additional small-sample corrections may not be necessary. This is somewhat surprising, as it is commonly recommended to use an appropriate sampling distribution in small samples to avoid overly narrow confidence intervals; a recommendation which also applies to multilevel analyses (Kenward & Roger, 1997; Schaalje et al., 2002) and within multiple imputation (Barnard & Rubin, 1999). However, confidence intervals are influenced by multiple sources of uncertainty beyond the degrees-of-freedom approximation alone. In particular, REML estimation was used for analyzing the imputed data sets. Compared to ML, REML typically yields larger estimates of level-2 variance components, which have been shown to improve variance estimation substantially in small samples (Elff et al., 2021; D. M. McNeish & Stapleton, 2016). As these variance components directly affect standard errors, this may have contributed to wider intervals and thus higher coverage. In addition, multiple imputation was performed using joint modeling via `jomo`. Previous work has shown that inverse Wishart priors with few degrees of freedom, as implemented in `jomo`, can introduce additional variability into fixed-effect estimates (Audigier et al., 2018). Overall, the present standard MI setup performed well in the random intercept model with group-mean predictors, which may be reassuring for applied researchers relying on default degrees-of-freedom specifications.

In terms of implementation, both fully Bayesian estimation and multiple imputation allow the inclusion of auxiliary variables, which were also part of the present setup. In multiple imputation, their inclusion is typically well-documented and supported by standard workflows. In fully Bayesian approaches, however, incorporating auxiliary variables without altering the analysis model requires additional modeling decisions that are less explicitly supported by standardized workflows and software guidance, and may therefore not be immediately apparent without prior methodological experience.

Furthermore, the choice between approaches is likely to depend not only on statistical performance but also on the research context and analytical goals. Multiple imputation may be advantageous in settings where standardized workflows and comparability across analyses are desired, as completed data sets can be shared and reused across different analyses. However, reproducibility can in principle be achieved with both approaches when data, code, and model specifications are transparently reported.

In contrast, fully Bayesian approaches may be particularly useful in cumulative research settings, where prior information can be updated across studies, allowing evidence to be built progressively.

5.2 Limitations and further research

`mdmb`’s default priors allow for even more variability in the parameter estimates than in `jomo`. Thus, it would be interesting to look at different prior specifications in both approaches to investigate this further.

Future research is needed to disentangle the relative contributions of these components. In particular, simulation studies that systematically vary estimation method (ML vs. REML), degrees-of-freedom corrections, prior specifications, and imputation frameworks (e.g., joint modeling vs. fully conditional specification) would help to

clarify their individual and combined effects on inferential performance of multiple imputation.

This study comes with many limitations. Firstly, only continuous variables were considered.

Furthermore, in some practical examples the number of clusters might be even smaller than 15.

The missing data mechanisms lacked the inclusion of missing not at random processes, which also are to be expected in practical research scenarios.

The simulation model was fairly simple, with no random slopes or cross-level interactions. This seemed a good start to investigate the performance of the methods of interest in a simple setting at first, but should be expanded from here.

Importantly, prior choice was not the focus of the present thesis, but is of great importance in small samples. As the sensitivity analysis revealed, it influences the results noticeably. The systematic manipulation of different priors to further compare multiple imputation and Bayesian techniques would be useful.

To note is also, that the present study only looked at artificial, simulated data that satisfy the supposed model assumptions (e.g., normality, homoscedasticity of residuals). It would be illuminating to look at the methods' behavior in a comparable real data set and/or look at the consequences of misspecifications of the imputation or analysis models.

nur geringes MAR, nur kontinuierliche Variablen, keine Random Slopes, einfaches Modell, keine geringsten Gruppengrößen, immer gleich große Gruppen (macht das was aus?), kein MNAR, keine priors angeguckt oder manipuliert, simulierte Daten die die Normalitätsannahmen erfüllen; manifeste vs. latente group means (lückte)

It is important to note that firstly, distinct software packages were used to emply the two approaches

Diverging behavior zwischen Bayes und MI würde vuielleicht auch Sinn machen in kleinen Stichproben, aber das könnte man nur untersuchen, wenn man alles andere konstant hält. Oder doch nicht? Wirken die Priors durch die imputierten Daten dann bei MI doch auf die Parameterschätzungen?

References

- Anderman, E. M. (2002). School effects on psychological outcomes during adolescence. *Journal of Educational Psychology*, 94(4), 795–809. <https://doi.org/10.1037/0022-0663.94.4.795>
- Asparouhov, T., & Muthen, B. (2010). *Bayesian Analysis Using Mplus: Technical Implementation*.
- Audigier, V., White, I. R., Jolani, S., Debray, T. P. A., Quartagno, M., Carpenter, J., Van Buuren, S., & Resche-Rigon, M. (2018). Multiple Imputation for Multilevel Data with Continuous and Binary Variables. *Statistical Science*, 33(2). <https://doi.org/10.1214/18-STS646>
- Barnard, J., & Rubin, D. (1999). Small-sample degrees of freedom with multiple imputation. *Biometrika*, 86(4), 948–955.
- Buuren, S. van. (2021). *Flexible imputation of missing data* (Second edition). Chapman & Hall/CRC.
- Collins, L. M., Schafer, J. L., & Kam, C.-M. (2001). A comparison of inclusive and restrictive strategies in modern missing data procedures. *Psychological Methods*, 6(4), 330–351. <https://doi.org/10.1037/1082-989X.6.4.330>
- Croon, M. A., & Van Veldhoven, M. J. P. M. (2007). Predicting group-level outcome variables from variables measured at the individual level: A latent variable multi-level model. *Psychological Methods*, 12(1), 45–57. <https://doi.org/10.1037/1082-989X.12.1.45>
- Daniel McNeish. (2016a). On Using Bayesian Methods to Address Small Sample Problems. *Structural Equation Modeling: A Multidisciplinary Journal*, 23(5), 750–773. <https://doi.org/10.1080/10705511.2016.1186549>
- Daniel McNeish. (2016b). Using Data-Dependent Priors to Mitigate Small Sample Bias in Latent Growth Models: A Discussion and Illustration Using M Plus. *Journal of Educational and Behavioral Statistics*, 41(1), 27–56. <https://doi.org/10.3102/1076998615621299>
- Daniel McNeish. (2017). Small Sample Methods for Multilevel Modeling: A Colloquial Elucidation of REML and the Kenward-Roger Correction. *Multivariate Behavioral Research*, 52(5), 661–670. <https://doi.org/10.1080/00273171.2017.1344538>
- Daniels, M. J., Wang, C., & Marcus, B. H. (2014). Fully Bayesian inference under ignorable missingness in the presence of auxiliary covariates. *Biometrics*, 70(1), 62–72. <https://doi.org/10.1111/biom.12121>
- Depaoli, S., & Clifton, J. P. (2015). A Bayesian Approach to Multilevel Structural Equation Modeling With Continuous and Dichotomous Outcomes. *Structural Equation Modeling: A Multidisciplinary Journal*, 22(3), 327–351. <https://doi.org/10.1080/10705511.2014.937849>
- Elff, M., Heisig, J. P., Schaeffer, M., & Shikano, S. (2021). Multilevel Analysis with Few Clusters: Improving Likelihood-Based Methods to Provide Unbiased Estimates and Accurate Inference. *British Journal of Political Science*, 51(1), 412–426. <https://doi.org/10.1017/S0007123419000097>
- Enders, C. K. (2022). *Applied Missing Data Analysis* (2nd ed). Guilford Press.

- Enders, C. K. (2025). Missing data: An update on the state of the art. *Psychological Methods*, 30(2), 322–339. <https://doi.org/10.1037/met0000563>
- Enders, C. K., Du, H., & Keller, B. T. (2020). A model-based imputation procedure for multilevel regression models with random coefficients, interaction effects, and nonlinear terms. *Psychological Methods*, 25(1), 88–112. <https://doi.org/10.1037/met0000228>
- Enders, C. K., Mistler, S. A., & Keller, B. T. (2016). Multilevel multiple imputation: A review and evaluation of joint modeling and chained equations imputation. *Psychological Methods*, 21(2), 222–240. <https://doi.org/10.1037/met0000063>
- Erler, N. S., Rizopoulos, D., Rosmalen, J. V., Jaddoe, V. W. V., Franco, O. H., & Lesaffre, E. M. E. H. (2016). Dealing with missing covariates in epidemiologic studies: A comparison between multiple imputation and a full Bayesian approach. *Statistics in Medicine*, 35(17), 2955–2974. <https://doi.org/10.1002/sim.6944>
- Fahrmeir, L., Kneib, T., Lang, S., & Marx, B. D. (2021). *Regression: Models, Methods and Applications*. Springer Berlin Heidelberg. <https://doi.org/10.1007/978-3-662-63882-8>
- Gelman, A. (2006). Prior distributions for variance parameters in hierarchical models (comment on article by Browne and Draper). *Bayesian Analysis*, 1(3). <https://doi.org/10.1214/06-BA117A>
- Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A., & Rubin, D. B. (2025). *Bayesian Data Analysis Third edition (with errors fixed as of 20 February 2025)* (3rd ed.).
- Gelman, A., & Rubin, D. B. (1992). Inference from Iterative Simulation Using Multiple Sequences. *Statistical Science*, 7(4). <https://doi.org/10.1214/ss/1177011136>
- Graham, J. W. (2003). Adding Missing-Data-Relevant Variables to FIML-Based Structural Equation Models. *Structural Equation Modeling: A Multidisciplinary Journal*, 10(1), 80–100. https://doi.org/10.1207/S15328007SEM1001_4
- Graham, J. W. (2009). Missing Data Analysis: Making It Work in the Real World. *Annual Review of Psychology*, 60(1), 549–576. <https://doi.org/10.1146/annurev.psych.58.110405.085530>
- Graham, J. W., Olchowski, A. E., & Gilreath, T. D. (2007). How Many Imputations are Really Needed? Some Practical Clarifications of Multiple Imputation Theory. *Prevention Science*, 8(3), 206–213. <https://doi.org/10.1007/s11121-007-0070-9>
- Grund, S., Lüdtke, O., & Robitzsch, A. (2016). Multiple Imputation of Multilevel Missing Data: An Introduction to the R Package pan. *Sage Open*, 6(4), 2158244016668220. <https://doi.org/10.1177/2158244016668220>
- Grund, S., Lüdtke, O., & Robitzsch, A. (2018). Multiple Imputation of Missing Data for Multilevel Models: Simulations and Recommendations. *Organizational Research Methods*, 21(1), 111–149. <https://doi.org/10.1177/1094428117703686>
- Grund, S., Lüdtke, O., & Robitzsch, A. (2019). Missing data in multilevel research. In S. E. Humphrey & J. M. LeBreton (Eds.), *The handbook of multilevel theory, measurement, and analysis*. (pp. 365–386). American Psychological Association. <https://doi.org/10.1037/0000115-017>
- Grund, S., Robitzsch, A., & Luedtke, O. (2015). *Mitml: Tools for Multiple Imputation in Multilevel Modeling* (pp. 0.4–5). Comprehensive R Archive Network. <https://doi.org/10.32614/CRAN.package.mitml>
- Holtmann, J., Koch, T., Lochner, K., & Eid, M. (2016). A Comparison of ML, WLSMV, and Bayesian Methods for Multilevel Structural Equation Models in Small Samples: A Simulation Study. *Multivariate Behavioral Research*, 51(5), 661–680. <https://doi.org/10.1080/00273171.2016.1208074>
- Kackar, R. N., & Harville, D. A. (1981). Unbiasedness of two-stage estimation and

- prediction procedures for mixed linear models. *Communications in Statistics - Theory and Methods*, 10(13), 1249–1261. <https://doi.org/10.1080/03610928108828108>
- Kackar, R. N., & Harville, D. A. (1984). Approximations for Standard Errors of Estimators of Fixed and Random Effect in Mixed Linear Models. *Journal of the American Statistical Association*, 79(388), 853. <https://doi.org/10.2307/2288715>
- Kahan, B. C., Forbes, G., Ali, Y., Jairath, V., Bremner, S., Harhay, M. O., Hooper, R., Wright, N., Eldridge, S. M., & Leyrat, C. (2016). Increased risk of type I errors in cluster randomised trials with small or medium numbers of clusters: A review, reanalysis, and simulation study. *Trials*, 17(1), 438. <https://doi.org/10.1186/s13063-016-1571-2>
- Kenward, M. G., & Roger, J. H. (1997). Small Sample Inference for Fixed Effects from Restricted Maximum Likelihood. *Biometrics*, 53(3), 983. <https://doi.org/10.2307/2533558>
- Leonhart, R. (2013). *Lehrbuch Statistik: Einstieg und Vertiefung* (3., überarbeitete und erweiterte Auflage). Verlag Hans Huber.
- Little, R. J. A., & Rubin, D. B. (1987). *Statistical analysis with missing data* (3. [Dr.]). Wiley.
- Little, R. J. A., & Rubin, D. B. (2020). *Statistical analysis with missing data* (3rd edition). Wiley. <https://doi.org/10.1002/9781119482260>
- Lüdtke, O., Marsh, H. W., Robitzsch, A., Trautwein, U., Asparouhov, T., & Muthén, B. (2008). The multilevel latent covariate model: A new, more reliable approach to group-level effects in contextual studies. *Psychological Methods*, 13(3), 203–229. <https://doi.org/10.1037/a0012869>
- Lüdtke, O., Robitzsch, A., & Grund, S. (2017). Multiple imputation of missing data in multilevel designs: A comparison of different strategies. *Psychological Methods*, 22(1), 141–165. <https://doi.org/10.1037/met0000096>
- Lüdtke, O., Robitzsch, A., Trautwein, U., & Köller, O. (2007). Umgang mit fehlenden Werten in der psychologischen Forschung. *Psychologische Rundschau*, 58(2), 103–117. <https://doi.org/10.1026/0033-3042.58.2.103>
- Lüdtke, O., Robitzsch, A., & West, S. G. (2020). Regression models involving nonlinear effects with missing data: A sequential modeling approach using Bayesian estimation. *Psychological Methods*, 25(2), 157–181. <https://doi.org/10.1037/met000233>
- McNeish, D. (2017). Missing data methods for arbitrary missingness with small samples. *Journal of Applied Statistics*, 44(1), 24–39. <https://doi.org/10.1080/02664763.2016.1158246>
- McNeish, D. (2018). Brief Research Report: Growth Models With Small Samples and Missing Data. *The Journal of Experimental Education*, 86(4), 690–701. <https://doi.org/10.1080/00220973.2017.1369384>
- McNeish, D. M., & Stapleton, L. M. (2016). The Effect of Small Sample Size on Two-Level Model Estimates: A Review and Illustration. *Educational Psychology Review*, 28(2), 295–314. <https://doi.org/10.1007/s10648-014-9287-x>
- McNeish, D., & Stapleton, L. M. (2016). Modeling Clustered Data with Very Few Clusters. *Multivariate Behavioral Research*, 51(4), 495–518. <https://doi.org/10.1080/00273171.2016.1167008>
- Morris, T. P., White, I. R., & Crowther, M. J. (2019). Using simulation studies to evaluate statistical methods. *Statistics in Medicine*, 38(11), 2074–2102. <https://doi.org/10.1002/sim.8086>
- Muthén, B., & Asparouhov, T. (2012). Bayesian structural equation modeling: A more flexible representation of substantive theory. *Psychological Methods*, 17(3),

- 313–335. <https://doi.org/10.1037/a0026802>
- Pandey, S., & Bright, C. L. (2008). What Are Degrees of Freedom? *Social Work Research*, 32(2), 119–128. <https://doi.org/10.1093/swr/32.2.119>
- Papaioannou, A., Marsh, H. W., & Theodorakis, Y. (2004). A Multilevel Approach to Motivational Climate in Physical Education and Sport Settings: An Individual or a Group Level Construct? *Journal of Sport and Exercise Psychology*, 26(1), 90–118. <https://doi.org/10.1123/jsep.26.1.90>
- Quartagno, M., & Carpenter, J. (2023). *jomo: A package for multilevel joint modelling multiple imputation* [Manual].
- R Core Team. (2025). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing.
- Raudenbush, S., & Bryk, A. S. (2010). *Hierarchical linear models: Applications and data analysis methods* (2. ed., [Nachdr.]). Sage Publ.
- Robitzsch, A., & Luedtke, O. (2024). *Mdmb: Model Based Treatment of Missing Data* (pp. 1.9–22) [Manual]. Comprehensive R Archive Network. <https://doi.org/10.32614/CRAN.package.mdmb>
- Rubin, D. B. (1976). Inference and missing data. *Biometrika*, 63(3), 581–592. <https://doi.org/10.1093/biomet/63.3.581>
- Rubin, D. B. (1987). *Multiple imputation for nonresponse in surveys*. Wiley. <https://doi.org/10.1002/9780470316696>
- Schaalje, G. B., McBride, J. B., & Fellingham, G. W. (2002). Adequacy of approximations to distributions of test statistics in complex mixed linear models. *Journal of Agricultural, Biological, and Environmental Statistics*, 7(4), 512–524. <https://doi.org/10.1198/108571102726>
- Schafer, J. L., & Graham, J. W. (2002). Missing data: Our view of the state of the art. *Psychological Methods*, 7(2), 147–177. <https://doi.org/10.1037/1082-989X.7.2.147>
- Shin, D., & Shin, Y. (2025). *Compatible Imputation for Hierarchical Linear Models with Incomplete Data: Interaction Effects of Continuous and Categorical Covariates MAR* (arXiv:2502.07033). arXiv. <https://doi.org/10.48550/arXiv.2502.07033>
- Shin, D., Shin, Y., & Hagiwara, N. (2025). *Bayesian Estimation of Hierarchical Linear Models from Incomplete Data: Cluster-Level Interaction Effects and Small Sample Sizes* (arXiv:2405.21020). arXiv. <https://doi.org/10.48550/arXiv.2405.21020>
- Siepe, B. S., Bartoš, F., Morris, T. P., Boulesteix, A.-L., Heck, D. W., & Pawel, S. (2024). Simulation studies for methodological research in psychology: A standardized template for planning, preregistration, and reporting. *Psychological Methods*. <https://doi.org/10.1037/met0000695>
- Snijders, T. A. B., & Bosker, R. J. (2012). *Multilevel analysis: An introduction to basic and advanced multilevel modeling* (2nd ed). Sage.
- Tanner, M. A., & Wong, W. H. (n.d.). *The Calculation of Posterior Distributions by Data Augmentation*.
- Van De Schoot, R., Kaplan, D., Denissen, J., Asendorpf, J. B., Neyer, F. J., & Van Aken, M. A. G. (2014). A Gentle Introduction to Bayesian Analysis: Applications to Developmental Research. *Child Development*, 85(3), 842–860. <https://doi.org/10.1111/cdev.12169>
- Vehtari, A., Gelman, A., Simpson, D., Carpenter, B., & Bürkner, P.-C. (2021). Rank-Normalization, Folding, and Localization: An Improved $\hat{R}$ for Assessing Convergence of MCMC (with Discussion). *Bayesian Analysis*, 16(2). <https://doi.org/10.1214/20-BA1221>
- Von Hippel, P. T. (2013). The Bias and Efficiency of Incomplete-Data Estimators in Small Univariate Normal Samples. *Sociological Methods & Research*, 42(4), 531–558. <https://doi.org/10.1177/0049124113494582>

- Yucel, R. M. (2008). Multiple imputation inference for multivariate multilevel continuous data with ignorable non-response. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 366(1874), 2389–2403. <https://doi.org/10.1098/rsta.2008.0038>
- Zitzmann, S., Lüdtke, O., & Robitzsch, A. (2015). A Bayesian Approach to More Stable Estimates of Group-Level Effects in Contextual Studies. *Multivariate Behavioral Research*, 50(6), 688–705. <https://doi.org/10.1080/00273171.2015.1090899>

Appendix A

Derivations of equations of data-generating model

A.1 Calculation of σ_w^2

The data-generating model for W_j is:

$$W_j = a \cdot \bar{X}_j + e_{Wj}, \quad e_{Wj} \sim N(0, \sigma_w^2) \quad (\text{A.1})$$

The variance of W_j is therefore:

$$\text{Var}(W_j) = \text{Var}(a \cdot \bar{X}_j + e_{Wj}) = a^2 \cdot \text{Var}(\bar{X}_j) + \text{Var}(e_{Wj}) \quad (\text{A.2})$$

Substituting the formula for slope calculation $a = \frac{\text{Cov}(W_j, \bar{X}_j)}{\text{Var}(\bar{X}_j)}$ into Equation [A.2](#) yields:

$$\begin{aligned} \text{Var}(W_j) &= \frac{\text{Cov}(W_j, \bar{X}_j)^2}{\text{Var}(\bar{X}_j)^2} \cdot \text{Var}(\bar{X}_j) + \text{Var}(e_{Wj}) \\ &= \frac{\text{Cov}(W_j, \bar{X}_j)^2}{\text{Var}(\bar{X}_j)} + \text{Var}(e_{Wj}) \end{aligned}$$

This term can be simplified using $\text{Cov}(A, B) = \text{Corr}(A, B) \cdot \text{SD}(A) \cdot \text{SD}(B)$:

$$\begin{aligned} \text{Var}(W_j) &= \frac{\text{Cov}(W_j, \bar{X}_j)^2}{\text{Var}(\bar{X}_j)} + \text{Var}(e_{Wj}) \\ &= \frac{(\text{Corr}(W_j, \bar{X}_j) \cdot \text{SD}(W_j) \cdot \text{SD}(\bar{X}_j))^2}{\text{Var}(\bar{X}_j)} + \text{Var}(e_{Wj}) \\ &= \frac{\text{Corr}(W_j, \bar{X}_j)^2 \cdot \text{Var}(W_j) \cdot \text{Var}(\bar{X}_j)}{\text{Var}(\bar{X}_j)} + \text{Var}(e_{Wj}) \\ &= \text{Corr}(W_j, \bar{X}_j)^2 \cdot \text{Var}(W_j) + \text{Var}(e_{Wj}) \end{aligned}$$

Thus,

$$\begin{aligned}
Var(W_j) &= Corr(W_j, \bar{X}_j)^2 \cdot Var(W_j) + Var(e_{Wj}) \\
Var(e_{Wj}) &= Var(W_j) - Corr(W_j, \bar{X}_j)^2 \cdot Var(W_j) \\
&= Var(W_j) (1 - Corr(W_j, \bar{X}_j)^2)
\end{aligned}$$

Since $Var(W_j) = 1$:

$$Var(e_{Wj}) = 1 - Corr(W_j, \bar{X}_j)^2$$

In short notation:

$$\sigma_w^2 = 1 - \rho_{\bar{X}W}^2$$

A.2 Calculation of the slope to generate W_j

$$a = \frac{Cov(W_j, \bar{X}_j)}{Var(\bar{X}_j)}$$

The variance of $\bar{X}_j$ can be derived as follows:

$$\begin{aligned}
Var(\bar{X}_j) &= Var\left(\frac{1}{n} \sum_{i=1}^n X_{ij}\right) \\
&= Var\left(\frac{1}{n} \sum_{i=1}^n (u_{Xj} + e_{Xij})\right) \\
&= Var\left(\frac{1}{n} \sum_{i=1}^n u_{Xj} + \frac{1}{n} \sum_{i=1}^n e_{Xij}\right) \\
&= Var\left(u_{Xj} + \frac{1}{n} \sum_{i=1}^n e_{Xij}\right) \\
&= Var(u_{Xj}) + Var\left(\frac{1}{n} \sum_{i=1}^n e_{Xij}\right) \\
&= Var(u_{Xj}) + \frac{1}{n^2} Var\left(\sum_{i=1}^n e_{Xij}\right) \\
&= Var(u_{Xj}) + \frac{1}{n^2} \sum_{i=1}^n Var(e_{Xij}) \\
&= \tau_x^2 + \frac{n \cdot \sigma_X^2}{n^2} \\
&= \tau_x^2 + \frac{\sigma_X^2}{n}
\end{aligned}$$

where n denotes the group size (cluster size), assumed to be constant across groups. The covariance $Cov(W_j, \bar{X}_j)$ is therefore:

$$\begin{aligned}
Cov(W_j, \bar{X}_j) &= \rho_{\bar{X}W} \cdot SD(\bar{X}_j) \cdot SD(W_j) \\
&= \rho_{\bar{X}W} \cdot \sqrt{Var(\bar{X}_j)} \cdot \sqrt{Var(W_j)} \\
&= \rho_{\bar{X}W} \cdot \sqrt{\tau_x^2 + \frac{\sigma_X^2}{n}} \cdot 1
\end{aligned}$$

A.3 Variances of random components of Y

Individual and group residual variances of Y ($Var(u_{0j}) = \psi^2$ and $Var(e_{ij}) = \sigma^2$) are determined to obtain a prespecified residual ICC of Y :

$$ICC_{Y|X,W} = \frac{\psi^2}{\psi^2 + \sigma^2}$$

Because $Var(Y) = 1$ and residuals are uncorrelated with predictors and with each other,

$$\psi^2 + \sigma^2 = Var(\text{Residual}) = 1 - Var(\text{fixed})$$

where $Var(\text{fixed})$ denotes the variance explained by the fixed effects of the model. The variance components can therefore be expressed as:

$$Var(u_{0j}) = ICC_{Y|X,W} \cdot (1 - Var(\text{fixed}))$$

$$Var(e_{ij}) = (1 - ICC_{Y|X,W}) \cdot (1 - Var(\text{fixed}))$$

$Var(\text{fixed})$ can be calculated as:

$$Var(\text{fixed}) = Var(\gamma_{10}(X_{ij} - \bar{X}_{.j}) + \gamma_{01}\bar{X}_{.j} + \gamma_{02}W_j)$$

The terms $(X_{ij} - \bar{X}_{.j})$ and $\bar{X}_{.j}$ are uncorrelated. W_j is correlated with $\bar{X}_{.j}$ but not with X_{ij} (see Equation Equation A.1), and $Var(W_j)$ is set to 1, which leads to:

$$Var(\text{fixed}) = \gamma_{10}^2 \cdot Var(X_{ij} - \bar{X}_{.j}) + \gamma_{01}^2 \cdot Var(\bar{X}_{.j}) + \gamma_{02}^2 + 2\gamma_{01}\gamma_{02}Cov(\bar{X}_{.j}, W_j)$$

Using the decomposition of X_{ij} introduced in Section Section 3.1.1,

$$X_{ij} = u_{Xj} + e_{Xij},$$

the variance of $(X_{ij} - \bar{X}_{.j})$ can be derived as follows:

$$\begin{aligned}
Var(X_{ij} - \bar{X}_{.j}) &= Var(X_{ij}) + Var(\bar{X}_{.j}) - 2 \cdot Cov(X_{ij}, \bar{X}_{.j}) \\
&= Var(X_{ij}) + Var(\bar{X}_{.j}) - 2 \cdot \left(Var(u_{Xj}) + \frac{1}{n} \sum_{i'=1}^n Cov(u_{Xj}, e_{Xi'j}) \right. \\
&\quad \left. + Cov(e_{Xij}, u_{Xj}) + \frac{1}{n} \sum_{i'=1}^n Cov(e_{Xij}, e_{Xi'j}) \right) \\
&= Var(X_{ij}) + Var(\bar{X}_{.j}) - 2 \cdot \left(Var(u_{Xj}) + \frac{1}{n} Var(e_{Xij}) \right) \\
&= Var(X_{ij}) + Var(\bar{X}_{.j}) - 2 \cdot Var(\bar{X}_{.j}) \\
&= Var(X_{ij}) - Var(\bar{X}_{.j}) \\
&= \tau_x^2 + \sigma_X^2 - \left(\tau_x^2 + \frac{\sigma_X^2}{n} \right) \\
&= \sigma_X^2 - \frac{\sigma_X^2}{n}
\end{aligned}$$

Because γ_{02} is set to 0, $Var(\text{fixed})$ can be calculated as

$$Var(\text{fixed}) = \gamma_{10}^2 \left(\sigma_X^2 - \frac{\sigma_X^2}{n} \right) + \gamma_{01}^2 \left(\tau_x^2 + \frac{\sigma_X^2}{n} \right).$$

The variances of the random components can thus be calculated like this:

$$\begin{aligned}
\psi^2 &= ICC_{Y|X,W} \cdot (1 - \gamma_{10}^2) \left(\sigma_X^2 - \frac{\sigma_X^2}{n_j} \right) + \gamma_{01}^2 \left(\tau_x^2 + \frac{\sigma_X^2}{n_j} \right) \\
\sigma^2 &= (1 - ICC_{Y|X,W}) \cdot (1 - \gamma_{10}^2) \left(\sigma_X^2 - \frac{\sigma_X^2}{n_j} \right) + \gamma_{01}^2 \left(\tau_x^2 + \frac{\sigma_X^2}{n_j} \right)
\end{aligned}$$

Appendix B

Convergence analysis for MI and fully Bayesian MCMC

Convergence behaviour was tested in one run of the most challenging condition ($N_2 = 15, \gamma_{01} = .4, \beta = .3, ICC = .1$).

B.1 Diagnostics of multiple imputation

Convergence of the imputation model was assessed using potential scale reduction factors (Gelman & Rubin, 1992). Scale reduction factors that are substantially larger than 1 indicate that convergence was not yet reached fully; an acceptable threshold can be set at 1.1 (Gelman et al., 2025, pp. 285–287), while a stricter recommendation was recently set to be 1.01 (Vehtari et al., 2021). Given the simulation setting with aggregated results over many replications, the first threshold was considered sufficient.

All values were close to 1, indicating satisfactory convergence across all parameters. A maximum $\hat{R}$ of 1.026 was observed for $\beta_{Y,X}$. Diagnostic plots are shown in Figure B.1 - @fig-MIdiag3 for this parameter and the most critical parameters, the level-2 residual variances of Y and X .

Trace plots showed no signs of substantial change after burn-in. Inspection of the autocorrelation plots indicated that autocorrelation diminished quickly across increasing lags and did not persist over longer distances between samples. The distance of 300 lags between imputation that was chosen in this simulation was therefore sufficient.

B.2 Diagnostics of fully Bayesian analysis

$\hat{R}$ -values were close to 1, except for the level-2 variance of X ($\hat{R} = 1.05$) and the regression coefficient of the group mean of X on W ($\hat{R} = 1.03$). Diagnostic plots of these parameters and the furthermore critical parameter τ_{y0}^2 are shown in Figure B.4. Trace plots showed no drifting, and autocorrelation plots indicated sufficiently low autocorrelation after few lags.




FIGURE B.1: Trace and autocorrelation plot in an exemplary simulation run for $\beta_{Y,X}$ (multiple imputation).




FIGURE B.2: Trace and autocorrelation plot in an exemplary simulation run for the level-2 variance of $Y \tau_{y0}^2$ (multiple imputation).




FIGURE B.3: Trace and autocorrelation plot in an exemplary simulation run for the level-2 variance of $X \tau_{x0}^2$ (multiple imputation).




FIGURE B.4: Trace and autocorrelation plot for the parameter $\beta_{wx}^{(imp)}$, representing the effect of the group mean of x on w in the imputation model (fully Bayesian estimation).




FIGURE B.5: Trace and autocorrelation plot in an exemplary simulation run for the level-2 variance of X τ_{x0}^2 (fully Bayesian analysis)




FIGURE B.6: Trace and autocorrelation plot in an exemplary simulation run for the level-2 variance of Y τ_{x0}^2 (fully Bayesian analysis)

Appendix C

Complete simulation results for all performance measures

TABLE C.1: Simulation results ($N_2 = 15$, $\gamma_{01} = 0.0$)

Parameter	CD				LD				MI-R				MI-a				Bayes			
	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW
MCAR																				
ICC = 0.1																				
γ_{01}	0.002	0.942	0.284	1.170	0.037	0.941	0.300	1.180	0.005	0.966	0.252	1.115	0.005	0.986	0.252	1.284	0.003	0.980	0.291	1.456
γ_{10}	-0.006	0.951	0.084	0.335	-0.006	0.944	0.104	0.414	-0.008	0.943	0.103	0.409	-0.008	0.944	0.103	0.414	-0.010	0.946	0.100	0.401
ICC = 0.3																				
γ_{01}	0.000	0.960	0.272	1.174	0.012	0.953	0.280	1.177	-0.001	0.942	0.278	1.126	-0.001	0.972	0.278	1.278	-0.003	0.980	0.288	1.384
γ_{10}	0.001	0.942	0.089	0.336	0.000	0.931	0.111	0.415	-0.005	0.938	0.108	0.411	-0.005	0.939	0.108	0.416	-0.006	0.934	0.108	0.405
MAR																				
ICC = 0.1																				
γ_{01}	-0.002	0.949	0.283	1.192	0.031	0.947	0.284	1.189	0.007	0.970	0.247	1.131	0.007	0.985	0.247	1.305	0.001	0.987	0.289	1.481
γ_{10}	0.000	0.942	0.087	0.336	-0.002	0.954	0.106	0.416	-0.007	0.949	0.104	0.412	-0.007	0.950	0.104	0.418	-0.007	0.954	0.102	0.403
ICC = 0.3																				
γ_{01}	-0.016	0.943	0.279	1.175	0.007	0.949	0.279	1.175	-0.014	0.943	0.275	1.130	-0.014	0.968	0.275	1.285	-0.016	0.975	0.291	1.390
γ_{10}	0.001	0.941	0.087	0.337	-0.001	0.948	0.105	0.414	-0.004	0.943	0.102	0.411	-0.004	0.945	0.102	0.417	-0.005	0.949	0.101	0.405

Note:

Bias = average deviation from the true parameter; Cov = empirical coverage probability; SD = empirical standard deviation; CIW = mean confidence/credible interval width. MCAR = missing completely at random; MAR = missing at random. CD = complete data; LD = listwise deletion; MI-R = multiple imputation with Rubin's rules; MI-a = multiple imputation with adjusted degrees of freedom; Bayes = fully Bayesian estimation.

TABLE C.2: Simulation results ($N_2 = 15$, $\gamma_{01} = 0.4$)

	CD				LD				MI-R				MI-a				Bayes			
	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW
MCAR																				
ICC = 0.1																				
γ_{01}	-0.008	0.953	0.275	1.174	-0.038	0.968	0.269	1.183	-0.075	0.965	0.239	1.151	-0.075	0.981	0.239	1.329	-0.036	0.985	0.281	1.516
γ_{10}	0.000	0.952	0.083	0.330	0.000	0.952	0.102	0.404	-0.001	0.950	0.101	0.401	-0.001	0.953	0.101	0.407	-0.003	0.954	0.099	0.393
ICC = 0.3																				
γ_{01}	0.003	0.955	0.272	1.154	-0.013	0.949	0.272	1.156	-0.016	0.951	0.275	1.122	-0.016	0.967	0.275	1.277	-0.008	0.971	0.290	1.385
γ_{10}	0.002	0.949	0.083	0.326	0.003	0.935	0.106	0.402	-0.001	0.944	0.103	0.397	-0.001	0.948	0.103	0.402	-0.003	0.942	0.101	0.391
MAR																				
ICC = 0.1																				
γ_{01}	-0.004	0.959	0.276	1.151	-0.050	0.942	0.283	1.169	-0.078	0.965	0.238	1.124	-0.078	0.977	0.238	1.299	-0.039	0.982	0.276	1.461
γ_{10}	-0.007	0.954	0.084	0.331	-0.005	0.952	0.105	0.408	-0.007	0.956	0.102	0.404	-0.007	0.959	0.102	0.409	-0.010	0.952	0.101	0.396
ICC = 0.3																				
γ_{01}	0.006	0.962	0.269	1.162	-0.019	0.956	0.275	1.165	-0.020	0.947	0.271	1.137	-0.020	0.976	0.271	1.296	-0.008	0.977	0.290	1.407
γ_{10}	-0.001	0.945	0.086	0.327	0.000	0.948	0.105	0.403	-0.003	0.946	0.102	0.398	-0.003	0.948	0.102	0.403	-0.006	0.951	0.102	0.393

Note:

Abbreviations as listed in Table C.1

TABLE C.3: Simulation results ($N_2 = 30$, $\gamma_{01} = 0.0$)

		CD				LD				MI-R				MI-a				Bayes			
		Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW
MCAR																					
ICC = 0.1																					
γ_{01}		0.003	0.945	0.188	0.755	0.042	0.939	0.186	0.757	0.014	0.962	0.173	0.766	0.014	0.976	0.173	0.812	0.009	0.963	0.199	0.864
γ_{10}		-0.005	0.953	0.059	0.236	-0.006	0.953	0.073	0.291	-0.007	0.954	0.073	0.290	-0.007	0.954	0.073	0.292	-0.008	0.950	0.072	0.283
ICC = 0.3																					
γ_{01}		-0.004	0.957	0.185	0.758	0.010	0.964	0.185	0.760	-0.003	0.955	0.190	0.769	-0.003	0.964	0.190	0.810	-0.003	0.965	0.192	0.826
γ_{10}		0.001	0.942	0.063	0.237	0.000	0.943	0.077	0.291	-0.003	0.939	0.077	0.291	-0.003	0.940	0.077	0.293	-0.003	0.943	0.076	0.284
MAR																					
ICC = 0.1																					
γ_{01}		0.006	0.941	0.191	0.757	0.039	0.947	0.192	0.761	0.010	0.965	0.177	0.767	0.010	0.972	0.177	0.813	0.003	0.965	0.203	0.870
γ_{10}		0.000	0.944	0.061	0.236	-0.002	0.945	0.075	0.290	-0.003	0.950	0.074	0.288	-0.003	0.951	0.074	0.290	-0.004	0.948	0.073	0.282
ICC = 0.3																					
γ_{01}		0.008	0.969	0.179	0.765	0.027	0.956	0.183	0.768	0.011	0.958	0.186	0.777	0.011	0.968	0.186	0.820	0.010	0.967	0.188	0.836
γ_{10}		0.000	0.944	0.062	0.238	0.000	0.948	0.076	0.293	-0.003	0.954	0.075	0.294	-0.003	0.954	0.075	0.296	-0.003	0.946	0.074	0.286

Note:

Abbreviations as listed in Table C.1

TABLE C.4: Simulation results ($N_2 = 30$, $\gamma_{01} = 0.4$)

	CD				LD				MI-R				MI-a				Bayes			
	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW
MCAR																				
ICC = 0.1																				
γ_{01}	0.005	0.950	0.189	0.744	-0.034	0.956	0.187	0.748	-0.052	0.956	0.175	0.767	-0.052	0.966	0.175	0.813	-0.009	0.970	0.204	0.865
γ_{10}	0.000	0.966	0.056	0.231	-0.001	0.967	0.068	0.284	0.001	0.966	0.066	0.283	0.001	0.969	0.066	0.285	-0.003	0.967	0.065	0.277
ICC = 0.3																				
γ_{01}	0.007	0.943	0.188	0.736	-0.008	0.941	0.191	0.740	-0.002	0.938	0.194	0.753	-0.002	0.953	0.194	0.794	0.004	0.951	0.201	0.811
γ_{10}	0.000	0.963	0.057	0.229	0.003	0.956	0.069	0.282	0.001	0.950	0.067	0.281	0.001	0.954	0.067	0.283	0.000	0.954	0.067	0.275
MAR																				
ICC = 0.1																				
γ_{01}	0.005	0.948	0.191	0.744	-0.033	0.944	0.190	0.749	-0.057	0.952	0.174	0.772	-0.057	0.963	0.174	0.819	-0.015	0.958	0.204	0.868
γ_{10}	0.001	0.947	0.060	0.233	0.001	0.946	0.073	0.285	0.003	0.948	0.073	0.285	0.003	0.950	0.073	0.287	0.000	0.946	0.071	0.277
ICC = 0.3																				
γ_{01}	0.002	0.941	0.182	0.737	-0.014	0.947	0.180	0.738	-0.006	0.942	0.185	0.754	-0.006	0.951	0.185	0.795	0.000	0.954	0.191	0.813
γ_{10}	-0.002	0.948	0.059	0.230	-0.001	0.947	0.075	0.284	-0.003	0.950	0.072	0.283	-0.003	0.951	0.072	0.285	-0.004	0.948	0.072	0.277

Note:

Abbreviations as listed in Table C.1

TABLE C.5: Simulation results ($N_2 = 60$, $\gamma_{01} = 0.0$)

		CD				LD				MI-R				MI-a				Bayes			
		Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW
MCAR																					
ICC = 0.1																					
γ_{01}		-0.006	0.955	0.130	0.516	0.027	0.954	0.128	0.517	-0.002	0.971	0.124	0.542	-0.002	0.979	0.124	0.556	-0.010	0.961	0.139	0.575
γ_{10}		0.000	0.944	0.043	0.167	-0.001	0.953	0.053	0.205	-0.002	0.939	0.053	0.205	-0.002	0.940	0.053	0.206	-0.002	0.945	0.052	0.199
ICC = 0.3																					
γ_{01}		-0.002	0.946	0.127	0.518	0.017	0.955	0.127	0.519	0.001	0.949	0.132	0.536	0.001	0.954	0.132	0.549	0.000	0.954	0.133	0.550
γ_{10}		0.000	0.947	0.042	0.167	0.002	0.947	0.052	0.206	0.000	0.948	0.051	0.206	0.000	0.948	0.051	0.206	0.000	0.946	0.051	0.201
MAR																					
ICC = 0.1																					
γ_{01}		0.000	0.951	0.131	0.517	0.037	0.949	0.129	0.520	0.008	0.960	0.126	0.545	0.008	0.965	0.126	0.559	0.001	0.953	0.142	0.582
γ_{10}		-0.001	0.954	0.043	0.167	0.000	0.954	0.052	0.204	-0.001	0.956	0.051	0.205	-0.001	0.957	0.051	0.205	-0.001	0.953	0.050	0.198
ICC = 0.3																					
γ_{01}		0.005	0.948	0.131	0.522	0.021	0.949	0.131	0.522	0.003	0.946	0.137	0.539	0.003	0.949	0.137	0.552	0.003	0.956	0.137	0.556
γ_{10}		0.001	0.954	0.041	0.167	0.000	0.949	0.052	0.205	-0.001	0.947	0.052	0.205	-0.001	0.947	0.052	0.206	-0.001	0.944	0.051	0.201

Note:

Abbreviations as listed in Table C.1

TABLE C.6: Simulation results ($N_2 = 60$, $\gamma_{01} = 0.4$)

	CD				LD				MI-R				MI-a				Bayes			
	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW
MCAR																				
ICC = 0.1																				
γ_{01}	0.001	0.946	0.126	0.509	-0.039	0.939	0.124	0.509	-0.044	0.955	0.121	0.541	-0.044	0.960	0.121	0.556	-0.008	0.952	0.136	0.577
γ_{10}	0.001	0.960	0.041	0.164	0.001	0.961	0.050	0.201	0.004	0.954	0.049	0.200	0.004	0.955	0.049	0.200	0.001	0.958	0.049	0.195
ICC = 0.3																				
γ_{01}	-0.002	0.952	0.129	0.501	-0.021	0.947	0.128	0.502	-0.007	0.954	0.132	0.521	-0.007	0.957	0.132	0.534	-0.005	0.962	0.133	0.536
γ_{10}	-0.001	0.954	0.040	0.162	-0.001	0.956	0.049	0.199	-0.002	0.956	0.048	0.199	-0.002	0.956	0.048	0.200	-0.002	0.961	0.048	0.194
MAR																				
ICC = 0.1																				
γ_{01}	0.000	0.954	0.129	0.509	-0.035	0.943	0.128	0.512	-0.044	0.960	0.125	0.544	-0.044	0.962	0.125	0.559	-0.008	0.957	0.144	0.580
γ_{10}	0.002	0.926	0.043	0.163	0.001	0.942	0.051	0.200	0.003	0.941	0.050	0.199	0.003	0.942	0.050	0.200	0.001	0.941	0.050	0.194
ICC = 0.3																				
γ_{01}	-0.005	0.950	0.127	0.503	-0.022	0.956	0.125	0.504	-0.009	0.958	0.130	0.523	-0.009	0.961	0.130	0.536	-0.008	0.962	0.132	0.540
γ_{10}	-0.001	0.953	0.042	0.162	-0.002	0.944	0.051	0.199	-0.003	0.944	0.050	0.199	-0.003	0.945	0.050	0.200	-0.003	0.942	0.050	0.195

Note:

Abbreviations as listed in Table C.1

TABLE C.7: Monte Carlo standard errors of performance measures ($N_2 = 15$, $\gamma_{01} = 0.0$)

		CD				LD				MI-R				MI-a				Bayes			
		Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW
MCAR																					
ICC = 0.1																					
γ_{01}		0.009	0.007	0.006	0.010	0.009	0.007	0.007	0.010	0.008	0.006	0.006	0.010	0.008	0.004	0.006	0.012	0.009	0.004	0.007	0.014
γ_{10}		0.003	0.007	0.002	0.001	0.003	0.007	0.002	0.001	0.003	0.007	0.002	0.002	0.003	0.007	0.002	0.002	0.003	0.007	0.002	0.001
ICC = 0.3																					
γ_{01}		0.009	0.006	0.006	0.010	0.009	0.007	0.006	0.010	0.009	0.007	0.006	0.011	0.009	0.005	0.006	0.012	0.009	0.004	0.006	0.013
γ_{10}		0.003	0.007	0.002	0.001	0.004	0.008	0.002	0.001	0.003	0.008	0.002	0.002	0.003	0.008	0.002	0.002	0.003	0.008	0.002	0.001
MAR																					
ICC = 0.1																					
γ_{01}		0.009	0.007	0.006	0.011	0.009	0.007	0.006	0.011	0.008	0.005	0.006	0.011	0.008	0.004	0.006	0.013	0.009	0.004	0.006	0.016
γ_{10}		0.003	0.007	0.002	0.001	0.003	0.007	0.002	0.001	0.003	0.007	0.002	0.002	0.003	0.007	0.002	0.002	0.003	0.007	0.002	0.001
ICC = 0.3																					
γ_{01}		0.009	0.007	0.006	0.011	0.009	0.007	0.006	0.011	0.009	0.007	0.006	0.011	0.009	0.006	0.006	0.013	0.009	0.005	0.007	0.014
γ_{10}		0.003	0.007	0.002	0.001	0.003	0.007	0.002	0.001	0.003	0.007	0.002	0.002	0.003	0.007	0.002	0.002	0.003	0.007	0.002	0.001

Note:

All entries are Monte Carlo standard errors (MCSEs). Abbreviations as listed in Table C.1

TABLE C.8: Monte Carlo standard errors of performance measures ($N_2 = 15$, $\gamma_{01} = 0.4$)

	CD				LD				MI-R				MI-a				Bayes			
	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW
MCAR																				
ICC = 0.1																				
γ_{01}	0.009	0.007	0.006	0.010	0.008	0.006	0.006	0.010	0.008	0.006	0.005	0.010	0.008	0.004	0.005	0.013	0.009	0.004	0.006	0.020
γ_{10}	0.003	0.007	0.002	0.001	0.003	0.007	0.002	0.001	0.003	0.007	0.002	0.002	0.003	0.007	0.002	0.002	0.003	0.007	0.002	0.001
ICC = 0.3																				
γ_{01}	0.009	0.007	0.006	0.011	0.009	0.007	0.006	0.011	0.009	0.007	0.006	0.011	0.009	0.006	0.006	0.013	0.009	0.005	0.006	0.016
γ_{10}	0.003	0.007	0.002	0.001	0.003	0.008	0.002	0.001	0.003	0.007	0.002	0.002	0.003	0.007	0.002	0.002	0.003	0.007	0.002	0.001
MAR																				
ICC = 0.1																				
γ_{01}	0.009	0.006	0.006	0.010	0.009	0.007	0.006	0.010	0.008	0.006	0.005	0.010	0.008	0.005	0.005	0.012	0.009	0.004	0.006	0.014
γ_{10}	0.003	0.007	0.002	0.001	0.003	0.007	0.002	0.001	0.003	0.006	0.002	0.002	0.003	0.006	0.002	0.002	0.003	0.007	0.002	0.001
ICC = 0.3																				
γ_{01}	0.008	0.006	0.006	0.010	0.009	0.006	0.006	0.010	0.009	0.007	0.006	0.011	0.009	0.005	0.006	0.013	0.009	0.005	0.006	0.014
γ_{10}	0.003	0.007	0.002	0.001	0.003	0.007	0.002	0.001	0.003	0.007	0.002	0.002	0.003	0.007	0.002	0.002	0.003	0.007	0.002	0.001

Note:

All entries are Monte Carlo standard errors (MCSEs). Abbreviations as listed in Table C.1

TABLE C.9: Monte Carlo standard errors of performance measures ($N_2 = 30$, $\gamma_{01} = 0.0$)

	CD				LD				MI-R				MI-a				Bayes			
	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW
MCAR																				
ICC = 0.1																				
γ_{01}	0.006	0.007	0.004	0.005	0.006	0.008	0.004	0.004	0.005	0.006	0.004	0.005	0.005	0.005	0.004	0.005	0.006	0.006	0.004	0.006
γ_{10}	0.002	0.007	0.001	0.000	0.002	0.007	0.002	0.001	0.002	0.007	0.002	0.001	0.002	0.007	0.002	0.001	0.002	0.007	0.002	0.001
ICC = 0.3																				
γ_{01}	0.006	0.006	0.004	0.005	0.006	0.006	0.004	0.004	0.006	0.007	0.004	0.005	0.006	0.006	0.004	0.005	0.006	0.006	0.004	0.005
γ_{10}	0.002	0.007	0.001	0.000	0.002	0.007	0.002	0.001	0.002	0.008	0.002	0.001	0.002	0.008	0.002	0.001	0.002	0.007	0.002	0.001
MAR																				
ICC = 0.1																				
γ_{01}	0.006	0.007	0.004	0.005	0.006	0.007	0.004	0.005	0.006	0.006	0.004	0.005	0.006	0.005	0.004	0.006	0.006	0.006	0.005	0.006
γ_{10}	0.002	0.007	0.001	0.000	0.002	0.007	0.002	0.001	0.002	0.007	0.002	0.001	0.002	0.007	0.002	0.001	0.002	0.007	0.002	0.001
ICC = 0.3																				
γ_{01}	0.006	0.005	0.004	0.005	0.006	0.006	0.004	0.005	0.006	0.006	0.004	0.005	0.006	0.006	0.004	0.006	0.006	0.006	0.004	0.006
γ_{10}	0.002	0.007	0.001	0.000	0.002	0.007	0.002	0.001	0.002	0.007	0.002	0.001	0.002	0.007	0.002	0.001	0.002	0.007	0.002	0.001

Note:

All entries are Monte Carlo standard errors (MCSEs). Abbreviations as listed in Table C.1

TABLE C.10: Monte Carlo standard errors of performance measures ($N_2 = 30$, $\gamma_{01} = 0.4$)

	CD				LD				MI-R				MI-a				Bayes			
	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW
MCAR																				
ICC = 0.1																				
γ_{01}	0.006	0.007	0.004	0.005	0.006	0.006	0.004	0.004	0.006	0.006	0.004	0.005	0.006	0.006	0.004	0.006	0.006	0.005	0.005	0.006
γ_{10}	0.002	0.006	0.001	0.000	0.002	0.006	0.002	0.001	0.002	0.006	0.001	0.001	0.002	0.005	0.001	0.001	0.002	0.006	0.001	0.001
ICC = 0.3																				
γ_{01}	0.006	0.007	0.004	0.004	0.006	0.007	0.004	0.004	0.006	0.008	0.004	0.005	0.006	0.007	0.004	0.005	0.006	0.007	0.005	0.005
γ_{10}	0.002	0.006	0.001	0.000	0.002	0.006	0.002	0.001	0.002	0.007	0.001	0.001	0.002	0.007	0.001	0.001	0.002	0.007	0.001	0.001
MAR																				
ICC = 0.1																				
γ_{01}	0.006	0.007	0.004	0.005	0.006	0.007	0.004	0.004	0.006	0.007	0.004	0.005	0.006	0.006	0.004	0.006	0.006	0.006	0.005	0.006
γ_{10}	0.002	0.007	0.001	0.000	0.002	0.007	0.002	0.001	0.002	0.007	0.002	0.001	0.002	0.007	0.002	0.001	0.002	0.007	0.002	0.001
ICC = 0.3																				
γ_{01}	0.006	0.007	0.004	0.005	0.006	0.007	0.004	0.005	0.006	0.007	0.004	0.005	0.006	0.007	0.004	0.006	0.006	0.007	0.004	0.006
γ_{10}	0.002	0.007	0.001	0.000	0.002	0.007	0.002	0.001	0.002	0.007	0.002	0.001	0.002	0.007	0.002	0.001	0.002	0.007	0.002	0.001

Note:

All entries are Monte Carlo standard errors (MCSEs). Abbreviations as listed in Table C.1

TABLE C.11: Monte Carlo standard errors of performance measures ($N_2 = 60$, $\gamma_{01} = 0.0$)

	CD				LD				MI-R				MI-a				Bayes			
	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW
MCAR																				
ICC = 0.1																				
γ_{01}	0.004	0.007	0.003	0.002	0.004	0.007	0.003	0.002	0.004	0.005	0.003	0.003	0.004	0.005	0.003	0.003	0.004	0.006	0.003	0.003
γ_{10}	0.001	0.007	0.001	0.000	0.002	0.007	0.001	0.000	0.002	0.008	0.001	0.001	0.002	0.008	0.001	0.001	0.002	0.007	0.001	0.000
ICC = 0.3																				
γ_{01}	0.004	0.007	0.003	0.002	0.004	0.007	0.003	0.002	0.004	0.007	0.003	0.002	0.004	0.007	0.003	0.003	0.004	0.007	0.003	0.002
γ_{10}	0.001	0.007	0.001	0.000	0.002	0.007	0.001	0.000	0.002	0.007	0.001	0.001	0.002	0.007	0.001	0.001	0.002	0.007	0.001	0.000
MAR																				
ICC = 0.1																				
γ_{01}	0.004	0.007	0.003	0.002	0.004	0.007	0.003	0.002	0.004	0.006	0.003	0.003	0.004	0.006	0.003	0.003	0.004	0.007	0.003	0.003
γ_{10}	0.001	0.007	0.001	0.000	0.002	0.007	0.001	0.000	0.002	0.006	0.001	0.001	0.002	0.006	0.001	0.001	0.002	0.007	0.001	0.000
ICC = 0.3																				
γ_{01}	0.004	0.007	0.003	0.002	0.004	0.007	0.003	0.002	0.004	0.007	0.003	0.002	0.004	0.007	0.003	0.003	0.004	0.006	0.003	0.003
γ_{10}	0.001	0.007	0.001	0.000	0.002	0.007	0.001	0.000	0.002	0.007	0.001	0.001	0.002	0.007	0.001	0.001	0.002	0.007	0.001	0.000

Note:

All entries are Monte Carlo standard errors (MCSEs). Abbreviations as listed in Table C.1

TABLE C.12: Monte Carlo standard errors of performance measures ($N_2 = 60$, $\gamma_{01} = 0.4$)

	CD				LD				MI-R				MI-a				Bayes			
	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW	Bias	Cov	SD	CIW
MCAR																				
ICC = 0.1																				
γ_{01}	0.004	0.007	0.003	0.002	0.004	0.008	0.003	0.002	0.004	0.007	0.003	0.003	0.004	0.006	0.003	0.003	0.004	0.007	0.003	0.003
γ_{10}	0.001	0.006	0.001	0.000	0.002	0.006	0.001	0.000	0.002	0.007	0.001	0.001	0.002	0.007	0.001	0.001	0.002	0.006	0.001	0.000
ICC = 0.3																				
γ_{01}	0.004	0.007	0.003	0.002	0.004	0.007	0.003	0.002	0.004	0.007	0.003	0.002	0.004	0.006	0.003	0.003	0.004	0.006	0.003	0.002
γ_{10}	0.001	0.007	0.001	0.000	0.002	0.006	0.001	0.000	0.002	0.006	0.001	0.001	0.002	0.006	0.001	0.001	0.002	0.006	0.001	0.000
MAR																				
ICC = 0.1																				
γ_{01}	0.004	0.007	0.003	0.002	0.004	0.007	0.003	0.002	0.004	0.006	0.003	0.003	0.004	0.006	0.003	0.003	0.005	0.006	0.003	0.003
γ_{10}	0.001	0.008	0.001	0.000	0.002	0.007	0.001	0.000	0.002	0.007	0.001	0.001	0.002	0.007	0.001	0.001	0.002	0.007	0.001	0.000
ICC = 0.3																				
γ_{01}	0.004	0.007	0.003	0.002	0.004	0.006	0.003	0.002	0.004	0.006	0.003	0.002	0.004	0.006	0.003	0.002	0.004	0.006	0.003	0.002
γ_{10}	0.001	0.007	0.001	0.000	0.002	0.007	0.001	0.000	0.002	0.007	0.001	0.001	0.002	0.007	0.001	0.001	0.002	0.007	0.001	0.000

Note:

All entries are Monte Carlo standard errors (MCSEs). Abbreviations as listed in Table C.1

Appendix D

Default priors and sensitivity analysis

D.1 Default priors in `mdmb`

Default priors in `mdmb` are noninformative. For regression coefficients, a uniform prior on the interval $(-\infty, \infty)$ is approximated by a normal distribution with large variance, i.e., $\mathcal{N}(0, 10^{10})$.

For variance–covariance matrices on level 1 and 2, inverse-Wishart priors are used:

$$\Sigma \sim \mathcal{IW}(\nu, \mathbf{S})$$

where

- Σ denotes a generic covariance matrix (σ at level 1 and τ at level 2),
- ν denotes the degrees of freedom, and
- $\mathbf{S}$ is the scale matrix.

The default in `mdmb` is set to improper inverse-Wishart priors with negative degrees of freedom, which can be written as $\mathcal{IW}(-p-1, \mathbf{0})$ (Asparouhov & Muthen, 2010), where p denotes the dimension of the covariance matrix. $\mathbf{S} = \mathbf{0}$ corresponds to a null matrix of appropriate dimension.

D.2 Default priors in `jomo`

In `jomo`, inverse-Wishart priors are used for the covariance matrices at level 1 and 2 for imputation. The default in the `mitml` implementation is set to “least informative” proper priors with largest dispersion, i.e., smallest possible degrees of freedom, scaled by the identity matrix (Grund et al., 2015; Quartagno & Carpenter, 2023). For the three variables of the imputation model (X, Y, Z) in this thesis the scale matrix is:

$$\mathbf{S} = \mathbf{I} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

D.3 Sensitivity analysis with weakly informative priors in `jomo`

In this analysis, for the conditions when multiple imputation produced the highest bias, i.e. when $ICC = 0.1$ and $\gamma_{01} = .4$, alternative inverse Wishart priors were specified

TABLE D.1: Bias reduction with alternative priors

N2	Bias.d	rel. Bias.d	Bias.a	rel. Bias.a	Bias red.
MCAR					
15	-0.075	18.7%	-0.048	12.1%	35.5%
30	-0.062	15.4%	-0.033	8.3%	46.2%
60	-0.040	9.9%	-0.014	3.6%	63.7%
MAR					
15	-0.068	17.1%	-0.037	9.1%	46.6%
30	-0.059	14.8%	-0.033	8.2%	44.3%
60	-0.045	11.2%	-0.021	5.3%	52.3%

Note:

d = default priors, a = alternative priors, Bias red. = Bias reduction

for the level-2 covariance matrix. This was done by specifying a plausible prior guess for the level-2 covariance matrix. In the simulation study, this guess was based on the true parameter values; in empirical applications, it could be derived from previous research. This guess for the covariance matrix corresponds to the two standardized level-1 variables X and Y ($ICC = .1$) and the standardized level-2 variable W :

$$\tau_{\text{guess}} = \begin{bmatrix} .1 & 0 & 0 \\ 0 & .1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

τ denotes the level-2 covariance matrix of the random effects. The level-2 covariance prior was then defined as

$$\tau \sim \mathcal{IW}(\nu, \mathbf{S}), \quad \mathbf{S} = 3\tau_{\text{guess}}$$

Only the scale matrix was modified, while the degrees of freedom were kept at their default value, such that the prior remained weakly informative but was centered around more plausible variance values. The level-1 covariance matrix was kept at the default specification.

Bias was substantially reduced across all investigated conditions, with reductions ranging from approximately 35% to over 60% and increasing with larger number of groups (see Table D.1). Coverage rates remained stable, while empirical standard deviations showed a small increase under the alternative prior specification.