

Linear Algebra

lonelyneutrino

December 2024

1 Vector Spaces

1.1 \mathbb{R}^n and \mathbb{C}^n

Definition: Complex Numbers

A complex number is an ordered pair of the form (a, b) denoted as $a + bi$. $i = \sqrt{-1}$

$$\mathbb{C} = \{(a, b) : a, b \in \mathbb{R}\}$$

$(+, \cdot)$ are defined on complex numbers as follows-

$$(a + bi) + (c + di) = (a + c) + (b + d)i$$

$$(a + bi)(c + di) = (ac - bd) + (ad + bc)i$$

Any real number $a \in \mathbb{R}$ can be defined as a complex number of the form $a + 0i$. Hence, it is clear that $\mathbb{R} \subset \mathbb{C}$.

The operators of complex numbers are well-defined.

For $\lambda, \alpha, \beta \in \mathbb{C}$,

commutativity

$$\alpha + \beta = \beta + \alpha$$

associativity

$$(\alpha + \beta) + \lambda = \alpha + (\beta + \lambda)$$

identities

$$\lambda + 0 = \lambda \text{ and } \lambda * (1) = \lambda$$

additive inverse

For every $\alpha \in \mathbb{C}$, there exists a unique $\beta : \alpha + \beta = 0$

multiplicative inverse

For every $\alpha \in \mathbb{C}$, there exists a unique $\beta : \alpha \cdot \beta = 1$

distributivity

$$\lambda(\alpha + \beta) = \lambda \cdot \alpha + \lambda \cdot \beta$$

Example: Proof of commutativity

To show that $\alpha\beta = \beta\alpha$ for all $\alpha, \beta \in \mathbb{C}$ Let $\alpha = a + bi$ and $\beta = c + di$, $a, b, c, d \in \mathbb{R}$

$$\alpha\beta = ac - bd + (ad + bc)i$$

$$\beta\alpha = ca - db + (da + cb)i$$

We know that multiplication is commutative over real numbers.

$$\therefore \alpha\beta = ca - db + (da + cb)i = ac - bd + (ad + bc)i = \beta\alpha$$

Hence, the commutativity of complex number multiplication is proven.

Additive and multiplicative inverses of complex numbers can be defined. By doing so, we can define subtraction and division.

additive inverse

For every $\alpha \in \mathbb{C}$, there exists an additive inverse $\beta : \alpha + \beta = 0$

It can be proven that $\beta = -\alpha$

$$\alpha + (-\alpha) = (1 - 1)\alpha = 0 \cdot \alpha = 0$$

subtraction

Hence, the subtraction of two complex numbers α and β is defined as the addition of α and the additive inverse of β .

$$\alpha - \beta = \alpha + (-\beta)$$

multiplicative inverse

For every $\alpha \neq 0 \in \mathbb{C}$, there exists a unique multiplicative inverse $\beta : \alpha\beta = 1$

It can be proven that $\beta = \frac{1}{\alpha}$.

$$\alpha\beta = \alpha\frac{1}{\alpha} = 1$$

division

Hence, the division of two complex numbers α and β is defined as the multiplication of α and the multiplicative inverse of β .

Definition: Lists

A list (also called an **n-tuple**) is a collection of objects in a particular order. Two lists are equal if and only if they have the same size and the same elements in the same order. Lists are commonly written in the following manner-

$$\mathbf{v} = (v_1, v_2, v_3 \cdots v_n)$$