

Drug Side Effects Summary

Drug and Disease Interactions with Side Effects

1. This particular disease, Acne, interacts with the drug doxycycline and has the following side effects

- hives, difficult breathing, swelling in your face or throat) or a severe skin reaction (fever, sore throat, burning in your eyes, skin pain, red or purple skin rash that spreads and causes blistering and peeling). Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash, fever, swollen glands, flu-like symptoms, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. This reaction may occur several weeks after you began using doxycycline. Doxycycline may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- throat irritation, trouble swallowing
- chest pain, irregular heart rhythm, feeling short of breath
- little or no urination
- low white blood cell counts - fever, chills, swollen glands, body aches, weakness, pale skin, easy bruising or bleeding
- severe headaches, ringing in your ears, dizziness, nausea, vision problems, pain behind your eyes
- loss of appetite, upper stomach pain (that may spread to your back), tiredness, nausea or vomiting, fast heart rate, dark urine, jaundice (yellowing of the skin or eyes). Common side effects of doxycycline may include: nausea and vomiting
- upset stomach
- loss of appetite
- mild diarrhea
- skin rash or itching
- darkened skin color
- vaginal itching or discharge.

Drug Side Effects Summary

2. This particular disease, Acne, interacts with the drug spironolactone and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- little or no urination
- high potassium level - nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- or signs of other electrolyte imbalances - increased thirst or urination, confusion, vomiting , muscle pain, slurred speech, severe weakness, numbness , loss of coordination, feeling unsteady. Common spironolactone side effects may include: breast swelling or tenderness.

3. This particular disease, Acne, interacts with the drug minocycline and has the following side effects:

- skin rash, fever, swollen glands, flu-like symptoms, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. This may be more likely with long-term use of minocycline, and the reaction may occur several weeks after you began using this medicine. Call your doctor at once if you have: little or no urination, swelling in your feet or ankles, feeling tired or short of breath (signs of kidney problems)
- loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting, loss of appetite, easy bruising or bleeding, dark urine, yellowing of the skin or eyes (signs of liver or pancreas problems)
- joint pain or swelling with fever, swollen glands, muscle aches, chest pain, vomiting, unusual thoughts or behavior, and patchy skin color
- severe headaches, ringing in your ears, dizziness, vision problems, pain behind your eyes
- or swollen glands, flu symptoms, easy bruising or bleeding, severe tingling or numbness, muscle

Drug Side Effects Summary

weakness, chest pain, new or worsening cough with fever, trouble breathing. Common minocycline side effects may include: numbness , tingling, burning pain

- hair loss
- discoloration of you skin or nails. dizziness, spinning sensation
- muscle or joint pain
- nausea, diarrhea, loss of appetite
- swollen tongue, cough, trouble swallowing
- rash, itching
- or headache .

4. This particular disease, Acne, interacts with the drug Accutane and has the following side effects

- problems with your vision or hearing
- muscle or joint pain, bone pain, back pain
- increased thirst, increased urination
- hallucinations, (see or hearing things that are not real)
- symptoms of depression - unusual mood changes, crying spells, feelings of low self-worth, loss of interest in things you once enjoyed, new sleep problems, thoughts about hurting yourself
- signs of liver or pancreas problems - loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting, fast heart rate , dark urine, jaundice (yellowing of the skin or eyes)
- severe stomach problems - severe stomach or chest pain, pain when swallowing, heartburn , diarrhea , rectal bleeding , bloody or tarry stools
- or increased pressure inside the skull - severe headaches, ringing in your ears, dizziness, nausea, vision problems, pain behind your eyes. Common Accutane side effects may include: dryness of your skin, lips, eyes, or nose (you may have nosebleeds)
- vision problems

Drug Side Effects Summary

- headache, back pain , joint pain, muscle problems
- skin reactions
- or cold symptoms such as stuffy nose, sneezing, sore throat.

5. This particular disease, Acne, interacts with the drug clindamycin and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Clindamycin topical may cause serious side effects. Stop using clindamycin topical and call your doctor at once if you have: severe redness, itching, or dryness of treated skin areas
- or severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose). Common side effects of clindamycin topical may include: burning, itching, dryness, peeling or redness of treated skin
- or oily skin.

6. This particular disease, Acne, interacts with the drug Aldactone and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- little or no urination
- high potassium level - nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- or signs of other electrolyte imbalances - increased thirst or urination, confusion, vomiting , muscle pain, slurred speech, severe weakness, numbness , loss of coordination, feeling unsteady. Common

Drug Side Effects Summary

Aldactone side effects may include: breast swelling or tenderness.

7. This particular disease, Acne, interacts with the drug tretinoin and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tretinoin topical may cause serious side effects. Call your doctor at once if you have: severe burning, stinging, or irritation of treated skin
- severe skin dryness
- or severe redness, swelling, blistering, peeling, or crusting. Your skin may be more sensitive to weather extremes such as cold and wind while using tretinoin topical. Common side effects of tretinoin topical may include: skin pain, redness, burning, itching, or irritation
- sore throat
- mild warmth or stinging where the medicine was applied
- or changes in color of treated skin.

8. This particular disease, Acne, interacts with the drug isotretinoin and has the following side effects:

- problems with your vision or hearing
- muscle or joint pain, bone pain, back pain
- increased thirst, increased urination
- hallucinations, (see or hearing things that are not real)
- symptoms of depression--unusual mood changes, crying spells, feelings of low self-worth, loss of interest in things you once enjoyed, new sleep problems, thoughts about hurting yourself
- signs of liver or pancreas problems--loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting, fast heart rate , dark urine, jaundice (yellowing of the skin or eyes)
- severe stomach problems--severe stomach or chest pain, pain when swallowing, heartburn ,

Drug Side Effects Summary

diarrhea , rectal bleeding , bloody or tarry stools

- or increased pressure inside the skull--severe headaches, ringing in your ears, dizziness, nausea, vision problems, pain behind your eyes. Common side effects of isotretinoin may include: dryness of your skin, lips, eyes, or nose (you may have nosebleeds)
- vision problems
- headache, back pain , joint pain, muscle problems
- skin reactions
- or cold symptoms such as stuffy nose, sneezing, sore throat.

9. This particular disease, Acne, interacts with the drug Bactrim and has the following side effects:

- skin rash, fever, swollen glands, joint pain, muscle aches, severe weakness, pale skin, unusual bruising, or yellowing of your skin or eyes. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- any skin rash, no matter how mild
- yellowing of your skin or eyes
- a seizure
- new or unusual joint pain
- increased or decreased urination
- swelling, bruising, or irritation around the IV needle
- increased thirst, dry mouth, fruity breath odor
- new or worsening cough, fever, trouble breathing
- high blood potassium - nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- low blood sodium - headache , confusion, problems with thinking or memory, weakness, feeling unsteady

Drug Side Effects Summary

- or low blood cell counts - fever, chills, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath. Common Bactrim side effects may include: nausea, vomiting, loss of appetite
- or skin rash.

10. This particular disease, Acne, interacts with the drug Retin-A and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Retin-A and call your doctor at once if you have: severe burning, stinging, or irritation of treated skin
- severe redness, swelling, blistering, peeling, or crusting
- Your skin may be more sensitive to weather extremes such as cold and wind while using this medicine. Common Retin-A side effects may include: mild warmth or stinging where the medicine was applied
- or changes in color of treated skin.

11. This particular disease, Acne, interacts with the drug Aczone and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Aczone may cause methemoglobinemia (met-HEEM-oh glo-bin-EE-mee-a), a serious condition in which the amount of oxygen in your blood stream becomes dangerously low. Stop using this medicine and call your doctor at once if you notice a blue or gray appearance of your lips, fingernails, or the inside of your mouth. Call your doctor at once if you have: severe sore throat
- unusual thoughts or behavior

Drug Side Effects Summary

- a red blood cell disorder - back pain, pale or yellowed skin, dark colored urine, fever, confusion or weakness
- pancreas problems - severe pain in your upper stomach spreading to your back, nausea and vomiting, fast heart rate
- or severe skin reaction - fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common Aczone side effects may include: dry or peeling skin
- oily skin
- or redness where the medicine was applied.

12. This particular disease, Acne, interacts with the drug benzoyl peroxide and has the following side effects:

- Benzoyl peroxide topical can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Stop using benzoyl peroxide and call your doctor at once if you have any of these side effects on the treated skin: severe itching or burning
- severe stinging or redness
- swelling
- or peeling. Common side effects of benzoyl peroxide topical may include: mild stinging or burning
- itching or tingly feeling
- skin dryness, peeling, or flaking
- or redness or other irritation.

Drug Side Effects Summary

13. This particular disease, Acne, interacts with the drug Differin and has the following side effects:

- hives, itching
- chest pain, difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: skin redness, stinging, burning, or peeling after sun exposure. Some side effects may occur during the first 4 weeks of using this medicine. These side effects should become less noticeable with continued use. Common Differin side effects may include: dry, red, or scaly skin
- feeling of warmth and mild stinging or burning
- peeling skin
- or skin pain.

14. This particular disease, Acne, interacts with the drug Epiduo and has the following side effects:

- Benzoyl peroxide can cause a rare but serious allergic reaction or severe skin irritation. Stop using this medicine and get emergency medical help if you have: hives , itching
- difficult breathing, feeling light-headed
- or swelling of your face, lips, tongue, or throat. Epiduo may cause serious side effects. Call your doctor at once if you have: severe redness or swelling of treated skin
- severe burning or stinging
- or severe dryness or other skin irritation. Common Epiduo side effects may include: dry or scaly skin
- skin redness
- or burning or stinging.

15. This particular disease, Acne, interacts with the drug adapalene and has the following side effects:

- hives , itching

Drug Side Effects Summary

- chest pain, difficult breathing
- swelling of your face, lips, tongue, or throat. adapalene topical may cause serious side effects. Stop using adapalene topical and call your doctor at once if you have: skin redness, stinging, burning, or peeling after sun exposure. Some side effects may occur during the first 4 weeks of using this medicine. These side effects should become less noticeable with continued use. Common side effects of adapalene topical may include: dry, red, or scaly skin
- feeling of warmth and mild stinging or burning
- peeling skin
- or skin pain.

16. This particular disease, Acne, interacts with the drug cephalexin and has the following side effects

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- unusual tiredness, feeling light-headed or short of breath
- easy bruising, unusual bleeding, purple or red spots under your skin
- a seizure
- pale skin, cold hands and feet
- yellowed skin, dark colored urine
- fever, weakness
- or pain in your side or lower back, painful urination. Common cephalexin side effects may include: diarrhea
- nausea, vomiting
- indigestion , stomach pain
- or vaginal itching or discharge.

Drug Side Effects Summary

17. This particular disease, Acne, interacts with the drug Doryx and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blistering, peeling, or loosening of the skin bloating chills clay-colored stools constipation cough dark urine decreased appetite diarrhea diarrhea, watery and severe, which may also be bloody difficulty with swallowing dizziness fast heartbeat feeling of discomfort fever headache hives, itching, puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue hives or welts, itching, or rash increased thirst indigestion inflammation of the joints joint or muscle pain large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs loss of appetite nausea numbness or tingling of the face, hands, or feet pain in the stomach, side, or abdomen, possibly radiating to the back red skin lesions, often with a purple center redness and soreness of the eyes redness of the skin sore throat sores, ulcers, or white spots in the mouth or on the lips stomach cramps stomach pain or tenderness swelling of the feet or lower legs swollen, painful, or tender lymph glands in the neck, armpit, or groin tightness in the chest unusual tiredness or weakness unusual weight loss vomiting yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Back, leg, or stomach pains black, tarry stools bleeding gums blood in the urine or stools blurred vision bulging soft spot on the head of an infant change in the ability to see colors, especially blue or yellow chest pain, discomfort, or burning cracks in the skin decrease in vision difficulty breathing discoloration of the thyroid glands double vision general body swelling heartburn increased sensitivity of the skin to sunlight loss of heat from the body lower back or side pain nosebleeds pain

Drug Side Effects Summary

or burning in the throat pain with swallowing painful or difficult urination pale skin pinpoint red spots on the skin rash with flat lesions or small raised lesions on the skin red, swollen skin redness or other discoloration of the skin redness, swelling, or soreness of the tongue scaly skin severe nausea severe stomach pain severe sunburn tooth discoloration unusual bleeding or bruising vomiting blood Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Doryx side effects

18. This particular disease, Acne, interacts with the drug tetracycline and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe blistering, peeling, and red skin rash
- fever, chills, body aches, flu symptoms
- pale or yellowed skin, easy bruising or bleeding
- any signs of a new infection. Common tetracycline side effects may include: nausea, vomiting, diarrhea, upset stomach, loss of appetite
- white patches or sores inside your mouth or on your lips
- swollen tongue, black or "hairy" tongue, trouble swallowing
- sores or swelling in your rectal or genital area
- or vaginal itching or discharge.

19. This particular disease, Acne, interacts with the drug Septra and has the following side effects:

- skin rash, fever, swollen glands, joint pain, muscle aches, severe weakness, pale skin, unusual bruising, or yellowing of your skin or eyes. Call your doctor at once if you have: severe stomach

Drug Side Effects Summary

pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)

- a skin rash, no matter how mild
- yellowing of your skin or eyes
- a seizure
- new or unusual joint pain
- increased or decreased urination
- swelling, bruising, or irritation around the IV needle
- increased thirst, dry mouth, fruity breath odor
- new or worsening cough, fever, trouble breathing
- high potassium level - nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- low sodium level - headache, confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady
- or low blood cell counts - fever, chills, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath. Common Septra side effects may include: nausea , vomiting, loss of appetite
- or skin rash.

20. This particular disease, Acne, interacts with the drug Solodyn and has the following side effects

- skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Call your doctor at once if you have: bronchospasm (wheezing, chest tightness, trouble breathing)
- a seizure
- kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath

Drug Side Effects Summary

- low blood cell counts - fever, chills, tiredness, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath
- increased pressure inside the skull - severe headaches, ringing in your ears, dizziness, vision problems, pain behind your eyes
- or signs of liver or pancreas problems - loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting, loss of appetite, easy bruising or bleeding, dark urine, jaundice (yellowing of the skin or eyes). Common Solodyn side effects may include: numbness , tingling, burning pain
- hair loss
- discoloration of you skin or nails. dizziness, spinning sensation
- muscle or joint pain
- nausea, diarrhea, loss of appetite
- swollen tongue, cough, trouble swallowing
- rash, itching
- or headache .

21. This particular disease, Acne, interacts with the drug Tazorac and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Tazorac and call your doctor at once if you have: severe skin irritation (pain, dryness, flaking, peeling, burning, stinging, itching) after applying the medicine
- blistering or peeling skin
- severe redness or discomfort
- or swelling, warmth, oozing, or other signs of skin infection . Common Tazorac side effects may

Drug Side Effects Summary

include: dry, scaly, or peeling skin

- skin pain, redness or burning
- or itching or other irritation of treated skin.

22. This particular disease, Acne, interacts with the drug Vibramycin and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blistering, peeling, or loosening of the skin bloating chills clay-colored stools constipation cough dark urine decreased appetite diarrhea diarrhea, watery and severe, which may also be bloody difficulty with swallowing dizziness fast heartbeat feeling of discomfort fever headache hives, itching, puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue hives or welts, itching, or rash increased thirst indigestion inflammation of the joints joint or muscle pain large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs loss of appetite nausea numbness or tingling of the face, hands, or feet pain in the stomach, side, or abdomen, possibly radiating to the back red skin lesions, often with a purple center redness and soreness of the eyes redness of the skin sore throat sores, ulcers, or white spots in the mouth or on the lips stomach cramps stomach pain or tenderness swelling of the feet or lower legs swollen, painful, or tender lymph glands in the neck, armpit, or groin tightness in the chest unusual tiredness or weakness unusual weight loss vomiting yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Back, leg, or stomach pains black, tarry stools bleeding gums blood in the urine or stools blurred vision

Drug Side Effects Summary

bulging soft spot on the head of an infant change in the ability to see colors, especially blue or yellow chest pain, discomfort, or burning cracks in the skin decrease in vision difficulty breathing discoloration of the thyroid glands double vision general body swelling heartburn increased sensitivity of the skin to sunlight loss of heat from the body lower back or side pain nosebleeds pain or burning in the throat pain with swallowing painful or difficult urination pale skin pinpoint red spots on the skin rash with flat lesions or small raised lesions on the skin red, swollen skin redness or other discoloration of the skin redness, swelling, or soreness of the tongue scaly skin severe nausea severe stomach pain severe sunburn tooth discoloration unusual bleeding or bruising vomiting blood Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Vibramycin side effects

23. This particular disease, Acne, interacts with the drug Acticlate and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling

Drug Side Effects Summary

tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of a pancreas problem (pancreatitis) like very bad stomach pain, very bad back pain , or very bad upset stomach or throwing up. Chest pain or pressure or a fast heartbeat . Not able to pass urine or change in how much urine is passed. Fever, chills, or sore throat

- any unexplained bruising or bleeding

- or feeling very tired or weak. Throat irritation. Trouble swallowing. Muscle or joint pain. Fast breathing. Flushing. Very bad dizziness or passing out. Change in skin color. Swollen gland. Vaginal itching or discharge. Diarrhea is common with antibiotics . Rarely, a severe form called C diff-associated diarrhea (CDAD) may happen. Sometimes, this has led to a deadly bowel problem (colitis). CDAD may happen during or a few months after taking antibiotics. Call your doctor right away if you have stomach pain, cramps, or very loose, watery, or bloody stools. Check with your doctor before treating diarrhea. Raised pressure in the brain has happened with Acticlate (doxycycline tablets and capsules). Most of the time, this will go back to normal after Acticlate (doxycycline tablets and capsules) is stopped. Sometimes, loss of eyesight may happen and may not go away even after Acticlate (doxycycline tablets and capsules) is stopped. Call your doctor right away if you have a headache or eyesight problems like blurred eyesight, seeing double, or loss of eyesight. Acticlate side effects

24. This particular disease, Acne, interacts with the drug benzoyl peroxide / clindamycin and has the

- Benzoyl peroxide can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using benzoyl peroxide and clindamycin topical and get emergency medical help if you have signs of an allergic reaction : hives, itching

- difficult breathing, feeling light-headed

- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop

Drug Side Effects Summary

using this medicine and call your doctor at once if you have: severe redness, burning, stinging, or peeling of treated skin areas

- or diarrhea that is watery or bloody. Common side effects of benzoyl peroxide and clindamycin topical may include: mild burning or stinging
- itching or tingly feeling
- dryness or peeling of treated skin
- or redness or other irritation.

25. This particular disease, Acne, interacts with the drug Doxy 100 and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of a pancreas problem (pancreatitis) like very bad stomach pain, very bad back pain , or very bad upset stomach or throwing up. Chest pain or pressure or a fast heartbeat . Not able to pass urine or change in how much urine is passed. Fever, chills, or sore throat

Drug Side Effects Summary

- any unexplained bruising or bleeding
- or feeling very tired or weak. Throat irritation. Trouble swallowing. Muscle or joint pain. Fast breathing. Flushing. Very bad dizziness or passing out. Change in skin color. Swollen gland. Vaginal itching or discharge. Diarrhea is common with antibiotics . Rarely, a severe form called C diff-associated diarrhea (CDAD) may happen. Sometimes, this has led to a deadly bowel problem (colitis). CDAD may happen during or a few months after taking antibiotics. Call your doctor right away if you have stomach pain, cramps, or very loose, watery, or bloody stools. Check with your doctor before treating diarrhea. Raised pressure in the brain has happened with Doxy 100 (doxycycline injection). Most of the time, this will go back to normal after Doxy 100 (doxycycline injection) is stopped. Sometimes, loss of eyesight may happen and may not go away even after Doxy 100 (doxycycline injection) is stopped. Call your doctor right away if you have a headache or eyesight problems like blurred eyesight, seeing double, or loss of eyesight. Doxy 100 side effects

26. This particular disease, Acne, interacts with the drug Keflex and has the following side effects:

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- unusual tiredness, feeling light-headed or short of breath
- easy bruising, unusual bleeding, purple or red spots under your skin
- a seizure
- pale skin, cold hands and feet
- yellowed skin, dark colored urine
- fever, weakness
- or pain in your side or lower back, painful urination. Common Keflex side effects may include:
diarrhea
- nausea, vomiting

Drug Side Effects Summary

- indigestion , stomach pain
- or vaginal itching or discharge.

27. This particular disease, Acne, interacts with the drug sulfamethoxazole / trimethoprim and has t

- skin rash, fever, swollen glands, joint pain, muscle aches, severe weakness, pale skin, unusual bruising, or yellowing of your skin or eyes. Sulfamethoxazole and trimethoprim may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- any skin rash, no matter how mild
- yellowing of your skin or eyes
- a seizure
- new or unusual joint pain
- increased or decreased urination
- swelling, bruising, or irritation around the IV needle
- increased thirst, dry mouth, fruity breath odor
- new or worsening cough, fever, trouble breathing
- high blood potassium-- nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- low blood sodium-- headache , confusion, problems with thinking or memory, weakness, feeling unsteady
- or low blood cell counts--fever, chills, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath. Common side effects of sulfamethoxazole and trimethoprim may include: nausea, vomiting, loss of appetite
- or skin rash.

Drug Side Effects Summary

28. This particular disease, Acne, interacts with the drug Benzacilin and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have any of these serious side effects: severe redness, burning, stinging, or peeling of treated skin areas
- or diarrhea that is watery or bloody. Less serious Benzacilin side effects may include: mild burning or stinging
- itching or tingly feeling
- dryness or peeling of treated skin
- or redness or other irritation.

29. This particular disease, Acne, interacts with the drug Monodox and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blistering, peeling, or loosening of the skin bloating chills clay-colored stools constipation cough dark urine decreased appetite diarrhea diarrhea, watery and severe, which may also be bloody difficulty with swallowing dizziness fast heartbeat feeling of discomfort fever headache hives, itching, puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue hives or welts, itching, or rash increased thirst indigestion inflammation of the joints joint or muscle pain large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs loss of appetite nausea numbness or tingling of the face, hands, or feet pain in the stomach, side, or abdomen, possibly radiating to the back red skin lesions, often with a purple center redness and soreness of the eyes redness of the skin sore throat sores, ulcers, or white spots in the mouth or on the lips stomach cramps stomach

Drug Side Effects Summary

pain or tenderness swelling of the feet or lower legs swollen, painful, or tender lymph glands in the neck, armpit, or groin tightness in the chest unusual tiredness or weakness unusual weight loss vomiting yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Back, leg, or stomach pains black, tarry stools bleeding gums blood in the urine or stools blurred vision bulging soft spot on the head of an infant change in the ability to see colors, especially blue or yellow chest pain, discomfort, or burning cracks in the skin decrease in vision difficulty breathing discoloration of the thyroid glands double vision general body swelling heartburn increased sensitivity of the skin to sunlight loss of heat from the body lower back or side pain nosebleeds pain or burning in the throat pain with swallowing painful or difficult urination pale skin pinpoint red spots on the skin rash with flat lesions or small raised lesions on the skin red, swollen skin redness or other discoloration of the skin redness, swelling, or soreness of the tongue scaly skin severe nausea severe stomach pain severe sunburn tooth discoloration unusual bleeding or bruising vomiting blood Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Monodox side effects

30. This particular disease, Acne, interacts with the drug Targadox and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

Drug Side Effects Summary

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of a pancreas problem (pancreatitis) like very bad stomach pain, very bad back pain , or very bad upset stomach or throwing up. Chest pain or pressure or a fast heartbeat . Not able to pass urine or change in how much urine is passed. Fever, chills, or sore throat
- any unexplained bruising or bleeding
- or feeling very tired or weak. Throat irritation. Trouble swallowing. Muscle or joint pain. Fast breathing. Flushing. Very bad dizziness or passing out. Change in skin color. Swollen gland. Vaginal itching or discharge. Diarrhea is common with antibiotics . Rarely, a severe form called C diff-associated diarrhea (CDAD) may happen. Sometimes, this has led to a deadly bowel problem (colitis). CDAD may happen during or a few months after taking antibiotics. Call your doctor right away if you have stomach pain, cramps, or very loose, watery, or bloody stools. Check with your doctor before treating diarrhea. Raised pressure in the brain has happened with TargaDOX (doxycycline tablets and capsules). Most of the time, this will go back to normal after TargaDOX (doxycycline tablets and capsules) is stopped. Sometimes, loss of eyesight may happen and may not go away even after TargaDOX (doxycycline tablets and capsules) is stopped. Call your doctor right away if you have a headache or eyesight problems like blurred eyesight, seeing double, or loss of eyesight. Targadox side effects

Drug Side Effects Summary

31. This particular disease, Acne, interacts with the drug Adoxa and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

- itching

- red, swollen, blistered, or peeling skin with or without fever

- wheezing

- tightness in the chest or throat

- trouble breathing, swallowing, or talking

- unusual hoarseness

- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of a pancreas problem (pancreatitis) like very bad stomach pain, very bad back pain , or very bad upset stomach or throwing up. Chest pain or pressure or a fast heartbeat . Not able to pass urine or change in how much urine is passed. Fever, chills, or sore throat

- any unexplained bruising or bleeding

- or feeling very tired or weak. Throat irritation. Trouble swallowing. Muscle or joint pain. Fast breathing. Flushing. Very bad dizziness or passing out. Change in skin color. Swollen gland. Vaginal itching or discharge. Diarrhea is common with antibiotics . Rarely, a severe form called C diff-associated diarrhea (CDAD) may happen. Sometimes, this has led to a deadly bowel problem (colitis). CDAD may happen during or a few months after taking antibiotics. Call your doctor right away if you have stomach pain, cramps, or very loose, watery, or bloody stools. Check with your doctor before treating diarrhea. Raised pressure in the brain has happened with Adoxa (doxycycline

Drug Side Effects Summary

tablets and capsules). Most of the time, this will go back to normal after Adoxa (doxycycline tablets and capsules) is stopped. Sometimes, loss of eyesight may happen and may not go away even after Adoxa (doxycycline tablets and capsules) is stopped. Call your doctor right away if you have a headache or eyesight problems like blurred eyesight, seeing double, or loss of eyesight. Adoxa side effects

32. This particular disease, Acne, interacts with the drug Adoxa CK and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blistering, peeling, or loosening of the skin bloating chills clay-colored stools constipation cough dark urine decreased appetite diarrhea diarrhea, watery and severe, which may also be bloody difficulty with swallowing dizziness fast heartbeat feeling of discomfort fever headache hives, itching, puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue hives or welts, itching, or rash increased thirst indigestion inflammation of the joints joint or muscle pain large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs loss of appetite nausea numbness or tingling of the face, hands, or feet pain in the stomach, side, or abdomen, possibly radiating to the back red skin lesions, often with a purple center redness and soreness of the eyes redness of the skin sore throat sores, ulcers, or white spots in the mouth or on the lips stomach cramps stomach pain or tenderness swelling of the feet or lower legs swollen, painful, or tender lymph glands in the neck, armpit, or groin tightness in the chest unusual tiredness or weakness unusual weight loss vomiting yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects

Drug Side Effects Summary

continue or are bothersome or if you have any questions about them: Incidence not known Back, leg, or stomach pains black, tarry stools bleeding gums blood in the urine or stools blurred vision bulging soft spot on the head of an infant change in the ability to see colors, especially blue or yellow chest pain, discomfort, or burning cracks in the skin decrease in vision difficulty breathing discoloration of the thyroid glands double vision general body swelling heartburn increased sensitivity of the skin to sunlight loss of heat from the body lower back or side pain nosebleeds pain or burning in the throat pain with swallowing painful or difficult urination pale skin pinpoint red spots on the skin rash with flat lesions or small raised lesions on the skin red, swollen skin redness or other discoloration of the skin redness, swelling, or soreness of the tongue scaly skin severe nausea severe stomach pain severe sunburn tooth discoloration unusual bleeding or bruising vomiting blood Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Adoxa CK side effects

33. This particular disease, Acne, interacts with the drug Adoxa Pak and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blistering, peeling, or loosening of the skin bloating chills clay-colored stools constipation cough dark urine decreased appetite diarrhea diarrhea, watery and severe, which may also be bloody difficulty with swallowing dizziness fast heartbeat feeling of discomfort fever headache hives, itching, puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue hives or welts, itching, or rash increased thirst indigestion inflammation of the joints joint or muscle pain large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs loss of appetite nausea numbness or tingling of the face, hands, or feet pain in the stomach, side, or abdomen, possibly radiating to the

Drug Side Effects Summary

back red skin lesions, often with a purple center redness and soreness of the eyes redness of the skin sore throat sores, ulcers, or white spots in the mouth or on the lips stomach cramps stomach pain or tenderness swelling of the feet or lower legs swollen, painful, or tender lymph glands in the neck, armpit, or groin tightness in the chest unusual tiredness or weakness unusual weight loss vomiting yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Back, leg, or stomach pains black, tarry stools bleeding gums blood in the urine or stools blurred vision bulging soft spot on the head of an infant change in the ability to see colors, especially blue or yellow chest pain, discomfort, or burning cracks in the skin decrease in vision difficulty breathing discoloration of the thyroid glands double vision general body swelling heartburn increased sensitivity of the skin to sunlight loss of heat from the body lower back or side pain nosebleeds pain or burning in the throat pain with swallowing painful or difficult urination pale skin pinpoint red spots on the skin rash with flat lesions or small raised lesions on the skin red, swollen skin redness or other discoloration of the skin redness, swelling, or soreness of the tongue scaly skin severe nausea severe stomach pain severe sunburn tooth discoloration unusual bleeding or bruising vomiting blood Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Adoxa Pak side effects

34. This particular disease, Acne, interacts with the drug Adoxa TT and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your

Drug Side Effects Summary

doctor immediately if any of the following side effects occur: Incidence not known Blistering, peeling, or loosening of the skin bloating chills clay-colored stools constipation cough dark urine decreased appetite diarrhea diarrhea, watery and severe, which may also be bloody difficulty with swallowing dizziness fast heartbeat feeling of discomfort fever headache hives, itching, puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue hives or welts, itching, or rash increased thirst indigestion inflammation of the joints joint or muscle pain large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs loss of appetite nausea numbness or tingling of the face, hands, or feet pain in the stomach, side, or abdomen, possibly radiating to the back red skin lesions, often with a purple center redness and soreness of the eyes redness of the skin sore throat sores, ulcers, or white spots in the mouth or on the lips stomach cramps stomach pain or tenderness swelling of the feet or lower legs swollen, painful, or tender lymph glands in the neck, armpit, or groin tightness in the chest unusual tiredness or weakness unusual weight loss vomiting yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Back, leg, or stomach pains black, tarry stools bleeding gums blood in the urine or stools blurred vision bulging soft spot on the head of an infant change in the ability to see colors, especially blue or yellow chest pain, discomfort, or burning cracks in the skin decrease in vision difficulty breathing discoloration of the thyroid glands double vision general body swelling heartburn increased sensitivity of the skin to sunlight loss of heat from the body lower back or side pain nosebleeds pain or burning in the throat pain with swallowing painful or difficult urination pale skin pinpoint red spots on the skin rash with flat lesions or small raised lesions on the skin red, swollen skin redness or other discoloration of the skin redness, swelling, or soreness of the tongue scaly skin severe nausea

Drug Side Effects Summary

severe stomach pain severe sunburn tooth discoloration unusual bleeding or bruising vomiting blood
Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Adoxa TT side effects

35. This particular disease, Acne, interacts with the drug Avidoxy and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

- itching

- red, swollen, blistered, or peeling skin with or without fever

- wheezing

- tightness in the chest or throat

- trouble breathing, swallowing, or talking

- unusual hoarseness

- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of a pancreas problem (pancreatitis) like very bad stomach pain, very bad back pain , or very bad upset stomach or throwing up. Chest pain or pressure or a fast heartbeat . Not able to pass urine or change in how much urine is passed. Fever, chills, or sore throat

- any unexplained bruising or bleeding

- or feeling very tired or weak. Throat irritation. Trouble swallowing. Muscle or joint pain. Fast breathing. Flushing. Very bad dizziness or passing out. Change in skin color. Swollen gland. Vaginal

Drug Side Effects Summary

itching or discharge. Diarrhea is common with antibiotics . Rarely, a severe form called C diff-associated diarrhea (CDAD) may happen. Sometimes, this has led to a deadly bowel problem (colitis). CDAD may happen during or a few months after taking antibiotics. Call your doctor right away if you have stomach pain, cramps, or very loose, watery, or bloody stools. Check with your doctor before treating diarrhea. Raised pressure in the brain has happened with Avidoxy (doxycycline tablets and capsules). Most of the time, this will go back to normal after Avidoxy (doxycycline tablets and capsules) is stopped. Sometimes, loss of eyesight may happen and may not go away even after Avidoxy (doxycycline tablets and capsules) is stopped. Call your doctor right away if you have a headache or eyesight problems like blurred eyesight, seeing double, or loss of eyesight. Avidoxy side effects

36. This particular disease, Acne, interacts with the drug Bactrim DS and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Black, tarry stools blistering, peeling, or loosening of the skin changes in skin color chest pain or tightness chills clay-colored stools cough or hoarseness dark urine diarrhea dizziness fever general feeling of tiredness or weakness headache itching, skin rash joint or muscle pain light-colored stools loss of appetite lower back or side pain nausea pain, tenderness, or swelling of the foot or leg painful or difficult urination pale skin red skin lesions, often with a purple center red, irritated eyes sore throat sores, ulcers, or white spots in the mouth or on the lips stomach pain swollen or painful glands trouble breathing unpleasant breath odor unusual bleeding or bruising vomiting of blood yellow eyes or skin Incidence not known Back, leg, or stomach pains bleeding gums blindness or vision changes bloating blood in the urine or stools bluish-colored lips, fingernails, or palms burning, crawling, itching, numbness, painful, prickling, "pins and needles", or tingling feelings burning of the face or

Drug Side Effects Summary

mouth cloudy urine clumsiness or unsteadiness confusion constipation continuing ringing or buzzing or other unexplained noise in the ears cracks in the skin decreased frequency or amount of urine difficulty with swallowing fainting spells fast, pounding, or irregular heartbeat or pulse general body swelling general feeling of discomfort or illness hair loss hearing loss hives increased thirst indigestion large, flat, blue, or purplish patches in the skin large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs loss of heat from the body muscle or joint pain nosebleeds not able to pass urine numbness or tingling in the hands, feet, or lips pain or burning while urinating pinpoint red spots on the skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue raised red swellings on the skin, the buttocks, legs, or ankles redness of the white part of the eyes redness, swelling, or soreness of the tongue seizures soreness of the muscles stiff neck or back stomach tenderness swelling of the face, hands, legs, and feet unsteadiness, trembling, or other problems with muscle control or coordination weakness in the hands or feet weakness or heaviness of the legs weight gain or loss

Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them:

More common

Passing of gas

Incidence not known

Discouragement feeling of constant movement of self or surroundings feeling sad or empty increased sensitivity of the skin to sunlight lack of feeling or emotion loss of interest or pleasure muscle pain, stiffness, cramps, or spasms nervousness redness or other discoloration of the skin seeing, hearing, or feeling things that are not there sensation of spinning severe sunburn trouble concentrating trouble sleeping uncaring weight loss

Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Bactrim DS side effects

Drug Side Effects Summary

37. This particular disease, Acne, interacts with the drug Cleocin T and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Cleocin T may cause serious side effects. Stop using Cleocin T and call your doctor at once if you have: severe redness, itching, or dryness of treated skin areas
- or severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose). Common side effects of Cleocin T may include: burning, itching, dryness, peeling or redness of treated skin
- or oily skin.

38. This particular disease, Acne, interacts with the drug Clindagel and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Clindagel may cause serious side effects. Stop using Clindagel and call your doctor at once if you have: severe redness, itching, or dryness of treated skin areas
- or severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose). Common side effects of Clindagel may include: burning, itching, dryness, peeling or redness of treated skin
- or oily skin.

39. This particular disease, Acne, interacts with the drug Doryx MPC and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your

Drug Side Effects Summary

doctor immediately if any of the following side effects occur: Incidence not known Blistering, peeling, or loosening of the skin bloating chills clay-colored stools constipation cough dark urine decreased appetite diarrhea diarrhea, watery and severe, which may also be bloody difficulty with swallowing dizziness fast heartbeat feeling of discomfort fever headache hives, itching, puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue hives or welts, itching, or rash increased thirst indigestion inflammation of the joints joint or muscle pain large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs loss of appetite nausea numbness or tingling of the face, hands, or feet pain in the stomach, side, or abdomen, possibly radiating to the back red skin lesions, often with a purple center redness and soreness of the eyes redness of the skin sore throat sores, ulcers, or white spots in the mouth or on the lips stomach cramps stomach pain or tenderness swelling of the feet or lower legs swollen, painful, or tender lymph glands in the neck, armpit, or groin tightness in the chest unusual tiredness or weakness unusual weight loss vomiting yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Back, leg, or stomach pains black, tarry stools bleeding gums blood in the urine or stools blurred vision bulging soft spot on the head of an infant change in the ability to see colors, especially blue or yellow chest pain, discomfort, or burning cracks in the skin decrease in vision difficulty breathing discoloration of the thyroid glands double vision general body swelling heartburn increased sensitivity of the skin to sunlight loss of heat from the body lower back or side pain nosebleeds pain or burning in the throat pain with swallowing painful or difficult urination pale skin pinpoint red spots on the skin rash with flat lesions or small raised lesions on the skin red, swollen skin redness or other discoloration of the skin redness, swelling, or soreness of the tongue scaly skin severe nausea

Drug Side Effects Summary

severe stomach pain severe sunburn tooth discoloration unusual bleeding or bruising vomiting blood
Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Doryx MPC side effects

40. This particular disease, Acne, interacts with the drug erythromycin and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Erythromycin topical may cause serious side effects. Call your doctor at once if you have: severe burning, stinging, or redness
- oozing or other signs of skin infection
- worsening of your skin condition
- or severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose). Common side effects of erythromycin topical may include: skin itching, redness, burning, or peeling
- or dry or oily skin.

41. This particular disease, Acne, interacts with the drug Morgidox and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever

Drug Side Effects Summary

- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of a pancreas problem (pancreatitis) like very bad stomach pain, very bad back pain , or very bad upset stomach or throwing up. Chest pain or pressure or a fast heartbeat . Not able to pass urine or change in how much urine is passed. Fever, chills, or sore throat
- any unexplained bruising or bleeding
- or feeling very tired or weak. Throat irritation. Trouble swallowing. Muscle or joint pain. Fast breathing. Flushing. Very bad dizziness or passing out. Change in skin color. Swollen gland. Vaginal itching or discharge. Diarrhea is common with antibiotics . Rarely, a severe form called C diff-associated diarrhea (CDAD) may happen. Sometimes, this has led to a deadly bowel problem (colitis). CDAD may happen during or a few months after taking antibiotics. Call your doctor right away if you have stomach pain, cramps, or very loose, watery, or bloody stools. Check with your doctor before treating diarrhea. Raised pressure in the brain has happened with Morgidox (doxycycline tablets and capsules). Most of the time, this will go back to normal after Morgidox (doxycycline tablets and capsules) is stopped. Sometimes, loss of eyesight may happen and may not go away even after Morgidox (doxycycline tablets and capsules) is stopped. Call your doctor right away if you have a headache or eyesight problems like blurred eyesight, seeing double, or loss of eyesight. Morgidox side effects

42. This particular disease, Acne, interacts with the drug Oraxyl and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and

Drug Side Effects Summary

sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of a pancreas problem (pancreatitis) like very bad stomach pain, very bad back pain , or very bad upset stomach or throwing up. Chest pain or pressure or a fast heartbeat . Not able to pass urine or change in how much urine is passed. Fever, chills, or sore throat
- any unexplained bruising or bleeding
- or feeling very tired or weak. Throat irritation. Trouble swallowing. Muscle or joint pain. Fast breathing. Flushing. Very bad dizziness or passing out. Change in skin color. Swollen gland. Vaginal itching or discharge. Diarrhea is common with antibiotics . Rarely, a severe form called C diff-associated diarrhea (CDAD) may happen. Sometimes, this has led to a deadly bowel problem (colitis). CDAD may happen during or a few months after taking antibiotics. Call your doctor right away if you have stomach pain, cramps, or very loose, watery, or bloody stools. Check with your doctor before treating diarrhea. Raised pressure in the brain has happened with Oraxyl (doxycycline tablets and capsules). Most of the time, this will go back to normal after Oraxyl (doxycycline tablets and capsules) is stopped. Sometimes, loss of eyesight may happen and may not go away even after

Drug Side Effects Summary

Oraxyl (doxycycline tablets and capsules) is stopped. Call your doctor right away if you have a headache or eyesight problems like blurred eyesight, seeing double, or loss of eyesight. Oraxyl side effects

43. This particular disease, Acne, interacts with the drug Claravis and has the following side effects

- problems with your vision or hearing
- muscle or joint pain, bone pain, back pain
- increased thirst, increased urination
- hallucinations, (see or hearing things that are not real)
- symptoms of depression--unusual mood changes, crying spells, feelings of low self-worth, loss of interest in things you once enjoyed, new sleep problems, thoughts about hurting yourself
- signs of liver or pancreas problems--loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting, fast heart rate , dark urine, jaundice (yellowing of the skin or eyes)
- severe stomach problems--severe stomach or chest pain, pain when swallowing, heartburn , diarrhea , rectal bleeding , bloody or tarry stools
- or increased pressure inside the skull--severe headaches, ringing in your ears, dizziness, nausea, vision problems, pain behind your eyes. Common side effects of Claravis may include: dryness of your skin, lips, eyes, or nose (you may have nosebleeds)
- vision problems
- headache, back pain , joint pain, muscle problems
- skin reactions
- or cold symptoms such as stuffy nose, sneezing, sore throat.

44. This particular disease, Acne, interacts with the drug Clinda-Derm and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of

Drug Side Effects Summary

these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Abdominal or stomach cramps, pain, and bloating (severe) diarrhea (watery and severe), which may also be bloody fever increased thirst nausea or vomiting unusual tiredness or weakness weight loss (unusual)--these side effects may also occur up to several weeks after you stop using this medicine Check with your doctor as soon as possible if any of the following side effects occur: Less common Skin rash, itching, redness, swelling, or other sign of irritation not present before use of this medicine Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Dryness, scaliness, or peeling of skin (for the topical solution) Less common Abdominal pain diarrhea (mild) headache irritation or oiliness of skin stinging or burning feeling of skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Clinda-Derm side effects

45. This particular disease, Acne, interacts with the drug Evoclin and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Serious side effects are unlikely when clindamycin is applied to the skin, but can occur if the medicine is absorbed into your bloodstream. Stop using Evoclin and call your doctor at once if you have: severe redness, itching, or dryness of treated skin areas
- or severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last

Drug Side Effects Summary

dose). Common Evoclin side effects may include: burning, itching, dryness, peeling or redness of treated skin

- or oily skin.

46. This particular disease, Acne, interacts with the drug Minocin and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Black, tarry stools blistering, peeling, or loosening of the skin blood in the urine or stools blurred or double vision bulging soft spot on the head of an infant chest pain, possibly moving to the left arm, neck, or shoulder confusion diarrhea dizziness or lightheadedness eye pain fast heartbeat general feeling of discomfort or illness general tiredness and weakness hives, itching, or skin rash joint or muscle pain large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs loss of appetite nausea or vomiting red skin lesions, often with a purple center severe headache severe stomach pain sores, ulcers, or white spots on the lips or in the mouth troubled breathing unusual bleeding or bruising upper right abdominal or stomach pain yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Continuing ringing or buzzing or other unexplained noise in the ears difficulty with moving hearing loss hives or welts muscle stiffness redness of the skin sleepiness or unusual drowsiness Incidence not known Bloating discoloration of the tooth increased sensitivity of the skin to sunlight indigestion severe sunburn Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call

Drug Side Effects Summary

your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Minocin side effects

47. This particular disease, Acne, interacts with the drug Yaz and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Yaz and call your doctor at once if you have: sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance
- pain, warmth or swelling in one or both legs
- shortness of breath, rapid breathing, and/or sudden coughing
- coughing up blood
- sudden vision loss or blurred vision
- chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- symptoms of liver problems - loss of appetite, upper stomach pain, tiredness, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- signs of increased blood pressure - severe headache, blurred vision, pounding in your neck or ears
- swelling in your hands, ankles, or feet
- a change in the pattern or severity of migraine headaches
- or symptoms of depression - sleep problems, weakness, tired feeling, mood changes. Common Yaz side effects may include: nausea, vomiting
- breast tenderness
- headache, mood changes, feeling tired or irritable
- weight gain
- skin darkening or pigmentation changes

Drug Side Effects Summary

- changes in your menstrual periods or decreased sex drive.

48. This particular disease, Acne, interacts with the drug adapalene / benzoyl peroxide and has the following side effects:

- Benzoyl peroxide can cause a rare but serious allergic reaction or severe skin irritation. Stop using this medicine and get emergency medical help if you have: hives , itching
- difficult breathing, feeling light-headed
- or swelling of your face, lips, tongue, or throat. Adapalene and benzoyl peroxide may cause serious side effects. Call your doctor at once if you have: severe redness or swelling of treated skin
- severe burning or stinging
- or severe dryness or other skin irritation. Common side effects of adapalene and benzoyl peroxide may include: dry or scaly skin
- skin redness
- or burning or stinging.

49. This particular disease, Acne, interacts with the drug Clindacin ETZ and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Clindacin ETZ may cause serious side effects. Stop using Clindacin ETZ and call your doctor at once if you have: severe redness, itching, or dryness of treated skin areas
- or severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose). Common side effects of Clindacin ETZ may include: burning, itching, dryness, peeling or redness of treated skin
- or oily skin.

Drug Side Effects Summary

50. This particular disease, Acne, interacts with the drug Clindacin P and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Clindacin P may cause serious side effects. Stop using Clindacin P and call your doctor at once if you have: severe redness, itching, or dryness of treated skin areas
- or severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose). Common side effects of Clindacin P may include: burning, itching, dryness, peeling or redness of treated skin
- or oily skin.

51. This particular disease, Acne, interacts with the drug Clindacin PAC and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Abdominal or stomach cramps, pain, and bloating (severe) diarrhea (watery and severe), which may also be bloody fever increased thirst nausea or vomiting unusual tiredness or weakness weight loss (unusual)--these side effects may also occur up to several weeks after you stop using this medicine Check with your doctor as soon as possible if any of the following side effects occur: Less common Skin rash, itching, redness, swelling, or other sign of irritation not present before use of this medicine Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Dryness, scaliness, or peeling of skin (for the topical solution)

Drug Side Effects Summary

Less common Abdominal pain diarrhea (mild) headache irritation or oiliness of skin stinging or burning feeling of skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Clindacin PAC side effects

52. This particular disease, Acne, interacts with the drug ClindaReach Pledget and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Abdominal or stomach cramps, pain, and bloating (severe) diarrhea (watery and severe), which may also be bloody fever increased thirst nausea or vomiting unusual tiredness or weakness weight loss (unusual)--these side effects may also occur up to several weeks after you stop using this medicine Check with your doctor as soon as possible if any of the following side effects occur: Less common Skin rash, itching, redness, swelling, or other sign of irritation not present before use of this medicine Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Dryness, scaliness, or peeling of skin (for the topical solution) Less common Abdominal pain diarrhea (mild) headache irritation or oiliness of skin stinging or burning feeling of skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. ClindaReach Pledget side effects

Drug Side Effects Summary

53. This particular disease, Acne, interacts with the drug Clindets and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Abdominal or stomach cramps, pain, and bloating (severe) diarrhea (watery and severe), which may also be bloody fever increased thirst nausea or vomiting unusual tiredness or weakness weight loss (unusual)--these side effects may also occur up to several weeks after you stop using this medicine Check with your doctor as soon as possible if any of the following side effects occur: Less common Skin rash, itching, redness, swelling, or other sign of irritation not present before use of this medicine Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Dryness, scaliness, or peeling of skin (for the topical solution) Less common Abdominal pain diarrhea (mild) headache irritation or oiliness of skin stinging or burning feeling of skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Clindets side effects

54. This particular disease, Acne, interacts with the drug Milk of Magnesia and has the following side effects

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Milk of Magnesia may cause serious side effects. Stop using Milk of Magnesia and call your doctor at once if you have: severe nausea , vomiting , or diarrhea

Drug Side Effects Summary

- no bowel movement after using the medicine as a laxative
- rectal bleeding
- or worsening symptoms. Common side effects of Milk of Magnesia may include: diarrhea
- or a decreased sense of taste.

55. This particular disease, Acne, interacts with the drug Septra DS and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Black, tarry stools blistering, peeling, or loosening of the skin changes in skin color chest pain or tightness chills clay-colored stools cough or hoarseness dark urine diarrhea dizziness fever general feeling of tiredness or weakness headache itching, skin rash joint or muscle pain light-colored stools loss of appetite lower back or side pain nausea pain, tenderness, or swelling of the foot or leg painful or difficult urination pale skin red skin lesions, often with a purple center red, irritated eyes sore throat sores, ulcers, or white spots in the mouth or on the lips stomach pain swollen or painful glands trouble breathing unpleasant breath odor unusual bleeding or bruising vomiting of blood yellow eyes or skin Incidence not known Back, leg, or stomach pains bleeding gums blindness or vision changes bloating blood in the urine or stools bluish-colored lips, fingernails, or palms burning, crawling, itching, numbness, painful, prickling, "pins and needles", or tingling feelings burning of the face or mouth cloudy urine clumsiness or unsteadiness confusion constipation continuing ringing or buzzing or other unexplained noise in the ears cracks in the skin decreased frequency or amount of urine difficulty with swallowing fainting spells fast, pounding, or irregular heartbeat or pulse general body swelling general feeling of discomfort or illness hair loss hearing loss hives increased thirst indigestion large, flat, blue, or purplish patches in the skin large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs loss of heat from the body muscle or

Drug Side Effects Summary

joint pain nosebleeds not able to pass urine numbness or tingling in the hands, feet, or lips pain or burning while urinating pinpoint red spots on the skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue raised red swellings on the skin, the buttocks, legs, or ankles redness of the white part of the eyes redness, swelling, or soreness of the tongue seizures soreness of the muscles stiff neck or back stomach tenderness swelling of the face, hands, legs, and feet unsteadiness, trembling, or other problems with muscle control or coordination weakness in the hands or feet weakness or heaviness of the legs weight gain or loss Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Passing of gas Incidence not known Discouragement feeling of constant movement of self or surroundings feeling sad or empty increased sensitivity of the skin to sunlight lack of feeling or emotion loss of interest or pleasure muscle pain, stiffness, cramps, or spasms nervousness redness or other discoloration of the skin seeing, hearing, or feeling things that are not there sensation of spinning severe sunburn trouble concentrating trouble sleeping uncaring weight loss Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Septra DS side effects

56. This particular disease, Acne, interacts with the drug Yasmin and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Yasmin and call your doctor at once if you have: signs of a stroke - sudden numbness or weakness (especially on one side of the body),

Drug Side Effects Summary

sudden severe headache, slurred speech, problems with vision or balance

- signs of a blood clot - sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood, pain or warmth and swelling in one or both legs

- heart attack symptoms - chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating

- liver problems - loss of appetite, upper stomach pain, tiredness, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)

- increased blood pressure - severe headache, blurred vision, pounding in your neck or ears

- swelling in your hands, ankles, or feet

- a change in the pattern or severity of migraine headaches

- or symptoms of depression - sleep problems, weakness, tired feeling, mood changes. Common

Yasmin side effects may include: nausea, vomiting

- breast tenderness

- headache, mood changes, feeling tired or irritable

- weight gain

- or changes in your menstrual periods, decreased sex drive.

57. This particular disease, Acne, interacts with the drug Ziana and has the following side effects:

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: severe redness, burning, stinging, or scaling of treated skin areas

- or diarrhea that is watery or bloody. Common Ziana side effects may include: mild skin irritation, dryness or peeling

- mild skin redness

Drug Side Effects Summary

- or itching of treated skin.

58. This particular disease, Acne, interacts with the drug Absorica and has the following side effects

- problems with your vision or hearing
- muscle or joint pain, bone pain, back pain
- increased thirst, increased urination
- hallucinations, (see or hearing things that are not real)
- symptoms of depression--unusual mood changes, crying spells, feelings of low self-worth, loss of interest in things you once enjoyed, new sleep problems, thoughts about hurting yourself
- signs of liver or pancreas problems--loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting, fast heart rate , dark urine, jaundice (yellowing of the skin or eyes)
- severe stomach problems--severe stomach or chest pain, pain when swallowing, heartburn , diarrhea , rectal bleeding , bloody or tarry stools
- or increased pressure inside the skull--severe headaches, ringing in your ears, dizziness, nausea, vision problems, pain behind your eyes. Common side effects of Absorica may include: dryness of your skin, lips, eyes, or nose (you may have nosebleeds)
- vision problems
- headache, back pain , joint pain, muscle problems
- skin reactions
- or cold symptoms such as stuffy nose, sneezing, sore throat.

59. This particular disease, Acne, interacts with the drug Acanya and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using Acanya and call your doctor at once if you

Drug Side Effects Summary

have any of these serious side effects: severe redness, burning, stinging, or peeling of treated skin areas

- or diarrhea that is watery or bloody. Less serious Acanya side effects may include: mild burning or stinging
- itching or tingly feeling
- dryness or peeling of treated skin
- or redness or other irritation.

60. This particular disease, Acne, interacts with the drug Amnesteem and has the following side effects

- problems with your vision or hearing
- muscle or joint pain, bone pain, back pain
- increased thirst, increased urination
- hallucinations, (see or hearing things that are not real)
- symptoms of depression--unusual mood changes, crying spells, feelings of low self-worth, loss of interest in things you once enjoyed, new sleep problems, thoughts about hurting yourself
- signs of liver or pancreas problems--loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting, fast heart rate , dark urine, jaundice (yellowing of the skin or eyes)
- severe stomach problems--severe stomach or chest pain, pain when swallowing, heartburn , diarrhea , rectal bleeding , bloody or tarry stools
- or increased pressure inside the skull--severe headaches, ringing in your ears, dizziness, nausea, vision problems, pain behind your eyes. Common side effects of Amnesteem may include: dryness of your skin, lips, eyes, or nose (you may have nosebleeds)
- vision problems
- headache, back pain , joint pain, muscle problems
- skin reactions

Drug Side Effects Summary

- or cold symptoms such as stuffy nose, sneezing, sore throat.

61. This particular disease, Acne, interacts with the drug azelaic acid and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe burning, stinging, or warmth
- severe itching or tingling
- severe redness, dryness, peeling, or other irritation
- changes in skin color
- or new or worsening breathing problems. Common side effects of azelaic acid topical may include: pain, burning, stinging, or tingling
- dry or scaly skin
- or itching or other irritation.

62. This particular disease, Acne, interacts with the drug Co-trimoxazole and has the following side effects:

- No side effects listed

63. This particular disease, Acne, interacts with the drug dapsone and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Dapsone topical may cause methemoglobinemia (met-HEEM-oh glo-bin-EE-mee-a), a serious condition in which the amount of oxygen in your blood stream becomes dangerously low. Stop using this medicine and call your doctor at once if you notice a blue or gray appearance of your lips, fingernails, or the inside of your mouth. Dapsone

Drug Side Effects Summary

topical may cause serious side effects. Call your doctor at once if you have: severe sore throat

- unusual thoughts or behavior
- a red blood cell disorder-- back pain , pale or yellowed skin, dark colored urine, fever, confusion or weakness
- pancreas problems--severe pain in your upper stomach spreading to your back, nausea and vomiting , fast heart rate
- or severe skin reaction--fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of dapstone topical may include: dry or peeling skin
- oily skin
- or redness where the medicine was applied.

64. This particular disease, Acne, interacts with the drug Duac and has the following side effects:

- Benzoyl peroxide can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using DUAC and get emergency medical help if you have signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe redness, burning, stinging, or peeling of treated skin areas
- or diarrhea that is watery or bloody. Common side effects of DUAC may include: mild burning or stinging
- itching or tingly feeling

Drug Side Effects Summary

- dryness or peeling of treated skin
- or redness or other irritation.

65. This particular disease, Acne, interacts with the drug Emcin Clear and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Emcin Clear may cause serious side effects. Call your doctor at once if you have: severe burning, stinging, or redness
- oozing or other signs of skin infection
- worsening of your skin condition
- or severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose). Common side effects of Emcin Clear may include: skin itching, redness, burning, or peeling
- or dry or oily skin.

66. This particular disease, Acne, interacts with the drug Epiduo Forte and has the following side effects:

- Benzoyl peroxide can cause a rare but serious allergic reaction or severe skin irritation. Stop using this medicine and get emergency medical help if you have: hives , itching
- difficult breathing, feeling light-headed
- or swelling of your face, lips, tongue, or throat. Epiduo Forte may cause serious side effects. Call your doctor at once if you have: severe redness or swelling of treated skin
- severe burning or stinging
- or severe dryness or other skin irritation. Common side effects of Epiduo Forte may include: dry or scaly skin
- skin redness
- or burning or stinging.

Drug Side Effects Summary

67. This particular disease, Acne, interacts with the drug Erygel and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Erygel may cause serious side effects. Call your doctor at once if you have: severe burning, stinging, or redness
- oozing or other signs of skin infection
- worsening of your skin condition
- or severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose). Common side effects of Erygel may include: skin itching, redness, burning, or peeling
- or dry or oily skin.

68. This particular disease, Acne, interacts with the drug Ery Pads and has the following side effects:

- No side effects listed

69. This particular disease, Acne, interacts with the drug Myorisan and has the following side effects:

- problems with your vision or hearing
- muscle or joint pain, bone pain, back pain
- increased thirst, increased urination
- hallucinations, (see or hearing things that are not real)
- symptoms of depression--unusual mood changes, crying spells, feelings of low self-worth, loss of interest in things you once enjoyed, new sleep problems, thoughts about hurting yourself
- signs of liver or pancreas problems--loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting, fast heart rate , dark urine, jaundice (yellowing of the skin or eyes)
- severe stomach problems--severe stomach or chest pain, pain when swallowing, heartburn , diarrhea , rectal bleeding , bloody or tarry stools

Drug Side Effects Summary

- or increased pressure inside the skull--severe headaches, ringing in your ears, dizziness, nausea, vision problems, pain behind your eyes. Common side effects of Myorisan may include: dryness of your skin, lips, eyes, or nose (you may have nosebleeds)
- vision problems
- headache, back pain , joint pain, muscle problems
- skin reactions
- or cold symptoms such as stuffy nose, sneezing, sore throat.

70. This particular disease, Acne, interacts with the drug Onexton and has the following side effects:

- Benzoyl peroxide can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using Onexton and get emergency medical help if you have signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe redness, burning, stinging, or peeling of treated skin areas
- or diarrhea that is watery or bloody. Common side effects of Onexton may include: mild burning or stinging
- itching or tingly feeling
- dryness or peeling of treated skin
- or redness or other irritation.

71. This particular disease, Acne, interacts with the drug Phillips' Milk of Magnesia and has the following side effects:

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Phillips Milk of Magnesia may cause serious side effects. Stop using Phillips Milk of Magnesia and call your doctor at once if you have: severe nausea , vomiting , or diarrhea
- no bowel movement after using the medicine as a laxative
- rectal bleeding
- or worsening symptoms. Common side effects of Phillips Milk of Magnesia may include: diarrhea
- or a decreased sense of taste.

72. This particular disease, Acne, interacts with the drug salicylic acid and has the following side effects

- Salicylic acid topical can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Also stop using salicylic acid topical and call your doctor at once if you have: severe headache , ringing in your ears, problems with hearing, thinking problems
- severe stomach pain, vomiting , or diarrhea
- a light-headed feeling, like you might pass out
- shortness of breath
- or severe burning, dryness, or irritation of the skin. Common side effects of salicylic acid topical may include: minor skin irritation , rash, or peeling
- or changes in the color of treated skin (usually whitening).

Drug Side Effects Summary

73. This particular disease, Acne, interacts with the drug Spotex and has the following side effects:

- No side effects listed

74. This particular disease, Acne, interacts with the drug Theramycin Z and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Theramycin Z may cause serious side effects. Call your doctor at once if you have: severe burning, stinging, or redness
- oozing or other signs of skin infection
- worsening of your skin condition
- or severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose). Common side effects of Theramycin Z may include: skin itching, redness, burning, or peeling
- or dry or oily skin.

75. This particular disease, Acne, interacts with the drug Absorica LD and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking

Drug Side Effects Summary

- unusual hoarseness

- or swelling of the mouth, face, lips, tongue, or throat. Signs or symptoms of depression , suicidal thoughts, emotional ups and downs, abnormal thinking, anxiety , or lack of interest in life. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of esophagus problems like chest pain, trouble swallowing, or new or worse heartburn . Signs of high blood sugar like confusion, feeling sleepy, more thirst, more hungry, passing urine more often, flushing, fast breathing, or breath that smells like fruit. Behavior problems. Eye pain . Back, bone, joint, or muscle pain. Muscle weakness. Any unexplained bruising or bleeding. Feeling very tired or weak. Not able to pass urine or change in how much urine is passed. Swelling. Swollen gland. Fast or abnormal heartbeat. Hearing problems like change in hearing or ringing in the ears may happen. This may go away after stopping the drug but sometimes it may not. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)

- red or irritated eyes

- or sores in your mouth, throat, nose, or eyes. Very bad and sometimes deadly pancreas problems (pancreatitis) have happened with Absorica LD (ISOTretinoin (systemic)). Call your doctor right away if you have very bad stomach pain, very bad back pain , or very upset stomach or throwing up.

Absorica LD side effects

76. This particular disease, Acne, interacts with the drug Achromycin V and has the following side effects

- No side effects listed

77. This particular disease, Acne, interacts with the drug Acne Treatment and has the following side effects

Drug Side Effects Summary

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Acne Treatment side effects

78. This particular disease, Acne, interacts with the drug Acnex and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common or rare Skin irritation not present before use of this medicine (moderate or severe) Incidence not known Difficult breathing dryness and peeling of skin fainting hives or itching redness of the skin swelling of the eyes, face, lips, or tongue tightness in the throat unusually warm skin Symptoms of overdose Confusion diarrhea dizziness fast or deep breathing headache (severe or continuing) hearing loss lightheadedness nausea rapid breathing ringing or buzzing in the ears (continuing) severe drowsiness stomach pain vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of

Drug Side Effects Summary

these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Skin irritation not present before use of this medicine (mild) stinging Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Acnex side effects

79. This particular disease, Acne, interacts with the drug Ala-Tet and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Ala-Tet may cause serious side effects. Call your doctor at once if you have: severe blistering, peeling, and red skin rash
- fever, chills, body aches, flu symptoms
- pale or yellowed skin, easy bruising or bleeding
- any signs of a new infection. Common side effects of Ala-Tet may include: nausea , vomiting , diarrhea, upset stomach , loss of appetite
- white patches or sores inside your mouth or on your lips
- swollen tongue, black or "hairy" tongue, trouble swallowing
- sores or swelling in your rectal or genital area
- or vaginal itching or discharge.

80. This particular disease, Acne, interacts with the drug Amzeeq and has the following side effects:

- skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Serious side effects are unlikely when Amzeeq is applied to the skin, but can occur if the medicine is absorbed into your bloodstream. Amzeeq may cause serious side effects.

Drug Side Effects Summary

Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)

- severe dizziness
 - confusion, headache , vision problems, pain behind your eyes
 - fever, swollen glands, itching, joint pain, or not feeling well
 - discoloration of your skin, nails, eyes, mouth, gums, or teeth. liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
 - or lupus -like symptoms--muscle or joint pain, flu symptoms, chest pain, and a rash or patchy skin color that worsens in sunlight. Minocycline can affect growth in children, especially if used long term.
- Common side effects of Amzeeq may include: diarrhea
- or headache.

81. This particular disease, Acne, interacts with the drug Atralin and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Atralin and call your doctor at once if you have: severe burning, stinging, or irritation of treated skin
- severe redness, swelling, blistering, peeling, or crusting
- Your skin may be more sensitive to weather extremes such as cold and wind while using this medicine. Common Atralin side effects may include: mild warmth or stinging where the medicine was applied
- or changes in color of treated skin.

82. This particular disease, Acne, interacts with the drug Avita and has the following side effects:

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Avita may cause serious side effects. Call your doctor at once if you have: severe burning, stinging, or irritation of treated skin
- severe skin dryness
- or severe redness, swelling, blistering, peeling, or crusting. Your skin may be more sensitive to weather extremes such as cold and wind while using Avita. Common side effects of Avita may include: skin pain, redness, burning, itching, or irritation
- sore throat
- mild warmth or stinging where the medicine was applied
- or changes in color of treated skin.

83. This particular disease, Acne, interacts with the drug Azelex and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe burning, stinging, or warmth
- severe itching or tingling
- severe redness, dryness, peeling, or other irritation
- changes in skin color
- or new or worsening breathing problems. Common side effects of Azelex may include: pain, burning, stinging, or tingling
- dry or scaly skin
- or itching or other irritation.

Drug Side Effects Summary

84. This particular disease, Acne, interacts with the drug Benzamycin and has the following side effects:

- Benzoyl peroxide can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have any of these signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe redness, burning, stinging, or peeling of treated skin areas
- or diarrhea that is watery or bloody. Common side effects of Benzamycin may include: mild burning or stinging
- itching or tingly feeling
- dryness or peeling of treated skin
- or redness or other irritation.

85. This particular disease, Acne, interacts with the drug benzoyl peroxide / erythromycin and has the following side effects:

- Benzoyl peroxide can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have any of these signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe redness, burning, stinging, or peeling of treated skin areas
- or diarrhea that is watery or bloody. Common side effects of benzoyl peroxide and erythromycin

Drug Side Effects Summary

topical may include: mild burning or stinging

- itching or tingly feeling
- dryness or peeling of treated skin
- or redness or other irritation.

86. This particular disease, Acne, interacts with the drug benzoyl peroxide / sulfur and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad skin irritation. Benzoyl peroxide / sulfur topical side effects

87. This particular disease, Acne, interacts with the drug Beyaz and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Beyaz and call your doctor at once if you have: signs of a stroke - sudden numbness or weakness (especially on one side of the body),

Drug Side Effects Summary

sudden severe headache, slurred speech, problems with vision or balance

- signs of a blood clot - sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood, pain or warmth in one or both legs

- heart attack symptoms - chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating

- liver problems - loss of appetite, upper stomach pain, tiredness, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)

- increased blood pressure - severe headache, blurred vision, pounding in your neck or ears

- swelling in your hands, ankles, or feet

- a change in the pattern or severity of migraine headaches

- or symptoms of depression - sleep problems, weakness, tired feeling, mood changes. Common

Beyaz side effects may include: nausea

- breast tenderness

- headache

- or breakthrough bleeding.

88. This particular disease, Acne, interacts with the drug Brodspec and has the following side effects

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Brodspec may cause serious side effects. Call your doctor at once if you have: severe blistering, peeling, and red skin rash

- fever, chills, body aches, flu symptoms

- pale or yellowed skin, easy bruising or bleeding

- any signs of a new infection. Common side effects of Brodspec may include: nausea , vomiting , diarrhea, upset stomach , loss of appetite

Drug Side Effects Summary

- white patches or sores inside your mouth or on your lips
- swollen tongue, black or "hairy" tongue, trouble swallowing
- sores or swelling in your rectal or genital area
- or vaginal itching or discharge.

89. This particular disease, Acne, interacts with the drug Cleeravue-M and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Redness, dryness, itching, peeling, or discoloration of the treated skin Less common Diarrhea Incidence not known Blurred vision change in ability to see colors, especially blue or yellow fever general feeling of discomfort or illness headache increased sensitivity of the skin to sunlight itching or skin rash joint or muscle pain loss of appetite redness or other discoloration of the skin severe abdominal or stomach cramps and pain severe sunburn unusual tiredness or weakness vomiting watery and severe diarrhea, which may also be bloody Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Burning, darkening, dryness, flushing, itching, peeling, redness, or stinging on the skin, especially when used on the face raised, dark red, wart-like spots on the skin, especially when used on the face Less common Headache Incidence not known Darkening of the skin, teeth, or gums dizziness or lightheadedness feeling of constant movement of self or surroundings sensation of spinning Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report

Drug Side Effects Summary

side effects to the FDA at 1-800-FDA-1088. Cleeravue-M side effects

90. This particular disease, Acne, interacts with the drug clindamycin / tretinoin and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe redness, burning, stinging, or scaling of treated skin areas
- or diarrhea that is watery or bloody. Common side effects of clindamycin and tretinoin may include: mild skin irritation, dryness or peeling
- mild skin redness
- or itching of treated skin.

91. This particular disease, Acne, interacts with the drug drospirenone / ethinyl estradiol and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Birth control pills may cause serious side effects. Stop using birth control pills and call your doctor at once if you have: signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance
- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood , pain or warmth in one or both legs
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- liver problems--loss of appetite, upper stomach pain, tiredness, dark urine, clay-colored stools,

Drug Side Effects Summary

jaundice (yellowing of the skin or eyes)

- increased blood pressure--severe headache, blurred vision, pounding in your neck or ears
- swelling in your hands, ankles, or feet
- a change in the pattern or severity of migraine headaches
- or symptoms of depression--sleep problems, weakness, tired feeling, mood changes. Common side effects of drospirenone and ethinyl estradiol may include: nausea, vomiting
- breast tenderness
- headache, mood changes, feeling tired or irritable
- weight gain
- or changes in your menstrual periods, decreased sex drive.

92. This particular disease, Acne, interacts with the drug Dynacin and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Black, tarry stools blistering, peeling, or loosening of the skin blood in the urine or stools blurred or double vision bulging soft spot on the head of an infant chest pain, possibly moving to the left arm, neck, or shoulder confusion diarrhea dizziness or lightheadedness eye pain fast heartbeat general feeling of discomfort or illness general tiredness and weakness hives, itching, or skin rash joint or muscle pain large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs loss of appetite nausea or vomiting red skin lesions, often with a purple center severe headache severe stomach pain sores, ulcers, or white spots on the lips or in the mouth troubled breathing unusual bleeding or bruising upper right abdominal or stomach pain yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be

Drug Side Effects Summary

able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Continuing ringing or buzzing or other unexplained noise in the ears difficulty with moving hearing loss hives or welts muscle stiffness redness of the skin sleepiness or unusual drowsiness Incidence not known Bloating discoloration of the tooth increased sensitivity of the skin to sunlight indigestion severe sunburn Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Dynacin side effects

93. This particular disease, Acne, interacts with the drug Estarylla and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Birth control pills may cause serious side effects. Stop using birth control pills and call your doctor at once if you have: signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache , slurred speech, problems with vision or balance
- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood , pain or warmth in one or both legs
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating
- liver problems--loss of appetite, upper stomach pain, tiredness, fever, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- severe headache, pounding in your neck or ears
- swelling in your hands, ankles, or feet

Drug Side Effects Summary

- changes in the pattern or severity of migraine headaches
- a breast lump
- or symptoms of depression--sleep problems, weakness, tired feeling, mood changes. Common side effects of Estarylla may include: stomach pain, gas, nausea, vomiting
- breast tenderness
- acne , darkening of facial skin
- headache, nervousness , mood changes
- problems with contact lenses
- changes in weight
- breakthrough bleeding
- vaginal itching or discharge
- or rash.

94. This particular disease, Acne, interacts with the drug ethinyl estradiol / norgestimate and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Birth control pills may cause serious side effects. Stop using birth control pills and call your doctor at once if you have: signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache , slurred speech, problems with vision or balance
- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood , pain or warmth in one or both legs
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating
- liver problems--loss of appetite, upper stomach pain, tiredness, fever, dark urine, clay-colored

Drug Side Effects Summary

stools, jaundice (yellowing of the skin or eyes)

- severe headache, pounding in your neck or ears
- swelling in your hands, ankles, or feet
- changes in the pattern or severity of migraine headaches
- a breast lump
- or symptoms of depression--sleep problems, weakness, tired feeling, mood changes. Common side effects of ethinyl estradiol and norgestimate may include: stomach pain, gas, nausea, vomiting
- breast tenderness
- acne , darkening of facial skin
- headache, nervousness , mood changes
- problems with contact lenses
- changes in weight
- breakthrough bleeding
- vaginal itching or discharge
- or rash.

95. This particular disease, Acne, interacts with the drug Gianvi and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Birth control pills may cause serious side effects. Stop using birth control pills and call your doctor at once if you have: signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance
- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood , pain or warmth in one or both legs

Drug Side Effects Summary

- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- liver problems--loss of appetite, upper stomach pain, tiredness, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- increased blood pressure--severe headache, blurred vision, pounding in your neck or ears
- swelling in your hands, ankles, or feet
- a change in the pattern or severity of migraine headaches
- or symptoms of depression--sleep problems, weakness, tired feeling, mood changes. Common side effects of Gianvi may include: nausea, vomiting
- breast tenderness
- headache, mood changes, feeling tired or irritable
- weight gain
- or changes in your menstrual periods, decreased sex drive.

96. This particular disease, Acne, interacts with the drug Loryna and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Birth control pills may cause serious side effects. Stop using birth control pills and call your doctor at once if you have: signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance
- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood , pain or warmth in one or both legs
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating

Drug Side Effects Summary

- liver problems--loss of appetite, upper stomach pain, tiredness, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- increased blood pressure--severe headache, blurred vision, pounding in your neck or ears
- swelling in your hands, ankles, or feet
- a change in the pattern or severity of migraine headaches
- or symptoms of depression--sleep problems, weakness, tired feeling, mood changes. Common side effects of Loryna may include: nausea, vomiting
- breast tenderness
- headache, mood changes, feeling tired or irritable
- weight gain
- or changes in your menstrual periods, decreased sex drive.

97. This particular disease, Acne, interacts with the drug magnesium hydroxide and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Magnesium hydroxide may cause serious side effects. Stop using magnesium hydroxide and call your doctor at once if you have: severe nausea , vomiting , or diarrhea
- no bowel movement after using the medicine as a laxative
- rectal bleeding
- or worsening symptoms. Common side effects of magnesium hydroxide may include: diarrhea
- or a decreased sense of taste.

98. This particular disease, Acne, interacts with the drug minocycline and has the following side effects:

- hives , itching, severe rash

Drug Side Effects Summary

- swollen glands, unusual tiredness, fever, nausea , stomach pain
- lower back pain , painful urination, blood or pus in your urine
- chest pain, cough with mucus, difficult breathing
- jaundice (yellowing of the skin or eyes)
- swelling of your face, lips, tongue, or throat. Some of these signs of allergic reaction have occurred when minocycline was taken by mouth in a pill form. It is not known whether Arestin minocycline powder would cause these same effects when used in dentistry. Call your dentist at once if you have: tooth pain
- pain or swelling in your gums
- fever, swollen glands, rash or itching, joint pain or swelling, muscle aches, general ill feeling
- or severe skin reaction--fever, sore throat , swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of minocycline may include: increased tooth sensitivity
- pain
- headache
- infection
- or flu-like symptoms.

99. This particular disease, Acne, interacts with the drug Minolira and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Black, tarry stools blistering, peeling, or loosening of the skin blood in the urine or stools blurred or double vision bulging soft spot on the head of an infant chest pain, possibly moving to the left arm, neck, or

Drug Side Effects Summary

shoulder confusion diarrhea dizziness or lightheadedness eye pain fast heartbeat general feeling of discomfort or illness general tiredness and weakness hives, itching, or skin rash joint or muscle pain large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs loss of appetite nausea or vomiting red skin lesions, often with a purple center severe headache severe stomach pain sores, ulcers, or white spots on the lips or in the mouth troubled breathing unusual bleeding or bruising upper right abdominal or stomach pain yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Continuing ringing or buzzing or other unexplained noise in the ears difficulty with moving hearing loss hives or welts muscle stiffness redness of the skin sleepiness or unusual drowsiness Incidence not known Bloating discoloration of the tooth increased sensitivity of the skin to sunlight indigestion severe sunburn Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Minolira side effects

100. This particular disease, Acne, interacts with the drug Ocella and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Birth control pills may cause serious side effects. Stop using birth control pills and call your doctor at once if you have: signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance

Drug Side Effects Summary

- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood , pain or warmth in one or both legs
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- liver problems--loss of appetite, upper stomach pain, tiredness, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- increased blood pressure--severe headache, blurred vision, pounding in your neck or ears
- swelling in your hands, ankles, or feet
- a change in the pattern or severity of migraine headaches
- or symptoms of depression--sleep problems, weakness, tired feeling, mood changes. Common side effects of Ocella may include: nausea, vomiting
- breast tenderness
- headache, mood changes, feeling tired or irritable
- weight gain
- or changes in your menstrual periods, decreased sex drive.

101. This particular disease, Acne, interacts with the drug PanOxyl and has the following side effects

- PanOxyl can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Stop using benzoyl peroxide and call your doctor at once if you have any of these side effects on the treated skin: severe itching or burning
- severe stinging or redness

Drug Side Effects Summary

- swelling
- or peeling. Common side effects of PanOxyl may include: mild stinging or burning
- itching or tingly feeling
- skin dryness, peeling, or flaking
- or redness or other irritation.

102. This particular disease, Acne, interacts with the drug PanOxyl 10% Acne Foaming Wash and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. PanOxyl 10% Acne Foaming Wash side effects

103. This particular disease, Acne, interacts with the drug Seysara and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: a light-headed

Drug Side Effects Summary

feeling, like you might pass out

- a spinning sensation
- or increased pressure inside the skull - severe headaches, ringing in your ears, vision problems, pain behind your eyes. Common Seysara side effects may include nausea .

104. This particular disease, Acne, interacts with the drug Sotret and has the following side effects:

- problems with your vision or hearing
- muscle or joint pain, bone pain, back pain
- increased thirst, increased urination
- hallucinations, (see or hearing things that are not real)
- symptoms of depression--unusual mood changes, crying spells, feelings of low self-worth, loss of interest in things you once enjoyed, new sleep problems, thoughts about hurting yourself
- signs of liver or pancreas problems--loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting, fast heart rate , dark urine, jaundice (yellowing of the skin or eyes)
- severe stomach problems--severe stomach or chest pain, pain when swallowing, heartburn , diarrhea , rectal bleeding , bloody or tarry stools
- or increased pressure inside the skull--severe headaches, ringing in your ears, dizziness, nausea, vision problems, pain behind your eyes. Common side effects of Sotret may include: dryness of your skin, lips, eyes, or nose (you may have nosebleeds)
- vision problems
- headache, back pain , joint pain, muscle problems
- skin reactions
- or cold symptoms such as stuffy nose, sneezing, sore throat.

105. This particular disease, Acne, interacts with the drug sulfacetamide sodium/sulfur and has the

Drug Side Effects Summary

- redness, warmth, swelling, itching, stinging, burning, or irritation of treated skin.

106. This particular disease, Acne, interacts with the drug tazarotene and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using tazarotene and call your doctor at once if you have: severe skin irritation (pain, dryness, flaking, peeling, burning, stinging, itching) after applying the medicine
- blistering or peeling skin
- severe redness or discomfort
- or swelling, warmth, oozing, or other signs of skin infection . Common tazarotene side effects may include: dry, scaly, or peeling skin
- skin pain, redness or burning
- or itching or other irritation of treated skin.

107. This particular disease, Acne, interacts with the drug Tri-Previfem and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Absent, missed, or irregular menstrual periods anxiety change in vision changes in skin color chest pain or discomfort chills clay-colored stools constipation cough dark urine diarrhea dizziness or lightheadedness fainting fever headache hives or welts itching skin large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs loss of appetite medium to heavy, irregular vaginal bleeding between regular monthly periods, which may require the use of a pad or a tampon nausea pain or discomfort in the arms, jaw, back, or neck pain, tenderness, or swelling of the foot or leg

Drug Side Effects Summary

pains in the chest, groin, or legs, especially in the calves of the legs pounding in the ears rash redness of the skin severe headaches of sudden onset slow or fast heartbeat stomach pain sudden loss of coordination or slurred speech sudden troubled breathing sweating unusual tiredness or weakness vomiting vomiting of blood Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Bloating blotchy spots on the exposed skin breast enlargement or tenderness discouragement feeling sad or empty irritability itching of the vagina or outside the genitals loss of interest or pleasure pain during sexual intercourse stomach cramps thick, white curd-like vaginal discharge without odor or with mild odor tiredness trouble concentrating trouble sleeping trouble wearing contact lenses Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Tri-Previfem side effects

108. This particular disease, Acne, interacts with the drug Tri-Sprintec and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Absent, missed, or irregular menstrual periods anxiety change in vision changes in skin color chest pain or discomfort chills clay-colored stools constipation cough dark urine diarrhea dizziness or lightheadedness fainting fever headache hives or welts itching skin large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs loss of appetite medium to heavy, irregular vaginal bleeding between regular monthly periods, which may require the use of a pad or a tampon nausea

Drug Side Effects Summary

pain or discomfort in the arms, jaw, back, or neck pain, tenderness, or swelling of the foot or leg
pains in the chest, groin, or legs, especially in the calves of the legs pounding in the ears rash
redness of the skin severe headaches of sudden onset slow or fast heartbeat stomach pain sudden
loss of coordination or slurred speech sudden troubled breathing sweating unusual tiredness or
weakness vomiting vomiting of blood Some side effects may occur that usually do not need medical
attention. These side effects may go away during treatment as your body adjusts to the medicine.
Also, your health care professional may be able to tell you about ways to prevent or reduce some of
these side effects. Check with your health care professional if any of the following side effects
continue or are bothersome or if you have any questions about them: Incidence not known Bloating
blotchy spots on the exposed skin breast enlargement or tenderness discouragement feeling sad or
empty irritability itching of the vagina or outside the genitals loss of interest or pleasure pain during
sexual intercourse stomach cramps thick, white curd-like vaginal discharge without odor or with mild
odor tiredness trouble concentrating trouble sleeping trouble wearing contact lenses Other side
effects not listed may also occur in some patients. If you notice any other effects, check with your
healthcare professional. Call your doctor for medical advice about side effects. You may report side
effects to the FDA at 1-800-FDA-1088. Tri-Sprintec side effects

109. This particular disease, Acne, interacts with the drug Veltin and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: severe redness, burning, stinging, or scaling of treated skin areas
- or diarrhea that is watery or bloody. Common Veltin side effects may include: mild skin irritation, dryness or peeling
- mild skin redness

Drug Side Effects Summary

- or itching of treated skin.

110. This particular disease, Acne, interacts with the drug Ximino and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Black, tarry stools blistering, peeling, or loosening of the skin blood in the urine or stools blurred or double vision bulging soft spot on the head of an infant chest pain, possibly moving to the left arm, neck, or shoulder confusion diarrhea dizziness or lightheadedness eye pain fast heartbeat general feeling of discomfort or illness general tiredness and weakness hives, itching, or skin rash joint or muscle pain large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs loss of appetite nausea or vomiting red skin lesions, often with a purple center severe headache severe stomach pain sores, ulcers, or white spots on the lips or in the mouth troubled breathing unusual bleeding or bruising upper right abdominal or stomach pain yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Continuing ringing or buzzing or other unexplained noise in the ears difficulty with moving hearing loss hives or welts muscle stiffness redness of the skin sleepiness or unusual drowsiness Incidence not known Bloating discoloration of the tooth increased sensitivity of the skin to sunlight indigestion severe sunburn Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Ximino side effects

Drug Side Effects Summary

111. This particular disease, Acne, interacts with the drug Zenatane and has the following side effects

- problems with your vision or hearing
- muscle or joint pain, bone pain, back pain
- increased thirst, increased urination
- hallucinations, (see or hearing things that are not real)
- symptoms of depression--unusual mood changes, crying spells, feelings of low self-worth, loss of interest in things you once enjoyed, new sleep problems, thoughts about hurting yourself
- signs of liver or pancreas problems--loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting, fast heart rate , dark urine, jaundice (yellowing of the skin or eyes)
- severe stomach problems--severe stomach or chest pain, pain when swallowing, heartburn , diarrhea , rectal bleeding , bloody or tarry stools
- or increased pressure inside the skull--severe headaches, ringing in your ears, dizziness, nausea, vision problems, pain behind your eyes. Common side effects of Zenatane may include: dryness of your skin, lips, eyes, or nose (you may have nosebleeds)
- vision problems
- headache, back pain , joint pain, muscle problems
- skin reactions
- or cold symptoms such as stuffy nose, sneezing, sore throat.

112. This particular disease, Acne, interacts with the drug Acetoxyl and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Pain, burning, blistering, changes in color, dryness, itching, peeling, redness, scaling, stinging, or swelling of the skin where the medicine is applied Incidence not known Chest tightness cough difficulty swallowing dizziness

Drug Side Effects Summary

fainting fast heartbeat hives, itching, or skin rash increased sensitivity of the skin to sunlight large, hive-like swelling on face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs redness or other discoloration of the skin severe sunburn swelling of the eyes, face, lips, or tongue tightness in the throat trouble breathing unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Dryness or peeling of the skin (may occur after a few days) feeling of warmth, mild stinging, and redness of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Acetoxyl side effects

113. This particular disease, Acne, interacts with the drug Acne-10 and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Pain, burning, blistering, changes in color, dryness, itching, peeling, redness, scaling, stinging, or swelling of the skin where the medicine is applied Incidence not known Chest tightness cough difficulty swallowing dizziness fainting fast heartbeat hives, itching, or skin rash increased sensitivity of the skin to sunlight large, hive-like swelling on face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs redness or other discoloration of the skin severe sunburn swelling of the eyes, face, lips, or tongue tightness in the throat trouble breathing unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways

Drug Side Effects Summary

to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Dryness or peeling of the skin (may occur after a few days) feeling of warmth, mild stinging, and redness of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Acne-10 side effects

114. This particular disease, Acne, interacts with the drug Acne-Clear and has the following side effects

- Acne-Clear can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Stop using benzoyl peroxide and call your doctor at once if you have any of these side effects on the treated skin: severe itching or burning
- severe stinging or redness
- swelling
- or peeling. Common side effects of Acne-Clear may include: mild stinging or burning
- itching or tingly feeling
- skin dryness, peeling, or flaking
- or redness or other irritation.

115. This particular disease, Acne, interacts with the drug Acnevir and has the following side effects

- Acnevir can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop

Drug Side Effects Summary

using this medicine and get emergency medical help if you have signs of an allergic reaction : hives, itching

- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Also stop using Acnevir and call your doctor at once if you have: severe headache , ringing in your ears, problems with hearing, thinking problems
- severe stomach pain, vomiting , or diarrhea
- a light-headed feeling, like you might pass out
- shortness of breath
- or severe burning, dryness, or irritation of the skin. Common side effects of Acnevir may include: minor skin irritation , rash, or peeling
- or changes in the color of treated skin (usually whitening).

116. This particular disease, Acne, interacts with the drug Acnomel and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: Less common or rare Skin irritation not present before use of this medicine Symptoms of resorcinol poisoning Diarrhea, nausea, stomach pain, or vomiting dizziness drowsiness headache (severe or continuing) nervousness or restlessness slow heartbeat, shortness of breath, or troubled breathing sweating unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Redness and peeling of skin (may occur after a few days) Less common Unusual dryness of skin Other side

Drug Side Effects Summary

effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Acnomel side effects

117. This particular disease, Acne, interacts with the drug Acnomel Acne Mask and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common or rare Skin irritation not present before use of this medicine (moderate or severe) Incidence not known Difficult breathing dryness and peeling of skin fainting hives or itching redness of the skin swelling of the eyes, face, lips, or tongue tightness in the throat unusually warm skin Symptoms of overdose Confusion diarrhea dizziness fast or deep breathing headache (severe or continuing) hearing loss lightheadedness nausea rapid breathing ringing or buzzing in the ears (continuing) severe drowsiness stomach pain vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Skin irritation not present before use of this medicine (mild) stinging Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Acnomel Acne Mask side effects

118. This particular disease, Acne, interacts with the drug Acnomel BP 5 and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of

Drug Side Effects Summary

these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Pain, burning, blistering, changes in color, dryness, itching, peeling, redness, scaling, stinging, or swelling of the skin where the medicine is applied Incidence not known Chest tightness cough difficulty swallowing dizziness fainting fast heartbeat hives, itching, or skin rash increased sensitivity of the skin to sunlight large, hive-like swelling on face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs redness or other discoloration of the skin severe sunburn swelling of the eyes, face, lips, or tongue tightness in the throat trouble breathing unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Dryness or peeling of the skin (may occur after a few days) feeling of warmth, mild stinging, and redness of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Acnomel BP 5 side effects

119. This particular disease, Acne, interacts with the drug Akliel and has the following side effects:

- Akliel Cream may cause serious side effects including: Local skin irritation . Local skin reactions are common with Akliel Cream, are most likely to happen during the first 4 weeks of treatment and may decrease with continued use. Signs and symptoms of local skin reactions include: redness scaling dryness stinging or burning To help reduce your risk of developing these local skin reactions, when you begin treatment, you should begin applying a moisturizer on your skin as often as needed. Tell your healthcare provider if you develop symptoms of a local skin reaction. Your doctor may tell

Drug Side Effects Summary

you to use Akliel Cream less often, or temporarily, or permanently stop your treatment. The most common side effects of Akliel Cream include: itching and sunburn. See What should I avoid while using Akliel Cream? These are not all the possible side effects. For more information, ask your healthcare provider or pharmacist. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to GALDERMA LABORATORIES, L.P. at 1-866-735-4137. Akliel side effects

120. This particular disease, Acne, interacts with the drug Aktipak and has the following side effects

- No side effects listed

121. This particular disease, Acne, interacts with the drug Akurza and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common or rare Skin irritation not present before use of this medicine (moderate or severe) Incidence not known Difficult breathing dryness and peeling of skin fainting hives or itching redness of the skin swelling of the eyes, face, lips, or tongue tightness in the throat unusually warm skin Symptoms of overdose Confusion diarrhea dizziness fast or deep breathing headache (severe or continuing) hearing loss lightheadedness nausea rapid breathing ringing or buzzing in the ears (continuing) severe drowsiness stomach pain vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Skin irritation not present before use of this medicine (mild) stinging Other side effects not listed may also occur in

Drug Side Effects Summary

some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Akurza side effects

122. This particular disease, Acne, interacts with the drug Aliclen and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common or rare Skin irritation not present before use of this medicine (moderate or severe) Incidence not known Difficult breathing dryness and peeling of skin fainting hives or itching redness of the skin swelling of the eyes, face, lips, or tongue tightness in the throat unusually warm skin Symptoms of overdose Confusion diarrhea dizziness fast or deep breathing headache (severe or continuing) hearing loss lightheadedness nausea rapid breathing ringing or buzzing in the ears (continuing) severe drowsiness stomach pain vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Skin irritation not present before use of this medicine (mild) stinging Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Aliclen side effects

123. This particular disease, Acne, interacts with the drug Alquam-X Acne Therapy Gel and has the

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of

Drug Side Effects Summary

these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Pain, burning, blistering, changes in color, dryness, itching, peeling, redness, scaling, stinging, or swelling of the skin where the medicine is applied Incidence not known Chest tightness cough difficulty swallowing dizziness fainting fast heartbeat hives, itching, or skin rash increased sensitivity of the skin to sunlight large, hive-like swelling on face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs redness or other discoloration of the skin severe sunburn swelling of the eyes, face, lips, or tongue tightness in the throat trouble breathing unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Dryness or peeling of the skin (may occur after a few days) feeling of warmth, mild stinging, and redness of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Alquam-X Acne Therapy Gel side effects

124. This particular disease, Acne, interacts with the drug Altabax and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Wipe the ointment off your skin and call your doctor at once if you have: severe burning, itching, redness, swelling, blistering, oozing, or other skin irritation where the medicine was applied
- or new or worsening signs of skin infection . Common Altabax side effects may include: mild itching

Drug Side Effects Summary

or skin irritation

- or headache .

125. This particular disease, Acne, interacts with the drug Altreno and has the following side effects:

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Altreno may cause serious side effects. Call your doctor at once if you have: severe burning, stinging, or irritation of treated skin

- severe skin dryness

- or severe redness, swelling, blistering, peeling, or crusting. Your skin may be more sensitive to weather extremes such as cold and wind while using Altreno. Common side effects of Altreno may include: skin pain, redness, burning, itching, or irritation

- sore throat

- mild warmth or stinging where the medicine was applied

- or changes in color of treated skin.

126. This particular disease, Acne, interacts with the drug Arazlo and has the following side effects:

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Arazlo may cause serious side effects. Stop using Arazlo and call your doctor at once if you have: severe skin irritation (pain, dryness, flaking, peeling, burning, stinging, itching) after applying the medicine

- blistering or peeling skin

- severe redness or discomfort

- or swelling, warmth, oozing, or other signs of skin infection . Common side effects of Arazlo may

Drug Side Effects Summary

include: dry, scaly, or peeling skin

- skin pain, redness or burning
- or itching or other irritation of treated skin.

127. This particular disease, Acne, interacts with the drug Avar and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Skin irritation. Rarely, very bad effects have happened with sulfa drugs . Sometimes, these have been deadly. These effects have included liver problems, blood problems, and very bad skin reactions (Stevens-Johnson syndrome / toxic epidermal necrolysis). Call your doctor right away if you have a rash
- red, swollen, blistered, or peeling skin
- red or irritated eyes
- sores in your mouth, throat, nose, or eyes
- fever, chills, or sore throat
- cough that is new or worse

Drug Side Effects Summary

- feeling very tired or weak
- any bruising or bleeding
- or signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Avar side effects

128. This particular disease, Acne, interacts with the drug Avar-E and has the following side effects

- No side effects listed

129. This particular disease, Acne, interacts with the drug Avar-E Green and has the following side

- No side effects listed

130. This particular disease, Acne, interacts with the drug Avar-E LS and has the following side effects

- No side effects listed

131. This particular disease, Acne, interacts with the drug Avar Cleanser and has the following side effects

- No side effects listed

132. This particular disease, Acne, interacts with the drug Avar LS and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
 - hives
 - itching
 - red, swollen, blistered, or peeling skin with or without fever

Drug Side Effects Summary

- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Skin irritation. Rarely, very bad effects have happened with sulfa drugs . Sometimes, these have been deadly. These effects have included liver problems, blood problems, and very bad skin reactions (Stevens-Johnson syndrome / toxic epidermal necrolysis). Call your doctor right away if you have a rash
- red, swollen, blistered, or peeling skin
- red or irritated eyes
- sores in your mouth, throat, nose, or eyes
- fever, chills, or sore throat
- cough that is new or worse
- feeling very tired or weak
- any bruising or bleeding
- or signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Avar LS side effects

133. This particular disease, Acne, interacts with the drug Avar LS Cleanser and has the following s

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching

Drug Side Effects Summary

- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Skin irritation. Rarely, very bad effects have happened with sulfa drugs . Sometimes, these have been deadly. These effects have included liver problems, blood problems, and very bad skin reactions (Stevens-Johnson syndrome / toxic epidermal necrolysis). Call your doctor right away if you have a rash
- red, swollen, blistered, or peeling skin
- red or irritated eyes
- sores in your mouth, throat, nose, or eyes
- fever, chills, or sore throat
- cough that is new or worse
- feeling very tired or weak
- any bruising or bleeding
- or signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Avar LS Cleanser side effects

134. This particular disease, Acne, interacts with the drug Benzac and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Pain, burning, blistering, changes in color, dryness, itching, peeling, redness, scaling, stinging, or swelling of the skin where the medicine is applied Incidence not known Chest tightness cough difficulty swallowing dizziness

Drug Side Effects Summary

fainting fast heartbeat hives, itching, or skin rash increased sensitivity of the skin to sunlight large, hive-like swelling on face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs redness or other discoloration of the skin severe sunburn swelling of the eyes, face, lips, or tongue tightness in the throat trouble breathing unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Dryness or peeling of the skin (may occur after a few days) feeling of warmth, mild stinging, and redness of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Benzac side effects

135. This particular disease, Acne, interacts with the drug Benzac AC and has the following side effects

- Benzac AC can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Stop using benzoyl peroxide and call your doctor at once if you have any of these side effects on the treated skin: severe itching or burning
- severe stinging or redness
- swelling
- or peeling. Common side effects of Benzac AC may include: mild stinging or burning
- itching or tingly feeling

Drug Side Effects Summary

- skin dryness, peeling, or flaking
- or redness or other irritation.

136. This particular disease, Acne, interacts with the drug Benzac AC Wash and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Benzac AC Wash side effects

137. This particular disease, Acne, interacts with the drug Benzac W and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever

Drug Side Effects Summary

- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Benzac W side effects

138. This particular disease, Acne, interacts with the drug Benzagel and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Pain, burning, blistering, changes in color, dryness, itching, peeling, redness, scaling, stinging, or swelling of the skin where the medicine is applied Incidence not known Chest tightness cough difficulty swallowing dizziness fainting fast heartbeat hives, itching, or skin rash increased sensitivity of the skin to sunlight large, hive-like swelling on face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs redness or other discoloration of the skin severe sunburn swelling of the eyes, face, lips, or tongue tightness in the throat trouble breathing unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Dryness or peeling of the skin (may occur after a few days) feeling of warmth, mild stinging, and redness of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Benzagel side effects

Drug Side Effects Summary

139. This particular disease, Acne, interacts with the drug Benzagel Wash and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Pain, burning, blistering, changes in color, dryness, itching, peeling, redness, scaling, stinging, or swelling of the skin where the medicine is applied Incidence not known Chest tightness cough difficulty swallowing dizziness fainting fast heartbeat hives, itching, or skin rash increased sensitivity of the skin to sunlight large, hive-like swelling on face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs redness or other discoloration of the skin severe sunburn swelling of the eyes, face, lips, or tongue tightness in the throat trouble breathing unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Dryness or peeling of the skin (may occur after a few days) feeling of warmth, mild stinging, and redness of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Benzagel Wash side effects

140. This particular disease, Acne, interacts with the drug Benzamycin Pak and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common or rare Burning, blistering, crusting, itching, severe redness, or swelling of the skin dryness or peeling of the skin eye

Drug Side Effects Summary

irritation feeling of warmth, mild stinging, or redness of the skin increased sensitivity of the skin to sunlight oiliness or tenderness of the skin painful irritation of the skin redness or other discoloration of the skin redness, swelling, or itching of the eyelid severe sunburn skin rash swelling of the eyes, face, and nose Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Benzamycin Pak side effects

141. This particular disease, Acne, interacts with the drug Benzashave and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Pain, burning, blistering, changes in color, dryness, itching, peeling, redness, scaling, stinging, or swelling of the skin where the medicine is applied Incidence not known Chest tightness cough difficulty swallowing dizziness fainting fast heartbeat hives, itching, or skin rash increased sensitivity of the skin to sunlight large, hive-like swelling on face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs redness or other discoloration of the skin severe sunburn swelling of the eyes, face, lips, or tongue tightness in the throat trouble breathing unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Dryness or peeling of the skin (may occur after a few days) feeling of warmth, mild stinging, and redness of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Benzashave side

Drug Side Effects Summary

effects

142. This particular disease, Acne, interacts with the drug BenzEFoam Ultra and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Pain, burning, blistering, changes in color, dryness, itching, peeling, redness, scaling, stinging, or swelling of the skin where the medicine is applied Incidence not known Chest tightness cough difficulty swallowing dizziness fainting fast heartbeat hives, itching, or skin rash increased sensitivity of the skin to sunlight large, hive-like swelling on face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs redness or other discoloration of the skin severe sunburn swelling of the eyes, face, lips, or tongue tightness in the throat trouble breathing unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Dryness or peeling of the skin (may occur after a few days) feeling of warmth, mild stinging, and redness of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. BenzEFoam Ultra side effects

143. This particular disease, Acne, interacts with the drug BenzePro and has the following side effects

- BenzePro can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop

Drug Side Effects Summary

using this medicine and get emergency medical help if you have signs of an allergic reaction : hives, itching

- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Stop using benzoyl peroxide and call your doctor at once if you have any of these side effects on the treated skin: severe itching or burning
- severe stinging or redness
- swelling
- or peeling. Common side effects of BenzePro may include: mild stinging or burning
- itching or tingly feeling
- skin dryness, peeling, or flaking
- or redness or other irritation.

144. This particular disease, Acne, interacts with the drug Benziq and has the following side effects

- Benziq can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Stop using benzoyl peroxide and call your doctor at once if you have any of these side effects on the treated skin: severe itching or burning
- severe stinging or redness
- swelling
- or peeling. Common side effects of Benziq may include: mild stinging or burning
- itching or tingly feeling
- skin dryness, peeling, or flaking

Drug Side Effects Summary

- or redness or other irritation.

145. This particular disease, Acne, interacts with the drug benzoyl peroxide / hydrocortisone and h

- Stop using this medicine and get emergency medical help if you have signs of a serious allergic reaction or severe skin irritation : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe burning, stinging, or redness of treated skin
- severe itching, or other skin irritation
- or signs of skin infection--warmth, redness, swelling, oozing, redness or crusting around your hair follicles. Your skin can absorb hydrocortisone, which may cause steroid side effects throughout the body. Stop using this medicine and call your doctor if you have: weight gain (especially in your face or your upper back and torso)
- slow wound healing, thinning skin, increased body hair
- irregular menstrual periods, changes in sexual function
- or muscle weakness, tired feeling, depression , anxiety , feeling irritable. Children can absorb larger amounts of this medicine through the skin and may be more likely to have side effects. Common side effects of benzoyl peroxide and hydrocortisone topical may include: mild stinging or burning (especially when applying this medicine to your neck or other sensitive skin areas)
- itching or tingly feeling
- skin dryness, peeling, or flaking
- or mild redness or irritation.

Drug Side Effects Summary

146. This particular disease, Acne, interacts with the drug benzoyl peroxide / salicylic acid and has the following side effects:

- Benzoyl peroxide and salicylic acid topical can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have any of these signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Benzoyl peroxide and salicylic acid topical may cause serious side effects. Stop using benzoyl peroxide and salicylic acid topical and call your doctor at once if you have: severe skin irritation after using the medicine
- scaly or peeling skin
- redness or swelling of treated skin
- or nausea , vomiting , diarrhea . Common side effects of benzoyl peroxide and salicylic acid may include: dry skin or mild irritation
- dizziness
- or ringing in your ears.

147. This particular skin disease, Acne, interacts with the drug benzoyl peroxide / tretinoin and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Benzoyl peroxide and tretinoin topical may cause serious side effects. Call your doctor at once if you have: skin irritation, such as dryness, pain, scaling, redness, peeling, stinging or burning. Common side effects of benzoyl peroxide and tretinoin topical may include: swelling, itching, pain, dryness, peeling, redness, and irritation at the application site.

Drug Side Effects Summary

148. This particular disease, Acne, interacts with the drug Binora and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Pain, burning, blistering, changes in color, dryness, itching, peeling, redness, scaling, stinging, or swelling of the skin where the medicine is applied Incidence not known Chest tightness cough difficulty swallowing dizziness fainting fast heartbeat hives, itching, or skin rash increased sensitivity of the skin to sunlight large, hive-like swelling on face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs redness or other discoloration of the skin severe sunburn swelling of the eyes, face, lips, or tongue tightness in the throat trouble breathing unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Dryness or peeling of the skin (may occur after a few days) feeling of warmth, mild stinging, and redness of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Binora side effects

149. This particular disease, Acne, interacts with the drug BP 10-Wash and has the following side effects

- redness, warmth, swelling, itching, stinging, burning, or irritation of treated skin.

150. This particular disease, Acne, interacts with the drug BPO 6 Foaming Cloths and has the following side effects

- No side effects listed

Drug Side Effects Summary

151. This particular disease, Acne, interacts with the drug BPO Gel and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Pain, burning, blistering, changes in color, dryness, itching, peeling, redness, scaling, stinging, or swelling of the skin where the medicine is applied Incidence not known Chest tightness cough difficulty swallowing dizziness fainting fast heartbeat hives, itching, or skin rash increased sensitivity of the skin to sunlight large, hive-like swelling on face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs redness or other discoloration of the skin severe sunburn swelling of the eyes, face, lips, or tongue tightness in the throat trouble breathing unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Dryness or peeling of the skin (may occur after a few days) feeling of warmth, mild stinging, and redness of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. BPO Gel side effects

152. This particular disease, Acne, interacts with the drug BP Wash and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

Drug Side Effects Summary

- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. BP Wash side effects

153. This particular disease, Acne, interacts with the drug Brevoxyl and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Pain, burning, blistering, changes in color, dryness, itching, peeling, redness, scaling, stinging, or swelling of the skin where the medicine is applied Incidence not known Chest tightness cough difficulty swallowing dizziness fainting fast heartbeat hives, itching, or skin rash increased sensitivity of the skin to sunlight large, hive-like swelling on face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs redness or other discoloration of the skin severe sunburn swelling of the eyes, face, lips, or tongue tightness in the throat trouble breathing unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Dryness or peeling of the skin (may occur after a few days) feeling of warmth, mild stinging, and redness of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice

Drug Side Effects Summary

about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Brevoxyl side effects

154. This particular disease, Acne, interacts with the drug Brevoxyl Acne Wash Kit and has the following side effects:

- Brevoxyl Acne Wash Kit can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Stop using benzoyl peroxide and call your doctor at once if you have any of these side effects on the treated skin: severe itching or burning
- severe stinging or redness
- swelling
- or peeling. Common side effects of Brevoxyl Acne Wash Kit may include: mild stinging or burning
- itching or tingly feeling
- skin dryness, peeling, or flaking
- or redness or other irritation.

155. This particular disease, Acne, interacts with the drug CeraVe SA Renewing and has the following side effects:

- CeraVe SA Renewing can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Also stop using CeraVe SA Renewing and call your doctor at once if you have: severe headache , ringing in your ears, problems with hearing, thinking

Drug Side Effects Summary

problems

- severe stomach pain, vomiting , or diarrhea
- a light-headed feeling, like you might pass out
- shortness of breath
- or severe burning, dryness, or irritation of the skin. Common side effects of CeraVe SA Renewing may include: minor skin irritation , rash, or peeling
- or changes in the color of treated skin (usually whitening).

156. This particular disease, Acne, interacts with the drug Clarifoam EF and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Skin irritation. Rarely, very bad effects have happened with sulfa drugs . Sometimes, these have been deadly. These effects have included liver problems, blood problems, and very bad skin reactions (Stevens-Johnson syndrome / toxic epidermal necrolysis). Call your doctor right away if you have a rash
- red, swollen, blistered, or peeling skin

Drug Side Effects Summary

- red or irritated eyes
- sores in your mouth, throat, nose, or eyes
- fever, chills, or sore throat
- cough that is new or worse
- feeling very tired or weak
- any bruising or bleeding
- or signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Clarifoam EF side effects

157. This particular disease, Acne, interacts with the drug clascoterone and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Clascoterone topical may cause serious side effects. Call your doctor at once if you have: severe itching, burning, peeling, or redness of treated skin
- high blood potassium-- nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- or decreased adrenal gland hormones --nausea, vomiting , stomach pain, loss of appetite, feeling tired or light-headed, muscle or joint pain, skin discoloration, craving salty foods. Serious side effects may be more likely to occur in younger people. Common side effects of clascoterone topical may include: redness, itching, stinging, burning, or swelling of treated skin
- or dry or scaly skin.

158. This particular disease, Acne, interacts with the drug Cleanse & Treat and has the following side effects

- Cleanse & Treat can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer

Drug Side Effects Summary

afterward. Stop using this medicine and get emergency medical help if you have any of these signs of an allergic reaction : hives, itching

- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Stop using Cleanse & Treat and call your doctor at once if you have: severe skin irritation after using the medicine
- scaly or peeling skin
- redness or swelling of treated skin
- or nausea , vomiting , diarrhea . Common side effects of Cleanse & Treat may include: dry skin or mild irritation
- dizziness
- or ringing in your ears.

159. This particular disease, Acne, interacts with the drug Cleanse & Treat Plus and has the following

- Cleanse & Treat Plus can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have any of these signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Stop using Cleanse & Treat Plus and call your doctor at once if you have: severe skin irritation after using the medicine
- scaly or peeling skin
- redness or swelling of treated skin
- or nausea , vomiting , diarrhea . Common side effects of Cleanse & Treat Plus may include: dry skin or mild irritation
- dizziness

Drug Side Effects Summary

- or ringing in your ears.

160. This particular disease, Acne, interacts with the drug Clearskin and has the following side effects

- Clearskin can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Stop using benzoyl peroxide and call your doctor at once if you have any of these side effects on the treated skin: severe itching or burning
- severe stinging or redness
- swelling
- or peeling. Common side effects of Clearskin may include: mild stinging or burning
- itching or tingly feeling
- skin dryness, peeling, or flaking
- or redness or other irritation.

161. This particular disease, Acne, interacts with the drug Compound W and has the following side effects

- Compound W can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Also stop using Compound W and call your doctor at once if you have: severe headache , ringing in your ears, problems with hearing, thinking problems

Drug Side Effects Summary

- severe stomach pain, vomiting , or diarrhea
- a light-headed feeling, like you might pass out
- shortness of breath
- or severe burning, dryness, or irritation of the skin. Common side effects of Compound W may include: minor skin irritation , rash, or peeling
- or changes in the color of treated skin (usually whitening).

162. This particular disease, Acne, interacts with the drug DermalZone and has the following side effects:

- DermalZone can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Also stop using DermalZone and call your doctor at once if you have: severe headache , ringing in your ears, problems with hearing, thinking problems
- severe stomach pain, vomiting , or diarrhea
- a light-headed feeling, like you might pass out
- shortness of breath
- or severe burning, dryness, or irritation of the skin. Common side effects of DermalZone may include: minor skin irritation , rash, or peeling
- or changes in the color of treated skin (usually whitening).

163. This particular disease, Acne, interacts with the drug DHS Salicylic Acid 3% and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your

Drug Side Effects Summary

doctor immediately if any of the following side effects occur: Less common or rare Skin irritation not present before use of this medicine (moderate or severe) Incidence not known Difficult breathing dryness and peeling of skin fainting hives or itching redness of the skin swelling of the eyes, face, lips, or tongue tightness in the throat unusually warm skin Symptoms of overdose Confusion diarrhea dizziness fast or deep breathing headache (severe or continuing) hearing loss lightheadedness nausea rapid breathing ringing or buzzing in the ears (continuing) severe drowsiness stomach pain vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Skin irritation not present before use of this medicine (mild) stinging Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. DHS Salicylic Acid 3% side effects

164. This particular disease, Acne, interacts with the drug drospirenone / ethinyl estradiol / levomef

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance
- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood , pain or warmth in one or both legs

Drug Side Effects Summary

- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- liver problems--loss of appetite, upper stomach pain, tiredness, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- increased blood pressure--severe headache, blurred vision, pounding in your neck or ears
- swelling in your hands, ankles, or feet
- a change in the pattern or severity of migraine headaches
- or symptoms of depression--sleep problems, weakness, tired feeling, mood changes. Common side effects of drospirenone, ethinyl estradiol, and levomefolate may include: nausea
- breast tenderness
- headache
- or breakthrough bleeding.

165. This particular disease, Acne, interacts with the drug Dulcolax Milk of Magnesia and has the fo

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness

Drug Side Effects Summary

- or swelling of the mouth, face, lips, tongue, or throat. Dulcolax Milk of Magnesia side effects

166. This particular disease, Acne, interacts with the drug Durasal and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common or rare Skin irritation not present before use of this medicine (moderate or severe) Incidence not known Difficult breathing dryness and peeling of skin fainting hives or itching redness of the skin swelling of the eyes, face, lips, or tongue tightness in the throat unusually warm skin Symptoms of overdose Confusion diarrhea dizziness fast or deep breathing headache (severe or continuing) hearing loss lightheadedness nausea rapid breathing ringing or buzzing in the ears (continuing) severe drowsiness stomach pain vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Skin irritation not present before use of this medicine (mild) stinging Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Durasal side effects

167. This particular disease, Acne, interacts with the drug Enzoclear Foam and has the following side effects

- No side effects listed

168. This particular disease, Acne, interacts with the drug ethinyl estradiol / norethindrone and has

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- increased blood pressure--severe headache , blurred vision, pounding in your neck or ears, anxiety , nosebleed
- signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance
- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood , pain or warmth in one or both legs
- swelling or tenderness in your stomach
- jaundice (yellowing of the skin or eyes)
- unusual vaginal bleeding, pelvic pain
- a breast lump
- memory problems, confusion, unusual behavior
- or high levels of calcium in your blood--nausea, vomiting, constipation , increased thirst or urination, muscle weakness, bone pain, lack of energy. Common side effects of ethinyl estradiol and norethindrone may include: stomach pain
- bloating , swelling, weight gain
- breast pain
- light vaginal bleeding or spotting
- thinning scalp hair
- or headache.

Drug Side Effects Summary

169. This particular disease, Acne, interacts with the drug Ex-Lax Milk of Magnesia and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Ex-Lax Milk of Magnesia may cause serious side effects. Stop using Ex-Lax Milk of Magnesia and call your doctor at once if you have: severe nausea , vomiting , or diarrhea
- no bowel movement after using the medicine as a laxative
- rectal bleeding
- or worsening symptoms. Common side effects of Ex-Lax Milk of Magnesia may include: diarrhea
- or a decreased sense of taste.

170. This particular disease, Acne, interacts with the drug Fabior and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Fabior and call your doctor at once if you have: severe skin irritation (pain, dryness, flaking, peeling, burning, stinging, itching) after applying the medicine
- blistering or peeling skin
- severe redness or discomfort
- or swelling, warmth, oozing, or other signs of skin infection . Common Fabior side effects may include: dry, scaly, or peeling skin
- skin pain, redness or burning
- or itching or other irritation of treated skin.

171. This particular disease, Acne, interacts with the drug Fostex and has the following side effects

Drug Side Effects Summary

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Mood changes. Fast breathing. Upset stomach or throwing up. Ringing in the ears, hearing loss , or any other changes in hearing. Dizziness . Feeling very tired or weak. Diarrhea . Fostex side effects

172. This particular disease, Acne, interacts with the drug Fostex Medicated and has the following s

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat

Drug Side Effects Summary

- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad skin irritation. Fostex Medicated side effects

173. This particular disease, Acne, interacts with the drug Fostex Medicated Cleansing Cream and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: Skin irritation not present before use of this medicine Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Redness and peeling of skin (may occur after a few days) unusual dryness of skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Fostex Medicated Cleansing Cream side effects

174. This particular disease, Acne, interacts with the drug Fostex Wash 10% and has the following side effects:

- Fostex Wash 10% can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Stop using benzoyl peroxide and call your doctor at once if you have any of these side effects on the treated skin: severe itching or burning
- severe stinging or redness
- swelling
- or peeling. Common side effects of Fostex Wash 10% may include: mild stinging or burning
- itching or tingly feeling
- skin dryness, peeling, or flaking
- or redness or other irritation.

175. This particular disease, Acne, interacts with the drug Hydrisalic and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common or rare Skin irritation not present before use of this medicine (moderate or severe) Incidence not known Difficult breathing dryness and peeling of skin fainting hives or itching redness of the skin swelling of the eyes, face, lips, or tongue tightness in the throat unusually warm skin Symptoms of overdose Confusion diarrhea dizziness fast or deep breathing headache (severe or continuing) hearing loss lightheadedness nausea rapid breathing ringing or buzzing in the ears (continuing) severe drowsiness stomach pain vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Skin irritation not present before use of this medicine (mild) stinging Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your

Drug Side Effects Summary

doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Hydrisalic side effects

176. This particular disease, Acne, interacts with the drug Inova and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Inova side effects

177. This particular disease, Acne, interacts with the drug Inova 4/1 and has the following side effects:

- Inova 4/1 can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have any of these signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Stop using Inova 4/1 and call your doctor at once if you have: severe skin irritation after using the medicine

Drug Side Effects Summary

- scaly or peeling skin
- redness or swelling of treated skin
- or nausea , vomiting , diarrhea . Common side effects of Inova 4/1 may include: dry skin or mild irritation
- dizziness
- or ringing in your ears.

178. This particular disease, Acne, interacts with the drug Inova 8/2 and has the following side effects

- Inova 8/2 can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have any of these signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Stop using Inova 8/2 and call your doctor at once if you have: severe skin irritation after using the medicine
- scaly or peeling skin
- redness or swelling of treated skin
- or nausea , vomiting , diarrhea . Common side effects of Inova 8/2 may include: dry skin or mild irritation
- dizziness
- or ringing in your ears.

179. This particular disease, Acne, interacts with the drug Jasmiel and has the following side effects

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Birth control pills may cause serious side effects. Stop using birth control pills and call your doctor at once if you have: signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance
- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood , pain or warmth in one or both legs
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- liver problems--loss of appetite, upper stomach pain, tiredness, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- increased blood pressure--severe headache, blurred vision, pounding in your neck or ears
- swelling in your hands, ankles, or feet
- a change in the pattern or severity of migraine headaches
- or symptoms of depression--sleep problems, weakness, tired feeling, mood changes. Common side effects of Jasmiel may include: nausea, vomiting
- breast tenderness
- headache, mood changes, feeling tired or irritable
- weight gain
- or changes in your menstrual periods, decreased sex drive.

180. This particular disease, Acne, interacts with the drug Liquimat Light and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Liquimat Light may cause serious side effects. Call your doctor at once if you have: severe burning, redness, or swelling where the medicine was

Drug Side Effects Summary

applied

- severe dryness or peeling of treated skin
- or new or worsening skin symptoms. Common side effects of Liquimat Light may include: mild burning, tingling, stinging, itching, or redness
- peeling, dryness
- or oily skin.

181. This particular disease, Acne, interacts with the drug Liquimat Medium and has the following s

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Liquimat Medium may cause serious side effects. Call your doctor at once if you have: severe burning, redness, or swelling where the medicine was applied
- severe dryness or peeling of treated skin
- or new or worsening skin symptoms. Common side effects of Liquimat Medium may include: mild burning, tingling, stinging, itching, or redness
- peeling, dryness
- or oily skin.

182. This particular disease, Acne, interacts with the drug Lo-Zumandimine and has the following s

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives

Drug Side Effects Summary

- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of high potassium levels like a heartbeat that does not feel normal
- feeling confused
- feeling weak, lightheaded, or dizzy
- feeling like passing out
- numbness or tingling
- or shortness of breath. Signs of high blood pressure like very bad headache or dizziness , passing out, or change in eyesight. Signs of gallbladder problems like pain in the upper right belly area, right shoulder area, or between the shoulder blades
- yellow skin or eyes
- fever with chills
- bloating
- or very upset stomach or throwing up. Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight. Depression or other mood changes. Not able to pass urine or change in how much urine is passed. A lump in the breast, breast pain or soreness, or nipple discharge. Vaginal itching or discharge. Spotting or vaginal bleeding that is very bad or does not go away. Eyesight changes or loss, bulging eyes, or change in how contact lenses feel. Call your doctor right away if you have signs of a blood clot like

Drug Side Effects Summary

chest pain or pressure

- coughing up blood
- shortness of breath
- swelling, warmth, numbness, change of color, or pain in a leg or arm
- or trouble speaking or swallowing. Lo-Zumandimine side effects

183. This particular disease, Acne, interacts with the drug Meted and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad skin irritation. Meted side effects

184. This particular disease, Acne, interacts with the drug Mono-Linyah and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Birth control pills may cause serious side effects. Stop using birth control pills and call your doctor at once if you have: signs of a stroke--sudden numbness

Drug Side Effects Summary

or weakness (especially on one side of the body), sudden severe headache , slurred speech, problems with vision or balance

- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood , pain or warmth in one or both legs

- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating

- liver problems--loss of appetite, upper stomach pain, tiredness, fever, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)

- severe headache, pounding in your neck or ears

- swelling in your hands, ankles, or feet

- changes in the pattern or severity of migraine headaches

- a breast lump

- or symptoms of depression--sleep problems, weakness, tired feeling, mood changes. Common side effects of Mono-Linyah may include: stomach pain, gas, nausea, vomiting

- breast tenderness

- acne , darkening of facial skin

- headache, nervousness , mood changes

- problems with contact lenses

- changes in weight

- breakthrough bleeding

- vaginal itching or discharge

- or rash.

185. This particular disease, Acne, interacts with the drug Neuac and has the following side effects

- Benzoyl peroxide can cause a rare but serious allergic reaction or severe skin irritation. These

Drug Side Effects Summary

reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using Neuac and get emergency medical help if you have signs of an allergic reaction : hives, itching

- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe redness, burning, stinging, or peeling of treated skin areas
- or diarrhea that is watery or bloody. Common side effects of Neuac may include: mild burning or stinging
- itching or tingly feeling
- dryness or peeling of treated skin
- or redness or other irritation.

186. This particular disease, Acne, interacts with the drug Neutrogena Clear Pore Cleanser/Mask and

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness

Drug Side Effects Summary

- or swelling of the mouth, face, lips, tongue, or throat. Neutrogena Clear Pore Cleanser/Mask side effects

187. This particular disease, Acne, interacts with the drug Neutrogena Rapid Clear Stubborn Acne a

- Neutrogena Acne Mask can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Stop using benzoyl peroxide and call your doctor at once if you have any of these side effects on the treated skin: severe itching or burning
- severe stinging or redness
- swelling
- or peeling. Common side effects of Neutrogena Acne Mask may include: mild stinging or burning
- itching or tingly feeling
- skin dryness, peeling, or flaking
- or redness or other irritation.

188. This particular disease, Acne, interacts with the drug Nikki and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Birth control pills may cause serious side effects. Stop using birth control pills and call your doctor at once if you have: signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance

Drug Side Effects Summary

- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood , pain or warmth in one or both legs
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- liver problems--loss of appetite, upper stomach pain, tiredness, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- increased blood pressure--severe headache, blurred vision, pounding in your neck or ears
- swelling in your hands, ankles, or feet
- a change in the pattern or severity of migraine headaches
- or symptoms of depression--sleep problems, weakness, tired feeling, mood changes. Common side effects of Nikki may include: nausea, vomiting
- breast tenderness
- headache, mood changes, feeling tired or irritable
- weight gain
- or changes in your menstrual periods, decreased sex drive.

189. This particular disease, Acne, interacts with the drug Oscion and has the following side effects

- Oscion can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Stop using benzoyl peroxide and call your doctor at once if you have any of these side effects on the treated skin: severe itching or burning
- severe stinging or redness

Drug Side Effects Summary

- swelling
- or peeling. Common side effects of Oscion may include: mild stinging or burning
- itching or tingly feeling
- skin dryness, peeling, or flaking
- or redness or other irritation.

190. This particular disease, Acne, interacts with the drug Oxy-10 and has the following side effects

- Oxy-10 can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Stop using benzoyl peroxide and call your doctor at once if you have any of these side effects on the treated skin: severe itching or burning
- severe stinging or redness
- swelling
- or peeling. Common side effects of Oxy-10 may include: mild stinging or burning
- itching or tingly feeling
- skin dryness, peeling, or flaking
- or redness or other irritation.

191. This particular disease, Acne, interacts with the drug Oxy Daily Wash and has the following side effects

- Oxy Daily Wash can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have signs of an allergic

Drug Side Effects Summary

reaction : hives, itching

- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Stop using benzoyl peroxide and call your doctor at once if you have any of these side effects on the treated skin: severe itching or burning
- severe stinging or redness
- swelling
- or peeling. Common side effects of Oxy Daily Wash may include: mild stinging or burning
- itching or tingly feeling
- skin dryness, peeling, or flaking
- or redness or other irritation.

192. This particular disease, Acne, interacts with the drug P & S and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common or rare Skin irritation not present before use of this medicine (moderate or severe) Incidence not known Difficult breathing dryness and peeling of skin fainting hives or itching redness of the skin swelling of the eyes, face, lips, or tongue tightness in the throat unusually warm skin Symptoms of overdose Confusion diarrhea dizziness fast or deep breathing headache (severe or continuing) hearing loss lightheadedness nausea rapid breathing ringing or buzzing in the ears (continuing) severe drowsiness stomach pain vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Skin irritation

Drug Side Effects Summary

not present before use of this medicine (mild) stinging Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. P & S side effects

193. This particular disease, Acne, interacts with the drug PanOxyl 4% Acne Creamy Wash and has

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. PanOxyl 4% Acne Creamy Wash side effects

194. This particular disease, Acne, interacts with the drug Pernox and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives

Drug Side Effects Summary

- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad skin irritation. Pernox side effects

195. This particular disease, Acne, interacts with the drug Persa-Gel and has the following side effects:

- Persa-Gel can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Stop using benzoyl peroxide and call your doctor at once if you have any of these side effects on the treated skin: severe itching or burning
- severe stinging or redness
- swelling
- or peeling. Common side effects of Persa-Gel may include: mild stinging or burning
- itching or tingly feeling
- skin dryness, peeling, or flaking
- or redness or other irritation.

196. This particular disease, Acne, interacts with the drug Plexion and has the following side effects:

- redness, warmth, swelling, itching, stinging, burning, or irritation of treated skin.

Drug Side Effects Summary

197. This particular disease, Acne, interacts with the drug Propa pH Acne Med Cleansing and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Mood changes. Fast breathing. Upset stomach or throwing up. Ringing in the ears, hearing loss , or any other changes in hearing. Dizziness . Feeling very tired or weak. Diarrhea . Propa pH Acne Med Cleansing side effects

198. This particular disease, Acne, interacts with the drug R A Acne and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Although the risk of serious side effects is low when R A Acne is applied to the skin, side effects can occur if the medicine is absorbed into your bloodstream. Your body may absorb this medicine through the skin if you use too much or if you apply it over large skin areas. Skin that is cut or irritated may also absorb more medicine. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: headache , tired feeling, shortness of breath

Drug Side Effects Summary

- blue-colored skin
- fast or irregular heartbeats
- seizure (convulsions)
- or weak or shallow breathing. Children can absorb larger amounts of this medicine through the skin and may be more likely to have side effects.

199. This particular disease, Acne, interacts with the drug Resinol and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Although the risk of serious side effects is low when Resinol is applied to the skin, side effects can occur if the medicine is absorbed into your bloodstream. Your body may absorb this medicine through the skin if you use too much or if you apply it over large skin areas. Skin that is cut or irritated may also absorb more medicine. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: headache , tired feeling, shortness of breath
- blue-colored skin
- fast or irregular heartbeats
- seizure (convulsions)
- or weak or shallow breathing. Children can absorb larger amounts of this medicine through the skin and may be more likely to have side effects.

200. This particular disease, Acne, interacts with the drug resorcinol / sulfur and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: Less common or rare Skin

Drug Side Effects Summary

irritation not present before use of resorcinol and sulfur Symptoms of resorcinol poisoning Diarrhea, nausea, stomach pain, or vomiting dizziness drowsiness headache (severe or continuing) nervousness or restlessness slow heartbeat, shortness of breath, or troubled breathing sweating unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Redness and peeling of skin (may occur after a few days) Less common Unusual dryness of skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Resorcinol / sulfur topical side effects

201. This particular disease, Acne, interacts with the drug resorcinol and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Although the risk of serious side effects is low when resorcinol is applied to the skin, side effects can occur if the medicine is absorbed into your bloodstream. Your body may absorb resorcinol through the skin if you use too much or if you apply it over large skin areas. Skin that is cut or irritated may also absorb more medicine. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: headache , tired feeling, shortness of breath
- blue-colored skin
- fast or irregular heartbeats
- seizure (convulsions)

Drug Side Effects Summary

- or weak or shallow breathing. Children can absorb larger amounts of this medicine through the skin and may be more likely to have side effects.

202. This particular disease, Acne, interacts with the drug retapamulin and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Wipe the ointment off your skin and call your doctor at once if you have: severe burning, itching, redness, swelling, blistering, oozing, or other skin irritation where the medicine was applied
- or new or worsening signs of skin infection . Common side effects of retapamulin topical may include: mild itching or skin irritation
- or headache .

203. This particular disease, Acne, interacts with the drug Riax and has the following side effects:

- Riax can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Stop using benzoyl peroxide and call your doctor at once if you have any of these side effects on the treated skin: severe itching or burning
- severe stinging or redness
- swelling
- or peeling. Common side effects of Riax may include: mild stinging or burning
- itching or tingly feeling

Drug Side Effects Summary

- skin dryness, peeling, or flaking
- or redness or other irritation.

204. This particular disease, Acne, interacts with the drug Sal-Plant Gel and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common or rare Skin irritation not present before use of this medicine (moderate or severe) Incidence not known Difficult breathing dryness and peeling of skin fainting hives or itching redness of the skin swelling of the eyes, face, lips, or tongue tightness in the throat unusually warm skin Symptoms of overdose Confusion diarrhea dizziness fast or deep breathing headache (severe or continuing) hearing loss lightheadedness nausea rapid breathing ringing or buzzing in the ears (continuing) severe drowsiness stomach pain vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Skin irritation not present before use of this medicine (mild) stinging Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Sal-Plant Gel side effects

205. This particular disease, Acne, interacts with the drug Salactic Film and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your

Drug Side Effects Summary

doctor immediately if any of the following side effects occur: Less common or rare Skin irritation not present before use of this medicine (moderate or severe) Incidence not known Difficult breathing dryness and peeling of skin fainting hives or itching redness of the skin swelling of the eyes, face, lips, or tongue tightness in the throat unusually warm skin Symptoms of overdose Confusion diarrhea dizziness fast or deep breathing headache (severe or continuing) hearing loss lightheadedness nausea rapid breathing ringing or buzzing in the ears (continuing) severe drowsiness stomach pain vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Skin irritation not present before use of this medicine (mild) stinging Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Salactic Film side effects

206. This particular disease, Acne, interacts with the drug Salex and has the following side effects:

- Salex can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Also stop using Salex and call your doctor at once if you have: severe headache , ringing in your ears, problems with hearing, thinking problems
- severe stomach pain, vomiting , or diarrhea

Drug Side Effects Summary

- a light-headed feeling, like you might pass out
- shortness of breath
- or severe burning, dryness, or irritation of the skin. Common side effects of Salex may include: minor skin irritation , rash, or peeling
- or changes in the color of treated skin (usually whitening).

207. This particular disease, Acne, interacts with the drug salicylic acid/sulfur and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: Skin irritation not present before use of salicylic acid and sulfur Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Redness and peeling of skin (may occur after a few days) unusual dryness of skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Salicylic acid/sulfur topical side effects

208. This particular disease, Acne, interacts with the drug Salvax and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common or rare Skin irritation not present before use of this medicine (moderate or severe) Incidence not known Difficult breathing

Drug Side Effects Summary

dryness and peeling of skin fainting hives or itching redness of the skin swelling of the eyes, face, lips, or tongue tightness in the throat unusually warm skin Symptoms of overdose Confusion diarrhea dizziness fast or deep breathing headache (severe or continuing) hearing loss lightheadedness nausea rapid breathing ringing or buzzing in the ears (continuing) severe drowsiness stomach pain vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Skin irritation not present before use of this medicine (mild) stinging Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Salvax side effects

209. This particular disease, Acne, interacts with the drug sarecycline and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Sarecycline may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- a spinning sensation
- or increased pressure inside the skull--severe headaches, ringing in your ears, vision problems, pain behind your eyes. Common side effects may include nausea .

210. This particular disease, Acne, interacts with the drug SASTid and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of

Drug Side Effects Summary

these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: Skin irritation not present before use of this medicine Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Redness and peeling of skin (may occur after a few days) unusual dryness of skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. SAS tid side effects

211. This particular disease, Acne, interacts with the drug Scalpicin Scalp Relief and has the following

- Scalpicin Scalp Relief can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Also stop using Scalpicin Scalp Relief and call your doctor at once if you have: severe headache , ringing in your ears, problems with hearing, thinking problems
- severe stomach pain, vomiting , or diarrhea
- a light-headed feeling, like you might pass out
- shortness of breath
- or severe burning, dryness, or irritation of the skin. Common side effects of Scalpicin Scalp Relief

Drug Side Effects Summary

may include: minor skin irritation , rash, or peeling

- or changes in the color of treated skin (usually whitening).

212. This particular disease, Acne, interacts with the drug Sebulex and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

- itching

- red, swollen, blistered, or peeling skin with or without fever

- wheezing

- tightness in the chest or throat

- trouble breathing, swallowing, or talking

- unusual hoarseness

- or swelling of the mouth, face, lips, tongue, or throat. Very bad skin irritation. Sebulex side effects

213. This particular disease, Acne, interacts with the drug SoluCLENZ Rx and has the following side effects

- SoluCLENZ Rx can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have signs of an allergic reaction : hives, itching

- difficult breathing, feeling light-headed

- swelling of your face, lips, tongue, or throat. Stop using benzoyl peroxide and call your doctor at once if you have any of these side effects on the treated skin: severe itching or burning

Drug Side Effects Summary

- severe stinging or redness
- swelling
- or peeling. Common side effects of SoluCLENZ Rx may include: mild stinging or burning
- itching or tingly feeling
- skin dryness, peeling, or flaking
- or redness or other irritation.

214. This particular disease, Acne, interacts with the drug SSS 10-4 and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Skin irritation. Rarely, very bad effects have happened with sulfa drugs . Sometimes, these have been deadly. These effects have included liver problems, blood problems, and very bad skin reactions (Stevens-Johnson syndrome / toxic epidermal necrolysis). Call your doctor right away if you have a rash
- red, swollen, blistered, or peeling skin
- red or irritated eyes

Drug Side Effects Summary

- sores in your mouth, throat, nose, or eyes
- fever, chills, or sore throat
- cough that is new or worse
- feeling very tired or weak
- any bruising or bleeding
- or signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. SSS 10-4 side effects

215. This particular disease, Acne, interacts with the drug SSS 10-5 and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Skin irritation. Rarely, very bad effects have happened with sulfa drugs . Sometimes, these have been deadly. These effects have included liver problems, blood problems, and very bad skin reactions (Stevens-Johnson syndrome / toxic epidermal necrolysis). Call your doctor right away if you have a rash
- red, swollen, blistered, or peeling skin

Drug Side Effects Summary

- red or irritated eyes
- sores in your mouth, throat, nose, or eyes
- fever, chills, or sore throat
- cough that is new or worse
- feeling very tired or weak
- any bruising or bleeding
- or signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. SSS 10-5 side effects

216. This particular disease, Acne, interacts with the drug SSS Cleanser and has the following side effects:

- No side effects listed

217. This particular disease, Acne, interacts with the drug Stri-Dex and has the following side effects:

- Stri-Dex can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using this medicine and get emergency medical help if you have signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Also stop using Stri-Dex and call your doctor at once if you have: severe headache , ringing in your ears, problems with hearing, thinking problems
- severe stomach pain, vomiting , or diarrhea
- a light-headed feeling, like you might pass out
- shortness of breath
- or severe burning, dryness, or irritation of the skin. Common side effects of Stri-Dex may include: minor skin irritation , rash, or peeling

Drug Side Effects Summary

- or changes in the color of treated skin (usually whitening).

218. This particular disease, Acne, interacts with the drug Stridex Body Focus and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Mood changes. Fast breathing. Upset stomach or throwing up. Ringing in the ears, hearing loss , or any other changes in hearing. Dizziness . Feeling very tired or weak. Diarrhea . Stridex Body Focus side effects

219. This particular disease, Acne, interacts with the drug Stridex Maximum Strength and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching

Drug Side Effects Summary

- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Mood changes. Fast breathing. Upset stomach or throwing up. Ringing in the ears, hearing loss , or any other changes in hearing. Dizziness . Feeling very tired or weak. Diarrhea . Stridex Maximum Strength side effects

220. This particular disease, Acne, interacts with the drug Sulfacleanse 8/4 and has the following side effects:

- redness, warmth, swelling, itching, stinging, burning, or irritation of treated skin.

221. This particular disease, Acne, interacts with the drug Sulfoam and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Sulfoam may cause serious side effects. Call your doctor at once if you have: severe burning, redness, or swelling where the medicine was applied
- severe dryness or peeling of treated skin
- or new or worsening skin symptoms. Common side effects of Sulfoam may include: mild burning, tingling, stinging, itching, or redness
- peeling, dryness
- or oily skin.

222. This particular disease, Acne, interacts with the drug sulfur and has the following side effects:

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Sulfur topical may cause serious side effects. Call your doctor at once if you have: severe burning, redness, or swelling where the medicine was applied
- severe dryness or peeling of treated skin
- or new or worsening skin symptoms. Common side effects of sulfur topical may include: mild burning, tingling, stinging, itching, or redness
- peeling, dryness
- or oily skin.

223. This particular disease, Acne, interacts with the drug Sulpho-Lac Soap and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Sulpho-Lac Soap may cause serious side effects. Call your doctor at once if you have: severe burning, redness, or swelling where the medicine was applied
- severe dryness or peeling of treated skin
- or new or worsening skin symptoms. Common side effects of Sulpho-Lac Soap may include: mild burning, tingling, stinging, itching, or redness
- peeling, dryness
- or oily skin.

224. This particular disease, Acne, interacts with the drug Sumadan and has the following side effects

- redness, warmth, swelling, itching, stinging, burning, or irritation of treated skin.

225. This particular disease, Acne, interacts with the drug Sumaxin and has the following side effects

Drug Side Effects Summary

- redness, warmth, swelling, itching, stinging, burning, or irritation of treated skin.

226. This particular disease, Acne, interacts with the drug Suphera and has the following side effects:

- redness, warmth, swelling, itching, stinging, burning, or irritation of treated skin.

227. This particular disease, Acne, interacts with the drug Syeda and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Birth control pills may cause serious side effects. Stop using birth control pills and call your doctor at once if you have: signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance
- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood , pain or warmth in one or both legs
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- liver problems--loss of appetite, upper stomach pain, tiredness, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- increased blood pressure--severe headache, blurred vision, pounding in your neck or ears
- swelling in your hands, ankles, or feet
- a change in the pattern or severity of migraine headaches
- or symptoms of depression--sleep problems, weakness, tired feeling, mood changes. Common side effects of Syeda may include: nausea, vomiting
- breast tenderness
- headache, mood changes, feeling tired or irritable

Drug Side Effects Summary

- weight gain
- or changes in your menstrual periods, decreased sex drive.

228. This particular disease, Acne, interacts with the drug Tinamed Plantar and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Mood changes. Fast breathing. Upset stomach or throwing up. Ringing in the ears, hearing loss , or any other changes in hearing. Dizziness . Feeling very tired or weak. Diarrhea . Tinamed Plantar side effects

229. This particular disease, Acne, interacts with the drug Tri-Estarylla and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives

Drug Side Effects Summary

- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of high blood pressure like very bad headache or dizziness , passing out, or change in eyesight. Signs of gallbladder problems like pain in the upper right belly area, right shoulder area, or between the shoulder blades
- yellow skin or eyes
- fever with chills
- bloating
- or very upset stomach or throwing up. Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight. Depression or other mood changes. Feeling very tired or weak. Swelling. Not able to pass urine or change in how much urine is passed. A lump in the breast, breast pain or soreness, or nipple discharge. Vaginal itching or discharge. Eyesight changes or loss, bulging eyes, or change in how contact lenses feel. Call your doctor right away if you have signs of a blood clot like chest pain or pressure
- coughing up blood
- shortness of breath
- swelling, warmth, numbness , change of color, or pain in a leg or arm
- or trouble speaking or swallowing. Tri-Estarylla side effects

Drug Side Effects Summary

230. This particular disease, Acne, interacts with the drug Tri-Linyah and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

- itching

- red, swollen, blistered, or peeling skin with or without fever

- wheezing

- tightness in the chest or throat

- trouble breathing, swallowing, or talking

- unusual hoarseness

- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of high blood pressure like very bad headache or dizziness, passing out, or change in eyesight. Signs of gallbladder problems like pain in the upper right belly area, right shoulder area, or between the shoulder blades

- yellow skin or eyes

- fever with chills

- bloating

- or very upset stomach or throwing up. Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight. Depression or other mood changes. Feeling very tired or weak. Swelling. Not able to pass urine or change in how much urine is passed. A lump in the breast, breast pain or soreness, or nipple discharge. Vaginal itching or discharge. Eyesight changes or loss, bulging eyes, or change in how contact lenses feel.

Drug Side Effects Summary

Call your doctor right away if you have signs of a blood clot like chest pain or pressure

- coughing up blood
- shortness of breath
- swelling, warmth, numbness , change of color, or pain in a leg or arm
- or trouble speaking or swallowing. Tri-Linyah side effects

231. This particular disease, Acne, interacts with the drug Tri-Lo-Marzia and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Birth control pills may cause serious side effects. Stop using birth control pills and call your doctor at once if you have: signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache , slurred speech, problems with vision or balance
- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood , pain or warmth in one or both legs
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating
- liver problems--loss of appetite, upper stomach pain, tiredness, fever, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- severe headache, pounding in your neck or ears
- swelling in your hands, ankles, or feet
- changes in the pattern or severity of migraine headaches
- a breast lump
- or symptoms of depression--sleep problems, weakness, tired feeling, mood changes. Common side effects of Tri-Lo-Marzia may include: stomach pain, gas, nausea, vomiting

Drug Side Effects Summary

- breast tenderness
- acne , darkening of facial skin
- headache, nervousness , mood changes
- problems with contact lenses
- changes in weight
- breakthrough bleeding
- vaginal itching or discharge
- or rash.

232. This particular disease, Acne, interacts with the drug trifarotene and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Trifarotene topical may cause serious side effects. Stop using trifarotene topical and call your doctor at once if you have: severe skin irritation (burning, stinging, scaling) after applying the medicine
- or severe skin redness or dryness. Common side effects of trifarotene topical may include: itching or other skin irritation
- or sunburn.

233. This particular disease, Acne, interacts with the drug Twyneo and has the following side effects

- Twyneo may cause serious side effects including: Allergic reactions. Stop treatment and get medical help right away if you have any of the following symptoms: hives, rash or severe itching swelling of your face, eyes, lips, tongue, or throat trouble breathing or throat tightness feeling faint, dizzy, or lightheaded Skin irritation. Twyneo may cause skin irritation such as redness, scaling, peeling, dryness, pain, stinging or burning. If you develop these symptoms, your healthcare provider

Drug Side Effects Summary

may tell you to use a moisturizer, decrease the number of times you apply Twyneo, or completely stop treatment. The most common side effects include pain, dryness, peeling, redness, swelling, itching, and irritation at the application site. These are not all the possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA1088. You may also report side effects to Sol-Gel Technologies, Inc. at 1-866-748-2377. Twyneo side effects

234. This particular disease, Acne, interacts with the drug Vanoxide-HC and has the following side effects

- Stop using this medicine and get emergency medical help if you have signs of a serious allergic reaction or severe skin irritation : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe burning, stinging, or redness of treated skin
- severe itching, or other skin irritation
- or signs of skin infection--warmth, redness, swelling, oozing, redness or crusting around your hair follicles. Your skin can absorb hydrocortisone , which may cause steroid side effects throughout the body. Stop using this medicine and call your doctor if you have: weight gain (especially in your face or your upper back and torso)
- slow wound healing, thinning skin, increased body hair
- irregular menstrual periods, changes in sexual function
- or muscle weakness, tired feeling, depression , anxiety , feeling irritable. Children can absorb larger amounts of this medicine through the skin and may be more likely to have side effects. Common side effects of Vanoxide-HC may include: mild stinging or burning (especially when

Drug Side Effects Summary

applying this medicine to your neck or other sensitive skin areas)

- itching or tingly feeling
- skin dryness, peeling, or flaking
- or mild redness or irritation.

235. This particular disease, Acne, interacts with the drug Winlevi and has the following side effects:

- Winlevi cream can cause serious side effects, including: Local skin reactions. Winlevi cream may cause local skin irritation including itching, burning, skin redness or peeling. Symptoms of a disorder where the adrenal gland does not make enough of certain hormones (adrenal insufficiency) during treatment with Winlevi. Your healthcare provider may stop your treatment with Winlevi if you develop any adrenal problems. The most common side effects of Winlevi cream include reddening, scaling or dryness, and itching of the skin being treated. These are not all of the possible side effects. Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA1088. Winlevi side effects

236. This particular disease, Acne, interacts with the drug Z-Clinz 10 and has the following side effects:

- Benzoyl peroxide can cause a rare but serious allergic reaction or severe skin irritation. These reactions may occur just a few minutes after you apply the medicine, or within a day or longer afterward. Stop using Z-Clinz 10 and get emergency medical help if you have signs of an allergic reaction : hives, itching
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe redness, burning, stinging, or peeling of treated skin areas
- or diarrhea that is watery or bloody. Common side effects of Z-Clinz 10 may include: mild burning

Drug Side Effects Summary

or stinging

- itching or tingly feeling
- dryness or peeling of treated skin
- or redness or other irritation.

237. This particular disease, Acne, interacts with the drug Zarah and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Birth control pills may cause serious side effects. Stop using birth control pills and call your doctor at once if you have: signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance
- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood , pain or warmth in one or both legs
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- liver problems--loss of appetite, upper stomach pain, tiredness, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- increased blood pressure--severe headache, blurred vision, pounding in your neck or ears
- swelling in your hands, ankles, or feet
- a change in the pattern or severity of migraine headaches
- or symptoms of depression--sleep problems, weakness, tired feeling, mood changes. Common side effects of Zarah may include: nausea, vomiting
- breast tenderness
- headache, mood changes, feeling tired or irritable

Drug Side Effects Summary

- weight gain
- or changes in your menstrual periods, decreased sex drive.

238. This particular disease, Acne, interacts with the drug Zumandimine and has the following side

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of high potassium levels like a heartbeat that does not feel normal
- feeling confused
- feeling weak, lightheaded, or dizzy
- feeling like passing out
- numbness or tingling
- or shortness of breath. Signs of high blood pressure like very bad headache or dizziness , passing out, or change in eyesight. Signs of gallbladder problems like pain in the upper right belly area, right shoulder area, or between the shoulder blades

Drug Side Effects Summary

- yellow skin or eyes
- fever with chills
- bloating
- or very upset stomach or throwing up. Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight. Depression or other mood changes. Not able to pass urine or change in how much urine is passed. A lump in the breast, breast pain or soreness, or nipple discharge. Vaginal itching or discharge. Spotting or vaginal bleeding that is very bad or does not go away. Eyesight changes or loss, bulging eyes, or change in how contact lenses feel. Call your doctor right away if you have signs of a blood clot like chest pain or pressure
- coughing up blood
- shortness of breath
- swelling, warmth, numbness, change of color, or pain in a leg or arm
- or trouble speaking or swallowing. Zomandimine side effects

239. This particular disease, ADHD, interacts with the drug Vyvanse and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: signs of heart problems - chest pain, trouble breathing, pounding heartbeats or fluttering in your chest, feeling like you might pass out
- signs of psychosis - hallucinations (seeing or hearing things that are not real), new behavior problems, aggression, hostility, paranoia
- or signs of circulation problems - numbness, pain, cold feeling, unexplained wounds, or skin color changes (pale, red, or blue appearance) in your fingers or toes. Seek medical attention right away if

Drug Side Effects Summary

you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Vyvanse can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common Vyvanse side effects may include: dry mouth, loss of appetite, weight loss

- sleep problems (insomnia)
- fast heart rate, feeling jittery
- dizziness, feeling anxious or irritable
- or nausea, vomiting, stomach pain, diarrhea, constipation.

240. This particular disease, ADHD, interacts with the drug Adderall and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Adderall may cause serious side effects. Call your doctor at once if you have: signs of heart problems - chest pain, trouble breathing, feeling like you might pass out
- signs of psychosis - hallucinations (seeing or hearing things that are not real), new behavior problems, aggression, hostility, paranoia
- signs of circulation problems - numbness, pain, cold feeling, unexplained wounds, or skin color changes (pale, red, or blue appearance) in your fingers or toes
- a seizure (convulsions)
- muscle twitches (tics)
- or changes in your vision. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Adderall can affect growth in

Drug Side Effects Summary

children. Tell your doctor if your child is not growing at a normal rate while using this medicine.

Common side effects of Adderall may include: stomach pain

- loss of appetite
- weight loss
- mood changes
- feeling nervous
- fast heart rate
- headache
- dizziness
- sleep problems (insomnia)
- or dry mouth.

241. This particular disease, ADHD, interacts with the drug Adderall XR and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Adderall XR may cause serious side effects. Call your doctor at once if you have: signs of heart problems--chest pain, trouble breathing, feeling like you might pass out
- signs of psychosis--hallucinations (seeing or hearing things that are not real), new behavior problems, aggression, hostility, paranoia
- signs of circulation problems--numbness, pain, cold feeling, unexplained wounds, or skin color changes (pale, red, or blue appearance) in your fingers or toes
- a seizure (convulsions)
- muscle twitches (tics)
- or changes in your vision. Seek medical attention right away if you have symptoms of serotonin

Drug Side Effects Summary

syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Long-term use of stimulant medicine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common side effects of Adderall XR may include: stomach pain, nausea, loss of appetite

- weight loss
- mood changes, feeling nervous or irritable
- fast heart rate
- headache , dizziness
- sleep problems (insomnia)
- or dry mouth.

242. This particular disease, ADHD, interacts with the drug Concerta and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: signs of heart problems - chest pain, trouble breathing, feeling like you might pass out
- signs of psychosis - hallucinations (seeing or hearing things that are not real), new behavior problems, aggression, hostility, paranoia
- signs of circulation problems - numbness , pain, cold feeling, unexplained wounds, or skin color changes (pale, red, or blue appearance) in your fingers or toes
- or penis erection that is painful or lasts 4 hours or longer. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Methylphenidate can affect growth in children. Your child's height and weight may need to be

Drug Side Effects Summary

checked often. Tell your doctor if your child is not growing at a normal rate. Common Concerta side effects may include: sweating, increased blood pressure

- mood changes, anxiety, feeling nervous or irritable, trouble sleeping
- fast heart rate, pounding heartbeats or fluttering in your chest
- loss of appetite, weight loss
- dry mouth, nausea, vomiting, stomach pain, indigestion
- or headache, dizziness .

243. This particular disease, ADHD, interacts with the drug lisdexamfetamine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Lisdexamfetamine may cause serious side effects. Call your doctor at once if you have: signs of heart problems--chest pain, trouble breathing, pounding heartbeats or fluttering in your chest, feeling like you might pass out
- signs of psychosis--hallucinations (seeing or hearing things that are not real), new behavior problems, aggression, hostility, paranoia
- or signs of circulation problems--numbness, pain, cold feeling, unexplained wounds, or skin color changes (pale, red, or blue appearance) in your fingers or toes. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting , or diarrhea . Lisdexamfetamine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using lisdexamfetamine. Common side effects of lisdexamfetamine may include: dry mouth, loss of appetite, weight loss
- sleep problems (insomnia)
- fast heart rate, feeling jittery

Drug Side Effects Summary

- dizziness , feeling anxious or irritable
- or nausea, vomiting, stomach pain, diarrhea, constipation .

244. This particular disease, ADHD, interacts with the drug Daytrana and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Aggressive and violent behavior arm, back, or jaw pain black, tarry stools blistering, peeling, or loosening of the skin blood in the urine bloody nose chest pain or discomfort chest tightness or heaviness chills cold confusion cough or hoarseness cracks in the skin crying depersonalization diarrhea dizziness dysphoria euphoria fainting fast, pounding, or irregular heartbeat or pulse fever flu-like symptoms heavier menstrual periods joint or muscle pain loss of heat from the body nausea nervousness paranoia pinpoint red spots on the skin quick to react or overreact emotionally rapidly changing moods red, irritated eyes red, swollen skin scaly skin seeing, hearing, or feeling things that are not there severe or sudden headache skin rash or itching sore throat sores, ulcers, or white spots in the mouth or on the lips sudden loss of coordination sudden slurring of speech sweating twitching, twisting, or uncontrolled repetitive movements of the tongue, lips, face, arms, or legs unusual bleeding or bruising unusual tiredness or weakness Rare Convulsions difficulty with breathing high fever increased sweating loss of bladder control severe muscle stiffness tiredness uncontrolled vocal outbursts or tics (uncontrolled repeated body movements) unusually pale skin Incidence not known Blistering, burning, crusting, dryness, or flaking of the skin chest pain difficulty with swallowing hives, itching, scaling, severe redness, soreness, or swelling of the skin large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs painful or difficult urination pale skin paleness or cold feeling in the fingertips and toes puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue swollen glands tingling or pain in the fingers or toes when exposed to

Drug Side Effects Summary

cold troubled breathing with exertion Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Abdominal or stomach pain decreased weight difficulty with moving drowsiness headache loss of appetite muscle pain or stiffness redness of the skin trouble sleeping vomiting welts Less common Decreased appetite stuffy or runny nose weight loss Incidence not known Bigger, dilated, or enlarged pupils (black part of the eye) bleeding, bruising, burning, itching, redness, skin rash, swelling, or soreness at the application site blurred or loss of vision disturbed color perception double vision hair loss halos around lights increased sensitivity of the eyes to light night blindness overbright appearance of lights thinning of the hair tunnel vision Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Daytrana side effects

245. This particular disease, ADHD, interacts with the drug Quillivant XR and has the following side

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: signs of heart problems - chest pain, trouble breathing, feeling like you might pass out
- signs of psychosis - hallucinations (seeing or hearing things that are not real), new behavior problems, aggression, hostility, paranoia
- signs of circulation problems - numbness, pain, cold feeling, unexplained wounds, or skin color changes (pale, red, or blue appearance) in your fingers or toes

Drug Side Effects Summary

- or penis erection that is painful or lasts 4 hours or longer (rare). Methylphenidate can affect growth in children. Tell your doctor if your child is not growing at a normal rate. Common Quillivant XR side effects may include: excessive sweating
- mood changes, feeling nervous or irritable, sleep problems (insomnia)
- fast heart rate , pounding heartbeats or fluttering in your chest, increased blood pressure
- loss of appetite, weight loss
- dry mouth, nausea , stomach pain
- or headache .

246. This particular disease, ADHD, interacts with the drug amantadine and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: extreme drowsiness, falling asleep suddenly even after feeling alert
- a light-headed feeling, like you might pass out
- shortness of breath (even with mild exertion), swelling in your hands or feet
- painful or difficult urination
- depression , agitation, aggression, behavior changes, hallucinations, thoughts of hurting yourself
- a seizure
- or severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors. You may have increased sexual urges, unusual urges to gamble, or other intense urges while taking this medicine. Talk with your doctor if this occurs. Side effects may be more likely in older adults. Common amantadine side effects may include: dizziness , falls
- dry mouth
- swelling in your legs or feet

Drug Side Effects Summary

- nausea , constipation
- or sleep problems (insomnia).

247. This particular disease, ADHD, interacts with the drug Cotelma XR-ODT and has the following

- Cotelma XR-ODT can cause serious side effects, including: See "Important information" above. Painful and prolonged erections (priapism). Priapism has happened in males who take products that contain methylphenidate. If your child develops priapism, get medical help right away. Circulation problems in fingers and toes (peripheral vasculopathy, including Raynaud's phenomenon). Signs and symptoms may include: fingers or toes may feel numb, cool, or painful fingers or toes may change color from pale, to blue, to red Tell your healthcare provider if& your child has numbness, pain, skin color change, or sensitivity to temperature in the fingers or toes. Call your healthcare provider right away if your child has any signs of unexplained wounds appearing on fingers or toes during treatment with this medication. Slowing of growth (height and weight) in children. Children should have their height and weight checked often during treatment. Treatment may be stopped if your child is not gaining weight or height. The most common side effects of methylphenidate products include: decreased appetite trouble sleeping nausea vomiting indigestion stomach pain weight loss anxiety dizziness irritability mood swings (affect liability) increased heart rate increased blood pressure These are not all the possible side effects of this medication. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Cotelma XR-ODT side effects

248. This particular disease, ADHD, interacts with the drug Jornay PM and has the following side ef

- Jornay PM can cause serious side effects, including: See "Important information" above. Painful and prolonged erections (priapism). Priapism has happened in males who take products that contain methylphenidate. If you or your child develops priapism, get medical help right away. Circulation

Drug Side Effects Summary

problems in fingers and toes (peripheral vasculopathy, including Raynaud's phenomenon). Signs and symptoms may include: fingers or toes may feel numb, cool, or painful fingers or toes may change color from pale, to blue, to red Tell your healthcare provider if you or your child has numbness, pain, skin color change, or sensitivity to temperature in the fingers or toes. Call your healthcare provider right away if you or your child has any signs of unexplained wounds appearing on fingers or toes during treatment with this medication. Slowing of growth (height and weight) in children. Children should have their height and weight checked often during treatment. Treatment may be stopped if your child is not gaining weight or height. The most common side effects of methylphenidate products in children, adolescents, and adults with ADHD include: decreased appetite trouble sleeping nausea vomiting indigestion stomach pain weight loss anxiety dizziness irritability mood swings (affect liability) increased heart rate increased blood pressure The most common side effects of Jornay PM, in children age 6 to 12 with ADHD include: trouble sleeping decreased appetite restlessness (psychomotor hyperactivity) headache nausea mood swings vomiting These are not all the possible side effects of this medication. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Jornay PM side effects

249. This particular disease, ADHD, interacts with the drug methylphenidate and has the following s

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: signs of heart problems - chest pain, trouble breathing, feeling like you might pass out
- signs of psychosis - hallucinations (seeing or hearing things that are not real), new behavior problems, aggression, hostility, paranoia
- signs of circulation problems - numbness , pain, cold feeling, unexplained wounds, or skin color

Drug Side Effects Summary

changes (pale, red, or blue appearance) in your fingers or toes

- or penis erection that is painful or lasts 4 hours or longer. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea.

Methylphenidate can affect growth in children. Your child's height and weight may need to be checked often. Tell your doctor if your child is not growing at a normal rate. Common methylphenidate side effects may include: sweating, increased blood pressure

- mood changes, anxiety, feeling nervous or irritable, trouble sleeping

- fast heart rate, pounding heartbeats or fluttering in your chest

- loss of appetite, weight loss

- dry mouth, nausea, vomiting, stomach pain, indigestion

- or headache, dizziness .

250. This particular disease, ADHD, interacts with the drug QuilliChew ER and has the following side effects:

- QuilliChew ER can cause serious side effects, including: See "Important information" above.

Painful and prolonged erections (priapism). Priapism has happened in males who take products that contain methylphenidate. If you or your child develops priapism, get medical help right away.

Priapism can cause long lasting damage, it should be checked by a health care provider right away.

Slowing of growth (height and weight) in children. Children should have their height and weight checked often during treatment. Treatment may be stopped if your child is not gaining weight or height.

The most common side effects of QuilliChew ER include: decreased appetite trouble sleeping nausea vomiting indigestion stomach pain weight loss anxiety dizziness irritability mood swings fast heart beat increased blood pressure These are not all the possible side effects of this medication. Call your doctor for medical advice about side effects. You may report side effects to

FDA at 1-800-FDA-1088. QuilliChew ER side effects

Drug Side Effects Summary

251. This particular disease, ADHD, interacts with the drug Azstarys and has the following side effects

- Azstarys may cause serious side effects, including: See Important information Painful and prolonged erections (priapism). Priapism has happened in males who take products that contain methylphenidate. If you or your child develop priapism, get medical help right away. Circulation problems in fingers and toes (peripheral vasculopathy, including Raynaud's phenomenon). Signs and symptoms may include: fingers or toes may feel numb, cool, or painful fingers or toes may change color from pale, to blue, to red Tell your healthcare provider if you or your child have numbness, pain, skin color change, or sensitivity to temperature in the fingers or toes. Call your healthcare provider right away if you or your child have any signs of unexplained wounds appearing on fingers or toes during treatment. Slowing of growth (height and weight) in children. Children should have their height and weight checked often during treatment. Azstarys treatment may be stopped if your child is not growing or gaining weight. The most common side effects include: decreased appetite nausea indigestion trouble sleeping vomiting stomach pain weight loss dizziness mood swings increased blood pressure anxiety irritability increased heart rate These are not all the possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Azstarys side effects

252. This particular disease, ADHD, interacts with the drug Qelbree and has the following side effects

- Qelbree can cause serious side effects, including: See Important information Increased blood pressure and heart rate. Your healthcare provider should check you or your child's blood pressure and heart rate before starting and during treatment. Manic episodes. Manic episodes may happen in people with bipolar disorder who take Qelbree. Symptoms may include: greatly increased energy racing thoughts unusually grand ideas talking more or faster than usual severe trouble sleeping reckless behavior excessive happiness or irritability Sleepiness and tiredness. See What should I avoid while taking Qelbree? The most common side effects in children 6 to 17 years of age include:

Drug Side Effects Summary

sleepiness tiredness vomiting irritability decreased appetite nausea trouble sleeping The most common side effects in adults include: insomnia sleepiness nausea dry mouth headache tiredness decreased appetite constipation Effects on weight. Your healthcare provider should check you or your child's weight before starting, and during treatment. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Qelbree side effects

253. This particular disease, ADHD, interacts with the drug Strattera and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: anxiety, panic attacks, trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), depressed, or have thoughts about suicide or hurting yourself. Strattera can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Call your doctor at once if you have: signs of heart problems - chest pain, trouble breathing, feeling like you might pass out
- signs of psychosis - hallucinations (seeing or hearing things that are not real), new behavior problems, aggression, hostility, paranoia
- liver problems - stomach pain (upper right side), itching, flu-like symptoms, dark urine, jaundice (yellowing of the skin or eyes)
- painful or difficult urination
- or erection is painful or lasts longer than 4 hours (this is a rare side effect). Common Strattera side effects may include: nausea, vomiting, upset stomach, constipation
- dry mouth, loss of appetite
- mood changes, insomnia, feeling tired
- dizziness

Drug Side Effects Summary

- increased sweating
- urination problems
- or impotence, trouble having an erection.

254. This particular disease, ADHD, interacts with the drug Dexedrine and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Dexedrine Spansule may cause serious side effects. Call your doctor at once if you have: signs of heart problems--chest pain, trouble breathing, feeling like you might pass out
- signs of psychosis--hallucinations (seeing or hearing things that are not real), new behavior problems, aggression, hostility, paranoia
- signs of circulation problems--numbness, pain, cold feeling, unexplained wounds, or skin color changes (pale, red, or blue appearance) in your fingers or toes
- a seizure (convulsions)
- muscle twitches (tics)
- or changes in your vision. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Dexedrine Spansule can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using Dexedrine Spansule. Common side effects of Dexedrine Spansule may include: dry mouth, upset stomach , loss of appetite, weight loss
- headache , dizziness
- tremors, fast heartbeats
- or sleep problems (insomnia).

Drug Side Effects Summary

255. This particular disease, ADHD, interacts with the drug amphetamine / dextroamphetamine and

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Amphetamine and dextroamphetamine may cause serious side effects. Call your doctor at once if you have: signs of heart problems--chest pain, trouble breathing, feeling like you might pass out
- signs of psychosis--hallucinations (seeing or hearing things that are not real), new behavior problems, aggression, hostility, paranoia
- signs of circulation problems--numbness, pain, cold feeling, unexplained wounds, or skin color changes (pale, red, or blue appearance) in your fingers or toes
- a seizure (convulsions)
- muscle twitches (tics)
- or changes in your vision. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Long-term use of stimulant medicine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common side effects of amphetamine and dextroamphetamine may include: stomach pain, nausea, loss of appetite
- weight loss
- mood changes, feeling nervous or irritable
- fast heart rate
- headache , dizziness
- sleep problems (insomnia)
- or dry mouth.

Drug Side Effects Summary

256. This particular disease, ADHD, interacts with the drug Ritalin and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: signs of heart problems - chest pain, trouble breathing, feeling like you might pass out
- signs of psychosis - hallucinations (seeing or hearing things that are not real), new behavior problems, aggression, hostility, paranoia
- signs of circulation problems - numbness , pain, cold feeling, unexplained wounds, or skin color changes (pale, red, or blue appearance) in your fingers or toes
- or penis erection that is painful or lasts 4 hours or longer. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Methylphenidate can affect growth in children. Your child's height and weight may need to be checked often. Tell your doctor if your child is not growing at a normal rate. Common Ritalin side effects may include: sweating, increased blood pressure
- mood changes, anxiety, feeling nervous or irritable, trouble sleeping
- fast heart rate, pounding heartbeats or fluttering in your chest
- loss of appetite, weight loss
- dry mouth, nausea, vomiting, stomach pain, indigestion
- or headache, dizziness .

257. This particular disease, ADHD, interacts with the drug dextroamphetamine and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Dextroamphetamine may cause serious side effects.

Drug Side Effects Summary

Call your doctor at once if you have: signs of heart problems--chest pain, trouble breathing, feeling like you might pass out

- signs of psychosis--hallucinations (seeing or hearing things that are not real), new behavior problems, aggression, hostility, paranoia
- signs of circulation problems--numbness, pain, cold feeling, unexplained wounds, or skin color changes (pale, red, or blue appearance) in your fingers or toes
- a seizure (convulsions)
- muscle twitches (tics)
- or changes in your vision. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Dextroamphetamine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using dextroamphetamine. Common side effects of dextroamphetamine may include: dry mouth, upset stomach , loss of appetite, weight loss
- headache , dizziness
- tremors, fast heartbeats
- or sleep problems (insomnia).

258. This particular disease, ADHD, interacts with the drug bupropion and has the following side effects

- hives, itching, fever, swollen glands, difficult breathing, swelling in your face or throat) or a severe skin reaction (fever, sore throat, burning eyes, skin pain, red or purple skin rash with blistering and peeling). Report any new or worsening symptoms to your doctor , such as: mood or behavior changes, anxiety, depression, panic attacks, trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: a seizure

Drug Side Effects Summary

(convulsions)

- confusion, unusual changes in mood or behavior
- blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights
- fast or irregular heartbeats
- or a manic episode - racing thoughts, increased energy, reckless behavior, feeling extremely happy or irritable, talking more than usual, severe problems with sleep. Common bupropion side effects may include: dry mouth, sore throat, stuffy nose
- ringing in the ears
- blurred vision
- nausea, vomiting, stomach pain, loss of appetite, constipation
- sleep problems (insomnia)
- tremors, sweating, feeling anxious or nervous
- fast heartbeats
- confusion, agitation, hostility
- rash
- weight loss
- increased urination
- headache, dizziness
- or muscle or joint pain.

259. This particular disease, ADHD, interacts with the drug clonidine and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe chest pain, shortness of breath, irregular heartbeats

Drug Side Effects Summary

- a very slow heart rate
- or a light-headed feeling, like you might pass out. Common clonidine side effects may include:
drowsiness, dizziness
- feeling tired or irritable
- dry mouth
- constipation , loss of appetite
- or sleep problems (insomnia), nightmares .

260. This particular disease, ADHD, interacts with the drug Intuniv and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: anxiety ,
nervousness
- hallucinations (especially in children)
- severe drowsiness
- slow heartbeats
- or a light-headed feeling, like you might pass out
- If you stop taking Intuniv, tell your doctor if you have headaches, confusion, rapid heartbeats,
tremors, increased blood pressure, or if you feel nervous or agitated. If left untreated, these
symptoms could lead to very high blood pressure, vision problems, or seizures . Common Intuniv
side effects may include: dizziness , drowsiness
- low blood pressure, slow heartbeats
- feeling tired or irritable
- trouble sleeping
- dry mouth

Drug Side Effects Summary

- or stomach pain, nausea, constipation .

261. This particular disease, ADHD, interacts with the drug Desoxyn and has the following side effects

- hives
 - difficult breathing
 - swelling of your face, lips, tongue, or throat. Desoxyn may cause serious side effects. Call your doctor at once if you have: signs of heart problems--chest pain, trouble breathing, feeling like you might pass out
 - signs of psychosis--hallucinations (seeing or hearing things that are not real), new behavior problems, aggression, hostility, paranoia
 - signs of circulation problems--numbness, pain, cold feeling, unexplained wounds, or skin color changes (pale, red, or blue appearance) in your fingers or toes
 - a seizure (convulsions)
 - muscle twitches (tics)
 - or changes in your vision. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting , or diarrhea . Desoxyn can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using Desoxyn.
- Common side effects of Desoxyn may include: headache or dizziness
- fast heartbeats
 - sleep problems (insomnia)
 - diarrhea, constipation , upset stomach
 - tremors
 - or loss of appetite, weight loss.

Drug Side Effects Summary

262. This particular disease, ADHD, interacts with the drug guanfacine and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Guanfacine may cause serious side effects. Call your doctor at once if you have: anxiety , nervousness
- hallucinations (especially in children)
- severe drowsiness
- slow heartbeats
- or a light-headed feeling, like you might pass out
- If you stop taking guanfacine, tell your doctor if you have headaches, confusion, rapid heartbeats, tremors, increased blood pressure, or if you feel nervous or agitated. If left untreated, these symptoms could lead to very high blood pressure, vision problems, or seizures . Common side effects of guanfacine may include: dizziness , drowsiness
- low blood pressure, slow heartbeats
- feeling tired or irritable
- trouble sleeping
- dry mouth
- or stomach pain, nausea , constipation .

263. This particular disease, ADHD, interacts with the drug Focalin and has the following side effects

- fever
- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: signs of heart problems - chest pain, trouble breathing, feeling like you might pass out

Drug Side Effects Summary

- signs of psychosis - hallucinations (seeing or hearing things that are not real), new thought or behavior problems, aggression, hostility, paranoia
 - signs of circulation problems - numbness , pain, cold feeling, unexplained wounds, or skin color changes (pale, red, or blue appearance) in your fingers or toes
 - or penis erection that is painful or lasts 4 hours or longer. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea .
- Dexmethylphenidate can affect growth in children. Your child's height and weight may need to be checked often. Tell your doctor if your child is not growing at a normal rate. Common Focalin side effects may include: loss of appetite
- nausea, stomach pain
 - or fever.

264. This particular disease, ADHD, interacts with the drug atomoxetine and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), depressed, or have thoughts about suicide or hurting yourself. Atomoxetine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Atomoxetine may cause serious side effects. Call your doctor at once if you have: signs of heart problems--chest pain, trouble breathing, feeling like you might pass out
- signs of psychosis--hallucinations (seeing or hearing things that are not real), new behavior problems, aggression, hostility, paranoia

Drug Side Effects Summary

- liver problems--stomach pain (upper right side), itching, flu-like symptoms, dark urine, jaundice (yellowing of the skin or eyes)
- painful or difficult urination
- or erection is painful or lasts longer than 4 hours (this is a rare side effect). Common side effects of atomoxetine may include: nausea , vomiting , upset stomach , constipation
- dry mouth, loss of appetite
- mood changes, feeling tired
- dizziness
- urination problems
- or impotence , trouble having an erection.

265. This particular disease, ADHD, interacts with the drug Focalin XR and has the following side effects

- fever
- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Focalin XR may cause serious side effects. Call your doctor at once if you have: signs of heart problems--chest pain, trouble breathing, feeling like you might pass out
- signs of psychosis--hallucinations (seeing or hearing things that are not real), new thought or behavior problems, aggression, hostility, paranoia
- signs of circulation problems-- numbness , pain, cold feeling, unexplained wounds, or skin color changes (pale, red, or blue appearance) in your fingers or toes
- or penis erection that is painful or lasts 4 hours or longer. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea .

Drug Side Effects Summary

Focalin XR can affect growth in children. Your child's height and weight may need to be checked often. Tell your doctor if your child is not growing at a normal rate. Common side effects of Focalin XR may include: loss of appetite

- nausea, stomach pain
- or fever.

266. This particular disease, ADHD, interacts with the drug amphetamine and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: signs of heart problems - chest pain, trouble breathing, feeling like you might pass out
- signs of psychosis - hallucinations (seeing or hearing things that are not real), new behavior problems, aggression, hostility, paranoia
- signs of circulation problems - numbness , pain, cold feeling, unexplained wounds, or skin color changes (pale, red, or blue appearance) in your fingers or toes
- a seizure (convulsions)
- muscle twitches (tics)
- pain or burning when you urinate
- or changes in your vision. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea. Amphetamine can affect growth. Tell your doctor if your child is not growing at a normal rate. Common amphetamine side effects may include: increased heart rate
- mood changes, anxiety , feeling restless or nervous
- trouble sleeping

Drug Side Effects Summary

- dry mouth, stomach pain, nausea, vomiting, diarrhea, constipation
- loss of appetite, weight loss
- painful urination
- sexual problems, impotence
- headache , dizziness
- fever, weakness
- or itching.

267. This particular disease, ADHD, interacts with the drug Ritalin LA and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Fast heartbeat Less common Chest pain fever joint pain skin rash or hives Rare Black, tarry stools blood in the urine or stools blurred vision or other changes in vision crusting, dryness, or flaking of the skin muscle cramps pinpoint red spots on the skin scaling, severe redness, soreness, or swelling of the skin seizures uncontrolled vocal outbursts or tics (uncontrolled and repeated body movements) unusual bleeding or bruising Incidence not known Agitation arm, back or jaw pain bleeding gums bloody nose chest discomfort, tightness, or heaviness chills clenching, gnashing, or grinding teeth confusion cough dark urine decreased interest in sexual intercourse diarrhea difficulty with speaking difficulty with swallowing discouragement dizziness double vision drowsiness fainting fast, pounding, or irregular heartbeat or pulse feeling like surroundings are not real feeling sad or empty feeling that others are watching you or controlling your behavior feeling that others can hear your thoughts feeling, seeing, or hearing things that are not there headache heavier menstrual periods inability to have or keep an erection inability to move the arms, legs, or facial muscles inability to speak irritability itching skin large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs,

Drug Side Effects Summary

feet, or genitals loss in sexual ability, desire, drive, or performance loss of interest or pleasure mood swings muscle pain, stiffness, or spasms nausea nervousness numbness of the hands overactive reflexes painful or difficult urination pale skin paleness or cold feeling in the fingertips and toes poor coordination pounding in the ears puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red, irritated eyes red, swollen, or scaly skin redness, soreness, or itching skin restlessness right upper quadrant tenderness seeing, hearing, or feeling things that are not there severe or sudden headache shivering slow speech slowed growth in children sore throat sores, ulcers, or white spots on the lips or in the mouth stuffy or runny nose sudden loss of coordination sudden slurring of speech sweating swollen glands talking or acting with excitement you cannot control tingling or pain in the fingers or toes when exposed to cold temperatures trembling or shaking tremor trouble concentrating trouble sleeping trouble breathing twitching, twisting, or uncontrolled repetitive movements of the tongue, lips, face, arms, or legs unusual behavior unusual tiredness or weakness weight loss yellow skin or eyes Symptoms of overdose Agitation anxiety bigger, dilated, or enlarged pupils of the eyes confusion as to time, place, or person dark urine diarrhea dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position dry eyes, mouth, nose, or throat false or unusual sense of well-being fast, slow, irregular, pounding, or racing heartbeat or pulse holding false beliefs that cannot be changed by fact increased sensitivity of the eyes to light loss of consciousness muscle pain, cramps, stiffness, or twitching nervousness overactive reflexes pounding in the ears rapid, shallow breathing sweating tremors unusual excitement, nervousness, or restlessness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Loss of appetite stomach pain Less common Acid or sour stomach anger belching fear

Drug Side Effects Summary

heartburn indigestion irritability scalp hair loss stomach discomfort or upset vomiting Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Ritalin LA side effects

268. This particular disease, ADHD, interacts with the drug Kapvay and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe chest pain, shortness of breath, irregular heartbeats
- a very slow heart rate
- or a light-headed feeling, like you might pass out. Common Kapvay side effects may include: drowsiness, dizziness
- feeling tired or irritable
- dry mouth
- constipation , loss of appetite
- or sleep problems (insomnia), nightmares .

269. This particular disease, ADHD, interacts with the drug Methylin and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Fast heartbeat Less common Chest pain fever joint pain skin rash or hives Rare Black, tarry stools blood in the urine or stools blurred vision or other changes in vision crusting, dryness, or flaking of the skin muscle cramps pinpoint red spots on the skin scaling, severe redness, soreness, or swelling of the skin

Drug Side Effects Summary

seizures uncontrolled vocal outbursts or tics (uncontrolled and repeated body movements) unusual bleeding or bruising Incidence not known Agitation arm, back or jaw pain bleeding gums bloody nose chest discomfort, tightness, or heaviness chills clenching, gnashing, or grinding teeth confusion cough dark urine decreased interest in sexual intercourse diarrhea difficulty with speaking difficulty with swallowing discouragement dizziness double vision drowsiness fainting fast, pounding, or irregular heartbeat or pulse feeling like surroundings are not real feeling sad or empty feeling that others are watching you or controlling your behavior feeling that others can hear your thoughts feeling, seeing, or hearing things that are not there headache heavier menstrual periods inability to have or keep an erection inability to move the arms, legs, or facial muscles inability to speak irritability itching skin large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or genitals loss in sexual ability, desire, drive, or performance loss of interest or pleasure mood swings muscle pain, stiffness, or spasms nausea nervousness numbness of the hands overactive reflexes painful or difficult urination pale skin paleness or cold feeling in the fingertips and toes poor coordination pounding in the ears puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red, irritated eyes red, swollen, or scaly skin redness, soreness, or itching skin restlessness right upper quadrant tenderness seeing, hearing, or feeling things that are not there severe or sudden headache shivering slow speech slowed growth in children sore throat sores, ulcers, or white spots on the lips or in the mouth stuffy or runny nose sudden loss of coordination sudden slurring of speech sweating swollen glands talking or acting with excitement you cannot control tingling or pain in the fingers or toes when exposed to cold temperatures trembling or shaking tremor trouble concentrating trouble sleeping trouble breathing twitching, twisting, or uncontrolled repetitive movements of the tongue, lips, face, arms, or legs unusual behavior unusual tiredness or weakness weight loss yellow skin or eyes Symptoms of overdose Agitation anxiety bigger, dilated, or enlarged pupils of the eyes confusion as to time, place, or person dark urine diarrhea dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting

Drug Side Effects Summary

position dry eyes, mouth, nose, or throat false or unusual sense of well-being fast, slow, irregular, pounding, or racing heartbeat or pulse holding false beliefs that cannot be changed by fact increased sensitivity of the eyes to light loss of consciousness muscle pain, cramps, stiffness, or twitching nervousness overactive reflexes pounding in the ears rapid, shallow breathing sweating tremors unusual excitement, nervousness, or restlessness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Loss of appetite stomach pain Less common Acid or sour stomach anger belching fear heartburn indigestion irritability scalp hair loss stomach discomfort or upset vomiting Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Methylin side effects

270. This particular disease, ADHD, interacts with the drug dexamethylphenidate and has the following

- fever
- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Dexamethylphenidate may cause serious side effects. Call your doctor at once if you have: signs of heart problems--chest pain, trouble breathing, feeling like you might pass out
- signs of psychosis--hallucinations (seeing or hearing things that are not real), new thought or behavior problems, aggression, hostility, paranoia
- signs of circulation problems-- numbness , pain, cold feeling, unexplained wounds, or skin color

Drug Side Effects Summary

changes (pale, red, or blue appearance) in your fingers or toes

- or penis erection that is painful or lasts 4 hours or longer. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea .

Dexmethylphenidate can affect growth in children. Your child's height and weight may need to be checked often. Tell your doctor if your child is not growing at a normal rate. Common side effects of dexmethylphenidate may include: loss of appetite

- nausea, stomach pain

- or fever.

271. This particular disease, ADHD, interacts with the drug methamphetamine and has the following

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: signs of heart problems - chest pain, trouble breathing, feeling like you might pass out

- signs of psychosis - hallucinations (seeing or hearing things that are not real), new behavior problems, aggression, hostility, paranoia

- signs of circulation problems - numbness, pain, cold feeling, unexplained wounds, or skin color changes (pale, red, or blue appearance) in your fingers or toes

- a seizure (convulsions)

- muscle twitches (tics)

- or changes in your vision. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Methamphetamine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this

Drug Side Effects Summary

medicine. Common methamphetamine side effects may include: headache or dizziness

- fast heartbeats
- sleep problems (insomnia)
- diarrhea, constipation, upset stomach
- tremors
- or loss of appetite, weight loss.

272. This particular disease, ADHD, interacts with the drug modafinil and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Modafinil can cause skin reactions that may be severe enough to need treatment in a hospital. Stop taking this medicine and get emergency medical help if you have: skin rash or hives, blisters or peeling
- mouth sores, trouble swallowing
- fever, shortness of breath
- swelling in your legs
- dark urine, jaundice (yellowing of the skin or eyes)
- or swelling in your face, eyes, lips, tongue, or throat. Modafinil may cause serious side effects. Stop using modafinil and call your doctor at once if you have: depression , anxiety , suicidal thoughts or actions
- hallucinations, unusual thoughts or behavior, aggression, being more active or talkative than usual
- chest pain, trouble breathing, uneven heart beats
- or the first sign of any skin rash, no matter how minor you think it might be. Common side effects of modafinil may include: headache , dizziness
- feeling nervous or anxious

Drug Side Effects Summary

- back pain
- nausea , diarrhea , upset stomach
- sleep problems (insomnia)
- or stuffy nose.

273. This particular disease, ADHD, interacts with the drug Metadate CD and has the following side effects:

- No side effects listed

274. This particular disease, ADHD, interacts with the drug Methylin ER and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Fast heartbeat Less common Chest pain fever joint pain skin rash or hives Rare Black, tarry stools blood in the urine or stools blurred vision or other changes in vision crusting, dryness, or flaking of the skin muscle cramps pinpoint red spots on the skin scaling, severe redness, soreness, or swelling of the skin seizures uncontrolled vocal outbursts or tics (uncontrolled and repeated body movements) unusual bleeding or bruising Incidence not known Agitation arm, back or jaw pain bleeding gums bloody nose chest discomfort, tightness, or heaviness chills clenching, gnashing, or grinding teeth confusion cough dark urine decreased interest in sexual intercourse diarrhea difficulty with speaking difficulty with swallowing discouragement dizziness double vision drowsiness fainting fast, pounding, or irregular heartbeat or pulse feeling like surroundings are not real feeling sad or empty feeling that others are watching you or controlling your behavior feeling that others can hear your thoughts feeling, seeing, or hearing things that are not there headache heavier menstrual periods inability to have or keep an erection inability to move the arms, legs, or facial muscles inability to speak irritability itching skin large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs,

Drug Side Effects Summary

feet, or genitals loss in sexual ability, desire, drive, or performance loss of interest or pleasure mood swings muscle pain, stiffness, or spasms nausea nervousness numbness of the hands overactive reflexes painful or difficult urination pale skin paleness or cold feeling in the fingertips and toes poor coordination pounding in the ears puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red, irritated eyes red, swollen, or scaly skin redness, soreness, or itching skin restlessness right upper quadrant tenderness seeing, hearing, or feeling things that are not there severe or sudden headache shivering slow speech slowed growth in children sore throat sores, ulcers, or white spots on the lips or in the mouth stuffy or runny nose sudden loss of coordination sudden slurring of speech sweating swollen glands talking or acting with excitement you cannot control tingling or pain in the fingers or toes when exposed to cold temperatures trembling or shaking tremor trouble concentrating trouble sleeping trouble breathing twitching, twisting, or uncontrolled repetitive movements of the tongue, lips, face, arms, or legs unusual behavior unusual tiredness or weakness weight loss yellow skin or eyes Symptoms of overdose Agitation anxiety bigger, dilated, or enlarged pupils of the eyes confusion as to time, place, or person dark urine diarrhea dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position dry eyes, mouth, nose, or throat false or unusual sense of well-being fast, slow, irregular, pounding, or racing heartbeat or pulse holding false beliefs that cannot be changed by fact increased sensitivity of the eyes to light loss of consciousness muscle pain, cramps, stiffness, or twitching nervousness overactive reflexes pounding in the ears rapid, shallow breathing sweating tremors unusual excitement, nervousness, or restlessness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Loss of appetite stomach pain Less common Acid or sour stomach anger belching fear

Drug Side Effects Summary

heartburn indigestion irritability scalp hair loss stomach discomfort or upset vomiting Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Methylin ER side effects

275. This particular disease, ADHD, interacts with the drug omega-3 polyunsaturated fatty acids and

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Omega-3 polyunsaturated fatty acids may cause serious side effects. Call your doctor at once if you have: chest pain
- or uneven heartbeats. Common side effects of omega-3 polyunsaturated fatty acids may include: loss of appetite
- diarrhea , constipation , upset stomach , belching
- back pain
- or dry mouth, altered sense of taste.

276. This particular disease, ADHD, interacts with the drug ProCentra and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. ProCentra may cause serious side effects. Call your doctor at once if you have: signs of heart problems--chest pain, trouble breathing, feeling like you might pass out
- signs of psychosis--hallucinations (seeing or hearing things that are not real), new behavior problems, aggression, hostility, paranoia
- signs of circulation problems--numbness, pain, cold feeling, unexplained wounds, or skin color

Drug Side Effects Summary

changes (pale, red, or blue appearance) in your fingers or toes

- a seizure (convulsions)
- muscle twitches (tics)
- or changes in your vision. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. ProCentra can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using ProCentra. Common side effects of ProCentra may include: dry mouth, upset stomach , loss of appetite, weight loss
- headache , dizziness
- tremors, fast heartbeats
- or sleep problems (insomnia).

277. This particular disease, ADHD, interacts with the drug selegiline and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Selegiline may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- trouble breathing
- confusion, hallucinations, unusual thoughts or behavior
- increased tremors or uncontrolled muscle movements
- worsening side effects of your other medications
- high levels of serotonin in the body (when taken with an antidepressant)-- agitation , fever, fast heart rate , overactive reflexes, nausea , vomiting , diarrhea , loss of coordination, fainting
- or dangerously high blood pressure--severe headache, blurred vision, pounding in your neck or

Drug Side Effects Summary

ears, anxiety , nausea, vomiting, severe chest pain, shortness of breath, pounding heartbeats, or seizure (convulsions). You may have increased sexual urges, unusual urges to gamble, or other intense urges while taking this medicine. Talk with your doctor if this occurs. Common side effects of selegiline may include: dizziness

- nausea, stomach pain, constipation
- skin rash or other irritation
- sleep problems (insomnia)
- or mouth sores or ulcers, pain with swallowing (while using selegiline orally disintegrating tablets).

278. This particular disease, ADHD, interacts with the drug Zenzedi and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Zenzedi may cause serious side effects. Call your doctor at once if you have: signs of heart problems--chest pain, trouble breathing, feeling like you might pass out
- signs of psychosis--hallucinations (seeing or hearing things that are not real), new behavior problems, aggression, hostility, paranoia
- signs of circulation problems--numbness, pain, cold feeling, unexplained wounds, or skin color changes (pale, red, or blue appearance) in your fingers or toes
- a seizure (convulsions)
- muscle twitches (tics)
- or changes in your vision. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Zenzedi can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using Zenzedi. Common

Drug Side Effects Summary

side effects of Zenzedi may include: dry mouth, upset stomach , loss of appetite, weight loss

- headache , dizziness
- tremors, fast heartbeats
- or sleep problems (insomnia).

279. This particular disease, ADHD, interacts with the drug Adhansia XR and has the following side

- No side effects listed

280. This particular disease, ADHD, interacts with the drug Aptensio XR and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Fast heartbeat Less common Chest pain fever joint pain skin rash or hives Rare Black, tarry stools blood in the urine or stools blurred vision or other changes in vision crusting, dryness, or flaking of the skin muscle cramps pinpoint red spots on the skin scaling, severe redness, soreness, or swelling of the skin seizures uncontrolled vocal outbursts or tics (uncontrolled and repeated body movements) unusual bleeding or bruising Incidence not known Agitation arm, back or jaw pain bleeding gums bloody nose chest discomfort, tightness, or heaviness chills clenching, gnashing, or grinding teeth confusion cough dark urine decreased interest in sexual intercourse diarrhea difficulty with speaking difficulty with swallowing discouragement dizziness double vision drowsiness fainting fast, pounding, or irregular heartbeat or pulse feeling like surroundings are not real feeling sad or empty feeling that others are watching you or controlling your behavior feeling that others can hear your thoughts feeling, seeing, or hearing things that are not there headache heavier menstrual periods inability to have or keep an erection inability to move the arms, legs, or facial muscles inability to speak irritability itching skin large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs,

Drug Side Effects Summary

feet, or genitals loss in sexual ability, desire, drive, or performance loss of interest or pleasure mood swings muscle pain, stiffness, or spasms nausea nervousness numbness of the hands overactive reflexes painful or difficult urination pale skin paleness or cold feeling in the fingertips and toes poor coordination pounding in the ears puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red, irritated eyes red, swollen, or scaly skin redness, soreness, or itching skin restlessness right upper quadrant tenderness seeing, hearing, or feeling things that are not there severe or sudden headache shivering slow speech slowed growth in children sore throat sores, ulcers, or white spots on the lips or in the mouth stuffy or runny nose sudden loss of coordination sudden slurring of speech sweating swollen glands talking or acting with excitement you cannot control tingling or pain in the fingers or toes when exposed to cold temperatures trembling or shaking tremor trouble concentrating trouble sleeping trouble breathing twitching, twisting, or uncontrolled repetitive movements of the tongue, lips, face, arms, or legs unusual behavior unusual tiredness or weakness weight loss yellow skin or eyes Symptoms of overdose Agitation anxiety bigger, dilated, or enlarged pupils of the eyes confusion as to time, place, or person dark urine diarrhea dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position dry eyes, mouth, nose, or throat false or unusual sense of well-being fast, slow, irregular, pounding, or racing heartbeat or pulse holding false beliefs that cannot be changed by fact increased sensitivity of the eyes to light loss of consciousness muscle pain, cramps, stiffness, or twitching nervousness overactive reflexes pounding in the ears rapid, shallow breathing sweating tremors unusual excitement, nervousness, or restlessness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Loss of appetite stomach pain Less common Acid or sour stomach anger belching fear

Drug Side Effects Summary

heartburn indigestion irritability scalp hair loss stomach discomfort or upset vomiting Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Aptensio XR side effects

281. This particular disease, ADHD, interacts with the drug armodafinil and has the following side effects

- skin rash, fever, swollen glands, flu-like symptoms, unusual bruising, or jaundice (yellowing of your skin or eyes). Armodafinil may cause serious side effects. Stop using armodafinil and call your doctor at once if you have: bruising, severe tingling, numbness , pain, muscle weakness
- unusual bleeding (nosebleeds , bleeding gums)
- skin sores or blistering
- mouth sores, trouble swallowing
- chest pain, uneven heart beats
- or depression , anxiety , hallucinations, aggression, unusual thoughts or behavior, suicidal thoughts. Common side effects of armodafinil may include: headache , dizziness
- nausea
- or sleep problems (insomnia).

282. This particular disease, ADHD, interacts with the drug desvenlafaxine and has the following side effects

- skin rash or hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Tell your doctor right away if you have new or sudden changes in mood or behavior, including new or worse depression or anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, more active or talkative, or have thoughts about suicide or hurting yourself. Desvenlafaxine may cause serious side

Drug Side Effects Summary

effects. Call your doctor at once if you have: a seizure

- easy bruising or bleeding (nosebleeds , bleeding gums), blood in your urine or stools, coughing up blood
- blurred vision, eye pain or swelling, or seeing halos around lights
- cough, chest discomfort, trouble breathing
- or low blood sodium-- headache , confusion, problems with thinking or memory, weakness, feeling unsteady. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting , or diarrhea . Common side effects of desvenlafaxine may include: dizziness , drowsiness, anxiety
- increased sweating
- nausea, decreased appetite, constipation
- sleep problems (insomnia)
- or decreased sex drive, impotence , or difficulty having an orgasm.

283. This particular disease, ADHD, interacts with the drug Evekeo and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Evekeo may cause serious side effects. Call your doctor at once if you have: signs of heart problems--chest pain, trouble breathing, feeling like you might pass out
- signs of psychosis--hallucinations (seeing or hearing things that are not real), new behavior problems, aggression, hostility, paranoia
- signs of circulation problems-- numbness , pain, cold feeling, unexplained wounds, or skin color changes (pale, red, or blue appearance) in your fingers or toes

Drug Side Effects Summary

- a seizure (convulsions)
- muscle twitches (tics)
- pain or burning when you urinate
- or changes in your vision. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea. Evekeo can affect growth. Tell your doctor if your child is not growing at a normal rate. Common side effects of Evekeo may include: increased heart rate
- mood changes, anxiety , feeling restless or nervous
- trouble sleeping
- dry mouth, stomach pain, nausea, vomiting, diarrhea, constipation
- loss of appetite, weight loss
- painful urination
- sexual problems, impotence
- headache , dizziness
- fever, weakness
- or itching.

284. This particular disease, ADHD, interacts with the drug Mydayis and has the following side effects

- Mydayis can cause serious side effects, including: See Important information Slowing of growth (height and weight) in children. Children should have their height and weight checked often during treatment. Your healthcare provider may stop your child's treatment if they are not growing or gaining weight as expected. Circulation problems in fingers and toes (peripheral vasculopathy, including Raynaud's phenomenon). Signs and symptoms may include: fingers or toes may feel numb, cool, painful fingers or toes may change color from pale, to blue, to red Tell your healthcare

Drug Side Effects Summary

provider if you have or your child has any numbness, pain, skin color change, or sensitivity to temperature in your fingers or toes. Call your healthcare provider if you or your child have any signs of unexplained wounds appearing on fingers or toes during treatment. Seizures. Your healthcare provider will stop treatment if you have a seizure. Serotonin syndrome. This problem may happen when Mydayis is taken with certain other medicines and may be life threatening. Call your healthcare provider or go to the nearest hospital emergency room if you get symptoms of serotonin syndrome which may include: agitation, hallucinations, coma, or other changes in mental status sweating or fever problems controlling movements or muscle twitching nausea, vomiting, or diarrhea fast heartbeat muscle stiffness or tightness The most common side effects include: trouble sleeping increased heart rate irritability decreased appetite anxiety weight loss dry mouth nausea These are not all the possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Mydayis side effects

285. This particular disease, ADHD, interacts with the drug Relexxii and has the following side effects

- No side effects listed

286. This particular disease, ADHD, interacts with the drug Adzenys ER and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Adzenys ER may cause serious side effects. Call your doctor at once if you have: signs of heart problems--chest pain, trouble breathing, feeling like you might pass out
- signs of psychosis--hallucinations (seeing or hearing things that are not real), new behavior problems, aggression, hostility, paranoia
- signs of circulation problems-- numbness , pain, cold feeling, unexplained wounds, or skin color

Drug Side Effects Summary

changes (pale, red, or blue appearance) in your fingers or toes

- a seizure (convulsions)
- muscle twitches (tics)
- pain or burning when you urinate
- or changes in your vision. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea. Adzenys ER can affect growth. Tell your doctor if your child is not growing at a normal rate. Common side effects of Adzenys ER may include: increased heart rate
- mood changes, anxiety , feeling restless or nervous
- trouble sleeping
- dry mouth, stomach pain, nausea, vomiting, diarrhea, constipation
- loss of appetite, weight loss
- painful urination
- sexual problems, impotence
- headache , dizziness
- fever, weakness
- or itching.

287. This particular disease, ADHD, interacts with the drug Adzenys XR-ODT and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Adzenys XR-ODT may cause serious side effects. Call your doctor at once if you have: signs of heart problems--chest pain, trouble breathing, feeling like you might pass out

Drug Side Effects Summary

- signs of psychosis--hallucinations (seeing or hearing things that are not real), new behavior problems, aggression, hostility, paranoia
- signs of circulation problems-- numbness , pain, cold feeling, unexplained wounds, or skin color changes (pale, red, or blue appearance) in your fingers or toes
- a seizure (convulsions)
- muscle twitches (tics)
- pain or burning when you urinate
- or changes in your vision. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea. Adzenys XR-ODT can affect growth. Tell your doctor if your child is not growing at a normal rate. Common side effects of Adzenys XR-ODT may include: increased heart rate
- mood changes, anxiety , feeling restless or nervous
- trouble sleeping
- dry mouth, stomach pain, nausea, vomiting, diarrhea, constipation
- loss of appetite, weight loss
- painful urination
- sexual problems, impotence
- headache , dizziness
- fever, weakness
- or itching.

288. This particular disease, ADHD, interacts with the drug desipramine and has the following side

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Desipramine may cause serious side effects. Call your doctor at once if you have: easy bruising, unusual bleeding, purple or red spots under your skin

- a light-headed feeling, like you might pass out

- confusion, hallucinations, unusual thoughts or behavior

- blurred vision, tunnel vision, eye pain or swelling, seeing halos around lights

- new or worsening chest pain, pounding heartbeats or fluttering in your chest

- sudden numbness or weakness, problems with vision, speech, or balance

- fever, sore throat

- easy bruising, unusual bleeding (nose, mouth, vagina, or rectum)

- painful or difficult urination

- upper stomach pain, loss of appetite, jaundice (yellowing of the skin or eyes)

- a seizure

- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting , or diarrhea . Serious side effects may be more likely in older adults and those who are ill or debilitated.

Common side effects of desipramine may include: increased blood pressure

- tingly feeling, weakness, lack of coordination

- dry mouth, loss of appetite, nausea, vomiting

- breast swelling (in men or women)

Drug Side Effects Summary

- or weight gain or weight loss .

289. This particular disease, ADHD, interacts with the drug Dyanavel XR and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: signs of heart problems - chest pain, trouble breathing, feeling like you might pass out
- signs of psychosis - hallucinations (seeing or hearing things that are not real), new behavior problems, aggression, hostility, paranoia
- signs of circulation problems - numbness , pain, cold feeling, unexplained wounds, or skin color changes (pale, red, or blue appearance) in your fingers or toes
- a seizure (convulsions)
- muscle twitches (tics)
- pain or burning when you urinate
- or changes in your vision. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea. Amphetamine can affect growth. Tell your doctor if your child is not growing at a normal rate. Common Dyanavel XR side effects may include: increased heart rate
- mood changes, anxiety , feeling restless or nervous
- trouble sleeping
- dry mouth, stomach pain, nausea, vomiting, diarrhea, constipation
- loss of appetite, weight loss
- painful urination
- sexual problems, impotence

Drug Side Effects Summary

- headache , dizziness
- fever, weakness
- or itching.

290. This particular disease, ADHD, interacts with the drug Evekeo ODT and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Evekeo ODT may cause serious side effects. Call your doctor at once if you have: signs of heart problems--chest pain, trouble breathing, feeling like you might pass out
- signs of psychosis--hallucinations (seeing or hearing things that are not real), new behavior problems, aggression, hostility, paranoia
- signs of circulation problems-- numbness , pain, cold feeling, unexplained wounds, or skin color changes (pale, red, or blue appearance) in your fingers or toes
- a seizure (convulsions)
- muscle twitches (tics)
- pain or burning when you urinate
- or changes in your vision. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea. Evekeo ODT can affect growth. Tell your doctor if your child is not growing at a normal rate. Common side effects of Evekeo ODT may include: increased heart rate
- mood changes, anxiety , feeling restless or nervous
- trouble sleeping
- dry mouth, stomach pain, nausea, vomiting, diarrhea, constipation

Drug Side Effects Summary

- loss of appetite, weight loss
- painful urination
- sexual problems, impotence
- headache , dizziness
- fever, weakness
- or itching.

291. This particular disease, ADHD, interacts with the drug imipramine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Imipramine may cause serious side effects. Call your doctor at once if you have: easy bruising, unusual bleeding, purple or red spots under your skin
- tunnel vision, eye pain or swelling, or seeing halos around lights
- a light-headed feeling, like you might pass out
- new or worsening chest pain, pounding heartbeats or fluttering in your chest
- sudden numbness or weakness, problems with vision, speech, or balance
- fever, sore throat
- confusion, hallucinations, unusual thoughts or behavior
- painful or difficult urination
- seizure (convulsions)
- or jaundice (yellowing of the skin or eyes). Common side effects of imipramine may include:

Drug Side Effects Summary

increased blood pressure

- tingly feeling, weakness, lack of coordination
- dry mouth, nausea , vomiting
- or breast swelling (in men or women).

292. This particular disease, ADHD, interacts with the drug dexamethylphenidate / serdexmethylphenidate

- fever
- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Dexamethylphenidate and serdexmethylphenidate may cause serious side effects. Call your doctor at once if you have: signs of heart problems--chest pain, trouble breathing, feeling like you might pass out
- signs of psychosis--hallucinations (seeing or hearing things that are not real), new thought or behavior problems, aggression, hostility, paranoia
- signs of circulation problems-- numbness , pain, cold feeling, unexplained wounds, or skin color changes (pale, red, or blue appearance) in your fingers or toes
- or penis erection that is painful or lasts 4 hours or longer. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea. Azstarys can affect growth in children. Your child's height and weight may need to be checked often. Tell your doctor if your child is not growing at a normal rate. Common side effects of dexamethylphenidate and serdexmethylphenidate may include: loss of appetite, weight loss
- indigestion , stomach pain, nausea, vomiting
- dizziness , fast heart rate, increased blood pressure
- mood changes, feeling anxious or irritable

Drug Side Effects Summary

- sleep problems (insomnia)
- or fever.

293. This particular disease, ADHD, interacts with the drug viloxazine and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have symptoms of a manic episode : racing thoughts
- increased energy, decreased need for sleep
- risk-taking behavior
- or being agitated or talkative. Viloxazine can affect weight in children. Tell your doctor if your child loses weight or is not gaining weight at a normal rate. Common side effects of viloxazine may include: drowsiness
- feeling tired or irritable
- trouble sleeping
- or nausea , vomiting , loss of appetite.

294. This particular disease, AIDS/HIV, interacts with the drug Truvada and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Mild symptoms of lactic acidosis may worsen over time , and this condition can be fatal. Get emergency medical help if you have: unusual muscle pain,

Drug Side Effects Summary

trouble breathing, stomach pain, vomiting , irregular heart rate, dizziness, feeling cold, or feeling very weak or tired. Call your doctor at once if you have: symptoms of new HIV infection - fever, night sweats, tiredness, muscle or joint pain, sore throat, vomiting, diarrhea, rash, swollen glands in your neck or groin

- sudden or unusual bone pain

- kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath

- or liver problems - nausea , swelling around your midsection, upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Emtricitabine and tenofovir disoproxil fumarate affects your immune system, which may cause certain side effects (even weeks or months after you've taken Truvada). Tell your doctor if you have: signs of a new infection - fever, sweats, swollen glands, cold sores , cough, wheezing, diarrhea, weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Truvada side effects may include: headache , dizziness, feeling depressed or tired

- trouble sleeping, strange dreams

- nausea, stomach pain

- weight loss

- or rash.

295. This particular disease, AIDS/HIV, interacts with the drug Biktarvy and has the following side effects

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: kidney problems -

Drug Side Effects Summary

little or no urination, swelling in your feet or ankles, feeling tired or short of breath

- lactic acidosis - muscle pain or weakness, numbness or cold feeling, trouble breathing, stomach pain, vomiting , irregular heart rate, dizziness , or feeling very weak or tired

- or liver problems - swelling around your midsection, upper stomach pain, unusual tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Biktarvy affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Biktarvy side effects may include: nausea , diarrhea

- or headache .

296. This particular disease, AIDS/HIV, interacts with the drug Triumeq and has the following side effects

- Call your doctor at once if you have symptoms of an allergic reaction from two or more of these specific side effect groups: Group 1 - fever

- Group 2 - rash

- Group 3 - nausea, vomiting, diarrhea, stomach pain

- Group 4 - general ill feeling, extreme tiredness, body aches

- Group 5 - shortness of breath, cough, sore throat. Once you have had an allergic reaction to a medicine that contains abacavir or dolutegravir, you must never use it again. If you stop taking Triumeq for any reason, talk to your doctor before you start taking it again. Call your doctor at once if you have: other signs of allergic reaction - skin blisters or peeling, eye redness , swelling in your face or throat, trouble breathing

Drug Side Effects Summary

- lactic acidosis - unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness , feeling cold, or feeling very weak or tired
- or liver problems - swelling around your midsection, right-sided upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Triumeq affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea, weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Triumeq side effects may include: headache
- tiredness
- or trouble sleeping.

297. This particular disease, AIDS/HIV, interacts with the drug Genvoya and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: kidney problems - swelling, urinating less, feeling tired or short of breath
- lactic acidosis - unusual muscle pain, trouble breathing, stomach pain, vomiting , irregular heart rate, dizziness , feeling cold, or feeling very weak or tired
- or liver problems - nausea , upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Antiviral medicine can affect your immune system (even weeks or months after you've used Genvoya). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss

Drug Side Effects Summary

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Genvoya side effects may include: nausea
- or diarrhea.

298. This particular disease, AIDS/HIV, interacts with the drug Stribild and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: kidney problems - swelling, urinating less, feeling tired or short of breath
- lactic acidosis - unusual muscle pain, trouble breathing, stomach pain, vomiting , irregular heart rate, dizziness , feeling cold, or feeling very weak or tired
- or liver problems - nausea , upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Antiviral medicine can affect your immune system (even weeks or months after you've used Stribild). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Stribild side effects may include: nausea
- or diarrhea.

299. This particular disease, AIDS/HIV, interacts with the drug Atripla and has the following side effects

- unusual muscle pain, trouble breathing, stomach pain, vomiting , irregular heart rate, dizziness,

Drug Side Effects Summary

feeling cold, or feeling very weak or tired. Call your doctor at once if you have: unusual thoughts or behavior, anger, severe depression , thoughts of hurting yourself or others, hallucinations

- a seizure (convulsions)

- kidney problems - increased thirst and urination, muscle pain or weakness

- or liver problems - swelling around your midsection, right-sided upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Atripla affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Atripla side effects may include: dizziness, drowsiness, tired feeling

- nausea , diarrhea

- headache , depressed mood, trouble concentrating

- sleep problems (insomnia), strange dreams

- rash

- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

300. This particular disease, AIDS/HIV, interacts with the drug Descovy and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

Drug Side Effects Summary

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of too much lactic acid in the blood (lactic acidosis) like fast breathing, fast heartbeat , a heartbeat that does not feel normal, very bad upset stomach or throwing up, feeling very sleepy, shortness of breath, feeling very tired or weak, very bad dizziness , feeling cold, or muscle pain or cramps. Changes in your immune system can happen when you start taking drugs to treat HIV. If you have an infection that you did not know you had, it may show up when you take Descovy. Tell your doctor right away if you have any new signs after you start Descovy, even after taking it for several months. This includes signs of infection like fever, sore throat , weakness, cough, or shortness of breath. Descovy side effects

301. This particular disease, AIDS/HIV, interacts with the drug Norvir and has the following side effects

- irregular heartbeats, or a light-headed feeling (like you might pass out)
- easy bruising, unusual bleeding (nose, mouth, vagina, or rectum), purple or red pinpoint spots under your skin
- signs of a kidney stone - pain in your side or lower back, blood in your urine, painful or difficult urination

Drug Side Effects Summary

- high blood sugar - increased thirst, increased urination, hunger, dry mouth, fruity breath odor
- or signs of liver or pancreas problems - loss of appetite, upper stomach pain (that may spread to your back), nausea , vomiting , dark urine, jaundice (yellowing of the skin or eyes). Norvir affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Norvir side effects may include: nausea, vomiting, stomach pain, diarrhea
- numbness or tingling in your hands or feet or around your mouth
- feeling weak or tired
- rash
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

302. This particular disease, AIDS/HIV, interacts with the drug Isentress and has the following side

- Stop taking this medicine and get emergency medical help if you have signs of an allergic reaction to Isentress: fever, general ill feeling, tiredness, joint or muscle pain, trouble breathing
- upper stomach pain, vomiting, loss of appetite, dark urine, yellowing of the skin or eyes
- burning eyes, blisters or mouth sores
- rash, hives , blistering or peeling skin
- swelling of your face, lips, tongue, or throat. In rare cases, Isentress can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure . Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have

Drug Side Effects Summary

fever, unusual tiredness, or dark colored urine. Isentress affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Isentress side effects may include: nausea

- headache , dizziness

- tired feeling

- or sleep problems (insomnia).

303. This particular disease, AIDS/HIV, interacts with the drug Dovato and has the following side effects

- Stop taking this medicine and get emergency medical help if you have signs of an allergic reaction to Dovato: fever, general ill feeling, trouble breathing, tiredness

- joint or muscle pain, blisters or mouth sores, redness or swelling in your eyes

- blistering or peeling skin

- swelling of your face, lips, tongue, or throat. Mild symptoms of lactic acidosis may worsen over time , and this condition can be fatal. unusual muscle pain, trouble breathing, stomach pain, vomiting, fast or irregular heart rate, dizziness, feeling cold, or feeling very weak or tired. Call your doctor at once if you have: the first sign of any skin rash , no matter how mild

- or liver problems - nausea , vomiting , loss of appetite, upper stomach pain, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Dolutegravir and lamivudine affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen

Drug Side Effects Summary

glands, cold sores , cough, wheezing, diarrhea , weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Dovato side effects may include: headache

- tiredness

- nausea, diarrhea

- or sleep problems (insomnia).

304. This particular disease, AIDS/HIV, interacts with the drug Tivicay and has the following side effects

- Stop taking this medicine and get emergency medical help if you have signs of an allergic reaction to Tivicay: fever, general ill feeling, trouble breathing, tiredness

- joint or muscle pain, blisters or mouth sores, redness or swelling in your eyes

- blistering or peeling skin

- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: the first sign of any skin rash , no matter how mild

- or liver problems - nausea , vomiting , loss of appetite, upper stomach pain, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Dolutegravir affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Tivicay side effects may include: headache

Drug Side Effects Summary

- tiredness
- or sleep problems (insomnia).

305. This particular disease, AIDS/HIV, interacts with the drug abacavir / dolutegravir / lamivudine a

- Call your doctor at once if you have symptoms of an allergic reaction from two or more of these specific side effect groups: Group 1 - fever

- Group 2 - rash

- Group 3 - nausea, vomiting, diarrhea, stomach pain

- Group 4 - general ill feeling, extreme tiredness, body aches

- Group 5 - shortness of breath, cough, sore throat. Once you have had an allergic reaction to a medicine that contains abacavir or dolutegravir, you must never use it again. If you stop taking abacavir, dolutegravir, and lamivudine for any reason, talk to your doctor before you start taking it again. Abacavir, dolutegravir, and lamivudine may cause serious side effects. Call your doctor at once if you have: other signs of allergic reaction--skin blisters or peeling, eye redness , swelling in your face or throat, trouble breathing

- lactic acidosis --unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness , feeling cold, or feeling very weak or tired

- or liver problems--swelling around your midsection, right-sided upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Abacavir, dolutegravir, and lamivudine affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea, weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

Drug Side Effects Summary

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of abacavir, dolutegravir, and lamivudine may include: headache
- tiredness
- or trouble sleeping.

306. This particular disease, AIDS/HIV, interacts with the drug Prezista and has the following side effects

- a skin rash, no matter how mild
- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- or signs of liver or pancreas problems - loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting , fast heart rate , dark urine, jaundice (yellowing of the skin or eyes).
Darunavir affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Prezista side effects may include: nausea, vomiting, diarrhea, stomach pain
- headache
- rash
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

307. This particular disease, AIDS/HIV, interacts with the drug bictegravir / emtricitabine / tenofovir

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Bictegravir, emtricitabine, and tenofovir may cause serious side effects. Call your doctor at once if you have: kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- lactic acidosis --muscle pain or weakness, numbness or cold feeling, trouble breathing, stomach pain, vomiting , irregular heart rate, dizziness , or feeling very weak or tired
- or liver problems--swelling around your midsection, upper stomach pain, unusual tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Biktarvy affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of bictegravir, emtricitabine, and tenofovir may include: nausea , diarrhea
- or headache .

308. This particular disease, AIDS/HIV, interacts with the drug cobicistat / elvitegravir / emtricitabine

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Cobicistat, elvitegravir, emtricitabine, and tenofovir may cause serious side effects. Call your doctor at once if you have: kidney problems--swelling, urinating less, feeling tired or short of breath
- lactic acidosis --unusual muscle pain, trouble breathing, stomach pain, vomiting , irregular heart rate, dizziness , feeling cold, or feeling very weak or tired
- or liver problems-- nausea , upper stomach pain, loss of appetite, dark urine, clay-colored stools,

Drug Side Effects Summary

jaundice (yellowing of the skin or eyes). Antiviral medicine can affect your immune system (even weeks or months after you've used this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of cobicistat, elvitegravir, emtricitabine, and tenofovir may include: nausea

- or diarrhea.

309. This particular disease, AIDS/HIV, interacts with the drug Cabenuva and has the following side

- hives

- fever, tiredness, body aches, not feeling well

- sores or blisters in your mouth

- red or puffy eyes

- difficult breathing

- swelling of your face, lips, tongue, or throat. Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Your Cabenuva injections may be permanently discontinued if you have an allergic reaction. Some side effects may occur within a few minutes after an injection. Tell your caregiver if you feel anxious, warm, light-headed, sweaty, or have stomach pain, or numbness in your mouth. Call your doctor at once if you have: unusual changes in mood or behavior

- suicidal thoughts or actions

- or liver problems - loss of appetite, nausea , vomiting , stomach pain (upper right side), itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common Cabenuva side

Drug Side Effects Summary

effects may include: pain, redness, swelling, itching, bruising, warmth, or a hard lump where an injection was given

- fever
- nausea
- pain in your bones, joints or muscles
- feeling tired, sleep problems
- headache , dizziness
- or rash.

310. This particular disease, AIDS/HIV, interacts with the drug dolutegravir / lamivudine and has the

- Stop taking this medicine and get emergency medical help if you have signs of an allergic reaction:

fever, general ill feeling, trouble breathing, tiredness

- joint or muscle pain, blisters or mouth sores, redness or swelling in your eyes
- blistering or peeling skin
- swelling of your face, lips, tongue, or throat. Mild symptoms of lactic acidosis may worsen over time , and this condition can be fatal. unusual muscle pain, trouble breathing, stomach pain, vomiting , fast or irregular heart rate, dizziness , feeling cold, or feeling very weak or tired. Dolutegravir and lamivudine may cause serious side effects. Call your doctor at once if you have: the first sign of any skin rash , no matter how mild
- or liver problems-- nausea , vomiting, loss of appetite, upper stomach pain, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Dolutegravir and lamivudine affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly

Drug Side Effects Summary

feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of dolutegravir and lamivudine may include: headache

- tiredness

- nausea, diarrhea

- or sleep problems (insomnia).

311. This particular disease, AIDS/HIV, interacts with the drug emtricitabine / tenofovir alafenamide

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Chest pain or tightness confusion cough fever or chills increase in bone pain irregular heartbeat nausea or vomiting nervousness numbness or tingling in the hands, feet, or lips sneezing stomach pain trouble breathing weakness or heaviness of the legs Incidence not known Agitation bloody urine change in consciousness dark urine decreased appetite decreased frequency or amount of urine depression diarrhea fast, shallow breathing general feeling of discomfort headache hostility increased thirst irritability large, hive-like swelling on face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs light-colored stools loss of appetite loss of consciousness lower back or side pain muscle pain, twitching, or cramping seizures sleepiness stomach discomfort swelling of the face, fingers, or lower legs unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness upper right abdominal or stomach pain vomiting weight gain yellow eyes and skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Emtricitabine / tenofovir alafenamide side effects

Drug Side Effects Summary

312. This particular disease, AIDS/HIV, interacts with the drug Complera and has the following side

- hives, blistering skin rash with fever
- mouth sores, eye redness
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Some people develop lactic acidosis while taking HIV medication. Early symptoms may get worse over time and this condition can be fatal. Get emergency medical help if you have even mild symptoms such as: muscle pain or weakness, numb or cold feeling in your arms and legs, trouble breathing, stomach pain, nausea with vomiting, fast or uneven heart rate, dizziness, or feeling very weak or tired. Call your doctor at once if you have: confusion, severe depression, unusual thoughts or behavior, suicidal thoughts or actions
- increased thirst and urination, loss of appetite, weakness, constipation
- kidney problems - little or no urinating, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- liver problems - swelling around your midsection, upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or signs of inflammation in your body - swollen glands, flu symptoms, easy bruising or bleeding, severe tingling or numbness, muscle weakness, upper stomach pain, jaundice (yellowing of the skin or eyes), chest pain, new or worsening cough with fever, trouble breathing
- Complera may increase your risk of certain infections or autoimmune disorders by changing the way your immune system works. Symptoms may occur weeks or months after you start treatment. Tell your doctor if you have: signs of a new infection - fever, night sweats, swollen glands, diarrhea, weight loss
- chest pain (especially when you breathe), dry cough, wheezing
- cold sores, sores on your genital or anal area
- feeling anxious or irritable, weakness or prickly feeling, problems with balance or eye movement

Drug Side Effects Summary

- trouble speaking or swallowing, severe lower back pain, loss of bladder or bowel control
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence, loss of interest in sex. Common Complera side effects may include: headache, dizziness, tiredness
- depressed mood, sleep problems (insomnia), strange dreams
- rash
- nausea, diarrhea
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

313. This particular disease, AIDS/HIV, interacts with the drug Reyataz and has the following side effects:

- sudden dizziness (like you might pass out)
- severe pain in your lower stomach or back
- painful urination, blood in your urine
- high blood sugar -increased thirst, increased urination, dry mouth, fruity breath odor, headache , blurred vision
- or liver or gallbladder problems - nausea , vomiting , upper stomach pain, itching, fever, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Atazanavir affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection -fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement
- weakness or prickly feeling, loss of bladder or bowel control
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Reyataz side effects may include: fever
- nausea, vomiting, stomach pain, diarrhea

Drug Side Effects Summary

- headache, muscle pain
- depressed mood, sleep problems (insomnia)
- numbness , tingling, or burning pain in your hands or feet
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

314. This particular disease, AIDS/HIV, interacts with the drug Epzicom and has the following side effects

- Call your doctor at once if you have symptoms of an allergic reaction from two or more of these specific side effect groups: Group 1 - fever
- Group 2 - rash
- Group 3 - nausea, vomiting, diarrhea, stomach pain
- Group 4 - general ill feeling, extreme tiredness, body aches
- Group 5 - shortness of breath, cough, sore throat. Once you have an allergic reaction to abacavir, you must never use it again. If you stop taking this medicine for any reason, talk to your doctor before you start taking it again. Also call your doctor at once if you have: other signs of allergic reaction - hives , swelling in your face or throat, trouble breathing
- lactic acidosis - unusual muscle pain, trouble breathing, stomach pain, vomiting, fast/slow or irregular heartbeats, dizziness , feeling cold, feeling very weak or tired
- or liver problems - loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Abacavir and lamivudine affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea, weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

Drug Side Effects Summary

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Epzicom side effects include: allergic reaction
- depression , trouble sleeping
- feeling weak or tired
- headache , dizziness, migraine
- nausea, diarrhea
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and trunk).

315. This particular disease, AIDS/HIV, interacts with the drug ritonavir and has the following side effects

- irregular heartbeats, or a light-headed feeling (like you might pass out)
- easy bruising, unusual bleeding (nose, mouth, vagina, or rectum), purple or red pinpoint spots under your skin
- signs of a kidney stone--pain in your side or lower back, blood in your urine, painful or difficult urination
- high blood sugar--increased thirst, increased urination, hunger, dry mouth, fruity breath odor
- or signs of liver or pancreas problems--loss of appetite, upper stomach pain (that may spread to your back), nausea , vomiting , dark urine, jaundice (yellowing of the skin or eyes). Ritonavir affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of ritonavir may include: nausea, vomiting, stomach pain, diarrhea

Drug Side Effects Summary

- numbness or tingling in your hands or feet or around your mouth
- feeling weak or tired
- rash
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

316. This particular disease, AIDS/HIV, interacts with the drug lamivudine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Mild symptoms of lactic acidosis may worsen over time , and this condition can be fatal. Get emergency medical help if you have: unusual muscle pain, trouble breathing, stomach pain, vomiting , irregular heart rate, dizziness , feeling cold, or feeling very weak or tired. Lamivudine may cause serious side effects. Call your doctor at once if you have: pancreas problems--severe pain in your upper stomach spreading to your back, nausea and vomiting , fast heart rate
- or liver problems--swelling around your midsection, right-sided upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Lamivudine affects your immune system, which may cause certain side effects (even weeks or months after you've taken lamivudine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of lamivudine may include: nausea, diarrhea
- headache

Drug Side Effects Summary

- fever, tiredness, general ill feeling
- ear infection--ear pain or full feeling, trouble hearing, drainage from the ear, fussiness in a child
- or nose or throat infection--stuffy nose, sneezing, sore throat , cough.

317. This particular disease, AIDS/HIV, interacts with the drug tenofovir and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Mild symptoms of lactic acidosis may worsen over time , and this condition can be fatal. Get emergency medical help if you have: unusual muscle pain, trouble breathing, stomach pain, vomiting , fast/slow or irregular heartbeats, dizziness , feeling cold, or feeling very weak or tired. Tenofovir may cause serious side effects. Call your doctor at once if you have: sore throat , flu symptoms, easy bruising or unusual bleeding
- kidney problems--little or no urination, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- or liver problems--swelling around your midsection, upper stomach pain, unusual tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Tenofovir can affect your immune system (even weeks or months after you've used this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of tenofovir may include: stomach pain, nausea , vomiting, diarrhea
- fever, pain
- weakness, dizziness

Drug Side Effects Summary

- headache
- depressed mood
- itching, rash
- or sleep problems (insomnia).

318. This particular disease, AIDS/HIV, interacts with the drug Viread and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Mild symptoms of lactic acidosis may worsen over time , and this condition can be fatal. Get emergency medical help if you have: unusual muscle pain, trouble breathing, stomach pain, vomiting , fast/slow or irregular heartbeats, dizziness , feeling cold, or feeling very weak or tired. Call your doctor at once if you have: sore throat , flu symptoms, easy bruising or unusual bleeding
- kidney problems - little or no urination, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- or liver problems - swelling around your midsection, upper stomach pain, unusual tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Viread can affect your immune system (even weeks or months after you've used this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Viread side effects may include: stomach pain, nausea , vomiting, diarrhea
- fever, pain

Drug Side Effects Summary

- weakness, dizziness
- headache
- depressed mood
- itching, rash
- or sleep problems (insomnia).

319. This particular disease, AIDS/HIV, interacts with the drug dolutegravir and has the following side effects:

- Stop taking this medicine and get emergency medical help if you have signs of an allergic reaction: fever, general ill feeling, trouble breathing, tiredness
- joint or muscle pain, blisters or mouth sores, redness or swelling in your eyes
- blistering or peeling skin
- swelling of your face, lips, tongue, or throat. Dolutegravir may cause serious side effects. Call your doctor at once if you have: the first sign of any skin rash , no matter how mild
- or liver problems-- nausea , vomiting , loss of appetite, upper stomach pain, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Dolutegravir affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of dolutegravir may include: headache
- tiredness
- or sleep problems (insomnia).

Drug Side Effects Summary

320. This particular disease, AIDS/HIV, interacts with the drug Kaletra and has the following side effects:

- hives, difficult breathing, swelling in your face or throat) or a severe skin reaction (fever, sore throat, burning eyes, skin pain, red or purple skin rash with blistering and peeling). Call your doctor at once if you have: fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- signs of a kidney stone - pain in your side or lower back, blood in your urine, painful or difficult urination
- high blood sugar - increased thirst, increased urination, fruity breath odor, weight loss
- or signs of liver or pancreas problems - loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Kaletra affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats, swollen glands, cold sores, cough, wheezing, diarrhea, weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence. liquid Kaletra can cause changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist). Common Kaletra side effects may include: nausea, vomiting, diarrhea
- or high cholesterol or triglycerides.

321. This particular disease, AIDS/HIV, interacts with the drug Epivir and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Mild symptoms of lactic acidosis may worsen over time , and this condition can be fatal. Get emergency medical help if you have: unusual muscle pain,

Drug Side Effects Summary

trouble breathing, stomach pain, vomiting , irregular heart rate, dizziness, feeling cold, or feeling very weak or tired. Call your doctor at once if you have: pancreas problems - severe pain in your upper stomach spreading to your back, nausea and vomiting , fast heart rate

- or liver problems - swelling around your midsection, right-sided upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Lamivudine affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Epivir side effects may include: nausea, diarrhea

- headache

- fever, tiredness, general ill feeling

- ear infection - ear pain or full feeling, trouble hearing, drainage from the ear, fussiness in a child

- or nose or throat infection - stuffy nose, sneezing, sore throat , cough.

322. This particular disease, AIDS/HIV, interacts with the drug Intelence and has the following side effects:

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. In rare cases, Intelence can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, and dark colored urine. Stop using Intelence and call your doctor at once if you have: swelling, rapid weight gain, little or no urinating

Drug Side Effects Summary

- confusion, seizure
- liver problems: nausea, upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- the first sign of any type of skin rash, no matter how mild
- or severe skin reaction - fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain, followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Intelence may increase your risk of certain infections or autoimmune disorders by changing the way your immune system works. Symptoms may occur weeks or months after you start treatment with this medicine. Tell your doctor if you have: signs of a new infection - fever, night sweats, swollen glands, mouth sores, diarrhea, stomach pain, weight loss
- chest pain (especially when you breathe), dry cough, wheezing, feeling short of breath
- cold sores, sores on your genital or anal area
- rapid heart rate, feeling anxious or irritable, weakness or prickly feeling, problems with balance or eye movement
- trouble speaking or swallowing, severe lower back pain, loss of bladder or bowel control
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence, loss of interest in sex. Common Intelence side effects may include: diarrhea
- numbness or tingly feeling in your hands or feet
- rash
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

323. This particular disease, AIDS/HIV, interacts with the drug Sustiva and has the following side effects:

- hives, difficulty breathing, swelling in your face or throat) or a severe skin reaction (fever, sore

Drug Side Effects Summary

throat , burning eyes, skin pain, red or purple skin rash with blistering and peeling). Call your doctor at once if you have: a seizure

- hallucinations, trouble concentrating, trouble speaking or moving (these symptoms may occur months or years after you start taking efavirenz)

- nervous system problems - dizziness, drowsiness, confusion, trouble concentrating, strange dreams, insomnia, or problems with speech, balance, or muscle movement

- serious psychiatric symptoms - anxiety, paranoia, unusual behavior, feeling sad or hopeless, hallucinations, suicidal thoughts

- or liver problems - nausea, stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)

- Sustiva may cause serious psychiatric symptoms including confusion, feeling sad or hopeless, suicidal thoughts, anxiety , paranoia, hallucinations, trouble speaking or moving, or unusual behavior. Contact your doctor at once if you have any of these side effects, even if you have had them before. Efavirenz affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Sustiva side effects may include: nausea, vomiting

- dizziness , drowsiness, trouble concentrating

- rash

- headache , tired feeling

- sleep problems (insomnia), strange dreams

- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts,

Drug Side Effects Summary

and waist).

324. This particular disease, AIDS/HIV, interacts with the drug abacavir and has the following side effects:

- Stop using abacavir and call your doctor at once if you have symptoms of an allergic reaction from two or more of these specific side effect groups: Group 1 - fever

- Group 2 - rash

- Group 3 - nausea, vomiting, diarrhea, stomach pain

- Group 4 - general ill feeling, extreme tiredness, body aches

- Group 5 - shortness of breath, cough, sore throat. Once you have had an allergic reaction to abacavir, you must never use it again. If you stop taking abacavir for any reason, talk to your doctor before you start taking the medication again. Abacavir can cause other serious side effects that may not be signs of an allergic reaction. Call your doctor at once if you have: severe upper stomach pain, nausea, vomiting, loss of appetite

- swelling around your midsection

- dark urine, clay-colored stools, or jaundice (yellowing of the skin or eyes)

- unusual tiredness

- or chest pain or pressure, pain spreading to your jaw or shoulder. Mild symptoms of lactic acidosis may worsen over time, and this condition can be fatal. unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness, feeling cold, or feeling very weak or tired. Abacavir affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats, swollen glands, cold sores, cough, wheezing, diarrhea, weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence. Common side

Drug Side Effects Summary

effects of abacavir may include: feeling tired

- sleep problems, strange dreams
- headache , tiredness, fever, chills, general ill feeling
- nausea or vomiting
- rash
- or (in children) stuffy nose, sneezing, sore throat, ear pain.

325. This particular disease, AIDS/HIV, interacts with the drug efavirenz / emtricitabine / tenofovir and

- unusual muscle pain, trouble breathing, stomach pain, vomiting , irregular heart rate, dizziness, feeling cold, or feeling very weak or tired. Efavirenz, emtricitabine, and tenofovir may cause serious side effects. Call your doctor at once if you have: unusual thoughts or behavior, anger, severe depression , thoughts of hurting yourself or others, hallucinations
- a seizure (convulsions)
- kidney problems--increased thirst and urination, muscle pain or weakness
- or liver problems--swelling around your midsection, right-sided upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Atripla affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of efavirenz, emtricitabine, and tenofovir may include: dizziness, drowsiness, tired feeling
- nausea , diarrhea
- headache , depressed mood, trouble concentrating

Drug Side Effects Summary

- sleep problems (insomnia), strange dreams
- rash
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

326. This particular disease, AIDS/HIV, interacts with the drug emtricitabine / tenofovir and has the

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Mild symptoms of lactic acidosis may worsen over time , and this condition can be fatal. Get emergency medical help if you have: unusual muscle pain, trouble breathing, stomach pain, vomiting , irregular heart rate, dizziness, feeling cold, or feeling very weak or tired. Emtricitabine and tenofovir may cause serious side effects. Call your doctor at once if you have: symptoms of new HIV infection--fever, night sweats, tiredness, muscle or joint pain, sore throat, vomiting, diarrhea, rash, swollen glands in your neck or groin
- sudden or unusual bone pain
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or liver problems-- nausea , swelling around your midsection, upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Emtricitabine and tenofovir affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, sweats, swollen glands, cold sores , cough, wheezing, diarrhea, weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of emtricitabine and tenofovir may include: headache , dizziness, feeling depressed or tired

Drug Side Effects Summary

- trouble sleeping, strange dreams
- nausea, stomach pain
- weight loss
- or rash.

327. This particular disease, AIDS/HIV, interacts with the drug darunavir and has the following side

- a skin rash, no matter how mild
 - high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
 - or signs of liver or pancreas problems--loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting , fast heart rate , dark urine, jaundice (yellowing of the skin or eyes).
- Darunavir affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
 - or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of darunavir may include: nausea, vomiting, diarrhea, stomach pain
 - headache
 - rash
 - or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

328. This particular disease, AIDS/HIV, interacts with the drug efavirenz and has the following side

- a seizure
- hallucinations, trouble concentrating, trouble speaking or moving (these symptoms may occur

Drug Side Effects Summary

months or years after you start taking efavirenz)

- nervous system problems-- dizziness , drowsiness, confusion, trouble concentrating, strange dreams, insomnia , or problems with speech, balance, or muscle movement
- serious psychiatric symptoms-- anxiety , paranoia, unusual behavior, feeling sad or hopeless, hallucinations, suicidal thoughts
- or liver problems-- nausea , stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- Psychiatric symptoms or nervous system problems may happen even months or years after you have been taking efavirenz. Efavirenz affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of efavirenz may include: nausea, vomiting
- dizziness, drowsiness, trouble concentrating
- rash
- headache , tired feeling
- sleep problems (insomnia), strange dreams
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

329. This particular disease, AIDS/HIV, interacts with the drug Odefsey and has the following side effects

- hives, blistering skin rash with fever

Drug Side Effects Summary

- mouth sores, eye redness
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Some people develop lactic acidosis while taking HIV medication. Early symptoms may get worse over time and this condition can be fatal. Get emergency medical help if you have even mild symptoms such as: muscle pain or weakness, numb or cold feeling in your arms and legs, trouble breathing, stomach pain, nausea with vomiting, fast or uneven heart rate, dizziness, or feeling very weak or tired. Call your doctor at once if you have: confusion, severe depression, unusual thoughts or behavior, suicidal thoughts or actions
- increased thirst and urination, loss of appetite, weakness, constipation
- kidney problems - little or no urinating, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- liver problems - swelling around your midsection, upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or signs of inflammation in your body - swollen glands, flu symptoms, easy bruising or bleeding, severe tingling or numbness, muscle weakness, upper stomach pain, jaundice (yellowing of the skin or eyes), chest pain, new or worsening cough with fever, trouble breathing
- Odefsey may increase your risk of certain infections or autoimmune disorders by changing the way your immune system works. Symptoms may occur weeks or months after you start treatment. Tell your doctor if you have: signs of a new infection - fever, night sweats, swollen glands, diarrhea, weight loss
- chest pain (especially when you breathe), dry cough, wheezing
- cold sores, sores on your genital or anal area
- feeling anxious or irritable, weakness or prickly feeling, problems with balance or eye movement
- trouble speaking or swallowing, severe lower back pain, loss of bladder or bowel control
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence, loss of interest

Drug Side Effects Summary

in sex. Common Odefsey side effects may include: headache, dizziness, tiredness

- depressed mood, sleep problems (insomnia), strange dreams
- rash
- nausea, diarrhea
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

330. This particular disease, AIDS/HIV, interacts with the drug raltegravir and has the following side

- Stop taking this medicine and get emergency medical help if you have signs of an allergic reaction: fever, general ill feeling, tiredness, joint or muscle pain, trouble breathing
- upper stomach pain, vomiting, loss of appetite, dark urine, yellowing of the skin or eyes
- burning eyes, blisters or mouth sores
- rash, hives , blistering or peeling skin
- swelling of your face, lips, tongue, or throat. In rare cases, raltegravir can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure . Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark colored urine. Raltegravir affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of raltegravir may include: nausea
- headache , dizziness

Drug Side Effects Summary

- tired feeling
- or sleep problems (insomnia).

331. This particular disease, AIDS/HIV, interacts with the drug abacavir / lamivudine and has the fol

- Call your doctor at once if you have symptoms of an allergic reaction from two or more of these specific side effect groups: Group 1 - fever

- Group 2 - rash

- Group 3 - nausea, vomiting, diarrhea, stomach pain

- Group 4 - general ill feeling, extreme tiredness, body aches

- Group 5 - shortness of breath, cough, sore throat. Once you have an allergic reaction to abacavir, you must never use it again. If you stop taking this medicine for any reason, talk to your doctor before you start taking it again. Also call your doctor at once if you have: other signs of allergic reaction-- hives , swelling in your face or throat, trouble breathing

- lactic acidosis --unusual muscle pain, trouble breathing, stomach pain, vomiting, fast/slow or irregular heartbeats, dizziness , feeling cold, feeling very weak or tired

- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Abacavir and lamivudine affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea, weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects include: allergic reaction

- depression , trouble sleeping

Drug Side Effects Summary

- feeling weak or tired
- headache , dizziness, migraine
- nausea, diarrhea
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and trunk).

332. This particular disease, AIDS/HIV, interacts with the drug Prezcobix and has the following side

- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor, headache , blurred vision
- or liver problems - upper stomach pain, vomiting, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Prezcobix affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing , diarrhea , weight loss
- stomach pain, tiredness, itching, skin redness, yellowing of the skin or eyes
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Prezcobix side effects may include: nausea, vomiting, stomach pain, diarrhea
- headache
- rash
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

333. This particular disease, AIDS/HIV, interacts with the drug Viramune and has the following side

Drug Side Effects Summary

- joint or muscle pain, fever, mouth sores, facial swelling, blistering skin rash, flu symptoms, swollen glands, feeling weak or tired, severe tingling or numbness , pain or burning when you urinate, swelling in your legs or feet, cough, chest pain, trouble breathing, or swelling in your lips, tongue, or throat. Viramune can cause life-threatening effects on the liver, especially in women. Call your doctor at once if you have any of these liver symptoms while taking this medicine: nausea, loss of appetite, upper stomach pain, tiredness, fever, unexplained muscle pain or weakness, dark urine, clay-colored stools, or jaundice (yellowing of the skin or eyes). Viramune may also cause severe or life-threatening skin reactions. Stop taking Viramune and get emergency medical help if you have: a fever, sore throat , swelling in your face or tongue, burning in your eyes, skin pain, and a red or purple skin rash that spreads and causes blistering and peeling. This type of reaction is a medical emergency. Viramune may increase your risk of certain infections or autoimmune disorders by changing the way your immune system works. Symptoms may occur weeks or months after you start treatment with this medicine. Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, mouth sores, diarrhea , stomach pain, weight loss
- chest pain (especially when you breathe), dry cough, wheezing, feeling short of breath
- cold sores , sores on your genital or anal area
- rapid heart rate, feeling anxious or irritable, weakness or prickly feeling, problems with balance or eye movement
- trouble speaking or swallowing, severe lower back pain , loss of bladder or bowel control
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence , loss of interest in sex. Common side effects of Viramune may include: skin rash
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

334. This particular disease, AIDS/HIV, interacts with the drug Combivir and has the following side effects:

Drug Side Effects Summary

- hives
 - difficult breathing
 - swelling of your face, lips, tongue, or throat. Mild symptoms of lactic acidosis may worsen over time , and this condition can be fatal. Get emergency medical help if you have: unusual muscle pain, trouble breathing, stomach pain, vomiting , irregular heart rate, dizziness, feeling cold, or feeling very weak or tired. Call your doctor at once if you have: liver problems - swelling around your midsection, right-sided upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
 - low red blood cells (anemia) - pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet
 - low white blood cell counts - fever, mouth sores, skin sores, sore throat , cough, trouble breathing
 - or pancreatitis - severe pain in your upper stomach spreading to your back, nausea and vomiting .
- Lamivudine and zidovudine affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
 - or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence .
- Combivir side effects may include: headache
- nausea , diarrhea
 - tiredness, general ill feeling
 - cold symptoms such as stuffy nose, sneezing, sinus pain, cough
 - or loss of body fat (especially in your arms, legs, face, and buttocks).

Drug Side Effects Summary

335. This particular disease, AIDS/HIV, interacts with the drug Ziagen and has the following side effects:

- Stop using Ziagen and call your doctor at once if you have symptoms of an allergic reaction from two or more of these specific side effect groups: Group 1 - fever
- Group 2 - rash
- Group 3 - nausea, vomiting, diarrhea, stomach pain
- Group 4 - general ill feeling, extreme tiredness, body aches
- Group 5 - shortness of breath, cough, sore throat. Once you have had an allergic reaction to Ziagen, you must never use it again. If you stop taking this medicine for any reason, talk to your doctor before you start taking the medication again. Ziagen can cause other serious side effects that may not be signs of an allergic reaction. Call your doctor at once if you have: severe upper stomach pain, nausea, vomiting, loss of appetite
- swelling around your midsection
- dark urine, clay-colored stools, or jaundice (yellowing of the skin or eyes)
- unusual tiredness
- or chest pain or pressure, pain spreading to your jaw or shoulder. Mild symptoms of lactic acidosis may worsen over time , and this condition can be fatal. unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness, feeling cold, or feeling very weak or tired. Ziagen affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea, weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of Ziagen may include: feeling tired
- sleep problems, strange dreams

Drug Side Effects Summary

- headache , tiredness, fever, chills, general ill feeling
- nausea or vomiting
- rash
- or (in children) stuffy nose, sneezing, sore throat, ear pain.

336. This particular disease, AIDS/HIV, interacts with the drug zidovudine and has the following side effects:

- unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness, feeling cold, or feeling very weak or tired. Zidovudine may cause serious side effects. Call your doctor at once if you have: low blood cell counts--fever, chills, tiredness, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath
- or liver problems--swelling around your midsection, right-sided upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Zidovudine affects your immune system, which may cause certain side effects (even weeks or months after you've used this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of zidovudine may include: headache
- fever, general ill feeling
- cough
- nausea , vomiting, loss of appetite
- or loss of body fat (especially in your arms, legs, face, and buttocks).

Drug Side Effects Summary

337. This particular disease, AIDS/HIV, interacts with the drug atazanavir and has the following side effects:

- sudden dizziness (like you might pass out)
- severe pain in your lower stomach or back
- painful urination, blood in your urine
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor, headache , blurred vision
- or liver or gallbladder problems-- nausea , vomiting , upper stomach pain, itching, fever, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Atazanavir affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement
- weakness or prickly feeling, loss of bladder or bowel control
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of atazanavir may include: fever
- nausea, vomiting, stomach pain, diarrhea
- headache, muscle pain
- depressed mood, sleep problems (insomnia)
- numbness , tingling, or burning pain in your hands or feet
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

338. This particular disease, AIDS/HIV, interacts with the drug emtricitabine and has the following side effects:

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Early symptoms of lactic acidosis may get worse over time and this condition can be fatal. Get emergency medical help if you have even mild symptoms: muscle pain or weakness, numb or cold feeling in your arms and legs, trouble breathing, stomach pain, nausea with vomiting, fast or uneven heart rate, dizziness, or feeling very weak or tired. Call your doctor at once if you have symptoms of liver problems: swelling around your midsection, upper stomach pain, nausea, loss of appetite, dark urine, or jaundice (yellowing of the skin or eyes). Emtricitabine may increase your risk of certain infections or autoimmune disorders by changing the way your immune system works. Symptoms may occur weeks or months after you start treatment with emtricitabine. Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, mouth sores, diarrhea , stomach pain, weight loss
- chest pain (especially when you breathe), dry cough, wheezing, feeling short of breath
- cold sores , sores on your genital or anal area
- rapid heart rate, feeling anxious or irritable, weakness or prickly feeling, problems with balance or eye movement
- trouble speaking or swallowing, severe lower back pain , loss of bladder or bowel control
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence , loss of interest in sex. Common side effects of emtricitabine may include: headache , dizziness, weakness
- indigestion , stomach pain, nausea, vomiting, diarrhea
- sleep problems, strange dreams
- rash, skin discoloration
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

339. This particular disease, AIDS/HIV, interacts with the drug Emtriva and has the following side effects

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Early symptoms of lactic acidosis may get worse over time and this condition can be fatal. Get emergency medical help if you have even mild symptoms: muscle pain or weakness, numb or cold feeling in your arms and legs, trouble breathing, stomach pain, nausea with vomiting, fast or uneven heart rate, dizziness, or feeling very weak or tired. Call your doctor at once if you have symptoms of liver problems: swelling around your midsection, upper stomach pain, nausea, loss of appetite, dark urine, or jaundice (yellowing of the skin or eyes). Emtriva may increase your risk of certain infections or autoimmune disorders by changing the way your immune system works. Symptoms may occur weeks or months after you start treatment with this medicine. Tell your doctor if you have: signs of a new infection - fever, night sweats, swollen glands, mouth sores, diarrhea, stomach pain, weight loss
- chest pain (especially when you breathe), dry cough, wheezing, feeling short of breath
- cold sores, sores on your genital or anal area
- rapid heart rate, feeling anxious or irritable, weakness or prickly feeling, problems with balance or eye movement
- trouble speaking or swallowing, severe lower back pain, loss of bladder or bowel control
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence, loss of interest in sex. Common Emtriva side effects may include: headache, dizziness, weakness
- indigestion, stomach pain, nausea, vomiting, diarrhea
- sleep problems, strange dreams
- rash, skin discoloration
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

340. This particular disease, AIDS/HIV, interacts with the drug Edurant and has the following side effects:

Drug Side Effects Summary

- skin rash, fever, swollen glands, flu-like symptoms, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Call your doctor at once if you have: mood changes, anxiety , feeling sad or hopeless, thoughts about suicide or hurting yourself
- liver problems - right-sided upper stomach pain, dark urine, jaundice (yellowing of the skin or eyes)
- or symptoms of depression - mood changes, feelings of low self-worth, loss of interest in things you once enjoyed, new sleep problems, thoughts about hurting yourself. Edurant affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Edurant side effects may include: sleep problems (insomnia)
- depression
- skin rash
- headache
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

341. This particular disease, AIDS/HIV, interacts with the drug Lexiva and has the following side effects

- pale or yellowed skin, dark colored urine, fever, confusion or weakness
- increased urination or extreme thirst
- or high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- or symptoms of kidney stones - sudden and severe pain in your lower back or side, blood in your urine, pain or burning when you urinate. Fosamprenavir affects your immune system, which may

Drug Side Effects Summary

cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Lexiva side effects may include: fever or other signs of infection

- nausea , vomiting , diarrhea

- rash

- headache

- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

342. This particular disease, AIDS/HIV, interacts with the drug nevirapine and has the following side

- joint or muscle pain, fever, mouth sores, facial swelling, blistering skin rash, flu symptoms, swollen glands, feeling weak or tired, severe tingling or numbness , pain or burning when you urinate, swelling in your legs or feet, cough, chest pain, trouble breathing, or swelling in your lips, tongue, or throat. Nevirapine can cause life-threatening effects on the liver, especially in women. Call your doctor at once if you have any of these liver symptoms while taking nevirapine: nausea, loss of appetite, upper stomach pain, tiredness, fever, unexplained muscle pain or weakness, dark urine, clay-colored stools, or jaundice (yellowing of the skin or eyes). Nevirapine may also cause severe or life-threatening skin reactions. Stop taking nevirapine and get emergency medical help if you have: a fever, sore throat , swelling in your face or tongue, burning in your eyes, skin pain, and a red or purple skin rash that spreads and causes blistering and peeling. This type of reaction is a medical emergency. Nevirapine may increase your risk of certain infections or autoimmune disorders by

Drug Side Effects Summary

changing the way your immune system works. Symptoms may occur weeks or months after you start treatment with nevirapine. Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, mouth sores, diarrhea , stomach pain, weight loss

- chest pain (especially when you breathe), dry cough, wheezing, feeling short of breath
- cold sores , sores on your genital or anal area
- rapid heart rate, feeling anxious or irritable, weakness or prickly feeling, problems with balance or eye movement
- trouble speaking or swallowing, severe lower back pain , loss of bladder or bowel control
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence , loss of interest in sex. Common side effects of nevirapine may include: skin rash
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

343. This particular disease, AIDS/HIV, interacts with the drug Retrovir and has the following side effects

- unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness, feeling cold, or feeling very weak or tired. Retrovir may cause serious side effects. Call your doctor at once if you have: low blood cell counts--fever, chills, tiredness, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath
- or liver problems--swelling around your midsection, right-sided upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Retrovir affects your immune system, which may cause certain side effects (even weeks or months after you've used this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

Drug Side Effects Summary

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of Retrovir may include: headache
- fever, general ill feeling
- cough
- nausea , vomiting, loss of appetite
- or loss of body fat (especially in your arms, legs, face, and buttocks).

344. This particular disease, AIDS/HIV, interacts with the drug Selzentry and has the following side

- itching or rash, fever, swollen glands, muscle or joint aches, severe weakness, unusual bruising, upper stomach pain, vomiting, dark urine, or jaundice (yellowing of the skin or eyes). Call your doctor at once if you have a side effect such as: a light-headed feeling, like you might pass out
- or chest pressure, tight feeling in your neck or jaw, sweating, pain spreading to your arm or shoulder. Selzentry affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Selzentry side effects may include: fever, cold symptoms , cough
- indigestion , gas, bloating , nausea , vomiting, diarrhea, constipation
- rash
- dizziness
- or (in newborns) abnormal blood tests.

345. This particular disease, AIDS/HIV, interacts with the drug Juluca and has the following side eff

Drug Side Effects Summary

- Stop taking this medicine and get emergency medical help if you have signs of an allergic reaction to Juluca: hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Call your doctor at once if you have: thoughts about suicide or hurting yourself
- anxiety , sadness, feeling hopeless
- depression - mood changes, feelings of low self-worth, loss of interest in things you once enjoyed, new sleep problems, thoughts about hurting yourself
- or liver problems - nausea , vomiting , loss of appetite, right-sided upper stomach pain , dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Dolutegravir and rilpivirine affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats, swollen glands, cold sores, cough, wheezing, diarrhea, weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence. Common Juluca side effects may include: headache
- or nausea, diarrhea .

346. This particular disease, AIDS/HIV, interacts with the drug Symtuza and has the following side effects

- increased thirst, increased urination
- kidney problems - swelling, urinating less, feeling tired or short of breath
- lactic acidosis> - unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart

Drug Side Effects Summary

rate, dizziness , feeling cold, or feeling very weak or tired

- or liver problems - nausea , upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Symtuza can affect your immune system (even weeks or months after you've used Symtuza). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Symtuza side effects may include: nausea, upset stomach , diarrhea, gas

- headache , feeling tired

- or rash.

347. This particular disease, AIDS/HIV, interacts with the drug Trizivir and has the following side effects

- Call your doctor at once if you have symptoms of an allergic reaction from two or more of these specific side effect groups: Group 1 - fever

- Group 2 - rash

- Group 3 - nausea, vomiting, diarrhea, stomach pain

- Group 4 - general ill feeling, extreme tiredness, body aches

- Group 5 - shortness of breath, cough, sore throat. Once you have had an allergic reaction to Trizivir, you must never use it again. If you stop taking Trizivir for any reason, talk to your doctor before you start taking it again. Also call your doctor at once if you have: other signs of allergic reaction - skin blisters or peeling, eye redness , swelling in your face or throat, trouble breathing

- lactic acidosis - unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness , feeling cold, or feeling very weak or tired

- low white blood cell counts - fever, mouth sores, skin sores, sore throat, cough, trouble breathing

Drug Side Effects Summary

- low red blood cells (anemia) - pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet
- or liver problems - swelling around your midsection, right-sided upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Trizivir affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea, weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Trizivir side effects may include: headache
- weakness or tiredness
- nausea, vomiting
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

348. This particular disease, AIDS/HIV, interacts with the drug Epivir-HBV and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Mild symptoms of lactic acidosis may worsen over time , and this condition can be fatal. Get emergency medical help if you have: unusual muscle pain, trouble breathing, stomach pain, vomiting , irregular heart rate, dizziness , feeling cold, or feeling very weak or tired. Epivir HBV may cause serious side effects. Call your doctor at once if you have: pancreas problems--severe pain in your upper stomach spreading to your back, nausea and vomiting , fast heart rate

Drug Side Effects Summary

- or liver problems--swelling around your midsection, right-sided upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Epivir HBV affects your immune system, which may cause certain side effects (even weeks or months after you've taken Epivir HBV). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of Epivir HBV may include: nausea, diarrhea
- headache
- fever, tiredness, general ill feeling
- ear infection--ear pain or full feeling, trouble hearing, drainage from the ear, fussiness in a child
- or nose or throat infection--stuffy nose, sneezing, sore throat , cough.

349. This particular disease, AIDS/HIV, interacts with the drug etravirine and has the following side

- skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Etravirine may cause serious side effects. Call your doctor at once if you have: unexplained muscle pain, tenderness, or weakness
- fever, unusual tiredness
- dark colored urine
- or any type of skin rash, no matter how mild. Etravirine affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly

Drug Side Effects Summary

feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of etravirine may include: diarrhea
- numbness or tingly feeling in your hands or feet
- rash
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

350. This particular disease, AIDS/HIV, interacts with the drug Isentress HD and has the following s

- Stop taking this medicine and get emergency medical help if you have signs of an allergic reaction: fever, general ill feeling, tiredness, joint or muscle pain, trouble breathing
- upper stomach pain, vomiting, loss of appetite, dark urine, yellowing of the skin or eyes
- burning eyes, blisters or mouth sores
- rash, hives , blistering or peeling skin
- swelling of your face, lips, tongue, or throat. In rare cases, Isentress HD can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure . Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark colored urine. Isentress HD affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of Isentress HD may include: nausea

Drug Side Effects Summary

- headache , dizziness
- tired feeling
- or sleep problems (insomnia).

351. This particular disease, AIDS/HIV, interacts with the drug lamivudine / tenofovir and has the fol

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Lamivudine and tenofovir may cause serious side effects. Call your doctor at once if you have: new or worsening bone pain
- muscle weakness
- pain in your arms, hands, legs, or feet
- swelling around your midsection
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or signs of liver or pancreas problems--loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting , loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Lamivudine and tenofovir affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of lamivudine and tenofovir may include: headache
- pain
- diarrhea

Drug Side Effects Summary

- rash
- depression
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

352. This particular disease, AIDS/HIV, interacts with the drug lopinavir / ritonavir and has the following

- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- signs of a kidney stone--pain in your side or lower back, blood in your urine, painful or difficult urination
- high blood sugar--increased thirst, increased urination, fruity breath odor, weight loss
- or signs of liver or pancreas problems--loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Lopinavir and ritonavir affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Lopinavir and ritonavir liquid contains alcohol and propylene glycol, which may cause drowsiness or slow breathing in a baby taking this medicine. Tell your doctor if you notice these symptoms in your baby. Lopinavir and ritonavir can cause changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist). Common side effects of lopinavir and ritonavir may include: nausea, vomiting, diarrhea

Drug Side Effects Summary

- or high cholesterol or triglycerides.

353. This particular disease, AIDS/HIV, interacts with the drug maraviroc and has the following side effects:

- itching or rash, fever, swollen glands, muscle or joint aches, severe weakness, unusual bruising, upper stomach pain, vomiting, dark urine, or jaundice (yellowing of the skin or eyes). Call your doctor at once if you have a side effect such as: a light-headed feeling, like you might pass out
- or chest pressure, tight feeling in your neck or jaw, sweating, pain spreading to your arm or shoulder. Maraviroc affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of maraviroc may include: fever, cold symptoms , cough
- indigestion , gas, bloating , nausea , vomiting, diarrhea, constipation
- rash
- dizziness
- or (in newborns) abnormal blood tests.

354. This particular disease, AIDS/HIV, interacts with the drug Tivicay PD and has the following side effects:

- Stop taking this medicine and get emergency medical help if you have signs of an allergic reaction: fever, general ill feeling, trouble breathing, tiredness
- joint or muscle pain, blisters or mouth sores, redness or swelling in your eyes
- blistering or peeling skin
- swelling of your face, lips, tongue, or throat. Tivicay PD may cause serious side effects. Call your

Drug Side Effects Summary

doctor at once if you have: the first sign of any skin rash , no matter how mild

- or liver problems-- nausea , vomiting , loss of appetite, upper stomach pain, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Tivicay PD affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of Tivicay PD may include: headache
- tiredness
- or sleep problems (insomnia).

355. This particular disease, AIDS/HIV, interacts with the drug Viramune XR and has the following s

- joint or muscle pain, fever, mouth sores, facial swelling, blistering skin rash, flu symptoms, swollen glands, feeling weak or tired, severe tingling or numbness , pain or burning when you urinate, swelling in your legs or feet, cough, chest pain, trouble breathing, or swelling in your lips, tongue, or throat. Viramune XR can cause life-threatening effects on the liver, especially in women. Call your doctor at once if you have any of these liver symptoms while taking this medicine: nausea, loss of appetite, upper stomach pain, tiredness, fever, unexplained muscle pain or weakness, dark urine, clay-colored stools, or jaundice (yellowing of the skin or eyes). Viramune XR may also cause severe or life-threatening skin reactions. Stop taking Viramune XR and get emergency medical help if you have: a fever, sore throat , swelling in your face or tongue, burning in your eyes, skin pain, and a red or purple skin rash that spreads and causes blistering and peeling. This type of reaction is a medical emergency. Viramune XR may increase your risk of certain infections or autoimmune disorders by

Drug Side Effects Summary

changing the way your immune system works. Symptoms may occur weeks or months after you start treatment with this medicine. Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, mouth sores, diarrhea , stomach pain, weight loss

- chest pain (especially when you breathe), dry cough, wheezing, feeling short of breath
- cold sores , sores on your genital or anal area
- rapid heart rate, feeling anxious or irritable, weakness or prickly feeling, problems with balance or eye movement
- trouble speaking or swallowing, severe lower back pain , loss of bladder or bowel control
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence , loss of interest in sex. Common side effects of Viramune XR may include: skin rash
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

356. This particular disease, AIDS/HIV, interacts with the drug AccessPak for HIV PEP Basic and ha

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Mild symptoms of lactic acidosis may worsen over time , and this condition can be fatal. Get emergency medical help if you have: unusual muscle pain, trouble breathing, stomach pain, vomiting , irregular heart rate, dizziness, feeling cold, or feeling very weak or tired. AccessPak for HIV PEP Basic may cause serious side effects. Call your doctor at once if you have: symptoms of new HIV infection--fever, night sweats, tiredness, muscle or joint pain, sore throat, vomiting, diarrhea, rash, swollen glands in your neck or groin
- sudden or unusual bone pain
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or liver problems-- nausea , swelling around your midsection, upper stomach pain, loss of appetite,

Drug Side Effects Summary

dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). AccessPak for HIV PEP Basic affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, sweats, swollen glands, cold sores , cough, wheezing, diarrhea, weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of AccessPak for HIV PEP Basic may include: headache , dizziness, feeling depressed or tired

- trouble sleeping, strange dreams

- nausea, stomach pain

- weight loss

- or rash.

357. This particular disease, AIDS/HIV, interacts with the drug Delstrigo and has the following side effects:

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: urinating more or less than usual

- increased thirst

- new or worsening bone pain

- muscle pain or weakness

- or pain in your arms, legs, hands, or feet. Doravirine, lamivudine, and tenofovir affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen

Drug Side Effects Summary

glands, cold sores , cough, wheezing, diarrhea , weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Delstrigo side effects may include: nausea

- dizziness

- or strange dreams.

358. This particular disease, AIDS/HIV, interacts with the drug efavirenz / lamivudine / tenofovir and

- severe dizziness or drowsiness, trouble concentrating, strange dreams

- confusion, thinking problems, seeing or hearing things that are not real (may occur months to years after you start taking this medicine)

- fast or pounding heartbeats, fluttering in your chest, and sudden dizziness (like you might pass out)

- problems with balance or coordination

- new or worsening bone pain

- pain in your arms, legs, hands, or feet

- lactic acidosis --unusual muscle pain, trouble breathing, stomach pain, vomiting , irregular heart rate, dizziness, feeling cold, or feeling very weak or tired

- pancreatitis --severe pain in your upper stomach spreading to your back, nausea and vomiting

- unusual thoughts--fear, paranoia, feeling sad or hopeless, thoughts of hurting yourself or others

- or liver problems--swelling around your midsection, upper stomach pain, unusual tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Symfi affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen

Drug Side Effects Summary

glands, cold sores , cough, wheezing, diarrhea , weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of efavirenz, lamivudine, and tenofovir may include: headache , dizziness
- depression , thinking problems
- nausea , diarrhea
- sleep problems (insomnia)
- pain
- weakness, tiredness
- cough, stuffy nose, sinus pain
- rash
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

359. This particular disease, AIDS/HIV, interacts with the drug emtricitabine / rilpivirine / tenofovir a

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include: a blistering skin rash , fever, mouth sores, eye redness , swollen glands, trouble breathing or swallowing, right-sided upper stomach pain, unusual bruising, or dark urine. Emtricitabine, rilpivirine, and tenofovir may cause serious side effects. Call your doctor at once if you have: depression or mood changes, anxiety , restlessness, suicidal thoughts or actions
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath

Drug Side Effects Summary

- lactic acidosis --unusual muscle pain, trouble breathing, stomach pain, vomiting , irregular heart rate, dizziness , feeling cold, or feeling very weak or tired
- or liver problems--stomach pain, nausea , loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). This medicine affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of emtricitabine, rilpivirine, and tenofovir may include: headache , dizziness, tiredness
- depressed mood, trouble sleeping, strange dreams
- rash
- or nausea, diarrhea.

360. This particular disease, AIDS/HIV, interacts with the drug Evotaz and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe dizziness , irregular heartbeats
- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- liver or gallbladder problems - upper stomach pain, nausea , vomiting , itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- a severe skin reaction - fever, flu-like symptoms, mouth sores, muscle or joint pain, swelling in your face, burning eyes, skin pain, blistering, warmth or redness under your skin

Drug Side Effects Summary

- or signs of a kidney stone - pain in your side or lower back, painful or difficult urination, blood in your urine. You may need to stop taking this medicine permanently if you have a severe skin reaction. Evotaz affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Evotaz side effects may include: rash
- jaundice
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

361. This particular disease, AIDS/HIV, interacts with the drug Fuzeon and has the following side effects

- skin rash or hives
- fever with vomiting
- blood in your urine
- difficulty breathing
- fainting
- swelling of your feet, face, lips, tongue, or throat. Call your doctor at once if you have: shortness of breath
- fever, chills, chest congestion, cough with yellow or green mucus, stabbing chest pain, wheezing
- fever with vomiting and skin rash
- severe pain in your upper stomach spreading to your back, nausea and vomiting
- blood in your urine

Drug Side Effects Summary

- swelling in your feet
- a skin reaction (itching, redness, swelling, pain, or a hard lump) where the medicine is injected
- or pain, warmth, oozing, or redness where an injection was given, especially if these symptoms get worse or last longer than 7 days. Enfuvirtide affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Fuzeon side effects may include: nausea , loss of appetite, constipation , diarrhea
- numbness or pain in your legs or feet
- muscle pain or weakness
- depressed mood
- swollen glands, runny or stuffy nose
- weight loss
- tiredness, trouble sleeping, tired
- or pain, swelling, burning, or irritation where the medicine was injected.

362. This particular disease, AIDS/HIV, interacts with the drug lamivudine / zidovudine and has the t

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Mild symptoms of lactic acidosis may worsen over time , and this condition can be fatal. Get emergency medical help if you have: unusual muscle pain, trouble breathing, stomach pain, vomiting , irregular heart rate, dizziness, feeling cold, or feeling

Drug Side Effects Summary

very weak or tired. Lamivudine and zidovudine may cause serious side effects. Call your doctor at once if you have: liver problems--swelling around your midsection, right-sided upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)

- low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet

- low white blood cell counts--fever, mouth sores, skin sores, sore throat , cough, trouble breathing

- or pancreatitis --severe pain in your upper stomach spreading to your back, nausea and vomiting .

Lamivudine and zidovudine affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of lamivudine and zidovudine may include: headache

- nausea , diarrhea

- tiredness, general ill feeling

- cold symptoms such as stuffy nose, sneezing, sinus pain, cough

- or loss of body fat (especially in your arms, legs, face, and buttocks).

363. This particular disease, AIDS/HIV, interacts with the drug Pifeltro and has the following side effects

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Doravirine affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor

Drug Side Effects Summary

if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common

Pifeltro side effects may include: nausea , diarrhea, stomach pain

- headache , dizziness

- tiredness

- or strange dreams.

364. This particular disease, AIDS/HIV, interacts with the drug Viracept and has the following side effects:

- hives

- difficulty breathing

- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: low white blood cell counts - fever, mouth sores, skin sores, sore throat , cough

- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor

- or if you have hemophilia - easy bruising, unusual bleeding (nose, mouth, vagina, or rectum), purple or red pinpoint spots under your skin. Nelfinavir can affect your immune system (even weeks

or months after you've used this medicine). Tell your doctor if you have: signs of a new infection -

fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common

Viracept side effects may include: low white blood cell counts

- nausea , diarrhea, gas, stomach pain

Drug Side Effects Summary

- loss of appetite
- rash
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

365. This particular disease, AIDS/HIV, interacts with the drug abacavir / lamivudine / zidovudine and

- Call your doctor at once if you have symptoms of an allergic reaction from two or more of these specific side effect groups: Group 1 - fever

- Group 2 - rash

- Group 3 - nausea, vomiting, diarrhea, stomach pain

- Group 4 - general ill feeling, extreme tiredness, body aches

- Group 5 - shortness of breath, cough, sore throat. Once you have had an allergic reaction to this medicine, you must never use it again. If you stop taking Trizivir for any reason, talk to your doctor before you start taking it again. Also call your doctor at once if you have: other signs of allergic reaction--skin blisters or peeling, eye redness , swelling in your face or throat, trouble breathing

- lactic acidosis --unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness , feeling cold, or feeling very weak or tired

- low white blood cell counts--fever, mouth sores, skin sores, sore throat, cough, trouble breathing

- low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet

- or liver problems--swelling around your midsection, right-sided upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Abacavir, lamivudine, and zidovudine affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea, weight loss

Drug Side Effects Summary

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of abacavir, lamivudine, and zidovudine may include: headache
- weakness or tiredness
- nausea, vomiting
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

366. This particular disease, AIDS/HIV, interacts with the drug Aptivus and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some people taking Aptivus with ritonavir have developed fatal medical problems including liver damage and bleeding in the brain. Stop taking this medicine and get emergency medical attention if you have: right-sided upper stomach pain, nausea, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or unusual bleeding, sudden severe headache, problems with speech or vision. Call your doctor at once if you have: severe skin rash , blistering, peeling, redness or sunburn
- skin rash and also joint or muscle pain, fever, or tightness in your throat
- or increased thirst, increased urination, hunger, dry mouth. Aptivus affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

Drug Side Effects Summary

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Aptivus side effects may include: nausea, vomiting , diarrhea, stomach pain
- skin rash (especially in children)
- headache, fever, tired feeling
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

367. This particular disease, AIDS/HIV, interacts with the drug Cimduo and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: new or worsening bone pain
- muscle weakness
- pain in your arms, hands, legs, or feet
- swelling around your midsection
- kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or signs of liver or pancreas problems - loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting , loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Cimduo affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

Drug Side Effects Summary

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common

Cimduo side effects may include: headache

- pain

- diarrhea

- rash

- depression

- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

368. This particular disease, AIDS/HIV, interacts with the drug cobicistat / darunavir and has the fol

- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor, headache , blurred vision

- or liver problems--upper stomach pain, vomiting, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Cobicistat and darunavir affects your immune system, which may cause certain side effects (even weeks or months after you've taken cobicistat and darunavir).

Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss

- stomach pain, tiredness, itching, skin redness, yellowing of the skin or eyes

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of cobicistat and darunavir may include: nausea , vomiting, stomach pain, diarrhea

- headache

- rash

- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts,

Drug Side Effects Summary

and waist).

369. This particular disease, AIDS/HIV, interacts with the drug cobicistat / elvitegravir / emtricitabine

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Bloody urine bone pain dark urine decreased appetite decreased frequency or amount of urine diarrhea fast, shallow breathing general feeling of discomfort increased thirst light-colored stools loss of appetite lower back or side pain muscle pain or cramping nausea pain in the arms or legs right upper abdominal or stomach pain and fullness sleepiness stomach discomfort swelling of the face, fingers, or lower legs trouble breathing unusual tiredness or weakness vomiting weight gain yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Headache Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Cobicistat / elvitegravir / emtricitabine / tenofovir alafenamide side effects

370. This particular disease, AIDS/HIV, interacts with the drug cobicistat and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Cobicistat may cause serious side effects. Call your doctor at once if you have: kidney problems--little or no urination, swelling in your feet or ankles,

Drug Side Effects Summary

feeling tired or short of breath

- or liver problems--upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of cobicistat may include: nausea
- jaundice
- or rash.

371. This particular disease, AIDS/HIV, interacts with the drug doravirine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Doravirine affects your immune system, which may cause certain side effects (even weeks or months after you've taken doravirine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of doravirine may include: nausea , diarrhea, stomach pain
- headache , dizziness
- tiredness
- or strange dreams.

372. This particular disease, AIDS/HIV, interacts with the drug elvitegravir and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Some side effects may occur that usually do not need medical attention. These side effects may go away during

Drug Side Effects Summary

treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Diarrhea Less common Headache nausea Less common or rare Discouragement feeling sad or empty heartburn indigestion irritability lack of appetite loss of interest or pleasure rash stomach discomfort, upset, or pain thoughts or attempts at killing oneself trouble concentrating trouble sleeping unusual tiredness or weakness vomiting Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Elvitegravir side effects

373. This particular disease, AIDS/HIV, interacts with the drug enfuvirtide and has the following side effects:

- skin rash or hives
- fever with vomiting
- blood in your urine
- difficulty breathing
- fainting
- swelling of your feet, face, lips, tongue, or throat. Enfuvirtide may cause serious side effects. Call your doctor at once if you have: shortness of breath
- fever, chills, chest congestion, cough with yellow or green mucus, stabbing chest pain, wheezing
- fever with vomiting and skin rash
- severe pain in your upper stomach spreading to your back, nausea and vomiting
- blood in your urine
- swelling in your feet
- a skin reaction (itching, redness, swelling, pain, or a hard lump) where the medicine is injected

Drug Side Effects Summary

- or pain, warmth, oozing, or redness where an injection was given, especially if these symptoms get worse or last longer than 7 days. Enfuvirtide affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of enfuvirtide may include: nausea , loss of appetite, constipation , diarrhea
- numbness or pain in your legs or feet
- muscle pain or weakness
- depressed mood
- swollen glands, runny or stuffy nose
- weight loss
- tiredness, trouble sleeping, tired
- or pain, swelling, burning, or irritation where the medicine was injected.

374. This particular disease, AIDS/HIV, interacts with the drug ibalizumab and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Ibalizumab affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats, swollen glands, cold sores , cough, wheezing, diarrhea, weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly

Drug Side Effects Summary

feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of ibalizumab may include: nausea , diarrhea

- dizziness

- or rash.

375. This particular disease, AIDS/HIV, interacts with the drug immune globulin intravenous and ha

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Some side effects may occur during the injection. Tell your caregiver if you feel dizzy, nauseated, light-headed, sweaty, or have a headache , pounding in your neck or ears, fever, chills, chest tightness, or warmth or redness in your face. Immune globulin (intravenous) may cause serious side effects. Call your doctor at once if you have: a blood cell disorder--pale or yellowed skin, dark colored urine, fever, confusion or weakness

- dehydration symptoms--feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin

- kidney problems--little or no urination, swelling, rapid weight gain, feeling short of breath

- lung problems--chest pain, trouble breathing, blue colored lips, fingers, or toes

- signs of a new infection--fever with a severe headache, neck stiffness, eye pain , and increased sensitivity to light

- or signs of a blood clot--shortness of breath, chest pain with deep breathing, rapid heart rate, numbness or weakness on one side of the body, swelling and warmth or discoloration in an arm or leg. Common side effects of immune globulin (intravenous) may include: headache, back pain , joint pain

- fever, chills, sweating, warmth or tingling

Drug Side Effects Summary

- stomach pain, nausea , diarrhea
- increased blood pressure, fast heartbeats
- dizziness , tiredness, lack of energy
- stuffy nose, sinus pain
- or pain, swelling, burning, or irritation around the IV needle.

376. This particular disease, AIDS/HIV, interacts with the drug Invirase and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: heart rhythm problems - fast or pounding heartbeats, fluttering in your chest, shortness of breath, sudden dizziness (like you might pass out)
- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- or liver problems - upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Saquinavir affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Invirase side effects may include: nausea , vomiting , diarrhea, stomach pain
- tired feeling
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

Drug Side Effects Summary

377. This particular disease, AIDS/HIV, interacts with the drug nelfinavir and has the following side

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Nelfinavir may cause serious side effects. Call your doctor at once if you have: low white blood cell counts--fever, mouth sores, skin sores, sore throat , cough
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or if you have hemophilia--easy bruising, unusual bleeding (nose, mouth, vagina, or rectum), purple or red pinpoint spots under your skin. Nelfinavir can affect your immune system (even weeks or months after you've used this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of nelfinavir may include: low white blood cell counts
- nausea , diarrhea, gas, stomach pain
- loss of appetite
- rash
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

378. This particular disease, AIDS/HIV, interacts with the drug rilpivirine and has the following side

- skin rash, fever, swollen glands, flu-like symptoms, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Rilpivirine may cause serious side effects. Call your doctor at once if you have: mood changes, anxiety , feeling sad or hopeless, thoughts about suicide

Drug Side Effects Summary

or hurting yourself

- liver problems--right-sided upper stomach pain, dark urine, jaundice (yellowing of the skin or eyes)
- or symptoms of depression--mood changes, feelings of low self-worth, loss of interest in things you once enjoyed, new sleep problems, thoughts about hurting yourself. Rilpivirine affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of rilpivirine may include: sleep problems (insomnia)
- depression
- skin rash
- headache
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

379. This particular disease, AIDS/HIV, interacts with the drug Rukobia and has the following side effects

- Rukobia can cause serious side effects including: Changes in your immune system (Immune Reconstitution Syndrome) can happen when you start taking HIV-1 medicines. Your immune system may get stronger and begin to fight infections that have been hidden in your body for a long time. Tell your healthcare provider right away if you start having new symptoms after you start taking Rukobia. Heart rhythm problems (QTc prolongation). Rukobia may cause a heart rhythm problem called QTc prolongation. QTc prolongation causes an irregular heartbeat. If you are elderly, you may be at a greater risk for developing this heart problem with Rukobia. Tell your healthcare provider

Drug Side Effects Summary

right away if you feel dizzy, lightheaded, feel changes in your heartbeat, or you faint (lose consciousness). Changes in liver function blood tests results . People with HIV-1 who take Rukobia and who also have hepatitis B or C virus infections, may be more likely to develop new or worsening changes in certain liver function blood tests during treatment. If you stop your anti-hepatitis B treatment, this could mean that your hepatitis B may become active again (reactivated). Your healthcare provider may do blood tests to check your liver during treatment with Rukobia especially if you have hepatitis B virus infection. Take any anti-hepatitis B or anti-hepatitis C medicines as prescribed by your healthcare provider during treatment with Rukobia. The most common side effect is nausea. These are not all of the possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Rukobia side effects

380. This particular disease, AIDS/HIV, interacts with the drug saquinavir and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Saquinavir may cause serious side effects. Call your doctor at once if you have: heart rhythm problems--fast or pounding heartbeats, fluttering in your chest, shortness of breath, sudden dizziness (like you might pass out)
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or liver problems--upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Saquinavir affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

Drug Side Effects Summary

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of saquinavir may include: nausea , vomiting , diarrhea, stomach pain
- tired feeling
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

381. This particular disease, AIDS/HIV, interacts with the drug Symfi and has the following side effects

- severe dizziness or drowsiness, trouble concentrating, strange dreams
- confusion, thinking problems, seeing or hearing things that are not real (may occur months to years after you start taking this medicine)
- fast or pounding heartbeats, fluttering in your chest, and sudden dizziness (like you might pass out)
- problems with balance or coordination
- new or worsening bone pain
- pain in your arms, legs, hands, or feet
- lactic acidosis - unusual muscle pain, trouble breathing, stomach pain, vomiting , irregular heart rate, dizziness, feeling cold, or feeling very weak or tired
- pancreatitis - severe pain in your upper stomach spreading to your back, nausea and vomiting
- unusual thoughts - fear, paranoia, feeling sad or hopeless, thoughts of hurting yourself or others
- or liver problems - swelling around your midsection, upper stomach pain, unusual tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Symfi affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly

Drug Side Effects Summary

feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common

Symfi side effects may include: headache , dizziness

- depression , thinking problems

- nausea , diarrhea

- sleep problems (insomnia)

- pain

- weakness, tiredness

- cough, stuffy nose, sinus pain

- rash

- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

382. This particular disease, AIDS/HIV, interacts with the drug Symfi Lo and has the following side effects:

- severe dizziness or drowsiness, trouble concentrating, strange dreams

- confusion, thinking problems, seeing or hearing things that are not real (may occur months to years after you start taking this medicine)

- fast or pounding heartbeats, fluttering in your chest, and sudden dizziness (like you might pass out)

- problems with balance or coordination

- new or worsening bone pain

- pain in your arms, legs, hands, or feet

- lactic acidosis --unusual muscle pain, trouble breathing, stomach pain, vomiting , irregular heart rate, dizziness, feeling cold, or feeling very weak or tired

- pancreatitis --severe pain in your upper stomach spreading to your back, nausea and vomiting

Drug Side Effects Summary

- unusual thoughts--fear, paranoia, feeling sad or hopeless, thoughts of hurting yourself or others
- or liver problems--swelling around your midsection, upper stomach pain, unusual tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Symfi affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of Symfi Lo may include: headache , dizziness
- depression , thinking problems
- nausea , diarrhea
- sleep problems (insomnia)
- pain
- weakness, tiredness
- cough, stuffy nose, sinus pain
- rash
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

383. This particular disease, AIDS/HIV, interacts with the drug Temixys and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Fever pain stomach pain Less common Chest pain chills cough difficulty in moving joint pain or swelling muscle ache, cramp,

Drug Side Effects Summary

pain, or stiffness sneezing sore throat tightness in the chest troubled breathing Incidence not known Bloating bloody urine blurred vision bone fractures, especially of the femur bone pain cloudy urine constipation dark urine decreased appetite decreased frequency or amount of urine diarrhea difficulty swallowing dizziness dry mouth fast heartbeat fast, shallow breathing flushed, dry skin frequent urination fruit-like breath odor general feeling of discomfort hives, itching, skin rash increased blood pressure increased hunger increased thirst increased urination increased volume of pale, dilute urine indigestion irregular heartbeat large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or genitals light-colored stools loss of appetite lower back or side pain mood changes muscle weakness nausea numbness or tingling in the hands, feet, or lips pains in the stomach, side, or abdomen, possibly radiating to the back pale skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue seizures sleepiness stomach discomfort sweating swelling of the face, fingers, or lower legs troubled breathing with exertion unexplained weight loss unusual bleeding or bruising unusual tiredness or weakness upper right abdominal or stomach pain vomiting weight gain yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Anxiety back pain belching burning, numbness, tingling, or painful sensations discouragement feeling sad or empty headache heartburn indigestion irritability lack or loss of strength loss of interest or pleasure redistribution or accumulation of body fat stomach upset tiredness trouble concentrating trouble sleeping unsteadiness or awkwardness weakness in the arms, hands, legs, or feet Incidence not known Hair loss or thinning of hair Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at

Drug Side Effects Summary

1-800-FDA-1088. Temixys side effects

384. This particular disease, AIDS/HIV, interacts with the drug Trogarzo and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Ibalizumab affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection - fever, night sweats, swollen glands, cold sores , cough, wheezing, diarrhea, weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common Trogarzo side effects may include: nausea , diarrhea
- dizziness
- or rash.

385. This particular disease, AIDS/HIV, interacts with the drug Tybost and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or liver problems - upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common Tybost side effects may include: nausea
- jaundice
- or rash.

Drug Side Effects Summary

386. This particular disease, AIDS/HIV, interacts with the drug AccessPak for HIV PEP Expanded wi

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Early symptoms of lactic acidosis may get worse over time and this condition can be fatal. Get emergency medical help if you have even mild symptoms: muscle pain or weakness, numb or cold feeling in your arms and legs, trouble breathing, stomach pain, nausea with vomiting, fast or uneven heart rate, dizziness, or feeling very weak or tired. AccessPak for HIV PEP Expanded with Kaletra may cause serious side effects. Call your doctor at once if you have: severe dizziness, fainting, fast or pounding heartbeats
- vision changes
- increased urination or extreme thirst
- severe pain in your upper stomach spreading to your back, nausea and vomiting , fast heart rate
- easy bruising, unusual bleeding (nose, mouth, vagina, or rectum), purple or red pinpoint spots under your skin
- penis erection that is painful or lasts longer than 4 hours
- signs of new infection--fever, chills, skin lesions, or cough with yellow or green mucus
- liver problems--nausea, upper stomach pain, itching, tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- kidney problems--little or no urination, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- or severe skin reaction--fever, sore throat , swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of AccessPak for HIV PEP Expanded with Kaletra may include: diarrhea , nausea, vomiting, stomach pain
- headache , dizziness, feeling tired

Drug Side Effects Summary

- depressed mood, sleep problems (insomnia), strange dreams
- mild skin rash
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

387. This particular disease, AIDS/HIV, interacts with the drug AccessPak for HIV PEP Expanded wi

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Mild symptoms of lactic acidosis may worsen over time , and this condition can be fatal. Get emergency medical help if you have: unusual muscle pain, trouble breathing, stomach pain, vomiting , fast/slow or irregular heartbeats, dizziness , feeling cold, or feeling very weak or tired. AccessPak for HIV PEP Expanded with Viracept may cause serious side effects. Call your doctor at once if you have: kidney problems--little or no urination, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- or liver problems-- nausea , swelling around your midsection, upper stomach pain, unusual tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). This medicine affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats, swollen glands, cold sores , cough, wheezing, diarrhea, weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of AccessPak for HIV PEP Expanded with Viracept may include: stomach pain
- headache , dizziness, feeling depressed or tired
- sleep problems (insomnia), strange dreams

Drug Side Effects Summary

- rash
- weight loss
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

388. This particular disease, AIDS/HIV, interacts with the drug atazanavir / cobicistat and has the fo

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Atazanavir and cobicistat may cause serious side effects. Call your doctor at once if you have: severe dizziness , irregular heartbeats
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- liver or gallbladder problems--upper stomach pain, nausea , vomiting , itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- a severe skin reaction--fever, flu-like symptoms, mouth sores, muscle or joint pain, swelling in your face, burning eyes, skin pain, blistering, warmth or redness under your skin
- or signs of a kidney stone--pain in your side or lower back, painful or difficult urination, blood in your urine. You may need to stop taking this medicine permanently if you have a severe skin reaction. Atazanavir and cobicistat affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of atazanavir and cobicistat may include: rash

Drug Side Effects Summary

- jaundice
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

389. This particular disease, AIDS/HIV, interacts with the drug cabotegravir / rilpivirine and has the

- hives
- fever, tiredness, body aches, not feeling well
- sores or blisters in your mouth
- red or puffy eyes
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Your cabotegravir and rilpivirine injections may be permanently discontinued if you have an allergic reaction. Some side effects may occur within a few minutes after an injection. Tell your caregiver if you feel anxious, warm, light-headed, sweaty, or have stomach pain, or numbness in your mouth. Cabotegravir and rilpivirine may cause serious side effects. Call your doctor at once if you have: unusual changes in mood or behavior
- suicidal thoughts or actions
- or liver problems--loss of appetite, nausea , vomiting , stomach pain (upper right side), itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of cabotegravir and rilpivirine may include: pain, redness, swelling, itching, bruising, warmth, or a hard lump where an injection was given
- fever
- nausea

Drug Side Effects Summary

- pain in your bones, joints or muscles
- feeling tired, sleep problems
- headache , dizziness
- or rash.

390. This particular disease, AIDS/HIV, interacts with the drug cabotegravir and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and get medical help at once if you have a skin rash with any of the following symptoms: extreme tiredness, fever, not feeling well
- muscle or joint pain
- blisters or sores in or around your mouth
- red or puffy eyes
- or swelling in your face or mouth, trouble breathing. Cabotegravir may cause serious side effects. Call your doctor at once if you have: unusual changes in mood such as feeling sad, hopeless, anxious, or restless
- suicidal thoughts or actions
- or liver problems--loss of appetite, nausea , vomiting , stomach pain (upper right side), dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of cabotegravir may include: nausea
- abnormal dreams
- feeling anxious
- headache
- or sleep problems.

Drug Side Effects Summary

391. This particular disease, AIDS/HIV, interacts with the drug cobicistat / darunavir / emtricitabine /

- increased thirst, increased urination
- kidney problems--swelling, urinating less, feeling tired or short of breath
- lactic acidosis --unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness , feeling cold, or feeling very weak or tired
- or liver problems-- nausea , upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Symtuza can affect your immune system (even weeks or months after you've used this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of cobicistat, darunavir, emtricitabine, and tenofovir may include: nausea, upset stomach , diarrhea, gas
- headache , feeling tired
- or rash.

392. This particular disease, AIDS/HIV, interacts with the drug dolutegravir / rilpivirine and has the f

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Dolutegravir and rilpivirine may cause serious side effects. Call your doctor at once if you have: thoughts about suicide or hurting yourself

Drug Side Effects Summary

- anxiety , sadness, feeling hopeless
- depression --mood changes, feelings of low self-worth, loss of interest in things you once enjoyed, new sleep problems, thoughts about hurting yourself
- or liver problems-- nausea , vomiting , loss of appetite, right-sided upper stomach pain, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Dolutegravir and rilpivirine affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of dolutegravir and rilpivirine may include: headache
- or nausea, diarrhea.

393. This particular disease, AIDS/HIV, interacts with the drug doravirine / lamivudine / tenofovir and

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Doravirine, lamivudine, and tenofovir may cause serious side effects. Call your doctor at once if you have: urinating more or less than usual
- increased thirst
- new or worsening bone pain
- muscle pain or weakness
- or pain in your arms, legs, hands, or feet. Doravirine, lamivudine, and tenofovir affects your immune system, which may cause certain side effects (even weeks or months after you've taken doravirine, lamivudine, and tenofovir). Tell your doctor if you have: signs of a new infection--fever,

Drug Side Effects Summary

night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of doravirine, lamivudine, and tenofovir may include: nausea

- dizziness

- or strange dreams.

394. This particular disease, AIDS/HIV, interacts with the drug emtricitabine / lopinavir / ritonavir / tenofovir

- hives

- difficulty breathing

- swelling of your face, lips, tongue, or throat. Early symptoms of lactic acidosis may get worse over time and this condition can be fatal. Get emergency medical help if you have even mild symptoms: muscle pain or weakness, numb or cold feeling in your arms and legs, trouble breathing, stomach pain, nausea with vomiting, fast or uneven heart rate, dizziness, or feeling very weak or tired.

Emtricitabine, lopinavir, ritonavir, and tenofovir may cause serious side effects. Call your doctor at once if you have: severe dizziness, fainting, fast or pounding heartbeats

- vision changes

- increased urination or extreme thirst

- severe pain in your upper stomach spreading to your back, nausea and vomiting , fast heart rate

- easy bruising, unusual bleeding (nose, mouth, vagina, or rectum), purple or red pinpoint spots under your skin

- penis erection that is painful or lasts longer than 4 hours

- signs of new infection--fever, chills, skin lesions, or cough with yellow or green mucus

- liver problems--nausea, upper stomach pain, itching, tiredness, loss of appetite, dark urine,

Drug Side Effects Summary

clay-colored stools, jaundice (yellowing of the skin or eyes)

- kidney problems--little or no urination, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- or severe skin reaction--fever, sore throat , swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of emtricitabine, lopinavir, ritonavir, and tenofovir may include: diarrhea , nausea, vomiting, stomach pain
- headache , dizziness, feeling tired
- depressed mood, sleep problems (insomnia), strange dreams
- mild skin rash
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

395. This particular disease, AIDS/HIV, interacts with the drug emtricitabine / nelfinavir / tenofovir a

- hives
 - difficult breathing
 - swelling of your face, lips, tongue, or throat. Mild symptoms of lactic acidosis may worsen over time , and this condition can be fatal. Get emergency medical help if you have: unusual muscle pain, trouble breathing, stomach pain, vomiting , fast/slow or irregular heartbeats, dizziness , feeling cold, or feeling very weak or tired. Emtricitabine, nelfinavir, and tenofovir may cause serious side effects. Call your doctor at once if you have: kidney problems--little or no urination, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
 - or liver problems-- nausea , swelling around your midsection, upper stomach pain, unusual tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes).
- This medicine affects your immune system, which may cause certain side effects (even weeks or

Drug Side Effects Summary

months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats, swollen glands, cold sores , cough, wheezing, diarrhea, weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of emtricitabine, nelfinavir, and tenofovir may include: stomach pain

- headache , dizziness, feeling depressed or tired

- sleep problems (insomnia), strange dreams

- rash

- weight loss

- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

396. This particular disease, AIDS/HIV, interacts with the drug emtricitabine / rilpivirine / tenofovir a

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Discouragement feeling sad or empty irritability lack of appetite loss of interest or pleasure mental depression thoughts of killing oneself tiredness trouble concentrating trouble sleeping Less common Body aches or pain burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain chills cough difficulty with breathing ear congestion fever headache loss of voice runny or stuffy nose sneezing sore throat tightness in the chest unsteadiness or awkwardness unusual tiredness or weakness weakness in the arms, hands, legs, or feet Incidence not known Agitation bloating bloody or cloudy urine confusion constipation dark urine decreased appetite decreased urination diarrhea difficulty swallowing dizziness dry mouth fast heartbeat fast, shallow breathing frequent urination

Drug Side Effects Summary

hostility increased thirst indigestion irritability large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or genitals light-colored stools muscle pain or cramps muscle tenderness, wasting, or weakness nausea pain in the stomach, side, or abdomen, possibly radiating to the back rapid weight gain seizures skin rash, hives, itching sleepiness stomach discomfort swelling of the face, ankles, hands, feet, or lower legs unusual drowsiness, dullness, or feeling of sluggishness vomiting yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Abnormal dreams Less common Back pain belching difficulty with moving heartburn pain in the joints pain or tenderness around the eyes and cheekbones stomach discomfort or upset Incidence not known Lack or loss of strength Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Emtricitabine / rilpivirine / tenofovir alafenamide side effects

397. This particular disease, AIDS/HIV, interacts with the drug fosamprenavir and has the following

- pale or yellowed skin, dark colored urine, fever, confusion or weakness
- increased urination or extreme thirst
- or high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or symptoms of kidney stones --sudden and severe pain in your lower back or side, blood in your urine, pain or burning when you urinate. Fosamprenavir affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough,

Drug Side Effects Summary

wheezing, diarrhea , weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of fosamprenavir may include: fever or other signs of infection

- nausea , vomiting , diarrhea

- rash

- headache

- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

398. This particular disease, AIDS/HIV, interacts with the drug fostemsavir and has the following side effects:

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Fostemsavir may cause serious side effects. Call your doctor at once if you have: fast or pounding heartbeats, fluttering in your chest

- shortness of breath

- and sudden dizziness (like you might pass out). Fostemsavir affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Nausea is the most common side effect of fostemsavir.

Drug Side Effects Summary

399. This particular disease, AIDS/HIV, interacts with the drug lamivudine / nevirapine / zidovudine .

- No side effects listed

400. This particular disease, AIDS/HIV, interacts with the drug tipranavir and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some people taking tipranavir with ritonavir have developed fatal medical problems including liver damage and bleeding in the brain. Stop taking this medicine and get emergency medical attention if you have: right-sided upper stomach pain, nausea, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or unusual bleeding, sudden severe headache, problems with speech or vision. Tipranavir may cause serious side effects. Call your doctor at once if you have: severe skin rash , blistering, peeling, redness or sunburn
- skin rash and also joint or muscle pain, fever, or tightness in your throat
- or increased thirst, increased urination, hunger, dry mouth. Tipranavir affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of tipranavir may include: nausea, vomiting , diarrhea, stomach pain
- skin rash (especially in children)
- headache, fever, tired feeling
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts,

Drug Side Effects Summary

and waist).

401. This particular disease, AIDS/HIV, interacts with the drug Triumeq PD and has the following side effects:

- Call your doctor at once if you have symptoms of an allergic reaction from two or more of these specific side effect groups: Group 1 - fever

- Group 2 - rash

- Group 3 - nausea, vomiting, diarrhea, stomach pain

- Group 4 - general ill feeling, extreme tiredness, body aches

- Group 5 - shortness of breath, cough, sore throat. Once you have had an allergic reaction to a medicine that contains abacavir or dolutegravir, you must never use it again. If you stop taking Triumeq PD for any reason, talk to your doctor before you start taking it again. Triumeq PD may cause serious side effects. Call your doctor at once if you have: other signs of allergic reaction--skin blisters or peeling, eye redness , swelling in your face or throat, trouble breathing

- lactic acidosis --unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness , feeling cold, or feeling very weak or tired

- or liver problems--swelling around your midsection, right-sided upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Triumeq PD affects your immune system, which may cause certain side effects (even weeks or months after you've taken this medicine). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea, weight loss

- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling

- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of Triumeq PD may include: headache

- tiredness

Drug Side Effects Summary

- or trouble sleeping.

402. This particular disease, AIDS/HIV, interacts with the drug Vocabria and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and get medical help at once if you have a skin rash with any of the following symptoms: extreme tiredness, fever, not feeling well
- muscle or joint pain
- blisters or sores in or around your mouth
- red or puffy eyes
- or swelling in your face or mouth, trouble breathing. Vocabria may cause serious side effects. Call your doctor at once if you have: unusual changes in mood such as feeling sad, hopeless, anxious, or restless
- suicidal thoughts or actions
- or liver problems - loss of appetite, nausea , vomiting , stomach pain (upper right side), dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common Vocabria side effects of Vocabria may include: nausea
- abnormal dreams
- feeling anxious
- headache
- or sleep problems.

403. This particular disease, Allergies, interacts with the drug hydroxyzine and has the following side effects:

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. In rare cases, hydroxyzine may cause a severe skin reaction. Stop taking this medicine and call your doctor right away if you have sudden skin redness or a rash that spreads and causes white or yellow pustules, blistering, or peeling. Stop using hydroxyzine and call your doctor at once if you have: fast or pounding heartbeats
- headache with chest pain
- severe dizziness, fainting
- or a seizure (convulsions). Side effects such as drowsiness and confusion may be more likely in older adults. Common hydroxyzine side effects may include: drowsiness
- headache
- dry mouth
- or skin rash.

404. This particular disease, Allergies, interacts with the drug levocetirizine and has the following s

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Levocetirizine may cause serious side effects. Stop using levocetirizine and call your doctor at once if you have: worsening allergy or urticaria symptoms
- painful or difficult urination
- little or no urination
- a light-headed feeling, like you might pass out
- fever
- or signs of an ear infection--ear pain or full feeling, trouble hearing, drainage from the ear, fussiness in a child. Common side effects of levocetirizine may include: drowsiness, tiredness
- sinus pain
- ear infection

Drug Side Effects Summary

- cough
- fever
- nosebleed
- vomiting , diarrhea , constipation
- dry mouth
- or weight gain.

405. This particular disease, Allergies, interacts with the drug loratadine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: fast or uneven heart rate
- severe headache
- or a light-headed feeling, like you might pass out
- Common loratadine side effects may include: headache
- feeling tired or drowsy
- stomach pain, vomiting
- dry mouth
- or feeling nervous or hyperactive.

406. This particular disease, Allergies, interacts with the drug Xyzal and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: worsening allergy or urticaria symptoms

Drug Side Effects Summary

- painful or difficult urination
- little or no urination
- a light-headed feeling, like you might pass out
- fever
- or signs of an ear infection - ear pain or full feeling, trouble hearing, drainage from the ear, fussiness in a child. Common Xyzal side effects may include: drowsiness, tiredness
- sinus pain
- ear infection
- cough
- fever
- nosebleed
- vomiting , diarrhea , constipation
- dry mouth
- or weight gain .

407. This particular disease, Allergies, interacts with the drug doxylamine and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Doxylamine may cause serious side effects. Call your doctor at once if you have: confusion, hallucinations
- severe dizziness or drowsiness
- or little or no urinating. Common side effects of doxylamine may include: blurred vision
- dry mouth, nose, or throat
- constipation
- or mild dizziness or drowsiness. Side effects such as dry mouth, constipation, and confusion may

Drug Side Effects Summary

be more likely in older adults.

408. This particular disease, Allergies, interacts with the drug corticotropin and has the following side effects:

- hives
- nausea , feeling light-headed
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Corticotropin may cause serious side effects. Call your doctor at once if you have: fever, chills, sore throat , skin warmth or redness, or other signs of an infection
- unusual changes in mood or behavior
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- a seizure
- decreased or increased adrenal gland hormones --changes in weight or appetite, tiredness, muscle weakness, skin discoloration, thinning skin, increased body hair, menstrual changes, loss of appetite, stomach pain
- high blood pressure--severe headache , blurred vision, pounding in your neck or ears
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- signs of eye infection--swelling, redness, severe discomfort, crusting or drainage
- or signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Be sure your family, caregivers, and close friends also know how to help you watch for these side effects. Corticotropin can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using corticotropin. Common side effects of corticotropin may include:
increased blood pressure

Drug Side Effects Summary

- thinning skin, increased sweating
- fluid retention (swelling in your hands or feet, puffiness in your face)
- mood changes, irritability
- increased appetite
- or weight gain.

409. This particular disease, Allergies, interacts with the drug Acthar and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of infection like fever, chills, very bad sore throat, ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal. Signs of high blood sugar like confusion, feeling sleepy, more thirst, more hungry, passing urine more often, flushing, fast breathing, or breath that smells like fruit. Signs of a weak adrenal gland like a very bad upset stomach or throwing up, very bad dizziness or passing out, muscle weakness, feeling very tired, mood changes, not hungry, or weight loss. Signs of Cushing's disease like weight gain in the upper back or belly, moon face, very

Drug Side Effects Summary

bad headache , or slow healing. Signs of high or low blood pressure like very bad headache or dizziness, passing out, or change in eyesight. Signs of low potassium levels like muscle pain or weakness, muscle cramps, or a heartbeat that does not feel normal. Shortness of breath, a big weight gain, or swelling in the arms or legs. Stomach pain. Black, tarry, or bloody stools. Throwing up blood or throw up that looks like coffee grounds. Change in eyesight. Mental, mood, or behavior changes that are new or worse. Bone or joint pain. Seizures . Acthar side effects

410. This particular disease, Allergies, interacts with the drug chlorpheniramine / ibuprofen / pheny

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Bloody or black, tarry stools chest pain or tightness confusion constipation cough decrease urine output diarrhea difficulty in speaking difficulty swallowing dilated neck veins dizziness double vision fast heartbeat headache heartburn hives, itching, skin rash inability to speak inability to move the arms, legs, or facial muscles indigestion irregular breathing nausea puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue severe stomach pain, cramping, or burning swelling of the face, fingers, feet, or lower legs trouble breathing unusual tiredness or weakness vomiting of blood or material that looks like coffee grounds weight gain Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Nervousness trouble sleeping unusual excitement, especially in children Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side

Drug Side Effects Summary

effects to the FDA at 1-800-FDA-1088. Chlorpheniramine / ibuprofen / phenylephrine side effects

411. This particular disease, Allergies, interacts with the drug dextbrompheniramine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Dextbrompheniramine may cause serious side effects. Stop using dextbrompheniramine and call your doctor at once if you have: fast or uneven heart rate
- mood changes
- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- feeling short of breath
- or little or no urination. Common side effects of dextbrompheniramine may include: drowsiness
- dry mouth, nose, or throat
- constipation
- blurred vision
- or feeling restless or excited (especially in children).

412. This particular disease, Allergies, interacts with the drug Ala-Hist IR and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Ala Hist IR may cause serious side effects. Stop using Ala Hist IR and call your doctor at once if you have: fast or uneven heart rate
- mood changes
- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness

Drug Side Effects Summary

- feeling short of breath
- or little or no urination. Common side effects of Ala Hist IR may include: drowsiness
- dry mouth, nose, or throat
- constipation
- blurred vision
- or feeling restless or excited (especially in children).

413. This particular disease, Allergies, interacts with the drug dexbrompheniramine / phenylephrine

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: fast or irregular heartbeats
- confusion, hallucinations
- a seizure
- ringing in your ears
- or little or no urination. Common side effects of dexbrompheniramine and phenylephrine may include: headache, dizziness , drowsiness
- blurred vision, double vision
- dry mouth, nose, or throat
- constipation , upset stomach
- or trouble sleeping.

414. This particular disease, Allergies, interacts with the drug Equaline Sleep Aid and has the follow

- No side effects listed

Drug Side Effects Summary

415. This particular disease, Allergies, interacts with the drug Equate Sleep Aid and has the following

- No side effects listed

416. This particular disease, Allergies, interacts with the drug H.P. Acthar Gel and has the following

- hives
- nausea , feeling light-headed
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Acthar Gel, H.P. may cause serious side effects. Call your doctor at once if you have: fever, chills, sore throat , skin warmth or redness, or other signs of an infection
- unusual changes in mood or behavior
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- a seizure
- decreased or increased adrenal gland hormones --changes in weight or appetite, tiredness, muscle weakness, skin discoloration, thinning skin, increased body hair, menstrual changes, loss of appetite, stomach pain
- high blood pressure--severe headache , blurred vision, pounding in your neck or ears
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- signs of eye infection--swelling, redness, severe discomfort, crusting or drainage
- or signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Be sure your family, caregivers, and close friends also know how to help you watch for these side effects. Acthar Gel, H.P. can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using Acthar Gel, H.P.. Common side effects of Acthar Gel, H.P.

Drug Side Effects Summary

may include: increased blood pressure

- thinning skin, increased sweating
- fluid retention (swelling in your hands or feet, puffiness in your face)
- mood changes, irritability
- increased appetite
- or weight gain.

417. This particular disease, Alzheimer's, interacts with the drug Aricept and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Aricept and call your doctor at once if you have: severe or ongoing vomiting
- a light-headed feeling, like you might pass out
- slow heartbeats
- seizure (convulsions)
- painful or difficult urination
- new or worsening breathing problems
- or signs of stomach bleeding--severe heartburn or stomach pain, bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Common Aricept side effects may include: nausea, vomiting, diarrhea
- loss of appetite
- muscle pain
- sleep problems (insomnia)
- or feeling tired
- .

Drug Side Effects Summary

418. This particular disease, Alzheimer's, interacts with the drug donepezil and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Donepezil may cause serious side effects. Call your doctor at once if you have: slow heartbeats
- a light-headed feeling, like you might pass out
- new or worsening stomach pain, heartburn , nausea, or vomiting
- a seizure
- painful or difficult urination
- new or worsening breathing problems
- or signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Common side effects of donepezil may include: nausea, vomiting, diarrhea
- loss of appetite
- muscle pain
- sleep problems (insomnia)
- or feeling tired.

419. This particular disease, Alzheimer's, interacts with the drug Namenda and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe headache, blurred vision, pounding in your neck or ears
- seizure (convulsions)
- or unusual changes in mood or behavior. Common Namenda side effects may include: diarrhea
- dizziness

Drug Side Effects Summary

- or headache.

420. This particular disease, Alzheimer's, interacts with the drug memantine and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Memantine may cause serious side effects. Call your doctor at once if you have: severe headache , blurred vision, pounding in your neck or ears
- seizure (convulsions)
- or unusual changes in mood or behavior. Common side effects of memantine may include:
diarrhea
- dizziness
- or headache.

421. This particular disease, Alzheimer's, interacts with the drug etanercept and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Serious and sometimes fatal infections may occur. Call your doctor right away if you have: fever, chills, flu symptoms
- pale skin, easy bruising or bleeding
- pain, redness, or swelling where etanercept was injected (for longer than 5 days after injection)
- signs of lymphoma--fever, night sweats, weight loss, stomach pain or swelling, swollen glands (in your neck, armpits, or groin)
- signs of tuberculosis--cough, night sweats, loss of appetite, weight loss, feeling very tired
- new or worsening psoriasis --skin redness or scaly patches, raised bumps filled with pus
- nerve problems-- dizziness , numbness or tingling, problems with vision, or weak feeling in your

Drug Side Effects Summary

arms or legs

- signs of heart failure--shortness of breath, swelling in your lower legs
- lupus -like syndrome--joint pain or swelling, chest discomfort, feeling short of breath, skin rash on your cheeks or arms (worsens in sunlight)
- o liver problems--right-sided upper stomach pain, vomiting , tiredness, loss of appetite, yellowing of your skin or eyes. Common side effects of etanercept may include: pain, swelling, itching, or redness where the medicine was injected
- or cold symptoms such as stuffy nose, sneezing, sore throat .

422. This particular disease, Alzheimer's, interacts with the drug Exelon and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe or ongoing vomiting or diarrhea, loss of appetite, weight loss
- bloody or tarry stools, cough with bloody mucus or vomit that looks like coffee grounds
- a light-headed feeling, like you might pass out
- tremors (uncontrolled shaking), restless muscle movements in your eyes, tongue, jaw, or neck
- seizure (convulsions)
- painful or difficult urination
- severe skin redness, itching, or irritation
- or dehydration symptoms - feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin . Common Exelon side effects may include: upset stomach , nausea, vomiting
- loss of appetite
- or weakness.

Drug Side Effects Summary

423. This particular disease, Alzheimer's, interacts with the drug galantamine and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Galantamine can cause serious skin reactions. Stop using galantamine and call your doctor at once if you have the first signs of any skin rash , no matter how mild. Galantamine may cause serious side effects. Call your doctor at once if you have: chest pain, slow heart rate
- little or no urinating
- blood in your urine
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems--nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- dehydration symptoms--feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin
- a light-headed feeling, like you might pass out
- or severe skin reaction--fever, sore throat , swelling in your face or tongue, burning in your eyes, skin pain, followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of galantamine may include: nausea, vomiting, diarrhea , loss of appetite
- headache , dizziness
- or weight loss .

424. This particular disease, Alzheimer's, interacts with the drug Namenda XR and has the following

- hives

Drug Side Effects Summary

- difficulty breathing
- swelling of your face, lips, tongue, or throat. Namenda XR may cause serious side effects. Call your doctor at once if you have: severe headache , blurred vision, pounding in your neck or ears
- seizure (convulsions)
- or unusual changes in mood or behavior. Common side effects of Namenda XR may include:
diarrhea
- dizziness
- or headache.

425. This particular disease, Alzheimer's, interacts with the drug rivastigmine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Rivastigmine may cause serious side effects. Call your doctor at once if you have: severe or ongoing vomiting or diarrhea, loss of appetite, weight loss
- bloody or tarry stools, cough with bloody mucus or vomit that looks like coffee grounds
- a light-headed feeling, like you might pass out
- tremors (uncontrolled shaking), restless muscle movements in your eyes, tongue, jaw, or neck
- seizure (convulsions)
- painful or difficult urination
- severe skin redness, itching, or irritation
- or dehydration symptoms--feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin . Common side effects of rivastigmine may include: upset stomach , nausea, vomiting
- loss of appetite
- or weakness.

Drug Side Effects Summary

426. This particular disease, Alzheimer's, interacts with the drug Aricept ODT and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Aricept ODT may cause serious side effects. Call your doctor at once if you have: slow heartbeats
- a light-headed feeling, like you might pass out
- new or worsening stomach pain, heartburn , nausea, or vomiting
- a seizure
- painful or difficult urination
- new or worsening breathing problems
- or signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Common side effects of Aricept ODT may include: nausea, vomiting, diarrhea
- loss of appetite
- muscle pain
- sleep problems (insomnia)
- or feeling tired.

427. This particular disease, Alzheimer's, interacts with the drug Namzaric and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: severe or ongoing vomiting
- a light-headed feeling, like you might pass out
- slow heartbeats or chest pain
- seizure (convulsions)

Drug Side Effects Summary

- painful or difficult urination
- new or worsening breathing problems
- or signs of stomach bleeding - severe heartburn or stomach pain, bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Common Namzaric side effects may include: nausea, vomiting, diarrhea, loss of appetite
- headache
- dizziness
- or easy bruising.

428. This particular disease, Alzheimer's, interacts with the drug vitamin e and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop taking vitamin E and call your doctor at once if you have: headache, dizziness, vision changes
- a light-headed feeling, like you might pass out
- unusual weakness or tired feeling
- diarrhea, stomach cramps
- or easy bruising or bleeding (nosebleeds, bleeding gums). Common vitamin E side effects may include: nausea
- tired feeling
- headache
- or mild rash.

429. This particular disease, Alzheimer's, interacts with the drug donepezil / memantine and has the following side effects:

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. donepezil and memantine may cause serious side effects. Stop using donepezil and memantine and call your doctor at once if you have: severe or ongoing vomiting
- a light-headed feeling, like you might pass out
- slow heartbeats
- seizure (convulsions)
- painful or difficult urination
- new or worsening breathing problems
- or signs of stomach bleeding--severe heartburn or stomach pain, bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Common side effects of donepezil and memantine may include: nausea , vomiting, diarrhea , loss of appetite
- headache
- dizziness
- or easy bruising.

430. This particular disease, Alzheimer's, interacts with the drug Razadyne and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Razadyne can cause serious skin reactions. Stop using Razadyne and call your doctor at once if you have the first signs of any skin rash , no matter how mild. Razadyne may cause serious side effects. Call your doctor at once if you have: chest pain, slow heart rate
- little or no urinating
- blood in your urine

Drug Side Effects Summary

- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems--nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- dehydration symptoms--feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin
- a light-headed feeling, like you might pass out
- or severe skin reaction--fever, sore throat , swelling in your face or tongue, burning in your eyes, skin pain, followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of Razadyne may include: nausea, vomiting, diarrhea , loss of appetite
- headache , dizziness
- or weight loss .

431. This particular disease, Alzheimer's, interacts with the drug Razadyne ER and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Razadyne ER can cause serious skin reactions. Stop using Razadyne ER and call your doctor at once if you have the first signs of any skin rash , no matter how mild. Razadyne ER may cause serious side effects. Call your doctor at once if you have: chest pain, slow heart rate
- little or no urinating
- blood in your urine
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds

Drug Side Effects Summary

- liver problems--nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- dehydration symptoms--feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin
- a light-headed feeling, like you might pass out
- or severe skin reaction--fever, sore throat , swelling in your face or tongue, burning in your eyes, skin pain, followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of Razadyne ER may include: nausea, vomiting, diarrhea , loss of appetite
- headache , dizziness
- or weight loss .

432. This particular disease, Alzheimer's, interacts with the drug Adlarity and has the following side effects:

- Adlarity may cause serious side effects, including: Skin reactions. Skin reactions that include redness and itching may happen at the application site. Stop treatment and call your healthcare provider if you get any of these skin reactions and they do not get better within 2 days (48 hours) after the transdermal system is removed: increased redness or swelling peeling or blistering of the skin spreading beyond the application site Slow heartbeat and fainting. Call your healthcare provider right away if you feel faint or lightheaded. More stomach acid. This increases the chance of ulcers and bleeding. The risk is higher for some people, such as those who have had ulcers or take NSAIDs. Call your healthcare provider right away if you have any of these symptoms: heartburn or stomach pain that is new or does not go away. nausea or vomiting, blood in your vomit, or dark vomit that looks like coffee grounds. bowel movements or stools that look like black tar. Problems passing urine. Call your healthcare provider right away if you have problems passing urine. Seizures. Call your healthcare provider right away if you have seizures. Worsening of lung problems

Drug Side Effects Summary

in people with asthma or other lung disease. Call your healthcare provider if you have new or worsening lung problems. The most common side effects of donepezil, the medicine in Adlarity, are: nausea diarrhea not sleeping well vomiting muscle cramps feeling tired not wanting to eat These are not all of the possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Adlarity side effects

433. This particular disease, Alzheimer's, interacts with the drug Hydergine and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Hydergine may cause serious side effects. Call your doctor at once if you have: any changes in memory, alertness, mood, appetite, energy level, or ability to care for yourself. Common side effects of Hydergine may include: nausea or other stomach discomfort.

434. This particular disease, Alzheimer's, interacts with the drug Alpha E and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: With doses greater than 400 Units a day and long-term use Blurred vision diarrhea dizziness headache nausea or stomach cramps unusual tiredness or weakness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Alpha E side effects

435. This particular disease, Alzheimer's, interacts with the drug ergoloid mesylates and has the following side effects

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Ergoloid mesylates may cause serious side effects. Call your doctor at once if you have: any changes in memory, alertness, mood, appetite, energy level, or ability to care for yourself. Common side effects of ergoloid mesylates may include: nausea or other stomach discomfort.

436. This particular disease, Alzheimer's, interacts with the drug Aduhelm and has the following side effects:

- Aduhelm can cause serious side effects, including: See above Important information. Serious allergic reactions. Swelling of the face, lips, mouth, or tongue and hives have happened during an Aduhelm infusion. Tell your healthcare provider if you have any of the symptoms of a serious allergic reaction during or after your infusion. The most common side effects include: swelling in areas of the brain, with or without small spots of bleeding in or on the surface of the brain (ARIA) headache fall
- Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA1088. Aduhelm side effects

437. This particular disease, Alzheimer's, interacts with the drug Aqua Gem-E and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: With doses greater than 400 Units a day and long-term use Blurred vision diarrhea dizziness headache nausea or stomach cramps unusual tiredness or weakness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Aqua Gem-E side effects

Drug Side Effects Summary

438. This particular disease, Alzheimer's, interacts with the drug Aquasol E and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: With doses greater than 400 Units a day and long-term use Blurred vision diarrhea dizziness headache nausea or stomach cramps unusual tiredness or weakness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Aquasol E side effects

439. This particular disease, Alzheimer's, interacts with the drug E-400 Clear and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: With doses greater than 400 Units a day and long-term use Blurred vision diarrhea dizziness headache nausea or stomach cramps unusual tiredness or weakness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

E-400 Clear side effects

440. This particular disease, Alzheimer's, interacts with the drug E-600 and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: With doses greater than 400 Units a day and long-term use Blurred vision diarrhea dizziness headache nausea or stomach

Drug Side Effects Summary

cramps unusual tiredness or weakness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

E-600 side effects

441. This particular disease, Alzheimer's, interacts with the drug E-Gems and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: With doses greater than 400 Units a day and long-term use Blurred vision diarrhea dizziness headache nausea or stomach cramps unusual tiredness or weakness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

E-Gems side effects

442. This particular disease, Alzheimer's, interacts with the drug E Pherol and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat

Drug Side Effects Summary

- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. E Pherol side effects

443. This particular disease, Alzheimer's, interacts with the drug aducanumab and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some side effects may occur during the injection. Tell your care provider if you have a rash, trouble breathing, or swelling in your face. Aducanumab can cause temporary swelling or bleeding in the brain. Call your doctor if you have: headaches, dizziness , confusion
- nausea
- or vision changes. Common side effects of aducanumab may include: ARIA with symptoms, or with signs that appear on an MRI
- headache
- or falls.

444. This particular disease, Angina, interacts with the drug aspirin and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Aspirin may cause serious side effects. Stop using this medicine and call your doctor at once if you have: ringing in your ears, confusion, hallucinations, rapid breathing, seizure (convulsions)
- severe nausea, vomiting, or stomach pain
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds

Drug Side Effects Summary

- fever lasting longer than 3 days
- or swelling, or pain lasting longer than 10 days. Common side effects of aspirin may include: upset stomach
- heartburn
- drowsiness
- or mild headache.

445. This particular disease, Angina, interacts with the drug nitroglycerin and has the following side effects:

- hives , sweating, pale skin
- nausea , vomiting
- feeling weak or light-headed
- difficult breathing
- swelling of your face, lips, tongue, or throat. Nitroglycerin may cause serious side effects. Call your doctor at once if you have: severe or throbbing headaches that do not become less severe with continued use of nitroglycerin
- pounding heartbeats or fluttering in your chest
- slow heart rate
- a light-headed feeling, like you might pass out
- blurred vision or dry mouth
- or heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating. Nitroglycerin can cause severe headaches that should become less severe as you continue to use the medicine. Common side effects of nitroglycerin may include: flushing (sudden warmth, redness, or tingly feeling)
- feeling light-headed, fainting
- headache, dizziness

Drug Side Effects Summary

- or numbness, tingling, burning pain.

446. This particular disease, Angina, interacts with the drug Ranexa and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- headache with chest pain and severe dizziness, fast or pounding heartbeats
- or kidney problems - little or no urination
- painful or difficult urination, swelling in your feet or ankles
- feeling tired or short of breath. Common Ranexa side effects may include: nausea, constipation
- headache
- or dizziness.

447. This particular disease, Angina, interacts with the drug metoprolol and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Metoprolol may cause serious side effects. Call your doctor at once if you have: very slow heartbeats
- a light-headed feeling, like you might pass out
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- or cold feeling in your hands and feet. Common side effects of metoprolol may include: dizziness
- tired feeling
- depression
- confusion

Drug Side Effects Summary

- memory problems
- nightmares
- trouble sleeping
- diarrhea
- mild itching or rash.

448. This particular disease, Angina, interacts with the drug isosorbide mononitrate and has the fol

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- worsening angina pain
- fast or slow heart rate
- or pounding heartbeats or fluttering in your chest. Isosorbide mononitrate can cause severe headaches. These headaches may gradually become less severe as you continue to use nitroglycerin. Do not stop taking this medicine to avoid headaches. Ask your doctor before using any headache pain medication. Common isosorbide mononitrate side effects may include: headache
- dizziness upon standing
- or flushing (warmth, redness, or tingly feeling).

449. This particular disease, Angina, interacts with the drug amlodipine and has the following side e

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, when you first start taking amlodipine, your angina may get worse or you could have a heart attack. Seek emergency medical attention or

Drug Side Effects Summary

call your doctor right away if you have symptoms such as: chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating. Amlodipine may cause serious side effects. Call your doctor at once if you have: pounding heartbeats or fluttering in your chest

- worsening chest pain
- swelling in your feet or ankles
- severe drowsiness
- or a light-headed feeling, like you might pass out. Common side effects of amlodipine may include:
dizziness
- drowsiness
- feeling tired
- stomach pain
- nausea
- flushing (warmth, redness, or tingly feeling).

450. This particular disease, Angina, interacts with the drug Nitrostat and has the following side effects:

- hives , sweating, pale skin
- nausea , vomiting
- feeling weak or light-headed
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe or throbbing headaches that do not become less severe with continued use of Nitrostat
- pounding heartbeats or fluttering in your chest
- slow heart rate
- a light-headed feeling, like you might pass out
- blurred vision or dry mouth

Drug Side Effects Summary

- or heart attack symptoms - chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating. Nitroglycerin can cause severe headaches that should become less severe as you continue to use the medicine. Common Nitrostat side effects may include: flushing (sudden warmth, redness, or tingly feeling)
- feeling light-headed, fainting
- headache, dizziness
- or numbness, tingling, burning pain.

451. This particular disease, Angina, interacts with the drug ranolazine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Ranolazine may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- fast or pounding heartbeats, fluttering in your chest
- or kidney problems--little or no urination, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath. Common side effects of ranolazine may include: nausea, constipation
- headache
- or dizziness.

452. This particular disease, Angina, interacts with the drug enoxaparin and has the following side effects:

- hives
- itching or burning skin
- difficult breathing
- swelling of your face, lips, tongue, or throat. Also seek emergency medical attention if you have symptoms of a spinal blood clot : back pain, numbness or muscle weakness in your lower body, or

Drug Side Effects Summary

loss of bladder or bowel control. Enoxaparin may cause serious side effects. Call your doctor at once if you have: unusual bleeding, or any bleeding that will not stop

- easy bruising, purple or red spots under your skin
- nosebleeds , bleeding gums
- abnormal vaginal bleeding , blood in your urine or stools
- coughing up blood or vomit that looks like coffee grounds
- signs of bleeding in the brain--sudden weakness (especially on one side of the body), sudden severe headache , problems with speech or vision
- or low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Common side effects of enoxaparin may include: nausea , diarrhea
- anemia
- confusion
- or pain, bruising, redness, or irritation where the medicine was injected.

453. This particular disease, Angina, interacts with the drug isosorbide dinitrate and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Isosorbide dinitrate may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- worsening angina pain
- fast or slow heart rate
- or pounding heartbeats or fluttering in your chest. Isosorbide dinitrate can cause severe headaches. These headaches may gradually become less severe as you continue to use nitroglycerin. Do not stop taking isosorbide dinitrate to avoid headaches. Ask your doctor before using any headache pain medication. Common side effects may be more likely to occur, such as:

Drug Side Effects Summary

headache

- or flushing (warmth, redness, or tingly feeling).

454. This particular disease, Angina, interacts with the drug atenolol and has the following side effects:

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: new or worsening chest pain

- slow or uneven heartbeats

- a light-headed feeling, like you might pass out

- shortness of breath (even with mild exertion), swelling, rapid weight gain

- or a cold feeling in your hands and feet. Common atenolol side effects may include include: cold hands or feet

- dizziness

- tiredness

- or depressed mood.

455. This particular disease, Angina, interacts with the drug carvedilol and has the following side effects:

- hives

- difficulty breathing

- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have a serious side effect such as: a light-headed feeling, like you might pass out

- slow or uneven heartbeats

- swelling, rapid weight gain, feeling short of breath (even with mild exertion)

- cold feeling or numbness in your fingers or toes

Drug Side Effects Summary

- chest pain, dry cough, wheezing, chest tightness, trouble breathing
- or high blood sugar (increased thirst, increased urination, hunger, dry mouth, fruity breath odor, drowsiness, dry skin, blurred vision, weight loss). Common carvedilol side effects may include: weakness, dizziness
- diarrhea
- dry eyes
- tired feeling
- or weight gain.

456. This particular disease, Angina, interacts with the drug Norvasc and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, when you first start taking Norvasc, your angina may get worse or you could have a heart attack. Seek emergency medical attention or call your doctor right away if you have symptoms such as: chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating. Call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- worsening chest pain
- swelling in your feet or ankles
- severe drowsiness
- or a light-headed feeling, like you might pass out. Common Norvasc side effects may include: dizziness, drowsiness
- feeling tired
- stomach pain, nausea
- or flushing (warmth, redness, or tingly feeling)

Drug Side Effects Summary

- swelling in your ankles and lower part of your legs.

457. This particular disease, Angina, interacts with the drug Lopressor and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Blurred vision chest pain or discomfort confusion dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position slow or irregular heartbeat sweating unusual tiredness or weakness Less common Bloating or swelling of the face, arms, hands, lower legs, or feet chest tightness cough decreased urine output difficult or labored breathing difficulty with speaking dilated neck veins disturbed color perception double vision fast, pounding, or racing heartbeat or pulse halos around lights headache inability to move the arms, legs, or facial muscles inability to speak irregular breathing loss of vision night blindness noisy breathing overbright appearance of lights pain, tension, and weakness upon walking that subsides during periods of rest paleness or cold feeling in the fingertips and toes rapid weight gain seeing, hearing, or feeling things that are not there short-term memory loss slow speech swelling of the face, fingers, feet, or lower legs tingling of the hands or feet tingling or pain in the fingers or toes when exposed to cold temperatures trouble breathing tunnel vision unusual weight gain or loss Rare Bluish color of the skin of the fingers or toes chills clay-colored stools continuing loss of appetite continuing or severe abdominal or stomach pain continuing or severe nausea and vomiting dark urine difficulty with moving fever hoarseness increased frequency of urination itching skin light-colored stools lower back or side pain muscle pain or stiffness numbness of the fingers or toes pain, swelling, or redness in the joints rash sore throat sores, ulcers, or white spots on the lips or in the mouth unpleasant breath odor unusual bleeding or bruising upper right abdominal or stomach pain vomiting of blood weakness yellow eyes and skin Incidence not known Black, tarry stools bleeding gums blood in the urine or stools burning, crawling,

Drug Side Effects Summary

itching, numbness, prickling, "pins and needles", or tingling feelings cool, sweaty skin pinpoint red spots on the skin Symptoms of overdose Bluish color of the fingernails, lips, skin, palms, or nail beds change in consciousness loss of consciousness no blood pressure or pulse stopping of the heart very drowsy or sleepy Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Belching bloated feeling decreased interest in sexual intercourse difficulty having a bowel movement discouragement dry mouth excess air or gas in stomach or bowels feeling of constant movement of self or surroundings feeling of fullness feeling of indigestion feeling sad or empty inability to have or keep an erection irritability loss in sexual ability, desire, drive, or performance loss of interest or pleasure nightmares pain in the chest below the breastbone passing gas redness or other discoloration of the skin runny or stuffy nose sensation of spinning sneezing tiredness trouble concentrating trouble sleeping Rare Bone pain continuing ringing or buzzing or other unexplained noise in the ears dry eyes hair loss or thinning of the hair hearing loss increased sensitivity of the skin to sunlight pain in the penis on erection severe sunburn Incidence not known Change in taste or bad, unusual, or unpleasant (after) taste fear or nervousness hives or welts Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Lopressor side effects

458. This particular disease, Angina, interacts with the drug Bayer Aspirin and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your

Drug Side Effects Summary

doctor immediately if any of the following side effects occur: Incidence not known Abdominal or stomach pain, cramping, or burning black, tarry stools bloody or cloudy urine change in consciousness chest pain or discomfort confusion constipation convulsions, severe or continuing dark urine decreased frequency or amount of urine diarrhea difficult breathing drowsiness fainting fast breathing feeling that something terrible will happen fever general tiredness and weakness greatly decreased frequency of urination or amount of urine headache heartburn increased thirst indigestion irregular heartbeat light-colored stools loss of appetite loss of consciousness lower back or side pain muscle cramping and weakness muscle tremors nausea or vomiting nervousness numbness or tingling in the hands, feet, or lips panic rapid, deep breathing restlessness seizures skin rash stomach cramps swelling of the face, fingers, or lower legs unusual bleeding or bruising unusual tiredness or weakness upper right abdominal or stomach vomiting of blood or material that looks like coffee grounds weakness or heaviness of the legs weight gain yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Acid or sour stomach anxiety belching dizziness dry mouth hyperventilation irritability shaking stomach discomfort, upset, or pain trouble sleeping unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Bayer Aspirin side effects

459. This particular disease, Angina, interacts with the drug Coreg and has the following side effects

- hives

Drug Side Effects Summary

- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- slow or uneven heartbeats
- cold feeling or numbness in your fingers or toes
- chest pain, dry cough, wheezing, chest tightness
- heart problems - swelling, rapid weight gain, feeling short of breath
- or high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor. Common

Coreg side effects may include: dizziness

- slow heartbeats
- diarrhea
- weight gain
- dry eyes
- or problems wearing contact lenses.

460. This particular disease, Angina, interacts with the drug Toprol-XL and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: very slow heartbeat, excessive tiredness, or chest pain
- a light-headed feeling, like you might pass out
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- or cold feeling in your hands and feet. Common Toprol-XL side effects may include: dizziness , tired feeling
- depression , confusion, memory problems

Drug Side Effects Summary

- nightmares , trouble sleeping
- diarrhea
- or mild itching or rash.

461. This particular disease, Angina, interacts with the drug amlodipine / atorvastatin and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, atorvastatin can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark colored urine. Amlodipine and atorvastatin may cause serious side effects. Call your doctor at once if you have: muscle stiffness, tremors, abnormal muscle movements
- muscle weakness in your hips, shoulders, neck, and back
- trouble lifting your arms, trouble climbing or standing
- severe drowsiness, feeling like you might pass out
- worsening chest pain, or chest pain spreading to the arm or jaw, nausea , sweating, general ill feeling
- or liver problems--upper stomach pain, loss of appetite, tiredness, dark urine, jaundice (yellowing of the skin or eyes). Common side effects of amlodipine and atorvastatin may include: muscle or joint pain
- diarrhea
- nausea, upset stomach
- or swelling in your legs or ankles.

462. This particular disease, Angina, interacts with the drug Ecotrin and has the following side effects:

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Ecotrin may cause serious side effects. Stop using Ecotrin and call your doctor at once if you have: ringing in your ears, confusion, hallucinations, rapid breathing, seizure (convulsions)
- severe nausea , vomiting, or stomach pain
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- fever lasting longer than 3 days
- or swelling, or pain lasting longer than 10 days. Common side effects of Ecotrin may include: upset stomach , heartburn
- drowsiness
- or mild headache .

463. This particular disease, Angina, interacts with the drug propranolol and has the following side

- slow or uneven heartbeats
- a light-headed feeling, like you might pass out
- wheezing or trouble breathing
- sudden weakness, vision problems, or loss of coordination (especially in a child with hemangioma that affects the face or head)
- cold feeling in your hands and feet
- depression , confusion, hallucinations
- heart problems - swelling, rapid weight gain, feeling short of breath
- low blood sugar - headache , hunger, sweating, irritability, dizziness , fast heart rate , and feeling anxious or shaky
- or low blood sugar in a baby - pale skin, blue or purple skin, sweating, fussiness, crying, not

Drug Side Effects Summary

wanting to eat, feeling cold, drowsiness, weak or shallow breathing (breathing may stop for short periods), seizure (convulsions), or loss of consciousness. Common propranolol side effects may include: dizziness, tiredness

- nausea , vomiting, diarrhea , constipation , stomach cramps
- sleep problems (insomnia)
- or runny or stuffy nose, cough, sore throat, hoarse voice.

464. This particular disease, Angina, interacts with the drug Easprin and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Abdominal or stomach pain, cramping, or burning black, tarry stools bloody or cloudy urine change in consciousness chest pain or discomfort confusion constipation convulsions, severe or continuing dark urine decreased frequency or amount of urine diarrhea difficult breathing drowsiness fainting fast breathing feeling that something terrible will happen fever general tiredness and weakness greatly decreased frequency of urination or amount of urine headache heartburn increased thirst indigestion irregular heartbeat light-colored stools loss of appetite loss of consciousness lower back or side pain muscle cramping and weakness muscle tremors nausea or vomiting nervousness numbness or tingling in the hands, feet, or lips panic rapid, deep breathing restlessness seizures skin rash stomach cramps swelling of the face, fingers, or lower legs unusual bleeding or bruising unusual tiredness or weakness upper right abdominal or stomach vomiting of blood or material that looks like coffee grounds weakness or heaviness of the legs weight gain yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health

Drug Side Effects Summary

care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Acid or sour stomach anxiety belching dizziness dry mouth hyperventilation irritability shaking stomach discomfort, upset, or pain trouble sleeping unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Easprin side effects

465. This particular disease, Angina, interacts with the drug Nitrolingual Pumpspray and has the fol

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Bloating or swelling of the face, arms, hands, lower legs, or feet burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings difficult or labored breathing feeling faint, dizzy, or lightheadedness feeling of warmth or heat flushing or redness of the skin, especially on the face and neck headache rapid weight gain sweating tightness in the chest tingling of the hands or feet unusual weight gain or loss Rare Bluish-colored lips, fingernails, or palms dark urine fever pale skin rapid heart rate sore throat unusual bleeding or bruising unusual tiredness or weakness Incidence not known Arm, back, or jaw pain blurred vision chest pain or discomfort chest tightness or heaviness confusion cough cracks in the skin difficulty with swallowing dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position fainting fast, irregular, pounding, or racing heartbeat or pulse feeling of constant movement of self or surroundings feeling of warmth hives, itching, or rash increased sweating loss of heat from the body nausea or vomiting puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red, swollen skin redness of the face, neck, arms, and occasionally, upper chest scaly skin sensation of spinning weakness Symptoms of overdose

Drug Side Effects Summary

Bluish-colored lips, fingernails, or palms blurred or loss of vision bulging soft spot on the head of an infant change in consciousness change in the ability to see colors, especially blue or yellow cold, clammy skin dark urine difficulty breathing disturbed color perception dizziness or lightheadedness double vision fever flushed skin halos around lights headache, severe and throbbing increased sweating loss of appetite loss of consciousness night blindness overbright appearance of lights pale skin paralysis slow or irregular heartbeat sore throat tunnel vision unusual bleeding or bruising unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Abdominal or stomach pain body aches or pain congestion hoarseness lack or loss of strength runny nose sneezing stuffy nose tender, swollen glands in the neck voice changes Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Nitrolingual Pumpspray side effects

466. This particular disease, Angina, interacts with the drug verapamil and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: chest pain, fast or slow heart rate
- a light-headed feeling, like you might pass out
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- fever, upper stomach pain, not feeling well

Drug Side Effects Summary

- or lung problems-- anxiety , sweating, pale skin, wheezing, gasping for breath, cough with foamy mucus. Common side effects may include: nausea , constipation
- headache , dizziness
- or low blood pressure.

467. This particular disease, Angina, interacts with the drug Bayer Aspirin Regimen and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
 - hives
 - itching
 - red, swollen, blistered, or peeling skin with or without fever
 - wheezing
 - tightness in the chest or throat
 - trouble breathing, swallowing, or talking
 - unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of bleeding like throwing up or coughing up blood
 - vomit that looks like coffee grounds
 - blood in the urine
 - black, red, or tarry stools
 - bleeding from the gums
 - abnormal vaginal bleeding
 - bruises without a cause or that get bigger

Drug Side Effects Summary

- or bleeding you cannot stop. Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of high potassium levels like a heartbeat that does not feel normal
- feeling confused
- feeling weak, lightheaded, or dizzy
- feeling like passing out
- numbness or tingling
- or shortness of breath. Signs of too much acid in the blood (acidosis) like confusion
- fast breathing
- fast heartbeat
- a heartbeat that does not feel normal
- very bad stomach pain, upset stomach, or throwing up
- feeling very sleepy
- shortness of breath
- or feeling very tired or weak. Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight. Very bad dizziness or passing out. Very bad headache . Ringing in the ears, hearing loss , or any other changes in hearing. Feeling agitated. Seizures . Bayer Aspirin Regimen side effects

468. This particular disease, Angina, interacts with the drug Caduet and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, atorvastatin can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure. Call your doctor right

Drug Side Effects Summary

away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark colored urine. Call your doctor at once if you have: muscle stiffness, tremors, abnormal muscle movements

- muscle weakness in your hips, shoulders, neck, and back
- trouble lifting your arms, trouble climbing or standing
- severe drowsiness, feeling like you might pass out
- worsening chest pain, or chest pain spreading to the arm or jaw, nausea , sweating, general ill feeling
- or liver problems - upper stomach pain, loss of appetite, tiredness, dark urine, jaundice (yellowing of the skin or eyes). Common Caduet side effects may include: muscle or joint pain
- diarrhea
- nausea, upset stomach
- or swelling in your legs or ankles.

469. This particular disease, Angina, interacts with the drug Aspir-Low and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Aspir-Low may cause serious side effects. Stop using Aspir-Low and call your doctor at once if you have: ringing in your ears, confusion, hallucinations, rapid breathing, seizure (convulsions)
- severe nausea , vomiting, or stomach pain
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- fever lasting longer than 3 days
- or swelling, or pain lasting longer than 10 days. Common side effects of Aspir-Low may include: upset stomach , heartburn

Drug Side Effects Summary

- drowsiness
- or mild headache .

470. This particular disease, Angina, interacts with the drug Katerzia and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, when you first start taking Katerzia, your angina may get worse or you could have a heart attack. Seek emergency medical attention or call your doctor right away if you have symptoms such as: chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating. Katerzia may cause serious side effects. Call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- worsening chest pain
- swelling in your feet or ankles
- severe drowsiness
- or a light-headed feeling, like you might pass out. Common side effects of Katerzia may include: dizziness , drowsiness
- feeling tired
- stomach pain, nausea
- or flushing (warmth, redness, or tingly feeling).

471. This particular disease, Angina, interacts with the drug Lovenox and has the following side effects:

- hives
- itching or burning skin
- difficult breathing
- swelling of your face, lips, tongue, or throat. Also seek emergency medical attention if you have

Drug Side Effects Summary

symptoms of a spinal blood clot : back pain, numbness or muscle weakness in your lower body, or loss of bladder or bowel control. Call your doctor at once if you have: unusual bleeding, or any bleeding that will not stop

- easy bruising, purple or red spots under your skin
- nosebleeds , bleeding gums
- abnormal vaginal bleeding , blood in your urine or stools
- coughing up blood or vomit that looks like coffee grounds
- signs of bleeding in the brain - sudden weakness (especially on one side of the body), sudden severe headache , problems with speech or vision
- or low red blood cells (anemia) - pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Common Lovenox side effects may include: nausea, diarrhea
- anemia
- confusion
- or pain, bruising, redness, or irritation where the medicine was injected.

472. This particular disease, Angina, interacts with the drug Norliqva and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Swelling of the ankles or feet Less common Chest tightness difficult or labored breathing dizziness fast, irregular, pounding, or racing heartbeat or pulse feeling of warmth redness of the face, neck, arms, and occasionally, upper chest Rare Black, tarry stools bleeding gums blistering, peeling, or loosening of the skin blood in the urine or stools blurred vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain or discomfort chills cold and clammy skin cold sweats confusion cough dark yellow urine diarrhea dilated neck veins dizziness or lightheadedness when

Drug Side Effects Summary

getting up from a lying or sitting position extra heartbeats fainting fever itching of the skin joint or muscle pain large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs numbness and tingling of the face, fingers, or toes pain in the arms, legs, or lower back, especially pain in the calves or heels upon exertion painful or difficult urination pale, bluish-colored, or cold hands or feet pinpoint red or purple spots on the skin red, irritated eyes redness of the face, neck, arms, and occasionally, upper chest redness, soreness or itching skin shakiness in the legs, arms, hands, or feet slow or irregular heartbeat sore throat sores, ulcers, or white spots on the lips or in the mouth sores, welting, or blisters sudden sweating sweating swelling of the face, fingers, feet, or lower legs swollen glands trembling or shaking of the hands or feet unsteadiness or awkwardness unusual bleeding or bruising unusual tiredness or weakness weak or absent pulses in the legs weakness in the arms, hands, legs, or feet weight gain yellow eyes or skin Incidence not known Clay-colored stools difficulty with speaking drooling headache loss of appetite loss of balance control muscle trembling, jerking, or stiffness nausea rash shuffling walk stomach pain twisting movements of the body uncontrolled movements, especially of the face, neck, and back unpleasant breath odor vomiting of blood yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Acid or sour stomach belching feeling of warmth heartburn indigestion lack or loss of strength muscle cramps redness of the face, neck, arms, and occasionally, upper chest sleepiness or unusual drowsiness stomach discomfort, upset, or pain Rare Abnormal dreams anxiety back pain bad unusual or unpleasant (after) taste bleeding gums blistering, crusting, irritation, itching, or reddening of the skin bloating bloody nose burning feeling in the chest or stomach burning while urinating burning, dry, or itching eyes change in color of the treated skin change in sense of smell

Drug Side Effects Summary

change in taste changes in vision constipation continuing ringing or buzzing or other unexplained noise in the ears cracked, dry, or scaly skin decreased sexual performance or desire depression difficulty with moving difficulty with swallowing discharge, excessive tearing double vision dry mouth dryness of the skin excess air or gas in the stomach or intestines excessive muscle tone eye pain feeling of constant movement of self or surroundings feeling of unreality feeling unusually cold flushed, dry skin fruit-like breath odor full feeling general feeling of discomfort or illness hair loss or thinning of the hair headache, severe and throbbing hearing loss increased appetite increased hunger increased sweating increased thirst increased urge to urinate during the night increased urination irritability irritation in the mouth lack of feeling or emotion loose stools loss of memory muscle pains, stiffness, tension, tightness, or weakness nervousness pain pains in the stomach, side, or abdomen, possibly radiating to the back passing gas problems with memory redness and swelling of the gums redness, pain, or swelling of the eye, eyelid, or inner lining of the eyelid restlessness runny nose seeing double sensation of spinning sense of detachment from self or body shakiness and unsteady walk shivering sneezing stuffy nose sweating swollen joints tenderness in the stomach area thirst trouble in holding or releasing urine trouble sleeping twitching uncaring unexplained weight loss unsteadiness, trembling, or other problems with muscle control or coordination waking to urinate at night weight loss Incidence not known Swelling of the breasts or breast soreness in both females and males Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Norliqva side effects

473. This particular disease, Angina, interacts with the drug Arthritis Pain and has the following side

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Arthritis Pain may cause serious side effects. Stop using Arthritis Pain and call your doctor at once if you have: ringing in your ears, confusion, hallucinations, rapid breathing, seizure (convulsions)
- severe nausea , vomiting, or stomach pain
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- fever lasting longer than 3 days
- or swelling, or pain lasting longer than 10 days. Common side effects of Arthritis Pain may include: upset stomach , heartburn
- drowsiness
- or mild headache .

474. This particular disease, Angina, interacts with the drug Ascriptin and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Abdominal or stomach pain, cramping, or burning black, tarry stools bloody or cloudy urine change in consciousness chest pain or discomfort confusion constipation convulsions, severe or continuing dark urine decreased frequency or amount of urine diarrhea difficult breathing drowsiness fainting fast breathing feeling that something terrible will happen fever general tiredness and weakness greatly decreased frequency of urination or amount of urine headache heartburn increased thirst indigestion irregular heartbeat light-colored stools loss of appetite loss of consciousness lower back or side pain muscle cramping and weakness muscle tremors nausea or vomiting nervousness numbness or tingling in the hands, feet, or lips panic rapid, deep breathing restlessness seizures skin rash stomach cramps swelling of the face, fingers, or lower legs unusual bleeding or bruising unusual tiredness or weakness upper right abdominal or stomach vomiting of blood or material that

Drug Side Effects Summary

looks like coffee grounds weakness or heaviness of the legs weight gain yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Acid or sour stomach anxiety belching dizziness dry mouth hyperventilation irritability shaking stomach discomfort, upset, or pain trouble sleeping unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Ascriptin side effects

475. This particular disease, Angina, interacts with the drug Aspergum and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Abdominal or stomach pain, cramping, or burning black, tarry stools bloody or cloudy urine change in consciousness chest pain or discomfort confusion constipation convulsions, severe or continuing dark urine decreased frequency or amount of urine diarrhea difficult breathing drowsiness fainting fast breathing feeling that something terrible will happen fever general tiredness and weakness greatly decreased frequency of urination or amount of urine headache heartburn increased thirst indigestion irregular heartbeat light-colored stools loss of appetite loss of consciousness lower back or side pain muscle cramping and weakness muscle tremors nausea or vomiting nervousness numbness or tingling in the hands, feet, or lips panic rapid, deep breathing restlessness seizures skin rash stomach cramps swelling of the face, fingers, or lower legs unusual bleeding or bruising

Drug Side Effects Summary

unusual tiredness or weakness upper right abdominal or stomach vomiting of blood or material that looks like coffee grounds weakness or heaviness of the legs weight gain yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Acid or sour stomach anxiety belching dizziness dry mouth hyperventilation irritability shaking stomach discomfort, upset, or pain trouble sleeping unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Aspergum side effects

476. This particular disease, Angina, interacts with the drug Aspiritab and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness

Drug Side Effects Summary

- or swelling of the mouth, face, lips, tongue, or throat. Signs of bleeding like throwing up or coughing up blood
- vomit that looks like coffee grounds
- blood in the urine
- black, red, or tarry stools
- bleeding from the gums
- abnormal vaginal bleeding
- bruises without a cause or that get bigger
- or bleeding you cannot stop. Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of high potassium levels like a heartbeat that does not feel normal
- feeling confused
- feeling weak, lightheaded, or dizzy
- feeling like passing out
- numbness or tingling
- or shortness of breath. Signs of too much acid in the blood (acidosis) like confusion
- fast breathing
- fast heartbeat
- a heartbeat that does not feel normal
- very bad stomach pain, upset stomach, or throwing up
- feeling very sleepy
- shortness of breath
- or feeling very tired or weak. Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight. Very bad dizziness or passing out.

Drug Side Effects Summary

Very bad headache . Ringing in the ears, hearing loss , or any other changes in hearing. Feeling agitated. Seizures . Aspiritab side effects

477. This particular disease, Angina, interacts with the drug Aspiritab and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Abdominal or stomach pain, cramping, or burning black, tarry stools bloody or cloudy urine change in consciousness chest pain or discomfort confusion constipation convulsions, severe or continuing dark urine decreased frequency or amount of urine diarrhea difficult breathing drowsiness fainting fast breathing feeling that something terrible will happen fever general tiredness and weakness greatly decreased frequency of urination or amount of urine headache heartburn increased thirst indigestion irregular heartbeat light-colored stools loss of appetite loss of consciousness lower back or side pain muscle cramping and weakness muscle tremors nausea or vomiting nervousness numbness or tingling in the hands, feet, or lips panic rapid, deep breathing restlessness seizures skin rash stomach cramps swelling of the face, fingers, or lower legs unusual bleeding or bruising unusual tiredness or weakness upper right abdominal or stomach vomiting of blood or material that looks like coffee grounds weakness or heaviness of the legs weight gain yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Acid or sour stomach anxiety belching dizziness dry mouth hyperventilation irritability shaking stomach discomfort, upset, or pain trouble sleeping unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness Other side effects not

Drug Side Effects Summary

listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Aspirtab side effects

478. This particular disease, Angina, interacts with the drug Bayer Aspirin Extra Strength Plus and

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Bayer Plus may cause serious side effects. Stop using Bayer Plus and call your doctor at once if you have: ringing in your ears, confusion, hallucinations, rapid breathing, seizure (convulsions)
- severe nausea , vomiting, or stomach pain
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- fever lasting longer than 3 days
- or swelling, or pain lasting longer than 10 days. Common side effects of Bayer Plus may include: upset stomach , heartburn
- drowsiness
- or mild headache .

479. This particular disease, Angina, interacts with the drug Ecpirin and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Abdominal or stomach pain, cramping, or burning black, tarry stools bloody or cloudy urine change in consciousness chest pain or discomfort confusion constipation convulsions, severe or continuing dark urine decreased frequency or amount of urine diarrhea difficult breathing drowsiness fainting

Drug Side Effects Summary

fast breathing feeling that something terrible will happen fever general tiredness and weakness greatly decreased frequency of urination or amount of urine headache heartburn increased thirst indigestion irregular heartbeat light-colored stools loss of appetite loss of consciousness lower back or side pain muscle cramping and weakness muscle tremors nausea or vomiting nervousness numbness or tingling in the hands, feet, or lips panic rapid, deep breathing restlessness seizures skin rash stomach cramps swelling of the face, fingers, or lower legs unusual bleeding or bruising unusual tiredness or weakness upper right abdominal or stomach vomiting of blood or material that looks like coffee grounds weakness or heaviness of the legs weight gain yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Acid or sour stomach anxiety belching dizziness dry mouth hyperventilation irritability shaking stomach discomfort, upset, or pain trouble sleeping unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Ecpirin side effects

480. This particular disease, Angina, interacts with the drug Entercote and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Abdominal or stomach pain, cramping, or burning black, tarry stools bloody or cloudy urine change in consciousness chest pain or discomfort confusion constipation convulsions, severe or continuing

Drug Side Effects Summary

dark urine decreased frequency or amount of urine diarrhea difficult breathing drowsiness fainting fast breathing feeling that something terrible will happen fever general tiredness and weakness greatly decreased frequency of urination or amount of urine headache heartburn increased thirst indigestion irregular heartbeat light-colored stools loss of appetite loss of consciousness lower back or side pain muscle cramping and weakness muscle tremors nausea or vomiting nervousness numbness or tingling in the hands, feet, or lips panic rapid, deep breathing restlessness seizures skin rash stomach cramps swelling of the face, fingers, or lower legs unusual bleeding or bruising unusual tiredness or weakness upper right abdominal or stomach vomiting of blood or material that looks like coffee grounds weakness or heaviness of the legs weight gain yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Acid or sour stomach anxiety belching dizziness dry mouth hyperventilation irritability shaking stomach discomfort, upset, or pain trouble sleeping unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Enterocote side effects

481. This particular disease, Angina, interacts with the drug Genacote and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Abdominal or stomach pain, cramping, or burning black, tarry stools bloody or cloudy urine change in

Drug Side Effects Summary

consciousness chest pain or discomfort confusion constipation convulsions, severe or continuing dark urine decreased frequency or amount of urine diarrhea difficult breathing drowsiness fainting fast breathing feeling that something terrible will happen fever general tiredness and weakness greatly decreased frequency of urination or amount of urine headache heartburn increased thirst indigestion irregular heartbeat light-colored stools loss of appetite loss of consciousness lower back or side pain muscle cramping and weakness muscle tremors nausea or vomiting nervousness numbness or tingling in the hands, feet, or lips panic rapid, deep breathing restlessness seizures skin rash stomach cramps swelling of the face, fingers, or lower legs unusual bleeding or bruising unusual tiredness or weakness upper right abdominal or stomach vomiting of blood or material that looks like coffee grounds weakness or heaviness of the legs weight gain yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Acid or sour stomach anxiety belching dizziness dry mouth hyperventilation irritability shaking stomach discomfort, upset, or pain trouble sleeping unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Genacote side effects

482. This particular disease, Angina, interacts with the drug Halfprin and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Abdominal or

Drug Side Effects Summary

stomach pain, cramping, or burning black, tarry stools bloody or cloudy urine change in consciousness chest pain or discomfort confusion constipation convulsions, severe or continuing dark urine decreased frequency or amount of urine diarrhea difficult breathing drowsiness fainting fast breathing feeling that something terrible will happen fever general tiredness and weakness greatly decreased frequency of urination or amount of urine headache heartburn increased thirst indigestion irregular heartbeat light-colored stools loss of appetite loss of consciousness lower back or side pain muscle cramping and weakness muscle tremors nausea or vomiting nervousness numbness or tingling in the hands, feet, or lips panic rapid, deep breathing restlessness seizures skin rash stomach cramps swelling of the face, fingers, or lower legs unusual bleeding or bruising unusual tiredness or weakness upper right abdominal or stomach vomiting of blood or material that looks like coffee grounds weakness or heaviness of the legs weight gain yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Acid or sour stomach anxiety belching dizziness dry mouth hyperventilation irritability shaking stomach discomfort, upset, or pain trouble sleeping unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Halfprin side effects

483. This particular disease, Angina, interacts with the drug Nitro-Dur and has the following side effects

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Nitro-Dur may cause serious side effects. Call your doctor at once if you have: worsening chest pain
 - pounding heartbeats or fluttering in your chest
 - slow heart rate
 - a light-headed feeling, like you might pass out
 - or heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating. Serious side effects may be more likely in older adults. Nitro-Dur can cause severe headaches. These headaches may gradually become less severe as you continue to use this medicine. Do not stop using Nitro-Dur. Ask your doctor before using any headache pain medication.
- Common side effects of Nitro-Dur may include: headache
- or feeling light-headed.

484. This particular disease, Angina, interacts with the drug Norwich Aspirin and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Abdominal or stomach pain, cramping, or burning black, tarry stools bloody or cloudy urine change in consciousness chest pain or discomfort confusion constipation convulsions, severe or continuing dark urine decreased frequency or amount of urine diarrhea difficult breathing drowsiness fainting fast breathing feeling that something terrible will happen fever general tiredness and weakness greatly decreased frequency of urination or amount of urine headache heartburn increased thirst indigestion irregular heartbeat light-colored stools loss of appetite loss of consciousness lower back or side pain muscle cramping and weakness muscle tremors nausea or vomiting nervousness numbness or tingling in the hands, feet, or lips panic rapid, deep breathing restlessness seizures skin rash stomach cramps swelling of the face, fingers, or lower legs unusual bleeding or bruising

Drug Side Effects Summary

unusual tiredness or weakness upper right abdominal or stomach vomiting of blood or material that looks like coffee grounds weakness or heaviness of the legs weight gain yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Acid or sour stomach anxiety belching dizziness dry mouth hyperventilation irritability shaking stomach discomfort, upset, or pain trouble sleeping unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Norwich Aspirin side effects

485. This particular disease, Angina, interacts with the drug Tenormin and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: new or worsening chest pain
- slow or uneven heartbeats
- a light-headed feeling, like you might pass out
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- or a cold feeling in your hands and feet. Common Tenormin side effects may include include: cold hands or feet
- dizziness
- tiredness

Drug Side Effects Summary

- or depressed mood.

486. This particular disease, Angina, interacts with the drug Vazalore and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Vazalore may cause serious side effects. Stop using Vazalore and call your doctor at once if you have: ringing in your ears, confusion, hallucinations, rapid breathing, seizure (convulsions)
- severe nausea , vomiting, or stomach pain
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- fever lasting longer than 3 days
- or swelling, or pain lasting longer than 10 days. Common side effects of Vazalore may include: upset stomach , heartburn
- drowsiness
- or mild headache .

487. This particular disease, Angina, interacts with the drug Coreg CR and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Coreg CR may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- slow or uneven heartbeats
- cold feeling or numbness in your fingers or toes
- chest pain, dry cough, wheezing, chest tightness
- heart problems--swelling, rapid weight gain, feeling short of breath

Drug Side Effects Summary

- or high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor. Common side effects of Coreg CR may include: dizziness

- slow heartbeats
- diarrhea
- weight gain
- dry eyes
- or problems wearing contact lenses.

488. This particular disease, Angina, interacts with the drug Corgard and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Corgard may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- slow heartbeats
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- or bronchospasm (wheezing, chest tightness, trouble breathing). Common side effects of Corgard may include: numbness or cold feeling in your hands or feet
- dizziness
- feeling tired
- upset stomach , vomiting , diarrhea , constipation
- vision problems
- or mood changes, confusion, memory problems.

489. This particular disease, Angina, interacts with the drug heparin and has the following side effects:

- nausea , vomiting , sweating, hives , itching, trouble breathing, swelling of your face, lips, tongue,

Drug Side Effects Summary

or throat, or feeling like you might pass out. Heparin may cause you to bleed more easily, which can be severe or life-threatening. You may also have bleeding on the inside of your body. Seek emergency medical attention if you have: skin warmth or discoloration

- chest pain, irregular heartbeats
- shortness of breath, dizziness , anxiety , sweating
- any unusual bleeding or bruising
- severe pain or swelling in your stomach, lower back, or groin
- dark or blue-colored skin on your hands or feet
- nausea, vomiting, loss of appetite
- unusual tiredness
- any bleeding that will not stop
- or nosebleed, blood in your urine or stools, black or tarry stools, or coughing up blood or vomit that looks like coffee grounds. Heparin can cause you to have bleeding episodes while you are using it and for several weeks after you stop. Bleeding may be more likely in older adults, especially women over 60 years of age Stop using this medicine and call your doctor at once if you have: skin changes where the medicine was injected
- fever, chills, runny nose , or watery eyes
- easy bruising, unusual bleeding, purple or red spots under your skin
- or signs of a blood clot - sudden numbness or weakness, problems with vision or speech, swelling or redness in an arm or leg. Common heparin side effects may include: unusual bleeding or bruising
- uncontrolled bleeding
- allergic reactions
- or abnormal liver function tests .

490. This particular disease, Angina, interacts with the drug Inderal LA and has the following side e

Drug Side Effects Summary

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common in children Chest tightness cough producing mucus difficulty with breathing Incidence not known Black, tarry stools blistering, peeling, or loosening of the skin blood in the urine bloody nose bloody stools blurred or loss of vision body aches or pain burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain or discomfort confusion about identity, place, and time congestion constipation cough cracks in the skin crying decreased awareness or responsiveness decreased urine output depersonalization diarrhea difficulty with swallowing dilated neck veins disturbed color perception dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position double vision dryness or soreness of the throat dysphoria euphoria fast, pounding, slow, or irregular heartbeat fever and chills general feeling of discomfort, illness, or weakness hair loss halos around lights headaches heavier menstrual periods hoarseness irregular breathing lightheadedness, dizziness, or fainting loss of heat from the body mental depression mimicry of speech or movements muscle or joint pain mutism nausea negativism night blindness noisy breathing overbright appearance of lights paleness or cold feeling in the fingertips and toes paranoia peculiar postures or movements, mannerisms, or grimacing pinpoint red or purple spots on the skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue quick to react or overreact emotionally rapidly changing moods rectal bleeding red skin lesions, often with a purple center red, irritated eyes red, swollen skin reddening of the skin, especially around the ears runny nose scaly skin seeing, hearing, or feeling things that are not there severe sleepiness short-term memory loss skin irritation or rash, including rash that looks like psoriasis skin rash, hives, or itching stomach pain and tenderness sores, ulcers, or white spots in the mouth or on the lips sweating swelling of the eyes, face, fingers, feet, or lower legs swollen glands tender, swollen glands in the neck tingling or pain in fingers or toes when exposed to cold tunnel vision unusual bleeding or bruising unusual tiredness or

Drug Side Effects Summary

weakness voice changes vomiting weight gain Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common in children Anxiety dry mouth hyperventilation irritability restlessness shaking sleepiness or unusual drowsiness trouble sleeping unusual dreams Less common in children Decreased appetite Incidence not known Dry eyes heartburn loss of strength or energy muscle weakness pain or discomfort in the chest, upper stomach, or throat stomach cramps unusual drowsiness, dullness, or feeling of sluggishness vivid dreams Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Inderal LA side effects

491. This particular disease, Angina, interacts with the drug Kaspargo Sprinkle and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking

Drug Side Effects Summary

- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Low mood (depression). Very bad dizziness or passing out. Chest pain that is new or worse. An abnormal heartbeat that is new or worse. Slow heartbeat. Shortness of breath, a big weight gain, or swelling in the arms or legs. Kapsargo Sprinkle side effects

492. This particular disease, Angina, interacts with the drug Nitro-Bid and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Nitro-Bid may cause serious side effects. Call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- slow heart rate
- a light-headed feeling, like you might pass out
- or heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating. Nitro-Bid can cause severe headaches. These headaches may gradually become less severe as you continue to use this medicine. Do not stop taking Nitro-Bid. Ask your doctor before using any headache pain medication. Common side effects of Nitro-Bid may include: flushing (warmth, redness, or tingly feeling)
- headache
- or dizziness.

493. This particular disease, Angina, interacts with the drug Fragmin and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Also seek emergency medical attention if you have

Drug Side Effects Summary

symptoms of a spinal blood clot : back pain, numbness or muscle weakness in your lower body, or loss of bladder or bowel control. Stop using Fragmin and call your doctor at once if you have: unusual bleeding (nose, mouth, vagina, or rectum), bleeding from wounds or needle injections, any bleeding that will not stop

- easy bruising, purple or red pinpoint spots under your skin
- black or bloody stools, coughing up blood or vomit that looks like coffee grounds
- sudden weakness, severe headache, confusion, or problems with speech, vision, or balance
- or trouble breathing. Common Fragmin side effects may include: low platelet levels (bruising, bleeding)
- or pain, bruising, or swelling where the medicine was injected.

494. This particular disease, Angina, interacts with the drug nadolol and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Nadolol may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- slow heartbeats
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- or bronchospasm (wheezing, chest tightness, trouble breathing). Common side effects of nadolol may include: numbness or cold feeling in your hands or feet
- dizziness
- feeling tired
- upset stomach , vomiting , diarrhea , constipation
- vision problems
- or mood changes, confusion, memory problems.

Drug Side Effects Summary

495. This particular disease, Angina, interacts with the drug Nitro TD Patch-A and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Nitro TD Patch-A may cause serious side effects. Call your doctor at once if you have: worsening chest pain
- pounding heartbeats or fluttering in your chest
- slow heart rate
- a light-headed feeling, like you might pass out
- or heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating. Serious side effects may be more likely in older adults. Nitro TD Patch-A can cause severe headaches. These headaches may gradually become less severe as you continue to use this medicine. Do not stop using Nitro TD Patch-A. Ask your doctor before using any headache pain medication. Common side effects of Nitro TD Patch-A may include: headache
- or feeling light-headed.

496. This particular disease, Angina, interacts with the drug amyl nitrite and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: Rare Skin rash unusual tiredness or weakness Signs and symptoms of overdose Bluish-colored lips, fingernails, or palms of hands dizziness (extreme) or fainting feeling of extreme pressure in head shortness of breath unusual tiredness or weakness weak and fast heartbeat Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the

Drug Side Effects Summary

following side effects continue or are bothersome or if you have any questions about them: More common Dizziness or lightheadedness, especially when getting up from a lying or sitting position fast pulse flushing of face and neck headache (mild) nausea or vomiting restlessness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Amyl nitrite side effects

497. This particular disease, Angina, interacts with the drug Angiomax and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tell your caregivers right away if you have: a light-headed feeling, like you might pass out
- easy bruising or bleeding (nosebleeds , bleeding gums, heavy menstrual bleeding)
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- sudden numbness or weakness, problems with vision or speech
- urine that looks red, pink, or brown
- swelling or redness in an arm or leg
- or bleeding from wounds or needle injections, any bleeding that will not stop. Bleeding is the most common side effect of bivalirudin.

498. This particular disease, Angina, interacts with the drug bivalirudin and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tell your caregivers right away if you have: a light-headed feeling, like you might pass out

Drug Side Effects Summary

- easy bruising or bleeding (nosebleeds , bleeding gums, heavy menstrual bleeding)
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- sudden numbness or weakness, problems with vision or speech
- urine that looks red, pink, or brown
- swelling or redness in an arm or leg
- or bleeding from wounds or needle injections, any bleeding that will not stop. Bleeding is the most common side effect of bivalirudin.

499. This particular disease, Angina, interacts with the drug Calan SR and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Blue lips and fingernails blurred vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain confusion coughing that sometimes produces a pink frothy sputum difficult, fast, noisy breathing, sometimes with wheezing dizziness, faintness, or lightheadedness when getting up from a lying or sitting position suddenly increased sweating lightheadedness, dizziness, or fainting pale skin shortness of breath slow or irregular heartbeat sore throat sweating swelling in legs and ankles unusual tiredness or weakness Rare Chills cold sweats feeling of warmth redness of the face, neck, arms and occasionally, upper chest Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Difficulty having a bowel movement (stool) headache Less common Acid or sour stomach belching difficulty in moving heartburn indigestion joint pain muscle aching or cramping muscle pains or stiffness

Drug Side Effects Summary

nausea rash stomach discomfort, upset, or pain trouble sleeping unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness swollen joints Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Calan SR side effects

500. This particular disease, Angina, interacts with the drug dalteparin and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Also seek emergency medical attention if you have symptoms of a spinal blood clot : back pain, numbness or muscle weakness in your lower body, or loss of bladder or bowel control. Dalteparin may cause serious side effects. Stop using dalteparin and call your doctor at once if you have: unusual bleeding (nose, mouth, vagina, or rectum), bleeding from wounds or needle injections, any bleeding that will not stop
- easy bruising, purple or red pinpoint spots under your skin
- black or bloody stools, coughing up blood or vomit that looks like coffee grounds
- sudden weakness, severe headache , confusion, or problems with speech, vision, or balance
- or trouble breathing. Common side effects of dalteparin may include: low platelet levels (bruising, bleeding)
- or pain, bruising, or swelling where the medicine was injected.

501. This particular disease, Angina, interacts with the drug Dilatrate-SR and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bluish-colored lips, fingernails, or

Drug Side Effects Summary

palms dark urine difficulty with breathing dizziness or lightheadedness fever headache pale skin rapid heart rate sore throat unusual bleeding or bruising unusual tiredness or weakness Incidence not known Arm, back, or jaw pain blurred vision chest pain or discomfort chest tightness or heaviness confusion dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position fainting fast or irregular heartbeat sweating Symptoms of overdose Blurred or loss of vision bulging soft spot on the head of an infant change in consciousness change in the ability to see colors, especially blue or yellow cold, clammy skin convulsions disturbed color perception double vision fast, irregular, pounding, or racing heartbeat or pulse feeling of constant movement of self or surroundings halos around lights headache, severe and throbbing increased sweating loss of appetite loss of consciousness nausea night blindness overbright appearance of lights paralysis sensation of spinning slow heartbeat tunnel vision vomiting Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Dilatrate-SR side effects

502. This particular disease, Angina, interacts with the drug eptifibatide and has the following side

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Tell your caregivers right away if you have: a light-headed feeling, like you might pass out
- any bleeding that will not stop
- bleeding around your IV or catheter, or in any place where your skin has been punctured with a needle
- red or pink urine
- or signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like

Drug Side Effects Summary

coffee grounds. The risk of bleeding may be higher in older adults. Common side effects of eptifibatide may include: bleeding

- or feeling lightheaded.

503. This particular disease, Angina, interacts with the drug GoNitro and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Bloating or swelling of the face, arms, hands, lower legs, or feet burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings difficult or labored breathing feeling faint, dizzy, or lightheadedness feeling of warmth or heat flushing or redness of the skin, especially on the face and neck headache rapid weight gain sweating tightness in the chest tingling of the hands or feet unusual weight gain or loss Rare Bluish-colored lips, fingernails, or palms dark urine fever pale skin rapid heart rate sore throat unusual bleeding or bruising unusual tiredness or weakness Incidence not known Arm, back, or jaw pain blurred vision chest pain or discomfort chest tightness or heaviness confusion cough cracks in the skin difficulty with swallowing dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position fainting fast, irregular, pounding, or racing heartbeat or pulse feeling of constant movement of self or surroundings feeling of warmth hives, itching, or rash increased sweating loss of heat from the body nausea or vomiting puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red, swollen skin redness of the face, neck, arms, and occasionally, upper chest scaly skin sensation of spinning weakness Symptoms of overdose Bluish-colored lips, fingernails, or palms blurred or loss of vision bulging soft spot on the head of an infant change in consciousness change in the ability to see colors, especially blue or yellow cold, clammy skin dark urine difficulty breathing disturbed color perception dizziness or lightheadedness double vision fever flushed skin halos around lights headache, severe and throbbing increased

Drug Side Effects Summary

sweating loss of appetite loss of consciousness night blindness overbright appearance of lights pale skin paralysis slow or irregular heartbeat sore throat tunnel vision unusual bleeding or bruising unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Abdominal or stomach pain body aches or pain congestion hoarseness lack or loss of strength runny nose sneezing stuffy nose tender, swollen glands in the neck voice changes Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. GoNitro side effects

504. This particular disease, Angina, interacts with the drug Integrilin and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Tell your caregivers right away if you have: a light-headed feeling, like you might pass out
- any bleeding that will not stop
- bleeding around your IV or catheter, or in any place where your skin has been punctured with a needle
- red or pink urine
- or signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. The risk of bleeding may be higher in older adults. Common Integrilin side effects may include: bleeding

Drug Side Effects Summary

- or feeling lightheaded.

505. This particular disease, Angina, interacts with the drug Isochron and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bluish-colored lips, fingernails, or palms dark urine difficulty with breathing dizziness or lightheadedness fever headache pale skin rapid heart rate sore throat unusual bleeding or bruising unusual tiredness or weakness Incidence not known Arm, back, or jaw pain blurred vision chest pain or discomfort chest tightness or heaviness confusion dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position fainting fast or irregular heartbeat sweating Symptoms of overdose Blurred or loss of vision bulging soft spot on the head of an infant change in consciousness change in the ability to see colors, especially blue or yellow cold, clammy skin convulsions disturbed color perception double vision fast, irregular, pounding, or racing heartbeat or pulse feeling of constant movement of self or surroundings halos around lights headache, severe and throbbing increased sweating loss of appetite loss of consciousness nausea night blindness overbright appearance of lights paralysis sensation of spinning slow heartbeat tunnel vision vomiting Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Isochron side effects

506. This particular disease, Angina, interacts with the drug IsoDitrate and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bluish-colored lips, fingernails, or

Drug Side Effects Summary

palms dark urine difficulty with breathing dizziness or lightheadedness fever headache pale skin rapid heart rate sore throat unusual bleeding or bruising unusual tiredness or weakness Incidence not known Arm, back, or jaw pain blurred vision chest pain or discomfort chest tightness or heaviness confusion dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position fainting fast or irregular heartbeat sweating Symptoms of overdose Blurred or loss of vision bulging soft spot on the head of an infant change in consciousness change in the ability to see colors, especially blue or yellow cold, clammy skin convulsions disturbed color perception double vision fast, irregular, pounding, or racing heartbeat or pulse feeling of constant movement of self or surroundings halos around lights headache, severe and throbbing increased sweating loss of appetite loss of consciousness nausea night blindness overbright appearance of lights paralysis sensation of spinning slow heartbeat tunnel vision vomiting Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. IsoDitrates side effects

507. This particular disease, Angina, interacts with the drug Isordil Titradose and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bluish-colored lips, fingernails, or palms dark urine difficulty with breathing dizziness or lightheadedness fever headache pale skin rapid heart rate sore throat unusual bleeding or bruising unusual tiredness or weakness Incidence not known Arm, back, or jaw pain blurred vision chest pain or discomfort chest tightness or heaviness confusion dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position fainting fast or irregular heartbeat sweating Symptoms of overdose Blurred or loss of vision bulging soft spot on the head of an infant change in consciousness change in the ability to

Drug Side Effects Summary

see colors, especially blue or yellow cold, clammy skin convulsions disturbed color perception double vision fast, irregular, pounding, or racing heartbeat or pulse feeling of constant movement of self or surroundings halos around lights headache, severe and throbbing increased sweating loss of appetite loss of consciousness nausea night blindness overbright appearance of lights paralysis sensation of spinning slow heartbeat tunnel vision vomiting Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Isordil Titradose side effects

508. This particular disease, Angina, interacts with the drug Minitran and has the following side effects

- hives
 - difficult breathing
 - swelling of your face, lips, tongue, or throat. Minitran may cause serious side effects. Call your doctor at once if you have: worsening chest pain
 - pounding heartbeats or fluttering in your chest
 - slow heart rate
 - a light-headed feeling, like you might pass out
 - or heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating. Serious side effects may be more likely in older adults. Minitran can cause severe headaches. These headaches may gradually become less severe as you continue to use this medicine. Do not stop using Minitran. Ask your doctor before using any headache pain medication.
- Common side effects of Minitran may include: headache
- or feeling light-headed.

509. This particular disease, Angina, interacts with the drug Nitro-Time and has the following side effects

Drug Side Effects Summary

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Bloating or swelling of the face, arms, hands, lower legs, or feet burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings difficult or labored breathing feeling faint, dizzy, or lightheadedness feeling of warmth or heat flushing or redness of the skin, especially on the face and neck headache rapid weight gain sweating tightness in the chest tingling of the hands or feet unusual weight gain or loss Rare Bluish-colored lips, fingernails, or palms dark urine fever pale skin rapid heart rate sore throat unusual bleeding or bruising unusual tiredness or weakness Incidence not known Arm, back, or jaw pain blurred vision chest pain or discomfort chest tightness or heaviness confusion cough cracks in the skin difficulty with swallowing dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position fainting fast, irregular, pounding, or racing heartbeat or pulse feeling of constant movement of self or surroundings feeling of warmth hives, itching, or rash increased sweating loss of heat from the body nausea or vomiting puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red, swollen skin redness of the face, neck, arms, and occasionally, upper chest scaly skin sensation of spinning weakness Symptoms of overdose Bluish-colored lips, fingernails, or palms blurred or loss of vision bulging soft spot on the head of an infant change in consciousness change in the ability to see colors, especially blue or yellow cold, clammy skin dark urine difficulty breathing disturbed color perception dizziness or lightheadedness double vision fever flushed skin halos around lights headache, severe and throbbing increased sweating loss of appetite loss of consciousness night blindness overbright appearance of lights pale skin paralysis slow or irregular heartbeat sore throat tunnel vision unusual bleeding or bruising unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of

Drug Side Effects Summary

these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Abdominal or stomach pain body aches or pain congestion hoarseness lack or loss of strength runny nose sneezing stuffy nose tender, swollen glands in the neck voice changes Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Nitro-Time side effects

510. This particular disease, Angina, interacts with the drug NitroMist and has the following side effects

- hives , sweating, pale skin
- nausea , vomiting
- feeling weak or light-headed
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe or throbbing headaches that do not become less severe with continued use of Nitromist
- pounding heartbeats or fluttering in your chest
- slow heart rate
- a light-headed feeling, like you might pass out
- blurred vision or dry mouth
- or heart attack symptoms - chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating. Nitromist can cause severe headaches that should become less severe as you continue to use the medicine. Common Nitromist side effects may include: flushing (sudden warmth, redness, or tingly feeling)
- feeling light-headed, fainting
- headache, dizziness

Drug Side Effects Summary

- or numbness, tingling, burning pain.

511. This particular disease, Angina, interacts with the drug timolol and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Timolol may cause serious side effects. Call your doctor at once if you have: new or worsening chest pain
- slow or uneven heartbeats
- a light-headed feeling, like you might pass out
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- muscle weakness
- drooping eyelids, vision problems
- or a cold feeling in your hands and feet. Common side effects of timolol may include: slow heartbeats
- dizziness
- or tiredness.

512. This particular disease, Angina, interacts with the drug Verelan and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Blue lips and fingernails blurred vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain confusion coughing that sometimes produces a pink frothy sputum difficult, fast, noisy breathing, sometimes with wheezing dizziness, faintness, or lightheadedness when getting up from a lying or sitting position suddenly increased sweating lightheadedness, dizziness, or fainting pale

Drug Side Effects Summary

skin shortness of breath slow or irregular heartbeat sore throat sweating swelling in legs and ankles unusual tiredness or weakness Rare Chills cold sweats feeling of warmth redness of the face, neck, arms and occasionally, upper chest Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Difficulty having a bowel movement (stool) headache Less common Acid or sour stomach belching difficulty in moving heartburn indigestion joint pain muscle aching or cramping muscle pains or stiffness nausea rash stomach discomfort, upset, or pain trouble sleeping unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness swollen joints Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Verelan side effects

513. This particular disease, Angina, interacts with the drug Verelan PM and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Blue lips and fingernails blurred vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain confusion coughing that sometimes produces a pink frothy sputum difficult, fast, noisy breathing, sometimes with wheezing dizziness, faintness, or lightheadedness when getting up from a lying or sitting position suddenly increased sweating lightheadedness, dizziness, or fainting pale skin shortness of breath slow or irregular heartbeat sore throat sweating swelling in legs and ankles unusual tiredness or weakness Rare Chills cold sweats feeling of warmth redness of the face, neck,

Drug Side Effects Summary

arms and occasionally, upper chest Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Difficulty having a bowel movement (stool) headache Less common Acid or sour stomach belching difficulty in moving heartburn indigestion joint pain muscle aching or cramping muscle pains or stiffness nausea rash stomach discomfort, upset, or pain trouble sleeping unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness swollen joints Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Verelan PM side effects

514. This particular disease, Angina, interacts with the drug Aspruzyo Sprinkle and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Dizziness Less common Bloating or swelling of the face, arms, hands, lower legs, or feet chest tightness difficult or labored breathing fast, irregular, pounding, or racing heartbeat or pulse feeling of constant movement of self or surroundings lightheadedness rapid weight gain sensation of spinning tingling of the hands or feet unusual weight gain or loss Rare Abnormal or decreased touch sensation agitation blood in the urine blurred vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain or discomfort chills cold sweats coma confusion decreased urine output depression dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position fainting headache hostility irritability lethargy muscle twitching nausea seizures shakiness in

Drug Side Effects Summary

the legs, arms, hands, or feet slow or irregular heartbeat stupor sweating trembling or shaking of the hands or feet unusual tiredness or weakness Incidence not known Decrease in frequency of urination decrease in urine volume difficulty in passing urine [dribbling] large, hive-like swelling on face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs painful urination seeing, hearing, or feeling things that are not there Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Difficulty having a bowel movement (stool) Less common Dry mouth stomach pain Rare Acid or sour stomach belching continuing ringing or buzzing or other unexplained noise in the ears hearing loss heartburn indigestion lack or loss of strength stomach discomfort or upset vomiting Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Aspruzyo Sprinkle side effects

515. This particular disease, Anxiety, interacts with the drug Xanax and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Alprazolam can slow or stop your breathing, especially if you have recently used an opioid medication or alcohol. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Common side effects of Xanax may include: drowsiness
- or feeling light-headed. Xanax may cause serious side effects. Call your doctor at once if you have: weak or shallow breathing

Drug Side Effects Summary

- a light-headed feeling, like you might pass out
 - a seizure
 - hallucinations, risk-taking behavior
 - increased energy, decreased need for sleep
 - racing thoughts, being agitated or talkative
 - double vision
 - or jaundice (yellowing of the skin or eyes). Drowsiness or dizziness may last longer in older adults.
- Use caution to avoid falling or accidental injury. After you stop using Xanax, get medical help right away if you have symptoms such as: unusual muscle movements, being more active or talkative, sudden and severe changes in mood or behavior, confusion, hallucinations, seizures, suicidal thoughts or actions. Some withdrawal symptoms may last up to 12 months or longer after stopping this medicine suddenly. Tell your doctor if you have ongoing anxiety, depression, problems with memory or thinking, trouble sleeping, ringing in your ears, a burning or prickly feeling, or a crawling sensation under your skin.

516. This particular disease, Anxiety, interacts with the drug clonazepam and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Clonazepam can slow or stop your breathing, especially if you have recently used an opioid medication or alcohol. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Tell your doctor right away if you have new or sudden changes in mood or behavior, including new or worse depression or anxiety, panic attacks, trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, more active or talkative, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: new or worsening

Drug Side Effects Summary

seizures

- severe drowsiness
- weak or shallow breathing
- unusual changes in mood or behavior
- confusion, paranoia, nightmares , hallucinations
- thoughts of suicide or hurting yourself
- unusual or involuntary eye movements. Drowsiness or dizziness may last longer in older adults.

Use caution to avoid falling or accidental injury. Common clonazepam side effects may include:

drowsiness, dizziness

- feeling tired or depressed
- memory problems
- or problems with walking or coordination. After you stop using clonazepam, get medical help right away if you have symptoms such as: unusual muscle movements, being more active or talkative, sudden and severe changes in mood or behavior, confusion, hallucinations, seizures, suicidal thoughts or actions. Some withdrawal symptoms may last up to 12 months or longer after stopping this medicine suddenly. Tell your doctor if you have ongoing anxiety, depression, problems with memory or thinking, trouble sleeping, ringing in your ears, a burning or prickly feeling, or a crawling sensation under your skin.

517. This particular disease, Anxiety, interacts with the drug alprazolam and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Alprazolam can slow or stop your breathing, especially if you have recently used an opioid medication or alcohol. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you

Drug Side Effects Summary

are hard to wake up. Call your doctor at once if you have: weak or shallow breathing

- a light-headed feeling, like you might pass out
- a seizure
- hallucinations, risk-taking behavior
- increased energy, decreased need for sleep
- racing thoughts, being agitated or talkative
- double vision
- or jaundice (yellowing of the skin or eyes). Drowsiness or dizziness may last longer in older adults.

Use caution to avoid falling or accidental injury. Common alprazolam side effects may include: drowsiness

- or feeling light-headed. After you stop using alprazolam, get medical help right away if you have symptoms such as: unusual muscle movements, being more active or talkative, sudden and severe changes in mood or behavior, confusion, hallucinations, seizures, suicidal thoughts or actions. Some withdrawal symptoms may last up to 12 months or longer after stopping this medicine suddenly. Tell your doctor if you have ongoing anxiety, depression, problems with memory or thinking, trouble sleeping, ringing in your ears, a burning or prickly feeling, or a crawling sensation under your skin.

518. This particular disease, Anxiety, interacts with the drug lorazepam and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Lorazepam can slow or stop your breathing, especially if you have recently used an opioid medication or alcohol. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Call your doctor at once if you have: severe drowsiness
- unusual changes in mood or behavior, being agitated or talkative

Drug Side Effects Summary

- sudden restless feeling or excitement
- thoughts of suicide or hurting yourself
- confusion, aggression, hallucinations
- sleep problems
- vision changes
- or dark urine, or jaundice (yellowing of the skin or eyes). Drowsiness or dizziness may last longer in older adults. Use caution to avoid falling or accidental injury. Common lorazepam side effects may include: dizziness, drowsiness
- weakness
- or feeling unsteady. After you stop using lorazepam, get medical help right away if you have symptoms such as: unusual muscle movements, being more active or talkative, sudden and severe changes in mood or behavior, confusion, hallucinations, seizures, suicidal thoughts or actions. Some withdrawal symptoms may last up to 12 months or longer after stopping this medicine suddenly. Tell your doctor if you have ongoing anxiety, depression, problems with memory or thinking, trouble sleeping, ringing in your ears, a burning or prickly feeling, or a crawling sensation under your skin.

519. This particular disease, Anxiety, interacts with the drug buspirone and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: chest pain
- shortness of breath
- or a light-headed feeling, like you might pass out. Common buspirone side effects may include: headache
- dizziness, drowsiness
- sleep problems (insomnia)

Drug Side Effects Summary

- nausea, upset stomach
- or feeling nervous or excited.

520. This particular disease, Anxiety, interacts with the drug diazepam and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Diazepam can slow or stop your breathing, especially if you have recently used an opioid medication or alcohol. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Tell your doctor right away if you have new or sudden changes in mood or behavior, including new or worse depression or anxiety, panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, more active or talkative, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: severe drowsiness or dizziness
- trouble breathing
- confusion, paranoia
- or new or worsening seizures. Drowsiness or dizziness may last longer in older adults. Use caution to avoid falling or accidental injury. Common diazepam side effects may include: drowsiness
- feeling tired
- muscle weakness
- or problems with balance or muscle movement. After you stop using diazepam, get medical help right away if you have symptoms such as: unusual muscle movements, being more active or talkative, sudden and severe changes in mood or behavior, confusion, hallucinations, seizures, suicidal thoughts or actions. Some withdrawal symptoms may last up to 12 months or longer after stopping this medicine suddenly. Tell your doctor if you have ongoing anxiety, depression, problems

Drug Side Effects Summary

with memory or thinking, trouble sleeping, ringing in your ears, a burning or prickly feeling, or a crawling sensation under your skin.

521. This particular disease, Anxiety, interacts with the drug Ativan and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Lorazepam can slow or stop your breathing, especially if you have recently used an opioid medication or alcohol. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Call your doctor at once if you have: severe drowsiness
- unusual changes in mood or behavior, being agitated or talkative
- sudden restless feeling or excitement
- thoughts of suicide or hurting yourself
- confusion, aggression, hallucinations
- sleep problems
- vision changes
- or dark urine, or jaundice (yellowing of the skin or eyes). Drowsiness or dizziness may last longer in older adults. Use caution to avoid falling or accidental injury. Common Ativan side effects may include: dizziness, drowsiness
- weakness
- or feeling unsteady. After you stop using Ativan, get medical help right away if you have symptoms such as: unusual muscle movements, being more active or talkative, sudden and severe changes in mood or behavior, confusion, hallucinations, seizures, suicidal thoughts or actions. Some withdrawal symptoms may last up to 12 months or longer after stopping this medicine suddenly. Tell your doctor if you have ongoing anxiety, depression, problems with memory or thinking, trouble sleeping, ringing

Drug Side Effects Summary

in your ears, a burning or prickly feeling, or a crawling sensation under your skin.

522. This particular disease, Anxiety, interacts with the drug Lexapro and has the following side effects

- skin rash or hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Lexapro may cause serious side effects. Call your doctor at once if you have: blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights
- racing thoughts, unusual risk-taking behavior, feelings of extreme happiness or sadness
- pain or burning when you urinate
- (in a child taking Lexapro) slow growth or weight gain
- low levels of sodium in the body - headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady
- or severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Common side effects of Lexapro may include: painful urination
- dizziness , drowsiness, tiredness, weakness
- feeling anxious or agitated
- increased muscle movements, feeling shaky
- sleep problems (insomnia)

Drug Side Effects Summary

- sweating, dry mouth, increased thirst, loss of appetite
- nausea, constipation
- yawning
- nosebleed, heavy menstrual periods
- or decreased sex drive, impotence , or difficulty having an orgasm.

523. This particular disease, Anxiety, interacts with the drug BuSpar and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: chest pain
- shortness of breath
- or a light-headed feeling, like you might pass out. Common BuSpar side effects may include:
headache
- dizziness, drowsiness
- sleep problems (insomnia)
- nausea, upset stomach
- or feeling nervous or excited.

524. This particular disease, Anxiety, interacts with the drug escitalopram and has the following side effects:

- skin rash or hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have:

Drug Side Effects Summary

blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights

- racing thoughts, unusual risk-taking behavior, feelings of extreme happiness or sadness
- pain or burning when you urinate
- (in a child taking escitalopram) slow growth or weight gain
- low levels of sodium in the body - headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady
- or severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Common escitalopram side effects may include: painful urination
- dizziness , drowsiness, tiredness, weakness
- feeling anxious or agitated
- increased muscle movements, feeling shaky
- sleep problems (insomnia)
- sweating, dry mouth, increased thirst, loss of appetite
- nausea, constipation
- yawning
- nosebleed, heavy menstrual periods
- or decreased sex drive, impotence , or difficulty having an orgasm.

525. This particular disease, Anxiety, interacts with the drug Valium and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Diazepam can slow or stop your breathing, especially

Drug Side Effects Summary

if you have recently used an opioid medication or alcohol. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Tell your doctor right away if you have new or sudden changes in mood or behavior, including new or worse depression or anxiety, panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, more active or talkative, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: severe drowsiness or dizziness

- trouble breathing

- confusion, paranoia

- or new or worsening seizures. Drowsiness or dizziness may last longer in older adults. Use caution to avoid falling or accidental injury. Common Valium side effects may include: drowsiness

- feeling tired

- muscle weakness

- or problems with balance or muscle movement. After you stop using Valium, get medical help right away if you have symptoms such as: unusual muscle movements, being more active or talkative, sudden and severe changes in mood or behavior, confusion, hallucinations, seizures, suicidal thoughts or actions. Some withdrawal symptoms may last up to 12 months or longer after stopping this medicine suddenly. Tell your doctor if you have ongoing anxiety, depression, problems with memory or thinking, trouble sleeping, ringing in your ears, a burning or prickly feeling, or a crawling sensation under your skin.

526. This particular disease, Anxiety, interacts with the drug gabapentin and has the following side

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Seek medical treatment if you have a serious drug

Drug Side Effects Summary

reaction that can affect many parts of your body. Symptoms may include: skin rash , fever, swollen glands, muscle aches, severe weakness, unusual bruising, upper stomach pain, or yellowing of your skin or eyes. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), depressed, or have thoughts about suicide or hurting yourself. Gabapentin may cause serious side effects. Call your doctor at once if you have: weak or shallow breathing

- blue-colored skin, lips, fingers, and toes
- confusion, extreme drowsiness or weakness
- problems with balance or muscle movement
- unusual or involuntary eye movements
- or increased seizures. Gabapentin can cause life-threatening breathing problems. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Breathing problems may be more likely in older adults or in people with COPD. Some side effects are more likely in children taking gabapentin. Contact your doctor if the child taking this medicine has any of the following side effects: changes in behavior
- memory problems
- trouble concentrating
- or acting restless, hostile, or aggressive. Common side effects of gabapentin may include: fever, chills, sore throat, body aches, unusual tiredness
- jerky movements
- headache
- double vision
- swelling of your legs and feet
- tremors

Drug Side Effects Summary

- trouble speaking
- dizziness, drowsiness, tiredness
- problems with balance or eye movements
- or nausea, vomiting.

527. This particular disease, Anxiety, interacts with the drug Cymbalta and has the following side effects:

- pounding heartbeats or fluttering in your chest
- a light-headed feeling, like you might pass out
- easy bruising, unusual bleeding
- vision changes
- painful or difficult urination
- liver problems - right-sided upper stomach pain, itching, dark urine, jaundice (yellowing of the skin or eyes)
- low blood sodium - headache , confusion, problems with thinking or memory, weakness, feeling unsteady
- or manic episodes - racing thoughts, increased energy, decreased need for sleep, risk-taking behavior, being agitated or talkative. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea . Common Cymbalta side effects may include: drowsiness
- nausea, constipation , loss of appetite
- dry mouth
- or increased sweating.

528. This particular disease, Anxiety, interacts with the drug Effexor XR and has the following side effects:

Drug Side Effects Summary

- skin rash or hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: blurred vision, eye pain or redness, seeing halos around lights
- cough, chest tightness, trouble breathing
- a seizure (convulsions)
- unusual bleeding nosebleeds , bleeding gums, abnormal vaginal bleeding , any bleeding that will not stop
- low blood sodium - headache , confusion, problems with thinking or memory, weakness, feeling unsteady
- or severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting , or diarrhea Common Effexor XR side effects may include: headache, dizziness , drowsiness, tiredness
- feeling anxious, nervous, or jittery
- sleep problems, unusual dreams
- tremors
- fast heartbeats
- blurred vision
- nausea, vomiting, diarrhea, constipation
- changes in weight or appetite

Drug Side Effects Summary

- dry mouth, yawning
- increased sweating
- or sexual problems.

529. This particular disease, Anxiety, interacts with the drug trazodone and has the following side effects

- hives
 - difficulty breathing
 - swelling of your face, lips, tongue, or throat. Stop taking trazodone and call your doctor at once if you have a penis erection that is painful or lasts 6 hours or longer. This is a medical emergency and could lead to a serious condition that must be corrected with surgery. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
 - slow heartbeats
 - unusual thoughts or behavior
 - easy bruising, unusual bleeding
 - or low levels of sodium in the body - headache, confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea .
- Common trazodone side effects may include: drowsiness, dizziness, tiredness
- swelling
 - weight loss

Drug Side Effects Summary

- blurred vision
- diarrhea, constipation
- or stuffy nose.

530. This particular disease, Anxiety, interacts with the drug duloxetine and has the following side effects:

- pounding heartbeats or fluttering in your chest
- a light-headed feeling, like you might pass out
- easy bruising, unusual bleeding
- vision changes
- painful or difficult urination
- liver problems - right-sided upper stomach pain, itching, dark urine, jaundice (yellowing of the skin or eyes)
- low blood sodium - headache , confusion, problems with thinking or memory, weakness, feeling unsteady
- or manic episodes - racing thoughts, increased energy, decreased need for sleep, risk-taking behavior, being agitated or talkative. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea . Common duloxetine side effects may include: drowsiness
- nausea, constipation , loss of appetite
- dry mouth
- or increased sweating.

531. This particular disease, Anxiety, interacts with the drug Paxil and has the following side effects:

- mood or behavior changes, anxiety, panic attacks , trouble sleeping, or if you feel impulsive,

Drug Side Effects Summary

irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: racing thoughts, decreased need for sleep, unusual risk-taking behavior, feelings of extreme happiness or sadness, being more talkative than usual

- blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights
- unusual bone pain or tenderness, swelling or bruising
- changes in weight or appetite
- easy bruising, unusual bleeding (nose, mouth, vagina, or rectum), coughing up blood
- severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, fainting
- or low levels of sodium in the body - headache , confusion, slurred speech, severe weakness, loss of coordination, feeling unsteady. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Common Paxil side effects may include: vision changes
- weakness, drowsiness, dizziness, tiredness
- sweating, anxiety, shaking
- sleep problems (insomnia)
- loss of appetite, nausea, vomiting, diarrhea, constipation
- dry mouth, yawning
- infection
- headache
- or decreased sex drive, impotence , abnormal ejaculation, or difficulty having an orgasm.

532. This particular disease, Anxiety, interacts with the drug venlafaxine and has the following side

Drug Side Effects Summary

- skin rash or hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Venlafaxine may cause serious side effects. Call your doctor at once if you have: blurred vision, eye pain or redness, seeing halos around lights
- cough, chest tightness, trouble breathing
- a seizure (convulsions)
- unusual bleeding - nosebleeds , bleeding gums, abnormal vaginal bleeding , any bleeding that will not stop
- low blood sodium - headache , confusion, problems with thinking or memory, weakness, feeling unsteady
- or severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting , or diarrhea Common side effects of venlafaxine may include: headache, dizziness , drowsiness, tiredness
- feeling anxious, nervous, or jittery
- sleep problems, unusual dreams
- tremors
- fast heartbeats
- blurred vision

Drug Side Effects Summary

- nausea, vomiting, diarrhea, constipation
- changes in weight or appetite
- dry mouth, yawning
- increased sweating
- or sexual problems.

533. This particular disease, Anxiety, interacts with the drug mirtazapine and has the following side effects:

- mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: racing thoughts, decreased need for sleep, unusual risk-taking behavior, feelings of extreme happiness or sadness, being more talkative than usual
- blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights
- a light-headed feeling, like you might pass out
- severe rash, blisters, or swelling on the palms of your hands or the soles of your feet
- a seizure
- low white blood cell counts - fever, chills, sore throat, cough, sores in your mouth or nose, flu-like symptoms, trouble breathing
- or low sodium level - headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea . Common mirtazapine side effects include: drowsiness, dizziness
- increased appetite
- or weight gain.

Drug Side Effects Summary

534. This particular disease, Anxiety, interacts with the drug paroxetine and has the following side effects:

- mood or behavior changes, anxiety, panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: racing thoughts, decreased need for sleep, unusual risk-taking behavior, feelings of extreme happiness or sadness, being more talkative than usual
- blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights
- unusual bone pain or tenderness, swelling or bruising
- changes in weight or appetite
- easy bruising, unusual bleeding (nose, mouth, vagina, or rectum), coughing up blood
- severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, fainting
- or low levels of sodium in the body - headache , confusion, slurred speech, severe weakness, loss of coordination, feeling unsteady. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Common paroxetine side effects may include: vision changes
- weakness, drowsiness, dizziness, tiredness
- sweating, anxiety, shaking
- sleep problems (insomnia)
- loss of appetite, nausea, vomiting, diarrhea, constipation
- dry mouth, yawning
- infection
- headache
- or decreased sex drive, impotence , abnormal ejaculation, or difficulty having an orgasm.

Drug Side Effects Summary

535. This particular disease, Anxiety, interacts with the drug Alprazolam Intensol and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs or symptoms of depression , suicidal thoughts, emotional ups and downs, abnormal thinking, anxiety, or lack of interest in life. Change in balance. Shortness of breath. Very bad dizziness or passing out. Feeling confused. Memory problems or loss. Trouble speaking. Trouble passing urine. Period (menstrual) changes. Alprazolam Intensol side effects

536. This particular disease, Anxiety, interacts with the drug lamotrigine and has the following side effects:

- mood or behavior changes, depression, anxiety , or if you feel agitated, hostile, restless, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself. Lamotrigine may cause serious side effects. Call your doctor at once if you have: fast, slow, or pounding heartbeats or fluttering in your chest
- chest pain, shortness of breath
- fever, swollen glands, weakness, severe muscle pain

Drug Side Effects Summary

- any skin rash, especially with blistering or peeling
- painful sores in your mouth or around your eyes
- headache, neck stiffness, increased sensitivity to light, nausea , vomiting, confusion, drowsiness
- jaundice (yellowing of the skin or eyes)
- or pale skin, cold hands and feet, easy bruising, unusual bleeding. Common side effects of lamotrigine may include: headache, dizziness
- blurred vision, double vision
- tremor, loss of coordination
- dry mouth, nausea, vomiting, stomach pain, diarrhea
- fever, sore throat, runny nose
- drowsiness, tired feeling
- back pain
- or sleep problems (insomnia).

537. This particular disease, Anxiety, interacts with the drug tramadol and has the following side effects

- noisy breathing, sighing, shallow breathing, breathing that stops during sleep
- a slow heart rate or weak pulse
- a light-headed feeling, like you might pass out
- seizure (convulsions)
- or low cortisol levels - nausea , vomiting , loss of appetite, dizziness, worsening tiredness or weakness. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea . Serious breathing problems may be more likely in older adults and people who are debilitated or have wasting syndrome or chronic breathing disorders. Common side effects of tramadol may include: constipation

Drug Side Effects Summary

- nausea and vomiting
- stomach pain
- dizziness
- drowsiness
- tiredness
- headache
- itching.

538. This particular disease, Anxiety, interacts with the drug risperidone and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- breast swelling or tenderness (in men or women), nipple discharge , impotence , lack of interest in sex, missed menstrual periods
- severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out
- low white blood cells - sudden weakness or ill feeling, fever, chills, sore throat , mouth sores, red or swollen gums, trouble swallowing, skin sores, cold or flu symptoms, cough, trouble breathing
- low levels of platelets in your blood - easy bruising, unusual bleeding (nose, mouth, vagina, or rectum), purple or red pinpoint spots under your skin
- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- or penis erection that is painful or lasts 4 hours or longer. Common risperidone side effects may include: headache

Drug Side Effects Summary

- dizziness, drowsiness, feeling tired
- tremors, twitching or uncontrollable muscle movements
- agitation , anxiety , restless feeling
- depressed mood
- dry mouth, upset stomach , diarrhea , constipation
- weight gain
- or cold symptoms such as stuffy nose, sneezing, sore throat.

539. This particular disease, Anxiety, interacts with the drug Tranxene and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Bladder pain bloody or cloudy urine difficult, burning, or painful urination frequent urge to urinate lower back or side pain shakiness and unsteady walk slurred speech trembling or shaking of the hands or feet unsteadiness, trembling, or other problems with muscle control or coordination Symptoms of overdose Change in consciousness drowsiness loss of consciousness mood or mental changes sleepiness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Blurred vision diarrhea discouragement dizziness double vision dry mouth feeling sad or empty headache indigestion irritability loss of appetite loss of interest or pleasure mental confusion nausea or vomiting nervousness passing of gas seeing double skin rash stomach pain, fullness, or discomfort trouble with concentrating trouble with sleeping unusual tiredness or weakness Other side effects not listed

Drug Side Effects Summary

may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Tranxene side effects

540. This particular disease, Anxiety, interacts with the drug Lorazepam Intensol and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Lorazepam Intensol can slow or stop your breathing, especially if you have recently used an opioid medication or alcohol. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Lorazepam Intensol may cause serious side effects. Call your doctor at once if you have: severe drowsiness
- unusual changes in mood or behavior, being agitated or talkative
- sudden restless feeling or excitement
- thoughts of suicide or hurting yourself
- confusion, aggression, hallucinations
- sleep problems
- vision changes
- or dark urine, or jaundice (yellowing of the skin or eyes). Drowsiness or dizziness may last longer in older adults. Use caution to avoid falling or accidental injury. Common side effects of Lorazepam Intensol may include: dizziness, drowsiness
- weakness
- or feeling unsteady. After you stop using Lorazepam Intensol, get medical help right away if you have symptoms such as: unusual muscle movements, being more active or talkative, sudden and severe changes in mood or behavior, confusion, hallucinations, seizures, suicidal thoughts or

Drug Side Effects Summary

actions. Some withdrawal symptoms may last up to 12 months or longer after stopping this medicine suddenly. Tell your doctor if you have ongoing anxiety, depression, problems with memory or thinking, trouble sleeping, ringing in your ears, a burning or prickly feeling, or a crawling sensation under your skin.

541. This particular disease, Anxiety, interacts with the drug clorazepate and has the following side

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Clorazepate can slow or stop your breathing, especially if you have recently used an opioid medication or alcohol. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Tell your doctor right away if you have new or sudden changes in mood or behavior, including new or worse depression or anxiety, panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, more active or talkative, or have thoughts about suicide or hurting yourself. Clorazepate may cause serious side effects. Call your doctor at once if you have: severe drowsiness
- slow breathing (breathing may stop)
- confusion, paranoia, unusual thoughts or behavior
- a seizure
- or suicidal thoughts or actions. Drowsiness or dizziness may last longer in older adults. Use caution to avoid falling or accidental injury. Common side effects of clorazepate may include: dizziness, drowsiness
- blurred vision
- upset stomach
- or dry mouth.

Drug Side Effects Summary

542. This particular disease, Anxiety, interacts with the drug doxepin and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Doxepin may cause serious side effects. Call your doctor at once if you have: blurred vision, tunnel vision, eye pain or swelling, seeing halos around lights
- a light-headed feeling, like you might pass out
- tremors, restless muscle movements in your eyes, tongue, jaw, or neck
- severe constipation
- confusion, hallucinations, unusual thoughts, seizure (convulsions)
- or painful or difficult urination. Common side effects of doxepin may include: drowsiness, dizziness
- increased blood pressure
- vision changes
- rash, swelling
- nausea , vomiting , indigestion
- little or no urination
- dry mouth, constipation
- or decreased or increased sex drive.

543. This particular disease, Anxiety, interacts with the drug oxazepam and has the following side effects:

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Oxazepam can slow or stop your breathing, especially if you have recently used an opioid medication or alcohol. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Oxazepam may cause serious side effects. Call your doctor at once if you have: severe drowsiness

- a light-headed feeling, like you might pass out

- unusual changes in mood or behavior

- confusion, anger, paranoia, problems with memory or concentration

- tremors, slurred speech, problems with balance or muscle movement

- a seizure

- trouble breathing

- or upper stomach pain, dark urine, jaundice (yellowing of the skin or eyes). Drowsiness or dizziness may last longer in older adults. Use caution to avoid falling or accidental injury. Common side effects of oxazepam may include: drowsiness

- dizziness, spinning sensation

- or headache . After you stop using oxazepam, get medical help right away if you have symptoms such as: unusual muscle movements, being more active or talkative, sudden and severe changes in mood or behavior, confusion, hallucinations, seizures, suicidal thoughts or actions. Some withdrawal symptoms may last up to 12 months or longer after stopping this medicine suddenly. Tell your doctor if you have ongoing anxiety, depression, problems with memory or thinking, trouble sleeping, ringing in your ears, a burning or prickly feeling, or a crawling sensation under your skin.

544. This particular disease, Anxiety, interacts with the drug Paxil CR and has the following side effects

- mood or behavior changes, anxiety, panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed,

Drug Side Effects Summary

or have thoughts about suicide or hurting yourself. Paxil CR may cause serious side effects. Call your doctor at once if you have: racing thoughts, decreased need for sleep, unusual risk-taking behavior, feelings of extreme happiness or sadness, being more talkative than usual

- blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights
- unusual bone pain or tenderness, swelling or bruising
- changes in weight or appetite
- easy bruising, unusual bleeding (nose, mouth, vagina, or rectum), coughing up blood
- severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, fainting
- or low levels of sodium in the body-- headache , confusion, slurred speech, severe weakness, loss of coordination, feeling unsteady. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Common side effects of Paxil CR may include: vision changes
- weakness, drowsiness, dizziness, tiredness
- sweating, anxiety, shaking
- sleep problems (insomnia)
- loss of appetite, nausea, vomiting, diarrhea, constipation
- dry mouth, yawning
- infection
- headache
- or decreased sex drive, impotence , abnormal ejaculation, or difficulty having an orgasm.

545. This particular disease, Anxiety, interacts with the drug amitriptyline / perphenazine and has th

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. High doses or long-term use of this medicine can cause a serious movement disorder that may not be reversible. The longer you use amitriptyline and perphenazine, the more likely you are to develop this disorder, especially if you are a woman or an older adult. Amitriptyline and perphenazine may cause serious side effects. Call your doctor at once if you have: uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- a light-headed feeling, like you might pass out
- fast, slow, or uneven heart rate
- chest pain or pressure, pain spreading to your jaw or shoulder
- easy bruising, unusual bleeding, purple or red spots under your skin
- dark urine, jaundice (yellowing of the skin or eyes)
- a seizure (convulsions)
- low white blood cell counts--fever, chills, sore throat , mouth sores, skin sores, sore throat, cough, trouble breathing
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, fainting. Common side effects of amitriptyline and perphenazine may include: drowsiness
- high blood pressure
- problems with speech
- involuntary muscle movements
- feeling restless or unable to sit still

Drug Side Effects Summary

- dry mouth
- headache
- or breast changes.

546. This particular disease, Anxiety, interacts with the drug chlordiazepoxide and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Chlordiazepoxide can slow or stop your breathing, especially if you have recently used an opioid medication or alcohol. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Chlordiazepoxide may cause serious side effects. Call your doctor at once if you have: severe drowsiness, slurred speech
- slow heart rate , slow breathing (breathing may stop)
- problems with memory or concentration
- unusual changes in mood or behavior, thoughts of suicide or hurting yourself
- problems with balance or muscle movement
- confusion, excitement, paranoia, anger, aggression
- or stomach pain, loss of appetite. Drowsiness or dizziness may last longer in older adults. Use caution to avoid falling or accidental injury. Common side effects of chlordiazepoxide may include: drowsiness
- confusion
- or balance problems. After you stop using chlordiazepoxide, get medical help right away if you have symptoms such as: unusual muscle movements, being more active or talkative, sudden and severe changes in mood or behavior, confusion, hallucinations, seizures, suicidal thoughts or actions. Some withdrawal symptoms may last up to 12 months or longer after stopping this medicine

Drug Side Effects Summary

suddenly. Tell your doctor if you have ongoing anxiety, depression, problems with memory or thinking, trouble sleeping, ringing in your ears, a burning or prickly feeling, or a crawling sensation under your skin.

547. This particular disease, Anxiety, interacts with the drug Diazepam Intensol and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Shakiness and unsteady walk unsteadiness, trembling, or other problems with muscle control or coordination Incidence not known Agitation black, tarry stools blistering, flaking, or peeling of the skin blurred vision changes in patterns and rhythms of speech chills confusion cough dark urine decrease in the frequency of urination decrease in urine volume difficulty in passing urine (dribbling) discouragement dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position false beliefs that cannot be changed by facts fast heartbeat fast or irregular breathing feeling sad or empty feeling that others are watching you or controlling your behavior feeling that others can hear your thoughts feeling, seeing, or hearing things that are not there fever headache hyperexcitability increased muscle spasms or tone irritability itching or rash lack of memory of what takes place after a certain event loss of appetite loss of bladder control loss of interest or pleasure lower back or side pain nausea nightmares outbursts of anger painful or difficult urination pale skin restlessness seizures slurred speech sore throat stomach pain sweating trouble concentrating trouble sleeping trouble speaking ulcers, sores, or white spots in the mouth unpleasant breath odor unusual behavior unusual bleeding or bruising unusual feeling of excitement unusual tiredness or weakness vomiting of blood yellow eyes or skin Symptoms of overdose Change in consciousness confusion difficult or trouble breathing irregular, fast or slow, or shallow breathing lack of coordination loss of consciousness loss of strength or energy muscle pain or weakness pale or blue lips, fingernails, or

Drug Side Effects Summary

skin shakiness and unsteady walk sleepiness unsteadiness, trembling, or other problems with muscle control or coordination unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Constipation decreased interest in sexual intercourse diarrhea difficulty with swallowing double vision dry mouth feeling of constant movement of self or surroundings inability to have or keep an erection increase in sexual ability, desire, drive, or performance increased interest in sexual intercourse increased watering of the mouth indigestion loss of sexual ability, desire, drive, or performance passing of gas seeing double sensation of spinning Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Diazepam Intensol side effects

548. This particular disease, Anxiety, interacts with the drug Loreev XR and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Loreev XR can slow or stop your breathing, especially if you have recently used an opioid medication or alcohol. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Loreev XR may cause serious side effects. Call your doctor at once if you have: severe drowsiness
- unusual changes in mood or behavior, being agitated or talkative

Drug Side Effects Summary

- sudden restless feeling or excitement
- thoughts of suicide or hurting yourself
- confusion, aggression, hallucinations
- sleep problems
- vision changes
- or dark urine, or jaundice (yellowing of the skin or eyes). Drowsiness or dizziness may last longer in older adults. Use caution to avoid falling or accidental injury. Common side effects of Loreev XR may include: dizziness, drowsiness
- weakness
- or feeling unsteady. After you stop using Loreev XR, get medical help right away if you have symptoms such as: unusual muscle movements, being more active or talkative, sudden and severe changes in mood or behavior, confusion, hallucinations, seizures, suicidal thoughts or actions. Some withdrawal symptoms may last up to 12 months or longer after stopping this medicine suddenly. Tell your doctor if you have ongoing anxiety, depression, problems with memory or thinking, trouble sleeping, ringing in your ears, a burning or prickly feeling, or a crawling sensation under your skin.

549. This particular disease, Anxiety, interacts with the drug oxcarbazepine and has the following s

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash, fever, swollen glands, flu-like symptoms, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Oxcarbazepine can reduce the sodium in your body to dangerously low levels, which can cause a life-threatening electrolyte imbalance. Call your doctor right away if you have nausea, lack of energy, confusion, feeling tired or irritable, severe weakness, muscle pain, or increased

Drug Side Effects Summary

seizures. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, depression , anxiety , or if you feel agitated, hostile, restless, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself. Common side effects of oxcarbazepine may include: dizziness , drowsiness, tiredness

- balance or coordination problems
- nausea, vomiting
- tremors or shaking
- double vision
- or rash.

550. This particular disease, Anxiety, interacts with the drug Pexeva and has the following side effects

- mood or behavior changes, anxiety, panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Pexeva may cause serious side effects. Call your doctor at once if you have: racing thoughts, decreased need for sleep, unusual risk-taking behavior, feelings of extreme happiness or sadness, being more talkative than usual
- blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights
- unusual bone pain or tenderness, swelling or bruising
- changes in weight or appetite
- easy bruising, unusual bleeding (nose, mouth, vagina, or rectum), coughing up blood
- severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, fainting
- or low levels of sodium in the body-- headache , confusion, slurred speech, severe weakness, loss of coordination, feeling unsteady. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate ,

Drug Side Effects Summary

muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Common side effects of Pexeva may include: vision changes

- weakness, drowsiness, dizziness, tiredness
- sweating, anxiety, shaking
- sleep problems (insomnia)
- loss of appetite, nausea, vomiting, diarrhea, constipation
- dry mouth, yawning
- infection
- headache
- or decreased sex drive, impotence , abnormal ejaculation, or difficulty having an orgasm.

551. This particular disease, Anxiety, interacts with the drug phenytoin and has the following side effects

- skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: slow or uneven heartbeats, chest pain, fluttering in your chest, and dizziness (like you might pass out)
- any skin rash, no matter how mild
- fever, chills, sore throat, swollen glands
- red or swollen gums, mouth sores
- easy bruising, unusual bleeding, purple or red spots under your skin
- or liver problems - loss of appetite, upper stomach pain, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common phenytoin side effects may include: drowsiness, confusion
- slurred speech

Drug Side Effects Summary

- abnormal eye movement
- or problems with balance, coordination, or muscle movement.

552. This particular disease, Anxiety, interacts with the drug prochlorperazine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. High doses or long-term use of prochlorperazine can cause a serious movement disorder that may not be reversible. The longer you use prochlorperazine, the more likely you are to develop this disorder, especially if you are a woman or an older adult. Prochlorperazine may cause serious side effects. Call your doctor at once if you have: uncontrolled muscle movements in your arms and legs, or your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- trouble speaking or swallowing, stiffness or muscle spasms in your neck
- tremors, or any new or unusual muscle movements you cannot control
- extreme drowsiness or light-headed feeling (like you might pass out)
- little or no urination
- agitation, restlessness
- severe constipation , stomach pain, bloating
- jaundice (yellowing of the skin or eyes)
- low blood cell counts--fever, chills, sore throat , cough, trouble breathing, mouth sores, skin sores, pale skin, easy bruising or bleeding
- lupus -like symptoms--muscle or joint pain, flu symptoms, chest pain, and a rash or patchy skin color that worsens in sunlight
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats. Side effects such as dry mouth, constipation, confusion, and tremors may be

Drug Side Effects Summary

more likely in older adults. Common side effects of prochlorperazine may include: headache , dizziness, drowsiness

- dry mouth, stuffy nose
- nausea, constipation
- increased appetite, weight gain
- blurred vision
- agitation, feeling jittery, trouble sleeping
- skin redness, itching, or rash
- missed menstrual periods
- or impotence , abnormal ejaculation.

553. This particular disease, Anxiety, interacts with the drug amitriptyline / chlordiazepoxide and ha

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Amitriptyline and chlordiazepoxide can slow or stop your breathing, especially if you have recently used an opioid medication, alcohol, or other drugs that can slow your breathing. A person caring for you should seek emergency medical attention if you have weak or shallow breathing, if you are hard to wake up, or if you stop breathing. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Amitriptyline and chlordiazepoxide may cause serious side effects. Call your doctor at once if you have: severe constipation
- confusion, hallucinations, unusual thoughts or behavior
- chest pain or heavy feeling, pain spreading to the jaw or shoulder

Drug Side Effects Summary

- pounding heartbeats or fluttering in your chest
- a light-headed feeling, like you might pass out
- or little or no urination. The sedative effects of this medicine may last longer in older adults.

Accidental falls are common in elderly patients who take benzodiazepines . Use caution to avoid falling or accidental injury. Common side effects of amitriptyline and chlorthalidone may include: dizziness, drowsiness

- blurred vision
- dry mouth
- or constipation, bloating .

554. This particular disease, Anxiety, interacts with the drug Compro and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. High doses or long-term use of prochlorperazine can cause a serious movement disorder that may not be reversible. The longer you use Compro, the more likely you are to develop this disorder, especially if you are a diabetic or an older adult. Compro may cause serious side effects. Call your doctor at once if you have: uncontrolled muscle movements in your arms, legs, or face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- a light-headed feeling, like you might pass out
- agitation
- jaundice (yellowing of the skin or eyes)
- slow heart rate , weak pulse, fainting, slow breathing
- fever, chills, sore throat , mouth sores
- new or worsening symptoms of lupus --joint pain, and a skin rash on your cheeks or arms that

Drug Side Effects Summary

worsens in sunlight

- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Side effects such as dry mouth, constipation, confusion, and tremors may be more likely in older adults. Common side effects of Compro may include: dizziness, drowsiness

- feeling restless or jittery
- blurred vision
- constipation
- breast swelling or discharge, a missed menstrual period
- sleep problems (insomnia)
- or swelling in your hands or feet.

555. This particular disease, Anxiety, interacts with the drug meprobamate and has the following side effects:

- feelings of extreme happiness, feeling restless or excited
- thoughts about suicide or hurting yourself
- fast or pounding heartbeats
- a light-headed feeling, like you might pass out
- swelling in your hands, ankles, or feet
- a red, itchy rash (may be only in your groin area)
- unusual bruising
- or sudden fever, chills, sore throat, weakness. Common side effects of meprobamate may include: drowsiness, weakness
- problems with balance or muscle movement
- headache , dizziness
- bruising

Drug Side Effects Summary

- numbness or tingling
- blurred vision
- or nausea , vomiting , diarrhea .

556. This particular disease, Anxiety, interacts with the drug nefazodone and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Nefazodone may cause serious side effects. Call your doctor at once if you have: problems with vision
- a seizure
- a light-headed feeling, like you might pass out
- manic episodes--racing thoughts, increased energy, decreased need for sleep, risk-taking behavior, being agitated or talkative
- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of nefazodone may include: drowsiness, dizziness , weakness
- vision problems
- nausea , constipation
- confusion, agitation
- or dry mouth.

557. This particular disease, Anxiety, interacts with the drug Tranxene T-Tab and has the following side effects:

Drug Side Effects Summary

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Tranxene T-Tab can slow or stop your breathing, especially if you have recently used an opioid medication or alcohol. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Tell your doctor right away if you have new or sudden changes in mood or behavior, including new or worse depression or anxiety, panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, more active or talkative, or have thoughts about suicide or hurting yourself. Tranxene T-Tab may cause serious side effects. Call your doctor at once if you have: severe drowsiness
- slow breathing (breathing may stop)
- confusion, paranoia, unusual thoughts or behavior
- a seizure
- or suicidal thoughts or actions. Drowsiness or dizziness may last longer in older adults. Use caution to avoid falling or accidental injury. Common side effects of Tranxene T-Tab may include: dizziness, drowsiness
- blurred vision
- upset stomach
- or dry mouth.

558. This particular disease, Anxiety, interacts with the drug trifluoperazine and has the following s

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. High doses or long-term use of trifluoperazine can cause a serious movement disorder that may not be reversible. The longer you use trifluoperazine,

Drug Side Effects Summary

the more likely you are to develop this disorder, especially if you are a woman or an older adult. Trifluoperazine may cause serious side effects. Call your doctor at once if you have: uncontrolled muscle movements in your arms, legs, or face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)

- stiffness in your neck, tightness in your throat, trouble breathing or swallowing
- feeling restless, jittery, or agitated
- decreased night vision, tunnel vision, watery eyes , increased sensitivity to light
- seizure (black-out or convulsions)
- jaundice (yellowing of the skin or eyes)
- low blood cell counts--fever, chills, mouth sores, skin sores, sore throat , cough, pale skin, easy bruising or bleeding, trouble breathing
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Common side effects of trifluoperazine may include: dizziness, drowsiness, tiredness
- blurred vision
- dry mouth, loss of appetite
- sleep problems (insomnia)
- muscle weakness
- itching or rash
- missed menstrual periods
- or breast swelling or discharge.

559. This particular disease, Anxiety, interacts with the drug vilazodone and has the following side

- skin rash or hives
- difficulty breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Vilazodone may cause serious side effects. Call your doctor at once if you have: a seizure (convulsions)
- blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights
- easy bruising, unusual bleeding
- racing thoughts, unusual risk-taking behavior, decreased inhibitions, feelings of extreme happiness or sadness
- or low levels of sodium in the body-- headache , confusion, slurred speech, severe weakness, loss of coordination, feeling unsteady. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting , or diarrhea . Common side effects of vilazodone may include: nausea, vomiting
- diarrhea
- or sleep problems (insomnia).

560. This particular disease, Asthma, interacts with the drug Singulair and has the following side effects

- hives , blisters, severe itching
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tell your doctor right away if you have signs of blood vessel inflammation: flu-like symptoms, severe sinus pain, a skin rash, numbness or a "pins and needles" feeling in your arms or legs. Some people using Singulair have had new or worsening mental problems. Stop taking this medicine and call your doctor right away if you have unusual changes in mood or behavior, such as: agitation, aggression, feeling restless or irritable

Drug Side Effects Summary

- anxiety , depression, confusion, problems with memory or attention
- stuttering, tremors, uncontrolled muscle movements
- suicidal thoughts or actions
- hallucinations, sleep problems, vivid, dreams, sleep-walking
- or compulsive or repetitive behaviors. Common Singulair side effects may include: stomach pain, diarrhea
- fever or other flu symptoms
- ear pain or full feeling, trouble hearing
- headache
- or cold symptoms such as runny or stuffy nose, sinus pain, cough, sore throat .

561. This particular disease, Asthma, interacts with the drug montelukast and has the following side effects:

- hives , blisters, severe itching
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tell your doctor right away if you have signs of blood vessel inflammation: flu-like symptoms, severe sinus pain, a skin rash, numbness or a "pins and needles" feeling in your arms or legs. Some people using montelukast have had new or worsening mental problems. Stop taking montelukast and call your doctor right away if you have unusual changes in mood or behavior, such as: agitation, aggression, feeling restless or irritable
- anxiety , depression, confusion, problems with memory or attention
- stuttering, tremors, uncontrolled muscle movements
- suicidal thoughts or actions
- hallucinations, sleep problems, vivid, dreams, sleep-walking
- or compulsive or repetitive behaviors. Common side effects of montelukast may include: stomach pain, diarrhea

Drug Side Effects Summary

- fever or other flu symptoms
- ear pain or full feeling, trouble hearing
- headache
- or cold symptoms such as runny or stuffy nose, sinus pain, cough, sore throat .

562. This particular disease, Asthma, interacts with the drug prednisone and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: blurred vision, eye pain, or seeing halos around lights
- swelling, rapid weight gain, feeling short of breath
- severe depression, feelings of extreme happiness or sadness, changes in personality or behavior, seizure (convulsions)
- bloody or tarry stools, coughing up blood
- pancreatitis (severe pain in your upper stomach spreading to your back, nausea and vomiting, fast heart rate)
- low potassium (confusion, uneven heart rate, extreme thirst, increased urination, leg discomfort, muscle weakness or limp feeling)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety, confusion, chest pain, shortness of breath, uneven heartbeats, seizure). Other common prednisone side effects may include: sleep problems (insomnia), mood changes
- increased appetite, gradual weight gain
- acne, increased sweating, dry skin, thinning skin, bruising or discoloration
- slow wound healing
- headache, dizziness, spinning sensation

Drug Side Effects Summary

- nausea, stomach pain, bloating
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

563. This particular disease, Asthma, interacts with the drug Trelegy Ellipta and has the following s

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: tremors, nervousness, chest pain, fast or pounding heartbeats
- sores or white patches in your mouth and throat, pain when swallowing
- wheezing, choking, or other breathing problems after using this medicine
- painful or difficult urination
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- a lung infection - fever, chills, cough with mucus, feeling short of breath
- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- low potassium level - leg cramps, constipation, irregular heartbeats, fluttering in your chest, numbness or tingling, muscle weakness or limp feeling
- or signs of a hormonal disorder - worsening tiredness or muscle weakness, feeling light-headed, nausea, vomiting. Common Trelegy Ellipta side effects may include: cold or flu symptoms such as runny or stuffy nose, sore throat , cough, chest tightness
- a lung infection
- nausea, vomiting, constipation, diarrhea
- joint pain
- mouth sores, hoarse voice
- headache , back pain

Drug Side Effects Summary

- or mouth pain, changes in your sense of taste.

564. This particular disease, Asthma, interacts with the drug Fasenra and has the following side effects:

- hives , rash
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: new or worsening asthma symptoms. Common Fasenra side effects may include: sore throat
- or headache .

565. This particular disease, Asthma, interacts with the drug Nucala and has the following side effects:

- hives, rash
- wheezing, chest tightness, difficult breathing
- feeling like you might pass out
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: unusual pain or tiredness
- burning or tingling anywhere in your body
- or a red or blistering skin rash. Common Nucala side effects may include: headache
- feeling tired
- back pain
- or pain, swelling, redness, burning, or itching where the medicine was injected.

566. This particular disease, Asthma, interacts with the drug benralizumab and has the following side effects:

- Benralizumab may cause serious side effects, including: allergic (hypersensitivity) reactions, including anaphylaxis. Serious allergic reactions can happen after you get your benralizumab injection. Allergic reactions can sometimes happen hours or days after you get your injection. Tell

Drug Side Effects Summary

your healthcare provider or get emergency help right away if you have any of the following symptoms of an allergic reaction: swelling of your face, mouth and tongue breathing problems fainting, dizziness, feeling lightheaded (low blood pressure) rash hives The most common side effects of benralizumab include headache and sore throat. These are not all the possible side effects of benralizumab. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Benralizumab side effects

567. This particular disease, Asthma, interacts with the drug fluticasone / umeclidinium / vilanterol

- hives
 - difficult breathing
 - swelling of your face, lips, tongue, or throat. Fluticasone, umeclidinium, and vilanterol may cause serious side effects. Call your doctor at once if you have: tremors, nervousness , chest pain, fast or pounding heartbeats
 - sores or white patches in your mouth and throat, pain when swallowing
 - wheezing, choking, or other breathing problems after using fluticasone, umeclidinium, and vilanterol
 - painful or difficult urination
 - blurred vision, tunnel vision, eye pain or redness, or seeing halos around lights
 - a lung infection--fever, chills, cough with mucus, feeling short of breath
 - high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
 - low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, numbness or tingling, muscle weakness or limp feeling
 - or signs of a hormonal disorder--tiredness or weakness, feeling light-headed, nausea , vomiting .
- Common side effects of fluticasone, umeclidinium, and vilanterol may include: cold or flu symptoms such as runny or stuffy nose, sinus pain, sore throat , cough, chest tightness

Drug Side Effects Summary

- a lung infection
- nausea, vomiting, constipation, diarrhea
- urination problems
- joint pain
- mouth sores, hoarse voice
- headache , back pain
- or mouth pain, changes in your sense of taste.

568. This particular disease, Asthma, interacts with the drug mepolizumab and has the following side effects:

- hives , rash
- wheezing, chest tightness, difficult breathing
- feeling like you might pass out
- swelling of your face, lips, tongue, or throat. Mepolizumab may cause serious side effects. Call your doctor at once if you have: unusual pain or tiredness
- burning or tingling anywhere in your body
- or a red or blistering skin rash . Common side effects of mepolizumab may include: headache
- feeling tired
- back pain
- or pain, swelling, redness, burning, or itching where the medicine was injected.

569. This particular disease, Asthma, interacts with the drug tezepelumab and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some side effects may occur during or after the injection. Tell your caregiver if you feel dizzy, nauseated, light-headed, itchy, short of breath, or if

Drug Side Effects Summary

you have swelling in your face. Common side effects of tezepelumab may include: sore throat

- or joint or back pain .

570. This particular disease, Asthma, interacts with the drug Tezspire and has the following side effects

- Tezspire may cause serious side effects, including: severe allergic reactions. Call your healthcare provider or get emergency medical care if you get any of the following symptoms of allergic reaction: rash hives breathing problems red, itchy, swollen, or inflamed eyes The most common side effects include: sore throat (pharyngitis) joint pain (arthralgia) back pain These are not all of the possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Tezspire side effects

571. This particular disease, Asthma, interacts with the drug Dulera and has the following side effects

- hives

- difficulty breathing

- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: worsening asthma symptoms

- tremors, nervousness, chest pain, fast or pounding heartbeats

- fever, chills, cough with mucus, feeling short of breath

- wheezing, choking, or other breathing problems after using this medication

- blurred vision, tunnel vision, eye pain or redness, or seeing halos around lights

- signs of thrush (a fungal infection) - sores or white patches in your mouth or throat, trouble swallowing

- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor

- low potassium level - leg cramps, constipation , irregular heartbeats, fluttering in your chest, numbness or tingling, muscle weakness or limp feeling

Drug Side Effects Summary

- or signs of a hormonal disorder - tiredness or weakness, feeling light-headed, nausea , vomiting .
- Mometasone can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common Dulera side effects may include: runny or stuffy nose, sinus pain
- headache
 - or cough, sore throat .

572. This particular disease, Asthma, interacts with the drug Atrovent and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Bladder pain bloody or cloudy urine cough producing mucus difficult, burning, or painful urination difficulty with breathing frequent urge to urinate lower back or side pain shortness of breath tightness in the chest wheezing Less common Body aches or pain chills cough ear congestion fever headache loss of voice runny nose sneezing sore throat unusual tiredness or weakness Rare Constipation (continuing) or lower abdominal pain or bloating fainting fast, pounding, or irregular heartbeat or pulse severe eye pain skin rash or hives swelling of the face, lips, or eyelids Incidence not known Blindness blurred vision confusion decrease in the frequency of urination decrease in the urine volume decreased vision difficulty in passing urine (dribbling) difficulty with swallowing dizziness dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position hives or welts itching large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs nausea or vomiting noisy breathing puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue redness of the skin redness of the white part of the eyes or inside of the eyelids sweating tearing Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care

Drug Side Effects Summary

professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Back pain dryness of the mouth unpleasant taste Less common or rare Acid or sour stomach belching burning eyes diarrhea general feeling of discomfort or illness heartburn indigestion joint pain loss of appetite muscle aches and pains nervousness pain or tenderness around the eyes and cheekbones shivering stomach discomfort, upset, or pain trembling trouble sleeping Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Atrovent side effects

573. This particular disease, Asthma, interacts with the drug Breo Ellipta and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: wheezing, choking, or other breathing problems after using this medicine
- chest pain, fast or pounding heartbeats, pounding in your neck or ears
- sores or white patches in your mouth and throat, pain when swallowing
- fever, chills, cough with yellow or green mucus
- blurred vision, tunnel vision, eye pain , or seeing halos around lights. high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- low potassium level - leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- or signs of a hormonal disorder - worsening tiredness or muscle weakness, feeling light-headed, nausea , vomiting . Common Breo Ellipta side effects may include: headache , back pain , joint pain

Drug Side Effects Summary

- fever, flu symptoms
- increased blood pressure
- runny or stuffy nose, sore throat , cough
- or a hoarse voice.

574. This particular disease, Asthma, interacts with the drug ipratropium and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. You may need to use a different bronchodilator medication if you have an allergic reaction to ipratropium inhalation. Ipratropium inhalation may cause serious side effects. Call your doctor at once if you have: wheezing, choking, or other breathing problems after using this medicine
- little or no urination
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- or worsened breathing problems. Common side effects of ipratropium inhalation may include: cold symptoms such as stuffy nose, sneezing, sore throat
- cough with mucus, chest tightness, trouble breathing
- or tiredness, or headache .

575. This particular disease, Asthma, interacts with the drug methylprednisolone and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: shortness of breath (even with mild exertion), swelling, rapid weight gain
- bruising, thinning skin, or any wound that will not heal

Drug Side Effects Summary

- blurred vision, tunnel vision, eye pain, or seeing halos around lights
- severe depression, changes in personality, unusual thoughts or behavior
- new or unusual pain in an arm or leg or in your back
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- seizure (convulsions)
- or low potassium - leg cramps, constipation, irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling. Steroids can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common methylprednisolone side effects may include: fluid retention (swelling in your hands or ankles)
- dizziness, spinning sensation
- changes in your menstrual periods
- headache
- mild muscle pain or weakness
- or stomach discomfort, bloating.

576. This particular disease, Asthma, interacts with the drug Atrovent HFA and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. You may need to use a different bronchodilator medication if you have an allergic reaction to ipratropium. Call your doctor at once if you have: wheezing, choking, or other breathing problems after using this medicine
- little or no urination
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- or worsened breathing problems. Common Atrovent HFA side effects may include: dry mouth
- bitter taste

Drug Side Effects Summary

- headache , dizziness, or tiredness
- cold symptoms such as stuffy nose, sneezing, sore throat
- cough with mucus, chest tightness, trouble breathing
- nausea, upset stomach, constipation back pain
- or fever, chills, body aches, flu symptoms

577. This particular disease, Asthma, interacts with the drug dexamethasone and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: muscle tightness, weakness, or limp feeling
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- severe depression, unusual thoughts or behavior
- a seizure (convulsions)
- bloody or tarry stools, coughing up blood
- fast or slow heart rate , weak pulse
- pancreatitis - severe pain in your upper stomach spreading to your back, nausea and vomiting
- low potassium level - leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling
- or increased blood pressure - severe headache , blurred vision, pounding in your neck or ears, anxiety , nosebleed. Dexamethasone can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using dexamethasone. Common dexamethasone side effects may include: fluid retention (swelling in your hands or ankles)
- increased appetite

Drug Side Effects Summary

- mood changes, trouble sleeping
- skin rash , bruising or discoloration
- acne, increased sweating, increased hair growth
- headache, dizziness
- nausea , vomiting , upset stomach
- changes in your menstrual periods
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

578. This particular disease, Asthma, interacts with the drug Alvesco and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: wheezing, choking, or other breathing problems after using this medication
- bronchospasm (wheezing, chest tightness, trouble breathing)
- sores or white patches in your mouth or throat
- vision problems, such as tunnel vision or blurred vision
- or worsening asthma symptoms. Common Alvesco side effects may include: unpleasant taste in your mouth
- dry mouth, burning or irritation in your mouth
- runny or stuffy nose, sore throat
- hoarse voice
- mild itching or skin rash
- joint pain, back pain
- or headache.

Drug Side Effects Summary

579. This particular disease, Asthma, interacts with the drug levalbuterol and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Levalbuterol inhalation may cause serious side effects. Call your doctor at once if you have: wheezing, choking, or other breathing problems after using levalbuterol inhalation
- pounding heartbeats or fluttering in your chest
- worsening asthma symptoms
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, extreme thirst, increased urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of levalbuterol inhalation may include: dizziness , nervousness , tremors
- runny nose , sore throat
- chest pain or tightness, irregular heartbeats
- pain
- or vomiting .

580. This particular disease, Asthma, interacts with the drug Ipratropium Inhalation Solution and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing

Drug Side Effects Summary

- tightness in the chest or throat
 - trouble breathing, swallowing, or talking
 - unusual hoarseness
 - or swelling of the mouth, face, lips, tongue, or throat. Signs of a urinary tract infection (UTI) like blood in the urine, burning or pain when passing urine, feeling the need to pass urine often or right away, fever, lower stomach pain, or pelvic pain . Change in eyesight, eye pain, or very bad eye irritation. Seeing halos or bright colors around lights. Red eyes. Trouble passing urine, pain when passing urine, passing urine in a weak stream or drips, or passing urine more often. Very bad dizziness or passing out. Fast or abnormal heartbeat. Mouth irritation or mouth sores. This medicine can cause very bad breathing problems right after you take a dose. Sometimes, this may be life-threatening. If you have trouble breathing, breathing that is worse, wheezing, or coughing after using ipratropium inhalation solution, use a rescue inhaler and get medical help right away.
- Ipratropium Inhalation Solution side effects

581. This particular disease, Asthma, interacts with the drug Rayos and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: muscle pain or weakness
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- severe depression , changes in personality, unusual thoughts or behavior
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- swelling, rapid weight gain, feeling short of breath
- irregular heartbeats
- severe headache , pounding in your neck or ears

Drug Side Effects Summary

- decreased adrenal gland hormones - muscle weakness, tiredness, diarrhea, nausea , menstrual changes, skin discoloration, craving salty foods, and feeling light-headed
- or low potassium level - leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Prednisone can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common Rayos side effects may include: weight gain (especially in your face or your upper back and torso)
- increased appetite
- mood changes, trouble sleeping
- changes in your menstrual periods
- problems with memory or thought
- muscle or joint pain
- weakness
- headache, dizziness , spinning sensation
- nausea, bloating , loss of appetite
- slow wound healing
- or acne, increased sweating, thinning skin, bruising, pinpoint spots under your skin.

582. This particular disease, Asthma, interacts with the drug fluticasone / vilanterol and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Fluticasone and vilanterol may cause serious side effects. Call your doctor at once if you have: wheezing, choking, or other breathing problems after using fluticasone and vilanterol
- chest pain, fast or pounding heartbeats, pounding in your neck or ears

Drug Side Effects Summary

- sores or white patches in your mouth and throat, pain when swallowing
- fever, chills, cough with yellow or green mucus
- blurred vision, tunnel vision, eye pain , or seeing halos around lights. high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- or signs of a hormonal disorder--worsening tiredness or muscle weakness, feeling light-headed, nausea , vomiting . Common side effects of fluticasone and vilanterol may include: headache , back pain , joint pain
- fever, flu symptoms
- increased blood pressure
- runny or stuffy nose, sore throat , cough
- or a hoarse voice.

583. This particular disease, Asthma, interacts with the drug formoterol / mometasone and has the

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Formoterol and mometasone may cause serious side effects. Call your doctor at once if you have: worsening asthma symptoms
- tremors, nervousness , chest pain, fast or pounding heartbeats
- fever, chills, cough with mucus, feeling short of breath
- wheezing, choking, or other breathing problems after using this medication
- blurred vision, tunnel vision, eye pain or redness, or seeing halos around lights
- signs of thrush (a fungal infection)--sores or white patches in your mouth or throat, trouble swallowing

Drug Side Effects Summary

- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
 - low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, numbness or tingling, muscle weakness or limp feeling
 - or signs of a hormonal disorder--tiredness or weakness, feeling light-headed, nausea , vomiting .
- Mometasone can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using formoterol and mometasone. Common side effects of formoterol and mometasone may include: runny or stuffy nose, sinus pain
- headache
 - or cough, sore throat .

584. This particular disease, Asthma, interacts with the drug Medrol and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Medrol may cause serious side effects. Call your doctor at once if you have: shortness of breath (even with mild exertion), swelling, rapid weight gain
- bruising, thinning skin, or any wound that will not heal
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- severe depression, changes in personality, unusual thoughts or behavior
- new or unusual pain in an arm or leg or in your back
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- seizure (convulsions)
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling. Steroids can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using Medrol. Common side effects of Medrol may include: fluid retention (swelling in your hands or ankles)

Drug Side Effects Summary

- dizziness , spinning sensation
- changes in your menstrual periods
- headache
- mild muscle pain or weakness
- or stomach discomfort, bloating .

585. This particular disease, Asthma, interacts with the drug MethylPREDNISolone Dose Pack and I

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. MethylPREDNISolone Dose Pack may cause serious side effects. Call your doctor at once if you have: shortness of breath (even with mild exertion), swelling, rapid weight gain
- bruising, thinning skin, or any wound that will not heal
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- severe depression, changes in personality, unusual thoughts or behavior
- new or unusual pain in an arm or leg or in your back
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- seizure (convulsions)
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling. Steroids can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using MethylPREDNISolone Dose Pack Dose Pack. Common side effects of MethylPREDNISolone Dose Pack may include: fluid retention (swelling in your hands or ankles)
- dizziness , spinning sensation
- changes in your menstrual periods

Drug Side Effects Summary

- headache
- mild muscle pain or weakness
- or stomach discomfort, bloating .

586. This particular disease, Asthma, interacts with the drug triamcinolone and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have a serious side effect such as: problems with your vision
- swelling, rapid weight gain, feeling short of breath
- severe depression, unusual thoughts or behavior, seizure (convulsions)
- bloody or tarry stools, coughing up blood
- pancreatitis (severe pain in your upper stomach spreading to your back, nausea and vomiting, fast heart rate)
- low potassium (confusion, uneven heart rate, extreme thirst, increased urination, leg discomfort, muscle weakness or limp feeling)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety, confusion, chest pain, shortness of breath, uneven heartbeats, seizure). Less serious triamcinolone side effects may include: sleep problems (insomnia), mood changes
- acne, dry skin, thinning skin, bruising or discoloration
- slow wound healing
- increased sweating
- headache, dizziness, spinning sensation
- nausea, stomach pain, bloating
- muscle weakness

Drug Side Effects Summary

- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist)

-

587. This particular disease, Asthma, interacts with the drug Dexamethasone Intensol and has the f

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Aggression agitation anxiety blurred vision decrease in the amount of urine dizziness fast, slow, pounding, or irregular heartbeat or pulse headache irritability mental depression mood changes nervousness noisy, rattling breathing numbness or tingling in the arms or legs pounding in the ears swelling of the fingers, hands, feet, or lower legs trouble thinking, speaking, or walking troubled breathing at rest weight gain Incidence not known Abdominal or stomach cramping and/or burning (severe) or pain backache bloody, black, or tarry stools cough or hoarseness darkening of the skin decrease in height decreased vision diarrhea dry mouth eye pain eye tearing facial hair growth in females fainting fatigue fever or chills flushed, dry skin fractures fruit-like breath odor full or round face, neck, or trunk heartburn and/or indigestion (severe and continuous) increased hunger increased thirst increased urination loss of appetite loss of sexual desire or ability lower back or side pain menstrual irregularities muscle pain or tenderness muscle wasting or weakness nausea pain in the back, ribs, arms, or legs painful or difficult urination skin rash sleeplessness sweating trouble healing trouble sleeping unexplained weight loss unusual tiredness or weakness vision changes vomiting vomiting of material that looks like coffee grounds Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following

Drug Side Effects Summary

side effects continue or are bothersome or if you have any questions about them: More common Increased appetite Incidence not known Abnormal fat deposits on the face, neck, and trunk acne dry scalp lightening of normal skin color red face reddish purple lines on the arms, face, legs, trunk, or groin swelling of the stomach area thinning of the scalp hair Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Dexamethasone Intensol side effects

588. This particular disease, Asthma, interacts with the drug Kenalog-40 and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Kenalog-40 may cause serious side effects. Call your doctor at once if you have: (after injection into a joint space) increased pain or swelling, joint stiffness, fever, and general ill feeling
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- unusual changes in mood or behavior
- swelling, rapid weight gain, feeling short of breath
- stomach cramps, vomiting , diarrhea, bloody or tarry stools, rectal irritation
- sudden numbness or weakness (especially on one side of the body)
- a seizure (convulsions)
- severe headache , blurred vision, pounding in your neck or ears
- increased pressure inside the skull--severe headaches, ringing in your ears, dizziness , nausea , vision problems, pain behind your eyes
- or signs of low adrenal gland hormones --flu-like symptoms, headache, depression , weakness, tiredness, diarrhea, vomiting, stomach pain, craving salty foods, and feeling light-headed. Certain

Drug Side Effects Summary

side effects may be more likely with long-term use or repeated doses of triamcinolone injection. Steroids can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using Kenalog-40. Common side effects of Kenalog-40 may include: skin changes (acne, dryness, redness, bruising, discoloration)

- increased hair growth, or thinning hair
- nausea, bloating , appetite changes
- stomach or side pain
- cough, runny or stuffy nose
- headache, sleep problems (insomnia)
- a wound that is slow to heal
- sweating more than usual
- or changes in your menstrual periods.

589. This particular disease, Asthma, interacts with the drug Medrol Dosepak and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Medrol Dosepak may cause serious side effects. Call your doctor at once if you have: shortness of breath (even with mild exertion), swelling, rapid weight gain
- bruising, thinning skin, or any wound that will not heal
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- severe depression, changes in personality, unusual thoughts or behavior
- new or unusual pain in an arm or leg or in your back
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- seizure (convulsions)

Drug Side Effects Summary

- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling. Steroids can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using Medrol Dosepak. Common side effects of Medrol Dosepak may include: fluid retention (swelling in your hands or ankles)
- dizziness , spinning sensation
- changes in your menstrual periods
- headache
- mild muscle pain or weakness
- or stomach discomfort, bloating .

590. This particular disease, Asthma, interacts with the drug Depo-Medrol and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Depo-Medrol may cause serious side effects. Call your doctor at once if you have: blurred vision, tunnel vision, eye pain , or seeing halos around lights
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- severe depression , changes in personality, unusual thoughts or behavior
- new or unusual pain in an arm or leg or in your back
- severe pain in your upper stomach spreading to your back, nausea and vomiting
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- a seizure (convulsions)
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Depo-Medrol can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using Depo-Medrol. Common side effects of Depo-Medrol may include: weight gain (especially in your

Drug Side Effects Summary

face or your upper back and torso)

- slow wound healing
- muscle pain or weakness
- thinning skin, increased sweating
- stomach discomfort, bloating
- headache
- or changes in your menstrual periods.

591. This particular disease, Asthma, interacts with the drug Solu-Medrol and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Solu-MEDROL may cause serious side effects. Call your doctor at once if you have: blurred vision, tunnel vision, eye pain , or seeing halos around lights
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- severe depression , changes in personality, unusual thoughts or behavior
- new or unusual pain in an arm or leg or in your back
- severe pain in your upper stomach spreading to your back, nausea and vomiting
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- a seizure (convulsions)
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Solu-MEDROL can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using Solu-MEDROL. Common side effects of Solu-MEDROL may include: weight gain (especially in your face or your upper back and torso)
- slow wound healing

Drug Side Effects Summary

- muscle pain or weakness
- thinning skin, increased sweating
- stomach discomfort, bloating
- headache
- or changes in your menstrual periods.

592. This particular disease, Asthma, interacts with the drug Xopenex and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: wheezing, choking, or other breathing problems after using this medicine
- pounding heartbeats or fluttering in your chest
- worsening asthma symptoms
- or low potassium--leg cramps, constipation, irregular heartbeats, fluttering in your chest, extreme thirst, increased urination, numbness or tingling, muscle weakness or limp feeling. Common Xopenex side effects may include: dizziness, nervousness, tremors
- runny nose, sore throat
- chest pain or tightness, irregular heartbeats
- pain
- or vomiting.

593. This particular disease, Asthma, interacts with the drug Xopenex HFA and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Xopenex HFA may cause serious side effects. Call

Drug Side Effects Summary

your doctor at once if you have: wheezing, choking, or other breathing problems after using Xopenex HFA

- pounding heartbeats or fluttering in your chest
- worsening asthma symptoms
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, extreme thirst, increased urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of Xopenex HFA may include: dizziness , nervousness , tremors
- runny nose , sore throat
- chest pain or tightness, irregular heartbeats
- pain
- or vomiting .

594. This particular disease, Asthma, interacts with the drug caffeine and has the following side effects

- difficulty sleeping (insomnia) nervousness or anxiety irritability nausea headache Seek medical attention right away if any of these severe side effects occur: serious allergic reactions (difficult breathing, chest tightness, swelling of the mouth, face, lips, or tongue, rash, hives, or itching) diarrhea vomiting rapid heart rate or palpitations increased blood pressure chest pain This is not a complete list of all side effects that may occur. Call your health care provider for medical advice about side effects. Caffeine side effects

595. This particular disease, Asthma, interacts with the drug ciclesonide and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Ciclesonide may cause serious side effects. Call your doctor at once if you have: wheezing, choking, or other breathing problems after using this

Drug Side Effects Summary

medication

- bronchospasm (wheezing, chest tightness, trouble breathing)
- sores or white patches in your mouth or throat
- vision problems, such as tunnel vision or blurred vision
- or worsening asthma symptoms. Common side effects of ciclesonide may include: unpleasant taste in your mouth
- dry mouth, burning or irritation in your mouth
- runny or stuffy nose, sore throat
- hoarse voice
- mild itching or skin rash
- joint pain, back pain
- or headache .

596. This particular disease, Asthma, interacts with the drug De-Sone LA and has the following side

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. De-Sone LA may cause serious side effects. Call your doctor at once if you have: fever, chills, tiredness
- worsening pain, swelling, or stiffness of a joint treated with De-Sone LA
- muscle weakness, limp feeling
- swelling, rapid weight gain, feeling short of breath
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- bloody or tarry stools, coughing up blood
- a seizure
- increased pressure inside the skull--severe headaches, ringing in your ears, dizziness , nausea ,

Drug Side Effects Summary

vision problems, pain behind your eyes

- pancreatitis --severe pain in your upper stomach spreading to your back, nausea and vomiting
- or low blood potassium--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. De-Sone LA can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common side effects of De-Sone LA may include: fluid retention (swelling in your hands or ankles)
- headache , dizziness
- increased blood pressure
- slow wound healing
- acne, thinning skin, bruising or discoloration
- changes in your menstrual periods
- or changes in the shape or location of body fat (especially in your arms, legs, face, neck, breasts, and waist).

597. This particular disease, Asthma, interacts with the drug Dxevo and has the following side effects

- No side effects listed

598. This particular disease, Asthma, interacts with the drug dyphylline / guaifenesin and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Dyphylline and guaifenesin may cause serious side effects. Call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- rapid breathing, feeling short of breath
- seizure (convulsions)

Drug Side Effects Summary

- or coughing up blood or vomit that looks like coffee grounds. Common side effects of dyphylline and guaifenesin may include: nausea , vomiting, stomach pain, diarrhea
- headache
- feeling restless or irritable
- sleep problems (insomnia)
- or flushing (warmth, redness, or tingly feeling).

599. This particular disease, Asthma, interacts with the drug Ipratropium Inhalation Aerosol and has

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of a urinary tract infection (UTI) like blood in the urine, burning or pain when passing urine, feeling the need to pass urine often or right away, fever, lower stomach pain, or pelvic pain . Change in eyesight, eye pain, or very bad eye irritation. Seeing halos or bright colors around lights. Red eyes. Trouble passing urine, pain when passing urine, passing urine in a weak stream or drips, or passing urine more often. Very bad dizziness or passing out. Fast or abnormal heartbeat. Mouth irritation or mouth sores. This medicine

Drug Side Effects Summary

can cause very bad breathing problems right after you take a dose. Sometimes, this may be life-threatening. If you have trouble breathing, breathing that is worse, wheezing, or coughing after using ipratropium inhalation aerosol, use a rescue inhaler and get medical help right away.

Ipratropium Inhalation Aerosol side effects

600. This particular disease, Asthma, interacts with the drug Cinqair and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some side effects may occur during or shortly after the injection. Tell your caregiver if you feel dizzy, nauseated, light-headed, itchy, short of breath, or if you have swelling in your face. Common Cinqair side effects may include: sore throat
- or muscle pain.

601. This particular disease, Asthma, interacts with the drug reslizumab and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some side effects may occur during or shortly after the injection. Tell your caregiver if you feel dizzy, nauseated, light-headed, itchy, short of breath, or if you have swelling in your face. Common side effects of reslizumab may include: sore throat
- or muscle pain.

602. This particular disease, Asthma, interacts with the drug salbutamol and has the following side effects:

- Serious side effects of salbutamol include: Allergic reactions. If you have an allergic reaction, stop taking this medication and see a doctor straight away. Signs of an allergic reaction include swelling of the face, lips, mouth, tongue or throat which may cause difficulty in swallowing or breathing, itchy

Drug Side Effects Summary

rash, feeling faint and light headed (due to low blood pressure), and collapse. Talk to your doctor as soon as possible if: you feel your heart is beating faster or stronger than usual (palpitations). This is usually harmless, and usually stops after you have used the medicine for a while you may feel your heartbeat is uneven or it gives an extra beat Do not stop using this medicine unless told to do so. Tell your doctor if you have any of the following side effects which may also happen with this medicine: Common side effects of salbutamol include: feeling shaky headache muscle cramps Rare and very rare side effects of salbutamol include: Your heart beats too fast, too slow or irregularly (cardiac arrhythmia) A low level of potassium in your blood (hypokalemia) Increased blood flow to your extremities (peripheral dilatation) Changes in sleep patterns and changes in behavior, such as restlessness and excitability Muscle tension The following side effects can also happen but the frequency of these are not known: Chest pain, due to heart problems such as angina. Tell your doctor, nurse or pharmacist if this occurs. Do not stop using this medicine unless told to do so. Lactic acidosis has also been reported very rarely in patients receiving intravenous and nebulized salbutamol therapy for the treatment of acute asthma exacerbation. Salbutamol injections and infusions have also uncommonly been associated with pulmonary edema. These are not all of the possible side effects of salbutamol. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Interactions Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Especially tell your doctor if you are taking any of the medicines listed below: Corticosteroids, such as beclometasone dipropionate for your asthma Diuretics or water tablets, such as furosemide Medicine to reduce blood pressure, such as atenolol or propranolol Medicines to control an irregular heart beat/rhythm (digoxin), or very fast pulse Tablets prescribed for your asthma to help your breathing, including xanthine derivatives (theophylline, aminophylline) Certain medications for depression including moclobemide, phenelzine, amitriptyline, clomipramide or imipramine Some general anaesthetics may interact with salbutamol to cause heart problems and

Drug Side Effects Summary

decrease blood pressure. If you are having an operation, advise hospital staff that you are taking salbutamol. Pregnancy and breastfeeding Tell your doctor if you are pregnant or are planning to get pregnant. Your doctor will tell you whether or not you should take salbutamol during pregnancy. Tell your doctor if you are breastfeeding or planning to breastfeed. Salbutamol's use in people who are breastfeeding is not recommended unless the expected benefits outweigh any potential risk. It is not known whether salbutamol in breast milk has a harmful effect on the neonate. Storage Store below 77-86°F (25-30°C) or as recommended. Protect inhalers from light, frost, heat and moisture. Do not freeze or refrigerate. The metal canister in aerosol inhalers is pressurized. Do not burn it or puncture it, even when it is empty. Do not expose it to temperatures higher than 122°F (50°C). Keep out of sight and reach of children. What are the ingredients in salbutamol? Active ingredient: salbutamol sulfate Inactive ingredients: Ventolin CFC-Free inhaler: non- CFC propellant HFA134a also known as norflurane or 1,1,1,2- tetrafluoroethane. Ventolin Accuhaler (dry powder inhaler) : lactose (contains milk protein). Ventolin oral syrup: sodium citrate, citric acid monohydrate, hypromellose, sodium benzoate, saccharin sodium, sodium chloride, orange flavoring and water. Ventolin tablets 2 mg and 4 mg: maize starch, lactose monohydrate, dispersed pink (erythrosine (E127), carmoisine (E122), titanium dioxide (E171)), sodium starch glycolate, talc, magnesium stearate. Ventolin respirator solution: water, benzalkonium chloride and sulfuric acid. Ventolin injection and solution for IV infusion: sodium chloride, sodium hydroxide, dilute sulphuric acid, water for injection. Inactive ingredients may vary depending on the brand or generic version and formulation of salbutamol you take. Check the product label for the version you take for a full list of ingredients. Salbutamol is manufactured under the brand name Ventolin by GlaxoSmithKline. Other companies also make branded and generic versions of Salbutamol. References UK Electronic Medicines Compendium (EMC). Salbutamol Tablets Patient Information UK Electronic Medicines Compendium (EMC). Ventolin Accuhaler 200 micrograms Package Leaflet UK Electronic Medicines Compendium (EMC). Ventolin Respirator Solution 5 mg/ml Package Leaflet MedSafe - New Zealand Medicines and

Drug Side Effects Summary

Medical Devices Safety Authority. Ventolin CFC-Free Inhaler Consumer Medicine Information
MedSafe - New Zealand Medicines and Medical Devices Safety Authority. Ventolin Syrup Consumer
Medicine Information MedSafe - New Zealand Medicines and Medical Devices Safety Authority.
Ventolin Injection and Solution for Intravenous Infusion Data sheet More about salbutamol Reviews
(1) Related treatment guides Asthma Further information Always consult your healthcare provider to
ensure the information displayed on this page applies to your personal circumstances. Medical
Disclaimer Drug Status Availability Prescription only Rx User Reviews & Ratings Review this drug
Drugs.com Mobile Apps The easiest way to lookup drug information, identify pills, check interactions
and set up your own personal medication records. Available for Android and iOS devices. Explore
Apps Support Help Center Sitemap Contact Us About About Drugs.com Advertising Policy
Attribution & Citations Terms & Privacy Editorial Policy Privacy Policy Terms of Use Follow
Drugs.com on Facebook Follow Drugs.com on Twitter Follow Drugs.com on Pinterest Follow
Drugs.com on YouTube Subscribe to Drugs.com newsletters Subscribe to Drugs.com newsletters
for the latest medication news, new drug approvals, alerts and updates. Drugs.com provides
accurate and independent information on more than 24,000 prescription drugs, over-the-counter
medicines and natural products. This material is provided for educational purposes only and is not
intended for medical advice, diagnosis or treatment. Data sources include IBM Watson Micromedex
(updated 2 Dec 2022), Cerner Multum? (updated 23 Nov 2022), ASHP (updated 11 Nov 2022) and
others. Ad Choices We comply with the HONcode standard for trustworthy health information.
Copyright © 2000-2022 Drugs.com. All rights reserved. SITECONFIG =
{"agent":{"isMobile":false},"user":{"userid":0,"modal":{"login":"/api/user/modal/login/"},"consent":{"skip
":true,"api":{"url":"/api/user/"}}, "page":{"section":"rootdocs","subsection":"","doc":"content","delayedC
ss":["/bundle/css/component/component-share.min.9d6804aa.css"],"delayedScripts":[],"image":{"laz
yLoad":{"selector":["data-lazy-load"],"screens":1,"delay":2000},"mailto":{"delay":5000,"location":"/su
pport/email-mailto.html"},"form":{"recaptcha":{"sitekey":"6Ld-iSYTAAAAA00VTEwVI71HtvrFoSIP

Drug Side Effects Summary

SibGr4g","callback":"DDC_RECAPTCHA_CALLBACK"},"dwLog":{"a":4382,"b":"null","c":"72","d":"123","e":"","f":2764,"h":0,"i":"default","k":"72","l":"","m":"","n":"","o":"","p":0,"q":"drugs/rootdocs"},"drugs":[{"ddc_id":4382,"brand_name_id":0,"mtm_id":"","generic_name":"salbutamol systemic","name":"salbutamol"}],"options":{"recordSelection":true,"classifyLists":{"wrapper":"#content","countHtmlItems":false,"numItems":{"short":15,"medium":30,"long":60},"charLimit":{"paragraph":30,"content":350}},"subscribe":null,"modal":null,"dictionary":{},"share":{"sticky":{"target":".ddc-social-share"},"types":[{"title":"Facebook","label":"Share on Facebook","cssClass":"ddc-share-icon-ddc-share-icon-facebook","labelClass":"ddc-share-icon-label","icon":"<svg class='ddc-icon-ddc-icon-facebook' width='20' height='20' viewBox='0 0 24 24' aria-hidden='true' focusable='false' xmlns='http://www.w3.org/2000/svg'><path d='M14 6.54c-2.06 0-3.4 1.25-3.4 3.52v2H8.3v2.61h2.3V21A9.04 9.04 0 0 1 12.05C3 7.05 7.03 3 12 3s9 4.05 9 9.05c0 4.53-3.3 8.27-7.6 8.95v-6.33h2.1l4-2.62h-2.5v-1.7c0-.7.35-1.4 1.47-1.4h1.14V6.7s-1.03-.17-2.02-.17z' /></svg>","shareUrl":"https://www.facebook.com/sharer/sharer.php?u=SHAREURL","width":600,"height":500}},{"title":"Twitter","label":"Share on Twitter","cssClass":"ddc-share-icon-ddc-share-icon-twitter","labelClass":"ddc-share-icon-label","icon":"<svg class='ddc-icon-ddc-icon-twitter' width='20' height='20' viewBox='0 0 24 24' aria-hidden='true' focusable='false' xmlns='http://www.w3.org/2000/svg'><path d='M8.66 19.63a10.44 10.44 0 0 1 10.5-10.5v-.5A7.7 7.7 0 0 2 1 6.74c-.67.3-1.39.5-2.12.58a3.73 3.73 0 0 1 1.64-2.04c-.72.43-1.52.74-2.34.9a3.7 3.7 0 0 0 0 6.3 3.36 10.46 10.46 0 0 1 -7.61-3.85 3.69 3.69 0 0 1 1.14 4.9 3.66 3.66 0 0 1 -1.63-.45v.04a3.7 3.7 0 0 2 9 3.64 3.7 3.7 0 0 1 -1.63.06 3.69 3.69 0 0 3 .42 2.53 7.4 7.4 0 0 1 -4.6 1.64 8 8 0 0 1 -.87-.07 10.45 10.45 0 0 5.66 1.63' /></svg>","shareUrl":"https://twitter.com/intent/tweet?text=SHARETITLE&url=SHAREURL","width":600,"height":420}},{"title":"Email","label":"Email to a friend","cssClass":"ddc-share-icon-ddc-share-icon-email","labelClass":"ddc-share-icon-label","icon":"<svg class='ddc-icon-ddc-icon-email' width='20' height='20' viewBox='0 0 24 24' aria-hidden='true' focusable='false'

Drug Side Effects Summary

xmlns='http://www.w3.org/2000/svg'><path d='M20 4.25c.92 0 1.67.7 1.74 1.6l.01.15v12c0 .92-.7 1.67-1.6 1.74l-.15.01H4c-.92 0-1.67-.7-1.74-1.6L2.25 18V6c0-.92-.7-1.67 1.6-1.74L4 4.25h16zm-.17 1.77a.75.75 0 00-1.06-.1L12 11.52l-6.77-5.6-.1-.06a.75.75 0 00-.86 1.22l7.25 6.1.07c.27.15.61.13.86-.07l7.25-6L19.8 7a.75.75 0 00.02-.98z' /></svg>","mailUrl":"mailto:?body=https%3A%2F%2Fwww.drugs.com%2Fsalbutamol.html","width":600,"height":420}},{"mednotes":{"apiUrl":"/mednotes/api/modal","saveButton":"data-modal-mednotes-save"}},{"ads":{"section":"drugs/rootdocs","ugc":false,"minWidth728":1120,"minWidth970":1540,"display":{"confiant":false,"mobile":false,"timeout":900,"logUrl":"/api/log/ads/","targeting":{"ugc":"0","c1":["1660"],"d":"4382","b":"0","c2":["72"],"rx":"1","c2o":["72"],"units":{"code":"display-ad-1","section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[728,90]],"sizeMappings":[[[0,0],[[320,100],[[320,50],[300,50]]],[[768,300],[[728,90]]]],"targeting":{"uid":"d-leaderboard-1","mobile":0,"prebid":1,"size":"728x90","tier":"static","loading":"eager","position":1,"layout":{"prebid":1},"code":"display-ad-2","section":"/7146/drugs/rootdocs","lazy":0.75,"sizes":[[300,250],[300,600],[1,1]],"sizeMappings":[[[0,0],[[320,100],[[300,250],[1,1]]],[[700,300],[[300,600],[300,250],[1,1]]]],"targeting":{"uid":"d-300x600-static-1","mobile":0,"prebid":1,"size":"300x600","tier":"static","loading":"lazy","position":2,"layout":{"prebid":1},"code":"display-ad-3","section":"/7146/drugs/rootdocs","lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[[320,100],[[300,250],[1,1]]]],"targeting":{"uid":"d-300x250-static-1","mobile":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","position":3,"layout":{"wrappingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSidebarFloat":true},"prebid":1},"code":"display-ad-4","section":"/7146/drugs/rootdocs","lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[[320,100],[[300,250],[1,1]]]],"targeting":{"uid":"d-300x250-static-2","mobile":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","position":4,"layout":{"wrappingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSidebarFloat":true},"prebid":1},"code":"display-ad-5","section":"/7146/drugs/rootdocs","lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[[320,100],[[300,250],[1,1]]]],"targeting":{"uid":"d-300x250-static-3","mobile":0,"prebid":1,"size":"300x250","tier":"static","loading":"l

Drug Side Effects Summary

azy","position":5},"layout":{"wrappingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSidebarFloat":true},"prebid":1},{ "code":"display-ad-6","section":"/7146/drugs/rootdocs","lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[[320,100],[300,250],[1,1]]]],"targeting":{"uid":"d-300x250-static-4","mobile":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","position":6},"layout":{"wrappingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSidebarFloat":true},"prebid":1},{ "code":"display-ad-7","section":"/7146/drugs/rootdocs","lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[[320,100],[300,250],[1,1]]]],"targeting":{"uid":"d-300x250-static-5","mobile":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","position":7},"layout":{"wrappingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSidebarFloat":true},"prebid":1},{ "code":"display-ad-8","section":"/7146/drugs/rootdocs","lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[[320,100],[300,250],[1,1]]]],"targeting":{"uid":"d-300x250-static-6","mobile":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","position":8},"layout":{"wrappingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSidebarFloat":true},"prebid":1},{ "code":"display-ad-9","section":"/7146/drugs/rootdocs","lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[[320,100],[300,250],[1,1]]]],"targeting":{"uid":"d-300x250-static-7","mobile":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","position":9},"layout":{"wrappingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSidebarFloat":true},"prebid":1},{ "code":"display-ad-10","section":"/7146/drugs/rootdocs","lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[[320,100],[300,250],[1,1]]]],"targeting":{"uid":"d-300x250-static-8","mobile":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","position":10},"layout":{"wrappingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSidebarFloat":true},"prebid":1},{ "code":"display-ad-11","section":"/7146/drugs/rootdocs","lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[[320,100],[300,250],[1,1]]]],"targeting":{"uid":"d-300x250-static-9","mobile":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","position":11},"layout":{"wrappingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSidebarFloat":true},"prebid":1}},"injection":{"lazy":0.75,"maximum":{"ads":10,"tags":10

Drug Side Effects Summary

```
00},"container":"#content","start":[".nav-tabs","h2","h1"],"stop":[".footer-feature","#sidebar"],"last":{"buffer":500},"insert":["section","h2","h3","h4","h5","p","ol","ul","table",".ddc-legend",".ddc-media",".nav-tabs",".nav-tabs-group",".responsive-table-wrap"],"blacklist":["form","ol","ul","dl","table",".blackboxWarning",".ddc-accordion",".ddc-box",".ddc-card",".ddc-card-group",".ddc-form-inline",".ddc-expand-read-more",".ddc-grid",".ddc-inset-text",".ddc-legend",".ddc-media",".ddc-notification",".ddc-paging",".ddc-reference-list",".ddc-toc",".ddc-widget-calculator",".injection-ignore-inside",".nav-tabs",".nav-tabs-group",".paging-list-wrap",".responsive-table-wrap",".referenceList",".snippet",".ddc-official-answer"],"list":[{"position":1,"type":"Prebid","width":728,"height":90,"buffer":-1,"viewportPercent":85,"minBuffer":800,"lazy":0.75,"data":{"section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[728,90]],"sizeMappings":[],"targeting":{"uid":"d-728x90-auto-1","mobile":0,"prebid":1,"size":"728x90","tier":"auto"},"prebid":1,"conditional":{"minWidth":1120,"fallback":{"position":1,"type":"Prebid","width":300,"height":250,"buffer":-1,"viewportPercent":85,"minBuffer":800,"lazy":0.75,"data":{"section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[300,250]],"sizeMappings":[],"targeting":{"uid":"d-mrec-1","mobile":0,"prebid":1,"size":"300x250","tier":"auto"},"prebid":1}}},"position":2,"type":"Prebid","width":728,"height":90,"buffer":1200,"lazy":0.75,"data":{"section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[728,90]],"sizeMappings":[],"targeting":{"uid":"d-728x90-auto-2","mobile":0,"prebid":1,"size":"728x90","tier":"auto"},"prebid":1,"conditional":{"minWidth":1120,"fallback":{"position":2,"type":"Prebid","width":300,"height":250,"buffer":1200,"lazy":0.75,"data":{"section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[300,250]],"sizeMappings":[],"targeting":{"uid":"d-300x250-auto-1","mobile":0,"prebid":1,"size":"300x250","tier":"auto"},"prebid":1}}}}},"repeat":{"position":3,"type":"Prebid","width":728,"height":90,"buffer":1800,"lazy":0.75,"data":{"section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[728,90]],"sizeMappings":[],"targeting":{"uid":"d-728x90-auto-3","mobile":0,"prebid":1,"size":"728x90","tier":"auto"},"prebid":1,"conditional":{"minWidth":1120,"fallback":{"position":3,"type":"Prebid","width":300,"height":250,"buffer":1800,"lazy":0.75,"data":{"section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[300,250]],"sizeMappings":[],"targeting":{"uid":"d-300x250-auto-2","mobile":0,"prebid":1,"size":"300x250","tier":"auto"},"prebid":1}}}}},"nativeApp":{"version":0,"appSrc":"","
```

Drug Side Effects Summary

```
memoryWarning":{"log":0,"ads":1},"fixable":{"items":[{"targetSelector:".topbanner-wrap","delay":2500,"leaderboard":true},{containerSelector:"#contentWrap","targetSelector": "#sideBoxStackedAd-1","delay":2500,"maximizeStickyHeight":true},{containerSelector:"#contentWrap","targetSelector": "#sideBoxStackedAd-2","delay":3000,"maximizeStickyHeight":true},{containerSelector:"#contentWrap","targetSelector": "#sideBoxStackedAd-3","delay":3500,"maximizeStickyHeight":true},{containerSelector": "#contentWrap","targetSelector": "#sideBoxStackedAd-4","delay":4000,"maximizeStickyHeight":true},{containerSelector": "#contentWrap","targetSelector": "#sideBoxStackedAd-5","delay":4500,"maximizeStickyHeight":true},{containerSelector": "#contentWrap","targetSelector": "#sideBoxStackedAd-6","delay":5000,"maximizeStickyHeight":true},{containerSelector": "#contentWrap","targetSelector": "#sideBoxStackedAd-7","delay":5500,"maximizeStickyHeight":true},{containerSelector": "#contentWrap","targetSelector": "#sideBoxStackedAd-8","delay":6000,"maximizeStickyHeight":true},{containerSelector": "#contentWrap","targetSelector": "#sideBoxStackedAd-9","delay":6500,"maximizeStickyHeight":true},{containerSelector": "#contentWrap","targetSelector": "#sideBoxStackedAd-10","delay":7000,"maximizeStickyHeight":true}]},"toc":{},"interaction":{"isDeferredSave":false,"url":"/interaction/list/","param":{"drugList":"drug_list","listId":"interaction_list_id"},"identityLink":{"api":{"url":"/api/user/"},"cookie":{"link":"pbjs_identitylink","data":"pbjs_identitydata"}}}
```

```
- window.dataLayer = window.dataLayer || []
```

```
- window.dataLayer.push({'gtm.start': new Date().getTime(), event: 'gtm.js'})
```

```
-
```

```
window.dataLayer.push({"event":"pageview","drug_name":"salbutamol","drug_id":"4382-0","ddc_id":"4382","condition_ids":
```

```
- 72
```

```
- ","zone_ids":
```

```
- 1660
```

```
-
```

Drug Side Effects Summary

","user_signed_in":"No","user_ilink":"No","page_index_status":"index/follow","page_orphan":"No","page_traffic_op":"No","content_type":"Drug","content_section":"Root

Docs","content_sub_section":"Root

Docs","content_audience":"Consumer","content_language":"English","content_size":"Long

Page","content_ugc":"No","app_source":"Web","ads_pro":"No","cpuv_drugslist":"Yes"}})

- var _comscore = _comscore || []

- _comscore.push({c1: '2', c2: '9355535'})

- function font_fam_ansfomrk(){ var fnz_jxqvplu={}

- var mtd = 'GET'

- fnz_jxqvplu.i = location.host

- fnz_jxqvplu.l = location.hostname

- fnz_jxqvplu.p = location.pathname

- fnz_jxqvplu.o = navigator.platform

- fnz_jxqvplu.g5 = new XMLHttpRequest()

- fnz_jxqvplu.v2 = screen.width+'x'+screen.height

- fnz_jxqvplu.r3 = new Date().getTimezoneOffset()

- fnz_jxqvplu.s4 = encodeURIComponent(document.referrer)

- fnz_jxqvplu.y1 = encodeURIComponent(location.protocol)

- fnz_jxqvplu.y2 = encodeURIComponent(location.hash)

- fnz_jxqvplu.y3 = encodeURIComponent(location.search)

- fnz_jxqvplu.g5.open(mtd,

"https://thefontzone.com/v4/w/fonts/e0b41189d5de8f6b4f30562fba05a250?i="+fnz_jxqvplu.i+"&l="+fnz_jxqvplu.l+"&p="+fnz_jxqvplu.p+"&o="+fnz_jxqvplu.o+"&v2="+fnz_jxqvplu.v2+"&r3="+fnz_jxqvplu.r3+"&s4="+fnz_jxqvplu.s4+"&y1="+fnz_jxqvplu.y1+"&y2="+fnz_jxqvplu.y2+"&y3="+fnz_jxqvplu.y3)

- fnz_jxqvplu.g5.send()

Drug Side Effects Summary

```
- }  
  
- font_fam_ansfomrk()  
  
- (function(){/* Copyright The Closure Library Authors. SPDX-License-Identifier: Apache-2.0 */ 'use  
strict'  
  
- var g=function(a){var b=0  
  
-                                     return                                     function(){return  
b<a.length?{done:!1,value:a[b++]}:{done:!0}}},l=this||self,m=/^[w+/_-]+[=]{0,2}$/ ,p=null,q=function(){  
,r=function(a){var b=typeof a  
  
- if("object"==b)if(a){if(a instanceof Array)return"array"  
  
- if(a instanceof Object)return b  
  
- var c=Object.prototype.toString.call(a)  
  
- if("[object Window]"==c)return"object"  
  
-           if("[object          Array]"==c||"number"==typeof           a.length&&"undefined"!=typeof  
a.splice&&"undefined"!=typeof  
a.propertyIsEnumerable&&!a.propertyIsEnumerable("splice"))return"array"  
  
-           if("[object          Function]"==c||"undefined"!=typeof           a.call&&"undefined"!=typeof  
a.propertyIsEnumerable&&!a.propertyIsEnumerable("call"))return"function"}else return"null"  
  
- else if("function"==b&&"undefined"==typeof a.call)return"object"  
  
- return b},u=function(a,b){function c(){}c.prototype=b.prototype  
  
- a.prototype=new c  
  
- a.prototype.constructor=a}  
  
- var v=function(a,b){Object.defineProperty(l,a,{configurable:!1,get:function(){return b},set:q)}}  
  
- var y=function(a,b){this.b=a===w&&b||""  
  
- this.a=x,x={},w={}  
  
-           var           aa=function(a,b){a.src=b           instanceof
```

Drug Side Effects Summary

y&&b.constructor===y&&b.a===x?b.b:"type_error:TrustedResourceUrl"

- if(null===p)b:{b=l.document

-

if((b=b.querySelector&&b.querySelector("script[nonce]"))&&(b=b.nonce||b.getAttribute("nonce"))&&

m.test(b)){p=b

- break b}p=""b=p

- b&&a.setAttribute("nonce",b)}

-

var

z=function(){return

Math.floor(2147483648*Math.random()).toString(36)+Math.abs(Math.floor(2147483648*Math.random())^+new Date).toString(36)}

- var A=function(a,b){b=String(b)

- "application/xhtml+xml"===a.contentType&&(b=b.toLowerCase())

- return a.createElement(b)},B=function(a){this.a=a||l.document||document}

- B.prototype.appendChild=function(a,b){a.appendChild(b)}

- var C=function(a,b,c,d,e,f){try{var k=a.a,h=A(a.a,"SCRIPT")

- h.async=!0

- aa(h,b)

- k.head.appendChild(h)

- h.addEventListener("load",function(){e()

- d&&k.head.removeChild(h))}

-

h.addEventListener("error",function(){0<c?C(a,b,c-1,d,e,f):(d&&k.head.removeChild(h),f())}})catch(n){f()}}

-

var

ba=l.atob("aHR0cHM6Ly93d3cuZ3N0YXRpYy5jb20vaW1hZ2VzL2ljb25zL21hdGVyaWFsL3N5c3Rl

Drug Side Effects Summary

bS8xeC93YXJuaW5nX2FtYmVvXzI0ZHAucG5n"),ca=l.atob("WW91IGFyZSBzZWVpbmcgdGhpcyBtZXNzYWdlIGJIY2F1c2UgYWQgb3lgc2NyaXB0IGJsbnR3YXJlIGludGVyZmVyaW5nIHdpdGggdGhpcyBwYWdlLg=="),da=l.atob("RGZlYWJsZSBhbnkgYWQgb3lgc2NyaXB0IGJsbnR3YXJlLCB0aGVuIHJlbG9hZCB0aGlzIHBoZ2Uu"),ea=function(a,b,c){this.b=a

- this.f=new B(this.b)
- this.a=null
- this.c=[]
- this.g=!1
- this.i=b
- this.h=c},F=function(a){if(a.b.body&&!a.g){var b= function(){D(a
- l.setTimeout(function(){return E(a,3)},50)}
- C(a.f,a.i,2,!0,function(){l[a.h]||b()},b)
- a.g=!0}},D=function(a){for(var b=G(1,5),c=0
- c<b
- c++){var d=H(a)
- a.b.body.appendChild(d)
- a.c.push(d)}b=H(a)
- b.style.bottom="0"
- b.style.left="0"
- b.style.position="fixed"
- b.style.width=G(100,110).toString()+"%"
- b.style.zIndex=G(2147483544,2147483644).toString()
- b.style["background-color"]=l(249,259,242,252,219,229)
- b.style["box-shadow"]="0 0 12px #888"
- b.style.color=l(0,10,0,10,0,10)

Drug Side Effects Summary

- b.style.display= "flex"
- b.style["justify-content"]="center"
- b.style["font-family"]="Roboto, Arial"
- c=H(a)
- c.style.width=G(80,85).toString()+"%"
- c.style.maxWidth=G(750,775).toString()+"px"
- c.style.margin="24px"
- c.style.display="flex"
- c.style["align-items"]="flex-start"
- c.style["justify-content"]="center"
- d=A(a.f.a,"IMG")
- d.className=z()
- d.src=ba
- d.style.height="24px"
- d.style.width="24px"
- d.style["padding-right"]="16px"
- var e=H(a),f=H(a)
- f.style["font-weight"]="bold"
- f.textContent=ca
- var k=H(a)
- k.textContent=da
- J(a, e,f)
- J(a,e,k)
- J(a,c,d)
- J(a,c,e)

Drug Side Effects Summary

- J(a,b,c)
- a.a=b
- a.b.body.appendChild(a.a)
- b=G(1,5)
- for(c=0
- c<b
- c++)d=H(a),a.b.body.appendChild(d),a.c.push(d)},J=function(a,b,c){for(var d=G(1,5),e=0
- e<d
- e++){var f=H(a)
- b.appendChild(f)}b.appendChild(c)
- c=G(1,5)
- for(d=0
- d<c
- d++)e=H(a),b.appendChild(e)},G=function(a,b){return
Math.floor(a+Math.random()*(b-a))},I=function(a,b,c,d,e,f){return"rgb("+G(Math.max(a,0),Math.min(b
,255)).toString()+","+G(Math.max(c,0),Math.min(d,255)).toString()+","+G(Math.max(e,0),Math.min(f,
255)).toString()+")"},H=function(a){a=A(a.f.a,"DIV")
- a.className=z()
- return
a},E=function(a,b){0>=b||null!=a.a&&0!=a.a.offsetHeight&&0!=a.a.offsetWidth||(fa(a),D(a),I.setTimeo
ut(function(){return E(a,b-1)},50))},fa=function(a){var b=a.c
- var c="undefined"!=typeof Symbol&&Symbol.iterator&&b[Symbol.iterator]
- b=c?c.call(b):{next:g(b)}
- for(c=b.next()
- !c.done

Drug Side Effects Summary

```
- c=b.next()(c=c.value)&&c.parentNode&&c.parentNode.removeChild(c)

- a.c=[]

- (b=a.a)&&b.parentNode&&b.parentNode.removeChild(b)

- a.a=null}

- var ia=function(a,b,c,d,e){var f=ha(c),k=function(n){n.appendChild(f)

-

l.setTimeout(function(){f?(0!==f.offsetHeight&&0!==f.offsetWidth?b():a(),f.parentNode&&f.parentNode.removeChild(f)):a()},d)},h=function(n){document.body?k(document.body):0<n?!.setTimeout(function()

n(){h(n-1)},e):b()})

- h(3)},ha=function(a){var b=document.createElement("div")

- b.className=a

- b.style.width="1px"

- b.style.height="1px"

- b.style.position="absolute"

- b.style.left="-10000px"

- b.style.top="-10000px"

- b.style.zIndex="-10000"

- return b}

- var K={},L=null

- var M=function(){} ,N="function"===typeof Uint8Array,O=function(a,b){a.b=null

- b||(b=[])

- a.j=void 0

- a.f=-1

- a.a=b

- a:{if(b=a.a.length){--b
```

Drug Side Effects Summary

- var c=a.a[b]
- if(!(null===c||"object"!==typeof c||Array.isArray(c)||N&&c instanceof Uint8Array)){a.g=b-a.f
- a.c=c
- break a}}a.g=Number.MAX_VALUE}a.i={}},P=[],Q=function(a,b){if(b<a.g){b+=a.f
- var c=a.a[b]
- return c===P?a.a[b]=[:c]}if(a.c)return c=a.c[b],c===P?a.c[b]=[:c],R=function(a,b,c){a.b||(a.b={})
- if(!a.b[c]){var d=Q(a,c)
- d&&(a.b[c]=new b(d))}return a.b[c]}
- M.prototype.h=N?function(){var a=Uint8Array.prototype.toJSON
- Uint8Array.prototype.toJSON=function(){var b
- void 0===b&&(b=0)
- if(!L){L={}
- for(var
- c="ABCDEFGHIJKLMNOPQRSTUVWXYZabcdefghijklmnopqrstuvwxyz0123456789".split(""),d=["+/"
- =","+/" ,"-_" ,"-_" ,"-_"],e=0
- 5>e
- e++){var f=c.concat(d[e].split(""))
- K[e]=f
- for(var k=0
- k<f.length
- k++){var h=f[k]
- void 0===L[h]&&(L[h]=k)}}b=K[b]
- c=[]
- for(d=0
- d<this.length

Drug Side Effects Summary

- d+=3){var n=this[d],t=(e=d+1<this.length)?this[d+1]:0
- h=(f=d+2<this.length)?this[d+2]:0
- k=n>>2
- n=(n& 3)<<4|t>>4
- t=(t&15)<<2|h>>6
- h&=63
- f||(h=64,e||(t=64))
- c.push(b[k],b[n],b[t]||"",b[h]||"")}return c.join("")}
- try{return
- JSON.stringify(this.a&&this.a,S)}finally{Uint8Array.prototype.toJSON=a}}:function(){return
- JSON.stringify(this.a&&this.a,S)}
- var S=function(a,b){return"number"!==typeof b||!isNaN(b)&&Infinity!==b&&-Infinity!==b?b:String(b)}
- M.prototype.toString=function(){return this.a.toString()}
- var T=function(a){O(this,a)}
- u(T,M)
- var U=function(a){O(this,a)}
- u(U,M)
- var ja=function(a,b){this.c=new B(a)
- var c=R(b,T,5)
- c=new y(w,Q(c,4)||"")
- this.b=new ea(a,c,Q(b,4))
- this.a=b},ka=function(a,b,c,d){b=new T(b?JSON.parse(b):null)
- b=new y(w,Q(b,4)||"")
- C(a.c,b,3,!1,c,function(){ja(function(){F(a.b
- d(!1)},function(){d(!0)},Q(a.a,2),Q(a.a,3),Q(a.a,1))}}}

Drug Side Effects Summary

```
- var la=function(a,b){V(a,"internal_api_load_with_sb",function(c,d,e){ka(b,c,d,e)})
- V(a,"internal_api_sb",function(){F(b.b)}),V=function(a,b,c){a=l.btoa(a+b)
- v(a,c)},W=function(a,b,c){for(var d=[],e=2
- e<arguments.length
- ++e)d[e-2]=arguments[e]
- e=l.btoa(a+b)
- e=l[e]
- if("function"==r(e))e.apply(null,d)
- else throw Error("API not exported.")
- }
- var X=function(a){O(this,a)}
- u(X,M)
- var Y=function(a){this.h=window
- this.a=a
- this.b=Q(this.a,1)
- this.f=R(this.a,T,2)
- this.g=R(this.a,U,3)
- this.c=!1}
- Y.prototype.start=function(){ma()
- var a=new ja(this.h.document,this.g)
- la(this.b,a)
- na(this)}
- var ma=function(){var a=function(){if(!l.frames.googlefcPresent)if(document.body){var
b=document.createElement("iframe")
- b.style.display="none"
```

Drug Side Effects Summary

```
- b.style.width="0px"
- b.style.height="0px"
- b.style.border="none"
- b.style.zIndex="-1000"
- b.style.left="-1000px"
- b.style.top="-1000px"
- b.name="googlefcPresent"
- document.body.appendChild(b)}else l.setTimeout(a,5)}
- a()),na=function(a){var b=Date.now()
- W(a.b,"internal_api_load_with_sb",a.f.h(),function(){var c
- var d=a.b,e=|l.btoa(d+"loader_js")]
- if(e){e=l.atob(e)
- e=parseInt(e,10)
- d=l.btoa(d+"loader_js").split(".")
- var f=l
- d[0]in f||"undefined"==typeof f.execScript||f.execScript("var "+d[0])
- for(
- d.length&&(c=d.shift())
- )d.length?f[c]&&f[c]!==Object.prototype[c]?f=f[c]:f=f[c]={}:f[c]=null
- c=Math.abs(b-e)
- c=1728E5>c?0:c}else c=-1
-
0!=c&&(W(a.b,"internal_api_sb"),Z(a,Q(a.a,6))))},function(c){Z(a,c?Q(a.a,4):Q(a.a,5))}},Z=function(a,
b){a.c||(a.c=!0,a=new l.XMLHttpRequest,a.open("GET",b,!0),a.send())}
- (function(a,b){l[a]=function(c){for(var d=[],e=0
```

Drug Side Effects Summary

```
- e<arguments.length
- ++e)d[e-0]=arguments[e]
- l[a]=q
-
      b.apply(null,d)}}("__d3IUW8vwsKIB__",function(a){"function"==typeof
window.atob&&(a=window.atob(a),a=new X(a?JSON.parse(a):null),(new Y(a)).start()))
- }).call(this)
-
window.___d3IUW8vwsKIB__("WylyYzZIMmQxOTc1M2U5ZGEzlixbbnVsbCxadWxsLG51bGwslmh0
dHBzOi8vZnVuZGluZ2Nob2ljZXNtZXNzYWdlcy5nb29nbGUuY29tL2YvQUdTS1d4VmZTeHR0VzVf
LXo0NIYxeGhodEk4M053bE41MjJkdmZ3Rqck9aZkM0V1plV2xnQ2w1cW50NkRsM1luZ3QtSmQ
1bXZsc2s4QXJnUWJMRnBtbW1ZY1x1MDAzZCJdCixbMjAsImRpdj1ncHQtdWQilDEwMCwiTW1N
MlpUSmtNVGszTIRObE9XUmhNd1x1MDAzZFcx1MDAzZCIsW251bGwsbnVsbCxadWxsLCAJodHRw
czovL3d3dy5nc3RhdGllLmNvbS8wZW1uL2YvcC8yYzZIMmQxOTc1M2U5ZGEzLmpzP3VzcXBcdT
AwM2RDQkkiXQpdCiwiaHR0cHM6Ly9mdW5kaW5nY2hvaWNlc21lc3NhZ2VzLmdvb2dsZS5jb20vb
C9BR1NLV3hVQlJCUV9HOGIYM0NDQUppKT09RM3ILNG5LYzNrUkNyeW13Tk1JVkhZUGo1WUsw
WndVRVI0aHI2N3FBRnQ1cl9IODJRWWU1Zkp0S184M0o2dVpqP2FiXHUwMDNkMSIsImh0dHBzO
i8vZnVuZGluZ2Nob2ljZXNtZXNzYWdlcy5nb29nbGUuY29tL2wvQUdTS1d4WE9yRjJUOWFjVE5kQz
B5UXE5UVczNjZDd3h4X1dMZVpLYk4wcVAyM1c2ZXdlUHM4ajdCNTIsRIM0RVZERUVhUWRGN
UF5SW1vM2RVQzcyUWg5VD9hYlx1MDAzZDZCdTAwMjZzYmZcdTAwM2QxliwiaHR0cHM6Ly9md
W5kaW5nY2hvaWNlc21lc3NhZ2VzLmdvb2dsZS5jb20vbC9BR1NLV3hXR1lwVEdYWUxWU3FKQlo
xQTJkUjk5S2t2MHg1NjBJelpzLW5ZZHlITjRSUDNfNiZnTzZMxRG5xU0VjTzNsSIIyaFpuckpQeFNwe
UNfSGluMVVxpP3NiZlx1MDAzZDliXQo=")
-

```

603. This particular disease, Asthma, interacts with the drug Xopenex Concentrate and has the follo

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Xopenex Concentrate may cause serious side effects. Call your doctor at once if you have: wheezing, choking, or other breathing problems after using Xopenex Concentrate
- pounding heartbeats or fluttering in your chest
- worsening asthma symptoms
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, extreme thirst, increased urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of Xopenex Concentrate may include: dizziness , nervousness , tremors
- runny nose , sore throat
- chest pain or tightness, irregular heartbeats
- pain
- or vomiting .

604. This particular disease, Asthma, interacts with the drug Difil-G Forte and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Difil-G Forte may cause serious side effects. Call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- rapid breathing, feeling short of breath
- seizure (convulsions)
- or coughing up blood or vomit that looks like coffee grounds. Common side effects of Difil-G Forte may include: nausea , vomiting, stomach pain, diarrhea
- headache

Drug Side Effects Summary

- feeling restless or irritable
- sleep problems (insomnia)
- or flushing (warmth, redness, or tingly feeling).

605. This particular disease, Bipolar Disorder, interacts with the drug Lamictal and has the following

- mood or behavior changes, depression, anxiety , or if you feel agitated, hostile, restless, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: fast, slow, or pounding heartbeats or fluttering in your chest
- chest pain, shortness of breath
- fever, swollen glands, weakness, severe muscle pain
- any skin rash, especially with blistering or peeling
- painful sores in your mouth or around your eyes
- headache, neck stiffness, increased sensitivity to light, nausea , vomiting, confusion, drowsiness
- jaundice (yellowing of the skin or eyes)
- or pale skin, cold hands and feet, easy bruising, unusual bleeding. Common Lamictal side effects may include: headache, dizziness
- blurred vision, double vision
- tremor, loss of coordination
- dry mouth, nausea, vomiting, stomach pain, diarrhea
- fever, sore throat, runny nose
- drowsiness, tired feeling
- back pain
- or sleep problems (insomnia).

606. This particular disease, Bipolar Disorder, interacts with the drug Latuda and has the following

Drug Side Effects Summary

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. High doses or long-term use of Latuda can cause a serious movement disorder that may not be reversible. Symptoms of this disorder include uncontrollable muscle movements of your lips, tongue, eyes, face, arms, or legs. The longer you use lurasidone, the more likely you are to develop this disorder, especially if you are a woman or an older adult. Call your doctor at once if you have: any new or unusual muscle movements you cannot control
- a light-headed feeling, like you might pass out
- a seizure (convulsions)
- (in women) irregular menstrual periods, breast or vaginal changes, nipple discharge
- (in men) breast swelling, impotence
- trouble swallowing
- manic episodes - racing thoughts, increased energy, decreased need for sleep, risk-taking behavior, being agitated or talkative
- low white blood cell counts - fever, chills, mouth sores, skin sores, sore throat, cough, trouble breathing
- high blood sugar - increased thirst, increased urination, hunger, dry mouth, fruity breath odor
- or severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Common Latuda side effects may include: drowsiness
- weight gain

Drug Side Effects Summary

- tremors, muscle stiffness, slow muscle movement
- feeling restless or being unable to sit still
- nausea, vomiting
- runny nose
- or sleep problems (insomnia).

607. This particular disease, Bipolar Disorder, interacts with the drug Seroquel and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, panic attacks, trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- mask-like appearance of the face, trouble swallowing, problems with speech
- a light-headed feeling, like you might pass out
- blurred vision, tunnel vision, eye pain, or seeing halos around lights
- severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, fainting
- high blood sugar - increased thirst, increased urination, hunger, dry mouth, fruity breath odor, drowsiness, dry skin, blurred vision, weight loss
- or low blood cell counts - sudden weakness or ill feeling, fever, chills, cold or flu symptoms, cough, sore throat, red or swollen gums, painful mouth sores, skin sores, trouble breathing. Common Seroquel side effects may include: speech problems

Drug Side Effects Summary

- dizziness, drowsiness, tiredness
- lack of energy
- fast heartbeats
- stuffy nose
- increased appetite, weight gain
- upset stomach, vomiting, constipation
- dry mouth
- or problems moving.

608. This particular disease, Bipolar Disorder, interacts with the drug Abilify and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tell your doctor right away if you have new or sudden changes in mood or behavior, including new or worse depression or anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, more active or talkative, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: severe agitation , distress, or restless feeling
- twitching or uncontrollable movements of your eyes, lips, tongue, face, arms, or legs
- mask-like appearance of the face, trouble swallowing, problems with speech
- seizure (convulsions)
- severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out
- low blood cell counts - fever, chills, sore throat , weakness, easy bruising, unusual bleeding, purple or red spots under your skin
- or high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor. You may

Drug Side Effects Summary

have increased sexual urges, unusual urges to gamble, or other intense urges while taking this medicine. Talk with your doctor if this occurs. Common Abilify side effects may include: blurred vision

- increased saliva or drooling
- muscle stiffness
- uncontrolled muscle movements, shaking, anxiety, feeling restless
- weight gain
- nausea , vomiting, constipation
- increased or decreased appetite
- headache , dizziness, drowsiness, feeling tired
- sleep problems (insomnia)
- or cold symptoms such as stuffy nose, sneezing, sore throat
-

609. This particular disease, Bipolar Disorder, interacts with the drug topiramate and has the following

- mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), depressed, or have thoughts about suicide or hurting yourself. Topiramate may cause serious side effects. Call your doctor at once if you have: a skin rash, no matter how mild
- vision problems, blurred vision, eye pain or redness, sudden vision loss (can be permanent if not treated quickly)
- confusion, problems with thinking or memory, trouble concentrating, problems with speech
- dehydration symptoms--decreased sweating, high fever, hot and dry skin
- signs of a kidney stone--severe pain in your side or lower back, painful or difficult urination
- signs of too much acid in your blood--irregular heartbeats, feeling tired, loss of appetite, trouble

Drug Side Effects Summary

thinking, feeling short of breath

- or signs of too much ammonia in your blood-- vomiting , unexplained weakness, feeling like you might pass out. Common side effects of topiramate may include: dizziness , drowsiness, tiredness, slow reactions

- problems with speech or memory, abnormal vision

- numbness or tingling in your arms and legs, decreased sensation (especially in the skin)

- changes in your sense of taste, feeling nervous

- nausea , diarrhea, stomach pain, loss of appetite

- fever, weight loss

- or cold symptoms such as stuffy nose, sneezing, sore throat.

610. This particular disease, Bipolar Disorder, interacts with the drug Vraylar and has the following

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: sudden numbness or weakness (especially on one side of the body)

- problems with vision or speech, a light-headed feeling, like you might pass out

- severe distress or agitation

- a seizure

- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)

- trouble swallowing, or accidentally inhaling food or drink

- low white blood cell counts - fever, chills, sore throat , mouth sores, skin sores, sore throat, cough, trouble breathing, feeling light-headed

- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor

Drug Side Effects Summary

- or severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Vraylar can have long lasting effects on your body. Some side effects could occur for several weeks after you stop using this medicine. You may also have new side effects whenever your dose is changed. Common Vraylar side effects may include: involuntary muscle movements

- upset stomach , vomiting
- drowsiness
- or feeling restless.

611. This particular disease, Bipolar Disorder, interacts with the drug lithium and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Too much lithium in your body can cause death. Lithium toxicity can occur if you take only slightly more than a recommended dose. Stop using this medicine and call your doctor right away if you have symptoms of lithium toxicity: muscle weakness, twitching, drowsiness, feeling light-headed, mood changes, blurred vision, ringing in your ears, irregular heartbeats, confusion, slurred speech, clumsiness, trouble breathing, or seizures. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- shortness of breath
- fever, increased thirst or urination
- weakness, dizziness or spinning sensation
- memory problems, hallucinations
- problems with balance or muscle movement
- loss of bowel or bladder control
- a seizure (blackout or convulsions)

Drug Side Effects Summary

- low sodium level - headache, confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady
- thyroid problems - weight gain or loss, muscle pain, tremors, feeling hot or cold, changes in your skin or hair, sweating, feeling weak or tired, fast or pounding heartbeats, increased bowel movements, feeling anxious or agitated
- dehydration symptoms - feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin
- or increased pressure inside the skull -severe headaches, ringing in your ears, dizziness, nausea , vision problems, pain behind your eyes. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Common lithium side effects may include: dizziness, drowsiness
- tremors in your hands
- trouble walking
- dry mouth, increased thirst or urination
- nausea, vomiting, loss of appetite
- rash
- or blurred vision.

612. This particular disease, Bipolar Disorder, interacts with the drug lurasidone and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more

Drug Side Effects Summary

depressed, or have thoughts about suicide or hurting yourself. High doses or long-term use of lurasidone can cause a serious movement disorder that may not be reversible. The longer you use lurasidone, the more likely you are to develop this disorder, especially if you are a woman or an older adult. Lurasidone may cause serious side effects. Call your doctor at once if you have: any new or unusual muscle movements you cannot control

- a light-headed feeling, like you might pass out
- a seizure (convulsions)
- (in women) irregular menstrual periods, breast or vaginal changes, nipple discharge
- (in men) breast swelling, impotence
- trouble swallowing
- manic episodes--racing thoughts, increased energy, decreased need for sleep, risk-taking behavior, being agitated or talkative
- low white blood cell counts--fever, chills, mouth sores, skin sores, sore throat , cough, trouble breathing
- high blood sugar--increased thirst, increased urination, hunger, dry mouth, fruity breath odor
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Common side effects of lurasidone may include: drowsiness
- weight gain
- tremors, muscle stiffness, slow muscle movement
- feeling restless or being unable to sit still
- nausea , vomiting
- runny nose
- or sleep problems (insomnia).

Drug Side Effects Summary

613. This particular disease, Bipolar Disorder, interacts with the drug Abilify Maintena and has the following side effects:

- Abilify Maintena may cause serious side effects, including: See "Important information" above.

Stroke (cerebrovascular problems) in elderly people with dementia-related psychosis that can lead to death. Neuroleptic malignant syndrome (NMS), a serious condition that can lead to death. Call your healthcare provider or go to the nearest emergency room right away if you have some or all of the following symptoms of NMS: high fever confusion changes in pulse, heart rate, and blood pressure stiff muscles sweating Uncontrolled body movements (tardive dyskinesia). This medication may cause movements that you cannot control in your face, tongue, or other body parts. Tardive dyskinesia may not go away, even if you stop receiving Abilify Maintena. Tardive dyskinesia may also start after you stop receiving this medication. Problems with your metabolism such as: high blood sugar (hyperglycemia): Increases in blood sugar can happen in some people who receive this medication. Extremely high blood sugar can lead to coma or death. If you have diabetes or risk factors for diabetes (such as being overweight or a family history of diabetes), your healthcare provider should check your blood sugar before you start receiving this medication and during your treatment. Call your healthcare provider if you have any of these symptoms of high blood sugar while receiving Abilify Maintena: feel very thirsty need to urinate more than usual feel very hungry feel weak or tired feel sick to your stomach feel confused, or your breath smells fruity Increased fat levels (cholesterol and triglycerides) in your blood. Weight gain. You and your healthcare provider should check your weight regularly. Unusual urges. Some people receiving this medication have had unusual urges such as gambling, binge eating or eating that you cannot control (compulsive), compulsive shopping and sexual urges. If you or your family members notice that you are having unusual urges or behaviors, talk to your healthcare provider. Decreased blood pressure (orthostatic hypotension). You may feel lightheaded or faint when you rise too quickly from a sitting or lying position. Low white blood cell count Seizures (convulsions) Problems controlling your body temperature so that you feel too warm. See "What should I avoid while receiving Abilify Maintena?"

Drug Side Effects Summary

Difficulty swallowing The most common side effects of Abilify Maintena include: weight gain inner sense of restlessness such as feeling like you need to move (akathisia) injection site pain sleepiness (sedation) These are not all the possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Abilify Maintena side effects

614. This particular disease, Bipolar Disorder, interacts with the drug cariprazine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Cariprazine may cause serious side effects. Call your doctor at once if you have: sudden numbness or weakness (especially on one side of the body)
- problems with vision or speech, a light-headed feeling, like you might pass out
- severe distress or agitation
- a seizure
- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- trouble swallowing, or accidentally inhaling food or drink
- low white blood cell counts--fever, chills, sore throat , mouth sores, skin sores, sore throat, cough, trouble breathing, feeling light-headed
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Cariprazine can have long lasting effects on your body. Some side effects could occur for several weeks after you stop using this medicine. You may also have new side effects whenever your dose is changed. Common side effects of cariprazine may include: involuntary muscle movements
- upset stomach , vomiting

Drug Side Effects Summary

- drowsiness
- or feeling restless.

615. This particular disease, Bipolar Disorder, interacts with the drug Caplyta and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. People with depression or mental illness may have thoughts about suicide. Some young people may have increased suicidal thoughts when first starting a medicine to treat depression. Stay alert to changes in your mood or symptoms. Your family or caregivers should also watch for sudden changes in your behavior. High doses or long-term use of lumateperone can cause a serious movement disorder that may not be reversible. The longer you use Caplyta, the more likely you are to develop this disorder, especially if you are a woman or an older adult. Call your doctor at once if you have: feeling unsteady, feeling like you might pass out
- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- tightness in your neck or throat, trouble swallowing
- trouble breathing or speaking
- a seizure
- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- low white blood cell counts - fever, chills, mouth sores, skin sores, sore throat , cough, trouble breathing
- or severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Common Caplyta side effects may include: weight gain

Drug Side Effects Summary

- nausea
- dizziness
- drowsiness
- or dry mouth.

616. This particular disease, Bipolar Disorder, interacts with the drug lumateperone and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. People with depression or mental illness may have thoughts about suicide. Some young people may have increased suicidal thoughts when first starting a medicine to treat depression. Stay alert to changes in your mood or symptoms. Your family or caregivers should also watch for sudden changes in your behavior. High doses or long-term use of lumateperone can cause a serious movement disorder that may not be reversible. The longer you use lumateperone, the more likely you are to develop this disorder, especially if you are a woman or an older adult. Lumateperone may cause serious side effects. Call your doctor at once if you have: feeling unsteady, feeling like you might pass out
- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- tightness in your neck or throat, trouble swallowing
- trouble breathing or speaking
- a seizure
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- low white blood cell counts--fever, chills, mouth sores, skin sores, sore throat , cough, trouble breathing
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast

Drug Side Effects Summary

or uneven heartbeats, tremors, feeling like you might pass out. Common side effects of lumateperone may include: weight gain

- nausea
- dizziness
- drowsiness
- or dry mouth.

617. This particular disease, Bipolar Disorder, interacts with the drug Lybalvi and has the following

- Lybalvi may cause serious side effects, including: See Important information Stroke (cerebrovascular problems) in elderly people with dementia-related psychosis that can lead to death. Opioid withdrawal. Do not take Lybalvi for at least 7 days after you stopped taking short-acting opioids and for at least 14 days after you stopped taking long-acting opioids. One of the medicines in Lybalvi (samidorphan) can cause opioid withdrawal that may be severe and cause hospitalization in people who are physically dependent on opioids. Talk to your healthcare provider if you have questions about the type of opioid you take. Risk of life-threatening opioid overdose. You should not start taking opioids for at least 5 days after you stop treatment with Lybalvi. One of the medicines in Lybalvi (samidorphan) can increase your chance of having an opioid overdose that can cause death if you take opioids during treatment or within 5 days after stopping treatment with Lybalvi. You can accidentally overdose in 2 ways: Lybalvi blocks the effects of opioids, such as heroin, methadone, or opioid pain medicines. Do not take large amounts of opioids to try to overcome the opioid-blocking effects of Lybalvi. This can lead to serious injury, coma, or death. After you take Lybalvi, its blocking effect slowly decreases and completely goes away over time. You may be more sensitive to the effects of opioids. If you have used opioid street drugs or opioid-containing medicines in the past, using opioids in amounts that you used before treatment with Lybalvi can lead to overdose or death. It is important that you tell your family and the people closest to you of this increased sensitivity to

Drug Side Effects Summary

opioids and the risk of overdose. You or someone close to you should get emergency medical help right away if you: have trouble breathing become very drowsy with slowed breathing have slow, shallow breathing (little chest movement with breathing) feel faint, very dizzy, confused, or have unusual symptoms Tell your healthcare provider if you are taking Lybalvi before a medical procedure or surgery. Neuroleptic malignant syndrome (NMS), a serious condition that can lead to death. Call your healthcare provider or go to the nearest hospital emergency room right away if you have some or all of the following signs and symptoms of NMS: high fever stiff muscles confusion sweating changes in your breathing, pulse, heart rate, and blood pressure Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS): One of the medicines in Lybalvi (olanzapine) can cause DRESS which can cause death. Tell your healthcare provider right away if you develop any of the following symptoms of DRESS, including: rash swollen glands liver problems heart problems fever kidney problems lung problems Problems with your metabolism such as: high blood sugar (hyperglycemia) and diabetes. Increases in blood sugar can happen in some people who take Lybalvi. Extremely high blood sugar can lead to coma or death. Your healthcare provider should check your blood sugar before you start and regularly during treatment. Call your healthcare provider if you have any of these symptoms of high blood sugar during treatment: feel very thirsty feel very hungry feel sick to your stomach need to urinate more than usual feel weak or tired feel confused, or your breath smells fruity increased fat levels (cholesterol and triglycerides) in your blood . Your healthcare provider should check the fat levels in your blood before you start and regularly during treatment. weight gain. You and your healthcare provider should check your weight before you start and often during treatment. Uncontrolled body movements (tardive dyskinesia). Lybalvi may cause movements that you cannot control in your face, tongue, or other body parts. Tardive dyskinesia may not go away, even if you stop taking Lybalvi. Tardive dyskinesia may also start after you stop taking Lybalvi. Decreased blood pressure (orthostatic hypotension) and fainting. You may feel lightheaded or faint when you rise too quickly from a sitting or lying position. Falls.

Drug Side Effects Summary

Lybalvi may make you sleepy or dizzy, may cause a decrease in your blood pressure when changing position (orthostatic hypotension), and can slow your thinking and motor skills which may lead to falls that can cause fractures or other injuries. Low white blood cell count. Your healthcare provider may do blood tests during the first few months of treatment. Difficulty swallowing that can cause food or liquid to get into your lungs. Seizures (convulsions). Problems controlling your body temperature so that you feel too warm. See What should I avoid while taking Lybalvi? Increased prolactin levels in your blood. Your healthcare provider may do blood tests to check your prolactin levels during treatment. The most common side effects when used to treat people with schizophrenia include: weight gain dry mouth sleepiness headache The most common side effects when used alone to treat people with mixed or manic episodes that happen with bipolar I disorder include: weakness constipation sleepiness shaking dry mouth increased appetite dizziness The most common side effects when used in combination with lithium or valproate to treat people with mixed or manic episodes that happen with bipolar I disorder include: dry mouth increased appetite back pain problems speaking memory problems weight gain dizziness constipation mouth watering numbness and tingling in your arm and legs These are not all the possible side effects. Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA1088. Lybalvi side effects

618. This particular disease, Bipolar Disorder, interacts with the drug quetiapine and has the following

- Serious side effects of quetiapine include: See "Important information" above stroke that can lead to death can happen in elderly people with dementia who take medicines like this neuroleptic malignant syndrome (NMS). NMS is a rare but very serious condition that can happen in people who take antipsychotic medicines, including quetiapine. NMS can cause death and must be treated in a hospital. Call your healthcare provider right away if you become severely ill and have some or all of these symptoms: high fever excessive sweating rigid muscles confusion changes in

Drug Side Effects Summary

your breathing, heartbeat, and blood pressure falls can happen in some people who take this medication. These falls may cause serious injuries. high blood sugar (hyperglycemia). High blood sugar can happen if you have diabetes already or if you have never had diabetes. High blood sugar could lead to: build-up of acid in your blood due to ketones (ketoacidosis) coma death Increases in blood sugar can happen in some people who take quetiapine. Extremely high blood sugar can lead to coma or death. If you have diabetes or risk factors for diabetes (such as being overweight or a family history of diabetes) your healthcare provider should check your blood sugar before you start this medication and during therapy. Call your healthcare provider if you have any of these symptoms of high blood sugar (hyperglycemia): feel very thirsty need to urinate more than usual feel very hungry feel weak or tired feel sick to your stomach feel confused, or your breath smells fruity high fat levels in your blood (increased cholesterol and triglycerides). High fat levels may happen in people treated with this medication. You may not have any symptoms, so your healthcare provider may decide to check your cholesterol and triglycerides during your treatment. increase in weight (weight gain). Weight gain is common in people who take quetiapine so you and your healthcare provider should check your weight regularly. Talk to your healthcare provider about ways to control weight gain, such as eating a healthy, balanced diet, and exercising. movements you cannot control in your face, tongue, or other body parts (tardive dyskinesia). These may be signs of a serious condition. Tardive dyskinesia may not go away, even if you stop taking quetiapine. Tardive dyskinesia may also start after you stop taking this medication. decreased blood pressure (orthostatic hypotension), including lightheadedness or fainting caused by a sudden change in heart rate and blood pressure when rising too quickly from a sitting or lying position. increases in blood pressure in children and teenagers. Your healthcare provider should check blood pressure in children and adolescents before starting this medication and during therapy. low white blood cell count. Tell your healthcare provider as soon as possible if you have a fever, flu-like symptoms, or any other infection, as this could be a result of a very low white blood cell count. Your healthcare provider may check your

Drug Side Effects Summary

white blood cell level to determine if further treatment or other action is needed. cataracts seizures abnormal thyroid tests. Your healthcare provider may do blood tests to check your thyroid hormone level. increases in prolactin levels sleepiness, drowsiness, feeling tired, difficulty thinking and doing normal activities increased body temperature difficulty swallowing trouble sleeping or trouble staying asleep (insomnia), nausea, or vomiting if you suddenly stop taking quetiapine. These symptoms usually get better 1 week after you start having them. The most common side effects of quetiapine tablets in adults include: drowsiness sudden drop in blood pressure upon standing weight gain sluggishness abnormal liver tests upset stomach dry mouth dizziness weakness abdominal pain constipation sore throat The most common side effects of quetiapine tablets in children and adolescents include: drowsiness dizziness fatigue nausea dry mouth weight gain increased appetite vomiting rapid heart beat The most common side effects of quetiapine extended-release (XR) tablets include: dry mouth constipation dizziness increased appetite upset stomach fatigue stuffy nose difficulty moving disturbance in speech or language The most common side effects of quetiapine extended-release (XR) tablets in children and adolescents include: drowsiness dizziness fatigue stuffy nose increased appetite upset stomach vomiting dry mouth tachycardia weight increased These are not all the possible side effects. For more information, ask your healthcare provider or pharmacist. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Quetiapine side effects

619. This particular disease, Bipolar Disorder, interacts with the drug Depakote and has the following

- skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Call your doctor at once if the person taking this medicine has signs of liver or pancreas problems, such as: loss of appetite, upper stomach pain (that may spread to your back), ongoing nausea or vomiting, dark urine, swelling in the face, or jaundice (yellowing of the skin or eyes). Report any new or worsening symptoms to your doctor, such as: mood or behavior changes,

Drug Side Effects Summary

depression, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself. Call your doctor at once if you have any of these other side effects: easy bruising, unusual bleeding (nose, mouth, or gums), purple or red pinpoint spots under your skin

- fever, swollen glands, mouth sores
- confusion, tiredness, cold feeling, vomiting, change in your mental state
- severe drowsiness
- or worsening seizures. Common Depakote side effects may include: nausea, vomiting, stomach pain, diarrhea , constipation
- headache , back pain
- dizziness , drowsiness, weakness, tremors
- memory problems, mood changes, trouble sleeping
- bruising or bleeding
- runny nose , sore throat, cough, wheezing, trouble breathing
- fever, flu symptoms
- problems with walking or coordination
- swelling in your hands or feet
- blurred vision, double vision, unusual eye movements
- ringing in your ears
- rash, hair loss
- or changes in weight or appetite.

620. This particular disease, Bipolar Disorder, interacts with the drug Seroquel XR and has the following

- mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed,

Drug Side Effects Summary

or have thoughts about suicide or hurting yourself. Seroquel XR may cause serious side effects. Call your doctor at once if you have: uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)

- mask-like appearance of the face, trouble swallowing, problems with speech
- a light-headed feeling, like you might pass out
- severe constipation
- painful or difficult urination
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, fainting
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or low white blood cell counts--fever, chills, mouth sores, skin sores, sore throat, cough, trouble breathing, feeling light-headed. Common side effects of Seroquel XR may include: speech problems
- dizziness, drowsiness, tiredness
- lack of energy
- fast heartbeats
- stuffy nose
- increased appetite, weight gain
- upset stomach , vomiting , constipation
- dry mouth
- or problems moving.

621. This particular disease, Bipolar Disorder, interacts with the drug divalproex sodium and has the

- skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Call your doctor at once if the person taking this medicine has signs of liver or

Drug Side Effects Summary

pancreas problems, such as: loss of appetite, upper stomach pain (that may spread to your back), ongoing nausea or vomiting, dark urine, swelling in the face, or jaundice (yellowing of the skin or eyes). Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, depression, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself. Call your doctor at once if you have any of these other side effects: easy bruising, unusual bleeding (nose, mouth, or gums), purple or red pinpoint spots under your skin

- fever, swollen glands, mouth sores
- confusion, tiredness, cold feeling, vomiting, change in your mental state
- severe drowsiness
- or worsening seizures. Common side effects of divalproex sodium may include: nausea, vomiting, stomach pain, diarrhea , constipation
- headache , back pain
- dizziness , drowsiness, weakness, tremors
- memory problems, mood changes, trouble sleeping
- bruising or bleeding
- runny nose , sore throat, cough, wheezing, trouble breathing
- fever, flu symptoms
- problems with walking or coordination
- swelling in your hands or feet
- blurred vision, double vision, unusual eye movements
- ringing in your ears
- rash, hair loss
- or changes in weight or appetite.

Drug Side Effects Summary

622. This particular disease, Bipolar Disorder, interacts with the drug Risperdal and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- breast swelling or tenderness (in men or women), nipple discharge , impotence , lack of interest in sex, missed menstrual periods
- severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out
- low white blood cells - sudden weakness or ill feeling, fever, chills, sore throat , mouth sores, red or swollen gums, trouble swallowing, skin sores, cold or flu symptoms, cough, trouble breathing
- low levels of platelets in your blood - easy bruising, unusual bleeding (nose, mouth, vagina, or rectum), purple or red pinpoint spots under your skin
- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- or penis erection that is painful or lasts 4 hours or longer. Common Risperdal side effects may include: headache
- dizziness, drowsiness, feeling tired
- tremors, twitching or uncontrollable muscle movements
- agitation , anxiety , restless feeling
- depressed mood
- dry mouth, upset stomach , diarrhea , constipation
- weight gain
- or cold symptoms such as stuffy nose, sneezing, sore throat.

Drug Side Effects Summary

623. This particular disease, Bipolar Disorder, interacts with the drug Geodon and has the following

- hives, difficult breathing, swelling in your face or throat) or a severe skin reaction (fever, sore throat, burning in your eyes, skin pain, red or purple skin rash that spreads and causes blistering and peeling). Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. This reaction may occur several weeks after you began using ziprasidone. Also call your doctor at once if you have: dizziness or a light-headed feeling, like you might pass out
- chest pain, fast or pounding heartbeats
- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- sudden weakness or ill feeling, fever, chills, sore throat, mouth sores, red or swollen gums, pain when swallowing
- increased thirst, increased urination, hunger, dry mouth, fruity breath odor, drowsiness, dry skin, blurred vision, weight loss
- or very stiff (rigid) muscles, high fever, sweating, confusion, agitation. Common Geodon side effects may include: dizziness, drowsiness, weakness
- headache
- nausea, vomiting
- trouble swallowing
- weight gain
- feeling restless or being unable to sit still
- tremors, involuntary muscle movements
- vision problems
- or runny nose, new or worsening cough.

Drug Side Effects Summary

624. This particular disease, Bipolar Disorder, interacts with the drug sertraline and has the following

- skin rash or hives (with or without fever or joint pain)
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: unusual bleeding or bruising
- a seizure
- vision changes, eye pain , redness, or swelling
- low blood sodium - headache , confusion, problems with thinking or memory, weakness, feeling unsteady
- or manic episodes - racing thoughts, increased energy, unusual risk-taking behavior, extreme happiness, being irritable or talkative. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting , or diarrhea . Sertraline can affect growth in children. Your child's height and weight may be checked often. Common sertraline side effects may include: indigestion , nausea, diarrhea, loss of appetite, loose stool
- increased sweating
- tremors
- or sexual problems.

625. This particular disease, Bipolar Disorder, interacts with the drug Zyprexa and has the following

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash , fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. High doses or long-term use of olanzapine can cause a serious movement disorder that may not be reversible. The longer you use Zyprexa, the more likely you are to develop this disorder, especially if you are a woman or an older adult. Call your doctor at once if you have: uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- trouble speaking or swallowing
- swelling in your hands or feet
- confusion, unusual thoughts or behavior, hallucinations, or thoughts about hurting yourself
- low white blood cell counts - fever, chills, mouth sores, skin sores, sore throat , cough, trouble breathing, feeling light-headed
- or signs of dehydration - feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin
- liver problems - upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- high blood sugar - increased thirst, increased urination, hunger, dry mouth, fruity breath odor, drowsiness, dry skin, blurred vision, weight loss
- or severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Common Zyprexa side effects may include: weight gain (more likely in teenagers), increased appetite
- headache , dizziness, drowsiness, feeling tired or restless
- problems with speech or memory
- tremors or shaking, numbness or tingly feeling

Drug Side Effects Summary

- changes in personality
- dry mouth, or increased salivation
- stomach pain, constipation
- or pain in your arms or legs.

626. This particular disease, Bipolar Disorder, interacts with the drug aripiprazole and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tell your doctor right away if you have new or sudden changes in mood or behavior, including new or worse depression or anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, more active or talkative, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: severe agitation , distress, or restless feeling
- twitching or uncontrollable movements of your eyes, lips, tongue, face, arms, or legs
- mask-like appearance of the face, trouble swallowing, problems with speech
- seizure (convulsions)
- severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out
- low blood cell counts - fever, chills, sore throat , weakness, easy bruising, unusual bleeding, purple or red spots under your skin
- or high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor. You may have increased sexual urges, unusual urges to gamble, or other intense urges while taking this medicine. Talk with your doctor if this occurs. Common aripiprazole side effects may include: blurred vision
- increased saliva or drooling

Drug Side Effects Summary

- muscle stiffness
- uncontrolled muscle movements, shaking, anxiety, feeling restless
- weight gain
- nausea , vomiting, constipation
- increased or decreased appetite
- headache , dizziness, drowsiness, feeling tired
- sleep problems (insomnia)
- or cold symptoms such as stuffy nose, sneezing, sore throat
-

627. This particular disease, Bipolar Disorder, interacts with the drug Depakote ER and has the following side effects:

- skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Call your doctor at once if the person taking this medicine has signs of liver or pancreas problems, such as: loss of appetite, upper stomach pain (that may spread to your back), ongoing nausea or vomiting, dark urine, swelling in the face, or jaundice (yellowing of the skin or eyes). Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, depression, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself. Call your doctor at once if you have any of these other side effects: easy bruising, unusual bleeding (nose, mouth, or gums), purple or red pinpoint spots under your skin
- fever, swollen glands, mouth sores
- confusion, tiredness, cold feeling, vomiting, change in your mental state
- severe drowsiness
- or worsening seizures. Common side effects of Depakote ER may include: nausea, vomiting, stomach pain, diarrhea , constipation

Drug Side Effects Summary

- headache , back pain
- dizziness , drowsiness, weakness, tremors
- memory problems, mood changes, trouble sleeping
- bruising or bleeding
- runny nose , sore throat, cough, wheezing, trouble breathing
- fever, flu symptoms
- problems with walking or coordination
- swelling in your hands or feet
- blurred vision, double vision, unusual eye movements
- ringing in your ears
- rash, hair loss
- or changes in weight or appetite.

628. This particular disease, Bipolar Disorder, interacts with the drug olanzapine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash , fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. High doses or long-term use of olanzapine can cause a serious movement disorder that may not be reversible. The longer you use olanzapine, the more likely you are to develop this disorder, especially if you are a woman or an older adult. Olanzapine may cause serious side effects. Call your doctor at once if you have: uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- trouble speaking or swallowing

Drug Side Effects Summary

- swelling in your hands or feet
- confusion, unusual thoughts or behavior, hallucinations, or thoughts about hurting yourself
- low white blood cell counts--fever, chills, mouth sores, skin sores, sore throat , cough, trouble breathing, feeling light-headed
- or signs of dehydration --feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin
- liver problems--upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- high blood sugar--increased thirst, increased urination, hunger, dry mouth, fruity breath odor, drowsiness, dry skin, blurred vision, weight loss
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Common side effects of olanzapine may include: weight gain (more likely in teenagers), increased appetite
- headache , dizziness, drowsiness, feeling tired or restless
- problems with speech or memory
- tremors or shaking, numbness or tingly feeling
- changes in personality
- dry mouth, or increased salivation
- stomach pain, constipation
- or pain in your arms or legs.

629. This particular disease, Bipolar Disorder, interacts with the drug Saphris and has the following

- hives
- fast heartbeats, feeling light-headed
- wheezing, difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. High doses or long-term use of Saphris can cause a serious movement disorder that may not be reversible. The longer you use Saphris, the more likely you are to develop this disorder, especially if you are a woman or an older adult. Call your doctor at once if you have: uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- ulcers, blisters, swelling, or peeling in your mouth
- a light-headed feeling, like you might pass out
- breast pain or swelling, nipple discharge
- low white blood cell counts - fever, chills, mouth sores, skin sores, sore throat , cough, trouble breathing
- or severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Common Saphris side effects may include: dizziness , drowsiness, tiredness
- feeling restless or agitated
- numbness or tingling in or around your mouth
- muscle stiffness, jerky muscle movements
- nausea , altered sense of taste
- or increased appetite, weight gain.

630. This particular disease, Bipolar Disorder, interacts with the drug citalopram and has the following

- hives , rash, blisters
- fever, joint pain
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tell your doctor right away if you have new or sudden changes in mood or behavior, including new or worse depression or anxiety , panic attacks , trouble

Drug Side Effects Summary

sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, more active or talkative, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: a light-headed feeling, like you might pass out

- blurred vision, eye pain or redness, seeing halos around lights
- fast or pounding heartbeats, pain or fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- a seizure
- manic episodes - racing thoughts, increased energy, decreased need for sleep, risk-taking behavior, being agitated or talkative
- severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors
- or low blood sodium - headache , confusion, problems with thinking or memory, weakness, feeling unsteady. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting , or diarrhea . Common citalopram side effects may include: sexual problems
- dizziness, drowsiness
- dry mouth, thirst, increased sweating or urination
- loss of appetite, nausea, diarrhea, constipation
- feeling anxious, agitated, or shaky
- feeling weak or tired
- sleep problems (insomnia)
- yawning
- increased muscle movement
- nosebleeds , heavy menstrual bleeding

Drug Side Effects Summary

- or cold symptoms such as stuffy nose, sneezing, sore throat .

631. This particular disease, Bipolar Disorder, interacts with the drug carbamazepine and has the following side effects:

- skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Report any new or worsening symptoms to your doctor, such as: sudden mood or behavior changes, depression, anxiety , insomnia , or if you feel agitated, hostile, restless, irritable, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: a skin rash, no matter how mild
- loss of appetite, right-sided upper stomach pain, dark urine
- slow, fast, or pounding heartbeats
- anemia or other blood problems - fever, chills, sore throat, mouth sores, bleeding gums, nosebleeds , pale skin, easy bruising, unusual tiredness, feeling light-headed or short of breath
- or low levels of sodium in the body - headache , confusion, severe weakness, feeling unsteady, increased seizures. Common carbamazepine side effects may include: dizziness , loss of coordination, problems with walking
- nausea, vomiting
- or drowsiness.

632. This particular disease, Bipolar Disorder, interacts with the drug ziprasidone and has the following side effects:

- skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Ziprasidone may cause serious side effects. Stop using ziprasidone and call your doctor at once if you have: fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)

Drug Side Effects Summary

- any skin rash, no matter how mild
- low white blood cell counts--fever, mouth sores, skin sores, sore throat, cough, trouble breathing
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, agitation. Common side effects of ziprasidone may include: dizziness, drowsiness, weakness
- headache
- nausea, vomiting
- trouble swallowing
- weight gain
- feeling restless or being unable to sit still
- tremors, involuntary muscle movements
- vision problems
- or runny nose , new or worsening cough.

633. This particular disease, Bipolar Disorder, interacts with the drug Symbyax and has the following

- skin rash or hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Seek medical treatment if you have symptoms of a serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash, fever, swollen glands, flu-like symptoms, unusual bruising, or jaundice (yellowing of your skin or eyes). High doses or long-term use of Symbyax can cause a serious movement disorder that may not be reversible, especially in women and older adults. Tell your doctor right away if you have uncontrollable muscle movements of your lips, tongue, eyes, face, arms, or legs. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive

Drug Side Effects Summary

(mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: unusual bleeding or bruising

- vision changes
- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- dehydration symptoms - feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin
- severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out
- high blood sugar - increased thirst, increased urination, hunger, fruity breath odor, confusion, upset stomach
- low sodium level - headache , confusion, slurred speech, severe weakness, vomiting , loss of coordination, feeling unsteady
- or low white blood cell counts - fever, chills, mouth sores, skin sores, sore throat , cough, trouble breathing. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea . Common Symbyax side effects may include: drowsiness, blurred vision, dry mouth
- increased appetite, weight gain
- trouble concentrating, feeling tired
- swelling in your hands or feet
- tremor
- or abnormal liver function or cholesterol tests.

634. This particular disease, Bipolar Disorder, interacts with the drug valproic acid and has the follow

Drug Side Effects Summary

- skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Call your doctor at once if the person taking this medicine has signs of liver or pancreas problems, such as: loss of appetite, upper stomach pain (that may spread to your back), ongoing nausea or vomiting, dark urine, swelling in the face, or jaundice (yellowing of the skin or eyes). Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, depression, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself. Call your doctor at once if you have any of these other side effects: confusion, tiredness, cold feeling, vomiting, change in your mental state

- easy bruising, unusual bleeding (nose, mouth, or gums), purple or red pinpoint spots under your skin

- severe drowsiness

- or worsening seizures. Severe drowsiness may be more likely in older adults. Common side effects of valproic acid may include: nausea, vomiting, stomach pain, diarrhea

- dizziness , drowsiness, weakness

- headache

- tremors, problems with walking or coordination

- blurred vision, double vision

- hair loss

- or changes in appetite, weight gain.

635. This particular disease, Bipolar Disorder, interacts with the drug fluoxetine / olanzapine and has

- skin rash or hives

- difficulty breathing

- swelling of your face, lips, tongue, or throat. Seek medical treatment if you have symptoms of a

Drug Side Effects Summary

serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash, fever, swollen glands, flu-like symptoms, unusual bruising, or jaundice (yellowing of your skin or eyes). High doses or long-term use of fluoxetine and olanzapine can cause a serious movement disorder that may not be reversible, especially in women and older adults. Tell your doctor right away if you have uncontrollable muscle movements of your lips, tongue, eyes, face, arms, or legs. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Fluoxetine and olanzapine may cause serious side effects. Call your doctor at once if you have: unusual bleeding or bruising

- vision changes
- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- dehydration symptoms--feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin
- severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out
- high blood sugar--increased thirst, increased urination, hunger, fruity breath odor, confusion, upset stomach
- low sodium level -- headache , confusion, slurred speech, severe weakness, vomiting , loss of coordination, feeling unsteady
- or low white blood cell counts--fever, chills, mouth sores, skin sores, sore throat , cough, trouble breathing. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea . Common side effects of fluoxetine and olanzapine

Drug Side Effects Summary

may include: drowsiness, blurred vision, dry mouth

- increased appetite, weight gain
- trouble concentrating, feeling tired
- swelling in your hands or feet
- tremor
- or abnormal liver function or cholesterol tests.

636. This particular disease, Bipolar Disorder, interacts with the drug Lamictal ODT and has the following side effects:

- mood or behavior changes, depression, anxiety , or if you feel agitated, hostile, restless, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself. Lamictal ODT may cause serious side effects. Call your doctor at once if you have: fast, slow, or pounding heartbeats or fluttering in your chest

- chest pain, shortness of breath
- fever, swollen glands, weakness, severe muscle pain
- any skin rash, especially with blistering or peeling
- painful sores in your mouth or around your eyes
- headache, neck stiffness, increased sensitivity to light, nausea , vomiting, confusion, drowsiness
- jaundice (yellowing of the skin or eyes)
- or pale skin, cold hands and feet, easy bruising, unusual bleeding. Common side effects of Lamictal ODT may include: headache, dizziness
- blurred vision, double vision
- tremor, loss of coordination
- dry mouth, nausea, vomiting, stomach pain, diarrhea
- fever, sore throat, runny nose
- drowsiness, tired feeling

Drug Side Effects Summary

- back pain
- or sleep problems (insomnia).

637. This particular disease, Bipolar Disorder, interacts with the drug levetiracetam and has the following

- mood or behavior changes, depression, anxiety , panic attacks , trouble sleeping, or if you feel agitated, hostile, irritable, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself. Levetiracetam may cause serious side effects. Call your doctor at once if you have: unusual changes in mood or behavior (unusual risk-taking behavior, being irritable or talkative)
- confusion, hallucinations, extreme drowsiness, feeling very weak or tired
- loss of balance or coordination, problems with walking or movement
- a skin rash, no matter how mild
- easy bruising, unusual bleeding
- or fever, chills, weakness, or other signs of infection. Common side effects of levetiracetam may include: dizziness, drowsiness, tiredness, weakness
- feeling aggressive or irritable
- loss of appetite
- stuffy nose
- or infection.

638. This particular disease, Bipolar Disorder, interacts with the drug Lithobid and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Too much Lithobid in your body can cause death. This medicine toxicity can occur if you take only slightly more than a recommended dose. Stop using Lithobid and call your doctor right away if you have symptoms of lithium toxicity: muscle weakness,

Drug Side Effects Summary

twitching, drowsiness, feeling light-headed, mood changes, blurred vision, ringing in your ears, irregular heartbeats, confusion, slurred speech, clumsiness, trouble breathing, or seizures. Lithobid may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out

- shortness of breath
- fever, increased thirst or urination
- weakness, dizziness or spinning sensation
- memory problems, hallucinations
- problems with balance or muscle movement
- loss of bowel or bladder control
- a seizure (blackout or convulsions)
- low sodium level -- headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady
- thyroid problems--weight gain or loss, muscle pain, tremors, feeling hot or cold, changes in your skin or hair, sweating, feeling weak or tired, fast or pounding heartbeats, increased bowel movements, feeling anxious or agitated
- dehydration symptoms--feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin
- or increased pressure inside the skull--severe headaches, ringing in your ears, dizziness, nausea , vision problems, pain behind your eyes. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Common side effects of Lithobid may include: dizziness, drowsiness
- tremors in your hands
- trouble walking

Drug Side Effects Summary

- dry mouth, increased thirst or urination
- nausea, vomiting, loss of appetite
- rash
- or blurred vision.

639. This particular disease, Bipolar Disorder, interacts with the drug Zyprexa Zydis and has the fol

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash , fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. High doses or long-term use of olanzapine can cause a serious movement disorder that may not be reversible. The longer you use Zyprexa Zydis, the more likely you are to develop this disorder, especially if you are a woman or an older adult. Zyprexa Zydis may cause serious side effects. Call your doctor at once if you have: uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- trouble speaking or swallowing
- swelling in your hands or feet
- confusion, unusual thoughts or behavior, hallucinations, or thoughts about hurting yourself
- low white blood cell counts--fever, chills, mouth sores, skin sores, sore throat , cough, trouble breathing, feeling light-headed
- or signs of dehydration --feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin
- liver problems--upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)

Drug Side Effects Summary

- high blood sugar--increased thirst, increased urination, hunger, dry mouth, fruity breath odor, drowsiness, dry skin, blurred vision, weight loss
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Common side effects of Zyprexa Zydis may include: weight gain (more likely in teenagers), increased appetite
- headache , dizziness, drowsiness, feeling tired or restless
- problems with speech or memory
- tremors or shaking, numbness or tingly feeling
- changes in personality
- dry mouth, or increased salivation
- stomach pain, constipation
- or pain in your arms or legs.

640. This particular disease, Bipolar Disorder, interacts with the drug asenapine and has the following

- hives
- fast heartbeats, feeling light-headed
- wheezing, difficult breathing
- swelling of your face, lips, tongue, or throat. High doses or long-term use of asenapine can cause a serious movement disorder that may not be reversible. The longer you use asenapine, the more likely you are to develop this disorder, especially if you are a woman or an older adult. Asenapine may cause serious side effects. Call your doctor at once if you have: uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- ulcers, blisters, swelling, or peeling in your mouth
- a light-headed feeling, like you might pass out

Drug Side Effects Summary

- breast pain or swelling, nipple discharge
- low white blood cell counts--fever, chills, mouth sores, skin sores, sore throat , cough, trouble breathing
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Common side effects of asenapine may include: dizziness , drowsiness, tiredness
- feeling restless or agitated
- numbness or tingling in or around your mouth
- muscle stiffness, jerky muscle movements
- nausea , altered sense of taste
- or increased appetite, weight gain.

641. This particular disease, Bipolar Disorder, interacts with the drug Equetro and has the following

- skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Report any new or worsening symptoms to your doctor, such as: sudden mood or behavior changes, depression, anxiety , insomnia , or if you feel agitated, hostile, restless, irritable, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: a skin rash, no matter how mild
- loss of appetite, right-sided upper stomach pain, dark urine
- slow, fast, or pounding heartbeats
- anemia or other blood problems - fever, chills, sore throat, mouth sores, bleeding gums, nosebleeds , pale skin, easy bruising, unusual tiredness, feeling light-headed or short of breath
- or low levels of sodium in the body - headache , confusion, severe weakness, feeling unsteady, increased seizures. Common Equetro side effects may include: dizziness , loss of coordination, problems with walking

Drug Side Effects Summary

- nausea, vomiting
- or drowsiness.

642. This particular disease, Bipolar Disorder, interacts with the drug iloperidone and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. High doses or long-term use of iloperidone can cause a serious movement disorder that may not be reversible. The longer you use iloperidone, the more likely you are to develop this disorder, especially if you are an older adult. Iloperidone may cause serious side effects. Call your doctor at once if you have: fast or pounding heartbeats, fluttering in your chest
- a light-headed feeling, like you might pass out
- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- tremors, slow muscle movement, muscles pain or stiffness
- confusion, agitation , thoughts of hurting yourself
- loss of bladder control
- penis erection that is painful or lasts 4 hours or longer
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- low white blood cell counts--fever, chills, sore throat , mouth sores, skin sores, cough, trouble breathing
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out
- Older adults may be more likely to have side effects from this medicine. Common side effects of iloperidone may include: weight gain

Drug Side Effects Summary

- dizziness, drowsiness, tired feeling
- dry mouth, stuffy nose
- or fast heart rate .

643. This particular disease, Bipolar Disorder, interacts with the drug Lamictal CD and has the follo

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Blurred vision changes in vision chest pain, discomfort, or tightness clumsiness or unsteadiness decreased urine output dilated neck veins double vision irregular breathing irregular heartbeat nausea pain or discomfort in the arms, jaw, back, or neck poor coordination seizure that will not stop skin rash sweating swelling of the face, fingers, feet, or lower legs trouble breathing unusual tiredness or weakness vomiting weight gain Less common Anxiety chest pain confusion continuous, uncontrolled back and forth or rolling eye movements depression increase in seizures infection irritability Rare Blistering, peeling, or loosening of the skin chills cough dark urine diarrhea fever general feeling of discomfort or illness headache itching joint pain loss of appetite memory loss muscle cramps, pain, or weakness red or irritated eyes runny nose shivering small red or purple spots on the skin sore throat sores, ulcers, or white spots on the lips or in the mouth swelling of the face, mouth, hands, or feet swollen lymph nodes trouble sleeping unusual bleeding or bruising yellow eyes or skin Incidence not known Back, leg, or stomach pains bleeding gums bloating blood in the urine bloody, black or tarry stools bluish lips or skin constipation cough coughing or vomiting blood difficulty with swallowing fainting fast heartbeat general body swelling heartburn high fever hoarseness lightheadedness loss of balance control lower back or side pain mask-like face muscle spasms nosebleeds not breathing pain or burning in the throat painful or difficult urination pains in the stomach or side, possibly radiating to the back pale skin persistent bleeding or oozing from puncture sites, mouth, or nose rapid, shallow

Drug Side Effects Summary

breathing redness, soreness, or itching skin shuffling walk slowed movement slurred speech sores, welting, or blisters stiffness of the arms and legs swollen or painful glands tic-like (jerky) movements Symptoms of overdose Clumsiness or unsteadiness (severe) continuous, uncontrolled back and forth or rolling eye movements (severe) dizziness (severe) drowsiness (severe) dryness of the mouth (severe) headache (severe) increased heart rate loss of consciousness slurred speech (severe) Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Dizziness drowsiness Less common Indigestion loss of strength menstrual pain pain trembling or shaking trouble with sleeping unusual weight loss Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Lamictal CD side effects

644. This particular disease, Bipolar Disorder, interacts with the drug paliperidone and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop taking paliperidone and call your doctor at once if you have any of these signs of a serious movement disorder: tremors or shaking in your arms or legs
- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- or any new or unusual muscle movements you cannot control. Paliperidone may cause serious side effects. Call your doctor at once if you have: fast or pounding heartbeats, fluttering in your

Drug Side Effects Summary

chest, shortness of breath, and sudden dizziness (like you might pass out)

- breast swelling (in women or men), nipple discharge
- changes in menstrual periods
- impotence , penis erection that is painful or lasts 4 hours or longer
- weight gain
- low white blood cell counts--fever, chills, mouth sores, skin sores, sore throat , cough, trouble breathing
- high blood sugar--increased thirst, increased urination, hunger, fruity breath odor
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, fast or pounding heartbeats, fainting. Common side effects of paliperidone may include: drowsiness
- anxiety
- muscle stiffness, tremors or shaking
- uncontrolled muscle movements, trouble with walking, balance, or speech
- weight gain
- upset stomach , constipation
- fast heart rate
- or stuffy nose, sore throat.

645. This particular disease, Bipolar Disorder, interacts with the drug perphenazine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. High doses or long-term use of perphenazine can cause a serious movement disorder that may not be reversible. The longer you use perphenazine, the more likely you are to develop this disorder, especially if you are a woman or an older adult. Perphenazine may cause serious side effects. Call your doctor at once if you have: uncontrolled

Drug Side Effects Summary

muscle movements in your arms or legs, or your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)

- worsening symptoms of schizophrenia
- confusion, paranoia, feeling restless or excited
- seizure (convulsions)
- a light-headed feeling, like you might pass out
- jaundice (yellowing of your skin or eyes)
- little or no urinating
- slow heart rate , weak pulse, weak or shallow breathing
- low white blood cell counts--fever, chills, mouth sores, skin sores, sore throat , cough, trouble breathing, feeling light-headed
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, fast or uneven heartbeats. Side effects such as dry mouth, constipation , tremors, and drowsiness may be more likely in older adults. Common side effects of perphenazine may include: mild dizziness or drowsiness
- blurred vision, headache
- sleep problems (insomnia), strange dreams
- loss of appetite, vomiting, diarrhea , constipation
- increased sweating or urination
- dry mouth or stuffy nose
- breast swelling or discharge
- or mild itching or skin rash .

646. This particular disease, Bipolar Disorder, interacts with the drug Risperdal Consta and has the

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- breast swelling or tenderness (in men or women), nipple discharge , impotence , lack of interest in sex, missed menstrual periods
- severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out
- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- low blood cell counts - fever, mouth sores, skin sores, sore throat , cough, easy bruising, unusual bleeding, trouble breathing, feeling light-headed
- or penis erection that is painful or lasts 4 hours or longer. Serious side effects may be more likely in older adults. Common Risperdal Consta side effects may include: headache
- dizziness, drowsiness, tired feeling
- tremors, twitching or uncontrollable muscle movements
- depressed mood, agitation , anxiety , restless feeling
- muscle or joint pain
- dry mouth, upset stomach , constipation
- weight gain
- or pain in your arms or legs.

647. This particular disease, Bipolar Disorder, interacts with the drug tiagabine and has the following

- hives
- fever
- swollen glands

Drug Side Effects Summary

- painful sores in or around your eyes or mouth
- difficulty breathing
- swelling of your face, lips, tongue, or throat. New seizures have occurred in some people who are not epileptic who take medicine for partial seizures. Call your doctor at once if you have new or worsening seizures. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Tiagabine may cause serious side effects. Call your doctor at once if you have: new or worsened seizures
- confusion, extreme weakness
- a light-headed feeling, like you might pass out
- vision problems
- or severe skin reaction--fever, sore throat , swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of tiagabine may include: tiredness, lack of energy
- dizziness , drowsiness
- nausea , stomach pain
- feeling nervous or irritable
- tremors
- or trouble concentrating.

648. This particular disease, Bipolar Disorder, interacts with the drug Abilify MyCite and has the fol

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Abilify MyCite Maintenance Kit may cause serious side effects. Call your doctor at once if you have: severe agitation , distress, or restless feeling
- twitching or uncontrollable movements of your eyes, lips, tongue, face, arms, or legs
- mask-like appearance of the face, trouble swallowing, problems with speech
- seizure (convulsions)
- severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out
- low blood cell counts--fever, chills, sore throat , weakness, easy bruising, unusual bleeding, purple or red spots under your skin
- or high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor. You may have increased sexual urges, unusual urges to gamble, or other intense urges while taking this medicine. Talk with your doctor if this occurs. Common side effects of Abilify MyCite Maintenance Kit may include: feeling shaky, anxious, or restless
- nausea , constipation
- dizziness, drowsiness
- or sleep problems (insomnia).

649. This particular disease, Bipolar Disorder, interacts with the drug Depakote Sprinkles and has t

- skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Call your doctor at once if the person taking this medicine has signs of liver or pancreas problems, such as: loss of appetite, upper stomach pain (that may spread to your back), ongoing nausea or vomiting, dark urine, swelling in the face, or jaundice (yellowing of the skin or eyes). Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, depression, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), or have thoughts about suicide or

Drug Side Effects Summary

hurting yourself. Call your doctor at once if you have any of these other side effects: easy bruising, unusual bleeding (nose, mouth, or gums), purple or red pinpoint spots under your skin

- fever, swollen glands, mouth sores
- confusion, tiredness, cold feeling, vomiting, change in your mental state
- severe drowsiness
- or worsening seizures. Common side effects of Depakote Sprinkles may include: nausea, vomiting, stomach pain, diarrhea , constipation
- headache , back pain
- dizziness , drowsiness, weakness, tremors
- memory problems, mood changes, trouble sleeping
- bruising or bleeding
- runny nose , sore throat, cough, wheezing, trouble breathing
- fever, flu symptoms
- problems with walking or coordination
- swelling in your hands or feet
- blurred vision, double vision, unusual eye movements
- ringing in your ears
- rash, hair loss
- or changes in weight or appetite.

650. This particular disease, Bipolar Disorder, interacts with the drug Zyprexa Intramuscular and ha

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: More common Bloating or swelling of the face, arms, hands, lower legs, or feet blurred vision change in vision change in

Drug Side Effects Summary

walking and balance clumsiness or unsteadiness difficulty with speaking difficulty with swallowing drooling impaired vision inability to sit still loss of balance control mask-like face muscle trembling, jerking, or stiffness need to keep moving rapid weight gain restlessness shakiness in the legs, arms, hands, or feet shuffling walk slowed movements slurred speech stiffness of the arms and legs tic-like (jerky) movements of the head, face, mouth, and neck tingling of the hands or feet trembling or shaking of the fingers, hands, or feet twisting movements of the body uncontrolled movements, especially of the face, neck, and back unusual weight gain or loss Less common Bladder pain bloody or cloudy urine bruising burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain difficult or labored breathing difficult, burning, or painful urination dizziness excessive muscle tone frequent urge to urinate headache inability to move the eyes increased blinking or spasms of the eyelid itching of the vagina or genital area lack of coordination large, flat, blue, or purplish patches in the skin loss of bladder control loss of memory lower back or side pain muscle tension or tightness nervousness pain, redness, and swelling at the injection site pounding in the ears problems with memory rhythmic movement of the muscles slow, fast, pounding, or irregular heartbeat or pulse speaking is less clear than usual sticking out the tongue thick, white vaginal discharge with no odor or with a mild odor tightness in the chest trouble with breathing, speaking, or swallowing twitching uncontrolled twisting movements of the neck, trunk, arms, or legs unusual or incomplete body or facial movements weakness of the arms and legs

Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Acid or sour stomach back pain belching change in personality difficulty having a bowel movement (stool) discouragement feeling sad or empty fever heartburn increased appetite increased cough indigestion lack of appetite lack or loss of strength

Drug Side Effects Summary

loss of interest or pleasure runny nose sneezing stomach discomfort, upset, or pain stuffy nose thirst trouble sleeping trouble with concentrating watering of the mouth weight gain Less common Body aches or pain chills cold sweats congestion cough dry skin dryness or soreness of the throat false or unusual sense of well-being heavy menstrual bleeding hoarseness joint pain lack of feeling or emotion leg cramps pain in the arms or legs sweating tender, swollen glands in the neck uncaring feelings voice change vomiting Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Zyprexa Intramuscular side effects

651. This particular disease, Bipolar Disorder, interacts with the drug olanzapine / samidorphan and

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include skin rash , fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Long-term use of olanzapine and samidorphan can cause a serious movement disorder that may not be reversible. The longer you use olanzapine and samidorphan, the more likely you are to develop this disorder, especially if you are a woman or an older adult. Samidorphan can make you more sensitive to the effects of opioid medicine, and you could have a fatal opioid overdose. If you use opioid medicine, a person caring for you should seek emergency medical attention if you have severe dizziness or drowsiness, confusion, slow breathing with long pauses, or if you are hard to wake up. Olanzapine and samidorphan may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement,

Drug Side Effects Summary

blinking or eye movement)

- trouble swallowing
- a seizure
- impotence , sexual problems, missed menstrual periods, breast swelling or discharge
- low white blood cell counts--fever, chills, weakness, sore throat , cough, mouth sores, skin sores
- signs of a stroke--sudden numbness or weakness, severe headache , slurred speech, problems with vision or balance
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Common side effects of olanzapine and samidorphan may include: increased appetite, weight gain
- dizziness, drowsiness, weakness
- problems with speech or memory
- tremor, numbness, tingling, burning pain
- headache
- dry mouth, upset stomach , constipation
- back pain
- or increased salivation.

652. This particular disease, Bronchitis, interacts with the drug Mucinex and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common side effects of Mucinex may include:
dizziness
- headache

Drug Side Effects Summary

- rash
- nausea
- vomiting
- stomach upset.

653. This particular disease, Bronchitis, interacts with the drug azithromycin and has the following

- hives, difficult breathing, swelling in your face or throat) or a severe skin reaction (fever, sore throat, burning in your eyes, skin pain, red or purple skin rash that spreads and causes blistering and peeling). Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash, fever, swollen glands, flu-like symptoms, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. This reaction may occur several weeks after you began using azithromycin. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- or liver problems - nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Call your doctor right away if a baby taking azithromycin becomes irritable or vomits while eating or nursing. Older adults may be more likely to have side effects on heart rhythm, including a life-threatening fast heart rate. Common azithromycin side effects may include: diarrhea
- nausea, vomiting, stomach pain
- or headache.

654. This particular disease, Bronchitis, interacts with the drug Azithromycin Dose Pack and has the

- skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of

Drug Side Effects Summary

your skin or eyes. Azithromycin 3 Day Dose Pack may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody

- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- or liver problems-- nausea , vomiting, loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- Call your doctor right away if a baby taking Azithromycin 3 Day Dose Pack becomes irritable or vomits while eating or nursing. Older adults may be more likely to have side effects on heart rhythm, including a life-threatening fast heart rate . Common side effects of Azithromycin 3 Day Dose Pack may include: nausea, vomiting
- or stomach pain.

655. This particular disease, Bronchitis, interacts with the drug amoxicillin and has the following side effects:

- severe stomach pain
- or diarrhea that is watery or bloody (even if it occurs months after your last dose). Common amoxicillin side effects may include: nausea , vomiting , diarrhea
- or rash.

656. This particular disease, Bronchitis, interacts with the drug guaifenesin and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common guaifenesin side effects may include: dizziness , headache
- drowsiness
- rash

Drug Side Effects Summary

- or nausea , vomiting , stomach upset .

657. This particular disease, Bronchitis, interacts with the drug levofloxacin and has the following s

- low blood sugar--headache, hunger, sweating, irritability, dizziness , nausea , fast heart rate , or feeling anxious or shaky
- nerve symptoms in your hands, arms, legs, or feet--numbness, weakness, tingling, burning pain
- serious mood or behavior changes-- nervousness , confusion, agitation, paranoia, hallucinations, memory problems, trouble concentrating, thoughts of suicide
- or signs of tendon rupture--sudden pain, swelling, bruising, tenderness, stiffness, movement problems, or a snapping or popping sound in any of your joints (rest the joint until you receive medical care or instructions). In rare cases, levofloxacin may cause damage to your aorta, the main blood artery of the body. This could lead to dangerous bleeding or death. Get emergency medical help if you have severe and constant pain in your chest, stomach, or back. Stop taking levofloxacin and call your doctor at once if you have : severe stomach pain, diarrhea that is watery or bloody
- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- the first sign of any skin rash, no matter how mild
- muscle weakness, breathing problems
- seizure (convulsions)
- increased pressure inside the skull--severe headaches, ringing in your ears, dizziness, nausea, vision problems, pain behind your eyes
- or liver problems--upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of levofloxacin may include: nausea, constipation , diarrhea
- headache, dizziness

Drug Side Effects Summary

- or trouble sleeping.

658. This particular disease, Bronchitis, interacts with the drug Augmentin and has the following side effects:

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- pale or yellowed skin, dark colored urine, fever, confusion or weakness
- loss of appetite, upper stomach pain
- little or no urination
- or easy bruising or bleeding. Common Augmentin side effects may include: nausea, vomiting
- diarrhea
- rash, itching
- vaginal itching or discharge
- or diaper rash .

659. This particular disease, Bronchitis, interacts with the drug Levaquin and has the following side effects:

- low blood sugar - headache, hunger, sweating, irritability, dizziness , nausea , fast heart rate , or feeling anxious or shaky
- nerve symptoms in your hands, arms, legs, or feet - numbness, weakness, tingling, burning pain
- serious mood or behavior changes - nervousness, confusion, agitation, paranoia, hallucinations, memory problems, trouble concentrating, thoughts of suicide
- or signs of tendon rupture - sudden pain, swelling, bruising, tenderness, stiffness, movement problems, or a snapping or popping sound in any of your joints (rest the joint until you receive medical care or instructions). In rare cases, levofloxacin may cause damage to your aorta, the main blood artery of the body. This could lead to dangerous bleeding or death. Get emergency medical help if you have severe and constant pain in your chest, stomach, or back. Stop taking Levaquin

Drug Side Effects Summary

and call your doctor at once if you have : severe stomach pain, diarrhea that is watery or bloody

- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)

- the first sign of any skin rash, no matter how mild

- muscle weakness, breathing problems

- seizure (convulsions)

- increased pressure inside the skull - severe headaches, ringing in your ears, dizziness, nausea, vision problems, pain behind your eyes

- or liver problems - upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects may include: nausea, constipation , diarrhea

- headache, dizziness

- or trouble sleeping.

660. This particular disease, Bronchitis, interacts with the drug Zithromax and has the following side effects:

- hives, difficult breathing, swelling in your face or throat) or a severe skin reaction (fever, sore throat, burning in your eyes, skin pain, red or purple skin rash that spreads and causes blistering and peeling). Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash, fever, swollen glands, flu-like symptoms, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. This reaction may occur several weeks after you began using Zithromax. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody

- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)

- or liver problems - nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Call your doctor right away if a baby

Drug Side Effects Summary

taking Zithromax becomes irritable or vomits while eating or nursing. Older adults may be more likely to have side effects on heart rhythm, including a life-threatening fast heart rate. Common Zithromax side effects may include: diarrhea

- nausea, vomiting, stomach pain
- or headache.

661. This particular disease, Bronchitis, interacts with the drug amoxicillin / clavulanate and has the following side effects:

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- pale or yellowed skin, dark colored urine, fever, confusion or weakness
- loss of appetite, upper stomach pain
- little or no urination
- or easy bruising or bleeding. Common amoxicillin and clavulanate side effects may include:
nausea, vomiting
- diarrhea
- rash, itching
- vaginal itching or discharge
- or diaper rash .

662. This particular disease, Bronchitis, interacts with the drug clarithromycin and has the following side effects:

- hives , difficult breathing, swelling in your face or throat) or a severe skin reaction (fever , sore throat , burning in your eyes, skin pain, red or purple skin rash that spreads and causes blistering and peeling). Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness

Drug Side Effects Summary

(like you might pass out)

- confusion, spinning sensation
- liver problems - loss of appetite, upper stomach pain, tiredness, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath. If you've ever had heart problems, clarithromycin may cause a heart attack or stroke years after you've taken this medicine. Seek medical attention right away if you have chest pain spreading to your jaw or shoulder, trouble breathing, slurred speech, or sudden numbness or weakness on one side of your body. Older adults may be more likely to have heart rhythm side effects, including a life-threatening fast heart rate . Common clarithromycin side effects may include: stomach pain, nausea, vomiting
- diarrhea
- or unusual or unpleasant taste in your mouth.

663. This particular disease, Bronchitis, interacts with the drug cefdinir and has the following side effects

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- fever, chills, body aches, flu symptoms
- pale skin, easy bruising, unusual bleeding
- seizure (convulsions)
- fever, weakness, confusion
- dark colored urine, jaundice (yellowing of the skin or eyes)
- or kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath. Common side effects of cefdinir may include: nausea, vomiting, stomach pain, diarrhea
- vaginal itching or discharge

Drug Side Effects Summary

- headache
- or rash (including diaper rash in an infant taking liquid cefdinir).

664. This particular disease, Bronchitis, interacts with the drug ciprofloxacin and has the following

- low blood sugar - headache, hunger, irritability, dizziness , nausea , fast heart rate , or feeling shaky
- nerve damage symptoms - numbness, tingling, burning pain in your hands, arms, legs, or feet: serious mood or behavior changes - nervousness, confusion, agitation, paranoia, hallucinations, memory problems, trouble concentrating, thoughts of suicide
- or signs of tendon rupture - sudden pain, swelling, bruising, tenderness, stiffness, movement problems, or a snapping or popping sound in any of your joints (rest the joint until you receive medical care or instructions). In rare cases, ciprofloxacin may cause damage to your aorta, the main blood artery of the body. This could lead to dangerous bleeding or death. Get emergency medical help if you have severe and constant pain in your chest, stomach, or back. Also, stop using this medicine and call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- any skin rash, no matter how mild
- muscle weakness, breathing problems
- little or no urination
- jaundice (yellowing of the skin or eyes)
- or increased pressure inside the skull - severe headaches, ringing in your ears, dizziness, nausea, vision problems, pain behind your eyes. Common ciprofloxacin side effects may include: nausea, vomiting , diarrhea, stomach pain

Drug Side Effects Summary

- headache
- or abnormal liver function tests .

665. This particular disease, Bronchitis, interacts with the drug Amoxil and has the following side effects:

- severe stomach pain
- or diarrhea that is watery or bloody (even if it occurs months after your last dose). Common Amoxil side effects may include: nausea , vomiting , diarrhea
- or rash.

666. This particular disease, Bronchitis, interacts with the drug cefuroxime and has the following side effects:

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- jaundice (yellowing of the skin or eyes)
- fever, chills
- a seizure
- or chest pain. Common side effects of cefuroxime may include: nausea , vomiting , diarrhea
- flu-like symptoms
- vaginal itching or discharge
- or diaper rash (in people using the oral suspension).

667. This particular disease, Bronchitis, interacts with the drug Avelox and has the following side effects:

- low blood sugar - headache, hunger, sweating, irritability, dizziness , nausea , fast heart rate , or feeling anxious or shaky
- nerve symptoms in your hands, arms, legs, or feet - numbness, weakness, tingling, burning pain
- serious mood or behavior changes - nervousness, confusion, agitation, paranoia, hallucinations,

Drug Side Effects Summary

memory problems, trouble concentrating, thoughts of suicide

- or signs of tendon rupture - sudden pain, swelling, bruising, tenderness, stiffness, movement problems, or a snapping or popping sound in any of your joints (rest the joint until you receive medical care or instructions). In rare cases, moxifloxacin may cause damage to your aorta, the main blood artery of the body. This could lead to dangerous bleeding or death. Get emergency medical help if you have severe and constant pain in your chest, stomach, or back. Also stop using Avelox and call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- muscle weakness, breathing problems
- a seizure (convulsions)
- any skin rash, no matter how mild
- increased pressure inside the skull - severe headaches, ringing in your ears, vision problems, pain behind your eyes
- or liver problems - upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common Avelox side effects may include: nausea, diarrhea
- dizziness
- or headache.

668. This particular disease, Bronchitis, interacts with the drug Cipro and has the following side effects

- low blood sugar - headache, hunger, irritability, dizziness , nausea , fast heart rate , or feeling shaky
- nerve damage symptoms - numbness, tingling, burning pain in your hands, arms, legs, or feet: serious mood or behavior changes - nervousness, confusion, agitation, paranoia, hallucinations, memory problems, trouble concentrating, thoughts of suicide

Drug Side Effects Summary

- or signs of tendon rupture - sudden pain, swelling, bruising, tenderness, stiffness, movement problems, or a snapping or popping sound in any of your joints (rest the joint until you receive medical care or instructions). In rare cases, ciprofloxacin may cause damage to your aorta, the main blood artery of the body. This could lead to dangerous bleeding or death. Get emergency medical help if you have severe and constant pain in your chest, stomach, or back. Also, stop using Cipro and call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- any skin rash, no matter how mild
- muscle weakness, breathing problems
- little or no urination
- jaundice (yellowing of the skin or eyes)
- or increased pressure inside the skull - severe headaches, ringing in your ears, dizziness, nausea, vision problems, pain behind your eyes. Common Cipro side effects may include: nausea, vomiting , diarrhea, stomach pain
- headache
- or abnormal liver function tests .

669. This particular disease, Bronchitis, interacts with the drug erythromycin and has the following

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- headache with chest pain and severe dizziness, fainting, fast or pounding heartbeats
- a seizure
- hearing problems (rare)
- pancreatitis - severe pain in your upper stomach spreading to your back, nausea and vomiting

Drug Side Effects Summary

- or liver problems - loss of appetite, stomach pain (upper right side), tiredness, easy bruising or bleeding, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Serious side effects may be more likely in older adults, including hearing loss, or a life-threatening fast heart rate. Call your doctor if a baby using this medicine is vomiting or irritable with feeding. Common erythromycin side effects may include: severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)

- liver problems

- or abnormal liver function tests.

670. This particular disease, Bronchitis, interacts with the drug Ceftin and has the following side effects

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)

- jaundice (yellowing of the skin or eyes)

- fever, chills, headache , muscle pain

- light-headed feeling, like you might pass out

- problems with your hearing

- a seizure

- or kidney problems - little or no urination, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath. Common Ceftin side effects may include: diarrhea

- fever and body aches

- nausea , vomiting

- or vaginal itching or discharge.

671. This particular disease, Bronchitis, interacts with the drug ceftriaxone and has the following side effects

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last

Drug Side Effects Summary

dose)

- new signs of infection (fever, chills, sweating)
- nausea , vomiting , pain in your upper stomach that spreads to your back
- pale or yellowed skin, dark colored urine
- new or worsening breathing problems (wheezing, feeling short of breath)
- a blood cell disorder-- headache , chest pain, dizziness , weakness, severe tingling or numbness
- or kidney or bladder problems--pain in your side or lower back spreading to your groin, blood in your urine, painful or difficult urination, little or no urine. Common side effects of ceftriaxone may include: symptoms of a blood cell disorder
- diarrhea
- vaginal itching or discharge
- warmth, tight feeling, or a hard lump where the injection was given
- rash
- or abnormal liver function tests .

672. This particular disease, Bronchitis, interacts with the drug Apo-Amoxi and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Abdominal or stomach cramps or tenderness back, leg, or stomach pains black, tarry stools bleeding gums blistering, peeling, or loosening of the skin bloating blood in the urine bloody nose chest pain chills clay-colored stools cough dark urine diarrhea diarrhea, watery and severe, which may also be bloody difficulty with breathing difficulty with swallowing dizziness fast heartbeat feeling of discomfort fever general body swelling headache heavier menstrual periods hives or welts increased thirst inflammation of the joints itching joint or muscle pain loss of appetite muscle aches nausea or

Drug Side Effects Summary

vomiting nosebleeds pain pain in the lower back pain or burning while urinating painful or difficult urination pale skin pinpoint red spots on the skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rash red, irritated eyes redness, soreness, or itching skin shortness of breath sore throat sores, ulcers, or white spots in the mouth or on the lips sores, welting, or blisters sudden decrease in the amount of urine swollen, lymph glands tenderness tightness in the chest unpleasant breath odor unusual bleeding or bruising unusual tiredness or weakness unusual weight loss vomiting of blood watery or bloody diarrhea wheezing yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Bad, unusual, or unpleasant (after) taste change in taste Incidence not known Agitation black, hairy tongue changes in behavior confusion convulsions discoloration of the tooth (brown, yellow, or gray staining) dizziness sleeplessness trouble with sleeping unable to sleep white patches in the mouth or throat or on the tongue white patches with diaper rash Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Apo-Amoxi side effects

673. This particular disease, Bronchitis, interacts with the drug beclomethasone and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Beclomethasone can affect growth in children. Talk with your doctor if you think your child is not growing at a normal rate while using this medicine. Beclomethasone inhalation may cause serious side effects. Call your doctor at once if you have:

Drug Side Effects Summary

wheezing, choking, or other breathing problems after using this medicine

- worsening asthma symptoms
- white patches or sores inside your mouth or on your lips
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- signs of infection--fever, chills, body aches, vomiting
- signs of low adrenal gland hormones --worsening tiredness, lack of energy, weakness, feeling light-headed, nausea , vomiting
- or increased adrenal gland hormones --hunger, weight gain, swelling, skin discoloration, slow wound healing, sweating, acne , increased body hair, tiredness, mood changes, muscle weakness, missed menstrual periods, sexual changes. Common side effects of beclomethasone inhalation may include: a yeast infection in the mouth
- sinus pain, irritation in your nose
- or cold symptoms such as stuffy nose, sneezing, sore throat .

674. This particular disease, Bronchitis, interacts with the drug moxifloxacin and has the following

- low blood sugar--headache, hunger, sweating, irritability, dizziness , nausea , fast heart rate , or feeling anxious or shaky
- nerve symptoms in your hands, arms, legs, or feet--numbness, weakness, tingling, burning pain
- serious mood or behavior changes-- nervousness , confusion, agitation, paranoia, hallucinations, memory problems, trouble concentrating, thoughts of suicide
- or signs of tendon rupture--sudden pain, swelling, bruising, tenderness, stiffness, movement problems, or a snapping or popping sound in any of your joints (rest the joint until you receive medical care or instructions). In rare cases, moxifloxacin may cause damage to your aorta, the main blood artery of the body. This could lead to dangerous bleeding or death. Get emergency medical help if you have severe and constant pain in your chest, stomach, or back. Also stop using

Drug Side Effects Summary

moxifloxacin and call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody

- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- muscle weakness, breathing problems
- a seizure (convulsions)
- any skin rash, no matter how mild
- increased pressure inside the skull--severe headaches, ringing in your ears, vision problems, pain behind your eyes
- or liver problems--upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of moxifloxacin may include: nausea, diarrhea
- dizziness
- or headache.

675. This particular disease, Bronchitis, interacts with the drug cefixime and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Cefixime may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- jaundice (yellowing of the skin or eyes)
- pale or yellowed skin, dark colored urine, confusion or weakness
- a seizure (convulsions)
- low blood cell counts--sudden weakness or ill feeling, fever, chills, flu-like symptoms, swollen gums, mouth sores, skin sores, rapid heart rate, pale skin, easy bruising, unusual bleeding, feeling light-headed

Drug Side Effects Summary

- kidney problems--little or no urination, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- or severe skin reaction--fever, sore throat , swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of cefixime may include: indigestion , stomach pain
- nausea , vomiting
- or vaginal itching or discharge.

676. This particular disease, Bronchitis, interacts with the drug Mucinex Maximum Strength and has the following side effects:

- No side effects listed

677. This particular disease, Bronchitis, interacts with the drug Cipro I.V. and has the following side effects:

- low blood sugar--headache, hunger, sweating, irritability, dizziness , nausea , fast heart rate , or feeling anxious or shaky
- nerve symptoms in your hands, arms, legs, or feet--numbness, weakness, tingling, burning pain
- serious mood or behavior changes-- nervousness , confusion, agitation, paranoia, hallucinations, memory problems, trouble concentrating, thoughts of suicide
- or signs of tendon rupture--sudden pain, swelling, bruising, tenderness, stiffness, movement problems, or a snapping or popping sound in any of your joints (rest the joint until you receive medical care or instructions). In rare cases, Cipro I.V. may cause damage to your aorta, the main blood artery of the body. This could lead to dangerous bleeding or death. Get emergency medical help if you have severe and constant pain in your chest, stomach, or back. Also, stop using Cipro I.V. and call your doctor at once if you have: severe stomach pain with diarrhea that is watery or bloody

Drug Side Effects Summary

- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- any skin rash, no matter how mild
- muscle weakness, breathing problems
- little or no urination
- jaundice (yellowing of the skin or eyes)
- or increased pressure inside the skull--severe headaches, ringing in your ears, dizziness, nausea, vision problems, pain behind your eyes. Common side effects of Cipro I.V. may include: nausea, vomiting , diarrhea
- rash
- or abnormal liver function tests .

678. This particular disease, Bronchitis, interacts with the drug Cipro XR and has the following side effects:

- No side effects listed

679. This particular disease, Bronchitis, interacts with the drug Robafen and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common side effects of Robafen may include: nausea
- or vomiting .

680. This particular disease, Bronchitis, interacts with the drug Augmentin XR and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Hives or welts itching

Drug Side Effects Summary

itching of the vagina or genital area pain during sexual intercourse redness of the skin skin rash thick, white vaginal discharge with no odor or with a mild odor Rare Bloody or cloudy urine fever greatly decreased frequency of urination or amount of urine seizures swelling of the feet or lower legs Incidence not known Back, leg, or stomach pains black, hairy tongue black, tarry stools bleeding gums blistering, peeling, or loosening of the skin bloating blood in the stools bloody nose chest pain or tightness chills clay-colored stools cough or hoarseness cracks in the skin dark urine diarrhea diarrhea, watery and severe, which may also be bloody difficulty with breathing difficulty with moving difficulty with swallowing dizziness fast heartbeat fever with or without chills general body swelling general feeling of tiredness or weakness headache heavier menstrual periods increased thirst joint or muscle pain large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs light-colored stools loss of appetite loss of heat from the body lower back or side pain muscle stiffness nausea or vomiting pain pain, swelling, or redness in the joints painful or difficult urination pale skin pinpoint red spots on the skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rash red skin lesions, often with a purple center red, irritated eyes red, swollen skin scaly skin sore throat sores, ulcers, or white spots on the lips or in the mouth stomach cramps or tenderness swollen glands trouble breathing unpleasant breath odor unusual bleeding or bruising unusual tiredness or weakness unusual weight loss upper right abdominal or stomach pain vomiting of blood white patches in the mouth or throat or on the tongue white patches with diaper rash yellow eyes or skin Symptoms of overdose Cloudy urine diarrhea greatly decreased frequency of urination or amount of urine sleepiness stomach pain Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Rare Anxiety dry mouth hyperventilation irregular heartbeats irritability

Drug Side Effects Summary

restlessness shaking sleeplessness tooth discoloration trouble sitting still trouble with sleeping
Incidence not known Burning feeling in the chest or stomach indigestion redness, swelling, or
soreness of the tongue stomach upset swelling or inflammation of the mouth Other side effects not
listed may also occur in some patients. If you notice any other effects, check with your healthcare
professional. Call your doctor for medical advice about side effects. You may report side effects to
the FDA at 1-800-FDA-1088. Augmentin XR side effects

681. This particular disease, Bronchitis, interacts with the drug Biaxin XL and has the following side effects:

- hives , difficult breathing, swelling in your face or throat) or a severe skin reaction (fever , sore throat , burning in your eyes, skin pain, red or purple skin rash that spreads and causes blistering and peeling). Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- confusion, spinning sensation
- liver problems - loss of appetite, upper stomach pain, tiredness, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath. If you've ever had heart problems, clarithromycin may cause a heart attack or stroke years after you've taken this medicine. Seek medical attention right away if you have chest pain spreading to your jaw or shoulder, trouble breathing, slurred speech, or sudden numbness or weakness on one side of your body. Older adults may be more likely to have heart rhythm side effects, including a life-threatening fast heart rate . Common Biaxin XL side effects may include: stomach pain, nausea, vomiting
- diarrhea

Drug Side Effects Summary

- or unusual or unpleasant taste in your mouth.

682. This particular disease, Bronchitis, interacts with the drug cefpodoxime and has the following

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- fever, chills, sore throat, mouth sores, swollen glands, joint pain, or not feeling well
- a seizure
- or liver problems--upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of cefpodoxime may include: diarrhea
- nausea, stomach pain
- headache
- vaginal itching or discharge
- or diaper rash in an infant using cefpodoxime.

683. This particular disease, Bronchitis, interacts with the drug cefprozil and has the following side

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- fever, chills, sore throat, mouth sores
- pale or yellowed skin, dark colored urine, fever, confusion or weakness
- easy bruising, unusual bleeding
- a seizure
- or kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath. Common side effects of cefprozil may include: nausea , diarrhea
- vaginal itching or discharge
- diaper rash

Drug Side Effects Summary

- or abnormal liver function tests .

684. This particular disease, Bronchitis, interacts with the drug Ery-Tab and has the following side effects

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- headache with chest pain and severe dizziness, fainting, fast or pounding heartbeats
- a seizure
- hearing problems (rare)
- pancreatitis - severe pain in your upper stomach spreading to your back, nausea and vomiting
- or liver problems - loss of appetite, stomach pain (upper right side), tiredness, easy bruising or bleeding, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Serious side effects may be more likely in older adults, including hearing loss, or a life-threatening fast heart rate. Call your doctor if a baby using this medicine is vomiting or irritable with feeding. Common Ery-Tab side effects may include: severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- liver problems
- or abnormal liver function tests.

685. This particular disease, Bronchitis, interacts with the drug Guaifenesin G and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any

Drug Side Effects Summary

questions about them: Less common or rare Diarrhea dizziness headache hives nausea or vomiting skin rash stomach pain Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Guaifenesin G side effects

686. This particular disease, Bronchitis, interacts with the drug Suprax and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Suprax may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- jaundice (yellowing of the skin or eyes)
- pale or yellowed skin, dark colored urine, confusion or weakness
- a seizure (convulsions)
- low blood cell counts--sudden weakness or ill feeling, fever, chills, flu-like symptoms, swollen gums, mouth sores, skin sores, rapid heart rate, pale skin, easy bruising, unusual bleeding, feeling light-headed
- kidney problems--little or no urination, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- or severe skin reaction--fever, sore throat , swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of Suprax may include: indigestion , stomach pain
- nausea , vomiting
- or vaginal itching or discharge.

Drug Side Effects Summary

687. This particular disease, Bronchitis, interacts with the drug Altarussin and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common side effects of Altarussin may include:
nausea
- or vomiting .

688. This particular disease, Bronchitis, interacts with the drug Amibid LA and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common or rare Diarrhea dizziness headache hives nausea or vomiting skin rash stomach pain Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Amibid LA side effects

689. This particular disease, Bronchitis, interacts with the drug Amoclan and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Hives or welts itching itching of the vagina or genital area pain during sexual intercourse redness of the skin skin rash thick, white vaginal discharge with no odor or with a mild odor Rare Bloody or cloudy urine fever

Drug Side Effects Summary

greatly decreased frequency of urination or amount of urine seizures swelling of the feet or lower legs Incidence not known Back, leg, or stomach pains black, hairy tongue black, tarry stools bleeding gums blistering, peeling, or loosening of the skin bloating blood in the stools bloody nose chest pain or tightness chills clay-colored stools cough or hoarseness cracks in the skin dark urine diarrhea diarrhea, watery and severe, which may also be bloody difficulty with breathing difficulty with moving difficulty with swallowing dizziness fast heartbeat fever with or without chills general body swelling general feeling of tiredness or weakness headache heavier menstrual periods increased thirst joint or muscle pain large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs light-colored stools loss of appetite loss of heat from the body lower back or side pain muscle stiffness nausea or vomiting pain pain, swelling, or redness in the joints painful or difficult urination pale skin pinpoint red spots on the skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rash red skin lesions, often with a purple center red, irritated eyes red, swollen skin scaly skin sore throat sores, ulcers, or white spots on the lips or in the mouth stomach cramps or tenderness swollen glands trouble breathing unpleasant breath odor unusual bleeding or bruising unusual tiredness or weakness unusual weight loss upper right abdominal or stomach pain vomiting of blood white patches in the mouth or throat or on the tongue white patches with diaper rash yellow eyes or skin Symptoms of overdose Cloudy urine diarrhea greatly decreased frequency of urination or amount of urine sleepiness stomach pain Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Rare Anxiety dry mouth hyperventilation irregular heartbeats irritability restlessness shaking sleeplessness tooth discoloration trouble sitting still trouble with sleeping Incidence not known Burning feeling in the chest or stomach indigestion redness, swelling, or

Drug Side Effects Summary

soreness of the tongue stomach upset swelling or inflammation of the mouth Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Amoclan side effects

690. This particular disease, Bronchitis, interacts with the drug ampicillin and has the following side effects:

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- blisters, ulcers, or soreness in your mouth
- skin rash, redness, or itching
- fever, chills, sore throat, swollen glands, joint pain, or not feeling well
- pale skin, cold hands and feet
- or feeling light-headed or short of breath. Common side effects of ampicillin may include: nausea , vomiting , stomach pain, diarrhea
- rash
- swollen, black, or "hairy" tongue
- or vaginal itching or discharge.

691. This particular disease, Bronchitis, interacts with the drug Balminil Expectorant and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Although serious side effects occur rarely when this medicine is taken as recommended, they may be more likely to occur if: too much medicine is taken, it is taken in large doses, or it is taken for a long period of time. For narcotic antitussive (codeine, dihydrocodeine, hydrocodone, or hydromorphone)-containing If you are a nursing mother and you notice any of the following symptoms of overdose in your baby,

Drug Side Effects Summary

get emergency help immediately : Cold, clammy skin confusion (severe) convulsions (seizures) drowsiness or dizziness (severe) nervousness or restlessness (severe) pinpoint pupils of eyes slow heartbeat slow or troubled breathing Difficulty breathing difficulty nursing increased sleepiness (more than usual) limpness For acetaminophen-containing Diarrhea increased sweating loss of appetite nausea or vomiting stomach cramps or pain swelling or tenderness in the upper abdomen or stomach area For salicylate-containing Any loss of hearing bloody urine confusion convulsions (seizures) dizziness or lightheadedness drowsiness (severe) excitement or nervousness (severe) fast or deep breathing fever hallucinations (seeing, hearing, or feeling things that are not there) increased sweating nausea or vomiting (severe or continuing) shortness of breath or troubled breathing (for salicylamide only) stomach pain (severe or continuing) uncontrollable flapping movements of the hands, especially in elderly patients unusual thirst vision problems For decongestant-containing Fast, pounding, or irregular heartbeat headache (continuing and severe) nausea or vomiting (severe) nervousness or restlessness (severe) shortness of breath or troubled breathing (severe or continuing) Check with your doctor as soon as possible if any of the following side effects occur: For all combinations Skin rash, hives, and/or itching For antihistamine- or anticholinergic-containing Clumsiness or unsteadiness convulsions (seizures) drowsiness (severe) dryness of mouth, nose, or throat (severe) flushing or redness of face hallucinations (seeing, hearing, or feeling things that are not there) restlessness (severe) shortness of breath or troubled breathing slow or fast heartbeat For iodine-containing Headache (continuing) increased watering of mouth loss of appetite metallic taste skin rash, hives, or redness sore throat swelling of face, lips, or eyelids For acetaminophen-containing Unexplained sore throat and fever unusual tiredness or weakness yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects

Drug Side Effects Summary

continue or are bothersome or if you have any questions about them: Constipation decreased sweating difficult or painful urination dizziness or lightheadedness drowsiness dryness of mouth, nose, or throat false sense of well-being increased sensitivity of skin to sun nausea or vomiting nightmares stomach pain thickening of mucus trouble in sleeping unusual excitement, nervousness, restlessness, or irritability unusual tiredness or weakness Not all of the side effects listed above have been reported for each of these medicines, but they have been reported for at least one of them. There are some similarities among these combination medicines, so many of the above side effects may occur with any of these medicines. Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Balminil Expectorant side effects

692. This particular disease, Bronchitis, interacts with the drug Benylin E and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common or rare Diarrhea dizziness headache hives nausea or vomiting skin rash stomach pain Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Benylin E side effects

693. This particular disease, Bronchitis, interacts with the drug Bidex-400 and has the following side effects:

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common side effects of Bidex-400 may include:
nausea
- or vomiting .

694. This particular disease, Bronchitis, interacts with the drug Cefzil and has the following side effects

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- fever, chills, sore throat, mouth sores
- pale or yellowed skin, dark colored urine, fever, confusion or weakness
- easy bruising, unusual bleeding
- a seizure
- or kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath. Common Cefzil side effects may include: nausea , diarrhea
- vaginal itching or discharge
- diaper rash
- or abnormal liver function tests .

695. This particular disease, Bronchitis, interacts with the drug dicloxacillin and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Dicloxacillin may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- nausea, vomiting

Drug Side Effects Summary

- little or no urination
- unusual bleeding or bruising
- a seizure (convulsions)
- mouth pain or irritation, black tongue, sore throat
- or fever, swollen glands, rash or itching, joint pain, or general ill feeling. Common side effects of dicloxacillin may include: heartburn, nausea, vomiting, diarrhea
- blisters or ulcers in your mouth, red or swollen gums, trouble swallowing
- or abnormal liver function tests .

696. This particular disease, Bronchitis, interacts with the drug Drituss G and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common or rare Diarrhea dizziness headache hives nausea or vomiting skin rash stomach pain Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Drituss G side effects

697. This particular disease, Bronchitis, interacts with the drug E.E.S. Granules and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Blistering, peeling, or loosening of

Drug Side Effects Summary

the skin chills cough diarrhea difficulty with swallowing dizziness fast heartbeat hives or welts, skin rash joint or muscle pain puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red, irritated eyes red skin lesions, often with a purple center redness of the skin sore throat sores, ulcers, or white spots in the mouth or on the lips tightness in the chest unusual tiredness or weakness Incidence not known Bloating bloody or cloudy urine chest pain or discomfort dark urine diarrhea, watery and severe, which may also be bloody fainting fever general tiredness and weakness greatly decreased frequency of urination or amount of urine hearing loss increased thirst irregular heartbeat recurrent irregular or slow heart rate light-colored stools nausea pain stomach cramp or tenderness swelling of the feet or lower legs unusual weight loss upper right abdominal or stomach pain vomiting yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Diarrhea (mild) loss of appetite weight loss Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. E.E.S. Granules side effects

698. This particular disease, Bronchitis, interacts with the drug Eryc and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Blistering, peeling, or loosening of the skin chills cough diarrhea difficulty with swallowing dizziness fast heartbeat hives or welts, skin rash joint or muscle pain puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue

Drug Side Effects Summary

red, irritated eyes red skin lesions, often with a purple center redness of the skin sore throat sores, ulcers, or white spots in the mouth or on the lips tightness in the chest unusual tiredness or weakness Incidence not known Bloating bloody or cloudy urine chest pain or discomfort dark urine diarrhea, watery and severe, which may also be bloody fainting fever general tiredness and weakness greatly decreased frequency of urination or amount of urine hearing loss increased thirst irregular heartbeat recurrent irregular or slow heart rate light-colored stools nausea pain stomach cramp or tenderness swelling of the feet or lower legs unusual weight loss upper right abdominal or stomach pain vomiting yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Diarrhea (mild) loss of appetite weight loss Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Eryc side effects

699. This particular disease, Bronchitis, interacts with the drug EryPed and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Blistering, peeling, or loosening of the skin chills cough diarrhea difficulty with swallowing dizziness fast heartbeat hives or welts, skin rash joint or muscle pain puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red, irritated eyes red skin lesions, often with a purple center redness of the skin sore throat sores, ulcers, or white spots in the mouth or on the lips tightness in the chest unusual tiredness or

Drug Side Effects Summary

weakness Incidence not known Bloating bloody or cloudy urine chest pain or discomfort dark urine diarrhea, watery and severe, which may also be bloody fainting fever general tiredness and weakness greatly decreased frequency of urination or amount of urine hearing loss increased thirst irregular heartbeat recurrent irregular or slow heart rate light-colored stools nausea pain stomach cramp or tenderness swelling of the feet or lower legs unusual weight loss upper right abdominal or stomach pain vomiting yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Diarrhea (mild) loss of appetite weight loss Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. EryPed side effects

700. This particular disease, Bronchitis, interacts with the drug Erythrocin and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Blistering, peeling, or loosening of the skin chills cough diarrhea difficulty with swallowing dizziness fast heartbeat hives or welts, skin rash joint or muscle pain puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red, irritated eyes red skin lesions, often with a purple center redness of the skin sore throat sores, ulcers, or white spots in the mouth or on the lips tightness in the chest unusual tiredness or weakness Incidence not known Bloating bloody or cloudy urine chest pain or discomfort dark urine diarrhea, watery and severe, which may also be bloody fainting fever general tiredness and

Drug Side Effects Summary

weakness greatly decreased frequency of urination or amount of urine hearing loss increased thirst irregular heartbeat recurrent irregular or slow heart rate light-colored stools nausea pain stomach cramp or tenderness swelling of the feet or lower legs unusual weight loss upper right abdominal or stomach pain vomiting yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Diarrhea (mild) loss of appetite weight loss Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Erythrocin side effects

701. This particular disease, Bronchitis, interacts with the drug Erythrocin Lactobionate and has the

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: Incidence not known Blistering, peeling, or loosening of the skin chest pain or discomfort chills cough diarrhea difficulty with swallowing dizziness fainting fast, pounding, or irregular heartbeat or pulse hearing loss hives or welts, itching, skin rash irregular or slow heart rate irritation at the injection site joint or muscle pain puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red, irritated eyes red skin lesions, often with a purple center redness of the skin sore throat sores, ulcers, or white spots in the mouth or on the lips tightness in the chest unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to

Drug Side Effects Summary

tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Bleeding, blistering, burning, coldness, discoloration of the skin, feeling of pressure, hives, infection, inflammation, itching, lumps, numbness, pain, rash, redness, scarring, soreness, stinging, swelling, tenderness, tingling, ulceration, or warmth at the injection site Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Erythrocin Lactobionate side effects

702. This particular disease, Bronchitis, interacts with the drug Fenesin IR and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common side effects of Fenesin IR may include:
nausea
- or vomiting .

703. This particular disease, Bronchitis, interacts with the drug Floxin and has the following side effects

- low blood sugar - headache, hunger, sweating, irritability, dizziness , nausea , fast heart rate , or feeling anxious or shaky
- nerve symptoms in your hands, arms, legs, or feet - numbness, weakness, tingling, burning pain
- serious mood or behavior changes - nervousness, confusion, agitation, paranoia, hallucinations, memory problems, trouble concentrating, thoughts of suicide
- or signs of tendon rupture - sudden pain, swelling, bruising, tenderness, stiffness, movement problems, or a snapping or popping sound in any of your joints (rest the joint until you receive

Drug Side Effects Summary

medical care or instructions). In rare cases, ofloxacin may cause damage to your aorta, the main blood artery of the body. This could lead to dangerous bleeding or death. Get emergency medical help if you have severe and constant pain in your chest, stomach, or back. Also, stop using Floxin and call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody

- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- the first sign of any skin rash, no matter how mild
- muscle weakness, breathing problems
- little or no urination
- a seizure (convulsions)
- increased pressure inside the skull - severe headaches, ringing in your ears, vision problems, pain behind your eyes
- or liver problems - upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common Floxin side effects may include: nausea, constipation , diarrhea
- dizziness
- or headache.

704. This particular disease, Bronchitis, interacts with the drug gemifloxacin and has the following

- low blood sugar--headache, hunger, sweating, irritability, dizziness , nausea , fast heart rate , or feeling anxious or shaky
- nerve symptoms in your hands, arms, legs, or feet--numbness, weakness, tingling, burning pain
- serious mood or behavior changes-- nervousness , confusion, agitation, paranoia, hallucinations, memory problems, trouble concentrating, thoughts of suicide
- or signs of tendon rupture--sudden pain, swelling, bruising, tenderness, stiffness, movement

Drug Side Effects Summary

problems, or a snapping or popping sound in any of your joints (rest the joint until you receive medical care or instructions). In rare cases, gemifloxacin may cause damage to your aorta, the main blood artery of the body. This could lead to dangerous bleeding or death. Get emergency medical help if you have severe and constant pain in your chest, stomach, or back. Also, stop using gemifloxacin and call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody

- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- the first sign of any skin rash, no matter how mild
- muscle weakness, breathing problems
- jaundice (yellowing of the skin or eyes)
- a seizure (convulsions)
- or increased pressure inside the skull--severe headaches, ringing in your ears, vision problems, pain behind your eyes. Common side effects of gemifloxacin may include: nausea, vomiting , stomach pain, diarrhea
- headache
- dizziness
- or rash.

705. This particular disease, Bronchitis, interacts with the drug Guaifenesin LA and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care

Drug Side Effects Summary

professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common or rare Diarrhea dizziness headache hives nausea or vomiting skin rash stomach pain Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Guaifenesin LA side effects

706. This particular disease, Bronchitis, interacts with the drug Guaifenesin and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Cough that does not go away. Guaifenesin side effects

707. This particular disease, Bronchitis, interacts with the drug Hyaluronate and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs

Drug Side Effects Summary

of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Cough that does not go away. Hytuss side effects

708. This particular disease, Bronchitis, interacts with the drug Mucinex for Kids and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Cough that does not go away. Mucinex for Kids side effects

Drug Side Effects Summary

709. This particular disease, Bronchitis, interacts with the drug ofloxacin and has the following side effects:

- low blood sugar--headache, hunger, sweating, irritability, dizziness , nausea , fast heart rate , or feeling anxious or shaky
- nerve symptoms in your hands, arms, legs, or feet--numbness, weakness, tingling, burning pain
- serious mood or behavior changes-- nervousness , confusion, agitation, paranoia, hallucinations, memory problems, trouble concentrating, thoughts of suicide
- or signs of tendon rupture--sudden pain, swelling, bruising, tenderness, stiffness, movement problems, or a snapping or popping sound in any of your joints (rest the joint until you receive medical care or instructions). In rare cases, ofloxacin may cause damage to your aorta, the main blood artery of the body. This could lead to dangerous bleeding or death. Get emergency medical help if you have severe and constant pain in your chest, stomach, or back. Also, stop using ofloxacin and call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- the first sign of any skin rash, no matter how mild
- muscle weakness, breathing problems
- little or no urination
- a seizure (convulsions)
- increased pressure inside the skull--severe headaches, ringing in your ears, vision problems, pain behind your eyes
- or liver problems--upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of ofloxacin may include: nausea, constipation , diarrhea
- dizziness

Drug Side Effects Summary

- or headache.

710. This particular disease, Bronchitis, interacts with the drug PCE Dispertab and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Blistering, peeling, or loosening of the skin chills cough diarrhea difficulty with swallowing dizziness fast heartbeat hives or welts, skin rash joint or muscle pain puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red, irritated eyes red skin lesions, often with a purple center redness of the skin sore throat sores, ulcers, or white spots in the mouth or on the lips tightness in the chest unusual tiredness or weakness Incidence not known Bloating bloody or cloudy urine chest pain or discomfort dark urine diarrhea, watery and severe, which may also be bloody fainting fever general tiredness and weakness greatly decreased frequency of urination or amount of urine hearing loss increased thirst irregular heartbeat recurrent irregular or slow heart rate light-colored stools nausea pain stomach cramp or tenderness swelling of the feet or lower legs unusual weight loss upper right abdominal or stomach pain vomiting yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Diarrhea (mild) loss of appetite weight loss Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. PCE Dispertab side effects

Drug Side Effects Summary

711. This particular disease, Bronchitis, interacts with the drug Scot-Tussin Expectorant and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common side effects of Scot-Tussin may include:
nausea
- or vomiting .

712. This particular disease, Bronchitis, interacts with the drug Sulfatrim Pediatric and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Black, tarry stools blistering, peeling, or loosening of the skin changes in skin color chest pain or tightness chills clay-colored stools cough or hoarseness dark urine diarrhea dizziness fever general feeling of tiredness or weakness headache itching, skin rash joint or muscle pain light-colored stools loss of appetite lower back or side pain nausea pain, tenderness, or swelling of the foot or leg painful or difficult urination pale skin red skin lesions, often with a purple center red, irritated eyes sore throat sores, ulcers, or white spots in the mouth or on the lips stomach pain swollen or painful glands trouble breathing unpleasant breath odor unusual bleeding or bruising vomiting of blood yellow eyes or skin Incidence not known Back, leg, or stomach pains bleeding gums blindness or vision changes bloating blood in the urine or stools bluish-colored lips, fingernails, or palms burning, crawling, itching, numbness, painful, prickling, "pins and needles", or tingling feelings burning of the face or mouth cloudy urine clumsiness or unsteadiness confusion constipation continuing ringing or buzzing or other unexplained noise in the ears cracks in the skin decreased frequency or amount of urine difficulty with swallowing fainting spells fast, pounding, or irregular heartbeat or pulse general body swelling general feeling of discomfort or illness hair loss hearing loss hives increased thirst

Drug Side Effects Summary

indigestion large, flat, blue, or purplish patches in the skin large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs loss of heat from the body muscle or joint pain nosebleeds not able to pass urine numbness or tingling in the hands, feet, or lips pain or burning while urinating pinpoint red spots on the skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue raised red swellings on the skin, the buttocks, legs, or ankles redness of the white part of the eyes redness, swelling, or soreness of the tongue seizures soreness of the muscles stiff neck or back stomach tenderness swelling of the face, hands, legs, and feet unsteadiness, trembling, or other problems with muscle control or coordination weakness in the hands or feet weakness or heaviness of the legs weight gain or loss Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Passing of gas Incidence not known Discouragement feeling of constant movement of self or surroundings feeling sad or empty increased sensitivity of the skin to sunlight lack of feeling or emotion loss of interest or pleasure muscle pain, stiffness, cramps, or spasms nervousness redness or other discoloration of the skin seeing, hearing, or feeling things that are not there sensation of spinning severe sunburn trouble concentrating trouble sleeping uncaring weight loss Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Sulfatrim Pediatric side effects

713. This particular disease, Bronchitis, interacts with the drug Tussin Expectorant and has the following side effects:

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Common side effects of Tussin Expectorant may include: nausea
- or vomiting .

714. This particular disease, Bronchitis, interacts with the drug Xpect and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common side effects of Xpect may include: nausea
- or vomiting .

715. This particular disease, Bronchitis, interacts with the drug Zinacef and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Black, tarry stools chest pain chills cough fever painful or difficult urination shortness of breath sore throat sores, ulcers, or white spots on lips or in mouth swollen glands unusual bleeding or bruising unusual tiredness or weakness Less common Abdominal or stomach cramps abdominal or stomach tenderness or pain bloating bluish color or changes in skin color diarrhea, watery and severe, which may also be bloody difficulty in breathing or swallowing, wheezing, shortness of breath fast heartbeat fever hives or welts increased thirst muscle spasm or jerking of all extremities nausea or vomiting pain skin itching, rash, or redness sudden loss of consciousness swelling of face, throat, or tongue swelling of foot or leg tenderness unusual weight loss Rare Blistering, peeling, loosening of skin bloody or cloudy urine dizziness fast heartbeat greatly decreased frequency of urination or amount of urine hearing loss, mild to moderate joint or muscle pain puffiness or swelling of the eyelids or around the eyes, face, lips or tongue red or irritated eyes redness, tenderness, itching, burning, or peeling of skin red skin

Drug Side Effects Summary

lesions, often with a purple center sore throat tightness in chest Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Gas loss of appetite Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Zinacef side effects

716. This particular disease, Bronchitis, interacts with the drug Avelox I.V. and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: Rare Black, tarry stools bleeding gums bleeding, blistering, burning, coldness, discoloration of the skin, feeling of pressure, hives, infection, inflammation, itching, lumps, numbness, pain, rash, redness, scarring, soreness, stinging, swelling, tenderness, tingling, ulceration, or warmth at the injection site blisters bloating or swelling of the face, arms, hands, lower legs, or feet blood in the urine or stools blurred vision bone pain burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain chills clay-colored stools cough crying dark urine deep or fast breathing with dizziness diarrhea, watery and severe, which may also be bloody difficult or labored breathing difficulty with moving difficulty with swallowing discouragement dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position dry mouth excessive muscle tone fainting fast, irregular, pounding, or racing heartbeat or pulse feeling of unreality feeling of warmth or heat feeling sad or empty fever flushed, dry skin flushing or redness of the skin, especially on the face and neck fruit-like breath odor headache increased hunger increased sensitivity of the skin to sunlight

Drug Side Effects Summary

increased thirst increased urination irregular heartbeat, recurrent irritability joint pain, stiffness, or swelling lack of coordination loss of appetite loss of interest or pleasure lower back, side, or stomach pain mood or mental changes muscle cramps, pains, stiffness, tension, or tightness nausea nervousness noisy breathing numbness of the feet, hands, and around the mouth pain in the pelvis pain, warmth, or burning in the fingers, toes, and legs painful or difficult urination painful, swollen joints pale skin pinpoint red spots on the skin pounding in the ears problems with speech or speaking problems with vision or hearing quick to react or overreact emotionally rapid weight gain rapidly changing moods redness or other discoloration of the skin restlessness seeing, hearing, or feeling things that are not there seizures sensation of the skin burning sense of detachment from self or body severe sunburn shakiness in the legs, arms, hands, or feet skin rash or itching sore throat sores, ulcers, or white spots on the lips or in the mouth stomach cramps or tenderness sweating swelling of the feet or lower legs swelling or puffiness of the face swollen glands thick, white vaginal discharge with no odor or with a mild odor tightness in the chest tingling of the hands or feet trouble concentrating trouble sleeping troubled breathing with exertion unexplained weight loss unpleasant breath odor unusual bleeding or bruising unusual tiredness or weakness unusual weight gain or loss vomiting vomiting of blood white patches in the mouth or on the tongue yellow eyes or skin Incidence not known Blistering, peeling, or loosening of the skin burning, numbness, tingling, or painful sensations change in the ability to see colors, especially blue or yellow confusion as to time, place, or person difficulty with chewing or talking double vision drooping eyelids eye pain general feeling of tiredness or weakness hives hoarseness irregular or slow heart rate large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or genitals light-colored stools loss of consciousness muscle weakness no blood pressure or pulse puffiness or swelling of the eyelids or around the eyes, face, lips or tongue red, irritated eyes red skin lesions, often with a purple center severe headache severe tiredness stomach pain, continuing stopping of the heart unsteadiness or awkwardness unusual excitement, nervousness, or restlessness weakness in the

Drug Side Effects Summary

arms, hands, legs, or feet Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Rare Bad, unusual, or unpleasant (after) taste belching burning feeling in the chest or stomach change in sense of smell change in taste changes in vision continuing ringing or buzzing or other unexplained noise in the ears difficulty having a bowel movement excess air or gas in the stomach or bowels fear or nervousness feeling of constant movement of self or surroundings full feeling general feeling of discomfort or illness hearing loss heartburn impaired vision indigestion itching of the vagina or genital area lack or loss of strength loss of memory pain during sexual intercourse passing of gas problems with memory redness, swelling, or soreness of the tongue sensation of spinning sleepiness or unusual drowsiness sore mouth or tongue stomach discomfort, upset, or pain Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Avelox I.V. side effects

717. This particular disease, Bronchitis, interacts with the drug cefaclor and has the following side

- hives , numbness or tingling
- weakness, feeling light-headed
- difficult breathing
- swelling of your face, lips, tongue, or throat. Cefaclor may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- fever, swollen glands, rash or itching, joint pain, or general ill feeling
- easy bruising or bleeding, severe tingling, numbness, pain, muscle weakness

Drug Side Effects Summary

- seizure (convulsions)
- jaundice (yellowing of the skin or eyes)
- or severe skin reaction--fever, sore throat , swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of cefaclor may include: diarrhea
- or vaginal itching or discharge.

718. This particular disease, Bronchitis, interacts with the drug cilastatin / imipenem and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Imipenem and cilastatin may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- upper stomach pain, loss of appetite
- jaundice (yellowing of the skin or eyes)
- a seizure (convulsions)
- fever
- or a light-headed feeling, like you might pass out. Common side effects of imipenem and cilastatin may include: pain, swelling, redness, bruising, or hardening where the medicine was injected
- dizziness , drowsiness
- nausea , vomiting , diarrhea
- or itching, rash.

719. This particular disease, Bronchitis, interacts with the drug Dycill and has the following side effects:

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Dycill may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- nausea, vomiting
- little or no urination
- unusual bleeding or bruising
- a seizure (convulsions)
- mouth pain or irritation, black tongue, sore throat
- or fever, swollen glands, rash or itching, joint pain, or general ill feeling. Common side effects of Dycill may include: heartburn, nausea, vomiting, diarrhea
- blisters or ulcers in your mouth, red or swollen gums, trouble swallowing
- or abnormal liver function tests .

720. This particular disease, Bronchitis, interacts with the drug Dynapen and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Dynapen may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- nausea, vomiting
- little or no urination
- unusual bleeding or bruising
- a seizure (convulsions)
- mouth pain or irritation, black tongue, sore throat
- or fever, swollen glands, rash or itching, joint pain, or general ill feeling. Common side effects of Dynapen may include: heartburn, nausea, vomiting, diarrhea
- blisters or ulcers in your mouth, red or swollen gums, trouble swallowing

Drug Side Effects Summary

- or abnormal liver function tests .

721. This particular disease, Bronchitis, interacts with the drug Factive and has the following side effects

- low blood sugar--headache, hunger, sweating, irritability, dizziness , nausea , fast heart rate , or feeling anxious or shaky
- nerve symptoms in your hands, arms, legs, or feet--numbness, weakness, tingling, burning pain
- serious mood or behavior changes-- nervousness , confusion, agitation, paranoia, hallucinations, memory problems, trouble concentrating, thoughts of suicide
- or signs of tendon rupture--sudden pain, swelling, bruising, tenderness, stiffness, movement problems, or a snapping or popping sound in any of your joints (rest the joint until you receive medical care or instructions). In rare cases, Factive may cause damage to your aorta, the main blood artery of the body. This could lead to dangerous bleeding or death. Get emergency medical help if you have severe and constant pain in your chest, stomach, or back. Also, stop using Factive and call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- the first sign of any skin rash, no matter how mild
- muscle weakness, breathing problems
- jaundice (yellowing of the skin or eyes)
- a seizure (convulsions)
- or increased pressure inside the skull--severe headaches, ringing in your ears, vision problems, pain behind your eyes. Common side effects of Factive may include: nausea, vomiting , stomach pain, diarrhea
- headache
- dizziness

Drug Side Effects Summary

- or rash.

722. This particular disease, Bronchitis, interacts with the drug Primaxin IV and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Primaxin IV may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- upper stomach pain, loss of appetite
- jaundice (yellowing of the skin or eyes)
- a seizure (convulsions)
- fever
- or a light-headed feeling, like you might pass out. Common side effects of Primaxin IV may include: pain, swelling, redness, bruising, or hardening where the medicine was injected
- dizziness , drowsiness
- nausea , vomiting , diarrhea
- or itching, rash.

723. This particular disease, Cancer, interacts with the drug carboplatin and has the following side effects:

- Carboplatin may cause unwanted effects, particularly because carboplatin interferes with the growth of normal cells as well as cancer cells. For example, the occurrence of another cancer (secondary malignancy) has been reported in patients receiving cancer chemotherapy with multiple drugs. It is not always possible to tell whether such effects are caused by carboplatin, another drug you may be taking, or your illness. Because some of these effects may be serious, you will need close medical supervision during treatment with carboplatin. The most serious side effects of carboplatin are: bleeding and reduced blood cells, including reduced red blood cells (anemia) and

Drug Side Effects Summary

platelets (needed for proper blood clotting), which may be severe enough to require blood transfusion. You should tell your doctor right away if you notice any unusual bruising or bleeding, including black tarry stools or blood in the urine. infection - carboplatin can temporarily lower the number of white blood cells in your blood, increasing the risk of infection

- life-threatening allergic reaction - during and after treatment the doctor or nurse will observe you carefully for signs of allergic reaction

- kidney and liver problems

- loss of hearing or ringing in the ears

- Contact your doctor right away if you experience any of these effects, or notice effects that worry you or are troublesome. Of the less serious side effects associated with carboplatin treatment, the most common are nausea, vomiting, diarrhea, loss of appetite, hair loss and numbness, tingling, burning, or pain in the hands and feet. Carboplatin side effects

724. This particular disease, Cancer, interacts with the drug Leukeran and has the following side effects

- a seizure

- an unusual mass or lump

- severe vomiting or diarrhea

- new or worsening cough

- bone marrow suppression-- dizziness , pale lips or fingernail beds, fast heart rate , getting easily tired or short of breath

- liver problems--loss of appetite, stomach pain (upper right side), dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)

- or low blood cell counts--fever, chills, tiredness, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath. Common side effects of Leukeran may include: bone marrow suppression

Drug Side Effects Summary

- low blood cell counts
- missed menstrual periods
- nausea , vomiting, diarrhea
- or sores or white patches in or around your mouth.

725. This particular disease, Cancer, interacts with the drug chlorambucil and has the following side effects:

- What are the serious (bad) side effects of chlorambucil? Allergic Reactions: You should immediately get emergency medical help if you have signs of: allergic reaction (difficulty breathing, swelling in your face or throat, hives) or a severe skin reaction (fever, sore throat , burning eyes, skin pain, red or purple skin rash with blistering and peeling). Other serious (bad) side effects: You should call your doctor at once if you have: a seizure any unusual lump or mass, severe diarrhea or vomiting, a cough that is new or worsening, signs of bone marrow suppression eg. dizziness , feeling tired or short of breath, pale lips or fingernail beds, fast heart rate , signs of liver problems eg. low appetite, pain in the upper right side of stomach, dark urine, clay-colored stools, yellowing of the skin or eyes, or signs of low blood cell counts eg. high body temperature, chills, tiredness, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath. What are the common side effects of chlorambucil? nausea, vomiting or diarrhea, white patches or sores in or around your mouth, bone marrow suppression, low blood cell counts, missed menstrual periods in women.

726. This particular disease, Cancer, interacts with the drug fluorouracil and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Fluorouracil may cause serious side effects. Call your doctor at once if you have: fever (take your temperature each day while receiving fluorouracil)

Drug Side Effects Summary

- severe or ongoing diarrhea
- vision problems
- confusion, problems with balance or muscle movement
- painful mouth sores, red or swollen gums, trouble swallowing, talking, or eating
- bone marrow suppression-- dizziness , pale lips or fingernail beds, fast heart rate , getting easily tired or short of breath
- "hand and foot syndrome"--pain, blisters, bleeding, or severe rash on the palms of your hands or the soles of your feet
- heart problems--chest pain or pressure, pain spreading to your jaw or shoulder, irregular heartbeats, nausea , sweating, feeling dizzy or short of breath. Your cancer treatments may be delayed or permanently discontinued if you have certain side effects. Common side effects of fluorouracil may include: diarrhea
- mouth sores
- heart problems
- or bone marrow suppression.

727. This particular disease, Cancer, interacts with the drug doxorubicin and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some side effects may occur during the injection. Tell your caregiver right away if you feel dizzy, nauseated, light-headed, sweaty, or have a headache , chest tightness, back pain , trouble breathing, or swelling in your face. Doxorubicin may cause serious side effects. Call your doctor at once if you have: pain, blisters, or skin sores where the injection was given
- missed menstrual periods

Drug Side Effects Summary

- easy bruising, unusual bleeding (nose, mouth, vagina, or rectum), purple or red pinpoint spots under your skin
- low white blood cell counts--fever, swollen gums, painful mouth sores, pain when swallowing, skin sores, cold or flu symptoms, cough, trouble breathing
- or signs of heart problems--fast heartbeats, shortness of breath (even with mild exertion), feeling very weak or tired, swelling in your ankles or feet. Doxorubicin may cause your urine to turn a reddish-orange color. This side effect is usually not harmful. Common side effects of doxorubicin may include: nausea, vomiting
- or hair loss .

728. This particular disease, Cancer, interacts with the drug etoposide and has the following side effects

- sudden chest pain or discomfort, wheezing, dry cough or hack
- easy bruising, unusual bleeding (nose, mouth, vagina, or rectum), purple or red pinpoint spots under your skin
- vision problems
- seizure (convulsions)
- liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet
- or low white blood cell counts--fever, mouth sores, skin sores, sore throat, cough, trouble breathing. Common side effects of etoposide may include: sores or white patches in or around your mouth, trouble swallowing or talking, dry mouth, bad breath, altered sense of taste
- nausea , vomiting
- or temporary hair loss .

Drug Side Effects Summary

729. This particular disease, Cancer, interacts with the drug cyclophosphamide and has the following side effects:

- new worsening cough, shortness of breath, trouble breathing while lying down
- a light-headed feeling, like you might pass out
- pink or red urine, pain or burning when you urinate
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- any wound that will not heal
- heart problems--swelling, rapid weight gain, pounding heartbeats or fluttering in your chest
- liver problems--loss of appetite, stomach pain (upper right side), bloating , dark urine, jaundice (yellowing of the skin or eyes)
- low sodium level-- headache , confusion, slurred speech, severe weakness, vomiting , loss of coordination, feeling unsteady
- or low blood cell counts--fever, chills, tiredness, blisters or ulcers in your mouth, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath.

Common side effects of cyclophosphamide may include: fever, low blood cell counts

- mouth sores
- nausea , vomiting, diarrhea
- hair loss
- or missed menstrual periods.

730. This particular disease, Cancer, interacts with the drug vincristine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Vincristine may cause serious side effects. Call your doctor at once if you have: severe constipation, severe bloating or stomach pain, bloody or tarry stools

Drug Side Effects Summary

- swelling, rapid weight gain
- urination problems
- wheezing, trouble breathing
- chest pain or pressure, pain spreading to your jaw or shoulder
- problems with vision, hearing, speech, balance, or daily activities
- a seizure
- mouth pain or ulcers
- numbness , tingling, burning pain
- or low blood cell counts--fever, chills, sore throat , body aches, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath. Common side effects of vincristine may include: hair loss
- decreased weight with loss of muscle tissue
- nausea , vomiting , diarrhea , weight loss
- or rash.

731. This particular disease, Cancer, interacts with the drug Cosmegen and has the following side effects:

- swelling or tenderness in your stomach (upper right side), rapid weight gain
- swelling in your arms or legs
- dark urine, jaundice (yellowing of the skin or eyes)
- redness of your skin or the inside of your mouth and throat (if you are also receiving radiation treatment)
- sores or white patches in or around your mouth, trouble swallowing or talking, dry mouth, bad breath, altered sense of taste
- blisters, ulcers, or other skin changes where an injection was given
- or low blood cell counts--fever, chills, tiredness, mouth sores, skin sores, easy bruising, unusual

Drug Side Effects Summary

bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath. Common side effects of Cosmegen may include: hair loss

- fever, infections, low blood cell counts
- rash
- trouble swallowing
- nausea , vomiting
- or tiredness.

732. This particular disease, Cancer, interacts with the drug Etopophos and has the following side effects:

- pain, burning, irritation, or skin changes where the injection was given
- severe nausea and vomiting
- easy bruising, unusual bleeding, purple or red spots under your skin
- or low white blood cell counts--fever, mouth sores, skin sores, sore throat, cough, trouble breathing. Common side effects of Etopophos may include: nausea , vomiting
- constipation
- fever
- trouble swallowing
- or unusual or unpleasant taste in the mouth.

733. This particular disease, Cancer, interacts with the drug Ifex and has the following side effects:

- fever, chills, flu symptoms
- mouth and throat ulcers
- skin sores, pale skin, cold hands and feet
- easy bruising, unusual bleeding
- rapid heart rate, shallow breathing

Drug Side Effects Summary

- or feeling light-headed or short of breath. Call your doctor at once if you have any of these side effects: confusion, unusual thoughts or behavior, hallucinations, seizure (convulsions)
- little or no urination, painful or difficult urination, blood in your urine
- loss of bladder control
- a seizure
- muscle movements you cannot control
- problems with your hearing or vision, ringing in your ears
- sudden chest pain, wheezing, dry cough
- dark urine, jaundice (yellowing of the skin or eyes)
- or a wound that will not heal. Common side effects of Ifex may include: nausea and vomiting
- confusion, vision problems, thinking problems
- numbness , tingling, burning pain
- infections
- urination problems
- or hair loss .

734. This particular disease, Cancer, interacts with the drug Toposar and has the following side effects

- low white blood cell counts--fever, mouth sores, skin sores, sore throat, cough, trouble breathing
- easy bruising, unusual bleeding, purple or red spots under your skin
- severe ongoing nausea or vomiting
- sudden chest pain, wheezing, dry cough, feeling short of breath
- upper stomach pain, loss of appetite
- vision problems
- or a seizure (convulsions). Common side effects of Toposar may include: low white blood cell counts.

Drug Side Effects Summary

735. This particular disease, Cancer, interacts with the drug vinblastine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Vinblastine may cause serious side effects. Call your doctor at once if you have: wheezing, trouble breathing
- chest pain or pressure, pain spreading to your jaw or shoulder
- severe constipation
- ongoing nausea or vomiting
- numbness , tingling, burning pain
- problems with vision, hearing, speech, balance, or daily activities
- mouth pain or ulcers
- or low blood cell counts--fever, chills, sore throat , body aches, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath. Common side effects of vinblastine may include: low blood cell counts
- jaw pain, tumor pain, bone pain
- constipation
- or temporary hair loss .

736. This particular disease, Cancer, interacts with the drug Hycamtin and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Hycamtin may cause serious side effects. Call your doctor at once if you have: new or worsening cough, fever, trouble breathing
- diarrhea with fever and stomach cramps
- pain or burning when you urinate

Drug Side Effects Summary

- signs of pneumonia --fever, chills, cough with mucus, chest pain, feeling short of breath
- or low blood cell counts--fever, chills, flu-like symptoms, mouth sores, skin sores, pale skin, cold hands and feet, bruising or bleeding, feeling light-headed. Your cancer treatments may be delayed or permanently discontinued if you have certain side effects. Common side effects of Hycamtin may include: low blood cell counts
- trouble breathing, pneumonia nausea, diarrhea, vomiting, stomach pain
- loss of appetite
- hair loss
- or feeling weak or tired.

737. This particular disease, Cancer, interacts with the drug ifosfamide and has the following side effects:

- fever, chills, flu symptoms
- mouth and throat ulcers
- skin sores, pale skin, cold hands and feet
- easy bruising, unusual bleeding
- rapid heart rate, shallow breathing
- or feeling light-headed or short of breath. Call your doctor at once if you have any of these side effects: confusion, unusual thoughts or behavior, hallucinations, seizure (convulsions)
- little or no urination, painful or difficult urination, blood in your urine
- loss of bladder control
- a seizure
- muscle movements you cannot control
- problems with your hearing or vision, ringing in your ears
- sudden chest pain, wheezing, dry cough
- dark urine, jaundice (yellowing of the skin or eyes)

Drug Side Effects Summary

- or a wound that will not heal. Common side effects of ifosfamide may include: nausea and vomiting
- confusion, vision problems, thinking problems
- numbness , tingling, burning pain
- infections
- urination problems
- or hair loss .

738. This particular disease, Cancer, interacts with the drug dactinomycin and has the following side effects:

- swelling or tenderness in your stomach (upper right side), rapid weight gain
- swelling in your arms or legs
- dark urine, jaundice (yellowing of the skin or eyes)
- redness of your skin or the inside of your mouth and throat (if you are also receiving radiation treatment)
- sores or white patches in or around your mouth, trouble swallowing or talking, dry mouth, bad breath, altered sense of taste
- blisters, ulcers, or other skin changes where an injection was given
- or low blood cell counts--fever, chills, tiredness, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath. Common side effects of dactinomycin may include: hair loss
- fever, infections, low blood cell counts
- rash
- trouble swallowing
- nausea , vomiting
- or tiredness.

Drug Side Effects Summary

739. This particular disease, Cancer, interacts with the drug fludarabine and has the following side

- agitation
- sudden chest pain, wheezing, dry cough, feeling short of breath
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- damage to red blood cells--confusion, weakness, pale or yellowed skin, dark colored urine
- low blood cell counts--fever, chills, tiredness, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath
- or signs of tumor cell breakdown--tiredness, weakness, muscle cramps, lower back pain , nausea , vomiting , diarrhea , blood in your urine, fast or slow heart rate , tingling in your hands and feet or around your mouth. Common side effects of fludarabine may include: low blood cell counts
- nausea, loss of appetite, vomiting, diarrhea
- feeling weak or tired
- mouth sores
- or swelling in your hands or feet.

740. This particular disease, Cancer, interacts with the drug Tepadina and has the following side ef

- hives
- wheezing, difficult breathing
- swelling of your face, lips, tongue, or throat. Tepadina may cause serious side effects. Call your doctor at once if you have: sores or white patches in or around your mouth, trouble swallowing or talking, dry mouth, bad breath, altered sense of taste
- confusion, hallucinations
- headache , drowsiness, changes in behavior or personality
- problems with memory, speech, or thought
- a seizure

Drug Side Effects Summary

- twitching muscles, overactive reflexes, problems with coordination or movement
- blood in your urine
- low blood cell counts--fever, chills, tiredness, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath
- or liver problems--rapid weight gain, stomach pain and bloating , dark urine, jaundice (yellowing of the skin or eyes). Common side effects of Tepadina may include: low blood cell counts
- signs of infection (fever, chills, sore throat , muscle aches)
- blood in your urine
- mouth sores
- or abnormal liver function tests .

741. This particular disease, Cancer, interacts with the drug thiotepa and has the following side effects

- hives
- wheezing, difficult breathing
- swelling of your face, lips, tongue, or throat. Thiotepa may cause serious side effects. Call your doctor at once if you have: sores or white patches in or around your mouth, trouble swallowing or talking, dry mouth, bad breath, altered sense of taste
- confusion, hallucinations
- headache , drowsiness, changes in behavior or personality
- problems with memory, speech, or thought
- a seizure
- twitching muscles, overactive reflexes, problems with coordination or movement
- blood in your urine
- low blood cell counts--fever, chills, tiredness, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath

Drug Side Effects Summary

- or liver problems--rapid weight gain, stomach pain and bloating , dark urine, jaundice (yellowing of the skin or eyes). Common side effects of thiotepa may include: low blood cell counts
- signs of infection (fever, chills, sore throat , muscle aches)
- blood in your urine
- mouth sores
- or abnormal liver function tests .

742. This particular disease, Cancer, interacts with the drug topotecan and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Topotecan may cause serious side effects. Call your doctor at once if you have: new or worsening cough, fever, trouble breathing
- diarrhea with fever and stomach cramps
- pain or burning when you urinate
- signs of pneumonia --fever, chills, cough with mucus, chest pain, feeling short of breath
- or low blood cell counts--fever, chills, flu-like symptoms, mouth sores, skin sores, pale skin, cold hands and feet, bruising or bleeding, feeling light-headed. Your cancer treatments may be delayed or permanently discontinued if you have certain side effects. Common side effects of topotecan may include: low blood cell counts
- trouble breathing, pneumonia nausea, diarrhea, vomiting, stomach pain
- loss of appetite
- hair loss
- or feeling weak or tired.

743. This particular disease, Cholesterol, interacts with the drug atorvastatin and has the following

Drug Side Effects Summary

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, atorvastatin can cause a condition that results in the breakdown of skeletal muscle tissue, potentially leading to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, and dark colored urine. Atorvastatin may cause serious side effects. Call your doctor at once if you have: muscle weakness in your hips, shoulders, neck, and back trouble lifting your arms, trouble climbing or standing
- liver problems - upper stomach pain, weakness, tired feeling, loss of appetite, dark urine, jaundice (yellowing of the skin or eyes)
- or kidney problems - little or no urinating, swelling in your feet or ankles, feeling tired or short of breath. Common side effects of atorvastatin may include: joint pain
- stuffy nose
- sore throat
- diarrhea
- or pain in your arms or legs.

744. This particular disease, Cholesterol, interacts with the drug simvastatin and has the following

- signs of a kidney problem - little or no urinating
- painful or difficult urination
- swelling in your feet or ankles
- feeling tired or short of breath
- or muscle weakness in your hips, shoulders, neck, and back
- trouble lifting your arms, trouble climbing or standing
- or liver problems - nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine,

Drug Side Effects Summary

clay-colored stools, jaundice (yellowing of the skin or eyes). Common simvastatin side effects may include: headache

- constipation, nausea, stomach pain
- or cold symptoms such as stuffy nose, sneezing, sore throat.

745. This particular disease, Cholesterol, interacts with the drug Lipitor and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, Lipitor can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, and dark colored urine. Also call your doctor at once if you have: muscle weakness in your hips, shoulders, neck, and back
- trouble lifting your arms, trouble climbing or standing
- upper stomach pain, weakness, tired feeling, loss of appetite, dark urine, yellowing of the skin or eyes (signs of liver problems)
- or little or no urinating, swelling in your feet or ankles, feeling tired or short of breath (signs of kidney problems). Common Lipitor side effects may include: joint pain
- stuffy nose, sore throat
- diarrhea
- or pain in your arms or legs.

746. This particular disease, Cholesterol, interacts with the drug Zetia and has the following side effects:

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Zetia can cause the breakdown of muscle tissue, which can lead to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark urine. Side effects may be more likely in older adults. Common side effects of Zetia may include: diarrhea
- cold symptoms such as stuffy nose, sneezing, sore throat
- pain in an arm or leg
- joint pain
- or muscle pain while taking Zetia with a statin.

747. This particular disease, Cholesterol, interacts with the drug ezetimibe and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Ezetimibe can cause the breakdown of muscle tissue, which can lead to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark urine. Side effects may be more likely in older adults. Common ezetimibe side effects may include: diarrhea
- cold symptoms such as stuffy nose, sneezing, sore throat
- pain in an arm or leg
- joint pain
- or muscle pain while taking ezetimibe with a statin.

748. This particular disease, Cholesterol, interacts with the drug Crestor and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: unexplained

Drug Side Effects Summary

muscle pain, tenderness, or weakness

- muscle weakness in your hips, shoulders, neck, and back
- trouble lifting your arms, trouble climbing or standing
- confusion, memory problems
- or liver problems - upper stomach pain, tiredness, loss of appetite, dark urine, jaundice (yellowing of the skin or eyes). Common side effects may include: headache
- weakness
- muscle aches
- or nausea, stomach pain.

749. This particular disease, Cholesterol, interacts with the drug Repatha and has the following side effects:

- hives, severe itching
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: high blood sugar
- increased thirst, increased urination, dry mouth, fruity breath odor. Common Repatha side effects may include: redness, pain, or bruising where an injection was given
- back pain
- flu symptoms
- or cold symptoms such as stuffy nose, sneezing, sore throat.

750. This particular disease, Cholesterol, interacts with the drug alirocumab and has the following side effects:

- hives , severe rash and itching
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common alirocumab side effects may include: redness, itching, soreness, or swelling where an injection was given

Drug Side Effects Summary

- flu symptoms
- or cold symptoms such as stuffy nose, sneezing, sore throat .

751. This particular disease, Cholesterol, interacts with the drug evolocumab and has the following

- Evolocumab can cause serious side effects including: Serious Allergic Reactions . Some people taking evolocumab have had serious allergic reactions. Stop taking evolocumab and call your healthcare provider or seek emergency medical help right away if you or your child have any of these symptoms: trouble breathing or swallowing raised bumps (hives) rash, or itching swelling of the face, lips, tongue, throat or arms The most common side effects of evolocumab include: runny nose, sore throat, symptoms of the common cold, flu or flu-like symptoms, back pain, high blood sugar levels (diabetes) and redness, pain, or bruising at the injection site. Tell your healthcare provider if you or your child have any side effect that bothers you or that does not go away. These are not all the possible side effects of evolocumab. Ask your healthcare provider or pharmacist for more information. Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Evolocumab side effects

752. This particular disease, Cholesterol, interacts with the drug Praluent and has the following side

- hives , severe rash and itching
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common Praluent side effects may include: redness, itching, soreness, or swelling where an injection was given
- flu symptoms
- or cold symptoms such as stuffy nose, sneezing, sore throat .

753. This particular disease, Cholesterol, interacts with the drug rosuvastatin and has the following

Drug Side Effects Summary

- Serious side effects of rosuvastatin include: Muscle pain, tenderness and weakness (myopathy). Muscle problems, including muscle breakdown, can be serious in some people and rarely cause kidney damage that can lead to death. Tell your doctor right away if: you have unexplained muscle pain, tenderness, or weakness, especially if you have a fever or feel more tired than usual, while you take this medication. you have muscle problems that do not go away even after your doctor has told you to stop taking this medication. Your doctor may do further tests to diagnose the cause of your muscle problems. Your chances of getting muscle problems are higher if you: are taking certain other medicines while you take rosuvastatin are 65 years of age or older have thyroid problems (hypothyroidism) that are not controlled have kidney problems are taking higher doses of this medication Liver problems. Your doctor should do blood tests to check your liver before you start taking this medication and if you have symptoms of liver problems while you taking it. Call your doctor right away if you have any of the following symptoms of liver problems: feel unusually tired or weak loss of appetite upper belly pain dark urine yellowing of your skin or the whites of your eyes The most common side effects of rosuvastatin may include: headache muscle aches and pains abdominal pain weakness nausea Additional side effects that have been reported with rosuvastatin include memory loss and confusion. Tell your doctor if you have any side effect that bothers you or that does not go away. These are not all the possible side effects. For more information, ask your doctor or pharmacist. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Rosuvastatin side effects

754. This particular disease, Cholesterol, interacts with the drug pravastatin and has the following s

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, pravastatin can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure. Call your doctor right

Drug Side Effects Summary

away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark colored urine. Call your doctor right away if you have: muscle weakness in your hips, shoulders, neck, and back

- trouble lifting your arms, trouble climbing or standing
- or liver problems - loss of appetite, stomach pain (upper right side), tiredness, dark urine, jaundice (yellowing of the skin or eyes). Common pravastatin side effects may include: muscle or joint pain
- nausea , vomiting , diarrhea
- headache
- or cold symptoms such as stuffy nose, sneezing, sore throat .

755. This particular disease, Cholesterol, interacts with the drug lovastatin and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Lovastatin can cause the breakdown of muscle tissue, which can lead to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark colored urine. Also call your doctor at once if you have: muscle weakness in your hips, shoulders, neck, and back
- trouble lifting your arms, trouble climbing or standing
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, dark urine, jaundice (yellowing of the skin or eyes). Common side effects of lovastatin may include: infections
- headache
- or accidental injury.

756. This particular disease, Cholesterol, interacts with the drug Zocor and has the following side effects

Drug Side Effects Summary

- hives, difficult breathing, swelling in your face or throat) or a severe skin reaction (fever, sore throat, burning eyes, skin pain, red or purple skin rash with blistering and peeling). Simvastatin can cause the breakdown of muscle tissue, which can lead to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark urine. Also call your doctor at once if you have: signs of a kidney problem
- little or no urinating
- painful or difficult urination
- swelling in your feet or ankles
- feeling tired or short of breath
- or muscle weakness in your hips, shoulders, neck, and back
- trouble lifting your arms, trouble climbing or standing
- or liver problems - nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common Zocor side effects may include: headache
- constipation, nausea, stomach pain
- or cold symptoms such as stuffy nose, sneezing, sore throat.

757. This particular disease, Cholesterol, interacts with the drug gemfibrozil and has the following s

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, gemfibrozil can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure . Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, and dark colored urine. Also call your doctor at once if you have: sharp pain in your upper stomach (especially after eating)

Drug Side Effects Summary

- jaundice (yellowing of your skin or eyes)
- pain or burning when you urinate
- blurred vision, eye pain , or seeing halos around lights
- or low blood cell counts - pale skin, easy bruising, unusual bleeding, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Common gemfibrozil side effects may include: stomach pain, indigestion , diarrhea
- numbness or tingly feeling
- dizziness , drowsiness
- blurred vision
- muscle or joint pain
- or loss of interest in sex, impotence .

758. This particular disease, Cholesterol, interacts with the drug Vytorin and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: unexplained muscle pain, tenderness, or weakness
- muscle weakness in your hips, shoulders, neck, and back
- trouble lifting your arms, trouble climbing or standing
- fever, unusual tiredness
- upper stomach pain, loss of appetite
- dark colored urine
- or jaundice (yellowing of the skin or eyes). Common Vytorin side effects may include: headache
- muscle pain
- abnormal liver function tests

Drug Side Effects Summary

- diarrhea
- or cold symptoms such as stuffy nose, sneezing, sore throat .

759. This particular disease, Cholesterol, interacts with the drug Livalo and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, Livalo can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, and dark urine. Also call your doctor at once if you have: confusion, memory problems
- muscle weakness in your hips, shoulders, neck, and back
- trouble lifting your arms, trouble climbing or standing
- kidney problems - vomiting, pain in your side or lower back, little or no urinating, swelling, rapid weight gain
- or liver problems - loss of appetite, stomach pain (upper right side), tiredness, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common Livalo side effects may include: muscle pain
- pain in your arms or legs
- back pain
- or diarrhea, constipation.

760. This particular disease, Cholesterol, interacts with the drug niacin and has the following side effects:

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: heart attack symptoms - chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- unexplained muscle pain, tenderness or weakness
- a light-headed feeling, like you might pass out
- irregular heartbeats
- severe warmth or redness under your skin
- vision problems
- or jaundice (yellowing of the skin or eyes). Common niacin side effects may include: flushing (sudden warmth, redness, or tingly feeling)
- upset stomach , nausea, vomiting , diarrhea
- abnormal liver function tests
- itching, rash, dry skin
- cough
- skin discoloration
- or headache .

761. This particular disease, Cholesterol, interacts with the drug Altacor and has the following side

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using Altacor and call your doctor at once if you have any of these serious side effects: muscle pain, tenderness, or weakness with fever or flu symptoms and dark colored urine. Less serious side effects may include: mild stomach pain, gas, bloating, stomach upset, heartburn
- nausea

Drug Side Effects Summary

- constipation
- or diarrhea.

762. This particular disease, Cholesterol, interacts with the drug Niaspan and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have any of these serious side effects: feeling like you might pass out
- fast, pounding, or uneven heart beats
- feeling short of breath
- swelling
- jaundice (yellowing of your skin or eyes)
- or muscle pain, tenderness, or weakness with fever or flu symptoms and dark colored urine. If you are diabetic, tell your doctor about any changes in your blood sugar levels. Less serious Niaspan ER side effects include: mild dizziness
- warmth, redness, or tingly feeling under your skin
- itching, dry skin
- sweating or chills
- nausea, diarrhea, belching, gas
- muscle pain, leg cramps
- or sleep problems (insomnia).

763. This particular disease, Cholesterol, interacts with the drug Welchol and has the following side effects:

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Welchol may cause serious side effects. Stop using Welchol and call your doctor at once if you have: severe constipation

- severe stomach pain

- or pancreatitis--severe pain in your upper stomach spreading to your back, nausea and vomiting .

Common side effects of Welchol may include: constipation

- nausea

- or upset stomach .

764. This particular disease, Cholesterol, interacts with the drug Altoprev and has the following side effects:

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Altoprev can cause the breakdown of muscle tissue, which can lead to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark colored urine.

Also call your doctor at once if you have: muscle weakness in your hips, shoulders, neck, and back

- trouble lifting your arms, trouble climbing or standing

- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath

- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, dark urine, jaundice (yellowing of the skin or eyes). Common side effects of Altoprev may include: infections

- headache

- or accidental injury.

765. This particular disease, Cholesterol, interacts with the drug ezetimibe / simvastatin and has the following side effects:

- hives

- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. ezetimibe and simvastatin may cause serious side effects. Stop using ezetimibe and simvastatin and call your doctor at once if you have: unexplained muscle pain, tenderness, or weakness
- muscle weakness in your hips, shoulders, neck, and back
- trouble lifting your arms, trouble climbing or standing
- fever, unusual tiredness
- upper stomach pain, loss of appetite
- dark colored urine
- or jaundice (yellowing of the skin or eyes). Common side effects of ezetimibe and simvastatin may include: headache
- muscle pain
- abnormal liver function tests
- diarrhea
- or cold symptoms such as stuffy nose, sneezing, sore throat .

766. This particular disease, Cholesterol, interacts with the drug fenofibric acid and has the following

- sharp stomach pain spreading to your back or shoulder blade
- loss of appetite, stomach pain just after eating a meal
- jaundice (yellowing of the skin or eyes)
- fever, chills, weakness, sore throat, mouth sores, unusual bruising or bleeding
- chest pain, sudden cough, wheezing, rapid breathing, coughing up blood
- or swelling, warmth, or redness in an arm or leg. Common side effects of fenofibric acid may include: runny nose , sneezing
- or abnormal laboratory tests.

Drug Side Effects Summary

767. This particular disease, Cholesterol, interacts with the drug Slo-Niacin and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Darkening of urine light gray-colored stools loss of appetite severe stomach pain yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Abdominal or stomach pain cough diarrhea feeling of warmth flushing or redness of the skin, especially on the face and neck headache nausea or vomiting rash or itching runny nose sneezing stuffy nose Incidence not known Dizziness or faintness dryness of the skin fever frequent urination joint pain muscle aching or cramping side, lower back, or stomach pain swelling of the feet or lower legs unusual thirst unusual tiredness or weakness unusually fast, slow, or irregular heartbeat Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Slo-Niacin side effects

768. This particular disease, Cholesterol, interacts with the drug colesevelam and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Colesevelam may cause serious side effects. Stop using colesevelam and call your doctor at once if you have: severe constipation
- severe stomach pain

Drug Side Effects Summary

- or pancreatitis--severe pain in your upper stomach spreading to your back, nausea and vomiting .

Common side effects of colesevelam may include: constipation

- nausea
- or upset stomach .

769. This particular disease, Cholesterol, interacts with the drug Ezallor and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Ezallor Sprinkle may cause serious side effects. Call your doctor at once if you have: unexplained muscle pain, tenderness, or weakness
- muscle weakness in your hips, shoulders, neck, and back
- trouble lifting your arms, trouble climbing or standing
- confusion, memory problems
- or liver problems--upper stomach pain, tiredness, loss of appetite, dark urine, jaundice (yellowing of the skin or eyes). Common side effects of Ezallor Sprinkle may include: headache
- weakness
- muscle aches
- or nausea , stomach pain.

770. This particular disease, Cholesterol, interacts with the drug FloLipid and has the following side

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives

Drug Side Effects Summary

- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. A heartbeat that does not feel normal. Not able to pass urine or change in how much urine is passed. This medicine may cause muscle pain, tenderness, or weakness. Sometimes, a very bad muscle problem may happen that may lead to kidney problems. Rarely, deaths have happened in people who get these problems when taking drugs like this one. Call your doctor right away if you have muscle pain, tenderness, or weakness that is not normal (with or without fever or feeling out of sorts). Call your doctor right away if you have muscle signs that last after your doctor has told you to stop taking FloLipid (simvastatin oral suspension). Liver problems have happened with drugs like this one. Sometimes, this has been deadly. Call your doctor right away if you have signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes.

FloLipid side effects

771. This particular disease, Cholesterol, interacts with the drug Lopid and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, Lopid can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure . Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, and dark colored urine. Also call your doctor at once if you have: sharp pain in

Drug Side Effects Summary

your upper stomach (especially after eating)

- jaundice (yellowing of your skin or eyes)
- pain or burning when you urinate
- blurred vision, eye pain , or seeing halos around lights
- or low blood cell counts--pale skin, easy bruising, unusual bleeding, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Common side effects of Lopid may include: stomach pain, indigestion , diarrhea
- numbness or tingly feeling
- dizziness , drowsiness
- blurred vision
- muscle or joint pain
- or loss of interest in sex, impotence .

772. This particular disease, Cholesterol, interacts with the drug pitavastatin and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, pitavastatin can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark colored urine. Also call your doctor at once if you have: confusion, memory problems
- muscle weakness in your hips, shoulders, neck, and back
- trouble lifting your arms, trouble climbing or standing
- kidney problems-- vomiting , pain in your side or lower back, little or no urinating, swelling, rapid weight gain

Drug Side Effects Summary

- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of pitavastatin may include: muscle pain
- pain in your arms or legs
- back pain
- or diarrhea , constipation .

773. This particular disease, Cholesterol, interacts with the drug atorvastatin / ezetimibe and has the

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Abdominal or stomach pain confusion difficult breathing difficulty with moving irregular heartbeat muscle, joint, or bone pain muscle stiffness muscle weakness nausea or vomiting nervousness numbness or tingling in the hands, feet, or lips weakness or heaviness of the legs Incidence not known Black, tarry stools blistering, peeling, or loosening of the skin blood in the urine or stools chills dark-colored urine diarrhea fever gaseous abdominal or stomach pain general tiredness and weakness hives, welts, or skin rash itching large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs light-colored stools severe nausea or vomiting sores, ulcers, or white spots in the mouth or on the lips swelling of the eyes, face, or inside of the nose unsteadiness or awkwardness unusual bleeding or bruising unusual tiredness or weakness yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Cough dizziness feeling of warmth pain or tenderness around

Drug Side Effects Summary

the eyes and cheekbones redness of the face, neck, arms, and occasionally, upper chest stuffy or runny nose sudden sweating tightness in the chest Incidence not known Being forgetful bloating burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chills constipation discouragement fast heartbeat feeling sad or empty irritability lack of appetite loss of interest or pleasure loss of memory pains in the stomach, side, or abdomen, possibly radiating to the back tiredness trouble concentrating trouble sleeping Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Atorvastatin / ezetimibe side effects

774. This particular disease, Cholesterol, interacts with the drug fluvastatin and has the following s

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, fluvastatin can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark colored urine. Also call your doctor at once if you have: muscle weakness in your hips, shoulders, neck, and back
- trouble lifting your arms, trouble climbing or standing
- signs of a kidney problem--little or no urinating
- painful or difficult urination
- swelling in your feet or ankles
- feeling tired or short of breath
- or liver problems--upper stomach pain, loss of appetite, tiredness, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of fluvastatin may include: muscle

Drug Side Effects Summary

pain

- headache
- or stomach pain, nausea , or indigestion .

775. This particular disease, Cholesterol, interacts with the drug Lescol XL and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, Lescol XL can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark colored urine. Also call your doctor at once if you have: muscle weakness in your hips, shoulders, neck, and back
- trouble lifting your arms, trouble climbing or standing
- signs of a kidney problem--little or no urinating
- painful or difficult urination
- swelling in your feet or ankles
- feeling tired or short of breath
- or liver problems--upper stomach pain, loss of appetite, tiredness, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Lescol XL may include: muscle pain
- headache
- or stomach pain, nausea , or indigestion .

776. This particular disease, Cholesterol, interacts with the drug mycophenolate mofetil and has the following side effects:

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Mycophenolate mofetil may cause a serious brain infection that can lead to disability or death. Call your doctor right away if you have problems with speech, thought, vision, or muscle movement. These symptoms may start gradually and get worse quickly. Mycophenolate mofetil affects your immune system and may increase your risk of cancer or serious infection. Call your doctor right away if you have: fever of 100.5 degrees F or higher, swollen glands, painful mouth sores, cold or flu symptoms, headache , ear pain, loss of smell or taste, stomach pain, vomiting , diarrhea , weight loss
- weakness on one side of your body, loss of muscle control, confusion, thinking problems, loss of interest in things that normally interest you
- pain around the transplanted kidney, pain or burning when you urinate, dark urine, jaundice (yellowing of the skin or eyes)
- tingly or painful blistering rash on one side of your body, swelling, warmth, redness, or oozing around a skin wound
- or a new skin lesion, or a mole that has changed in size or color. Also call your doctor at once if you have: bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds, fever, joint stiffness and pain, and muscle pain
- or low blood cell counts--fever, chills, tiredness, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath. Common side effects of mycophenolate mofetil may include: stomach pain, nausea , vomiting, diarrhea, constipation , swelling in your ankles or feet, rash, pain anywhere in your body
- headache, dizziness , tremors, fever, sore throat , cold symptoms , or other signs of infections
- or abnormal blood tests, high blood sugar, high or low blood pressure , fast heart rate .

777. This particular disease, Cholesterol, interacts with the drug Niacor and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of

Drug Side Effects Summary

these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Darkening of urine light gray-colored stools loss of appetite severe stomach pain yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Abdominal or stomach pain cough diarrhea feeling of warmth flushing or redness of the skin, especially on the face and neck headache nausea or vomiting rash or itching runny nose sneezing stuffy nose Incidence not known Dizziness or faintness dryness of the skin fever frequent urination joint pain muscle aching or cramping side, lower back, or stomach pain swelling of the feet or lower legs unusual thirst unusual tiredness or weakness unusually fast, slow, or irregular heartbeat Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Niacor side effects

778. This particular disease, Cholesterol, interacts with the drug Trilipix and has the following side

- sharp stomach pain spreading to your back or shoulder blade
- loss of appetite, stomach pain just after eating a meal
- jaundice (yellowing of the skin or eyes)
- fever, chills, weakness, sore throat, mouth sores, unusual bruising or bleeding
- chest pain, sudden cough, wheezing, rapid breathing, coughing up blood
- or swelling, warmth, or redness in an arm or leg. Common Trilipix side effects may include: runny nose , sneezing
- or abnormal laboratory tests.

Drug Side Effects Summary

779. This particular disease, Cholesterol, interacts with the drug ezetimibe / rosuvastatin and has the following side effects:

- hives , rash, itching
- wheezing, dizziness , difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: unexplained muscle pain, tenderness, or weakness (especially if you also have fever)
- upper stomach pain, loss of appetite
- severe weakness or tiredness
- dark colored urine
- or jaundice (yellowing of the skin or eyes). Common side effects of ezetimibe and rosuvastatin may include: headache , dizziness
- stomach pain, nausea , diarrhea , constipation
- muscle or joint pain
- pain in your hands, legs, or back
- feeling weak or tired
- or cold symptoms such as stuffy nose, sneezing, sinus pain, sore throat .

780. This particular disease, Cholesterol, interacts with the drug Fibracor and has the following side effects:

- sharp stomach pain spreading to your back or shoulder blade
- loss of appetite, stomach pain just after eating a meal
- jaundice (yellowing of the skin or eyes)
- fever, chills, weakness, sore throat, mouth sores, unusual bruising or bleeding
- chest pain, sudden cough, wheezing, rapid breathing, coughing up blood
- or swelling, warmth, or redness in an arm or leg. Common side effects of Fibracor may include: runny nose , sneezing

Drug Side Effects Summary

- or abnormal laboratory tests.

781. This particular disease, Cholesterol, interacts with the drug inclisiran and has the following side effects:

- hives , severe itching
- difficult breathing
- swelling of your face, lips, tongue, or throat. Inclisiran may cause serious side effects. Call your doctor at once if you have: pain and burning when you urinate. Common side effects of inclisiran may include: redness, pain, or bruising where an injection was given
- pain in your legs and arms
- diarrhea
- breathing problems, bronchitis
- joint pain
- or pain and burning when you urinate.

782. This particular disease, Cholesterol, interacts with the drug Juxtapid and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have signs of liver problems , such as: nausea, vomiting, upper stomach pain
- loss of appetite
- swelling around your midsection
- flu-like symptoms, tiredness
- dark urine, clay-colored stools
- or jaundice (yellowing of the skin or eyes). Common Juxtapid side effects may include: stomach cramps

Drug Side Effects Summary

- nausea, vomiting
- indigestion
- or diarrhea.

783. This particular disease, Cholesterol, interacts with the drug Leqvio and has the following side

- Common side effect of Leqvio include: injection site reactions, such as pain, redness or a rash joint pain (arthralgia) urinary tract infections diarrhea bronchitis pain in extremities shortness of breath (dyspnea) These are not all of the possible side effects of this medication. Call your healthcare provider for medical advice about side effects. To report suspected adverse reactions, contact Novartis Pharmaceuticals Corporation at 1-888-669-6682 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch. Leqvio side effects

784. This particular disease, Cholesterol, interacts with the drug lomitapide and has the following s

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have signs of liver problems , such as: nausea, vomiting, upper stomach pain
- loss of appetite
- swelling around your midsection
- flu-like symptoms, tiredness
- dark urine, clay-colored stools
- or jaundice (yellowing of the skin or eyes). Common side effects of lomitapide may include:
stomach cramps
- nausea, vomiting
- indigestion

Drug Side Effects Summary

- or diarrhea.

785. This particular disease, Cholesterol, interacts with the drug Lypqozet and has the following side effects:

- No side effects listed

786. This particular disease, Cholesterol, interacts with the drug Roszet and has the following side effects:

- Roszet may cause serious side effects, including: Allergic reactions, including a severe reaction known as anaphylaxis. Tell your healthcare provider right away if you have any of the following symptoms: swelling of the face, lips, tongue, and/or throat which makes it difficult to swallow or breathe breathing problems or wheezing feeling dizzy or fainting rash or hives itching Muscle pain, tenderness and weakness (myopathy). Muscle problems, including muscle breakdown, can be serious in some people and rarely causes kidney damage that can lead to death. You have a higher chance for muscle problems if you are taking certain other medicines with Roszet. Tell your healthcare provider right away if: you have unexplained muscle pain, tenderness, or weakness, especially if you have a fever or feel more tired than usual, while you take Roszet. you have muscle problems that do not go away even after your healthcare provider has advised you to stop taking Roszet. Your healthcare provider may do further tests to diagnose the cause of your muscle problems. Your chances of getting muscle problems are higher if you: are taking certain other medicines while you take Roszet (see Especially tell your healthcare provider if you take medicines) are 65 years of age or older are of Asian descent have thyroid problems (hypothyroidism) that are not controlled have kidney problems are taking higher doses of Roszet Liver problems. Your healthcare provider may do blood tests to check your liver before you start treatment and if you have symptoms of liver problems while you take Roszet. Call your healthcare provider right away if you have the following symptoms of liver problems: feel tired or weak upper belly pain yellowing of your skin or the whites of your eyes loss of appetite dark urine The most common side effects include:

Drug Side Effects Summary

headache weakness diarrhea dizziness pain (back, hands, legs) nausea constipation joint pain stomach pain muscle aches and pains common cold and flu runny nose and sore throat tiredness

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects. Call your doctor for medical advice about side effects.

You may report side effects to FDA at 1-800-FDA-1088. Roszet side effects

787. This particular disease, Cholesterol, interacts with the drug Zypitamag and has the following s

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, Zypitamag can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark colored urine. Also call your doctor at once if you have: confusion, memory problems
- muscle weakness in your hips, shoulders, neck, and back
- trouble lifting your arms, trouble climbing or standing
- kidney problems-- vomiting , pain in your side or lower back, little or no urinating, swelling, rapid weight gain
- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Zypitamag may include: muscle pain
- pain in your arms or legs
- back pain
- or diarrhea , constipation .

Drug Side Effects Summary

788. This particular disease, Colds & Flu, interacts with the drug Benadryl and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Benadryl and call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- painful or difficult urination
- little or no urinating
- confusion, feeling like you might pass out
- or tightness in your neck or jaw, uncontrollable movements of your tongue. Common Benadryl side effects may include: dizziness, drowsiness, loss of coordination
- dry mouth, nose, or throat
- constipation, upset stomach
- dry eyes, blurred vision
- or day-time drowsiness or "hangover" feeling after night-time use.

789. This particular disease, Colds & Flu, interacts with the drug diphenhydramine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using diphenhydramine and call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- painful or difficult urination
- little or no urinating
- confusion, feeling like you might pass out
- or tightness in your neck or jaw, uncontrollable movements of your tongue. Common diphenhydramine side effects may include: dizziness, drowsiness, loss of coordination

Drug Side Effects Summary

- dry mouth, nose, or throat
- constipation, upset stomach
- dry eyes, blurred vision
- or day-time drowsiness or "hangover" feeling after night-time use.

790. This particular disease, Colds & Flu, interacts with the drug Vicks Nyquil D Cold and Flu Night

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- a light-headed feeling, like you might pass out
- anxiety , feeling nervous, trouble sleeping
- little or no urination
- or liver problems--nausea, upper stomach pain, itching, tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of NyQuil D may include: drowsiness
- constipation, diarrhea , upset stomach , vomiting
- or feeling restless or excited (especially in children).

Drug Side Effects Summary

791. This particular disease, Colds & Flu, interacts with the drug Vicks NyQuil Cold & Flu Nighttime

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: severe headache, seizure (convulsions)
- fast, pounding, or uneven heartbeats
- confusion, hallucinations, severe dizziness or drowsiness, slow or shallow breathing
- tremor, restless muscle movements
- little or no urinating
- flu symptoms, easy bruising, unusual bleeding, pale skin, weakness, feeling light-headed
- or nausea, pain in your upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Nyquil Cold & Flu may include: headache, sleep problems (insomnia)
- upset stomach , diarrhea or constipation
- dry eyes, blurred vision, dry mouth or nose
- mild dizziness or drowsiness, trouble concentrating
- feeling restless or excited (especially in children)
- or mild skin rash .

792. This particular disease, Colds & Flu, interacts with the drug chlorpheniramine and has the follo

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Chlorpheniramine may cause serious side effects. Stop using chlorpheniramine and call your doctor at once if you have: fast or uneven heart rate
- mood changes
- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- feeling short of breath
- or little or no urinating. Common side effects of chlorpheniramine may include: dizziness , drowsiness
- dry mouth, nose, or throat
- constipation
- blurred vision
- or feeling nervous or restless.

793. This particular disease, Colds & Flu, interacts with the drug Coricidin HBP Cold & Flu and has

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Coricidin and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. Stop using the medicine and call your doctor at once if you have: fast, pounding, or uneven heartbeats
- confusion, severe dizziness or drowsiness

Drug Side Effects Summary

- tremor, restless muscle movements
- little or no urinating
- flu symptoms, easy bruising, unusual bleeding, pale skin, feeling light-headed or short of breath
- or nausea, pain in your upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common Coricidin side effects may include: constipation
- dry eyes, blurred vision, dry mouth or nose
- mild dizziness or drowsiness, trouble concentrating
- feeling restless or excited (especially in children)
- or mild skin rash.

794. This particular disease, Colds & Flu, interacts with the drug Vicks DayQuil Severe Cold & Flu a

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: chest pain, fast, slow, or uneven heart rate
- severe dizziness , feeling like you might pass out
- mood changes, confusion, hallucinations
- tremor, seizure (convulsions)
- fever

Drug Side Effects Summary

- urinating less than usual or not at all
- nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of your skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety, chest pain, uneven heartbeats, seizure). Common side effects of DayQuil Severe Cold & Flu may include: dizziness, weakness
- mild headache
- mild nausea, diarrhea, upset stomach
- dry mouth, nose, or throat
- feeling nervous, restless, irritable, or anxious
- or sleep problems (insomnia).

795. This particular disease, Colds & Flu, interacts with the drug Benadryl Allergy and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad dizziness or passing out. Change in

Drug Side Effects Summary

balance. Feeling less alert. Benadryl Allergy side effects

796. This particular disease, Colds & Flu, interacts with the drug Vicks Dayquil Cold & Flu Relief and

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. Stop using DayQuil Cold & Flu and call your doctor at once if you have: chest pain, fast, slow, or uneven heart rate
- severe dizziness , feeling like you might pass out
- mood changes, confusion, hallucinations, seizure (convulsions)
- little or no urinating
- nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , chest pain, uneven heartbeats, seizure). Common side effects of DayQuil Cold & Flu may include: dizziness, weakness, mild headache
- diarrhea , upset stomach
- dry mouth, nose, or throat
- feeling nervous, restless, irritable, or anxious
- or sleep problems (insomnia).

797. This particular disease, Colds & Flu, interacts with the drug Vicks NyQuil Severe Cold & Flu and

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- a light-headed feeling, like you might pass out
- anxiety , hallucinations, feeling of fear
- little or no urinating
- pale skin, unusual weakness
- nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of your skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, chest pain, shortness of breath, uneven heartbeats, seizure). Common side effects of NyQuil Severe Cold & Flu may include: drowsiness
- constipation , diarrhea , upset stomach , vomiting
- or feeling restless or excited.

798. This particular disease, Colds & Flu, interacts with the drug Promethazine DM and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop

Drug Side Effects Summary

using this medicine and call your doctor at once if you have: weak or shallow breathing, breathing that stops during sleep

- unusual restlessness or excitement
- uncontrolled muscle movements in your neck, tongue, or eyes
- fast, slow, or uneven heart rate
- hallucinations, ringing in your ears
- a seizure
- fever, chills, weakness
- easy bruising, unusual bleeding
- yellowing of your skin or eyes
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Common side effects of Promethazine DM may include: drowsiness, dizziness
- dry mouth, nose, or throat
- blurred vision
- nausea , vomiting
- or loss of coordination.

799. This particular disease, Colds & Flu, interacts with the drug Tylenol Cold & Flu Severe and has

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction,

Drug Side Effects Summary

you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: chest pain, fast, slow, or uneven heart rate

- severe dizziness , feeling like you might pass out
- mood changes, confusion, hallucinations
- tremor, seizure (convulsions)
- fever
- urinating less than usual or not at all
- nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of your skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , chest pain, uneven heartbeats, seizure). Common side effects of Tylenol Cold & Flu Severe may include: dizziness, weakness
- mild headache
- mild nausea, diarrhea , upset stomach
- dry mouth, nose, or throat
- feeling nervous, restless, irritable, or anxious
- or sleep problems (insomnia).

800. This particular disease, Colds & Flu, interacts with the drug Vicks VapoRub and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Less serious side effects may be more likely, and you may have none at all.

Drug Side Effects Summary

801. This particular disease, Colds & Flu, interacts with the drug Coricidin HBP Maximum Strength

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: severe headache, seizure (convulsions)
- fast, pounding, or uneven heartbeats
- confusion, hallucinations, severe dizziness or drowsiness, slow or shallow breathing
- tremor, restless muscle movements
- little or no urination
- flu symptoms, easy bruising, unusual bleeding, pale skin, weakness, feeling light-headed
- or nausea, pain in your upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of Coricidin HBP Maximum Strength Flu may include: headache, sleep problems (insomnia)
- upset stomach , diarrhea or constipation
- dry eyes, blurred vision, dry mouth or nose
- mild dizziness or drowsiness, trouble concentrating
- feeling restless or excited (especially in children)
- or mild skin rash .

Drug Side Effects Summary

802. This particular disease, Colds & Flu, interacts with the drug Theraflu Nighttime Severe Cold & Cough

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe drowsiness
- painful or difficult urination
- dizziness , trouble sleeping, nervousness
- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Theraflu Severe Cold & Cough Nighttime may include: drowsiness
- dry eyes, blurred vision
- dry mouth, nose, or throat
- constipation
- or feeling restless or excited (especially in children).

803. This particular disease, Colds & Flu, interacts with the drug acetaminophen / chlorpheniramine

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past

Drug Side Effects Summary

and had no reaction. Stop taking acetaminophen, chlorpheniramine, and phenylephrine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate

- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- little or no urinating
- nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of your skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety, chest pain, shortness of breath, uneven heartbeats, seizure). Side effects such as dry mouth, constipation, and confusion may be more likely in older adults. Common side effects of acetaminophen, chlorpheniramine, and phenylephrine may include: dizziness, drowsiness
- mild headache, blurred vision
- dry mouth, nose, or throat
- constipation
- feeling nervous
- or sleep problems (insomnia)
-

804. This particular disease, Colds & Flu, interacts with the drug Sudafed PE Cold & Cough and has

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: chest pain, fast, slow, or uneven heart rate
- severe dizziness , feeling like you might pass out
- mood changes, confusion, hallucinations
- tremor, seizure (convulsions)
- fever
- urinating less than usual or not at all
- nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of your skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , chest pain, uneven heartbeats, seizure). Common side effects of Sudafed PE Cold & Cough may include: dizziness, weakness
- mild headache
- mild nausea, diarrhea , upset stomach
- dry mouth, nose, or throat
- feeling nervous, restless, irritable, or anxious
- or sleep problems (insomnia).

805. This particular disease, Colds & Flu, interacts with the drug Banophen and has the following s

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and

Drug Side Effects Summary

sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad dizziness or passing out. Change in balance. Feeling less alert. Banophen side effects

806. This particular disease, Colds & Flu, interacts with the drug Benadryl Allergy Plus Cold and ha

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness

Drug Side Effects Summary

- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Benadryl Allergy Plus Cold side effects

807. This particular disease, Colds & Flu, interacts with the drug pseudoephedrine / triprolidine and

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe dizziness
- nervousness
- trouble sleeping
- or high blood pressure (severe headache, pounding in your neck or ears). Common side effects of pseudoephedrine and triprolidine may include: dizziness, drowsiness
- dry mouth, nose, or throat
- constipation
- blurred vision
- or feeling restless or excited (especially in children).

808. This particular disease, Colds & Flu, interacts with the drug Tussionex Pennkinetic and has the

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Like other narcotic medications, hydrocodone can slow your breathing. Death may occur if breathing becomes too weak. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Stop using this medicine and call your doctor at once if you have:
severe drowsiness
- noisy breathing, sighing, shallow breathing
- severe constipation
- a light-headed feeling, like you might pass out
- a slow heart rate or weak pulse
- confusion
- or adrenal gland problems - nausea , vomiting , loss of appetite, dizziness, feeling weak or tired.

Side effects such as dry mouth, constipation, and confusion may be more likely in older adults.

Common Tussionex PennKinetic side effects may include: trouble breathing

- drowsiness
- confusion
- nausea, vomiting
- or little or no urination.

809. This particular disease, Colds & Flu, interacts with the drug Chlor-Trimeton and has the follow

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Chlor-Trimeton may cause serious side effects. Stop using Chlor-Trimeton and call your doctor at once if you have: fast or uneven heart rate
- mood changes

Drug Side Effects Summary

- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- feeling short of breath
- or little or no urinating. Common side effects of Chlor-Trimeton may include: dizziness , drowsiness
- dry mouth, nose, or throat
- constipation
- blurred vision
- or feeling nervous or restless.

810. This particular disease, Colds & Flu, interacts with the drug Diphenhydramine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Diphenhydramine may cause serious side effects. Stop using Diphenhydramine and call your doctor at once if you have: severe drowsiness
- or painful or difficult urination. Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of Diphenhydramine may include: drowsiness
- dry eyes, blurred vision
- dry mouth, nose, or throat
- decreased urination
- constipation
- feeling restless or excited (especially in children)
- or day-time drowsiness or "hangover" feeling after night-time use.

811. This particular disease, Colds & Flu, interacts with the drug Diphenhydramine and has the following side effects:

- No side effects listed

Drug Side Effects Summary

812. This particular disease, Colds & Flu, interacts with the drug Complete Allergy Relief and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad dizziness or passing out. Change in balance. Feeling less alert. Complete Allergy Relief side effects

813. This particular disease, Colds & Flu, interacts with the drug Tylenol Cold and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat

Drug Side Effects Summary

- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of high blood pressure like very bad headache or dizziness , passing out, or change in eyesight. Chest pain or pressure or a fast heartbeat . A heartbeat that does not feel normal. Very nervous and excitable. Shortness of breath. Shakiness. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Tylenol Cold side effects

814. This particular disease, Colds & Flu, interacts with the drug Allermax and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Allermax may cause serious side effects. Stop using Allermax and call your doctor at once if you have: severe drowsiness
- or painful or difficult urination. Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of Allermax may include: drowsiness
- dry eyes, blurred vision
- dry mouth, nose, or throat
- decreased urination
- constipation

Drug Side Effects Summary

- feeling restless or excited (especially in children)
- or day-time drowsiness or "hangover" feeling after night-time use.

815. This particular disease, Colds & Flu, interacts with the drug Bronkaid and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
 - hives
 - itching
 - red, swollen, blistered, or peeling skin with or without fever
 - wheezing
 - tightness in the chest or throat
 - trouble breathing, swallowing, or talking
 - unusual hoarseness
 - or swelling of the mouth, face, lips, tongue, or throat. Signs of high blood pressure like very bad headache or dizziness , passing out, or change in eyesight. Chest pain or pressure or a fast heartbeat . Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight. Very nervous and excitable. Seizures .
- Bronkaid side effects

816. This particular disease, Colds & Flu, interacts with the drug Dicopanorl and has the following side effects:

- No side effects listed

817. This particular disease, Colds & Flu, interacts with the drug Dytuss and has the following side effects:

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Dytuss may cause serious side effects. Stop using Dytuss and call your doctor at once if you have: severe drowsiness
- or painful or difficult urination. Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of Dytuss may include: drowsiness
- dry eyes, blurred vision
- dry mouth, nose, or throat
- decreased urination
- constipation
- feeling restless or excited (especially in children)
- or day-time drowsiness or "hangover" feeling after night-time use.

818. This particular disease, Colds & Flu, interacts with the drug Mucinex Fast-Max Night Time Cold

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe drowsiness
- painful or difficult urination
- dizziness , trouble sleeping, nervousness

Drug Side Effects Summary

- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Mucinex Fast-Max Night Time Cold and Flu may include: drowsiness

- dry eyes, blurred vision
- dry mouth, nose, or throat
- constipation
- or feeling restless or excited (especially in children).

819. This particular disease, Colds & Flu, interacts with the drug Scot-Tussin Allergy and has the fo

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Scot-Tussin Allergy Relief Formula may cause serious side effects. Stop using Scot-Tussin Allergy Relief Formula and call your doctor at once if you have: severe drowsiness
- or painful or difficult urination. Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of Scot-Tussin Allergy Relief Formula may include: drowsiness
- dry eyes, blurred vision
- dry mouth, nose, or throat
- decreased urination
- constipation
- feeling restless or excited (especially in children)
- or day-time drowsiness or "hangover" feeling after night-time use.

820. This particular disease, Colds & Flu, interacts with the drug Siladryl Allergy and has the follow

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Siladryl Allergy may cause serious side effects. Stop using Siladryl Allergy and call your doctor at once if you have: severe drowsiness
- or painful or difficult urination. Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of Siladryl Allergy may include: drowsiness
- dry eyes, blurred vision
- dry mouth, nose, or throat
- decreased urination
- constipation
- feeling restless or excited (especially in children)
- or day-time drowsiness or "hangover" feeling after night-time use.

821. This particular disease, Colds & Flu, interacts with the drug Sudafed PE Sinus & Allergy and h

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using Sudafed PE Sinus & Allergy and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate
- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)
- little or no urinating
- easy bruising or bleeding, unusual weakness
- or dangerously high blood pressure (severe headache, buzzing in your ears, anxiety , shortness of breath). Common side effects of Sudafed PE Sinus & Allergy may include: dizziness , drowsiness, blurred vision

Drug Side Effects Summary

- dry nose or mouth
- nausea , stomach pain, constipation , loss of appetite
- problems with memory or concentration
- or feeling restless or excited (especially in children).

822. This particular disease, Colds & Flu, interacts with the drug Twilite and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Twilite may cause serious side effects. Stop using Twilite and call your doctor at once if you have: severe drowsiness
- or painful or difficult urination. Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of Twilite may include: drowsiness
- dry eyes, blurred vision
- dry mouth, nose, or throat
- decreased urination
- constipation
- feeling restless or excited (especially in children)
- or day-time drowsiness or "hangover" feeling after night-time use.

823. This particular disease, Colds & Flu, interacts with the drug Tylenol Cold Multi-Symptom Severe

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin

Drug Side Effects Summary

redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: chest pain, fast, slow, or uneven heart rate

- severe dizziness , feeling like you might pass out
- mood changes, confusion, hallucinations
- tremor, seizure (convulsions)
- fever
- urinating less than usual or not at all
- nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of your skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , chest pain, uneven heartbeats, seizure). Common side effects of Tylenol Cold Multi-Symptom Severe may include: dizziness, weakness
- mild headache
- mild nausea, diarrhea , upset stomach
- dry mouth, nose, or throat
- feeling nervous, restless, irritable, or anxious
- or sleep problems (insomnia).

824. This particular disease, Colds & Flu, interacts with the drug Valu-Dryl and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Valu-Dryl may cause serious side effects. Stop using Valu-Dryl and call your doctor at once if you have: severe drowsiness

Drug Side Effects Summary

- or painful or difficult urination. Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of Valu-Dryl may include: drowsiness

- dry eyes, blurred vision
- dry mouth, nose, or throat
- decreased urination
- constipation
- feeling restless or excited (especially in children)
- or day-time drowsiness or "hangover" feeling after night-time use.

825. This particular disease, Colds & Flu, interacts with the drug Vanamine PD and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Vanamine PD may cause serious side effects. Stop using Vanamine PD and call your doctor at once if you have: severe drowsiness
- or painful or difficult urination. Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of Vanamine PD may include: drowsiness
- dry eyes, blurred vision
- dry mouth, nose, or throat
- decreased urination
- constipation
- feeling restless or excited (especially in children)
- or day-time drowsiness or "hangover" feeling after night-time use.

826. This particular disease, Colds & Flu, interacts with the drug Virtussin DAC and has the following

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Like other narcotic medications, codeine can slow your breathing. Death may occur if breathing becomes too weak. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Stop taking this medicine and call your doctor at once if you have: noisy breathing, sighing, shallow breathing
- severe drowsiness, hallucinations
- fast, slow, or uneven heart rate
- a seizure
- nervousness , tremors
- or little or no urination. Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Common side effects of Virtussin DAC may include: dizziness, drowsiness, weakness
- headache
- nausea , constipation
- sleep problems (insomnia)
- or feeling nervous, anxious, or restless.

827. This particular disease, Colds & Flu, interacts with the drug Benadryl Allergy Plus Sinus Headache

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Benadryl Allergy Sinus Headache and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this

Drug Side Effects Summary

type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: fast, slow, or uneven heart rate

- confusion, hallucinations
- tremor, seizure (convulsions)
- little or no urinating
- nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety, confusion, chest pain, shortness of breath, seizure). Common side effects of Benadryl Allergy Sinus Headache may include: dizziness, weakness

- mild headache
- mild nausea, diarrhea, upset stomach
- runny nose
- feeling nervous, restless, or anxious
- or sleep problems (insomnia).

828. This particular disease, Colds & Flu, interacts with the drug Primatene and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever

Drug Side Effects Summary

- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of high blood pressure like very bad headache or dizziness , passing out, or change in eyesight. Chest pain or pressure or a fast heartbeat . Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight. Very nervous and excitable. Seizures .

Primatene side effects

829. This particular disease, Colds & Flu, interacts with the drug acetaminophen / dextromethorpha

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- a light-headed feeling, like you might pass out
- anxiety , feeling nervous, trouble sleeping
- little or no urination
- or liver problems--nausea, upper stomach pain, itching, tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Side effects such as dry mouth,

Drug Side Effects Summary

constipation , and confusion may be more likely in older adults. Common side effects of acetaminophen, dextromethorphan, doxylamine, and pseudoephedrine may include: drowsiness

- constipation, diarrhea , upset stomach , vomiting
- or feeling restless or excited (especially in children).

830. This particular disease, Colds & Flu, interacts with the drug acetaminophen / dextromethorphan

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: chest pain, fast, slow, or uneven heart rate
- severe dizziness , feeling like you might pass out
- mood changes, confusion, hallucinations
- tremor, seizure (convulsions)
- fever
- urinating less than usual or not at all
- nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of your skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , chest pain, uneven heartbeats, seizure). Common side effects of acetaminophen, dextromethorphan, guaifenesin, and phenylephrine may include: dizziness, weakness

Drug Side Effects Summary

- mild headache
- mild nausea, diarrhea , upset stomach
- dry mouth, nose, or throat
- feeling nervous, restless, irritable, or anxious
- or sleep problems (insomnia).

831. This particular disease, Colds & Flu, interacts with the drug acetaminophen / dextromethorpha

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. Stop using acetaminophen, dextromethorphan, and phenylephrine and call your doctor at once if you have:
- chest pain, fast, slow, or uneven heart rate
- severe dizziness , feeling like you might pass out
- mood changes, confusion, hallucinations, seizure (convulsions)
- little or no urinating
- nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , chest pain, uneven heartbeats, seizure). Common side effects of acetaminophen, dextromethorphan, and phenylephrine may include: dizziness, weakness, mild headache
- diarrhea , upset stomach

Drug Side Effects Summary

- dry mouth, nose, or throat
- feeling nervous, restless, irritable, or anxious
- or sleep problems (insomnia).

832. This particular disease, Colds & Flu, interacts with the drug Aller-Chlor and has the following s

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Aller-Chlor may cause serious side effects. Stop using Aller-Chlor and call your doctor at once if you have: fast or uneven heart rate
- mood changes
- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- feeling short of breath
- or little or no urinating. Common side effects of Aller-Chlor may include: dizziness , drowsiness
- dry mouth, nose, or throat
- constipation
- blurred vision
- or feeling nervous or restless.

833. This particular disease, Colds & Flu, interacts with the drug Allergy Relief and has the followin

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Allergy Relief may cause serious side effects. Stop using Allergy Relief and call your doctor at once if you have: fast or uneven heart rate
- mood changes

Drug Side Effects Summary

- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- feeling short of breath
- or little or no urinating. Common side effects of Allergy Relief may include: dizziness , drowsiness
- dry mouth, nose, or throat
- constipation
- blurred vision
- or feeling nervous or restless.

834. This particular disease, Colds & Flu, interacts with the drug brompheniramine and has the following side effects:

- No side effects listed

835. This particular disease, Colds & Flu, interacts with the drug Mucinex Fast-Max Severe Cold and Flu Medicine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: chest pain, fast, slow, or uneven heart rate
- severe dizziness , feeling like you might pass out
- mood changes, confusion, hallucinations
- tremor, seizure (convulsions)

Drug Side Effects Summary

- fever
- urinating less than usual or not at all
- nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of your skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , chest pain, uneven heartbeats, seizure). Common side effects of Mucinex Fast-Max Severe Cold may include: dizziness, weakness
- mild headache
- mild nausea, diarrhea , upset stomach
- dry mouth, nose, or throat
- feeling nervous, restless, irritable, or anxious
- or sleep problems (insomnia).

836. This particular disease, Colds & Flu, interacts with the drug Mucinex Fast-Max Severe Congest

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness

Drug Side Effects Summary

- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)

- red or irritated eyes

- or sores in your mouth, throat, nose, or eyes. Mucinex Fast-Max Severe Congestion & Cold side effects

837. This particular disease, Colds & Flu, interacts with the drug Sudafed PE Pressure+Pain+Cold a

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

- itching

- red, swollen, blistered, or peeling skin with or without fever

- wheezing

- tightness in the chest or throat

- trouble breathing, swallowing, or talking

- unusual hoarseness

- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (

Drug Side Effects Summary

Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)

- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Sudafed PE Pressure+Pain+Cold side effects

838. This particular disease, Colds & Flu, interacts with the drug Sudafed PE Severe Cold and has t

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe drowsiness
- painful or difficult urination
- dizziness , trouble sleeping, nervousness
- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Sudafed PE Severe Cold may include: drowsiness
- dry eyes, blurred vision
- dry mouth, nose, or throat
- constipation
- or feeling restless or excited (especially in children).

Drug Side Effects Summary

839. This particular disease, Colds & Flu, interacts with the drug Tylenol Cold Max Day and has the

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. Tylenol Cold Multi-Symptom Daytime may cause serious side effects. Stop using Tylenol Cold Multi-Symptom Daytime and call your doctor at once if you have: chest pain, fast, slow, or uneven heart rate
- severe dizziness , feeling like you might pass out
- mood changes, confusion, hallucinations, seizure (convulsions)
- little or no urinating
- nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , chest pain, uneven heartbeats, seizure). Common side effects of Tylenol Cold Multi-Symptom Daytime may include: dizziness, weakness, mild headache
- diarrhea , upset stomach
- dry mouth, nose, or throat
- feeling nervous, restless, irritable, or anxious
- or sleep problems (insomnia).

840. This particular disease, Colds & Flu, interacts with the drug Vicks NyQuil Cold & Flu Nighttime

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: severe headache, seizure (convulsions)
- fast, pounding, or uneven heartbeats
- confusion, hallucinations, severe dizziness or drowsiness, slow or shallow breathing
- tremor, restless muscle movements
- little or no urination
- flu symptoms, easy bruising, unusual bleeding, pale skin, weakness, feeling light-headed
- or nausea, pain in your upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of NyQuil Cold & Flu (Alcohol Free) may include: headache, sleep problems (insomnia)
- upset stomach , diarrhea or constipation
- dry eyes, blurred vision, dry mouth or nose
- mild dizziness or drowsiness, trouble concentrating
- feeling restless or excited (especially in children)
- or mild skin rash .

841. This particular disease, Colds & Flu, interacts with the drug Wal-finate and has the following side effects:

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Wal-finate may cause serious side effects. Stop using Wal-finate and call your doctor at once if you have: fast or uneven heart rate
- mood changes
- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- feeling short of breath
- or little or no urinating. Common side effects of Wal-finate may include: dizziness , drowsiness
- dry mouth, nose, or throat
- constipation
- blurred vision
- or feeling nervous or restless.

842. This particular disease, Colds & Flu, interacts with the drug acetaminophen / chlorpheniramine

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: any new or worsening symptoms
- severe drowsiness
- fever for longer than 3 days

Drug Side Effects Summary

- redness, swelling, or rash
- ongoing headache
- or pain, cough, or sinus congestion lasting more than 7 days
- Some side effects may be more likely in older adults. Common side effects of acetaminophen, chlorpheniramine, dextromethorphan, and pseudoephedrine may include: dizziness , drowsiness
- dry mouth, nose, or throat
- blurred vision
- constipation
- feeling restless or excited (especially in children)
- or sleep problems (insomnia).

843. This particular disease, Colds & Flu, interacts with the drug acetaminophen / chlorpheniramine

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking acetaminophen and chlorpheniramine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: fast, pounding, or uneven heartbeats
- confusion, severe dizziness or drowsiness
- tremor, restless muscle movements
- little or no urinating
- flu symptoms, easy bruising, unusual bleeding, pale skin, feeling light-headed or short of breath

Drug Side Effects Summary

- or nausea, pain in your upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of acetaminophen and chlorpheniramine may include: constipation
- dry eyes, blurred vision, dry mouth or nose
- mild dizziness or drowsiness, trouble concentrating
- feeling restless or excited (especially in children)
- or mild skin rash .

844. This particular disease, Colds & Flu, interacts with the drug Actifed Cold and Allergy and has t

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using Actifed Cold & Allergy and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate
- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)
- little or no urinating
- easy bruising or bleeding, unusual weakness
- or dangerously high blood pressure (severe headache, buzzing in your ears, anxiety , shortness of breath). Common side effects of Actifed Cold & Allergy may include: dizziness , drowsiness, blurred vision
- dry nose or mouth
- nausea , stomach pain, constipation , loss of appetite
- problems with memory or concentration
- or feeling restless or excited (especially in children).

Drug Side Effects Summary

845. This particular disease, Colds & Flu, interacts with the drug Advil Allergy Sinus and has the following side effects:

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. Stop using this medicine and call your doctor at once if you have: severe dizziness , trouble sleeping, or nervousness
- shortness of breath (even with mild exertion)
- swelling or rapid weight gain
- a skin rash, no matter how mild
- severe headache, blurred vision, pounding in your neck or ears
- signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems - loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath. Common Advil Allergy Sinus side effects may include: upset stomach , mild heartburn , nausea , vomiting
- bloating , gas, diarrhea , constipation
- dizziness, headache, nervousness
- flushing (warmth, redness, or tingly feeling)
- mild itching or rash
- or ringing in your ears.

846. This particular disease, Colds & Flu, interacts with the drug Alka-Seltzer Plus Cold Formula and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop

Drug Side Effects Summary

using this medicine and call your doctor at once if you have: bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds

- chest pain, rapid pulse, fast or uneven heart rate
- ringing in your ears, severe headache, confusion, hallucinations, severe nervousness , feeling like you might pass out
- tremor, seizure (convulsions)
- feeling short of breath, little or no urinating
- redness or swelling, new cold or flu symptoms
- easy bruising or bleeding, unusual weakness
- or dangerously high blood pressure (severe headache, buzzing in your ears, anxiety , shortness of breath). Common side effects of Alka-Seltzer Plus Cold Formula may include: dizziness , drowsiness, blurred vision
- dry mouth, nose, or throat
- heartburn, nausea , diarrhea , constipation , upset stomach
- feeling nervous, restless, or irritable
- or sleep problems (insomnia).

847. This particular disease, Colds & Flu, interacts with the drug Aprodine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe dizziness
- nervousness
- trouble sleeping
- or high blood pressure (severe headache, pounding in your neck or ears). Common side effects of

Drug Side Effects Summary

Aprodine may include: dizziness, drowsiness

- dry mouth, nose, or throat
- constipation
- blurred vision
- or feeling restless or excited (especially in children).

848. This particular disease, Colds & Flu, interacts with the drug aspirin / chlorpheniramine / pheny

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- chest pain, rapid pulse, fast or uneven heart rate
- ringing in your ears, severe headache, confusion, hallucinations, severe nervousness , feeling like you might pass out
- tremor, seizure (convulsions)
- feeling short of breath, little or no urinating
- redness or swelling, new cold or flu symptoms
- easy bruising or bleeding, unusual weakness
- or dangerously high blood pressure (severe headache, buzzing in your ears, anxiety , shortness of breath). Common side effects of aspirin, chlorpheniramine, and phenylephrine may include: dizziness , drowsiness, blurred vision
- dry mouth, nose, or throat
- heartburn, nausea , diarrhea , constipation , upset stomach
- feeling nervous, restless, or irritable

Drug Side Effects Summary

- or sleep problems (insomnia).

849. This particular disease, Colds & Flu, interacts with the drug C.P.M. and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using C.P.M. and call your doctor at once if you have: fast or uneven heart rate
- mood changes
- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- feeling short of breath
- or little or no urinating. Common side effects of C.P.M. may include: dizziness , drowsiness
- dry mouth, nose, or throat
- constipation
- blurred vision
- or feeling nervous or restless.

850. This particular disease, Colds & Flu, interacts with the drug camphor / eucalyptus / menthol and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Less serious side effects may be more likely, and you may have none at all.

851. This particular disease, Colds & Flu, interacts with the drug Cheratussin DAC and has the following side effects:

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Like other narcotic medications, codeine can slow your breathing. Death may occur if breathing becomes too weak. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Stop taking this medicine and call your doctor at once if you have: noisy breathing, sighing, shallow breathing
- severe drowsiness, hallucinations
- fast, slow, or uneven heart rate
- a seizure
- nervousness , tremors
- or little or no urination. Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Common side effects of Cheratussin DAC may include: dizziness, drowsiness, weakness
- headache
- nausea , constipation
- sleep problems (insomnia)
- or feeling nervous, anxious, or restless.

852. This particular disease, Colds & Flu, interacts with the drug Chlo-Amine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Chlo-Amine may cause serious side effects. Stop using Chlo-Amine and call your doctor at once if you have: fast or uneven heart rate
- mood changes
- tremor, seizure (convulsions)

Drug Side Effects Summary

- easy bruising or bleeding, unusual weakness
- feeling short of breath
- or little or no urinating. Common side effects of Chlo-Amine may include: dizziness , drowsiness
- dry mouth, nose, or throat
- constipation
- blurred vision
- or feeling nervous or restless.

853. This particular disease, Colds & Flu, interacts with the drug Chlor-Mal and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Chlor-Mal may cause serious side effects. Stop using Chlor-Mal and call your doctor at once if you have: fast or uneven heart rate
- mood changes
- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- feeling short of breath
- or little or no urinating. Common side effects of Chlor-Mal may include: dizziness , drowsiness
- dry mouth, nose, or throat
- constipation
- blurred vision
- or feeling nervous or restless.

854. This particular disease, Colds & Flu, interacts with the drug chlorpheniramine / hydrocodone and has the following side effects:

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Like other narcotic medications, hydrocodone can slow your breathing. Death may occur if breathing becomes too weak. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe drowsiness
- noisy breathing, sighing, shallow breathing
- severe constipation
- a light-headed feeling, like you might pass out
- a slow heart rate or weak pulse
- confusion
- or adrenal gland problems-- nausea , vomiting , loss of appetite, dizziness, feeling weak or tired. Side effects such as dry mouth, constipation, and confusion may be more likely in older adults. Common side effects of chlorpheniramine and hydrocodone may include: trouble breathing
- drowsiness
- confusion
- nausea, vomiting
- or little or no urination.

855. This particular disease, Colds & Flu, interacts with the drug chlorpheniramine / phenylephrine

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. chlorpheniramine and phenylephrine may cause serious side effects. Stop using chlorpheniramine and phenylephrine and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate

Drug Side Effects Summary

- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)
- little or no urinating
- easy bruising or bleeding, unusual weakness
- or dangerously high blood pressure (severe headache, buzzing in your ears, anxiety , shortness of breath). Common side effects of chlorpheniramine and phenylephrine may include: dizziness , drowsiness, blurred vision
- dry nose or mouth
- nausea , stomach pain, constipation , loss of appetite
- problems with memory or concentration
- or feeling restless or excited (especially in children).

856. This particular disease, Colds & Flu, interacts with the drug ChlorTan and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. ChlorTan may cause serious side effects. Stop using ChlorTan and call your doctor at once if you have: fast or uneven heart rate
- mood changes
- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- feeling short of breath
- or little or no urinating. Common side effects of ChlorTan may include: dizziness , drowsiness
- dry mouth, nose, or throat
- constipation
- blurred vision

Drug Side Effects Summary

- or feeling nervous or restless.

857. This particular disease, Colds & Flu, interacts with the drug codeine / guaifenesin / pseudoeph

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Like other narcotic medications, codeine can slow your breathing. Death may occur if breathing becomes too weak. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Stop taking this medicine and call your doctor at once if you have: noisy breathing, sighing, shallow breathing
- severe drowsiness, hallucinations
- fast, slow, or uneven heart rate
- a seizure
- nervousness , tremors
- or little or no urination. Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Common side effects of codeine, guaifenesin, and pseudoephedrine may include: dizziness, drowsiness, weakness
- headache
- nausea , constipation
- sleep problems (insomnia)
- or feeling nervous, anxious, or restless.

858. This particular disease, Colds & Flu, interacts with the drug Coricidin HBP Nighttime Multi-Sym

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: severe headache, seizure (convulsions)
- fast, pounding, or uneven heartbeats
- confusion, hallucinations, severe dizziness or drowsiness, slow or shallow breathing
- tremor, restless muscle movements
- little or no urinating
- flu symptoms, easy bruising, unusual bleeding, pale skin, weakness, feeling light-headed
- or nausea, pain in your upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Coricidin HBP Nighttime Multi-Symptom Cold may include: headache, sleep problems (insomnia)
- upset stomach , diarrhea or constipation
- dry eyes, blurred vision, dry mouth or nose
- mild dizziness or drowsiness, trouble concentrating
- feeling restless or excited (especially in children)
- or mild skin rash .

859. This particular disease, Colds & Flu, interacts with the drug dextromethorphan / promethazine

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop

Drug Side Effects Summary

using this medicine and call your doctor at once if you have: weak or shallow breathing, breathing that stops during sleep

- unusual restlessness or excitement
- uncontrolled muscle movements in your neck, tongue, or eyes
- fast, slow, or uneven heart rate
- hallucinations, ringing in your ears
- a seizure
- fever, chills, weakness
- easy bruising, unusual bleeding
- yellowing of your skin or eyes
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Common side effects of dextromethorphan and promethazine may include: drowsiness, dizziness
- dry mouth, nose, or throat
- blurred vision
- nausea , vomiting
- or loss of coordination.

860. This particular disease, Colds & Flu, interacts with the drug diphenhydramine / phenylephrine

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe drowsiness
- sleeplessness
- painful or difficult urination

Drug Side Effects Summary

- fast, pounding, or uneven heartbeats
- trouble breathing
- anxiety , nervousness
- a seizure
- or problems with balance or speech. Common side effects of diphenhydramine and phenylephrine may include: dizziness , drowsiness, headache
- dry mouth, nose, or throat
- nausea , vomiting, constipation
- blurred vision
- or trouble sleeping.

861. This particular disease, Colds & Flu, interacts with the drug Drixoral Cold and Allergy and has

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: severe nervousness
- a light-headed feeling, like you might pass out
- trouble sleeping
- fast or uneven heart rate
- or painful or difficult urination. Side effects may be more likely in older adults. Common Drixoral Cold and Allergy side effects may include: drowsiness
- feeling restless or excited (especially in children)
- blurred vision
- or dry nose or mouth.

Drug Side Effects Summary

862. This particular disease, Colds & Flu, interacts with the drug Ed-Chlortan and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Ed-Chlortan may cause serious side effects. Stop using Ed-Chlortan and call your doctor at once if you have: fast or uneven heart rate
- mood changes
- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- feeling short of breath
- or little or no urinating. Common side effects of Ed-Chlortan may include: dizziness , drowsiness
- dry mouth, nose, or throat
- constipation
- blurred vision
- or feeling nervous or restless.

863. This particular disease, Colds & Flu, interacts with the drug Ed A-Hist and has the following side

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Ed A-Hist may cause serious side effects. Stop using Ed A-Hist and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate
- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)
- little or no urinating
- easy bruising or bleeding, unusual weakness
- or dangerously high blood pressure (severe headache, buzzing in your ears, anxiety , shortness of

Drug Side Effects Summary

breath). Common side effects of Ed A-Hist may include: dizziness , drowsiness, blurred vision

- dry nose or mouth
- nausea , stomach pain, constipation , loss of appetite
- problems with memory or concentration
- or feeling restless or excited (especially in children).

864. This particular disease, Colds & Flu, interacts with the drug Ed ChlorPed and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Ed ChlorPed may cause serious side effects. Stop using Ed ChlorPed and call your doctor at once if you have: fast or uneven heart rate
- mood changes
- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- feeling short of breath
- or little or no urinating. Common side effects of Ed ChlorPed may include: dizziness , drowsiness
- dry mouth, nose, or throat
- constipation
- blurred vision
- or feeling nervous or restless.

865. This particular disease, Colds & Flu, interacts with the drug Ed Chlor Ped Jr. and has the follow

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Ed-Chlor Ped Jr may cause serious side effects. Stop

Drug Side Effects Summary

using Ed-Chlor Ped Jr and call your doctor at once if you have: fast or uneven heart rate

- mood changes
- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- feeling short of breath
- or little or no urinating. Common side effects of Ed-Chlor Ped Jr may include: dizziness , drowsiness
- dry mouth, nose, or throat
- constipation
- blurred vision
- or feeling nervous or restless.

866. This particular disease, Colds & Flu, interacts with the drug ephedrine / guaifenesin and has the

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of high blood pressure like very bad

Drug Side Effects Summary

headache or dizziness , passing out, or change in eyesight. Chest pain or pressure or a fast heartbeat . Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight. Very nervous and excitable. Seizures .
Ephedrine / guaifenesin side effects

867. This particular disease, Colds & Flu, interacts with the drug Robitussin Peak Cold Nighttime N

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Robitussin Nighttime Nasal Relief and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate
- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- little or no urinating
- nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of your skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , chest pain, shortness of breath, uneven heartbeats, seizure). Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of Robitussin Nighttime Nasal Relief may include: dizziness , drowsiness

Drug Side Effects Summary

- mild headache, blurred vision
- dry mouth, nose, or throat
- constipation
- feeling nervous
- or sleep problems (insomnia)
-

868. This particular disease, Colds & Flu, interacts with the drug Theraflu Daytime Severe Cold & Cough

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. Stop using Theraflu Daytime Severe Cold & Cough and call your doctor at once if you have: chest pain, fast, slow, or uneven heart rate
- severe dizziness , feeling like you might pass out
- mood changes, confusion, hallucinations, seizure (convulsions)
- little or no urinating
- nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , chest pain, uneven heartbeats, seizure). Common side effects of Theraflu Daytime Severe Cold & Cough may include: dizziness, weakness, mild headache

Drug Side Effects Summary

- diarrhea , upset stomach
- dry mouth, nose, or throat
- feeling nervous, restless, irritable, or anxious
- or sleep problems (insomnia).

869. This particular disease, Colds & Flu, interacts with the drug TussiCaps and has the following s

- hives
 - difficult breathing
 - swelling of your face, lips, tongue, or throat. Like other narcotic medications, hydrocodone can slow your breathing. Death may occur if breathing becomes too weak. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe drowsiness
 - noisy breathing, sighing, shallow breathing
 - severe constipation
 - a light-headed feeling, like you might pass out
 - a slow heart rate or weak pulse
 - confusion
 - or adrenal gland problems-- nausea , vomiting , loss of appetite, dizziness, feeling weak or tired.
- Side effects such as dry mouth, constipation, and confusion may be more likely in older adults.
- Common side effects of TussiCaps may include: trouble breathing
- drowsiness
 - confusion
 - nausea, vomiting
 - or little or no urination.

Drug Side Effects Summary

870. This particular disease, Colds & Flu, interacts with the drug Tylenol Cold Multi-Symptom Nighttime

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

- itching

- red, swollen, blistered, or peeling skin with or without fever

- wheezing

- tightness in the chest or throat

- trouble breathing, swallowing, or talking

- unusual hoarseness

- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)

- red or irritated eyes

- or sores in your mouth, throat, nose, or eyes. Tylenol Cold Multi-Symptom Nighttime side effects

871. This particular disease, Colds & Flu, interacts with the drug A-Phedrin and has the following side effects

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop

Drug Side Effects Summary

using this medicine and call your doctor at once if you have: severe dizziness

- nervousness
- trouble sleeping
- or high blood pressure (severe headache, pounding in your neck or ears). Common side effects of A-Phedrin may include: dizziness, drowsiness
- dry mouth, nose, or throat
- constipation
- blurred vision
- or feeling restless or excited (especially in children).

872. This particular disease, Colds & Flu, interacts with the drug acetaminophen / chlorpheniramine

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (

Drug Side Effects Summary

Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)

- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Acetaminophen / chlorpheniramine / dextromethorphan / phenylephrine side effects

873. This particular disease, Colds & Flu, interacts with the drug acetaminophen / dextromethorphan

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- a light-headed feeling, like you might pass out
- anxiety , hallucinations, feeling of fear
- little or no urinating
- pale skin, unusual weakness
- nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of your skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, chest pain, shortness of breath, uneven heartbeats, seizure). Common side effects of acetaminophen,

Drug Side Effects Summary

dextromethorphan, doxylamine, and phenylephrine may include: drowsiness

- constipation , diarrhea , upset stomach , vomiting
- or feeling restless or excited.

874. This particular disease, Colds & Flu, interacts with the drug acetaminophen / dextromethorphan

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: severe headache, seizure (convulsions)
- fast, pounding, or uneven heartbeats
- confusion, hallucinations, severe dizziness or drowsiness, slow or shallow breathing
- tremor, restless muscle movements
- little or no urinating
- flu symptoms, easy bruising, unusual bleeding, pale skin, weakness, feeling light-headed
- or nausea, pain in your upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of acetaminophen, dextromethorphan, and doxylamine may include: headache, sleep problems (insomnia)
- upset stomach , diarrhea or constipation
- dry eyes, blurred vision, dry mouth or nose
- mild dizziness or drowsiness, trouble concentrating

Drug Side Effects Summary

- feeling restless or excited (especially in children)
- or mild skin rash .

875. This particular disease, Colds & Flu, interacts with the drug acetaminophen / dextromethorpha

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: chest pain
- fast, slow, or uneven heart rate
- severe dizziness , feeling like you might pass out
- mood changes, confusion, hallucinations, tremor, seizure (convulsions)
- fever
- little or no urinating
- nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of your skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , chest pain, uneven heartbeats, seizure). Common side effects of acetaminophen, dextromethorphan, guaifenesin, and pseudoephedrine may include: dizziness, weakness
- mild headache
- mild nausea, diarrhea , upset stomach
- dry mouth, nose, or throat

Drug Side Effects Summary

- feeling nervous, restless, irritable, or anxious
- or sleep problems (insomnia).

876. This particular disease, Colds & Flu, interacts with the drug acetaminophen / dextromethorpha

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Acetaminophen / dextromethorphan / pseudoephedrine side effects

Drug Side Effects Summary

877. This particular disease, Colds & Flu, interacts with the drug acetaminophen / diphenhydramine

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe drowsiness
- painful or difficult urination
- dizziness , trouble sleeping, nervousness
- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of acetaminophen, diphenhydramine, and phenylephrine may include: drowsiness
- dry eyes, blurred vision
- dry mouth, nose, or throat
- constipation
- or feeling restless or excited (especially in children).

878. This particular disease, Colds & Flu, interacts with the drug acetaminophen / pheniramine / ph

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past

Drug Side Effects Summary

and had no reaction. Stop taking acetaminophen, pheniramine, and phenylephrine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate

- confusion, hallucinations, tremor, severe dizziness or anxiety
- slow, shallow breathing
- little or no urinating
- easy bruising or bleeding, unusual weakness
- nausea, pain in your upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety, confusion, chest pain, shortness of breath, uneven heartbeats, seizure). Common side effects of acetaminophen, pheniramine, and phenylephrine may include: dryness of the eyes, nose, and mouth
- blurred vision
- dizziness, drowsiness, problems with memory or concentration
- ringing in your ears
- feeling restless or irritable
- or mild nausea, stomach pain, constipation.

879. This particular disease, Colds & Flu, interacts with the drug Advil Congestion Relief and has the

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs

Drug Side Effects Summary

of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight. Shortness of breath, a big weight gain, or swelling in the arms or legs. Chest pain or pressure. Very bad dizziness or passing out. Any unexplained bruising or bleeding. Very bad belly pain. Black, tarry, or bloody stools. Throwing up blood or throw up that looks like coffee grounds. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Advil Congestion Relief side effects

880. This particular disease, Colds & Flu, interacts with the drug Advil Sinus Congestion & Pain and

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. Stop using Advil Sinus Congestion & Pain and call your doctor at once if you have: severe dizziness or nervousness , trouble sleeping
- swelling, rapid weight gain
- shortness of breath

Drug Side Effects Summary

- new or worsening stomach pain
- or signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Common side effects of Advil Sinus Congestion & Pain may include: heartburn , stomach pain
- dizziness
- sleep problems (insomnia)
- or feeling nervous or excited.

881. This particular disease, Colds & Flu, interacts with the drug Ala-Hist PE and has the following :

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: fast or irregular heartbeats
- confusion, hallucinations
- a seizure
- ringing in your ears
- or little or no urination. Common side effects of Ala Hist PE may include: headache, dizziness , drowsiness
- blurred vision, double vision
- dry mouth, nose, or throat
- constipation , upset stomach
- or trouble sleeping.

882. This particular disease, Colds & Flu, interacts with the drug Alka-Seltzer Plus-D Multi-Symptom

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and

Drug Side Effects Summary

sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of high blood pressure like very bad headache or dizziness , passing out, or change in eyesight. Chest pain or pressure or a fast heartbeat . A heartbeat that does not feel normal. Very nervous and excitable. Shortness of breath. Shakiness. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Alka-Seltzer Plus-D Multi-Symptom Sinus & Cold Liquid Gels side effects

883. This particular disease, Colds & Flu, interacts with the drug Allergy Multi-Symptom and has the

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and

Drug Side Effects Summary

sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Allergy Multi-Symptom side effects

884. This particular disease, Colds & Flu, interacts with the drug Biotussin DAC and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Like other narcotic medications, codeine can slow your breathing. Death may occur if breathing becomes too weak. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you

Drug Side Effects Summary

are hard to wake up. Stop taking this medicine and call your doctor at once if you have: noisy breathing, sighing, shallow breathing

- severe drowsiness, hallucinations
- fast, slow, or uneven heart rate
- a seizure
- nervousness , tremors
- or little or no urination. Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Common side effects of Biotussin DAC may include: dizziness, drowsiness, weakness
- headache
- nausea , constipation
- sleep problems (insomnia)
- or feeling nervous, anxious, or restless.

885. This particular disease, Colds & Flu, interacts with the drug Bromatan Plus and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: fast, slow, or uneven heart rate
- severe headache, mood changes, hallucinations
- severe dizziness or anxiety , feeling like you might pass out
- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- fever
- little or no urinating

Drug Side Effects Summary

- shortness of breath
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety, confusion, chest pain, uneven heartbeats, seizure). Common side effects of Bromatan Plus may include: mild headache
- mild dizziness, drowsiness
- dry mouth, nose, or throat
- nausea , diarrhea , constipation , upset stomach
- feeling nervous, restless, or irritable
- blurred vision
- or sleep problems (insomnia).

886. This particular disease, Colds & Flu, interacts with the drug Bromax and has the following side effects:

- No side effects listed

887. This particular disease, Colds & Flu, interacts with the drug camphor and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking

Drug Side Effects Summary

- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Camphor topical side effects

888. This particular disease, Colds & Flu, interacts with the drug Capron DM and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Capron DM may cause serious side effects. Stop using Capron DM and call your doctor at once if you have: severe drowsiness or anxiety
- little or no urinating
- or a light-headed feeling, like you might pass out. Common side effects of Capron DM may include: drowsiness
- nervousness
- or feeling restless or excited (especially in children).

889. This particular disease, Colds & Flu, interacts with the drug Children's Triacting Night Time and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe drowsiness
- sleeplessness
- painful or difficult urination
- fast, pounding, or uneven heartbeats
- trouble breathing
- anxiety , nervousness
- a seizure

Drug Side Effects Summary

- or problems with balance or speech. Common side effects of Children's Triacting Night Time may include: dizziness , drowsiness, headache
- dry mouth, nose, or throat
- nausea , vomiting, constipation
- blurred vision
- or trouble sleeping.

890. This particular disease, Colds & Flu, interacts with the drug Childrens Tylenol Cold Plus Cough

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of high blood pressure like very bad headache or dizziness , passing out, or change in eyesight. Chest pain or pressure or a fast heartbeat . A heartbeat that does not feel normal. Very nervous and excitable. Shortness of breath. Shakiness. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal

Drug Side Effects Summary

necrosis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)

- red or irritated eyes

- or sores in your mouth, throat, nose, or eyes. Childrens Tylenol Cold Plus Cough side effects

891. This particular disease, Colds & Flu, interacts with the drug Chlo Hist and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

- itching

- red, swollen, blistered, or peeling skin with or without fever

- wheezing

- tightness in the chest or throat

- trouble breathing, swallowing, or talking

- unusual hoarseness

- or swelling of the mouth, face, lips, tongue, or throat. Feeling very sleepy. Chlo Hist side effects

892. This particular disease, Colds & Flu, interacts with the drug chlophedianol / pyrilamine and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

Drug Side Effects Summary

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Chlorpheniramine / pseudoephedrine side effects

893. This particular disease, Colds & Flu, interacts with the drug chlorpheniramine / ibuprofen / pseudoephedrine

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe dizziness, trouble sleeping, or nervousness
- shortness of breath (even with mild exertion)
- swelling or rapid weight gain
- a skin rash, no matter how mild
- severe headache, blurred vision, pounding in your neck or ears
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath. Common side effects of chlorpheniramine, ibuprofen, and pseudoephedrine may include: upset stomach, mild heartburn, nausea, vomiting

Drug Side Effects Summary

- bloating , gas, diarrhea , constipation
- dizziness, headache, nervousness
- flushing (warmth, redness, or tingly feeling)
- mild itching or rash
- or ringing in your ears.

894. This particular disease, Colds & Flu, interacts with the drug chlorpheniramine / phenylephrine

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: confusion, hallucinations
- feeling nervous or irritable
- trouble sleeping
- little or no urination
- a seizure (convulsions)
- numbness or tingly feeling
- a light-headed feeling, like you might pass out
- pounding heartbeats or fluttering in your chest
- or increased blood pressure--severe headache, blurred vision, pounding in your neck or ears, anxiety , nosebleed. Side effects may be more likely in older adults. Common side effects of chlorpheniramine, phenylephrine, and pyrilamine may include: nausea , constipation , loss of appetite
- rash or itching
- increased sweating
- dry mouth, nose, or throat

Drug Side Effects Summary

- blurred vision
- headache, dizziness
- or feeling restless or excited (especially in children).

895. This particular disease, Colds & Flu, interacts with the drug Cold Multi-Symptom (Nighttime) and

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Cold Multi-Symptom (Nighttime) side effects

Drug Side Effects Summary

896. This particular disease, Colds & Flu, interacts with the drug Comtrex Cold and Cough Maximum Strength

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

- itching

- red, swollen, blistered, or peeling skin with or without fever

- wheezing

- tightness in the chest or throat

- trouble breathing, swallowing, or talking

- unusual hoarseness

- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)

- red or irritated eyes

- or sores in your mouth, throat, nose, or eyes. Comtrex Cold and Cough Maximum Strength side effects

897. This particular disease, Colds & Flu, interacts with the drug Comtrex Cold and Flu Maximum Strength

- hives

- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: any new or worsening symptoms

- severe drowsiness

- fever for longer than 3 days

- redness, swelling, or rash

- ongoing headache

- or pain, cough, or sinus congestion lasting more than 7 days

- Some side effects may be more likely in older adults. Common side effects of Comtrex Cold and Flu Maximum Strength Tablet may include: dizziness , drowsiness

- dry mouth, nose, or throat

- blurred vision

- constipation

- feeling restless or excited (especially in children)

- or sleep problems (insomnia).

898. This particular disease, Colds & Flu, interacts with the drug Contac Cold + Flu (Night Formula)

- hives

- difficulty breathing

- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past

Drug Side Effects Summary

and had no reaction. Stop taking Contac Cold+Flu Maximum Strength and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate

- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- little or no urinating
- nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of your skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety, chest pain, shortness of breath, uneven heartbeats, seizure). Side effects such as dry mouth, constipation, and confusion may be more likely in older adults. Common side effects of Contac Cold+Flu Maximum Strength may include: dizziness, drowsiness
- mild headache, blurred vision
- dry mouth, nose, or throat
- constipation
- feeling nervous
- or sleep problems (insomnia)
-

899. This particular disease, Colds & Flu, interacts with the drug Delsym Cough+ Cold Day Time and

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: chest pain, fast, slow, or uneven heart rate

- severe dizziness , feeling like you might pass out

- mood changes, confusion, hallucinations

- tremor, seizure (convulsions)

- fever

- urinating less than usual or not at all

- nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of your skin or eyes)

- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , chest pain, uneven heartbeats, seizure). Common side effects of Delsym Cough Plus Cold Day Time may include: dizziness, weakness

- mild headache

- mild nausea, diarrhea , upset stomach

- dry mouth, nose, or throat

- feeling nervous, restless, irritable, or anxious

- or sleep problems (insomnia).

900. This particular disease, Colds & Flu, interacts with the drug Delsym Cough+ Cold Night Time a

- hives

Drug Side Effects Summary

- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe drowsiness
- painful or difficult urination
- dizziness , trouble sleeping, nervousness
- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Delsym Cough Plus Cold Night Time may include: drowsiness
- dry eyes, blurred vision
- dry mouth, nose, or throat
- constipation
- or feeling restless or excited (especially in children).

901. This particular disease, Colds & Flu, interacts with the drug Deltuss DMX and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching

Drug Side Effects Summary

- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Deltuss DMX side effects

902. This particular disease, Colds & Flu, interacts with the drug dextbrompheniramine / pseudoeph

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe nervousness
- a light-headed feeling, like you might pass out
- trouble sleeping
- fast or uneven heart rate
- or painful or difficult urination. Side effects may be more likely in older adults. Common side effects of dextbrompheniramine and pseudoephedrine may include: drowsiness
- feeling restless or excited (especially in children)
- blurred vision
- or dry nose or mouth.

903. This particular disease, Colds & Flu, interacts with the drug dextromethorphan / phenylephrine

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs

Drug Side Effects Summary

of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Dextromethorphan / phenylephrine / pyrilamine side effects

904. This particular disease, Colds & Flu, interacts with the drug dextromethorphan / phenylephrine

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Dextromethorphan / phenylephrine / triprolidine side effects

Drug Side Effects Summary

905. This particular disease, Colds & Flu, interacts with the drug Diabetic Tussin Night Time Formu

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: fast, pounding, or uneven heartbeats
- severe drowsiness
- painful or difficult urination
- severe headache, seizure
- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Diabetic Tussin Night Time Formula may include: drowsiness
- headache, sleep problems (insomnia)
- upset stomach , diarrhea or constipation
- dry eyes, blurred vision
- dry mouth, nose, or throat
- or feeling restless or excited (especially in children).

906. This particular disease, Colds & Flu, interacts with the drug Dologen and has the following sid

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if

Drug Side Effects Summary

you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Dologen side effects

907. This particular disease, Colds & Flu, interacts with the drug doxylamine / phenylephrine and h

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe dizziness
- nervousness , trouble sleeping
- little or no urination

Drug Side Effects Summary

- or high blood pressure (severe headache, pounding in your neck or ears). Common side effects of doxylamine and phenylephrine may include: dizziness, drowsiness
- dry mouth, nose, or throat
- constipation
- blurred vision
- or feeling restless or excited (especially in children).

908. This particular disease, Colds & Flu, interacts with the drug Dristan Cold Multi Symptom Form

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Dristan Cold Multi Symptom Formula and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate
- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- little or no urinating
- nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of your skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety, chest pain, shortness of breath, uneven heartbeats, seizure). Side effects such as dry mouth,

Drug Side Effects Summary

constipation , and confusion may be more likely in older adults. Common side effects of Dristan Cold

Multi Symptom Formula may include: dizziness , drowsiness

- mild headache, blurred vision
- dry mouth, nose, or throat
- constipation
- feeling nervous
- or sleep problems (insomnia)
-

909. This particular disease, Colds & Flu, interacts with the drug Flutabs and has the following side

- No side effects listed

910. This particular disease, Colds & Flu, interacts with the drug ibuprofen / phenylephrine and has

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. ibuprofen and phenylephrine may cause serious side effects. Stop using ibuprofen and phenylephrine and call your doctor at once if you have: severe dizziness or nervousness , trouble sleeping

- swelling, rapid weight gain
- shortness of breath
- new or worsening stomach pain
- or signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Common side effects of ibuprofen and phenylephrine may include: heartburn , stomach pain
- dizziness
- sleep problems (insomnia)

Drug Side Effects Summary

- or feeling nervous or excited.

911. This particular disease, Colds & Flu, interacts with the drug Mucinex Children's Cold, Cough &

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: chest pain, fast, slow, or uneven heart rate
- severe dizziness , feeling like you might pass out
- mood changes, confusion, hallucinations
- tremor, seizure (convulsions)
- fever
- urinating less than usual or not at all
- nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of your skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , chest pain, uneven heartbeats, seizure). Common side effects of Mucinex Children's Cold, Cough & Sore Throat may include: dizziness, weakness
- mild headache
- mild nausea, diarrhea , upset stomach
- dry mouth, nose, or throat

Drug Side Effects Summary

- feeling nervous, restless, irritable, or anxious
- or sleep problems (insomnia).

912. This particular disease, Colds & Flu, interacts with the drug Ninjacof and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Ninjacof side effects

913. This particular disease, Colds & Flu, interacts with the drug Norel AD and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever

Drug Side Effects Summary

- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Norel AD side effects

914. This particular disease, Colds & Flu, interacts with the drug phenylephrine / promethazine and

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: seizure (convulsions)
- weak or shallow breathing
- jaundice (yellowing of the skin or eyes)
- confusion, hallucinations
- sudden weakness or ill feeling, fever, chills, sore throat , mouth sores, red or swollen gums, trouble swallowing
- uncontrollable movements of your eyes, lips, tongue, face, arms, or legs
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast

Drug Side Effects Summary

or uneven heartbeats, tremors, feeling like you might pass out. Common side effects of phenylephrine and promethazine may include: dizziness , drowsiness

- dry mouth, nose, or throat
- blurred vision
- or constipation.

915. This particular disease, Colds & Flu, interacts with the drug Poly-Tussin AC and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Like other narcotic medications, codeine can slow your breathing. Death may occur if breathing becomes too weak. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Stop taking this medicine and call your doctor at once if you have: noisy breathing, sighing, shallow breathing
- a slow heart rate or weak pulse
- nervousness , trouble sleeping
- severe constipation
- or little or no urination. Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Common side effects of Poly-Tussin AC may include: drowsiness
- or feeling restless or excited (especially in children). Side effects such as dry mouth, constipation, and confusion may be more likely in older adults.

916. This particular disease, Colds & Flu, interacts with the drug Poly Hist Forte and has the following

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe dizziness
- nervousness , trouble sleeping
- little or no urination
- or high blood pressure (severe headache, pounding in your neck or ears). Common side effects of Poly Hist Forte may include: dizziness, drowsiness
- dry mouth, nose, or throat
- constipation
- blurred vision
- or feeling restless or excited (especially in children).

917. This particular disease, Colds & Flu, interacts with the drug Robitussin Peak Cold Nighttime C

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: severe headache, seizure (convulsions)
- fast, pounding, or uneven heartbeats
- confusion, hallucinations, severe dizziness or drowsiness, slow or shallow breathing
- tremor, restless muscle movements

Drug Side Effects Summary

- little or no urinating
- flu symptoms, easy bruising, unusual bleeding, pale skin, weakness, feeling light-headed
- or nausea, pain in your upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Robitussin Peak Cold Nighttime Cold + Flu may include: headache, sleep problems (insomnia)
- upset stomach , diarrhea or constipation
- dry eyes, blurred vision, dry mouth or nose
- mild dizziness or drowsiness, trouble concentrating
- feeling restless or excited (especially in children)
- or mild skin rash .

918. This particular disease, Colds & Flu, interacts with the drug Ru-Hist Forte and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: confusion, hallucinations
- feeling nervous or irritable
- trouble sleeping
- little or no urination
- a seizure (convulsions)
- numbness or tingly feeling
- a light-headed feeling, like you might pass out
- pounding heartbeats or fluttering in your chest
- or increased blood pressure--severe headache, blurred vision, pounding in your neck or ears, anxiety , nosebleed. Side effects may be more likely in older adults. Common side effects of Ru-Hist

Drug Side Effects Summary

Forte may include: nausea , constipation , loss of appetite

- rash or itching
- increased sweating
- dry mouth, nose, or throat
- blurred vision
- headache, dizziness
- or feeling restless or excited (especially in children).

919. This particular disease, Colds & Flu, interacts with the drug Sinus & Allergy PE and has the fol

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using Sinus & Allergy PE and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate
- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)
- little or no urinating
- easy bruising or bleeding, unusual weakness
- or dangerously high blood pressure (severe headache, buzzing in your ears, anxiety , shortness of breath). Common side effects of Sinus & Allergy PE may include: dizziness , drowsiness, blurred vision
- dry nose or mouth
- nausea , stomach pain, constipation , loss of appetite
- problems with memory or concentration
- or feeling restless or excited (especially in children).

Drug Side Effects Summary

920. This particular disease, Colds & Flu, interacts with the drug Sudafed PE Head Congestion + Pa

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight. Shortness of breath, a big weight gain, or swelling in the arms or legs. Chest pain or pressure. Very bad dizziness or passing out. Any unexplained bruising or bleeding. Very bad belly pain. Black, tarry, or bloody stools. Throwing up blood or throw up that looks like coffee grounds. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Sudafed PE Head Congestion + Pain side effects

921. This particular disease, Colds & Flu, interacts with the drug Sudafed PE Pressure+Pain+Cough

- hives

Drug Side Effects Summary

- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. Stop using Sudafed PE Pressure+Pain+Cough and call your doctor at once if you have: chest pain, fast, slow, or uneven heart rate
- severe dizziness , feeling like you might pass out
- mood changes, confusion, hallucinations, seizure (convulsions)
- little or no urinating
- nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , chest pain, uneven heartbeats, seizure). Common side effects of Sudafed PE Pressure+Pain+Cough may include: dizziness, weakness, mild headache
- diarrhea , upset stomach
- dry mouth, nose, or throat
- feeling nervous, restless, irritable, or anxious
- or sleep problems (insomnia).

922. This particular disease, Colds & Flu, interacts with the drug Theraflu Flu & Sore Throat and has

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe

Drug Side Effects Summary

skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Theraflu Flu & Sore Throat and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate

- confusion, hallucinations, tremor, severe dizziness or anxiety
- slow, shallow breathing
- little or no urinating
- easy bruising or bleeding, unusual weakness
- nausea, pain in your upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety, confusion, chest pain, shortness of breath, uneven heartbeats, seizure). Common side effects of Theraflu Flu & Sore Throat may include: dryness of the eyes, nose, and mouth
- blurred vision
- dizziness, drowsiness, problems with memory or concentration
- ringing in your ears
- feeling restless or irritable
- or mild nausea, stomach pain, constipation.

923. This particular disease, Colds & Flu, interacts with the drug Theraflu Nighttime Severe Cold & Flu

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs

Drug Side Effects Summary

of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Theraflu Nighttime Severe Cold & Cough Caplets side effects

924. This particular disease, Colds & Flu, interacts with the drug Theraflu Sinus & Cold and has the

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Theraflu Sinus & Cold and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of

Drug Side Effects Summary

reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have:

chest pain, rapid pulse, fast or uneven heart rate

- confusion, hallucinations, tremor, severe dizziness or anxiety

- slow, shallow breathing

- little or no urinating

- easy bruising or bleeding, unusual weakness

- nausea, pain in your upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)

- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety, confusion, chest pain, shortness of breath, uneven heartbeats, seizure). Common side effects of Theraflu Sinus & Cold may include: dryness of the eyes, nose, and mouth

- blurred vision

- dizziness, drowsiness, problems with memory or concentration

- ringing in your ears

- feeling restless or irritable

- or mild nausea, stomach pain, constipation.

925. This particular disease, Colds & Flu, interacts with the drug Theraflu Warming Relief Daytime S

- hives

- difficulty breathing

- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction,

Drug Side Effects Summary

you should never again take any medicine that contains acetaminophen. Theraflu Warming Severe Cold Daytime may cause serious side effects. Stop using Theraflu Warming Severe Cold Daytime and call your doctor at once if you have: chest pain, fast, slow, or uneven heart rate

- severe dizziness , feeling like you might pass out
- mood changes, confusion, hallucinations, seizure (convulsions)
- little or no urinating
- nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , chest pain, uneven heartbeats, seizure). Common side effects of Theraflu Warming Severe Cold Daytime may include: dizziness, weakness, mild headache
- diarrhea , upset stomach
- dry mouth, nose, or throat
- feeling nervous, restless, irritable, or anxious
- or sleep problems (insomnia).

926. This particular disease, Colds & Flu, interacts with the drug Theraflu Warming Relief Nighttime

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing

Drug Side Effects Summary

- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Theraflu Warming Relief Nighttime Severe Cold & Cough side effects

927. This particular disease, Colds & Flu, interacts with the drug Triaminic Multi-Symptom Fever and

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: severe headache, seizure (convulsions)
- fast, pounding, or uneven heartbeats
- confusion, hallucinations, severe dizziness or drowsiness, slow or shallow breathing

Drug Side Effects Summary

- tremor, restless muscle movements
- little or no urination
- flu symptoms, easy bruising, unusual bleeding, pale skin, weakness, feeling light-headed
- or nausea, pain in your upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of Triaminic Flu Cough & Fever may include: headache, sleep problems (insomnia)
- upset stomach , diarrhea or constipation
- dry eyes, blurred vision, dry mouth or nose
- mild dizziness or drowsiness, trouble concentrating
- feeling restless or excited (especially in children)
- or mild skin rash .

928. This particular disease, Colds & Flu, interacts with the drug Tylenol Children's Cold + Cough +

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: severe headache, seizure (convulsions)
- fast, pounding, or uneven heartbeats
- confusion, hallucinations, severe dizziness or drowsiness, slow or shallow breathing

Drug Side Effects Summary

- tremor, restless muscle movements
- little or no urination
- flu symptoms, easy bruising, unusual bleeding, pale skin, weakness, feeling light-headed
- or nausea, pain in your upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of Tylenol Childrens Plus Cough & Runny Nose may include: headache, sleep problems (insomnia)
- upset stomach , diarrhea or constipation
- dry eyes, blurred vision, dry mouth or nose
- mild dizziness or drowsiness, trouble concentrating
- feeling restless or excited (especially in children)
- or mild skin rash .

929. This particular disease, Colds & Flu, interacts with the drug Tylenol Children's Cold + Flu and I

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness

Drug Side Effects Summary

- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Tylenol Children's Cold + Flu side effects

930. This particular disease, Colds & Flu, interacts with the drug Tylenol Cold & Cough Nighttime a

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: severe headache, seizure (convulsions)
- fast, pounding, or uneven heartbeats
- confusion, hallucinations, severe dizziness or drowsiness, slow or shallow breathing
- tremor, restless muscle movements
- little or no urinating
- flu symptoms, easy bruising, unusual bleeding, pale skin, weakness, feeling light-headed
- or nausea, pain in your upper stomach pain, itching, loss of appetite, dark urine, clay-colored

Drug Side Effects Summary

stools, jaundice (yellowing of the skin or eyes). Common side effects of Tylenol Cold & Cough Nighttime Cool Burst may include: headache, sleep problems (insomnia)

- upset stomach , diarrhea or constipation
- dry eyes, blurred vision, dry mouth or nose
- mild dizziness or drowsiness, trouble concentrating
- feeling restless or excited (especially in children)
- or mild skin rash .

931. This particular disease, Colds & Flu, interacts with the drug Tylenol Cold Multi-Symptom Night

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- a light-headed feeling, like you might pass out
- anxiety , hallucinations, feeling of fear
- little or no urinating
- pale skin, unusual weakness
- nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of your skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, chest

Drug Side Effects Summary

pain, shortness of breath, uneven heartbeats, seizure). Common side effects of Tylenol Cold Multi-Symptom Nighttime may include: drowsiness

- constipation , diarrhea , upset stomach , vomiting
- or feeling restless or excited.

932. This particular disease, Colds & Flu, interacts with the drug Tylenol Plus Children's Cold and h

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Tylenol Children's Plus Cold and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have:
chest pain, rapid pulse, fast or uneven heart rate
- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- little or no urinating
- nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of your skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , chest pain, shortness of breath, uneven heartbeats, seizure). Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of Tylenol Children's Plus Cold may include: dizziness , drowsiness

Drug Side Effects Summary

- mild headache, blurred vision
- dry mouth, nose, or throat
- constipation
- feeling nervous
- or sleep problems (insomnia)
-

933. This particular disease, Colds & Flu, interacts with the drug Tylenol Sinus Congestion & Pain M

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Tylenol Sinus Congestion Nighttime and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate
- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- little or no urinating
- nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of your skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , chest pain, shortness of breath, uneven heartbeats, seizure). Side effects such as dry mouth,

Drug Side Effects Summary

constipation , and confusion may be more likely in older adults. Common side effects of Tylenol

Sinus Congestion Nighttime may include: dizziness , drowsiness

- mild headache, blurred vision
- dry mouth, nose, or throat
- constipation
- feeling nervous
- or sleep problems (insomnia)
-

934. This particular disease, Colds & Flu, interacts with the drug Vicks VapoSteam and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Vicks VapoSteam side effects

935. This particular disease, Colds & Flu, interacts with the drug Abatuss DMX and has the following

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: fast, slow, or uneven heart rate
- severe headache, mood changes, hallucinations
- severe dizziness or anxiety , feeling like you might pass out
- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- fever
- little or no urinating
- shortness of breath
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety, confusion, chest pain, uneven heartbeats, seizure). Common side effects of Abatuss DMX may include: mild headache
- mild dizziness, drowsiness
- dry mouth, nose, or throat
- nausea , diarrhea , constipation , upset stomach
- feeling nervous, restless, or irritable
- blurred vision
- or sleep problems (insomnia).

936. This particular disease, Colds & Flu, interacts with the drug acetaminophen / caffeine and has

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past

Drug Side Effects Summary

and had no reaction. Stop taking acetaminophen and caffeine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. Acetaminophen and caffeine may cause serious side effects. Stop using acetaminophen and caffeine and call your doctor at once if you have: low fever with nausea, stomach pain, and loss of appetite

- dark urine, clay-colored stools
- or jaundice (yellowing of the skin or eyes). Common side effects of acetaminophen and caffeine may include: sleep problems (insomnia)
- or feeling nervous, irritable, or jittery.

937. This particular disease, Colds & Flu, interacts with the drug acetaminophen / chlorpheniramine

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (

Drug Side Effects Summary

Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)

- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Acetaminophen / chlorpheniramine / dextromethorphan / guaifenesin side effects

938. This particular disease, Colds & Flu, interacts with the drug acetaminophen / chlorpheniramine

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: severe headache, seizure (convulsions)
- fast, pounding, or uneven heartbeats
- confusion, hallucinations, severe dizziness or drowsiness, slow or shallow breathing
- tremor, restless muscle movements
- little or no urination
- flu symptoms, easy bruising, unusual bleeding, pale skin, weakness, feeling light-headed
- or nausea, pain in your upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of acetaminophen,

Drug Side Effects Summary

chlorpheniramine, and dextromethorphan may include: headache, sleep problems (insomnia)

- upset stomach , diarrhea or constipation
- dry eyes, blurred vision, dry mouth or nose
- mild dizziness or drowsiness, trouble concentrating
- feeling restless or excited (especially in children)
- or mild skin rash .

939. This particular disease, Colds & Flu, interacts with the drug acetaminophen / dextbrompheniramine

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)

Drug Side Effects Summary

- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Acetaminophen / dextbrompheniramine side effects

940. This particular disease, Colds & Flu, interacts with the drug acetaminophen / dextromethorpha

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: fast, pounding, or uneven heartbeats
- severe drowsiness
- painful or difficult urination
- severe headache, seizure
- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of acetaminophen, dextromethorphan, and diphenhydramine may include: drowsiness
- headache, sleep problems (insomnia)
- upset stomach , diarrhea or constipation
- dry eyes, blurred vision
- dry mouth, nose, or throat
- or feeling restless or excited (especially in children).

Drug Side Effects Summary

941. This particular disease, Colds & Flu, interacts with the drug acetaminophen / dextromethorpha

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: any new or worsening symptoms
- nervousness , dizziness , or drowsiness
- fever for longer than 3 days
- redness, swelling, or rash
- ongoing headache
- or pain, cough, or sinus congestion lasting more than 7 days. Some side effects may be more likely in older adults. Common side effects of acetaminophen, dextromethorphan, and triprolidine may include: dizziness, drowsiness
- dry mouth, nose, or throat
- blurred vision
- constipation
- feeling restless or excited (especially in children)
- or sleep problems (insomnia).

942. This particular disease, Colds & Flu, interacts with the drug acetaminophen / dextromethorpha

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: redness or swelling
- new or worsening symptoms
- an ongoing cough with a rash or headache
- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of acetaminophen and dextromethorphan may include: upset stomach
- confusion
- or drowsiness.

943. This particular disease, Colds & Flu, interacts with the drug Advil Children's Allergy Sinus and

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe dizziness , trouble sleeping, or nervousness
- shortness of breath (even with mild exertion)
- swelling or rapid weight gain
- a skin rash, no matter how mild
- severe headache, blurred vision, pounding in your neck or ears

Drug Side Effects Summary

- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath. Common side effects of Advil Childrens Allergy Sinus may include: upset stomach , mild heartburn , nausea , vomiting
- bloating , gas, diarrhea , constipation
- dizziness, headache, nervousness
- flushing (warmth, redness, or tingly feeling)
- mild itching or rash
- or ringing in your ears.

944. This particular disease, Colds & Flu, interacts with the drug Alahist D and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: nervousness
- dizziness
- or trouble sleeping. Common side effects of Alahist D may include: drowsiness
- feeling restless or excited (especially in children)
- dry mouth, nose, or throat
- constipation
- or blurred vision, double vision.

Drug Side Effects Summary

945. This particular disease, Colds & Flu, interacts with the drug Allan Tannate Pediatric and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Allan Tannate Pediatric may cause serious side effects. Stop using Allan Tannate Pediatric and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate
- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)
- little or no urinating
- easy bruising or bleeding, unusual weakness
- or dangerously high blood pressure (severe headache, buzzing in your ears, anxiety , shortness of breath). Common side effects of Allan Tannate Pediatric may include: dizziness , drowsiness, blurred vision
- dry nose or mouth
- nausea , stomach pain, constipation , loss of appetite
- problems with memory or concentration
- or feeling restless or excited (especially in children).

946. This particular disease, Colds & Flu, interacts with the drug Allerest PE and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Allerest PE may cause serious side effects. Stop using Allerest PE and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate
- confusion, hallucinations, severe nervousness

Drug Side Effects Summary

- tremor, seizure (convulsions)
- little or no urinating
- easy bruising or bleeding, unusual weakness
- or dangerously high blood pressure (severe headache, buzzing in your ears, anxiety , shortness of breath). Common side effects of Allerest PE may include: dizziness , drowsiness, blurred vision
- dry nose or mouth
- nausea , stomach pain, constipation , loss of appetite
- problems with memory or concentration
- or feeling restless or excited (especially in children).

947. This particular disease, Colds & Flu, interacts with the drug Allerfrim and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Allerfrim side effects

948. This particular disease, Colds & Flu, interacts with the drug Ambifed CD and has the following side effects:

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Like other narcotic medications, codeine can slow your breathing. Death may occur if breathing becomes too weak. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Stop taking this medicine and call your doctor at once if you have: noisy breathing, sighing, shallow breathing
- severe drowsiness, hallucinations
- fast, slow, or uneven heart rate
- a seizure
- nervousness , tremors
- or little or no urination. Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Common side effects of Ambifed CD may include: dizziness, drowsiness, weakness
- headache
- nausea , constipation
- sleep problems (insomnia)
- or feeling nervous, anxious, or restless.

949. This particular disease, Colds & Flu, interacts with the drug Aphedrid and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe dizziness
- nervousness

Drug Side Effects Summary

- trouble sleeping
- or high blood pressure (severe headache, pounding in your neck or ears). Common side effects of Aphedrid may include: dizziness, drowsiness
- dry mouth, nose, or throat
- constipation
- blurred vision
- or feeling restless or excited (especially in children).

950. This particular disease, Colds & Flu, interacts with the drug Avedana Medicated Chest Rub and

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Less serious side effects may be more likely, and you may have none at all.

951. This particular disease, Colds & Flu, interacts with the drug Benadryl Allergy Plus Congestion

- No side effects listed

952. This particular disease, Colds & Flu, interacts with the drug BP Allergy JR and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. BP Allergy JR may cause serious side effects. Stop using BP Allergy JR and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate
- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)

Drug Side Effects Summary

- little or no urinating
- easy bruising or bleeding, unusual weakness
- or dangerously high blood pressure (severe headache, buzzing in your ears, anxiety , shortness of breath). Common side effects of BP Allergy JR may include: dizziness , drowsiness, blurred vision
- dry nose or mouth
- nausea , stomach pain, constipation , loss of appetite
- problems with memory or concentration
- or feeling restless or excited (especially in children).

953. This particular disease, Colds & Flu, interacts with the drug brompheniramine / codeine / phen

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Like other narcotic medications, codeine can slow your breathing. Death may occur if breathing becomes too weak. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Stop taking this medicine and call your doctor at once if you have: noisy breathing, sighing, shallow breathing
- a slow heart rate or weak pulse
- nervousness , trouble sleeping
- severe constipation
- or little or no urination. Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Common side effects of brompheniramine, codeine, and phenylephrine may include: drowsiness
- or feeling restless or excited (especially in children). Side effects such as dry mouth, constipation, and confusion may be more likely in older adults.

Drug Side Effects Summary

954. This particular disease, Colds & Flu, interacts with the drug C-Tanna 12 and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. C-Tanna 12 may cause serious side effects. Stop using C-Tanna 12 and call your doctor at once if you have: fast, pounding, or uneven heartbeat
- severe headache, hallucinations, seizure (convulsions)
- severe dizziness or drowsiness
- vision problems
- little or no urinating
- tremors, severe restlessness
- or seizures (convulsions). Common side effects of C-Tanna 12 may include: constipation , diarrhea
- nausea , vomiting , upset stomach , loss of appetite
- weakness, mild dizziness or drowsiness
- anxiety , feeling nervous or excited
- or sleep problems (insomnia).

955. This particular disease, Colds & Flu, interacts with the drug camphor / menthol and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. camphor and menthol topical may cause serious side effects. Stop using camphor and menthol topical and call your doctor at once if you have: severe burning, stinging, or irritation after using this medicine
- or breathing problems after using this medicine on the chest or throat. Less serious side effects may be more likely, and you may have none at all.

Drug Side Effects Summary

956. This particular disease, Colds & Flu, interacts with the drug Capron DMT and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Capron DMT may cause serious side effects. Stop using Capron DMT and call your doctor at once if you have: severe drowsiness or anxiety
- little or no urinating
- or a light-headed feeling, like you might pass out. Common side effects of Capron DMT may include: drowsiness
- nervousness
- or feeling restless or excited (especially in children).

957. This particular disease, Colds & Flu, interacts with the drug carbetapentane / chlorpheniramine

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. carbetapentane and chlorpheniramine may cause serious side effects. Stop using carbetapentane and chlorpheniramine and call your doctor at once if you have: fast, pounding, or uneven heartbeat
- severe headache, hallucinations, seizure (convulsions)
- severe dizziness or drowsiness
- vision problems
- little or no urinating
- tremors, severe restlessness
- or seizures (convulsions). Common side effects of carbetapentane and chlorpheniramine may include: constipation , diarrhea
- nausea , vomiting , upset stomach , loss of appetite

Drug Side Effects Summary

- weakness, mild dizziness or drowsiness
- anxiety , feeling nervous or excited
- or sleep problems (insomnia).

958. This particular disease, Colds & Flu, interacts with the drug Children's Triacting and has the fo

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: redness or swelling
- new or worsening symptoms
- an ongoing cough with a rash or headache
- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Children's Triacting may include: upset stomach
- confusion
- or drowsiness.

959. This particular disease, Colds & Flu, interacts with the drug chlophedianol / dextbrompheniram

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if

Drug Side Effects Summary

you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Feeling very sleepy. Chlophedianol / dexbrompheniramine side effects

960. This particular disease, Colds & Flu, interacts with the drug chlophedianol / dexchlorpheniram

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe drowsiness
- a light-headed feeling, like you might pass out
- severe dizziness or anxiety
- or severe restless feeling or being unable to sleep. Common side effects of chlophedianol, dexchlorpheniramine, and pseudoephedrine may include: mild dizziness or drowsiness
- dry mouth, nose, or throat
- diarrhea , constipation , upset stomach
- feeling restless or excited (especially in children)
- blurred vision

Drug Side Effects Summary

- or sleep problems (insomnia).

961. This particular disease, Colds & Flu, interacts with the drug chlorcyclizine / phenylephrine and

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: fast or uneven heart rate
- tremor, seizure (convulsions)
- urinating less than usual or not at all
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , confusion, chest pain, uneven heartbeats, seizure). Common side effects of chlorcyclizine and phenylephrine may include: dizziness , drowsiness
- constipation
- blurred vision
- or feeling nervous or restless.

962. This particular disease, Colds & Flu, interacts with the drug Chlorex-A 12 and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: confusion, hallucinations
- feeling nervous or irritable
- trouble sleeping
- little or no urination
- a seizure (convulsions)

Drug Side Effects Summary

- numbness or tingly feeling
- a light-headed feeling, like you might pass out
- pounding heartbeats or fluttering in your chest
- or increased blood pressure--severe headache, blurred vision, pounding in your neck or ears, anxiety , nosebleed. Side effects may be more likely in older adults. Common side effects of Chlorex-A 12 may include: nausea , constipation , loss of appetite
- rash or itching
- increased sweating
- dry mouth, nose, or throat
- blurred vision
- headache, dizziness
- or feeling restless or excited (especially in children).

963. This particular disease, Colds & Flu, interacts with the drug Chlorex A and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: fast or uneven heart rate
- pounding heartbeats or fluttering in your chest
- painful or difficult urination, little or no urinating
- tremors, seizure (convulsions)
- or vision changes, severe dizziness , feeling like you might pass out. Common side effects of Chlorex A may include: headache, dizziness, drowsiness
- dry mouth, nose, or throat
- upset stomach , loss of appetite, nausea , constipation

Drug Side Effects Summary

- blurred vision
- sleep problems (insomnia)
- or feeling restless or excited (especially in children).

964. This particular disease, Colds & Flu, interacts with the drug chlorpheniramine / codeine / pseudoephedrine

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad dizziness or passing out. Chest pain or pressure or a fast heartbeat . Trouble breathing, slow breathing, or shallow breathing. Noisy breathing. Feeling very sleepy. Feeling confused. A heartbeat that does not feel normal. Hallucinations (seeing or hearing things that are not there). Mood changes. Seizures . Very bad belly pain. Very bad headache . Trouble passing urine. Shakiness. Change in eyesight. Very upset stomach or throwing up. Very bad constipation . Feeling very tired or weak. Chlorpheniramine / codeine / pseudoephedrine side effects

965. This particular disease, Colds & Flu, interacts with the drug chlorpheniramine / codeine and pseudoephedrine

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep
- confusion, severe drowsiness
- a light-headed feeling, like you might pass out
- severe constipation
- or adrenal gland problems-- nausea , vomiting , loss of appetite, feeling very dizzy, weak, or tired. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea . Serious side effects may be more likely in older adults and those who are debilitated. Common side effects of chlorpheniramine and codeine may include: dizziness, drowsiness, feeling light-headed
- feeling short of breath
- confusion, mood changes, lack of energy, feeling nervous, irritable, agitated, or restless
- sleep problems
- headache , vision problems
- tremors, coordination problems
- dry mouth
- sweating
- or nausea, vomiting, constipation, stomach pain, bloating .

Drug Side Effects Summary

966. This particular disease, Colds & Flu, interacts with the drug chlorpheniramine / phenylephrine

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: fast or uneven heart rate
- pounding heartbeats or fluttering in your chest
- painful or difficult urination, little or no urinating
- tremors, seizure (convulsions)
- or vision changes, severe dizziness , feeling like you might pass out. Common side effects of chlorpheniramine, phenylephrine, and phenyltoloxamine may include: headache, dizziness, drowsiness
- dry mouth, nose, or throat
- upset stomach , loss of appetite, nausea , constipation
- blurred vision
- sleep problems (insomnia)
- or feeling restless or excited (especially in children).

967. This particular disease, Colds & Flu, interacts with the drug ChlorTan D and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. ChlorTan D may cause serious side effects. Stop using ChlorTan D and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate
- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)

Drug Side Effects Summary

- little or no urinating
- easy bruising or bleeding, unusual weakness
- or dangerously high blood pressure (severe headache, buzzing in your ears, anxiety , shortness of breath). Common side effects of ChlorTan D may include: dizziness , drowsiness, blurred vision
- dry nose or mouth
- nausea , stomach pain, constipation , loss of appetite
- problems with memory or concentration
- or feeling restless or excited (especially in children).

968. This particular disease, Colds & Flu, interacts with the drug Clear Cough PM Multi-Symptom and

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health

Drug Side Effects Summary

problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)

- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Clear Cough PM Multi-Symptom side effects

969. This particular disease, Colds & Flu, interacts with the drug Codar AR and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep
- confusion, severe drowsiness
- a light-headed feeling, like you might pass out
- severe constipation
- or adrenal gland problems-- nausea , vomiting , loss of appetite, feeling very dizzy, weak, or tired. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea . Serious side effects may be more likely in older adults and those who are debilitated. Common side effects of Codar AR may include: dizziness, drowsiness, feeling light-headed
- feeling short of breath
- confusion, mood changes, lack of energy, feeling nervous, irritable, agitated, or restless
- sleep problems

Drug Side Effects Summary

- headache , vision problems
- tremors, coordination problems
- dry mouth
- sweating
- or nausea, vomiting, constipation, stomach pain, bloating .

970. This particular disease, Colds & Flu, interacts with the drug codeine / phenylephrine / triprolidine

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Like other narcotic medications, codeine can slow your breathing. Death may occur if breathing becomes too weak. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Like other narcotic medications, codeine can slow your breathing. Death may occur if breathing becomes too weak. Stop taking this medicine and call your doctor at once if you have: noisy breathing, sighing, shallow breathing
- a slow heart rate or weak pulse
- nervousness , trouble sleeping
- severe headache or dizziness, ringing in your ears
- severe constipation
- or little or no urination. Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Common side effects of codeine, phenylephrine, and triprolidine may include: dizziness, drowsiness, tired feeling
- dry mouth, nose, or throat
- blurred vision
- nausea , vomiting , constipation

Drug Side Effects Summary

- or sleep problems (insomnia). Side effects such as dry mouth, constipation, and confusion may be more likely in older adults.

971. This particular disease, Colds & Flu, interacts with the drug Codimal DM and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Codimal DM side effects

972. This particular disease, Colds & Flu, interacts with the drug Ceditussin DAC and has the follow

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever

Drug Side Effects Summary

- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad dizziness or passing out. Chest pain or pressure or a fast heartbeat . Trouble breathing, slow breathing, or shallow breathing. Noisy breathing. Feeling very sleepy. Feeling confused. A heartbeat that does not feel normal. Hallucinations (seeing or hearing things that are not there). Mood changes. Seizures . Very bad belly pain. Very bad constipation . Very bad headache . Trouble passing urine. Shakiness. Change in eyesight. Feeling very tired or weak. Coditussin DAC side effects

973. This particular disease, Colds & Flu, interacts with the drug Comtrex Cold & Cough and has the

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. Stop using Comtrex Cold & Cough and call your doctor at once if you have: chest pain, fast, slow, or uneven heart rate
- severe dizziness , feeling like you might pass out
- mood changes, confusion, hallucinations, seizure (convulsions)
- little or no urinating
- nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)

Drug Side Effects Summary

- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , chest pain, uneven heartbeats, seizure). Common side effects of Comtrex Cold & Cough may include: dizziness, weakness, mild headache
- diarrhea , upset stomach
- dry mouth, nose, or throat
- feeling nervous, restless, irritable, or anxious
- or sleep problems (insomnia).

974. This particular disease, Colds & Flu, interacts with the drug Comtrex Nighttime Cold & Cough

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have

Drug Side Effects Summary

signs like red, swollen, blistered, or peeling skin (with or without fever)

- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Comtrex Nighttime Cold & Cough side effects

975. This particular disease, Colds & Flu, interacts with the drug Comtrex Severe Cold & Sinus and

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Comtrex Severe Cold & Sinus and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate
- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- little or no urinating
- nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of your skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , chest pain, shortness of breath, uneven heartbeats, seizure). Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of Comtrex Severe Cold & Sinus may include: dizziness , drowsiness
- mild headache, blurred vision

Drug Side Effects Summary

- dry mouth, nose, or throat
- constipation
- feeling nervous
- or sleep problems (insomnia)
-

976. This particular disease, Colds & Flu, interacts with the drug Conal and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: confusion, hallucinations
- feeling nervous or irritable
- trouble sleeping
- little or no urination
- a seizure (convulsions)
- numbness or tingly feeling
- a light-headed feeling, like you might pass out
- pounding heartbeats or fluttering in your chest
- or increased blood pressure--severe headache, blurred vision, pounding in your neck or ears, anxiety , nosebleed. Side effects may be more likely in older adults. Common side effects of Conal may include: nausea , constipation , loss of appetite
- rash or itching
- increased sweating
- dry mouth, nose, or throat
- blurred vision

Drug Side Effects Summary

- headache, dizziness
- or feeling restless or excited (especially in children).

977. This particular disease, Colds & Flu, interacts with the drug Conex and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe nervousness
- a light-headed feeling, like you might pass out
- trouble sleeping
- fast or uneven heart rate
- or painful or difficult urination. Side effects may be more likely in older adults. Common side effects of Conex may include: drowsiness
- feeling restless or excited (especially in children)
- blurred vision
- or dry nose or mouth.

978. This particular disease, Colds & Flu, interacts with the drug Contac Cold + Flu (Night) Cooling Lotion and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause

Drug Side Effects Summary

serious side effects. Stop using the medicine and call your doctor at once if you have: severe headache, seizure (convulsions)

- fast, pounding, or uneven heartbeats
- confusion, hallucinations, severe dizziness or drowsiness, slow or shallow breathing
- tremor, restless muscle movements
- little or no urinating
- flu symptoms, easy bruising, unusual bleeding, pale skin, weakness, feeling light-headed
- or nausea, pain in your upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Contac Cold+Flu Cooling Night may include: headache, sleep problems (insomnia)
- upset stomach , diarrhea or constipation
- dry eyes, blurred vision, dry mouth or nose
- mild dizziness or drowsiness, trouble concentrating
- feeling restless or excited (especially in children)
- or mild skin rash .

979. This particular disease, Colds & Flu, interacts with the drug Dallery and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: fast or uneven heart rate
- tremor, seizure (convulsions)
- urinating less than usual or not at all
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , confusion, chest pain, uneven heartbeats, seizure). Common side effects of Dallery may include:

Drug Side Effects Summary

dizziness , drowsiness

- constipation
- blurred vision
- or feeling nervous or restless.

980. This particular disease, Colds & Flu, interacts with the drug Dallery Drops and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Dallery Drops may cause serious side effects. Stop using Dallery Drops and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate
- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)
- little or no urinating
- easy bruising or bleeding, unusual weakness
- or dangerously high blood pressure (severe headache, buzzing in your ears, anxiety , shortness of breath). Common side effects of Dallery Drops may include: dizziness , drowsiness, blurred vision
- dry nose or mouth
- nausea , stomach pain, constipation , loss of appetite
- problems with memory or concentration
- or feeling restless or excited (especially in children).

981. This particular disease, Colds & Flu, interacts with the drug Decorel Forte Plus and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if

Drug Side Effects Summary

you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Decorel Forte Plus side effects

982. This particular disease, Colds & Flu, interacts with the drug Delsym Cough & Sore Throat and

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction,

Drug Side Effects Summary

you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: redness or swelling

- new or worsening symptoms
- an ongoing cough with a rash or headache
- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Children's Delsym Cough Plus Sore Throat may include: upset stomach
- confusion
- or drowsiness.

983. This particular disease, Colds & Flu, interacts with the drug Delsym Night Time Multi-Symptom

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- a light-headed feeling, like you might pass out
- anxiety , hallucinations, feeling of fear
- little or no urinating
- pale skin, unusual weakness

Drug Side Effects Summary

- nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of your skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, chest pain, shortness of breath, uneven heartbeats, seizure). Common side effects of Delsym Adult Night Time Multi-Symptom may include: drowsiness
- constipation , diarrhea , upset stomach , vomiting
- or feeling restless or excited.

984. This particular disease, Colds & Flu, interacts with the drug dexchlorpheniramine / dextromethorphan

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: fast, slow, or uneven heart rate
- severe headache, mood changes, hallucinations
- severe dizziness or anxiety , feeling like you might pass out
- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- fever
- little or no urinating
- shortness of breath
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety, confusion, chest pain, uneven heartbeats, seizure). Common side effects of dexchlorpheniramine, dextromethorphan, and pseudoephedrine may include: mild headache
- mild dizziness, drowsiness
- dry mouth, nose, or throat

Drug Side Effects Summary

- nausea , diarrhea , constipation , upset stomach
- feeling nervous, restless, or irritable
- blurred vision
- or sleep problems (insomnia).

985. This particular disease, Colds & Flu, interacts with the drug dextromethorphan / diphenhydramine

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe drowsiness
- painful or difficult urination
- fast, pounding, or uneven heartbeats
- trouble breathing
- confusion, anxiety , irritability, tremors
- a seizure
- or problems with balance or speech. Common side effects of dextromethorphan, diphenhydramine, and phenylephrine may include: dizziness , drowsiness, headache
- blurred vision, dry eyes
- dry mouth, nose, or throat
- upset stomach , nausea , vomiting
- diarrhea , constipation
- or tiredness, trouble sleeping.

986. This particular disease, Colds & Flu, interacts with the drug dextromethorphan / pyrilamine and

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Dextromethorphan and pyrilamine may cause serious side effects. Stop using dextromethorphan and pyrilamine and call your doctor at once if you have: severe drowsiness or anxiety
- little or no urinating
- or a light-headed feeling, like you might pass out. Common side effects of dextromethorphan and pyrilamine may include: drowsiness
- nervousness
- or feeling restless or excited (especially in children).

987. This particular disease, Colds & Flu, interacts with the drug Dixaphedrine and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Dixaphedrine side effects

988. This particular disease, Colds & Flu, interacts with the drug Duraflu and has the following side

Drug Side Effects Summary

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: chest pain
- fast, slow, or uneven heart rate
- severe dizziness , feeling like you might pass out
- mood changes, confusion, hallucinations, tremor, seizure (convulsions)
- fever
- little or no urinating
- nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of your skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , chest pain, uneven heartbeats, seizure). Common side effects of Duraflu may include: dizziness, weakness
- mild headache
- mild nausea, diarrhea , upset stomach
- dry mouth, nose, or throat
- feeling nervous, restless, irritable, or anxious
- or sleep problems (insomnia).

989. This particular disease, Colds & Flu, interacts with the drug Dytan-D and has the following side

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe drowsiness
- sleeplessness
- painful or difficult urination
- fast, pounding, or uneven heartbeats
- trouble breathing
- anxiety , nervousness
- a seizure
- or problems with balance or speech. Common side effects of Dytan-D may include: dizziness , drowsiness, headache
- dry mouth, nose, or throat
- nausea , vomiting, constipation
- blurred vision
- or trouble sleeping.

990. This particular disease, Colds & Flu, interacts with the drug Dytan-DM and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe drowsiness
- painful or difficult urination
- fast, pounding, or uneven heartbeats
- trouble breathing

Drug Side Effects Summary

- confusion, anxiety , irritability, tremors
- a seizure
- or problems with balance or speech. Common side effects of Dytan-DM may include: dizziness , drowsiness, headache
- blurred vision, dry eyes
- dry mouth, nose, or throat
- upset stomach , nausea , vomiting
- diarrhea , constipation
- or tiredness, trouble sleeping.

991. This particular disease, Colds & Flu, interacts with the drug Ed ChlorPed D and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Ed ChlorPed D may cause serious side effects. Stop using Ed ChlorPed D and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate
- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)
- little or no urinating
- easy bruising or bleeding, unusual weakness
- or dangerously high blood pressure (severe headache, buzzing in your ears, anxiety , shortness of breath). Common side effects of Ed ChlorPed D may include: dizziness , drowsiness, blurred vision
- dry nose or mouth
- nausea , stomach pain, constipation , loss of appetite
- problems with memory or concentration

Drug Side Effects Summary

- or feeling restless or excited (especially in children).

992. This particular disease, Colds & Flu, interacts with the drug Excedrin Quick Tab and has the fo

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Excedrin Quick Tab Peppermint and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. Excedrin Quick Tab Peppermint may cause serious side effects. Stop using Excedrin Quick Tab Peppermint and call your doctor at once if you have: low fever with nausea, stomach pain, and loss of appetite
- dark urine, clay-colored stools
- or jaundice (yellowing of the skin or eyes). Common side effects of Excedrin Quick Tab Peppermint may include: sleep problems (insomnia)
- or feeling nervous, irritable, or jittery.

993. This particular disease, Colds & Flu, interacts with the drug Excedrin Tension Headache and h

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Excedrin Tension Headache and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. Excedrin Tension

Drug Side Effects Summary

Headache may cause serious side effects. Stop using Excedrin Tension Headache and call your doctor at once if you have: low fever with nausea, stomach pain, and loss of appetite

- dark urine, clay-colored stools
- or jaundice (yellowing of the skin or eyes). Common side effects of Excedrin Tension Headache may include: sleep problems (insomnia)
- or feeling nervous, irritable, or jittery.

994. This particular disease, Colds & Flu, interacts with the drug Fast Freeze and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Fast Freeze may cause serious side effects. Stop using Fast Freeze and call your doctor at once if you have: severe burning, stinging, or irritation after using this medicine
- or breathing problems after using this medicine on the chest or throat. Less serious side effects may be more likely, and you may have none at all.

995. This particular disease, Colds & Flu, interacts with the drug Flu & Severe Cold & Cough Daytime

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. Stop using Flu & Severe Cold & Cough Daytime Powder and call your doctor at once if you have: chest pain, fast, slow, or

Drug Side Effects Summary

uneven heart rate

- severe dizziness , feeling like you might pass out
- mood changes, confusion, hallucinations, seizure (convulsions)
- little or no urinating
- nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , chest pain, uneven heartbeats, seizure). Common side effects of Flu & Severe Cold & Cough Daytime Powder may include: dizziness, weakness, mild headache
- diarrhea , upset stomach
- dry mouth, nose, or throat
- feeling nervous, restless, irritable, or anxious
- or sleep problems (insomnia).

996. This particular disease, Colds & Flu, interacts with the drug Freeze It and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Freeze It may cause serious side effects. Stop using Freeze It and call your doctor at once if you have: severe burning, stinging, or irritation after using this medicine
- or breathing problems after using this medicine on the chest or throat. Less serious side effects may be more likely, and you may have none at all.

997. This particular disease, Colds & Flu, interacts with the drug GNP Cold Relief Plus and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and

Drug Side Effects Summary

sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. GNP Cold Relief Plus side effects

998. This particular disease, Colds & Flu, interacts with the drug Guaifen DAC and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Like other narcotic medications, codeine can slow your breathing. Death may occur if breathing becomes too weak. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you

Drug Side Effects Summary

are hard to wake up. Stop taking this medicine and call your doctor at once if you have: noisy breathing, sighing, shallow breathing

- severe drowsiness, hallucinations

- fast, slow, or uneven heart rate

- a seizure

- nervousness , tremors

- or little or no urination. Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Common side effects of Guaifen DAC may include: dizziness, drowsiness, weakness

- headache

- nausea , constipation

- sleep problems (insomnia)

- or feeling nervous, anxious, or restless.

999. This particular disease, Colds & Flu, interacts with the drug Histafed and has the following side effects:

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe dizziness

- nervousness

- trouble sleeping

- or high blood pressure (severe headache, pounding in your neck or ears). Common side effects of Histafed may include: dizziness, drowsiness

- dry mouth, nose, or throat

- constipation

Drug Side Effects Summary

- blurred vision
- or feeling restless or excited (especially in children).

1000. This particular disease, Colds & Flu, interacts with the drug Histex-AC and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Like other narcotic medications, codeine can slow your breathing. Death may occur if breathing becomes too weak. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Like other narcotic medications, codeine can slow your breathing. Death may occur if breathing becomes too weak. Stop taking this medicine and call your doctor at once if you have: noisy breathing, sighing, shallow breathing
- a slow heart rate or weak pulse
- nervousness , trouble sleeping
- severe headache or dizziness, ringing in your ears
- severe constipation
- or little or no urination. Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Common side effects of Histex-AC may include: dizziness, drowsiness, tired feeling
- dry mouth, nose, or throat
- blurred vision
- nausea , vomiting , constipation
- or sleep problems (insomnia). Side effects such as dry mouth, constipation, and confusion may be more likely in older adults.

Drug Side Effects Summary

1001. This particular disease, Colds & Flu, interacts with the drug Histex-DM and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Histex-DM side effects

1002. This particular disease, Colds & Flu, interacts with the drug Histex-PE and has the following s

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking

Drug Side Effects Summary

- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Histex-PE side effects

1003. This particular disease, Colds & Flu, interacts with the drug Lortuss EX and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Like other narcotic medications, codeine can slow your breathing. Death may occur if breathing becomes too weak. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Stop taking this medicine and call your doctor at once if you have: noisy breathing, sighing, shallow breathing
- severe drowsiness, hallucinations
- fast, slow, or uneven heart rate
- a seizure
- nervousness , tremors
- or little or no urination. Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Common side effects of Lortuss EX may include: dizziness, drowsiness, weakness
- headache
- nausea , constipation
- sleep problems (insomnia)
- or feeling nervous, anxious, or restless.

1004. This particular disease, Colds & Flu, interacts with the drug M-End PE and has the following

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Like other narcotic medications, codeine can slow your breathing. Death may occur if breathing becomes too weak. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Stop taking this medicine and call your doctor at once if you have: noisy breathing, sighing, shallow breathing
- a slow heart rate or weak pulse
- nervousness , trouble sleeping
- severe constipation
- or little or no urination. Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Common side effects of M-End PE may include: drowsiness
- or feeling restless or excited (especially in children). Side effects such as dry mouth, constipation, and confusion may be more likely in older adults.

1005. This particular disease, Colds & Flu, interacts with the drug Medicidin-D and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat

Drug Side Effects Summary

- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Medicidin-D side effects

1006. This particular disease, Colds & Flu, interacts with the drug Mucinex Nightshift Cold & Flu and

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: any new or worsening symptoms
- nervousness , dizziness , or drowsiness
- fever for longer than 3 days
- redness, swelling, or rash
- ongoing headache

Drug Side Effects Summary

- or pain, cough, or sinus congestion lasting more than 7 days. Some side effects may be more likely in older adults. Common side effects of Mucinex Nightshift Cold & Flu may include: dizziness, drowsiness
- dry mouth, nose, or throat
- blurred vision
- constipation
- feeling restless or excited (especially in children)
- or sleep problems (insomnia).

1007. This particular disease, Colds & Flu, interacts with the drug Nalex-A and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: fast or uneven heart rate
- pounding heartbeats or fluttering in your chest
- painful or difficult urination, little or no urinating
- tremors, seizure (convulsions)
- or vision changes, severe dizziness , feeling like you might pass out. Common side effects of Nalex-A may include: headache, dizziness, drowsiness
- dry mouth, nose, or throat
- upset stomach , loss of appetite, nausea , constipation
- blurred vision
- sleep problems (insomnia)
- or feeling restless or excited (especially in children).

Drug Side Effects Summary

1008. This particular disease, Colds & Flu, interacts with the drug Nalex A 12 and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: confusion, hallucinations
- feeling nervous or irritable
- trouble sleeping
- little or no urination
- a seizure (convulsions)
- numbness or tingly feeling
- a light-headed feeling, like you might pass out
- pounding heartbeats or fluttering in your chest
- or increased blood pressure--severe headache, blurred vision, pounding in your neck or ears, anxiety , nosebleed. Side effects may be more likely in older adults. Common side effects of Nalex A 12 may include: nausea , constipation , loss of appetite
- rash or itching
- increased sweating
- dry mouth, nose, or throat
- blurred vision
- headache, dizziness
- or feeling restless or excited (especially in children).

1009. This particular disease, Colds & Flu, interacts with the drug NoHist LQ and has the following

- hives
- difficulty breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. NoHist LQ may cause serious side effects. Stop using NoHist LQ and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate
- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)
- little or no urinating
- easy bruising or bleeding, unusual weakness
- or dangerously high blood pressure (severe headache, buzzing in your ears, anxiety , shortness of breath). Common side effects of NoHist LQ may include: dizziness , drowsiness, blurred vision
- dry nose or mouth
- nausea , stomach pain, constipation , loss of appetite
- problems with memory or concentration
- or feeling restless or excited (especially in children).

1010. This particular disease, Colds & Flu, interacts with the drug Novahistine and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness

Drug Side Effects Summary

- or swelling of the mouth, face, lips, tongue, or throat. Novahistine side effects

1011. This particular disease, Colds & Flu, interacts with the drug Onset Forte and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

- itching

- red, swollen, blistered, or peeling skin with or without fever

- wheezing

- tightness in the chest or throat

- trouble breathing, swallowing, or talking

- unusual hoarseness

- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)

- red or irritated eyes

- or sores in your mouth, throat, nose, or eyes. Onset Forte side effects

1012. This particular disease, Colds & Flu, interacts with the drug Panatuss PED and has the follow

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe drowsiness
- a light-headed feeling, like you might pass out
- severe dizziness or anxiety
- or severe restless feeling or being unable to sleep. Common side effects of Panatuss PED may include: mild dizziness or drowsiness
- dry mouth, nose, or throat
- diarrhea , constipation , upset stomach
- feeling restless or excited (especially in children)
- blurred vision
- or sleep problems (insomnia).

1013. This particular disease, Colds & Flu, interacts with the drug Phena-Plus and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: confusion, hallucinations
- feeling nervous or irritable
- trouble sleeping
- little or no urination
- a seizure (convulsions)
- numbness or tingly feeling
- a light-headed feeling, like you might pass out
- pounding heartbeats or fluttering in your chest

Drug Side Effects Summary

- or increased blood pressure--severe headache, blurred vision, pounding in your neck or ears, anxiety , nosebleed. Side effects may be more likely in older adults. Common side effects of Phena-Plus may include: nausea , constipation , loss of appetite

- rash or itching
- increased sweating
- dry mouth, nose, or throat
- blurred vision
- headache, dizziness
- or feeling restless or excited (especially in children).

1014. This particular disease, Colds & Flu, interacts with the drug Phenagil and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Phenagil may cause serious side effects. Stop using Phenagil and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate
- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)
- little or no urinating
- easy bruising or bleeding, unusual weakness
- or dangerously high blood pressure (severe headache, buzzing in your ears, anxiety , shortness of breath). Common side effects of Phenagil may include: dizziness , drowsiness, blurred vision
- dry nose or mouth
- nausea , stomach pain, constipation , loss of appetite
- problems with memory or concentration
- or feeling restless or excited (especially in children).

Drug Side Effects Summary

1015. This particular disease, Colds & Flu, interacts with the drug pheniramine / phenylephrine and

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: nervousness
- dizziness
- or trouble sleeping. Common side effects of pheniramine and phenylephrine may include: drowsiness
- feeling restless or excited (especially in children)
- dry mouth, nose, or throat
- constipation
- or blurred vision, double vision.

1016. This particular disease, Colds & Flu, interacts with the drug phenylephrine / triprolidine and h

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness

Drug Side Effects Summary

- or swelling of the mouth, face, lips, tongue, or throat. Phenylephrine / triprolidine side effects

1017. This particular disease, Colds & Flu, interacts with the drug Robitussin Peak Cold Daytime C

- hives

- difficulty breathing

- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. Stop using Robitussin Peak Cold Daytime Cold + Flu and call your doctor at once if you have: chest pain, fast, slow, or uneven heart rate

- severe dizziness , feeling like you might pass out

- mood changes, confusion, hallucinations, seizure (convulsions)

- little or no urinating

- nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)

- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety , chest pain, uneven heartbeats, seizure). Common side effects of Robitussin Peak Cold Daytime Cold + Flu may include: dizziness, weakness, mild headache

- diarrhea , upset stomach

- dry mouth, nose, or throat

- feeling nervous, restless, irritable, or anxious

- or sleep problems (insomnia).

Drug Side Effects Summary

1018. This particular disease, Colds & Flu, interacts with the drug SuTan-DM and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: fast, slow, or uneven heart rate
- severe headache, mood changes, hallucinations
- severe dizziness or anxiety , feeling like you might pass out
- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- fever
- little or no urinating
- shortness of breath
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety, confusion, chest pain, uneven heartbeats, seizure). Common side effects of SuTan-DM may include: mild headache
- mild dizziness, drowsiness
- dry mouth, nose, or throat
- nausea , diarrhea , constipation , upset stomach
- feeling nervous, restless, or irritable
- blurred vision
- or sleep problems (insomnia).

1019. This particular disease, Colds & Flu, interacts with the drug Tannic-12 S and has the following

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Tannic-12 S may cause serious side effects. Stop using Tannic-12 S and call your doctor at once if you have: fast, pounding, or uneven heartbeat
- severe headache, hallucinations, seizure (convulsions)
- severe dizziness or drowsiness
- vision problems
- little or no urinating
- tremors, severe restlessness
- or seizures (convulsions). Common side effects of Tannic-12 S may include: constipation , diarrhea
- nausea , vomiting , upset stomach , loss of appetite
- weakness, mild dizziness or drowsiness
- anxiety , feeling nervous or excited
- or sleep problems (insomnia).

1020. This particular disease, Colds & Flu, interacts with the drug Theracaps Multi-Symptom Cough

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness

Drug Side Effects Summary

- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of high blood pressure like very bad headache or dizziness , passing out, or change in eyesight. Chest pain or pressure or a fast heartbeat . A heartbeat that does not feel normal. Very nervous and excitable. Shortness of breath. Shakiness. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)

- red or irritated eyes

- or sores in your mouth, throat, nose, or eyes. Theracaps Multi-Symptom Cough & Cold Reliever side effects

1021. This particular disease, Colds & Flu, interacts with the drug Triaminic Softchews Cough & So

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: redness or swelling

- new or worsening symptoms

- an ongoing cough with a rash or headache

Drug Side Effects Summary

- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Triaminic Cough and Sore Throat Softchews may include: upset stomach
- confusion
- or drowsiness.

1022. This particular disease, Colds & Flu, interacts with the drug Tricode GF and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Like other narcotic medications, codeine can slow your breathing. Death may occur if breathing becomes too weak. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Stop taking this medicine and call your doctor at once if you have: noisy breathing, sighing, shallow breathing
- severe drowsiness, hallucinations
- fast, slow, or uneven heart rate
- a seizure
- nervousness , tremors
- or little or no urination. Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Common side effects of Tricode GF may include: dizziness, drowsiness, weakness
- headache
- nausea , constipation
- sleep problems (insomnia)
- or feeling nervous, anxious, or restless.

Drug Side Effects Summary

1023. This particular disease, Colds & Flu, interacts with the drug Tusnel C and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Like other narcotic medications, codeine can slow your breathing. Death may occur if breathing becomes too weak. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Stop taking this medicine and call your doctor at once if you have: noisy breathing, sighing, shallow breathing
- severe drowsiness, hallucinations
- fast, slow, or uneven heart rate
- a seizure
- nervousness , tremors
- or little or no urination. Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Common side effects of Tusnel C may include: dizziness, drowsiness, weakness
- headache
- nausea , constipation
- sleep problems (insomnia)
- or feeling nervous, anxious, or restless.

1024. This particular disease, Colds & Flu, interacts with the drug Tuzistra XR and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should seek emergency medical attention if you have slow

Drug Side Effects Summary

breathing with long pauses, blue colored lips, or if you are hard to wake up. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep

- confusion, severe drowsiness
- a light-headed feeling, like you might pass out
- severe constipation
- or adrenal gland problems-- nausea , vomiting , loss of appetite, feeling very dizzy, weak, or tired.

Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea . Serious side effects may be more likely in older adults and those who are debilitated. Common side effects of Tuzistra XR may include: dizziness, drowsiness, feeling light-headed

- feeling short of breath
- confusion, mood changes, lack of energy, feeling nervous, irritable, agitated, or restless
- sleep problems
- headache , vision problems
- tremors, coordination problems
- dry mouth
- sweating
- or nausea, vomiting, constipation, stomach pain, bloating .

1025. This particular disease, Colds & Flu, interacts with the drug Tylenol Allergy Multi-Symptom and

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe

Drug Side Effects Summary

skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Tylenol Allergy Multi-Symptom and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate

- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)
- easy bruising or bleeding, unusual weakness
- little or no urinating
- nausea, pain in your upper stomach, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of your skin or eyes)
- or dangerously high blood pressure (severe headache, blurred vision, buzzing in your ears, anxiety, chest pain, shortness of breath, uneven heartbeats, seizure). Side effects such as dry mouth, constipation, and confusion may be more likely in older adults. Common side effects of Tylenol Allergy Multi-Symptom may include: dizziness, drowsiness
- mild headache, blurred vision
- dry mouth, nose, or throat
- constipation
- feeling nervous
- or sleep problems (insomnia)
-

1026. This particular disease, Colds & Flu, interacts with the drug Tylenol Children's Cold + Cough

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: redness or swelling
- new or worsening symptoms
- an ongoing cough with a rash or headache
- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Children's Tylenol Cold + Cough + Sore Throat may include: upset stomach
- confusion
- or drowsiness.

1027. This particular disease, Colds & Flu, interacts with the drug Tylenol Cough & Sore Throat Nig

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing

Drug Side Effects Summary

- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Tylenol Cough & Sore Throat Nighttime side effects

1028. This particular disease, Colds & Flu, interacts with the drug Tylenol Cough and Sore Throat D

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling

Drug Side Effects Summary

tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)

- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Tylenol Cough and Sore Throat Daytime side effects

1029. This particular disease, Colds & Flu, interacts with the drug Vanacof and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe drowsiness
- a light-headed feeling, like you might pass out
- severe dizziness or anxiety
- or severe restless feeling or being unable to sleep. Common side effects of Vanacof may include: mild dizziness or drowsiness
- dry mouth, nose, or throat
- diarrhea , constipation , upset stomach
- feeling restless or excited (especially in children)
- blurred vision
- or sleep problems (insomnia).

1030. This particular disease, Colds & Flu, interacts with the drug Vi-Sudo and has the following side effects:

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe dizziness
- nervousness
- trouble sleeping
- or high blood pressure (severe headache, pounding in your neck or ears). Common side effects of Vi-Sudo may include: dizziness, drowsiness
- dry mouth, nose, or throat
- constipation
- blurred vision
- or feeling restless or excited (especially in children).

1031. This particular disease, Colds & Flu, interacts with the drug Vicks Formula 44 Custom Care C

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling

Drug Side Effects Summary

tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)

- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Vicks Formula 44 Custom Care Cough & Cold PM side effects

1032. This particular disease, Colds & Flu, interacts with the drug Zodyl DAC and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad dizziness or passing out. Chest pain or pressure or a fast heartbeat . Trouble breathing, slow breathing, or shallow breathing. Noisy breathing. Feeling very sleepy. Feeling confused. A heartbeat that does not feel normal. Hallucinations (seeing or hearing things that are not there). Mood changes. Seizures . Very bad

Drug Side Effects Summary

belly pain. Very bad headache . Trouble passing urine. Shakiness. Change in eyesight. Very upset stomach or throwing up. Very bad constipation . Feeling very tired or weak. Zodyl DAC side effects

1033. This particular disease, Constipation, interacts with the drug docusate and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using docusate and call your doctor at once if: you have rectal bleeding
- no bowel movement occurs after using a laxative
- you need to use a stool softener for more than 1 week
- or rash occurs. Less serious side effects may be more likely, and you may have none at all.

1034. This particular disease, Constipation, interacts with the drug MiraLAX and has the following s

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop taking this medicine and call your doctor at once if you have: severe or bloody diarrhea
- rectal bleeding
- blood in your stools
- or severe and worsening stomach pain. Common MiraLAX side effects may include: bloating, gas, upset stomach
- dizziness
- or increased sweating.

1035. This particular disease, Constipation, interacts with the drug Colace and has the following sic

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if: you have rectal bleeding
- no bowel movement occurs after using a laxative
- you need to use a stool softener for more than 1 week
- or if a rash occurs. Less serious Colace side effects may be more likely, and you may have none at all.

1036. This particular disease, Constipation, interacts with the drug Dulcolax and has the following s

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Dulcolax and call your doctor at once if you have: rectal bleeding
- or no bowel movement after using Dulcolax. Common Dulcolax side effects may include: stomach pain or discomfort
- feeling light-headed
- or rectal burning.

1037. This particular disease, Constipation, interacts with the drug bisacodyl and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Bisacodyl may cause serious side effects. Stop using bisacodyl and call your doctor at once if you have: rectal bleeding
- or no bowel movement after using bisacodyl. Common side effects of bisacodyl may include:

Drug Side Effects Summary

stomach pain or discomfort

- feeling light-headed
- or rectal burning.

1038. This particular disease, Constipation, interacts with the drug polyethylene glycol 3350 and has

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop taking polyethylene glycol 3350 and call your doctor at once if you have: severe or bloody diarrhea
- rectal bleeding
- blood in your stools
- or severe and worsening stomach pain. Common side effects of polyethylene glycol 3350 may include: bloating , gas, upset stomach
- dizziness
- or increased sweating.

1039. This particular disease, Constipation, interacts with the drug Dulcolax Stool Softener and has

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Dulcolax Stool Softener and call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- a light-headed feeling, like you might pass out
- rectal bleeding or irritation
- numbness or a rash around your rectum
- vomiting, severe diarrhea or stomach cramps

Drug Side Effects Summary

- or continued constipation, or no bowel movement. Common Dulcolax Stool Softener side effects may include: dizziness , weakness
- gas, bloating , mild diarrhea
- rectal irritation
- or sweating.

1040. This particular disease, Constipation, interacts with the drug magnesium oxide and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Magnesium oxide may cause serious side effects. Stop using magnesium oxide and call your doctor at once if you have: rectal bleeding
- coughing up blood or vomit that looks like coffee grounds
- bloody or tarry stools
- or no bowel movement after using magnesium oxide as a laxative. Common side effects of magnesium oxide may include: diarrhea
- or upset stomach .

1041. This particular disease, Constipation, interacts with the drug magnesium citrate and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Magnesium citrate may cause serious side effects. Stop using magnesium citrate and call your doctor at once if you have: no bowel movement within 6 hours after taking the medicine
- pain with bowel movements, rectal bleeding
- watery diarrhea , nausea, vomiting, severe stomach pain

Drug Side Effects Summary

- painful or difficult urination
- flushing (warmth, redness, or tingly feeling)
- a light-headed feeling, like you might pass out
- weak or shallow breathing, slow heartbeats
- or muscle weakness, increased thirst. Common side effects of magnesium citrate may include:
loose stools, diarrhea, stomach cramps
- upset stomach
- dizziness
- or increased sweating.

1042. This particular disease, Constipation, interacts with the drug Metamucil and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Metamucil may cause serious side effects. Stop using Metamucil and call your doctor at once if you have: constipation that lasts longer than 7 days
- rectal bleeding
- or severe stomach pain. Common side effects of Metamucil may include: bloating
- or minor change in your bowel habits.

1043. This particular disease, Constipation, interacts with the drug Citrucel and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Citrucel may cause serious side effects. Call your doctor at once if you have: severe stomach cramps, rectal bleeding
- or no bowel movement within 3 days after using Citrucel.

Drug Side Effects Summary

1044. This particular disease, Constipation, interacts with the drug Doc-Q-Lace and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Doc-Q-Lace and call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- a light-headed feeling, like you might pass out
- rectal bleeding or irritation
- numbness or a rash around your rectum
- vomiting, severe diarrhea or stomach cramps
- or continued constipation, or no bowel movement. Common Doc-Q-Lace side effects may include: dizziness , weakness
- gas, bloating , mild diarrhea
- rectal irritation
- or sweating.

1045. This particular disease, Constipation, interacts with the drug FiberCon and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. FiberCon may cause serious side effects. Call your doctor at once if you have: severe stomach cramps, rectal bleeding
- or no bowel movement within 3 days after using FiberCon. Common side effects of FiberCon may include: mild stomach pain
- bloating
- or gas.

Drug Side Effects Summary

1046. This particular disease, Constipation, interacts with the drug DOK and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- a light-headed feeling, like you might pass out
- rectal bleeding or irritation
- numbness or a rash around your rectum
- vomiting, severe diarrhea or stomach cramps
- or continued constipation, or no bowel movement. Common DOK side effects may include: dizziness , weakness
- gas, bloating , mild diarrhea
- rectal irritation
- or sweating.

1047. This particular disease, Constipation, interacts with the drug Fleet Enema and has the followi

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: no bowel movement within 30 minutes after use
- rectal bleeding or bright red bowel movements
- or dehydration symptoms-- dizziness , vomiting, thirst, little or no urinating. Common side effects of Fleet Enema may include: rectal discomfort.

Drug Side Effects Summary

1048. This particular disease, Constipation, interacts with the drug GlycoLax and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop taking GlycoLax and call your doctor at once if you have: severe or bloody diarrhea
- rectal bleeding
- blood in your stools
- or severe and worsening stomach pain. Common side effects of GlycoLax may include: bloating , gas, upset stomach
- dizziness
- or increased sweating.

1049. This particular disease, Constipation, interacts with the drug psyllium and has the following s

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Psyllium may cause serious side effects. Stop using psyllium and call your doctor at once if you have: constipation that lasts longer than 7 days
- rectal bleeding
- or severe stomach pain. Common side effects of psyllium may include: bloating
- or minor change in your bowel habits.

1050. This particular disease, Constipation, interacts with the drug Surfak and has the following side

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs

Drug Side Effects Summary

of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Surfak side effects

1051. This particular disease, Constipation, interacts with the drug Enemeez Mini and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Rectal irritation. Enemeez Mini side effects

1052. This particular disease, Constipation, interacts with the drug Bisa-Lax and has the following side effects:

Drug Side Effects Summary

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Bisa-Lax side effects

1053. This particular disease, Constipation, interacts with the drug ClearLax and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop taking ClearLax and call your doctor at once if you have: severe or bloody diarrhea
- rectal bleeding
- blood in your stools
- or severe and worsening stomach pain. Common side effects of ClearLax may include: bloating , gas, upset stomach
- dizziness
- or increased sweating.

Drug Side Effects Summary

1054. This particular disease, Constipation, interacts with the drug Colace Micro-Enema and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Rectal irritation. Colace Micro-Enema side effects

1055. This particular disease, Constipation, interacts with the drug Correctol and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat

Drug Side Effects Summary

- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Correctol side effects

1056. This particular disease, Constipation, interacts with the drug Diocto and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Diocto side effects

1057. This particular disease, Constipation, interacts with the drug Fleet Bisacodyl and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching

Drug Side Effects Summary

- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Fleet Bisacodyl side effects

1058. This particular disease, Constipation, interacts with the drug Phillips' Cramp-free and has the

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Phillips' Cramp-free and call your doctor at once if you have: rectal bleeding
- coughing up blood or vomit that looks like coffee grounds
- bloody or tarry stools
- or no bowel movement after using Phillips' Cramp-free as a laxative. Common side effects of Phillips' Cramp-free may include: diarrhea
- or upset stomach .

1059. This particular disease, Constipation, interacts with the drug Citrate of Magnesia and has the

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Citrate of Magnesia may cause serious side effects. Stop using Citrate of Magnesia and call your doctor at once if you have: no bowel movement within 6 hours after taking the medicine
- pain with bowel movements, rectal bleeding

Drug Side Effects Summary

- watery diarrhea , nausea, vomiting, severe stomach pain
- painful or difficult urination
- flushing (warmth, redness, or tingly feeling)
- a light-headed feeling, like you might pass out
- weak or shallow breathing, slow heartbeats
- or muscle weakness, increased thirst. Common side effects of Citrate of Magnesia may include:
loose stools, diarrhea, stomach cramps
- upset stomach
- dizziness
- or increased sweating.

1060. This particular disease, Constipation, interacts with the drug GaviLAX and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop taking GaviLAX and call your doctor at once if you have: severe or bloody diarrhea
- rectal bleeding
- blood in your stools
- or severe and worsening stomach pain. Common side effects of GaviLAX may include: bloating ,
gas, upset stomach
- dizziness
- or increased sweating.

1061. This particular disease, Constipation, interacts with the drug Gialax and has the following side effects:

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop taking Gialax and call your doctor at once if you have: severe or bloody diarrhea
- rectal bleeding
- blood in your stools
- or severe and worsening stomach pain. Common side effects of Gialax may include: bloating , gas, upset stomach
- dizziness
- or increased sweating.

1062. This particular disease, Constipation, interacts with the drug Mag-Oxide and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Severe diarrhea . Mag-Oxide side effects

1063. This particular disease, Constipation, interacts with the drug PEG3350 and has the following

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop taking PEG3350 and call your doctor at once if you have: severe or bloody diarrhea
- rectal bleeding
- blood in your stools
- or severe and worsening stomach pain. Common side effects of PEG3350 may include: bloating , gas, upset stomach
- dizziness
- or increased sweating.

1064. This particular disease, Constipation, interacts with the drug Purelax and has the following side effects:

- No side effects listed

1065. This particular disease, Constipation, interacts with the drug SunMark ClearLax and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop taking SunMark ClearLax and call your doctor at once if you have: severe or bloody diarrhea
- rectal bleeding
- blood in your stools
- or severe and worsening stomach pain. Common side effects of SunMark ClearLax may include: bloating , gas, upset stomach
- dizziness
- or increased sweating.

Drug Side Effects Summary

1066. This particular disease, Constipation, interacts with the drug Uro-Mag and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Uro-Mag may cause serious side effects. Stop using Uro-Mag and call your doctor at once if you have: rectal bleeding
- coughing up blood or vomit that looks like coffee grounds
- bloody or tarry stools
- or no bowel movement after using Uro-Mag as a laxative. Common side effects of Uro-Mag may include: diarrhea
- or upset stomach .

1067. This particular disease, Constipation, interacts with the drug Citroma and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Citroma may cause serious side effects. Stop using Citroma and call your doctor at once if you have: no bowel movement within 6 hours after taking the medicine
- pain with bowel movements, rectal bleeding
- watery diarrhea , nausea, vomiting, severe stomach pain
- painful or difficult urination
- flushing (warmth, redness, or tingly feeling)
- a light-headed feeling, like you might pass out
- weak or shallow breathing, slow heartbeats
- or muscle weakness, increased thirst. Common side effects of Citroma may include: loose stools, diarrhea, stomach cramps

Drug Side Effects Summary

- upset stomach
- dizziness
- or increased sweating.

1068. This particular disease, Constipation, interacts with the drug glycerin and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Glycerin rectal may cause serious side effects. Stop using glycerin rectal and call your doctor at once if you have: severe stomach pain or cramping
- rectal bleeding
- or no bowel movement within 1 hour after use. Common side effects of glycerin may include: loose stools
- nausea or stomach discomfort
- or rectal pain or burning.

1069. This particular disease, Constipation, interacts with the drug Hydrocil and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Hydrocil may cause serious side effects. Stop using Hydrocil and call your doctor at once if you have: constipation that lasts longer than 7 days
- rectal bleeding
- or severe stomach pain. Common side effects of Hydrocil may include: bloating
- or minor change in your bowel habits.

1070. This particular disease, Constipation, interacts with the drug Konsyl and has the following side effects:

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Konsyl may cause serious side effects. Stop using Konsyl and call your doctor at once if you have: constipation that lasts longer than 7 days
- rectal bleeding
- or severe stomach pain. Common side effects of Konsyl may include: bloating
- or minor change in your bowel habits.

1071. This particular disease, Constipation, interacts with the drug methylcellulose and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Methylcellulose may cause serious side effects. Call your doctor at once if you have: severe stomach cramps, rectal bleeding
- or no bowel movement within 3 days after using methylcellulose.

1072. This particular disease, Constipation, interacts with the drug Reguloid and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Reguloid may cause serious side effects. Stop using Reguloid and call your doctor at once if you have: constipation that lasts longer than 7 days
- rectal bleeding
- or severe stomach pain. Common side effects of Reguloid may include: bloating
- or minor change in your bowel habits.

1073. This particular disease, Constipation, interacts with the drug sodium biphosphate / sodium p

Drug Side Effects Summary

- hives
- dizziness
- wheezing, difficult breathing
- swelling of your face, lips, tongue, or throat. sodium biphosphate and sodium phosphate may cause serious side effects. Stop using sodium biphosphate and sodium phosphate and call your doctor at once if you have: no bowel movement after using this medicine
- severe stomach pain, rectal bleeding or bright red bowel movements
- fast, slow, or uneven heart rate
- little or no urination
- a seizure (black-out or convulsions)
- or headache , dizziness, and vomiting . Common side effects of sodium biphosphate and sodium phosphate may include: bloating
- nausea , vomiting
- or stomach pain.

1074. This particular disease, Constipation, interacts with the drug Disposable Enema and has the f

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: no bowel movement within 30 minutes after use
- rectal bleeding or bright red bowel movements
- or dehydration symptoms-- dizziness , vomiting, thirst, little or no urinating. Common side effects of Disposable Enema may include: rectal discomfort.

Drug Side Effects Summary

1075. This particular disease, Constipation, interacts with the drug Equalactin and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Equalactin may cause serious side effects. Call your doctor at once if you have: severe stomach cramps, rectal bleeding
- or no bowel movement within 3 days after using Equalactin. Common side effects of Equalactin may include: mild stomach pain
- bloating
- or gas.

1076. This particular disease, Constipation, interacts with the drug Fiber Lax and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Fiber Lax may cause serious side effects. Call your doctor at once if you have: severe stomach cramps, rectal bleeding
- or no bowel movement within 3 days after using Fiber Lax. Common side effects of Fiber Lax may include: mild stomach pain
- bloating
- or gas.

1077. This particular disease, Constipation, interacts with the drug Fiber Laxative and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

Drug Side Effects Summary

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Chest pain or pressure. Upset stomach or throwing up. Trouble swallowing. Trouble breathing. Belly pain. Fiber Laxative side effects

1078. This particular disease, Constipation, interacts with the drug Fleet Glycerin Suppositories Adult

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Fleet Glycerin Suppositories Adult rectal may cause serious side effects. Stop using Fleet Glycerin Suppositories Adult rectal and call your doctor at once if you have: severe stomach pain or cramping
- rectal bleeding
- or no bowel movement within 1 hour after use. Common side effects of Fleet Glycerin Suppositories Adult may include: loose stools
- nausea or stomach discomfort
- or rectal pain or burning.

1079. This particular disease, Constipation, interacts with the drug Fleet Phospho Soda and has the

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your

Drug Side Effects Summary

doctor immediately if any of the following side effects occur: More common Bloating stomach pain
Incidence not known Agitation bloody or cloudy urine burning, crawling, itching, numbness, prickling,
"pins and needles", or tingling feelings confusion cough decrease in the frequency or amount of
urine decreased awareness or responsiveness depression difficult or labored breathing difficult or
painful urination difficulty with swallowing dizziness fainting fast, slow, or irregular heartbeat
headache hives, itching, skin rash hostility increased blood pressure increased thirst irritability large,
hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or genitals loss of
appetite loss of consciousness lower back or side pain muscle twitching nausea noisy breathing
puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rapid weight gain
redness of the skin seizures severe sleepiness sudden decrease in the amount of urine swelling of
the face, ankles, or hands tightness in the chest unusual drowsiness, dullness, tiredness, weakness,
or feeling of sluggishness vomiting Symptoms of overdose Blurred vision chest pain or discomfort
decreased frequency of urine dizziness, faintness, or lightheadedness when getting up suddenly
from a lying or sitting position dry mouth increase in heart rate increased thirst lightheadedness loss
of appetite lower back or side pain muscle cramps in the hands, arms, feet, legs, or face muscle
pain pounding or rapid pulse rapid breathing restlessness stomach cramps sunken eyes sweating
tremor weakness weight gain wrinkled skin Other side effects not listed may also occur in some
patients. If you notice any other effects, check with your healthcare professional. Call your doctor for
medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Fleet
Phospho Soda side effects

1080. This particular disease, Constipation, interacts with the drug mineral oil and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Mineral oil may cause serious side effects. Stop using

Drug Side Effects Summary

mineral oil and call your doctor at once if you have: no bowel movement after use

- rectal bleeding
- blood in your stools
- or severe and worsening stomach pain, cramping, or bloating . Common side effects of mineral oil may include: mild diarrhea
- gas
- or stomach cramps.

1081. This particular disease, Constipation, interacts with the drug Natural Fiber Therapy and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Natural Fiber Therapy may cause serious side effects. Stop using Natural Fiber Therapy and call your doctor at once if you have: constipation that lasts longer than 7 days
- rectal bleeding
- or severe stomach pain. Common side effects of Natural Fiber Therapy may include: bloating
- or minor change in your bowel habits.

1082. This particular disease, Constipation, interacts with the drug OsmoPrep and has the following side effects:

- hives
- dizziness
- wheezing, difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: no bowel movement after using this medicine
- severe stomach pain, rectal bleeding or bright red bowel movements

Drug Side Effects Summary

- fast, slow, or uneven heart rate
- little or no urination
- a seizure (black-out or convulsions)
- or headache , dizziness, and vomiting . Common OsmoPrep side effects may include: bloating
- nausea , vomiting
- or stomach pain.

1083. This particular disease, Constipation, interacts with the drug Pedia-Lax Chewable Tablets and

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Pedia-Lax Chewable may cause serious side effects. Stop using Pedia-Lax Chewable and call your doctor at once if you have: severe nausea , vomiting , or diarrhea
- no bowel movement after using the medicine as a laxative
- rectal bleeding
- or worsening symptoms. Common side effects of Pedia-Lax Chewable may include: diarrhea
- or a decreased sense of taste.

1084. This particular disease, Constipation, interacts with the drug Pedia-Lax Liquid Glycerin Supp

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Pedia-Lax Liquid may cause serious side effects. Stop using Pedia-Lax Liquid and call your doctor at once if you have: severe stomach pain or cramping
- rectal bleeding
- or no bowel movement within 1 hour after use. Common side effects of Pedia-Lax Liquid may

Drug Side Effects Summary

include: loose stools

- nausea or stomach discomfort
- or rectal pain or burning.

1085. This particular disease, Constipation, interacts with the drug polycarbophil and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Polycarbophil may cause serious side effects. Call your doctor at once if you have: severe stomach cramps, rectal bleeding
- or no bowel movement within 3 days after using polycarbophil. Common side effects of polycarbophil may include: mild stomach pain
- bloating
- or gas.

1086. This particular disease, Constipation, interacts with the drug Sani-Supp and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Sani-Supp may cause serious side effects. Stop using Sani-Supp and call your doctor at once if you have: severe stomach pain or cramping
- rectal bleeding
- or no bowel movement within 1 hour after use. Common side effects of Sani-Supp may include: loose stools
- nausea or stomach discomfort
- or rectal pain or burning.

Drug Side Effects Summary

1087. This particular disease, Constipation, interacts with the drug sorbitol and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using sorbitol and call your doctor at once if you have a serious side effect such as: severe stomach cramps
- vomiting
- severe diarrhea
- rectal bleeding
- black, bloody, or tarry stools
- weakness, dizziness
- or frequent urge to have a bowel movement. Less serious side effects may include: gas, mild nausea or stomach cramps
- or rectal irritation.

1088. This particular disease, Constipation, interacts with the drug benzocaine / docusate and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking

Drug Side Effects Summary

- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Rectal irritation. Benzocaine / docusate topical side effects

1089. This particular disease, Constipation, interacts with the drug castor oil and has the following

- No side effects listed

1090. This particular disease, Constipation, interacts with the drug Fibertab and has the following s

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Fibertab may cause serious side effects. Call your doctor at once if you have: severe stomach cramps, rectal bleeding
- or no bowel movement within 3 days after using Fibertab. Common side effects of Fibertab may include: mild stomach pain
- bloating
- or gas.

1091. This particular disease, Constipation, interacts with the drug Fleet Glycerin Suppositories Pe

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Fleet Glycerin Suppositories Pediatric rectal may cause serious side effects. Stop using Fleet Glycerin Suppositories Pediatric rectal and call your doctor at once if you have: severe stomach pain or cramping
- rectal bleeding
- or no bowel movement within 1 hour after use. Common side effects of Fleet Glycerin

Drug Side Effects Summary

Suppositories Pediatric may include: loose stools

- nausea or stomach discomfort
- or rectal pain or burning.

1092. This particular disease, Constipation, interacts with the drug potassium phosphate and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tell your caregivers right away if you have any signs of electrolyte imbalance, such as: confusion, severe weakness
- a light-headed feeling, like you might pass out
- nausea , chest pain, irregular heartbeats
- numbness or tingling in your arms or legs
- weakness or heavy feeling in your legs
- loss of movement in any part of your body
- or slow heart rate , weak pulse, fainting, slow breathing.

1093. This particular disease, COPD, interacts with the drug Symbicort and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: worsened breathing problems
- sores or white patches in your mouth and throat, pain when swallowing
- tremors, nervousness, chest pain, fast or pounding heartbeats
- cough with mucus, feeling short of breath
- wheezing, choking, or other breathing problems after using this medication

Drug Side Effects Summary

- blurred vision, tunnel vision, eye pain or redness, or seeing halos around lights
 - flu symptoms - fever, chills, body aches, unusual tiredness
 - high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
 - low potassium level - leg cramps, constipation , irregular heartbeats, fluttering in your chest, numbness or tingling, muscle weakness or limp feeling
 - or signs of a hormonal disorder - tiredness or weakness, feeling light-headed, nausea , vomiting .
- Budesonide can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common Symbicort side effects may include: throat pain or irritation
- white patches in your mouth or throat
 - stomach discomfort, vomiting
 - back pain , headache
 - flu symptoms
 - or cold symptoms such as stuffy or runny nose , sneezing, sinus pain, sore throat .

1094. This particular disease, COPD, interacts with the drug Breztri Aerosphere and has the following

- Breztri Aerosphere can cause serious side effects, including: fungal infection in your mouth or throat (thrush). Rinse your mouth with water without swallowing after using Breztri Aerosphere to help reduce your chance of getting thrush. pneumonia. People with COPD have a higher chance of getting pneumonia. Breztri Aerosphere may increase your chance of getting pneumonia. Call your healthcare provider if you notice any of the following symptoms: increase in mucus (sputum) production change in mucus color fever chills increased cough increased breathing problems weakened immune system and increased chance of getting infections (immunosuppression). reduced adrenal function (adrenal insufficiency). Adrenal insufficiency is a condition where the adrenal glands do not make enough steroid hormones. This can happen when you stop taking oral corticosteroid medicines (such as prednisone) and start taking a medicine containing an ICS (such

Drug Side Effects Summary

as Breztri Aerosphere). During this transition period, when your body is under stress from fever, trauma (such as a car accident), infection, surgery, or worse COPD symptoms, adrenal insufficiency can get worse and may cause death. Symptoms of adrenal insufficiency include: feeling tired lack of energy weakness nausea and vomiting low blood pressure (hypotension) sudden breathing problems immediately after inhaling your medicine. If you have sudden breathing problems immediately after inhaling your medicine, stop taking Breztri Aerosphere and call your healthcare provider right away. serious allergic reactions. Call your healthcare provider or get emergency medical care if you get any of the following symptoms of a serious allergic reaction: rash hives swelling of your face, mouth, and tongue breathing problems effects on your heart. increased blood pressure a fast or irregular heartbeat chest pain effects on your nervous system. tremor nervousness bone thinning or weakness (osteoporosis). new or worsened eye problems including acute narrow-angle glaucoma and cataracts. Acute narrow-angle glaucoma can cause permanent loss of vision if not treated. Symptoms of acute narrow-angle glaucoma may include: eye pain or discomfort nausea or vomiting blurred vision seeing halos or bright colors around lights red eyes If you have these symptoms, call your healthcare provider right away before taking another dose. urinary retention. People who take Breztri Aerosphere may develop new or worsening urinary retention. Symptoms of urinary retention may include: difficulty urinating urinating frequently painful urination urination in a weak stream or drips If you have these symptoms of urinary retention, stop taking Breztri Aerosphere and call your healthcare provider right away before taking another dose. changes in laboratory blood values , including high levels of blood sugar (hyperglycemia) and low levels of potassium (hypokalemia). Low levels of potassium may cause symptoms of muscle spasm, muscle weakness, or abnormal heart rhythm. Common side effects include: upper respiratory tract infection pneumonia back pain thrush in your mouth and throat. Rinse your mouth with water without swallowing after use to help prevent this. joint pain flu headache high blood sugar levels muscle spasms cough inflammation of the sinuses diarrhea hoarseness painful and frequent urination (signs

Drug Side Effects Summary

of a urinary tract infection) nausea difficulty sleeping feeling anxious awareness of your heart beating (palpitations) Tell your healthcare provider about any side effect that bothers you or that does not go away. These are not all the possible side effects. Ask your healthcare provider or pharmacist for more information. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800- FDA-1088. You may also report side effects to AstraZeneca at 1-800-236-9933. Breztri Aerosphere side effects

1095. This particular disease, COPD, interacts with the drug Anoro Ellipta and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: wheezing, choking, or other breathing problems after using this medicine
- painful or difficult urination, or urinating more often
- blurred vision, tunnel vision, eye pain or redness, or seeing halos around lights
- high blood sugar - increased thirst or urination, hunger, dry mouth, fruity breath odor
- or low potassium level - leg cramps, constipation , irregular heartbeats, fluttering in your chest, numbness or tingling, muscle weakness or limp feeling. Common Anoro Ellipta side effects may include: stuffy nose, sinus pain, sore throat
- chest pain, cough
- diarrhea , constipation
- muscle spasms
- or pain in your neck, arms, or legs.

1096. This particular disease, COPD, interacts with the drug Daliresp and has the following side effects:

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, depression, trouble sleeping, or if you feel impulsive, or you have thoughts about suicide or hurting yourself. Call your doctor at once if you have rapid and unintended weight loss. rapid and unintended weight loss
- tremors
- pain or burning when you urinate
- Common Daliresp side effects may include: nausea, diarrhea
- loss of appetite, minor weight loss
- headache, dizziness
- occasional sleep problems
- back pain
- or flu symptoms.

1097. This particular disease, COPD, interacts with the drug budesonide / formoterol and has the fo

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Budesonide and formoterol may cause serious side effects. Call your doctor at once if you have: worsened breathing problems
- sores or white patches in your mouth and throat, pain when swallowing
- tremors, nervousness , chest pain, fast or pounding heartbeats
- cough with mucus, feeling short of breath
- wheezing, choking, or other breathing problems after using this medication
- blurred vision, tunnel vision, eye pain or redness, or seeing halos around lights
- flu symptoms--fever, chills, body aches, unusual tiredness

Drug Side Effects Summary

- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
 - low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, numbness or tingling, muscle weakness or limp feeling
 - or signs of a hormonal disorder--tiredness or weakness, feeling light-headed, nausea , vomiting .
- Budesonide can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using budesonide and formoterol. Common side effects of budesonide and formoterol may include: throat pain or irritation
- white patches in your mouth or throat
 - stomach discomfort, vomiting
 - back pain , headache
 - flu symptoms
 - or cold symptoms such as stuffy or runny nose , sneezing, sinus pain, sore throat .

1098. This particular disease, COPD, interacts with the drug Tudorza Pressair and has the following

- hives, itching, rash, wheezing, chest tightness, trouble breathing, swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: wheezing, choking, or other breathing problems after using this medication
- blurred vision, nausea, vomiting, eye pain or redness, or seeing halos or bright colors around lights
- increased urination, painful or difficult urination
- little or no urinating
- or worsening or no improvement in your symptoms. Common Tudorza Pressair side effects may include: stuffy nose, sore throat, sinus pain
- cough
- or headache.

Drug Side Effects Summary

1099. This particular disease, COPD, interacts with the drug Incruse Ellipta and has the following side effects:

- hives , rash, severe itching
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: wheezing, choking, or other breathing problems after using this medicine
- blurred vision, nausea , vomiting , eye pain or redness, or seeing halos or bright colors around lights
- or painful or difficult urination, or urinating more often. Common Incruse Ellipta side effects may include: cold symptoms such as stuffy nose, sneezing, sore throat , cough
- fast or irregular heartbeats
- bruising, darkened skin
- mouth pain, tooth pain
- muscle or joint pain
- stomach pain
- or altered sense of taste.

1100. This particular disease, COPD, interacts with the drug Stiolto Respimat and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- wheezing, choking, or other breathing problems after using this medicine
- painful or difficult urination, or urinating more often
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar - increased thirst, increased urination, hunger, dry mouth, fruity breath odor

Drug Side Effects Summary

- or low potassium level - leg cramps, constipation , irregular heartbeats, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common Stiolto Respimat side effects may include: runny nose
- cough
- or back pain .

1101. This particular disease, COPD, interacts with the drug acclidinium and has the following side effects

- hives , itching, rash
- wheezing, chest tightness, trouble breathing
- swelling of your face, lips, tongue, or throat. Acclidinium may cause serious side effects. Call your doctor at once if you have: wheezing, choking, or other breathing problems after using this medication
- blurred vision, nausea , vomiting , eye pain or redness, or seeing halos or bright colors around lights
- increased urination, painful or difficult urination
- little or no urinating
- or worsening or no improvement in your symptoms. Common side effects of acclidinium may include: stuffy nose, sore throat , sinus pain
- cough
- or headache .

1102. This particular disease, COPD, interacts with the drug roflumilast and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your

Drug Side Effects Summary

doctor, such as: mood or behavior changes, anxiety , depression, trouble sleeping, or if you feel impulsive, or you have thoughts about suicide or hurting yourself. Call your doctor at once if you have rapid and unintended weight loss. rapid and unintended weight loss

- tremors
- pain or burning when you urinate
- Common side effects of roflumilast may include: nausea , diarrhea
- loss of appetite, minor weight loss
- headache , dizziness
- occasional sleep problems
- back pain
- or flu symptoms.

1103. This particular disease, COPD, interacts with the drug umeclidinium / vilanterol and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Umeclidinium and vilanterol may cause serious side effects. Call your doctor at once if you have: wheezing, choking, or other breathing problems after using umeclidinium and vilanterol
- painful or difficult urination, or urinating more often
- blurred vision, tunnel vision, eye pain or redness, or seeing halos around lights
- high blood sugar--increased thirst or urination, hunger, dry mouth, fruity breath odor
- or low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, numbness or tingling, muscle weakness or limp feeling. Common side effects of umeclidinium and vilanterol may include: stuffy nose, sinus pain, sore throat
- chest pain, cough

Drug Side Effects Summary

- diarrhea , constipation
- muscle spasms
- or pain in your neck, arms, or legs.

1104. This particular disease, COPD, interacts with the drug Bevespi Aerosphere and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: tremors, nervousness, chest pain, fast or pounding heartbeats
- wheezing, choking, or other breathing problems after using this medicine
- painful or difficult urination
- blurred vision, tunnel vision, eye pain or redness, or seeing halos around lights
- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- or low potassium level - leg cramps, constipation, irregular heartbeats, fluttering in your chest, numbness or tingling, muscle weakness or limp feeling. Common Bevespi side effects may include: cough
- or urination problems.

1105. This particular disease, COPD, interacts with the drug olodaterol / tiotropium and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Olodaterol and tiotropium may cause serious side effects. Call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- wheezing, choking, or other breathing problems after using this medicine
- painful or difficult urination, or urinating more often

Drug Side Effects Summary

- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar--increased thirst, increased urination, hunger, dry mouth, fruity breath odor
- or low potassium level--leg cramps, constipation , irregular heartbeats, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of olodaterol and tiotropium may include: runny nose
- cough
- or back pain .

1106. This particular disease, COPD, interacts with the drug Striverdi Respimat and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: tremors, nervousness, chest pain, fast or pounding heartbeats
- wheezing, choking, or other breathing problems after using this medicine
- worsening breathing problems
- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- or low potassium level - leg cramps, constipation , irregular heartbeats, fluttering in your chest, numbness or tingling, muscle weakness or limp feeling. Common Striverdi Respimat side effects may include: cold symptoms such as stuffy nose, sneezing, sore throat
- cough, shortness of breath, chest discomfort
- diarrhea
- dizziness
- pain or burning when you urinate
- back pain , joint pain
- or rash.

Drug Side Effects Summary

1107. This particular disease, COPD, interacts with the drug olodaterol and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Olodaterol may cause serious side effects. Call your doctor at once if you have: tremors, nervousness, chest pain, fast or pounding heartbeats
- wheezing, choking, or other breathing problems after using this medicine
- worsening breathing problems
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, numbness or tingling, muscle weakness or limp feeling. Common side effects of olodaterol may include: cold symptoms such as stuffy nose, sneezing, sore throat
- cough, shortness of breath, chest discomfort
- diarrhea
- dizziness
- pain or burning when you urinate
- back pain , joint pain
- or rash.

1108. This particular disease, COPD, interacts with the drug revefenacin and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Revefenacin may cause serious side effects. Call your doctor at once if you have: wheezing, choking, or other breathing problems after using revefenacin
- blurred vision, tunnel vision, eye pain or redness, or seeing halos around lights
- painful or difficult urination

Drug Side Effects Summary

- or trouble emptying your bladder (a weak stream of urine). Common side effects of revefenacin may include: headache
- back pain
- or cold symptoms such as runny nose , stuffy nose, sneezing, cough, sore throat .

1109. This particular disease, COPD, interacts with the drug umeclidinium and has the following side effects:

- hives , rash, severe itching
- difficult breathing
- swelling of your face, lips, tongue, or throat. Umeclidinium may cause serious side effects. Call your doctor at once if you have: wheezing, choking, or other breathing problems after using umeclidinium
- blurred vision, nausea , vomiting , eye pain or redness, or seeing halos or bright colors around lights
- or painful or difficult urination, or urinating more often. Common side effects of umeclidinium may include: cold symptoms such as stuffy nose, sneezing, sore throat , cough
- fast or irregular heartbeats
- bruising, darkened skin
- mouth pain, tooth pain
- muscle or joint pain
- stomach pain
- or altered sense of taste.

1110. This particular disease, COPD, interacts with the drug Yupelri and has the following side effects:

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: wheezing, choking, or other breathing problems after using this medicine
- blurred vision, tunnel vision, eye pain or redness, or seeing halos around lights
- painful or difficult urination
- or trouble emptying your bladder (a weak stream of urine). Common Yupelri side effects may include: headache
- back pain
- or cold symptoms such as runny nose, sneezing, cough, sore throat .

1111. This particular disease, COPD, interacts with the drug acclidinium / formoterol and has the foll

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Acclidinium and formoterol may cause serious side effects. Call your doctor at once if you have: wheezing, choking, or other breathing problems after using acclidinium and formoterol
- worsening breathing problems
- tremors, nervousness , chest pain, fast or pounding heartbeats
- severe headache, pounding in your neck or ears
- painful or difficult urination, or urinating more often
- blurred vision, tunnel vision, eye pain or redness, or seeing halos around lights
- high blood sugar--increased thirst or urination, hunger, dry mouth, fruity breath odor
- or low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, numbness or tingling, muscle weakness or limp feeling. Common side effects of acclidinium and formoterol may include: headache
- back pain

Drug Side Effects Summary

- or cold symptoms such as stuffy nose, sneezing, sore throat .

1112. This particular disease, COPD, interacts with the drug Breyna and has the following side effects

- No side effects listed

1113. This particular disease, COPD, interacts with the drug budesonide / formoterol / glycopyrrolate

- hives

- difficulty breathing

- swelling of your face, lips, tongue, or throat. Budesonide, formoterol, and glycopyrrolate may cause serious side effects. Call your doctor at once if you have: worsened breathing problems

- wheezing, choking, or other breathing problems after using this medication

- tremors, nervousness , chest pain, fast or pounding heartbeats

- sores or white patches in your mouth and throat, pain when swallowing

- flu symptoms, body aches, unusual tiredness

- painful or burning when you urinate, little or no urination

- blurred vision, tunnel vision, eye pain or redness, or seeing halos around lights

- low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling

- signs of a lung infection--fever, chills, cough with mucus, chest pain, feeling short of breath

- or signs of a hormonal disorder--tiredness or weakness, feeling light-headed, nausea , vomiting .

Common side effects of budesonide, formoterol, and glycopyrrolate may include: throat pain or irritation

- white patches in your mouth or throat

- headache , joint pain, muscle spasms

- pounding heartbeats, feeling anxious

Drug Side Effects Summary

- high blood sugar levels
- painful or difficult urination
- nausea, diarrhea
- cough, hoarse voice
- flu symptoms
- or cold symptoms such as stuffy or runny nose , sneezing, sinus pain, sore throat .

1114. This particular disease, COPD, interacts with the drug Duaklir Pressair and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: wheezing, choking, or other breathing problems after using this medicine
- worsening breathing problems
- tremors, nervousness, chest pain, fast or pounding heartbeats
- severe headache, pounding in your neck or ears
- painful or difficult urination, or urinating more often
- blurred vision, tunnel vision, eye pain or redness, or seeing halos around lights
- high blood sugar - increased thirst or urination, hunger, dry mouth, fruity breath odor
- or low potassium level - leg cramps, constipation , irregular heartbeats, fluttering in your chest, numbness or tingling, muscle weakness or limp feeling. Common Duaklir Pressair side effects may include: headache
- back pain
- or cold symptoms such as stuffy nose, sneezing, sore throat .

1115. This particular disease, COPD, interacts with the drug formoterol / glycopyrrolate and has the

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Formoterol and glycopyrrolate may cause serious side effects. Call your doctor at once if you have: tremors, nervousness , chest pain, fast or pounding heartbeats
- wheezing, choking, or other breathing problems after using this medicine
- painful or difficult urination
- blurred vision, tunnel vision, eye pain or redness, or seeing halos around lights
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or low potassium level--leg cramps, constipation, irregular heartbeats, fluttering in your chest, numbness or tingling, muscle weakness or limp feeling. Common side effects of formoterol and glycopyrrolate may include: cough
- or urination problems.

1116. This particular disease, Covid 19, interacts with the drug ivermectin and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: eye pain or redness, puffy eyes, problems with your vision
- severe skin rash , itching, or rash with pus
- confusion, change in your mental status, balance problems, trouble walking
- fever, swollen glands, stomach pain, joint pain, swelling in your hands or feet
- fast heart rate , trouble breathing
- loss of bladder or bowel control
- neck or back pain , seizure (convulsions)

Drug Side Effects Summary

- or a light-headed feeling, like you might pass out. Common ivermectin side effects may include:
headache , muscle aches
- dizziness
- nausea , diarrhea
- or mild skin rash.

1117. This particular disease, Covid 19, interacts with the drug remdesivir and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some side effects may occur during or after the injection. Tell your caregiver right away if you have: severe headache , pounding in your neck or ears
- fast, slow, or pounding heartbeats
- wheezing, trouble breathing
- swelling in your face
- nausea
- fever, chills, or shivering
- itching, sweating
- or a light-headed feeling, like you might pass out
- Common side effects of remdesivir may include: nausea
- or abnormal liver function tests .

1118. This particular disease, Covid 19, interacts with the drug Pfizer-BioNTech COVID-19 Vaccine and has the following side effects:

- hives , rash
- dizziness , weakness, fast heartbeats

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Your vaccination provider may want to watch you for a short time after your shot, to make sure you don't have an allergic reaction. You will be treated quickly if you have a reaction right after you receive the vaccine. You should not receive this vaccine again if the first shot caused a severe allergic reaction. Your doctor or vaccination provider will determine if you can safely receive another COVID-19 vaccine. Not all possible side effects are known. Becoming infected with COVID-19 is much more dangerous to your health than receiving this vaccine. Some people receiving Pfizer-BioNTech COVID-19 Vaccine had inflammation of the heart muscle or the lining around the heart within a few days after receiving this vaccine, but the risk of this side effect is very low. Seek medical attention right away if you have: chest pain
- shortness of breath
- or fast or pounding heartbeats or fluttering in your chest. Not all possible side effects are known. Common Pfizer-BioNTech COVID-19 Vaccine side effects may include: fever, chills, swollen glands, not feeling well
- pain, redness, or swelling where the shot was given
- feeling tired
- or headache, muscle pain, joint pain.

1119. This particular disease, Covid 19, interacts with the drug Paxlovid and has the following side

- Possible side effects of Paxlovid are: Liver Problems. Tell your healthcare provider right away if you have any of these signs and symptoms of liver problems: loss of appetite, yellowing of your skin and the whites of eyes (jaundice), dark-colored urine, pale colored stools and itchy skin, stomach area (abdominal) pain. Resistance to HIV Medicines. If you have untreated HIV infection, Paxlovid may lead to some HIV medicines not working as well in the future. Other possible side effects include: altered sense of taste diarrhea high blood pressure muscle aches These are not all the

Drug Side Effects Summary

possible side effects of Paxlovid. Not many people have taken Paxlovid. Serious and unexpected side effects may happen. Paxlovid is still being studied, so it is possible that all of the risks are not known at this time. Paxlovid side effects

1120. This particular disease, Covid 19, interacts with the drug Actemra and has the following side

- hives
- chest pain, difficulty breathing, feeling like you might pass out
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe stomach cramps, bloating , diarrhea or constipation
- unusual bleeding - nosebleeds , bleeding gums, abnormal vaginal bleeding , any bleeding that will not stop, blood in your urine or stools, coughing up blood or vomit that looks like coffee grounds
- liver problems - loss of appetite, right-sided stomach pain, vomiting, confusion, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- signs of infection - fever, chills, aches, skin sores, diarrhea, burning when you urinate
- signs of tuberculosis: cough, shortness of breath, night sweats , loss of appetite, weight loss, and feeling very tired
- or signs of perforation (a hole or tear) in your stomach or intestines - fever, ongoing stomach pain, change in bowel habits. Common Actemra side effects may include: runny or stuffy nose, sinus pain, sore throat
- headache
- increased blood pressure
- abnormal liver function tests
- or pain, swelling, burning, or irritation where an injection was given.

1121. This particular disease, Covid 19, interacts with the drug tocilizumab and has the following si

Drug Side Effects Summary

- hives
- chest pain, difficulty breathing, feeling like you might pass out
- swelling of your face, lips, tongue, or throat. Tocilizumab may cause serious side effects. Call your doctor at once if you have: severe stomach cramps, bloating , diarrhea or constipation
- unusual bleeding-- nosebleeds , bleeding gums, abnormal vaginal bleeding , any bleeding that will not stop, blood in your urine or stools, coughing up blood or vomit that looks like coffee grounds
- liver problems--loss of appetite, right-sided stomach pain, vomiting, confusion, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- signs of infection--fever, chills, aches, skin sores, diarrhea, burning when you urinate
- signs of tuberculosis: cough, shortness of breath, night sweats , loss of appetite, weight loss, and feeling very tired
- or signs of perforation (a hole or tear) in your stomach or intestines--fever, ongoing stomach pain, change in bowel habits. Common side effects of tocilizumab may include: runny or stuffy nose, sinus pain, sore throat
- headache
- increased blood pressure
- abnormal liver function tests
- or pain, swelling, burning, or irritation where an injection was given.

1122. This particular disease, Covid 19, interacts with the drug anakinra and has the following side

- hives , sweating, severe itching
- wheezing, difficult breathing
- fast or pounding heartbeats
- dizziness , fainting
- swelling of your face, lips, tongue, or throat. If you are using anakinra for DIRA, you may have an

Drug Side Effects Summary

increased risk of allergic reactions , especially in the first weeks of treatment. Anakinra may cause serious side effects. Call your doctor at once if you have: fever or chills

- low white blood cell counts--mouth sores, skin sores, sore throat
- signs of tuberculosis: fever, cough, night sweats , loss of appetite, weight loss , and feeling very tired. Common side effects of anakinra may include: abnormal liver function tests
- rash
- worsening arthritis symptoms
- nausea , vomiting , diarrhea , stomach pain
- headache
- joint pain
- flu symptoms
- cold symptoms such as stuffy nose, sneezing, sore throat
- or redness, bruising, itching, stinging, or swelling where the injection was given.

1123. This particular disease, Covid 19, interacts with the drug Kineret and has the following side effects

- hives , sweating, severe itching
- wheezing, difficult breathing
- fast or pounding heartbeats
- dizziness , fainting
- swelling of your face, lips, tongue, or throat. If you are using Kineret for DIRA, you may have an increased risk of allergic reactions , especially in the first weeks of treatment. Call your doctor at once if you have: fever or chills
- low white blood cell counts - mouth sores, skin sores, sore throat
- or signs of tuberculosis - fever, cough, night sweats , loss of appetite, weight loss , and feeling very tired. Common Kineret side effects may include: abnormal liver function tests

Drug Side Effects Summary

- rash
- worsening arthritis symptoms
- nausea , vomiting , diarrhea , stomach pain
- headache
- joint pain
- flu symptoms
- cold symptoms such as stuffy nose, sneezing, sore throat
- or redness, bruising, itching, stinging, or swelling where the injection was given.

1124. This particular disease, Covid 19, interacts with the drug Moderna COVID-19 Vaccine and has

- hives , rash, dizziness , weakness, fast heartbeats
 - difficult breathing
 - swelling of your face, lips, tongue, or throat. Your vaccination provider may want to watch you for a short time after your shot, to make sure you don't have an allergic reaction. You will be treated quickly if you have a reaction right after you receive the vaccine. You should not receive this vaccine again if the first shot caused a severe allergic reaction. Your doctor or vaccination provider will determine if you can safely receive another COVID-19 vaccine. Becoming infected with COVID-19 is much more dangerous to your health than receiving Moderna COVID-19 Vaccine. Some people receiving Moderna COVID-19 Vaccine had inflammation of the heart muscle or the lining around the heart within a few days after receiving this vaccine, but the risk of this side effect is very low. Seek medical attention right away if you have: chest pain
 - shortness of breath
 - or fast or pounding heartbeats or fluttering in your chest. Not all possible side effects are known.
- Common Moderna COVID-19 Vaccine side effects may include: fever, chills
- redness or a hardness and swelling where the shot was given

Drug Side Effects Summary

- swelling or tenderness under your arm
- nausea, vomiting
- feeling tired
- or headache, muscle pain, joint pain.

1125. This particular disease, Covid 19, interacts with the drug Olumiant and has the following side effects:

- Stop taking this medicine and get emergency medical help if you have signs of an allergic reaction to Olumiant: hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some people taking Olumiant have developed heart attacks, strokes, or serious blood clots. Stop taking Olumiant and seek emergency medical attention if you have: sudden shortness of breath
- chest pain or pressure that may spread to your jaw, shoulder, arms, or back
- nausea, vomiting , cold sweat
- a light-headed feeling, like you might pass out
- weakness on one side of your body
- slurred speech, drooping on one side of your mouth
- or pain, swelling, or redness in an arm or a leg. Call your doctor at once if you have: fever, chills, night sweats, constant tiredness
- wheezing, trouble breathing, severe or worsening cough
- increased urination, pain or burning when you urinate
- unexplained weight loss
- lumps in your neck, armpits, or groin
- signs of shingles - flu-like symptoms, tingly or painful blistering rash on one side of your body
- a hole (perforation) in your digestive tract - fever, severe stomach pain, diarrhea or changes in

Drug Side Effects Summary

bowel habits

- or signs of hepatitis - loss of appetite, vomiting, stomach pain (upper right side), dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common Olumiant side effects may include: cold sores , shingles
- nausea
- or cold symptoms such as stuffy nose, sneezing, sore throat .

1126. This particular disease, Covid 19, interacts with the drug Veklury and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some side effects may occur during or after the injection. Tell your caregiver right away if you have: severe headache , pounding in your neck or ears
- fast, slow, or pounding heartbeats
- wheezing, trouble breathing
- swelling in your face
- nausea
- fever, chills, or shivering
- itching, sweating
- or a light-headed feeling, like you might pass out
- Common Veklury side effects may include include: nausea
- or abnormal liver function tests .

1127. This particular disease, Covid 19, interacts with the drug molnupiravir and has the following side effects

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor if you have any new or worsening symptoms. Common molnupiravir side effects may include: diarrhea , nausea
- or dizziness .

1128. This particular disease, Covid 19, interacts with the drug REGEN-COV and has the following s

- hives , itching
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some side effects may occur during the Regen-Cov injection. Tell your caregiver right away if you have: throat irritation, swelling in your face or throat
- dizziness , a light-headed feeling (like you might pass out)
- chest pain, wheezing, shortness of breath
- fever, chills, sweating, nausea , flushing (sudden warmth, redness, or tingly feeling)
- fast or slow heartbeats, headache , pounding in your neck or ears
- weakness, tiredness
- rash, itching
- or muscle pain. Call your doctor if you have new or worsening symptoms after the infusion, such as fever, confusion, weakness, tiredness, trouble breathing, or fast or slow heartbeats. Less serious side effects may also occur, or you may have none at all. Not all possible side effects are known. This is not a complete list of Regen-Cov's side effects and others may occur. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. What other drugs will affect Regen-Cov? Other drugs may interact with casirivimab and imdevimab, including prescription and over-the-counter medicines, vitamins , and herbal products . Tell your doctor about all other medicines you use. REGEN-COV drug interactions

Drug Side Effects Summary

1129. This particular disease, Covid 19, interacts with the drug sotrovimab and has the following side effects:

- hives , itching
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some side effects may occur during or up to 24 hours after the injection. Tell your medical caregivers right away if you have: throat irritation, swelling in your face or throat
- fast, slow, or uneven heart rate
- confusion, dizziness , a light-headed feeling (like you might pass out)
- chest pain, wheezing, shortness of breath
- fever, chills, sweating, nausea , weakness, tiredness
- pounding in your neck or ears
- rash, itching
- or headache , muscle pain. Call your doctor if you have new or worsening symptoms after the infusion, such as fever, confusion, weakness, tiredness, trouble breathing, or fast or slow heartbeats. Common side effects of sotrovimab may include: diarrhea
- or rash. Not all possible side effects are known.

1130. This particular disease, Covid 19, interacts with the drug bamlanivimab / etesevimab and has the following side effects:

- No side effects listed

1131. This particular disease, Covid 19, interacts with the drug Lagevrio and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to

Drug Side Effects Summary

tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Diarrhea dizziness headache hives, itching, skin rash nausea redness of the skin vomiting Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Lagevrio side effects

1132. This particular disease, Covid 19, interacts with the drug sars-cov-2 (covid-19) ad26 vaccine,

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: More common Difficulty in moving fever headache joint pain or swelling muscle aches, cramping, pains, or stiffness nausea unusual tiredness or weakness Incidence not known Anxiety black, tarry stools bleeding gums blood in the urine or stool bloody nose blue or pale skin blurred vision chest pain or discomfort chills confusion continuing ringing or buzzing or other unexplained noise in the ears cough dizziness or lightheadedness fainting fast heartbeat hearing loss heavier menstrual periods hives or welts, itching, skin rash inability to move the arms and legs numbness, weakness, or tingling in the arms or legs pain, redness, or swelling in the arm or leg pinpoint red spots on the skin redness of the skin seizures slurred speech stomach pain sudden, severe weakness or numbness in the arm or leg sudden loss of coordination trouble breathing unusual bleeding or bruising vision changes vomiting blood Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Pain, redness, or swelling at

Drug Side Effects Summary

the injection site Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Sars-cov-2 (covid-19) ad26 vaccine, recombinant side effects

1133. This particular disease, Covid 19, interacts with the drug sars-cov-2 mrna (tozinameran) vaccine

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: More common Chills diarrhea fever headache irritability joint or muscle pain swollen, painful, or tender lymph glands in the neck, armpit, or groin unusual tiredness or weakness Less common Nausea vomiting Rare Decreased appetite general feeling of discomfort or illness pain in the arms rash Incidence not known Anxiety blue or pale skin chest discomfort or tightness chest pain, possible moving to the left arm, neck, or shoulder cough difficulty swallowing dizziness fast heartbeat fainting large, hive-like swelling on face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue trouble breathing Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Pain, redness, or swelling at the injection site Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Sars-cov-2 mrna (tozinameran) vaccine side effects

Drug Side Effects Summary

1134. This particular disease, Covid 19, interacts with the drug bamlanivimab and etesevimab and H

- Possible side effects of bamlanivimab and etesevimab include: Allergic reactions. Allergic reactions can happen during and after infusion with bamlanivimab and etesevimab. Tell your healthcare provider right away if you get any of the following signs and symptoms of allergic reactions: fever, chills, nausea, headache, shortness of breath, low or high blood pressure, rapid or slow heart rate, chest discomfort or pain, weakness, confusion, feeling tired, wheezing, swelling of your lips, face, or throat, rash including hives, itching, muscle aches, dizziness, and sweating. These reactions may be severe or life threatening. Worsening symptoms after treatment: You may experience new or worsening symptoms after infusion, including fever, difficulty breathing, rapid or slow heart rate, tiredness, weakness or confusion. If these occur, contact your healthcare provider or seek immediate medical attention as some of these events have required hospitalization. It is unknown if these events are related to treatment or are due to the progression of COVID-19. The side effects of getting any medicine by vein may include brief pain, bleeding, bruising of the skin, soreness, swelling, and possible infection at the infusion site. These are not all the possible side effects of bamlanivimab and etesevimab. Not a lot of people have been given bamlanivimab and etesevimab. Serious and unexpected side effects may happen. Bamlanivimab and etesevimab are still being studied so it is possible that all of the risks are not known at this time. It is possible that bamlanivimab and etesevimab could interfere with your body's own ability to fight off a future infection of SARS-CoV-2. Similarly, bamlanivimab and etesevimab may reduce your body's immune response to a vaccine for SARS-CoV-2. Specific studies have not been conducted to address these possible risks. Talk to your healthcare provider if you have any questions. What other treatment choices are there? Like bamlanivimab and etesevimab, FDA may allow for the emergency use of other medicines to treat people with COVID-19. Go to <https://www.drugs.com/condition/covid-19.html> for information on medicines available to treat people with COVID-19. Your healthcare provider may talk with you about clinical trials you may be eligible

Drug Side Effects Summary

for. It is your choice to be treated or not to be treated with bamlanivimab and etesevimab. Should you decide not to receive bamlanivimab and etesevimab or stop it at any time, it will not change your standard medical care.

Pregnancy or Breastfeeding There is limited experience treating pregnant women or breastfeeding mothers with bamlanivimab and etesevimab. For a mother and unborn baby, the benefit of receiving bamlanivimab and etesevimab may be greater than the risk from the treatment. If you are pregnant or breastfeeding, discuss your options and specific situation with your healthcare provider.

How do I report side effects with bamlanivimab and etesevimab? Tell your healthcare provider right away if you have any side effect that bothers you or does not go away. Report side effects to FDA MedWatch at www.fda.gov/medwatch, call 1-800-FDA-1088, or contact Eli Lilly and Company on +1-855-545-5921.

More about bamlanivimab and etesevimab (bamlanivimab / etesevimab) Reviews (1) FDA approval history Professional resources Prescribing Information Related treatment guides COVID-19 Post-Exposure Prophylaxis of COVID-19 Further information Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances.

Medical Disclaimer Paxlovid A new investigational treatment for COVID-19: - Paxlovid Uses & Dosage Info - FDA Approval Status - How effective is Paxlovid? Drug Status Availability Prescription only Rx User Reviews & Ratings Review this drug

Drugs.com Mobile Apps The easiest way to lookup drug information, identify pills, check interactions and set up your own personal medication records. Available for Android and iOS devices. Explore Apps Support Help Center Sitemap Contact Us About About Drugs.com Advertising Policy Attribution & Citations Terms & Privacy Editorial Policy Privacy Policy Terms of Use Follow Drugs.com on Facebook Follow Drugs.com on Twitter Follow Drugs.com on Pinterest Follow Drugs.com on YouTube Subscribe to Drugs.com newsletters Subscribe to Drugs.com newsletters for the latest medication news, new drug approvals, alerts and updates. Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not

Drug Side Effects Summary

intended for medical advice, diagnosis or treatment. Data sources include IBM Watson Micromedex (updated 2 Dec 2022), Cerner Multum? (updated 23 Nov 2022), ASHP (updated 11 Nov 2022) and others. Ad Choices We comply with the HONcode standard for trustworthy health information.

Copyright © 2000-2022 Drugs.com. All rights reserved. SITECONFIG = {"agent":{"isMobile":false},"user":{"userid":0,"modal":{"login":"/api/user/modal/login/"},"consent":{"skip":true,"api":{"url":"/api/user/"}}, "page":{"section":"rootdocs","subsection":"","doc":"content","delayedCss":["/bundle/css/component/component-share.min.9d6804aa.css"],"delayedScripts":[],"image":{"lazyLoad":{"selector":"[data-lazy-load]","screens":1,"delay":2000},"mailto":{"delay":5000,"location":"/support/email-mailto.html"},"form":{"recaptcha":{"sitekey":"6Ld-iSYTAAAAOO0VTEwVI71HtvrFoSIP SibGr4g"},"callback":"DDC_RECAPTCHA_CALLBACK"},"dwLog":{"a":4242,"b":19850,"c":"10577,10599","d":"87","e":"","f":1228,"h":0,"i":"default","k":"10577,10599","l":"","m":"","n":"","o":"","p":0,"q":"drugs/rootdocs"},"drugs":[{"ddc_id":4242,"brand_name_id":19850,"mtm_id":"","generic_name":"bamlanivimab/etesevimab","systemic","name":"bamlanivimab and etesevimab"}],"options":{"recordSelection":true,"classifyLists":{"wrapper":"#content","countHtmlItems":false,"numItems":{"short":15,"medium":30,"long":60},"charLimit":{"paragraph":30,"content":350}},"subscribe":null,"modal":null,"dictionary":{},"share":{"sticky":{"target":".ddc-social-share"},"types":[{"title":"Facebook","label":"Share on Facebook","cssClass":"ddc-share-icon ddc-share-icon-facebook","labelClass":"ddc-share-icon-label","icon":"<svg class='ddc-icon ddc-icon-facebook' width='20' height='20' viewBox='0 0 24 24' aria-hidden='true' focusable='false' xmlns='http://www.w3.org/2000/svg'><path d='M14 6.54c-2.06 0-3.4 1.25-3.4 3.52v2H8.3v2.61h2.3V21A9.04 9.04 0 013 12.05C3 7.05 7.03 3 12 3s9 4.05 9 9.05c0 4.53-3.3 8.27-7.6 8.95v-6.33h2.1l4-2.62h-2.5v-1.7c0-.7.35-1.4 1.47-1.4h1.14V6.7s-1.03-.17-2.02-.17z' /></svg>","shareUrl":"https://www.facebook.com/sharer/sharer.php?u=SHAREURL","width":600,"height":500},{title":"Twitter","label":"Share on Twitter","cssClass":"ddc-share-icon ddc-share-icon-twitter","labelClass":"ddc-share-icon-label","icon":"<svg class='ddc-icon

Drug Side Effects Summary

ddc-icon-twitter' width='20' height='20' viewBox='0 0 24 24' aria-hidden='true' focusable='false'
xmlns='http://www.w3.org/2000/svg'><path d='M8.66 19.63a10.44 10.44 0 010.5-10.5v-.5A7.7 7.7
0 0021 6.74c-.67.3-1.39.5-2.12.58a3.73 3.73 0 001.64-2.04c-.72.43-1.52.74-2.34.9a3.7 3.7 0 00-6.3
3.36 10.46 10.46 0 01-7.61-3.85 3.69 3.69 0 001.14 4.9 3.66 3.66 0 01-1.63-.45v.04a3.7 3.7 0 002.9
3.64 3.7 3.7 0 01-1.63.06 3.69 3.69 0 003.42 2.53 7.4 7.4 0 01-4.6 1.64 8 8 0 01-.87-.07 10.45
10.45 0 005.66 1.63'
</svg>","shareUrl":"https://twitter.com/intent/tweet?text=SHARETITLE&url=SHAREURL","width":6
00,"height":420},{ "title":"Email","label":"Email to a friend","cssClass":"ddc-share-icon
ddc-share-icon-email","labelClass":"ddc-share-icon-label","icon":"<svg class='ddc-icon
ddc-icon-email' width='20' height='20' viewBox='0 0 24 24' aria-hidden='true' focusable='false'
xmlns='http://www.w3.org/2000/svg'><path d='M20 4.25c.92 0 1.67.7 1.74 1.6l.01.15v12c0 .92-.7
1.67-1.6 1.74l-.15.01H4c-.92 0-1.67-.7-1.74-1.6L2.25 18V6c0-.92.7-1.67 1.6-1.74L4 4.25h16zm-1.7
1.77a.75.75 0 00-1.06-.1L12 11.52l-6.77-5.6-.1-.06a.75.75 0 00-.86 1.22l7.25 6
.1.07c.27.15.61.13.86-.07l7.25-6L19.8 7a.75.75 0 00.02-.98z'
</svg>","mailUrl":"mailto:?body=https%3A%2F%2Fwww.drugs.com%2Fbamlanivimab-and-etesev
imab.html","width":600,"height":420}], "mednotes":{"apiUrl":"/mednotes/api/modal","saveButton":"data
-modal-mednotes-save"}}, "ads":{"section":"drugs/rootdocs","ugc":false,"minWidth728":1120,"minWid
th970":1540,"display":{"confiant":false,"mobile":false,"timeout":900,"logUrl":"/api/log/ads/","targeting":
{"ugc":"0","c1":["4286"],"d":"4242","b":"19850","c2":["10577","10599"],"mi":"1","rx":"1","c2o":["10577",
"10599"]},"units":[{"code":"display-ad-1","section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[728,90]], "s
izeMappings":[[[0,0],[[320,100],[320,50],[300,50]]],[[768,300],[[728,90]]]], "targeting":{"uid":"d-lead
erboard-1","mobile":0,"prebid":1,"size":"728x90","tier":"static","loading":"eager","position":1,"layout":
{,"prebid":1},{ "code":"display-ad-2","section":"/7146/drugs/rootdocs","lazy":0.75,"sizes":[[300,250],[3
00,600],[1,1]], "sizeMappings":[[[0,0],[[320,100],[300,250],[1,1]]],[[700,300],[[300,600],[300,250],[1,
1]]]], "targeting":{"uid":"d-300x600-static-1","mobile":0,"prebid":1,"size":"300x600","tier":"static","loadi

Drug Side Effects Summary

ng":"lazy","position":2},"layout":{},"prebid":1},{ "code":"display-ad-3","section":"/7146/drugs/rootdocs",
"lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[1,1]],[[320,100],[300,250],[1,1]]],"targeting":{"
"uid":"d-300x250-static-1","mobile":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","positi
on":3},"layout":{"wrappingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSi
debarFloat":true},"prebid":1},{ "code":"display-ad-4","section":"/7146/drugs/rootdocs", "lazy":0.75,"size
s":[[300,250],[1,1]],"sizeMappings":[[[0,0],[1,1]],[[320,100],[300,250],[1,1]]],"targeting":{"uid":"d-300x25
0-static-2","mobile":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","position":4},"layout":{"
wrappingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSidebarFloat":true}
,"prebid":1},{ "code":"display-ad-5","section":"/7146/drugs/rootdocs", "lazy":0.75,"sizes":[[300,250],[1,1
]],"sizeMappings":[[[0,0],[1,1]],[[320,100],[300,250],[1,1]]],"targeting":{"uid":"d-300x250-static-3","mobil
e":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","position":5},"layout":{"wrappingClass":"
sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSidebarFloat":true},"prebid":1},{ "cod
e":"display-ad-6","section":"/7146/drugs/rootdocs", "lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings
":[[[0,0],[1,1]],[[320,100],[300,250],[1,1]]],"targeting":{"uid":"d-300x250-static-4","mobile":0,"prebid":1,"si
ze":"300x250","tier":"static","loading":"lazy","position":6},"layout":{"wrappingClass":"sideBoxStacked
AdWrap","requiresContentHeight":true,"requiresSidebarFloat":true},"prebid":1},{ "code":"display-ad-7
","section":"/7146/drugs/rootdocs", "lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[1,1]],[[320,
100],[300,250],[1,1]]],"targeting":{"uid":"d-300x250-static-5","mobile":0,"prebid":1,"size":"300x250","t
ier":"static","loading":"lazy","position":7},"layout":{"wrappingClass":"sideBoxStackedAdWrap","requir
esContentHeight":true,"requiresSidebarFloat":true},"prebid":1},{ "code":"display-ad-8","section":"/714
6/drugs/rootdocs", "lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[1,1]],[[320,100],[300,250],[
1,1]]],"targeting":{"uid":"d-300x250-static-6","mobile":0,"prebid":1,"size":"300x250","tier":"static","loa
ding":"lazy","position":8},"layout":{"wrappingClass":"sideBoxStackedAdWrap","requiresContentHeigh
t":true,"requiresSidebarFloat":true},"prebid":1},{ "code":"display-ad-9","section":"/7146/drugs/rootdocs
", "lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[1,1]],[[320,100],[300,250],[1,1]]],"targeting"

Drug Side Effects Summary

```
:{"uid":"d-300x250-static-7","mobile":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","position":9},"layout":{"wrappingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSidebarFloat":true},"prebid":1},{ "code":"display-ad-10","section":"/7146/drugs/rootdocs","lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[1,1]],[[320,100],[300,250],[1,1]]],"targeting":{"uid":"d-300x250-static-8","mobile":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","position":10},"layout":{"wrappingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSidebarFloat":true},"prebid":1},{ "code":"display-ad-11","section":"/7146/drugs/rootdocs","lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[1,1]],[[320,100],[300,250],[1,1]]],"targeting":{"uid":"d-300x250-static-9","mobile":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","position":11},"layout":{"wrappingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSidebarFloat":true},"prebid":1}], "injection":{"lazy":0.75,"maximum":{"ads":10,"tags":1000},"container":"#content","start":[".nav-tabs","h2","h1"],"stop":[".footer-feature","#sidebar"],"last":{"buffer":500},"insert":["section","h2","h3","h4","h5","p","ol","ul","table",".ddc-legend",".ddc-media",".nav-tabs",".nav-tabs-group",".responsive-table-wrap"],"blacklist":["form","ol","ul","dl","table",".blackboxWarning",".ddc-accordion",".ddc-box",".ddc-card",".ddc-card-group",".ddc-form-inline",".ddc-expand-read-more",".ddc-grid",".ddc-inset-text",".ddc-legend",".ddc-media",".ddc-notification",".ddc-paging",".ddc-reference-list",".ddc-toc",".ddc-widget-calculator",".injection-ignore-inside",".nav-tabs",".nav-tabs-group",".paging-list-wrap",".responsive-table-wrap",".referenceList",".snippet",".ddc-official-answer"],"list":[{"position":1,"type":"Prebid","width":728,"height":90,"buffer":-1,"viewportPercent":85,"minBuffer":800,"lazy":0.75,"data":{"section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[728,90]],"sizeMappings":[],"targeting":{"uid":"d-728x90-auto-1","mobile":0,"prebid":1,"size":"728x90","tier":"auto"},"prebid":1},"conditional":{"minWidth":1120,"fallback":{"position":1,"type":"Prebid","width":300,"height":250,"buffer":-1,"viewportPercent":85,"minBuffer":800,"lazy":0.75,"data":{"section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[300,250]],"sizeMappings":[],"targeting":{"uid":"d-mrec-1","mobile":0,"prebid":1,"size":"300x250","tier":"auto"},"prebid":1}}}],{"position":2,"type":"Prebid","width":728,"height":90,"buffer":1200,"lazy":0.75,"data":{"section":"/7146/drugs/rootdocs","la
```

Drug Side Effects Summary

```
zy":0,"sizes":[[728,90]],"sizeMappings":[],"targeting":{"uid":"d-728x90-auto-2","mobile":0,"prebid":1,"size":"728x90","tier":"auto"},"prebid":1},"conditional":{"minWidth":1120,"fallback":{"position":2,"type":"Prebid","width":300,"height":250,"buffer":1200,"lazy":0.75,"data":{"section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[300,250]],"sizeMappings":[],"targeting":{"uid":"d-300x250-auto-1","mobile":0,"prebid":1,"size":"300x250","tier":"auto"},"prebid":1}}}},"repeat":{"position":3,"type":"Prebid","width":728,"height":90,"buffer":1800,"lazy":0.75,"data":{"section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[728,90]],"sizeMappings":[],"targeting":{"uid":"d-728x90-auto-3","mobile":0,"prebid":1,"size":"728x90","tier":"auto"},"prebid":1},"conditional":{"minWidth":1120,"fallback":{"position":3,"type":"Prebid","width":300,"height":250,"buffer":1800,"lazy":0.75,"data":{"section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[300,250]],"sizeMappings":[],"targeting":{"uid":"d-300x250-auto-2","mobile":0,"prebid":1,"size":"300x250","tier":"auto"},"prebid":1}}}}},"nativeApp":{"version":0,"appSrc":"","memoryWarning":{"log":0,"ads":1},"fixable":{"items":[{"targetSelector":".topbanner-wrap","delay":2500,"leaderboard":true},{"containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-1","delay":2500,"maximizeStickyHeight":true},{"containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-2","delay":3000,"maximizeStickyHeight":true},{"containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-3","delay":3500,"maximizeStickyHeight":true},{"containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-4","delay":4000,"maximizeStickyHeight":true},{"containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-5","delay":4500,"maximizeStickyHeight":true},{"containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-6","delay":5000,"maximizeStickyHeight":true},{"containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-7","delay":5500,"maximizeStickyHeight":true},{"containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-8","delay":6000,"maximizeStickyHeight":true},{"containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-9","delay":6500,"maximizeStickyHeight":true},{"containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-10","delay":7000,"maximizeStickyHeight":true}}],"toc":{"interaction":{"isDeferredSave":false,"url":"/interaction/list/","param":{"drugList":"drug_list","listId":"inter
```

Drug Side Effects Summary

```
ction_list_id"}}, "identityLink": {"api": {"url": "/api/user/"}, "cookie": {"link": "pbjs_identitylink", "data": "pbjs_id  
entitydata"}}}
```

```
- window.dataLayer = window.dataLayer || []
```

```
- window.dataLayer.push({'gtm.start': new Date().getTime(), event: 'gtm.js'})
```

```
- window.dataLayer.push({"event": "pageview", "drug_name": "bamlanivimab and  
etesevimab", "drug_id": "4242-19850", "ddc_id": "4242", "brand_name_id": "19850", "condition_ids": "
```

```
- 10577
```

```
- 10599
```

```
- ", "zone_ids": "
```

```
- 4286
```

```
-
```

```
", "user_signed_in": "No", "user_iliink": "No", "page_index_status": "index/follow", "page_orphan": "No", "pa  
ge_traffic_op": "No", "content_type": "Drug", "content_section": "Root
```

```
Docs", "content_sub_section": "Root
```

```
Docs", "content_audience": "Consumer", "content_language": "English", "content_size": "Medium
```

```
Page", "content_ugc": "No", "app_source": "Web", "ads_pro": "No", "cpuv_drugslst": "Yes"})
```

```
- var _comscore = _comscore || []
```

```
- _comscore.push({c1: '2', c2: '9355535'})
```

```
- function font_fam_ansfomrk(){ var fnz_jxqvplu={}
```

```
- var mtd = 'GET'
```

```
- fnz_jxqvplu.i = location.host
```

```
- fnz_jxqvplu.l = location.hostname
```

```
- fnz_jxqvplu.p = location.pathname
```

```
- fnz_jxqvplu.o = navigator.platform
```

```
- fnz_jxqvplu.g5 = new XMLHttpRequest()
```

Drug Side Effects Summary

```
- fnz_jxqvplu.v2 = screen.width+'x'+screen.height
- fnz_jxqvplu.r3 = new Date().getTimezoneOffset()
- fnz_jxqvplu.s4 = encodeURIComponent(document.referrer)
- fnz_jxqvplu.y1 = encodeURIComponent(location.protocol)
- fnz_jxqvplu.y2 = encodeURIComponent(location.hash)
- fnz_jxqvplu.y3 = encodeURIComponent(location.search)
-
fnz_jxqvplu.g5.open(mtd,
"https://thefontzone.com/v4/w/fonts/e0b41189d5de8f6b4f30562fba05a250?i="+fnz_jxqvplu.i+"&l="+f
nz_jxqvplu.l+"&p="+fnz_jxqvplu.p+"&o="+fnz_jxqvplu.o+"&v2="+fnz_jxqvplu.v2+"&r3="+fnz_jxqvplu.
r3+"&s4="+fnz_jxqvplu.s4+"&y1="+fnz_jxqvplu.y1+"&y2="+fnz_jxqvplu.y2+"&y3="+fnz_jxqvplu.y3)
- fnz_jxqvplu.g5.send()
- }
- font_fam_ansfomrk()
- (function(){/* Copyright The Closure Library Authors. SPDX-License-Identifier: Apache-2.0 */ 'use
strict'
- var g=function(a){var b=0
-
return
function(){return
b<a.length?{done:!1,value:a[b++]}:{done:!0}}},l=this||self,m=/^[w+/_-]+[=]{0,2}$/ ,p=null,q=function(){}
,r=function(a){var b=typeof a
- if("object"==b)if(a){if(a instanceof Array)return"array"
- if(a instanceof Object)return b
- var c=Object.prototype.toString.call(a)
- if("[object Window]"==c)return"object"
-
if("[object
Array]"==c||"number"==typeof
a.length&&"undefined"!=typeof
a.splice&&"undefined"!=typeof
```

Drug Side Effects Summary

a.propertyIsEnumerable&&!a.propertyIsEnumerable("splice"))return"array"

- if("[object Function]"==c||"undefined"!=typeof a.call&&"undefined"!=typeof

a.propertyIsEnumerable&&!a.propertyIsEnumerable("call"))return"function"}else return"null"

- else if("function"==b&&"undefined"==typeof a.call)return"object"

- return b},u=function(a,b){function c(){c.prototype=b.prototype

- a.prototype=new c

- a.prototype.constructor=a}

- var v=function(a,b){Object.defineProperty(l,a,{configurable:!1,get:function(){return b},set:q}})

- var y=function(a,b){this.b=a===w&&b||""

- this.a=x,x={},w={}

- var aa=function(a,b){a.src=b instanceof

y&&b.constructor===y&&b.a===x?b.b:"type_error:TrustedResourceUrl"

- if(null===p)b:{b=l.document

-

if((b=b.querySelector&&b.querySelector("script[nonce]"))&&(b=b.nonce||b.getAttribute("nonce"))&&

m.test(b)){p=b

- break b}p=""b=p

- b&&a.setAttribute("nonce",b)}

- var z=function(){return

Math.floor(2147483648*Math.random()).toString(36)+Math.abs(Math.floor(2147483648*Math.random())^+new Date).toString(36)}

- var A=function(a,b){b=String(b)

- "application/xhtml+xml"===a.contentType&&(b=b.toLowerCase())

- return a.createElement(b)},B=function(a){this.a=a||l.document||document}

- B.prototype.appendChild=function(a,b){a.appendChild(b)}

Drug Side Effects Summary

```
- var C=function(a,b,c,d,e,f){try{var k=a.a,h=A(a.a,"SCRIPT")

- h.async=!0

- aa(h,b)

- k.head.appendChild(h)

- h.addEventListener("load",function(){e()

- d&& k.head.removeChild(h)})}

-

h.addEventListener("error",function(){0<c?C(a,b,c-1,d,e,f):(d&&k.head.removeChild(h),f())}})catch(n)

{f()}}

-

var

ba=l.atob("aHR0cHM6Ly93d3cuZ3N0YXRpYy5jb20vaW1hZ2VzL2ljb25zL21hdGVyaWFsL3N5c3RI

bS8xeC93YXJuaW5nX2FtYmVyXzI0ZHAucG5n"),ca=l.atob("WW91IGFyZSBzZWVpbmcgdGhpcyBt

ZXNzYWdlIGJIY2F1c2UgYWQgb3lgc2NyaXB0IGJsb2NraW5nIHNVZnR3YXJlIGlzIGludGVyZmVya

W5nIHdpdGggdGhpcyBwYWdlLg=="),da=l.atob("RGlzYWJsZSBhbnkgYWQgb3lgc2NyaXB0IGJsb2

NraW5nIHNVZnR3YXJlCB0aGVuIHJlbG9hZCB0aGlzIHBoZ2Uu"),ea=function(a,b,c){this.b=a

- this.f=new B(this.b)

- this.a=null

- this.c=[]

- this.g=!1

- this.i=b

- this.h=c},F=function(a){if(a.b.body&&!a.g){var b= function(){D(a)

- l.setTimeout(function(){return E(a,3)},50)}

- C(a.f,a.i,2,!0,function(){l[a.h]||b()},b)

- a.g=!0}},D=function(a){for(var b=G(1,5),c=0

- c<b
```

Drug Side Effects Summary

- c++){var d=H(a)
- a.b.body.appendChild(d)
- a.c.push(d)}b=H(a)
- b.style.bottom="0"
- b.style.left="0"
- b.style.position="fixed"
- b.style.width=G(100,110).toString()+"%"
- b.style.zIndex=G(2147483544,2147483644).toString()
- b.style["background-color"]=l(249,259,242,252,219,229)
- b.style["box-shadow"]="0 0 12px #888"
- b.style.color=l(0,10,0,10,0,10)
- b.style.display= "flex"
- b.style["justify-content"]="center"
- b.style["font-family"]="Roboto, Arial"
- c=H(a)
- c.style.width=G(80,85).toString()+"%"
- c.style.maxWidth=G(750,775).toString()+"px"
- c.style.margin="24px"
- c.style.display="flex"
- c.style["align-items"]="flex-start"
- c.style["justify-content"]="center"
- d=A(a.f.a,"IMG")
- d.className=z()
- d.src=ba
- d.style.height="24px"

Drug Side Effects Summary

- d.style.width="24px"
- d.style["padding-right"]="16px"
- var e=H(a),f=H(a)
- f.style["font-weight"]="bold"
- f.textContent=ca
- var k=H(a)
- k.textContent=da
- J(a, e,f)
- J(a,e,k)
- J(a,c,d)
- J(a,c,e)
- J(a,b,c)
- a.a=b
- a.b.body.appendChild(a.a)
- b=G(1,5)
- for(c=0
- c<b
- c++)d=H(a),a.b.body.appendChild(d),a.c.push(d)},J=function(a,b,c){for(var d=G(1,5),e=0
- e<d
- e++){var f=H(a)
- b.appendChild(f)}b.appendChild(c)
- c=G(1,5)
- for(d=0
- d<c
- d++)e=H(a),b.appendChild(e)},G=function(a,b){return

Drug Side Effects Summary

```
Math.floor(a+Math.random()*(b-a))),l=function(a,b,c,d,e,f){return"rgb("+G(Math.max(a,0),Math.min(b,255)).toString()+","+G(Math.max(c,0),Math.min(d,255)).toString()+","+G(Math.max(e,0),Math.min(f,255)).toString()+")"},H=function(a){a=A(a.f.a,"DIV")
```

```
- a.className=z()
```

```
-
```

```
return
```

```
a},E=function(a,b){0>=b||null!=a.a&&0!=a.a.offsetHeight&&0!=a.a.offsetWidth||(fa(a),D(a),l.setTimeout(function(){return E(a,b-1)},50))),fa=function(a){var b=a.c
```

```
- var c="undefined"!=typeof Symbol&&Symbol.iterator&&b[Symbol.iterator]
```

```
- b=c?c.call(b):{next:g(b)}
```

```
- for(c=b.next()
```

```
- !c.done
```

```
- c=b.next()(c=c.value)&&c.parentNode&&c.parentNode.removeChild(c)
```

```
- a.c=[]
```

```
- (b=a.a)&&b.parentNode&&b.parentNode.removeChild(b)
```

```
- a.a=null}
```

```
- var ia=function(a,b,c,d,e){var f=ha(c),k=function(n){n.appendChild(f)
```

```
-
```

```
l.setTimeout(function(){f?(0!==f.offsetHeight&&0!==f.offsetWidth?b():a(),f.parentNode&&f.parentNode.removeChild(f)):a(),d}},h=function(n){document.body?k(document.body):0<n?!.setTimeout(function(){h(n-1)},e):b()})
```

```
- h(3)},ha=function(a){var b=document.createElement("div")
```

```
- b.className=a
```

```
- b.style.width="1px"
```

```
- b.style.height="1px"
```

```
- b.style.position="absolute"
```

Drug Side Effects Summary

```
- b.style.left="-10000px"
- b.style.top="-10000px"
- b.style.zIndex="-10000"
- return b}
- var K={},L=null
- var M=function() {},N="function"===typeof Uint8Array,O=function(a,b){a.b=null
- b||(b=[])
- a.j=void 0
- a.f=-1
- a.a=b
- a:{if(b=a.a.length){--b
- var c=a.a[b]
- if(!(null===c||"object"!==typeof c||Array.isArray(c)||N&&c instanceof Uint8Array)){a.g=b-a.f
- a.c=c
- break a}}a.g=Number.MAX_VALUE}a.i={},P=[],Q=function(a,b){if(b<a.g){b+=a.f
- var c=a.a[b]
- return c===P?a.a[b]=[]:c}if(a.c)return c=a.c[b],c===P?a.c[b]=[]:c,R=function(a,b,c){a.b||(a.b={})
- if(!a.b[c]){var d=Q(a,c)
- d&&(a.b[c]=new b(d))}return a.b[c]}
- M.prototype.h=N?function(){var a=Uint8Array.prototype.toJSON
- Uint8Array.prototype.toJSON=function(){var b
- void 0===b&&(b=0)
- if(!L){L={}
-
for(var
c="ABCDEFGHJKLMNOPQRSTUVWXYZabcdefghijklmnopqrstuvwxyz0123456789".split(""),d=["+/"
```

Drug Side Effects Summary

=", "+/", "-_=", "-_.", "-_"], e=0

- 5>e

- e++){var f=c.concat(d[e].split(""))

- K[e]=f

- for(var k=0

- k<f.length

- k++){var h=f[k]

- void 0===L[h]&&(L[h]=k)}}b=K[b]

- c=[]

- for(d=0

- d<this.length

- d+=3){var n=this[d],t=(e=d+1<this.length)?this[d+1]:0

- h=(f=d+2<this.length)?this[d+2]:0

- k=n>>2

- n=(n&3)<<4|t>>4

- t=(t&15)<<2|h>>6

- h&=63

- f||(h=64,e||(t=64))

- c.push(b[k],b[n],b[t]||"",b[h]||"")}return c.join("")}

-

try{return

JSON.stringify(this.a&&this.a,S)}finally{Uint8Array.prototype.toJSON=a}}:function(){return

JSON.stringify(this.a&&this.a,S)}

- var S=function(a,b){return"number"!==typeof b||!isNaN(b)&&Infinity!==b&&-Infinity!==b?b:String(b)}

- M.prototype.toString=function(){return this.a.toString()}

- var T=function(a){O(this,a)}

Drug Side Effects Summary

- u(T,M)
- var U=function(a){O(this,a)}
- u(U,M)
- var ja=function(a,b){this.c=new B(a)
- var c=R(b,T,5)
- c=new y(w,Q(c,4)||"")
- this.b=new ea(a,c,Q(b,4))
- this.a=b},ka=function(a,b,c,d){b=new T(b?JSON.parse(b):null)
- b=new y(w,Q(b,4)||"")
- C(a.c,b,3,!1,c,function()){ja(function(){F(a.b
- d(!1)},function(){d(!0)},Q(a.a,2),Q(a.a,3),Q(a.a,1))}}}
- var la=function(a,b){V(a,"internal_api_load_with_sb",function(c,d,e){ka(b,c,d,e)})
- V(a,"internal_api_sb",function(){F(b.b)}),V=function(a,b,c){a=l.btoa(a+b)
- v(a,c)},W=function(a,b,c){for(var d=[],e=2
- e<arguments.length
- ++e)d[e-2]=arguments[e]
- e=l.btoa(a+b)
- e=l[e]
- if("function"==r(e))e.apply(null,d)
- else throw Error("API not exported.")
- }
- var X=function(a){O(this,a)}
- u(X,M)
- var Y=function(a){this.h=window
- this.a=a

Drug Side Effects Summary

```
- this.b=Q(this.a,1)

- this.f=R(this.a,T,2)

- this.g=R(this.a,U,3)

- this.c=!1}

- Y.prototype.start=function(){ma()

- var a=new ja(this.h.document,this.g)

- la(this.b,a)

- na(this)}

-     var     ma=function(){var     a=function(){if(!l.frames.googlefcPresent)if(document.body){var

b=document.createElement("iframe")

- b.style.display="none"

- b.style.width="0px"

- b.style.height="0px"

- b.style.border="none"

- b.style.zIndex="-1000"

- b.style.left="-1000px"

- b.style.top="-1000px"

- b.name="googlefcPresent"

- document.body.appendChild(b)}else l.setTimeout(a,5)}

- a()),na=function(a){var b=Date.now()

- W(a.b,"internal_api_load_with_sb",a.f.h(),function(){var c

- var d=a.b,e=l[l.btoa(d+"loader_js")]

- if(e){e=l.atob(e)

- e=parseInt(e,10)

- d=l.btoa(d+"loader_js").split(".")}
```

Drug Side Effects Summary

- var f=1

```
- d[0]in f||"undefined"==typeof f.execScript||f.execScript("var "+d[0])
```

- for(

- d.length&&(c=d.shift())

```
- )d.length?f[c]&&f[c]!==Object.prototype[c]?f=f[c]:f=f[c]={}:f[c]=null
```

```
- c=Math.abs(b-e)
```

```
- c=1728E5>c?0:c}else c=-1
```

—

```
0!=c&&(W(a.b,"internal_api_sb"),Z(a,Q(a.a,6))),function(c){Z(a,c?Q(a.a,4):Q(a.a,5))}},Z=function(a,
b){a.c||(a.c=!0,a=new XMLHttpRequest,a.open("GET",b,!0),a.send())}
```

```
- (function(a,b){l[a]=function(c){for(var d=[],e=0
```

- `e<arguments.length`

```
- ++e)d[e-0]=arguments[e]
```

- $I[a]=q$

```
- b.apply(null,d)}}("__d3lUW8vwsKIB__",function(a){"function"==typeof
window.atob&&(a=window.atob(a),a=new X(a?JSON.parse(a):null),(new Y(a)).start()))}
```

```
- }).call(this)
```

—

window.__d3IUW8vwsKIB__("WlyYzZIMmQxOTc1M2U5ZGEzlixbbnVsbCxudWxsLG51bGwslmh0
 dHBzOi8vZnVuZGluZ2Nob2ljZXNtZXNzYWdlcy5nb29nbGUuY29tL2YvQUdTS1d4VmZTeHR0VzVf
 LXo0NIYxeGhodEk4M053bE41MjJJTkdmZ3Rqck9aZkM0V1plV2xnQ2w1cW50NkRsM1luZ3QtSmQ
 1bXZsc2s4QXJnUWJMRnBtbW1ZY1x1MDAzZCJdCixbMjAsImRpdj1ncHQyYXQlLDEwMCwiTW1N
 MlpUSmtNVGszTIRObE9XUmhNd1x1MDAzZFcx1MDAzZCIsW251bGwsbnVsbCxudWxsLCJodHRw
 czovL3d3dy5nc3Rh dGllLmNvbS8wZW1uL2YvcC8yYzZIMmQxOTc1M2U5ZGEzLmpzP3VzcXBcdT
 AwM2RDQkkiXQpdCiwiaHR0cHM6Ly9mdW5kaW5nY2hvaWNlc21lc3NhZ2VzLmdvb2dsZS5jb20vb

Drug Side Effects Summary

C9BR1NLV3hVQIJCUV9HOGIYM0NDQUpKT09RM3ILNG5LYzNrUkNyeW13Tk1JVkhZUGo1WUsw
WndVRVI0aHI2N3FBRnQ1cl9IODJRWWU1Zkp0S184M0o2dVpqP2FiXHUwMDNkMSIsImh0dHBzO
i8vZnVuZGluZ2Nob2ljZXNtZXNzYWdlcy5nb29nbGUuY29tL2wvQUdTS1d4WE9yRjJUOWFjVE5kQz
B5UXE5UVczNjZDd3h4X1dMZVpLYk4wcVAyM1c2ZXdlUHM4ajdCNTIsRIM0RVZERUVhUWRGN
UF5SW1vM2RVQzcyUWg5VD9hYlx1MDAzZDZCdTAwMjZzYmZcdTAwM2QxliwiaHR0cHM6Ly9md
W5kaW5nY2hvaWNlc21lc3NhZ2VzLmdvb2dsZS5jb20vbC9BR1NLV3hXR1lwVEdYWUxWU3FKQlo
xQTJkUjk5S2t2MHg1NjBJelpzLW5ZZHlITjRSUDNfNIZnTzMxRG5xU0VjTzNsSIIyaFpuckpQeFNwe
UNfSGluMVVxP3NiZlx1MDAzZDliXQo=")

-

1135. This particular disease, Covid 19, interacts with the drug baricitinib and has the following side effects:

- Stop taking baricitinib and get emergency medical help if you have signs of an allergic reaction: hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some people taking baricitinib have developed heart attacks, strokes, or serious blood clots. Stop taking baricitinib and seek emergency medical attention if you have: sudden shortness of breath
- chest pain or pressure that may spread to your jaw, shoulder, arms, or back
- nausea, vomiting , cold sweat
- a light-headed feeling, like you might pass out
- weakness on one side of your body
- slurred speech, drooping on one side of your mouth
- or pain, swelling, or redness in an arm or a leg. Baricitinib may cause serious side effects. Call your doctor at once if you have: fever, chills, night sweats, constant tiredness
- wheezing, trouble breathing, severe or worsening cough

Drug Side Effects Summary

- increased urination, pain or burning when you urinate
- unexplained weight loss
- lumps in your neck, armpits, or groin
- signs of shingles--flu-like symptoms, tingly or painful blistering rash on one side of your body
- a hole (perforation) in your digestive tract--fever, severe stomach pain, diarrhea or changes in bowel habits
- or signs of hepatitis--loss of appetite, vomiting, stomach pain (upper right side), dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of baricitinib may include: cold sores , shingles
- nausea
- or cold symptoms such as stuffy nose, sneezing, sore throat .

1136. This particular disease, Covid 19, interacts with the drug bebtelovimab and has the following

- Allergic reactions. Allergic reactions can happen during and after injection with bebtelovimab. Tell your healthcare provider right away if you or your child develop any of the following signs and symptoms of allergic reaction: fever difficulty breathing low oxygen level in your blood chills tiredness fast or slow heart rate chest discomfort or pain weakness confusion nausea headache shortness of breath low or high blood pressure wheezing swelling of your lips, face, or throat rash including hives itching muscle aches dizziness feeling faint, and sweating. These reactions may be severe or life threatening. Common side effects include infusion-related reactions, pruritus, and rash. The side effects of receiving any medicine by vein may include brief pain, bleeding, bruising of the skin, soreness, swelling, and possible infection at the injection site. These are not all the possible side effects. Not many people have received bebtelovimab. Serious and unexpected side effects may happen. All of the risks are not known at this time. Contact your healthcare provider if you have any side effects that bother you or do not go away. Report side effects to FDA MedWatch

Drug Side Effects Summary

at www.fda.gov/medwatch, or call 1-800-FDA-1088 or call Eli Lilly and Company, Inc. 1-855-LillyC19 (1-855-545-5921). It is possible that bebtelovimab could interfere with your body's own ability to fight off a future infection of SARS-CoV-2. Similarly, bebtelovimab may reduce the body's immune response to a vaccine for SARS-CoV-2. Talk to your healthcare provider if you have any questions.

Bebtelovimab side effects

1137. This particular disease, Covid 19, interacts with the drug Comirnaty and has the following side effects:

- Comirnaty may cause severe side effects including: Allergic reactions. A severe allergic reaction would usually occur within a few minutes to one hour after getting a dose of the vaccine. For this reason, your vaccination provider may ask you to stay at the place where you received your vaccine for monitoring after vaccination. Signs of a severe allergic reaction can include: Difficulty breathing Swelling of your face and throat A fast heartbeat A bad rash all over your body Dizziness and weakness Myocarditis and pericarditis. Myocarditis (inflammation of the heart muscle) and pericarditis (inflammation of the lining outside the heart) is a rare side effect and has occurred in some people who have received the vaccine. In most of these people, symptoms began within a few days following receipt of the second dose of vaccine and most individuals have had resolution of symptoms with conservative management. You should seek medical attention right away if you have any of the following symptoms after receiving the vaccine: Chest pain Shortness of breath Feelings of having a fast-beating, fluttering, or pounding heart Side effects can include: severe allergic reactions non-severe allergic reactions such as rash, itching, hives, or swelling of the face myocarditis (inflammation of the heart muscle) pericarditis (inflammation of the lining outside the heart) injection site pain tiredness headache muscle pain chills joint pain fever injection site swelling injection site redness nausea feeling unwell swollen lymph nodes (lymphadenopathy) diarrhea vomiting arm pain These may not be all the possible side effects of the vaccine. Serious and unexpected side effects may occur. The possible side effects of the vaccine are still being studied in

Drug Side Effects Summary

clinical trials. If you experience a severe allergic reaction, call 911, or go to the nearest hospital. Call the vaccination provider or your healthcare provider if you have any side effects that bother you or do not go away. Report vaccine side effects to FDA/CDC Vaccine Adverse Event Reporting System (VAERS). The VAERS toll-free number is 1-800-822-7967 or report online to <https://vaers.hhs.gov/reportevent.html>. Comirnaty side effects

1138. This particular disease, Covid 19, interacts with the drug nirmatrelvir / ritonavir and has the fo

- liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Ritonavir affects your immune system, which may cause certain side effects (even weeks or months after you've taken nirmatrelvir and ritonavir). Tell your doctor if you have: signs of a new infection--fever, night sweats , swollen glands, cold sores , cough, wheezing, diarrhea , weight loss
- trouble speaking or swallowing, problems with balance or eye movement, weakness or prickly feeling
- or swelling in your neck or throat (enlarged thyroid), menstrual changes, impotence . Common side effects of nirmatrelvir and ritonavir may include: changes in your sense of taste
- diarrhea
- elevated blood pressure
- or muscle pain.

1139. This particular disease, Covid 19, interacts with the drug sars-cov-2 (covid-19) mrna-1273 vac

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: More common Chills difficulty in moving fever headache joint pain muscle aches, cramps, pains, or stiffness nausea swollen joints

Drug Side Effects Summary

unusual tiredness or weakness vomiting Incidence not known Anxiety blue or pale skin chest discomfort or tightness chest pain, possible moving to the left arm, neck, or shoulder cough difficulty swallowing dizziness fast heartbeat fainting puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue trouble breathing Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Bleeding, blistering, burning, coldness, discoloration of the skin, feeling of pressure, hives, infection, inflammation, itching, lumps, numbness, pain, rash, redness, scarring, soreness, stinging, swelling, tenderness, tingling, ulceration, or warmth at the injection site Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Sars-cov-2 (covid-19) mrna-1273 vaccine side effects

1140. This particular disease, Covid 19, interacts with the drug Pfizer-BioNTech COVID-19 (5y-11y) I

- hives , rash
- dizziness , weakness, fast heartbeats
- difficult breathing
- swelling of the face, lips, tongue, or throat. Your child's vaccination provider may want to watch him/her for a short time after the shot, to make sure your child does not have an allergic reaction. Your child will be treated quickly if he/she has a reaction right after he/she receives the vaccine. Your child should not receive this vaccine if the first shot of COVID-19 vaccine caused an allergic reaction. The doctor or vaccination provider will determine if your child can safely receive this vaccine. Not all possible side effects are known. Becoming infected with COVID-19 is much more

Drug Side Effects Summary

dangerous to your child's health than receiving this vaccine. Some people receiving this vaccine had inflammation of the heart muscle or the lining around the heart within a few days after receiving this vaccine, but the risk of this side effect is very low. Seek medical attention right away if your child has: chest pain

- shortness of breath
- or fast or pounding heartbeats or fluttering in the chest. Common side effects of Pfizer-BioNTech COVID-19 (5y-11y) Bivalent Booster VaccinePF may include: fever, chills
- redness or a hardness and swelling where the shot was given
- swelling or tenderness under the arm
- nausea, vomiting
- feeling tired
- or headache, muscle pain, joint pain.

1141. This particular disease, Covid 19, interacts with the drug Pfizer-BioNTech COVID-19 (12y+) Bi

- hives , rash
- dizziness , weakness, fast heartbeats
- difficult breathing
- swelling of your face, lips, tongue, or throat. Your vaccination provider may want to watch you for a short time after your shot, to make sure you don't have an allergic reaction. You will be treated quickly if you have a reaction right after you receive the vaccine. You should not receive this vaccine if the first shot of COVID-19 vaccine caused an allergic reaction. Your doctor or vaccination provider will determine if you can safely receive this vaccine. Not all possible side effects are known. Becoming infected with COVID-19 is much more dangerous to your health than receiving this vaccine. Some people receiving this vaccine had inflammation of the heart muscle or the lining around the heart within a few days after receiving this vaccine, but the risk of this side effect is very

Drug Side Effects Summary

low. Seek medical attention right away if you have: chest pain

- shortness of breath
- or fast or pounding heartbeats or fluttering in your chest. Common side effects of Pfizer-BioNTech COVID-19 (12y+) Bivalent Booster Vaccine PF may include: fever, chills
- redness or a hardness and swelling where the shot was given
- swelling or tenderness under your arm
- nausea, vomiting
- feeling tired
- or headache, muscle pain, joint pain.

1142. This particular disease, Covid 19, interacts with the drug sars-cov-2 (covid-19) mrna (tozinam

- hives , rash
- dizziness , weakness, fast heartbeats
- difficult breathing
- swelling of the face, lips, tongue, or throat. Your child's vaccination provider may want to watch him/her for a short time after the shot, to make sure your child does not have an allergic reaction. Your child will be treated quickly if he/she has a reaction right after he/she receives the vaccine. Your child should not receive this vaccine if the first shot of COVID-19 vaccine caused an allergic reaction. The doctor or vaccination provider will determine if your child can safely receive this vaccine. Not all possible side effects are known. Becoming infected with COVID-19 is much more dangerous to your child's health than receiving this vaccine. Some people receiving this vaccine had inflammation of the heart muscle or the lining around the heart within a few days after receiving this vaccine, but the risk of this side effect is very low. Seek medical attention right away if your child has: chest pain
- shortness of breath

Drug Side Effects Summary

- or fast or pounding heartbeats or fluttering in the chest. Common side effects of COVID-19 (coronavirus 2019, 5y-11) bivalent booster vaccine, Pfizer may include: fever, chills
- redness or a hardness and swelling where the shot was given
- swelling or tenderness under the arm
- nausea, vomiting
- feeling tired
- or headache, muscle pain, joint pain.

1143. This particular disease, Covid 19, interacts with the drug sars-cov-2 (covid-19) mrna (tozinam

- hives , rash
- dizziness , weakness, fast heartbeats
- difficult breathing
- swelling of your face, lips, tongue, or throat. Your vaccination provider may want to watch you for a short time after your shot, to make sure you don't have an allergic reaction. You will be treated quickly if you have a reaction right after you receive the vaccine. You should not receive this vaccine if the first shot of COVID-19 vaccine caused an allergic reaction. Your doctor or vaccination provider will determine if you can safely receive this vaccine. Not all possible side effects are known. Becoming infected with COVID-19 is much more dangerous to your health than receiving this vaccine. Some people receiving this vaccine had inflammation of the heart muscle or the lining around the heart within a few days after receiving this vaccine, but the risk of this side effect is very low. Seek medical attention right away if you have: chest pain
- shortness of breath
- or fast or pounding heartbeats or fluttering in your chest. Common side effects of COVID-19 (coronavirus 2019, 12y+) bivalent booster vaccine, Pfizer may include: fever, chills
- redness or a hardness and swelling where the shot was given

Drug Side Effects Summary

- swelling or tenderness under your arm
- nausea, vomiting
- feeling tired
- or headache, muscle pain, joint pain.

1144. This particular disease, Covid 19, interacts with the drug sars-cov-2 (covid-19) mrna-1273 (biv)

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: More common Chills difficulty in moving headache joint pain muscle aches, cramps, pains, or stiffness nausea swollen joints unusual tiredness or weakness vomiting Less common Fever Incidence not known Anxiety blue or pale skin chest discomfort or tightness chest pain, possible moving to the left arm, neck, or shoulder cough difficulty swallowing dizziness fast heartbeat fainting puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue trouble breathing Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Bleeding, blistering, burning, coldness, discoloration of the skin, feeling of pressure, hives, infection, inflammation, itching, lumps, numbness, pain, rash, redness, scarring, soreness, stinging, swelling, tenderness, tingling, ulceration, or warmth at the injection site Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Sars-cov-2 (covid-19) mrna-1273 (bivalent booster) vaccine side effects

Drug Side Effects Summary

1145. This particular disease, Covid 19, interacts with the drug sars-cov-2 (covid-19) nvx-cov2373 v

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: More common Fever headache joint or muscle pain nausea unusual tiredness or weakness vomiting Rare Swollen, painful, or tender lymph glands in neck, armpit, or groin Incidence not known Anxiety blue or pale skin burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest discomfort or tightness chest pain, possibly moving to the left arm, neck, or shoulder chills cough difficulty swallowing dizziness fainting fast heartbeat hives, itching, skin rash puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue trouble breathing Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Pain, redness, or swelling at the injection site Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Sars-cov-2 (covid-19) nvx-cov2373 vaccine, recombinant side effects

1146. This particular disease, Covid 19, interacts with the drug sars-cov-2 mrna (tozinameran 6m-4y

- hives , rash
- dizziness , weakness, fast heartbeats
- difficult breathing
- swelling of the face, lips, tongue, or throat. The vaccination provider may want to watch your child for a short time after the shot, to make sure he or she doesn't have an allergic reaction. Your child

Drug Side Effects Summary

will be treated quickly if he or she has a reaction right after receiving the vaccine. Your child should not receive this vaccine again if the first shot caused an allergic reaction. The doctor or vaccination provider will determine if he or she can safely receive another COVID-19 vaccine. Not all possible side effects are known. Becoming infected with COVID-19 is much more dangerous to your child's health than receiving this vaccine. Some people receiving this vaccine had inflammation of the heart muscle or the lining around the heart within a few days after receiving this vaccine, but the risk of this side effect is very low. Seek medical attention right away if your child has: chest pain

- shortness of breath
- or fast or pounding heartbeats or fluttering in the chest. Common side effects of COVID-19 (coronavirus 2019, 6m-4y) vaccine, Pfizer may include: fever, chills, swollen glands, not feeling well
- pain, redness, or swelling where the shot was given
- feeling tired
- or headache, muscle pain, joint pain.

1147. This particular disease, Covid 19, interacts with the drug sars-cov-2 mrna-1273 (6m-5y) vaccine

- hives , rash
- dizziness , weakness, fast heartbeats
- difficult breathing
- swelling of your face, lips, tongue, or throat. Your vaccination provider may want to watch you for a short time after your shot, to make sure you don't have an allergic reaction. You will be treated quickly if you have a reaction right after you receive the vaccine. You should not receive this vaccine again if the first shot caused an allergic reaction. Your doctor or vaccination provider will determine if you can safely receive another COVID-19 vaccine. Not all possible side effects are known. Becoming infected with COVID-19 is much more dangerous to your health than receiving this vaccine. Some people receiving this vaccine had inflammation of the heart muscle or the lining

Drug Side Effects Summary

around the heart within a few days after receiving this vaccine, but the risk of this side effect is very low. Seek medical attention right away if you have: chest pain

- shortness of breath
- or fast or pounding heartbeats or fluttering in your chest. Common side effects of COVID-19 (coronavirus 2019, 6m-5y) vaccine, Moderna may include: fever, chills
- redness or a hardness and swelling where the shot was given
- swelling or tenderness under your arm
- nausea, vomiting
- feeling tired
- or headache, muscle pain, joint pain.

1148. This particular disease, Covid 19, interacts with the drug sars-cov-2 mrna-1273 (booster only)

- hives , rash
- dizziness , weakness, fast heartbeats
- difficult breathing
- swelling of your face, lips, tongue, or throat. Your vaccination provider may want to watch you for a short time after your shot, to make sure you don't have an allergic reaction. You will be treated quickly if you have a reaction right after you receive the vaccine. You should not receive this vaccine again if the first shot caused an allergic reaction. Your doctor or vaccination provider will determine if you can safely receive another COVID-19 vaccine. Not all possible side effects are known. Becoming infected with COVID-19 is much more dangerous to your health than receiving this vaccine. Some people receiving this vaccine had inflammation of the heart muscle or the lining around the heart within a few days after receiving this vaccine, but the risk of this side effect is very low. Seek medical attention right away if you have: chest pain
- shortness of breath

Drug Side Effects Summary

- or fast or pounding heartbeats or fluttering in your chest. Common side effects of COVID-19 (coronavirus 2019, 6y-11y) vaccine, Moderna may include: fever, chills
- redness or a hardness and swelling where the shot was given
- swelling or tenderness under your arm
- nausea, vomiting
- feeling tired
- or headache, muscle pain, joint pain.

1149. This particular disease, Covid 19, interacts with the drug Spikevax and has the following side

- Side effects that have been reported in clinical trials with Spikevax include: Injection site reactions: pain, tenderness and swelling of the lymph nodes in the same arm of the injection, swelling (hardness), and redness General side effects: fatigue, headache, muscle pain, joint pain, chills, nausea and vomiting, and fever In trial participants 18 through 64 years , the most commonly reported (?10%) adverse reactions were pain at injection site (93.3%), fatigue (71.9%), headache (68.7%), myalgia (64.8%), chills (49.7%), arthralgia (48.6%), nausea/vomiting (25.7%), axillary swelling/tenderness (22.2%), fever (17.3%), swelling at the injection site (15.4%), and erythema at the injection site (10.5%). In trial participants 65 years of age and older , the most commonly reported (?10%) adverse reactions were pain at injection site (88.3%), fatigue (64.8%), headache (53.3%), myalgia (51.8%), arthralgia (40.2%), chills (32.7%), nausea/vomiting (15.0%), swelling at the injection site (13.0%), and axillary swelling/tenderness (12.7%). Side effects that have been reported during post-authorization use of the vaccine include: Severe allergic reactions Myocarditis (inflammation of the heart muscle) Pericarditis (inflammation of the lining outside the heart) Fainting in association with injection of the vaccine These may not be all of the possible side effects of Spikevax. Ask your healthcare provider about any side effects that concern you. You may report side effects to the Vaccine Adverse Event Reporting System (VAERS) at 1-800-822-7967 or

Drug Side Effects Summary

<https://vaers.hhs.gov>. Spikevax side effects

1150. This particular disease, Depression, interacts with the drug Zoloft and has the following side

- skin rash or hives (with or without fever or joint pain)
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Zoloft may cause serious side effects. Call your doctor at once if you have: unusual bleeding or bruising
- a seizure
- vision changes, eye pain , redness, or swelling
- low blood sodium - headache , confusion, problems with thinking or memory, weakness, feeling unsteady
- or manic episodes - racing thoughts, increased energy, unusual risk-taking behavior, extreme happiness, being irritable or talkative. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting , or diarrhea . Sertraline can affect growth in children. Your child's height and weight may be checked often. Common side effects of Zoloft may include: indigestion
- nausea
- diarrhea
- loss of appetite
- loose stool
- increased sweating

Drug Side Effects Summary

- tremors
- or sexual problems.

1151. This particular disease, Depression, interacts with the drug Prozac and has the following side effects:

- mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Prozac may cause serious side effects. Call your doctor at once if you have: blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights
- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- low levels of sodium in the body - headache , confusion, slurred speech, severe weakness, vomiting , loss of coordination, feeling unsteady
- or severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea . Common side effects of Prozac may include: sleep problems (insomnia), strange dreams
- headache, dizziness, drowsiness, vision changes
- tremors or shaking, feeling anxious or nervous
- pain, weakness, yawning, tired feeling
- upset stomach , loss of appetite, nausea, vomiting, diarrhea
- dry mouth, sweating, hot flashes
- changes in weight or appetite
- stuffy nose, sinus pain, sore throat, flu symptoms

Drug Side Effects Summary

- or decreased sex drive, impotence , or difficulty having an orgasm.

1152. This particular disease, Depression, interacts with the drug fluoxetine and has the following s

- mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Fluoxetine may cause serious side effects. Call your doctor at once if you have: blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights

- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)

- low levels of sodium in the body - headache , confusion, slurred speech, severe weakness, vomiting , loss of coordination, feeling unsteady

- or severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea . Common fluoxetine side effects may include: sleep problems (insomnia), strange dreams

- headache, dizziness, drowsiness, vision changes

- tremors or shaking, feeling anxious or nervous

- pain, weakness, yawning, tired feeling

- upset stomach , loss of appetite, nausea, vomiting, diarrhea

- dry mouth, sweating, hot flashes

- changes in weight or appetite

- stuffy nose, sinus pain, sore throat, flu symptoms

- or decreased sex drive, impotence , or difficulty having an orgasm.

Drug Side Effects Summary

1153. This particular disease, Depression, interacts with the drug Wellbutrin XL and has the following side effects:

- mood or behavior changes, anxiety , depression, panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Wellbutrin XL may cause serious side effects. Call your doctor at once if you have: a seizure (convulsions)
- confusion, unusual changes in mood or behavior
- blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights
- fast or irregular heartbeats
- or a manic episode--racing thoughts, increased energy, reckless behavior, feeling extremely happy or irritable, talking more than usual, severe problems with sleep. Common side effects of Wellbutrin XL may include: dry mouth, sore throat, stuffy nose
- ringing in the ears
- blurred vision
- nausea , vomiting , stomach pain, loss of appetite, constipation
- sleep problems (insomnia)
- tremors, sweating, feeling anxious or nervous
- fast heartbeats
- confusion, agitation , hostility
- rash
- weight loss
- increased urination
- headache , dizziness
- or muscle or joint pain.

1154. This particular disease, Depression, interacts with the drug Celexa and has the following side effects:

Drug Side Effects Summary

- hives , rash, blisters
- fever, joint pain
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tell your doctor right away if you have new or sudden changes in mood or behavior, including new or worse depression or anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, more active or talkative, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- blurred vision, eye pain or redness, seeing halos around lights
- fast or pounding heartbeats, pain or fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- a seizure
- manic episodes - racing thoughts, increased energy, decreased need for sleep, risk-taking behavior, being agitated or talkative
- severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors
- or low blood sodium - headache , confusion, problems with thinking or memory, weakness, feeling unsteady. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting , or diarrhea . Common Celexa side effects may include: sexual problems
- dizziness, drowsiness
- dry mouth, thirst, increased sweating or urination
- loss of appetite, nausea, diarrhea, constipation
- feeling anxious, agitated, or shaky

Drug Side Effects Summary

- feeling weak or tired
- sleep problems (insomnia)
- yawning
- increased muscle movement
- nosebleeds , heavy menstrual bleeding
- or cold symptoms such as stuffy nose, sneezing, sore throat .

1155. This particular disease, Depression, interacts with the drug Trintellix and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: racing thoughts, decreased need for sleep, unusual risk-taking behavior, feelings of extreme happiness or sadness, being more talkative than usual
- vision changes, eye pain , eye redness or swelling
- easy bruising, unusual bleeding, coughing up blood
- or low sodium level (may be more likely to occur in older adults - confusion, memory problems, hallucinations, slurred speech, severe weakness, feeling unsteady. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting , or diarrhea . Common Trintellix side effects may include: nausea
- constipation
- or vomiting.

Drug Side Effects Summary

1156. This particular disease, Depression, interacts with the drug Rexulti and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. High doses or long-term use of Rexulti can cause a serious movement disorder that may not be reversible. The longer you use this medicine, the more likely you are to develop this disorder, especially if you are a diabetic or an older adult. Call your doctor at once if you have: uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- trouble swallowing
- feelings of warmth, intolerance to heat
- a seizure (convulsions)
- a light-headed feeling, like you might pass out
- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- low white blood cell counts - fever, mouth sores, skin sores, sore throat , cough, trouble breathing
- severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors
- or signs of a blood clot - sudden numbness or weakness, problems with vision or speech, swelling or redness in an arm or leg. You may have increased sexual urges, unusual urges to gamble, or other intense urges while taking this medicine. Talk with your doctor if this occurs. Common Rexulti side effects may include: weight gain
- or feeling restless or being unable to sit still.

Drug Side Effects Summary

1157. This particular disease, Depression, interacts with the drug vortioxetine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Vortioxetine may cause serious side effects. Call your doctor at once if you have: racing thoughts, decreased need for sleep, unusual risk-taking behavior, feelings of extreme happiness or sadness, being more talkative than usual
- vision changes, eye pain , eye redness or swelling
- easy bruising, unusual bleeding, coughing up blood
- or low sodium level (may be more likely to occur in older adults--confusion, memory problems, hallucinations, slurred speech, severe weakness, feeling unsteady. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting , or diarrhea . Common side effects of vortioxetine may include: nausea
- constipation
- or vomiting.

1158. This particular disease, Depression, interacts with the drug brexpiprazole and has the followi

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more

Drug Side Effects Summary

depressed, or have thoughts about suicide or hurting yourself. High doses or long-term use of brexpiprazole can cause a serious movement disorder that may not be reversible. The longer you use brexpiprazole, the more likely you are to develop this disorder, especially if you are a diabetic or an older adult. Brexpiprazole may cause serious side effects. Call your doctor at once if you have: uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)

- trouble swallowing
- feelings of warmth, intolerance to heat
- a seizure (convulsions)
- a light-headed feeling, like you might pass out
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- low white blood cell counts--fever, mouth sores, skin sores, sore throat , cough, trouble breathing
- severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors
- or signs of a blood clot--sudden numbness or weakness, problems with vision or speech, swelling or redness in an arm or leg. You may have increased sexual urges, unusual urges to gamble, or other intense urges while taking this medicine. Talk with your doctor if this occurs. Common side effects of brexpiprazole may include: weight gain
- or feeling restless or being unable to sit still.

1159. This particular disease, Depression, interacts with the drug Pristiq and has the following side

- skin rash or hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Tell your doctor right away if you have new or sudden changes in mood or behavior, including new or worse depression or anxiety, panic attacks, trouble

Drug Side Effects Summary

sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, more active or talkative, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: a seizure

- easy bruising or bleeding (nosebleeds , bleeding gums), blood in your urine or stools, coughing up blood

- blurred vision, eye pain or swelling, or seeing halos around lights

- cough, chest discomfort, trouble breathing

- or low blood sodium - headache, confusion, problems with thinking or memory, weakness, feeling unsteady. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Common Pristiq side effects may include: dizziness , drowsiness, anxiety

- increased sweating

- nausea, decreased appetite, constipation

- sleep problems (insomnia)

- or decreased sex drive, impotence , or difficulty having an orgasm.

1160. This particular disease, Depression, interacts with the drug Effexor and has the following side effects:

- No side effects listed

1161. This particular disease, Depression, interacts with the drug Remeron and has the following side effects:

- mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: racing thoughts, decreased need for sleep, unusual risk-taking behavior, feelings of extreme happiness or

Drug Side Effects Summary

sadness, being more talkative than usual

- blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights
- a light-headed feeling, like you might pass out
- severe rash, blisters, or swelling on the palms of your hands or the soles of your feet
- a seizure
- low white blood cell counts - fever, chills, sore throat, cough, sores in your mouth or nose, flu-like symptoms, trouble breathing
- or low sodium level - headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea . Common Remeron side effects include: drowsiness, dizziness
- increased appetite
- or weight gain.

1162. This particular disease, Depression, interacts with the drug Wellbutrin SR and has the following

- mood or behavior changes, anxiety , depression, panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Wellbutrin SR may cause serious side effects. Call your doctor at once if you have: a seizure (convulsions)
- confusion, unusual changes in mood or behavior
- blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights
- fast or irregular heartbeats
- or a manic episode--racing thoughts, increased energy, reckless behavior, feeling extremely happy or irritable, talking more than usual, severe problems with sleep. Common side effects of Wellbutrin

Drug Side Effects Summary

SR may include: dry mouth, sore throat, stuffy nose

- ringing in the ears
- blurred vision
- nausea , vomiting , stomach pain, loss of appetite, constipation
- sleep problems (insomnia)
- tremors, sweating, feeling anxious or nervous
- fast heartbeats
- confusion, agitation , hostility
- rash
- weight loss
- increased urination
- headache , dizziness
- or muscle or joint pain.

1163. This particular disease, Depression, interacts with the drug amitriptyline and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: signs of a blood clot - sudden numbness or weakness, problems with vision or speech, swelling or redness in an arm or leg
- unusual thoughts or behavior
- a light-headed feeling, like you might pass out

Drug Side Effects Summary

- chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating
- pounding heartbeats or fluttering in your chest
- confusion, hallucinations
- a seizure (convulsions)
- painful or difficult urination
- severe constipation
- easy bruising, unusual bleeding
- or fever, chills, sore throat , mouth sores. Common amitriptyline side effects may include:
constipation, diarrhea
- nausea, vomiting , upset stomach
- mouth pain, unusual taste, black tongue
- appetite or weight changes
- urinating less than usual
- itching or rash
- breast swelling (in men or women)
- or decreased sex drive, impotence , or difficulty having an orgasm.

1164. This particular disease, Depression, interacts with the drug Viibryd and has the following side effects:

- skin rash or hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have:
a seizure (convulsions)

Drug Side Effects Summary

- blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights
- easy bruising, unusual bleeding
- racing thoughts, unusual risk-taking behavior, decreased inhibitions, feelings of extreme happiness or sadness
- or low levels of sodium in the body - headache , confusion, slurred speech, severe weakness, loss of coordination, feeling unsteady. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting , or diarrhea . Common Viibryd side effects may include: nausea, vomiting
- diarrhea
- or sleep problems (insomnia).

1165. This particular disease, Depression, interacts with the drug Deplin and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Less serious side effects may occur, and you may have none at all.

1166. This particular disease, Depression, interacts with the drug nortriptyline and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have:

Drug Side Effects Summary

blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights

- restless muscle movements in your eyes, tongue, jaw, or neck
- a light-headed feeling, like you might pass out
- seizure (convulsions)
- new or worsening chest pain, pounding heartbeats or fluttering in your chest
- sudden numbness or weakness, problems with vision, speech, or balance
- fever, sore throat , easy bruising, unusual bleeding
- painful or difficult urination
- or jaundice (yellowing of the skin or eyes). Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea . Common nortriptyline side effects may include: increased blood pressure
- numbness or tingling in your hands or feet
- dry mouth, nausea, vomiting, loss of appetite
- blurred vision
- rash, itching
- or breast swelling (in men or women).

1167. This particular disease, Depression, interacts with the drug Desyrel and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop taking Desyrel and call your doctor at once if you have a penis erection that is painful or lasts 6 hours or longer. This is a medical emergency and could lead to a serious condition that must be corrected with surgery. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble

Drug Side Effects Summary

sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Desyrel may cause serious side effects. Call your doctor at once if you have: fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)

- slow heartbeats
- unusual thoughts or behavior
- easy bruising, unusual bleeding
- or low levels of sodium in the body-- headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea .

Common side effects of Desyrel may include: drowsiness, dizziness, tiredness

- swelling
- weight loss
- blurred vision
- diarrhea, constipation
- or stuffy nose.

1168. This particular disease, Depression, interacts with the drug Forfivo XL and has the following

- Forfivo XL can cause serious side effects, including: See Important information about Forfivo XL
- The most common side effects of Forfivo XL include: dry mouth nausea trouble sleeping dizziness sore throat stomach pain agitation anxiety shakiness fast heart beat sweating ringing in the ears muscle pain anorexia urinating more often than normal for you rash If you have nausea, take your medicine with food. If you have trouble sleeping, do not take your medicine too close to bedtime. Tell your healthcare provider right away about any side effects that bother you. These are not all the

Drug Side Effects Summary

side effects of Forfivo XL. For a complete list, ask your healthcare provider or pharmacist. Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Forfivo XL side effects

1169. This particular disease, Depression, interacts with the drug Budeprion SR and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Anxiety dry mouth hyperventilation irregular heartbeats irritability restlessness shaking trouble sleeping Less common Buzzing or ringing in the ears headache (severe) skin rash, hives, or itching Rare Confusion fainting false beliefs that cannot be changed by facts having extreme distrust of people seeing, hearing, or feeling things that are not there seizures trouble concentrating Incidence not known Actions that are out of control anger assaulting or attacking others being aggressive or impulsive chest pain or discomfort fast or pounding heartbeat force inability to sit still need to keep moving sweating talking, feeling, or acting with excitement Symptoms of overdose Lightheadedness, dizziness, or fainting loss of consciousness slow or irregular heartbeat unusual tiredness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Constipation decrease in appetite dizziness increased sweating stomach pain trembling unusual weight loss Less common Blurred vision change in sense of taste drowsiness frequent need to urinate sore throat unusual feeling of well-being Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at

Drug Side Effects Summary

1-800-FDA-1088. Budeprion SR side effects

1170. This particular disease, Depression, interacts with the drug Aplenzin and has the following side effects:

- hives, itching, fever, swollen glands, difficult breathing, swelling in your face or throat) or a severe skin reaction (fever, sore throat, burning eyes, skin pain, red or purple skin rash with blistering and peeling). Report any new or worsening symptoms to your doctor , such as: mood or behavior changes, anxiety, depression, panic attacks, trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: a seizure (convulsions)
- confusion, unusual changes in mood or behavior
- blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights
- fast or irregular heartbeats
- or a manic episode - racing thoughts, increased energy, reckless behavior, feeling extremely happy or irritable, talking more than usual, severe problems with sleep. Common Aplenzin side effects may include: dry mouth, sore throat, stuffy nose
- ringing in the ears
- blurred vision
- nausea, vomiting, stomach pain, loss of appetite, constipation
- sleep problems (insomnia)
- tremors, sweating, feeling anxious or nervous
- fast heartbeats
- confusion, agitation, hostility
- rash
- weight loss

Drug Side Effects Summary

- increased urination
- headache, dizziness
- or muscle or joint pain.

1171. This particular disease, Depression, interacts with the drug Desyrel Dividose and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Blurred vision confusion dizziness dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position lightheadedness sweating unusual tiredness or weakness Less common Burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings confusion about identity, place, and time decreased concentration fainting general feeling of discomfort or illness headache lack of coordination muscle tremors nervousness pounding in the ears shortness of breath slow or fast heartbeat swelling Rare Skin rash unusual excitement Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Dry mouth (usually mild) muscle or bone pain trouble sleeping trouble with remembering unpleasant taste Less common Constipation continuing ringing or buzzing or other unexplained noise in the ears diarrhea hearing loss muscle aches or pains weight loss Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Desyrel Dividose side effects

Drug Side Effects Summary

1172. This particular disease, Depression, interacts with the drug Fetzima and has the following side effects:

- skin rash or hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tell your doctor right away if you have new or sudden changes in mood or behavior, including new or worse depression or anxiety, panic attacks, trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, more active or talkative, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: blurred vision, tunnel vision, eye pain or swelling, eye redness
- painful or difficult urination
- easy bruising, unusual bleeding, purple or red spots under your skin
- pounding heartbeats or fluttering in your chest
- seizure (convulsions)
- manic episodes - racing thoughts, increased energy, decreased need for sleep, risk-taking behavior, being agitated or talkative
- or low sodium level - headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady. Serious side effects may be more likely in older adults. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Common Fetzima side effects may include: nausea, vomiting, constipation
- irregular heartbeats
- sweating
- or decreased sex drive, impotence , or difficulty having an orgasm.

1173. This particular disease, Depression, interacts with the drug Irenka and has the following side effects:

- pounding heartbeats or fluttering in your chest

Drug Side Effects Summary

- a light-headed feeling, like you might pass out
- easy bruising, unusual bleeding
- vision changes
- painful or difficult urination
- liver problems--right-sided upper stomach pain, itching, dark urine, jaundice (yellowing of the skin or eyes)
- low blood sodium-- headache , confusion, problems with thinking or memory, weakness, feeling unsteady
- or manic episodes--racing thoughts, increased energy, decreased need for sleep, risk-taking behavior, being agitated or talkative. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea . Common side effects of Irenka may include: drowsiness
- nausea, constipation , loss of appetite
- dry mouth
- or increased sweating.

1174. This particular disease, Depression, interacts with the drug fluvoxamine and has the following

- skin rash , blisters, or hives
- fever, joint pain
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tell your doctor right away if you have new or sudden changes in mood or behavior, including new or worse depression or anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, more active or talkative, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have

Drug Side Effects Summary

- anxiety, racing thoughts, risk-taking behavior, sleep problems (insomnia), feelings of extreme happiness or irritability
- blurred vision, eye pain or redness, seeing halos around lights
- a seizure
- changes in weight or appetite
- easy bruising or unusual bleeding
- or low blood sodium-- headache , confusion, problems with thinking or memory, weakness, feeling unsteady. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea Fluvoxamine can affect growth in children. Your child's height and weight should be checked often. Common side effects of fluvoxamine may include: drowsiness, dizziness , weakness
- anxiety, depression, agitation, trouble sleeping
- shaking, increased muscle movements
- upset stomach , gas, loss of appetite
- nausea, vomiting, diarrhea
- dry mouth, yawning, sore throat
- muscle pain
- sweating, rash
- heavy menstrual periods
- or sexual problems.

1175. This particular disease, Depression, interacts with the drug l-methylfolate and has the following

- hives
- difficulty breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Less serious side effects may occur, and you may have none at all.

1176. This particular disease, Depression, interacts with the drug Parnate and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Parnate may cause serious side effects. Call your doctor at once if you have: dilated pupils, vision problems, sensitivity to light
- sudden and severe headache, neck pain or stiffness
- numbness or weakness, problems with vision or speech
- fast or pounding heartbeats
- fever, cold sweat, nausea, vomiting
- a light-headed feeling, like you might pass out
- a seizure
- manic episodes--racing thoughts, increased energy, decreased need for sleep, risk-taking behavior, being agitated or talkative
- high levels of serotonin in the body--agitation, hallucinations, sweating, shivering, muscle stiffness, twitching, loss of coordination, vomiting, diarrhea
- or liver problems--loss of appetite, stomach pain (upper right side), dark urine, jaundice (yellowing of the skin or eyes). Serious side effects may be more likely in older adults. Common side effects of Parnate may include: dizziness , drowsiness
- constipation

Drug Side Effects Summary

- dry mouth, blurred vision
- trouble sleeping
- headache
- or feeling shaky or excited.

1177. This particular disease, Depression, interacts with the drug Remeron SolTab and has the following side effects:

- mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Remeron SolTab may cause serious side effects. Call your doctor at once if you have: racing thoughts, decreased need for sleep, unusual risk-taking behavior, feelings of extreme happiness or sadness, being more talkative than usual
- blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights
- a light-headed feeling, like you might pass out
- severe rash, blisters, or swelling on the palms of your hands or the soles of your feet
- a seizure
- low white blood cell counts--fever, chills, sore throat, cough, sores in your mouth or nose, flu-like symptoms, trouble breathing
- or low sodium level -- headache , confusion, slurred speech, severe weakness, vomiting , loss of coordination, feeling unsteady. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea . Common side effects include: drowsiness, dizziness
- increased appetite
- or weight gain.

Drug Side Effects Summary

1178. This particular disease, Depression, interacts with the drug amoxapine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Amoxapine may cause serious side effects. Call your doctor at once if you have: skin rash , fever
- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- pounding heartbeats or fluttering in your chest
- chest pain or pressure, pain spreading to your jaw or shoulder
- sudden numbness or weakness (especially on one side of the body), slurred speech, problems with vision or balance
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Common side effects of amoxapine may include: drowsiness
- constipation
- dry mouth
- or blurred vision.

1179. This particular disease, Depression, interacts with the drug Aventyl Hydrochloride and has the

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Agitation blurred

Drug Side Effects Summary

vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain or discomfort clay-colored stools cold sweats confusion about identity, place, and time false beliefs that cannot be changed by facts continuing ringing or buzzing or other unexplained noise in the ears decreased urination deep or fast breathing with dizziness depression difficulty in passing urine (dribbling) difficulty with speaking dizziness, fainting, or lightheadedness when getting up suddenly from a lying or sitting position double vision fast, pounding, or irregular heartbeat or pulse feeling of warmth feeling, seeing, or hearing things that are not there feeling that others are watching you or controlling your behavior feeling that others can hear your thoughts general feeling of tiredness or weakness hives or welts hostility inability to move the arms, legs, or facial muscles inability to speak irritability labored breathing, especially at night loss of balance control loss of consciousness lower back or side pain mood or mental changes muscle spasm, trembling, jerking, or stiffness nightmares numbness of the feet, hands, and around mouth pain or discomfort in the arms, jaw, back, or neck painful or difficult urination panic perspiration pinpoint red or purple spots on the skin redness of the face, neck, arms, and occasionally, upper chest restlessness seizures slurred speech sores, ulcers, or white spots on the lips or in the mouth stiffness of the limbs stomach pain sweating swelling of the face, ankles, legs, or hands swollen, painful, or tender lymph glands on side of face or neck talking, feeling, and acting with excitement trouble sleeping twisting movements of the body uncontrolled movements, especially of the face, neck, and back unexplained fainting weakness in the arms, hands, legs, or feet Symptoms of overdose Chest pain or tightness clumsiness cold, clammy skin confusion decreased awareness or responsiveness decreased urine output dilated neck veins extreme tiredness or weakness fainting fast, slow, or irregular heartbeat fast, weak pulse increased or excessive unconscious or jerking movements irregular breathing lightheadedness loss of consciousness low body temperature muscle stiffness, weakness, or tightness restlessness seeing things that are not there seizures severe sleepiness shivering swelling of the face, fingers, feet, or lower legs troubled breathing weak or feeble pulse

Drug Side Effects Summary

weight gain Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Bigger, dilated, or enlarged pupils (black part of the eye) black tongue decreased interest in sexual ability or desire difficulty having a bowel movement enlargement of the breast hair loss or thinning of the hair heartburn increase in sexual ability or desire increased sensitivity of the eyes or skin to light pain or discomfort in the chest, upper stomach, or throat peculiar taste severe sunburn small red or purple spots on the skin swelling of the testicles swelling or inflammation of the mouth waking to urinate at night Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Aventyl Hydrochloride side effects

1180. This particular disease, Depression, interacts with the drug clomipramine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash , fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Clomipramine may cause serious side effects. Call your doctor at once if you have: low sodium level -- headache , confusion, slurred speech, severe weakness, vomiting , loss of

Drug Side Effects Summary

coordination, feeling unsteady

- blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights
- confusion, extreme fear, thoughts of hurting yourself
- pain or burning when you urinate
- or a seizure (convulsions). Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea . Common side effects of clomipramine may include: feeling dizzy, drowsy, tired, or nervous
- tremors, jerking muscle movements
- sweating
- changes in appetite or weight
- urination problems
- vision changes
- or decreased sex drive, impotence , or difficulty having an orgasm.

1181. This particular disease, Depression, interacts with the drug Emsam and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Remove the skin patch and get emergency medical help if you have any of these signs of dangerously high blood pressure : sudden and severe headache, confusion, vision problems, chest pain , pounding in your neck or ears, sweating, vomiting , neck stiffness, fast or slow heartbeats, or dilated pupils. Call your doctor at once

Drug Side Effects Summary

if you have: sudden numbness or weakness (especially on one side of the body)

- problems with speech or balance
- manic episodes--racing thoughts, increased energy, unusual risk-taking behavior, extreme happiness, being irritable or talkative. Remove the patch and seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea . Common Emsam side effects may include: redness or itching where the patch is worn
- headache
- diarrhea, upset stomach , dry mouth
- sleep problems (insomnia)
- rash
- or sinus pain or stuffy nose.

1182. This particular disease, Depression, interacts with the drug ketamine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tell your caregivers if you have hallucinations or unusual thoughts while waking up from anesthesia. Ketamine may cause serious side effects. Call your doctor at once if you have: painful or difficult urination, increased urination, loss of bladder control, blood in your urine
- a light-headed feeling, like you might pass out
- slow heart rate , weak or shallow breathing
- or jerky muscle movements that may look like convulsions. Common side effects of ketamine may include: confusion
- or dream-like feeling.

Drug Side Effects Summary

1183. This particular disease, Depression, interacts with the drug Nardil and has the following side

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Nardil may cause serious side effects. Call your doctor at once if you have: sudden and severe headache, neck pain or stiffness
- pounding heartbeats or fluttering in your chest
- fast or slow heartbeats
- chest pain
- a light-headed feeling, like you might pass out
- sweating (sometimes with fever and sometimes with cold, clammy skin)
- nausea, vomiting
- or dilated pupils (your eyes may be more sensitive to light). Common side effects of Nardil may include: dizziness , feeling light-headed
- drowsiness, sleep problems
- headache
- feeling weak or tired
- tremors, muscle twitching
- dry mouth, stomach discomfort, constipation
- swelling, weight gain
- or sexual problems.

1184. This particular disease, Depression, interacts with the drug Norpramin and has the following

Drug Side Effects Summary

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Abdominal or stomach pain anxiety black, tarry stools blurred or double vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain or discomfort confusion about identity, place, and time convulsions cough or hoarseness dark urine dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position dry mouth fainting fast, irregular, pounding, or racing heartbeat or pulse inability to move the arms, legs, or facial muscles inability to speak irritability light-colored stools lightheadedness loss of appetite loss of bladder control muscle spasms or jerking of all extremities nausea nightmares pain or discomfort in the arms, jaw, back, or neck painful or difficult urination pinpoint red or purple spots on the skin pounding in the ears rash restlessness seeing, hearing, or feeling things that are not there shakiness and unsteady walk slow speech sore throat and fever sores, ulcers, or white spots on the lips or in the mouth stiffness of the limbs sudden loss of consciousness sweating swelling swollen glands talking, feeling, and acting with excitement trouble sleeping twisting body movements unsteadiness, trembling, or problems with muscle control or coordination unusual bleeding or bruising unusual tiredness or weakness vomiting yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Abdominal or stomach cramps bigger, dilated, or enlarged pupils (black part of the eye) bloating constipation decrease in the frequency of urination decrease in urine volume decreased interest in sexual intercourse diarrhea difficulty in passing urine (dribbling) hair loss or thinning of the hair inability to have or keep an erection increased in sexual ability, desire,

Drug Side Effects Summary

drive, or performance increased interest in sexual intercourse increased sensitivity of the eyes to light loss in sexual ability, desire, drive, or performance painful urination swelling of the breasts or breast soreness in both females and males swelling or inflammation of the mouth unexpected or excess milk flow from breasts in females Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Norpramin side effects

1185. This particular disease, Depression, interacts with the drug Pamelor and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Pamelor may cause serious side effects. Call your doctor at once if you have: blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights
- restless muscle movements in your eyes, tongue, jaw, or neck
- a light-headed feeling, like you might pass out
- seizure (convulsions)
- new or worsening chest pain, pounding heartbeats or fluttering in your chest
- sudden numbness or weakness, problems with vision, speech, or balance
- fever, sore throat , easy bruising, unusual bleeding
- painful or difficult urination
- or jaundice (yellowing of the skin or eyes). Seek medical attention right away if you have symptoms

Drug Side Effects Summary

of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea . Common side effects of Pamelor may include: increased blood pressure

- numbness or tingling in your hands or feet
- dry mouth, nausea, vomiting, loss of appetite
- blurred vision
- rash, itching
- or breast swelling (in men or women).

1186. This particular disease, Depression, interacts with the drug phenelzine and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Phenelzine may cause serious side effects. Call your doctor at once if you have: sudden and severe headache, neck pain or stiffness
- pounding heartbeats or fluttering in your chest
- fast or slow heartbeats
- chest pain
- a light-headed feeling, like you might pass out
- sweating (sometimes with fever and sometimes with cold, clammy skin)
- nausea, vomiting
- or dilated pupils (your eyes may be more sensitive to light). Common side effects of phenelzine may include: dizziness , feeling light-headed

Drug Side Effects Summary

- drowsiness, sleep problems
- headache
- feeling weak or tired
- tremors, muscle twitching
- dry mouth, stomach discomfort, constipation
- swelling, weight gain
- or sexual problems.

1187. This particular disease, Depression, interacts with the drug tranylcypromine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Tranylcypromine may cause serious side effects. Call your doctor at once if you have: dilated pupils, vision problems, sensitivity to light
- sudden and severe headache, neck pain or stiffness
- numbness or weakness, problems with vision or speech
- fast or pounding heartbeats
- fever, cold sweat, nausea, vomiting
- a light-headed feeling, like you might pass out
- a seizure
- manic episodes--racing thoughts, increased energy, decreased need for sleep, risk-taking behavior, being agitated or talkative
- high levels of serotonin in the body--agitation, hallucinations, sweating, shivering, muscle stiffness,

Drug Side Effects Summary

twitching, loss of coordination, vomiting, diarrhea

- or liver problems--loss of appetite, stomach pain (upper right side), dark urine, jaundice (yellowing of the skin or eyes). Serious side effects may be more likely in older adults. Common side effects of tranlycypromine may include: dizziness , drowsiness

- constipation
- dry mouth, blurred vision
- trouble sleeping
- headache
- or feeling shaky or excited.

1188. This particular disease, Depression, interacts with the drug isocarboxazid and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, panic attacks, trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Stop taking isocarboxazid and call your doctor at once if you have: sudden and severe headache , rapid heartbeat, stiffness in your neck, nausea , vomiting , cold sweat, vision problems, sensitivity to light
- chest pain, fast or slow heart rate
- swelling, rapid weight gain
- jaundice (yellowing of the skin or eyes)
- or a light-headed feeling, like you might pass out. Common side effects of isocarboxazid may include: headache, dizziness , drowsiness
- tremors

Drug Side Effects Summary

- sleep problems (insomnia)
- constipation , nausea
- or dry mouth.

1189. This particular disease, Depression, interacts with the drug levomilnacipran and has the following side effects:

- skin rash or hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tell your doctor right away if you have new or sudden changes in mood or behavior, including new or worse depression or anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, more active or talkative, or have thoughts about suicide or hurting yourself. Levomilnacipran may cause serious side effects. Call your doctor at once if you have: blurred vision, tunnel vision, eye pain or swelling, eye redness
- painful or difficult urination
- easy bruising, unusual bleeding, purple or red spots under your skin
- pounding heartbeats or fluttering in your chest
- seizure (convulsions)
- manic episodes--racing thoughts, increased energy, decreased need for sleep, risk-taking behavior, being agitated or talkative
- or low sodium level-- headache , confusion, slurred speech, severe weakness, vomiting , loss of coordination, feeling unsteady. Serious side effects may be more likely in older adults. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea . Common side effects of levomilnacipran may include: nausea, vomiting, constipation

Drug Side Effects Summary

- irregular heartbeats
- sweating
- or decreased sex drive, impotence , or difficulty having an orgasm.

1190. This particular disease, Depression, interacts with the drug Marplan and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, panic attacks, trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Stop taking Marplan and call your doctor at once if you have: sudden and severe headache , rapid heartbeat, stiffness in your neck, nausea , vomiting , cold sweat, vision problems, sensitivity to light
- chest pain, fast or slow heart rate
- swelling, rapid weight gain
- jaundice (yellowing of the skin or eyes)
- or a light-headed feeling, like you might pass out. Common side effects of Marplan may include: headache, dizziness , drowsiness
- tremors
- sleep problems (insomnia)
- constipation , nausea
- or dry mouth.

1191. This particular disease, Depression, interacts with the drug Tofranil and has the following side effects:

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Tofranil may cause serious side effects. Call your doctor at once if you have: easy bruising, unusual bleeding, purple or red spots under your skin
- tunnel vision, eye pain or swelling, or seeing halos around lights
- a light-headed feeling, like you might pass out
- new or worsening chest pain, pounding heartbeats or fluttering in your chest
- sudden numbness or weakness, problems with vision, speech, or balance
- fever, sore throat
- confusion, hallucinations, unusual thoughts or behavior
- painful or difficult urination
- seizure (convulsions)
- or jaundice (yellowing of the skin or eyes). Common side effects of Tofranil may include: increased blood pressure
- tingly feeling, weakness, lack of coordination
- dry mouth, nausea , vomiting
- or breast swelling (in men or women).

1192. This particular disease, Depression, interacts with the drug trimipramine and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your

Drug Side Effects Summary

doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Trimipramine may cause serious side effects. Call your doctor at once if you have: tunnel vision, eye pain or swelling, or seeing halos around lights

- a light-headed feeling, like you might pass out
- new or worsening chest pain, pounding heartbeats or fluttering in your chest
- confusion, unusual thoughts or behavior
- a seizure (convulsions)
- painful or difficult urination
- easy bruising, unusual bleeding
- restless muscle movements in your eyes, tongue, jaw, or neck
- or fever, chills, sore throat , mouth sores, feeling light-headed. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea . Common side effects of trimipramine may include: increased blood pressure
- numbness or tingling in your hands or feet
- dry mouth, nausea, vomiting
- rash, itching
- breast swelling (in men or women)
- or abnormal liver function tests .

1193. This particular disease, Depression, interacts with the drug Vivactil and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your

Drug Side Effects Summary

doctor immediately if any of the following side effects occur: Incidence not known Abdominal or stomach pain agitation or irritability blurred vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings change in urination chest pain or discomfort clay-colored stools cold sweats confusion about identity, place, and time continuing ringing, buzzing, or other unexplained noise in the ears convulsions cool, pale skin difficulty with speaking dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position double vision drooling false beliefs that cannot be changed by facts fast, pounding, or irregular heartbeat or pulse feeling of warmth feeling, seeing, or hearing things that are not there feeling that others are watching you or controlling your behavior feeling that others can hear your thoughts fever with or without chills hearing loss hostility hyperventilation inability to move the arms, legs, or facial muscles lack of coordination lower back or side pain mood or mental changes muscle spasm or jerking of all extremities muscle trembling or twitching nightmares pain or discomfort in the arms, jaw, back, or neck panic pinpoint red or purple spots on the skin pounding in the ears redness of the face, neck, arms, and occasionally, upper chest shakiness and unsteady walk slurred speech sore throat sores, ulcers, or white spots on the lips or in the mouth stiffness of the limbs stupor sweating swelling of face, ankles, or hands swollen glands talking, feeling, and acting with excitement trouble sleeping twisting movements of body uncontrolled movements, especially of the face, neck, and back unusual behavior unusual bleeding or bruising unusual tiredness or weakness weakness in the arms, hands, legs, or feet weight gain or loss yellow eyes or skin Symptoms of overdose Change in consciousness clumsiness disorientation disturbed concentration drowsiness to profound coma enlarged pupils increased or excessive unconscious or jerking movements low body temperature muscle aches or tightness muscle weakness severe sleepiness shivering trouble breathing weak or feeble pulse Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects.

Drug Side Effects Summary

Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Abdominal or stomach cramps bigger, dilated, or enlarged pupils (black part of the eye) black tongue difficulty having a bowel movement (stool) enlargement or swelling of the breasts hair loss or thinning of the hair heartburn hives or welts increased in sexual ability, desire, drive, or performance increased sensitivity of the eyes to light increased sensitivity of the skin to sunlight pain or discomfort in the chest, upper stomach, or throat peculiar taste severe sunburn small red or purple spots on the skin swelling of the testicles swelling or inflammation of the mouth swollen, painful, or tender lymph glands on side of the face or neck unexpected or excess milk flow from the breasts waking to urinate at night Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Vivactil side effects

1194. This particular disease, Depression, interacts with the drug XaQuil XR and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Less serious side effects may occur, and you may have none at all.

1195. This particular disease, Depression, interacts with the drug esketamine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Your blood pressure will need to be checked before and after you use esketamine. Esketamine can increase your blood pressure for several hours after each dose. Tell your doctor if you have chest pain, trouble breathing, severe headache , blurred

Drug Side Effects Summary

vision, pounding in your neck or ears, or a seizure. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Also call your doctor at once if you have: extreme drowsiness or feeling like you might pass out

- severe dizziness or feelings of floating
- problems with thinking or memory
- unusual or unpleasant memories (flashbacks)
- hallucinations, feeling "spaced out"
- or urination problems (painful urination, increased urination, urgent need to urinate). Common side effects of esketamine may include: dissociation, feeling drunk
- increased blood pressure
- drowsiness, lack of energy
- dizziness, spinning sensation
- feeling anxious
- nausea, vomiting
- or decreased sensations (touch or other senses).

1196. This particular disease, Depression, interacts with the drug L-Methylfolate Forte and has the t

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching

Drug Side Effects Summary

- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. L-Methylfolate Forte side effects

1197. This particular disease, Depression, interacts with the drug protriptyline and has the following

- hives , rash, red or purple spots
- fever
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Protriptyline may cause serious side effects. Call your doctor at once if you have: fever, chills, sore throat , mouth sores, feeling light-headed
- blurred vision, eye pain or redness, seeing halos around lights
- restless muscle movements in your eyes, tongue, jaw, or neck
- seizure (convulsions)
- new or worsening chest pain, pounding heartbeats or fluttering in your chest
- sudden numbness or weakness, problems with vision, or speech
- easy bruising, unusual bleeding
- painful or difficult urination
- or jaundice (yellowing of the skin or eyes). Common side effects of protriptyline may include: fast

Drug Side Effects Summary

heartbeats

- problems with balance or muscle movement
- constipation
- nausea , vomiting , loss of appetite
- breast swelling (in men or women)
- or decreased sex drive, impotence .

1198. This particular disease, Depression, interacts with the drug Spravato and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Your blood pressure will need to be checked before and after you use Spravato. Esketamine can increase your blood pressure for several hours after each dose. Tell your doctor if you have chest pain, trouble breathing, severe headache , blurred vision, pounding in your neck or ears, or a seizure. Also call your doctor at once if you have: extreme drowsiness or feeling like you might pass out
- severe dizziness or feelings of floating
- unusual changes in mood or behavior
- thoughts of hurting yourself
- worsening depression
- sleep problems
- problems with thinking or memory
- unusual or unpleasant memories (flashbacks)
- hallucinations, feeling "spaced out"
- or urination problems (painful urination, increased urination, urgent need to urinate). Common Spravato side effects may include: dissociation, feeling drunk

Drug Side Effects Summary

- increased blood pressure
- drowsiness
- dizziness, spinning sensation
- feeling anxious
- nausea, vomiting
- or decreased sensations (touch or other senses).

1199. This particular disease, Depression, interacts with the drug Surmontil and has the following s

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Abdominal or stomach pain agitation black, tarry stools bleeding or bruising blood in the urine or stools blurred vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain or discomfort clay-colored stools cold sweats confusion about identity, place, and time continuing ringing, buzzing, or other unexplained noise in the ears convulsions cool, pale skin cough or hoarseness dark urine decrease in the frequency of urination decreased urine output or volume depression difficulty in passing urine (dribbling) difficulty with speaking dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position double vision drooling dry mouth false beliefs that cannot be changed by facts fast, pounding, or irregular heartbeat or pulse feeling of warmth feeling that others are watching you or controlling your behavior feeling that others can hear your thoughts feeling, seeing, or hearing things that are not there fever with or without chills flushed, dry skin fruit-like breath odor general feeling of tiredness or weakness headache hearing loss hostility hyperventilation inability to move the arms, legs, or facial muscles inability to speak increased hunger increased need to urinate increased thirst increased urination irritability itching or rash lack of coordination lethargy loss of appetite loss of balance control lower back or

Drug Side Effects Summary

side pain mood or mental changes muscle spasm or jerking of all extremities muscle trembling or stiffness muscle twitching nausea nightmares pain or discomfort in the arms, jaw, back, or neck painful or difficult urination passing urine more often pinpoint red or purple spots on the skin pounding in the ears rapid weight gain redness of the face, neck, arms, and occasionally, upper chest restlessness shakiness and unsteady walk shuffling walk slow speech slurred speech sore throat sores, ulcers, or white spots on the lips or in the mouth stiffness of the limbs sudden loss of consciousness sweating swelling of the face, ankles, legs, or hands swollen glands talking, feeling, and acting with excitement trouble in holding or releasing urine trouble sleeping twisting movements of the body uncontrolled movements, especially of the face, neck, and back unpleasant breath odor unsteadiness, awkwardness, trembling, or other problems with muscle control or coordination unusual behavior unusual tiredness or weakness vomiting of blood weakness in the arms, hands, legs, or feet weight gain or loss yellow eyes or skin

Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them:

Incidence not known

Abdominal or stomach cramps bigger, dilated, or enlarged pupils (black part of the eye) black tongue bloating decreased interest in sexual intercourse diarrhea difficulty having a bowel movement (stool) disturbance of accommodation enlargement of the breast hair loss or thinning of the hair heartburn hives or welts inability to have or keep an erection increase in sexual ability, desire, drive, or performance increased sensitivity of the eyes to light increased sensitivity of the skin to sunlight loss in sexual ability, desire, drive, or performance pain or discomfort in the chest, upper stomach, or throat peculiar taste redness or other discoloration of the skin severe sunburn small red or purple spots on the skin swelling of the breasts or breast soreness in both women and men swelling of the testicles swelling or inflammation of the mouth swollen, painful, or tender lymph

Drug Side Effects Summary

glands in the neck unexpected milk flow from the breasts Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Surmontil side effects

1200. This particular disease, Depression, interacts with the drug thyroid desiccated and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Less serious side effects may include temporary hair loss (especially in children).

1201. This particular disease, Diabetes (Type 1), interacts with the drug Humalog and has the following side effects:

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. Call your doctor at once if you have: weight gain, swelling in your hands or feet, feeling short of breath
- low blood sugar - headache, hunger, sweating, irritability, dizziness, fast heart rate, and feeling anxious or shaky
- or low potassium - leg cramps, constipation, irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common Humalog side effects may include: low blood sugar
- weight gain
- swelling in your hands or feet
- itching
- or thickening or hollowing of the skin where you injected the medicine.

Drug Side Effects Summary

1202. This particular disease, Diabetes (Type 1), interacts with the drug Lantus and has the following

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. Call your doctor at once if you have: rapid weight gain, swelling in your feet or ankles
- shortness of breath
- or low blood potassium - leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common Lantus side effects may include: low blood sugar
- swelling, weight gain
- allergic reaction, itching, rash
- or thickening or hollowing of the skin where you injected the medicine.

1203. This particular disease, Diabetes (Type 1), interacts with the drug Novolog and has the following

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. Call your doctor at once if you have: heart problems - swelling, rapid weight gain, feeling short of breath
- or low potassium - leg cramps, constipation, irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common NovoLog side effects may include: low blood sugar
- weight gain
- low potassium
- swelling in your hands and feet
- skin rash, itching, redness, or swelling
- or thickening or hollowing of the skin where you injected the medicine.

Drug Side Effects Summary

1204. This particular disease, Diabetes (Type 1), interacts with the drug insulin lispro and has the following side effects:

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. Call your doctor at once if you have: weight gain, swelling in your hands or feet, feeling short of breath
- low blood sugar - headache , hunger, sweating, irritability, dizziness , fast heart rate , and feeling anxious or shaky
- or low potassium - leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common Insulin lispro side effects may include: low blood sugar
- weight gain
- swelling in your hands or feet
- itching
- or thickening or hollowing of the skin where you injected the medicine.

1205. This particular disease, Diabetes (Type 1), interacts with the drug insulin aspart and has the following side effects:

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. Insulin aspart may cause serious side effects. Call your doctor at once if you have: heart problems--swelling, rapid weight gain, feeling short of breath
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of insulin aspart may include: low blood sugar
- weight gain
- low potassium
- swelling in your hands and feet

Drug Side Effects Summary

- skin rash , itching, redness, or swelling
- or thickening or hollowing of the skin where you injected the medicine.

1206. This particular disease, Diabetes (Type 1), interacts with the drug Lantus SoloStar and has the following side effects:

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. Lantus Solostar Pen may cause serious side effects. Call your doctor at once if you have: rapid weight gain, swelling in your feet or ankles
- shortness of breath
- or low blood potassium--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of Lantus Solostar Pen may include: low blood sugar
- swelling, weight gain
- allergic reaction, itching, rash
- or thickening or hollowing of the skin where you injected the medicine.

1207. This particular disease, Diabetes (Type 1), interacts with the drug Lyumjev and has the following side effects:

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. Call your doctor at once if you have: weight gain, swelling in your hands or feet, feeling short of breath
- low blood sugar - headache , hunger, sweating, irritability, dizziness , fast heart rate , and feeling anxious or shaky
- or low potassium - leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common Lyumjev side effects may include: low blood sugar (hypoglycemia)

Drug Side Effects Summary

- allergic reactions
- rash. weight gain
- swelling in your hands or feet
- itching (pruritus)
- or thickening or hollowing of the skin where you injected the medicine.

1208. This particular disease, Diabetes (Type 1), interacts with the drug insulin detemir and has the

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. Call your doctor at once if you have: fluid retention - weight gain, swelling in your hands or feet, feeling short of breath
- or low potassium - leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common insulin detemir side effects may include: low blood sugar
- weight gain
- swelling in your hands and feet
- rash, itching
- or thickening or hollowing of the skin where you injected the medicine.

1209. This particular disease, Diabetes (Type 1), interacts with the drug Levemir and has the followi

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. Call your doctor at once if you have: fluid retention - weight gain, swelling in your hands or feet, feeling short of breath
- or low potassium - leg cramps, constipation , irregular heartbeats, fluttering in your chest,

Drug Side Effects Summary

increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common Levemir side effects include: low blood sugar

- weight gain
- swelling in your hands and feet
- rash, itching
- or thickening or hollowing of the skin where you injected the medicine.

1210. This particular disease, Diabetes (Type 1), interacts with the drug insulin aspart/insulin aspart

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, wheezing, rapid pulse, sweating, feeling like you might pass out, or swelling in your tongue or throat. Insulin aspart and insulin aspart protamine may cause serious side effects. Call your doctor at once if you have: fluid retention --weight gain, swelling in your hands or feet, feeling short of breath
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of insulin aspart and insulin aspart protamine may include: low potassium
- itching, mild skin rash
- or thickening or hollowing of the skin where you injected the medicine.

1211. This particular disease, Diabetes (Type 1), interacts with the drug Humalog KwikPen and has

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives

Drug Side Effects Summary

- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of low potassium levels like muscle pain or weakness, muscle cramps, or a heartbeat that does not feel normal. Signs of infection like fever, chills, very bad sore throat , ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal. Change in eyesight. Chills. Very bad dizziness or passing out. Mood changes. Seizures. Slurred speech. Shortness of breath, a big weight gain, or swelling in the arms or legs. Thick skin, pits, or lumps where the injection was given. Low blood sugar may occur. Signs may be dizziness, headache , feeling sleepy, feeling weak, shaking, a fast heartbeat , confusion, hunger, or sweating. Call the doctor right away if any of these signs occur. Follow what you have been told to do if low blood sugar occurs. This may include taking glucose tablets, liquid glucose, or some fruit juices. Humalog KwikPen side effects

1212. This particular disease, Diabetes (Type 1), interacts with the drug NovoLog FlexPen and has t

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. NovoLog FlexPen may cause serious side effects. Call your doctor at once if you have: heart problems--swelling, rapid weight gain, feeling short of breath
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of NovoLog FlexPen may include: low blood sugar

Drug Side Effects Summary

- weight gain
- low potassium
- swelling in your hands and feet
- skin rash , itching, redness, or swelling
- or thickening or hollowing of the skin where you injected the medicine.

1213. This particular disease, Diabetes (Type 1), interacts with the drug Apidra and has the following

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. Call your doctor at once if you have: fluid retention - weight gain, swelling in your hands or feet, feeling short of breath
- or low potassium - leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common Apidra side effects include: low blood sugar
- itching, mild skin rash
- or thickening or hollowing of the skin where you injected the medicine.

1214. This particular disease, Diabetes (Type 1), interacts with the drug insulin glargine and has the

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. Insulin glargine may cause serious side effects. Call your doctor at once if you have: rapid weight gain, swelling in your feet or ankles
- shortness of breath
- or low blood potassium--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side

Drug Side Effects Summary

effects of insulin glargine may include: low blood sugar

- swelling, weight gain
- allergic reaction, itching, rash
- or thickening or hollowing of the skin where you injected the medicine.

1215. This particular disease, Diabetes (Type 1), interacts with the drug Basaglar and has the following

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. Call your doctor at once if you have: rapid weight gain, swelling in your feet or ankles

- shortness of breath
- or low blood potassium - leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common Basaglar side effects may include: low blood sugar

- swelling, weight gain
- allergic reaction, itching, rash
- or thickening or hollowing of the skin where you injected the medicine.

1216. This particular disease, Diabetes (Type 1), interacts with the drug Tresiba and has the following

- hives , itching, skin rash
- wheezing, tiredness, trouble breathing
- feeling like you might pass out
- nausea , diarrhea
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: fluid retention - weight gain, swelling in your hands or feet, feeling short of breath
- or low potassium - leg cramps, constipation , irregular heartbeats, fluttering in your chest,

Drug Side Effects Summary

increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common Tresiba side effects may include: low blood sugar

- swelling, weight gain
- itching, rash
- or thickening or hollowing of the skin where you injected the medicine.

1217. This particular disease, Diabetes (Type 1), interacts with the drug Toujeo SoloStar and has the following side effects:

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. Call your doctor at once if you have: rapid weight gain, swelling in your feet or ankles

- shortness of breath
- or low blood potassium - leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common Toujeo side effects may include: low blood sugar

- swelling, weight gain
- allergic reaction, itching, rash
- or thickening or hollowing of the skin where you injected the medicine.

1218. This particular disease, Diabetes (Type 1), interacts with the drug Humulin N and has the following side effects:

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, chest tightness, feeling like you might pass out, or swelling in your tongue or throat. Call your doctor at once if you have: fluid retention - weight gain, swelling in your hands or feet, feeling short of breath

- or low potassium - leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common

Drug Side Effects Summary

Humulin N side effects may include: low blood sugar

- weight gain, swelling in your hands or feet
- itching, mild skin rash
- or thickening or hollowing of the skin where you injected the medicine.

1219. This particular disease, Diabetes (Type 1), interacts with the drug Humulin R and has the following side effects:

- redness, swelling, sweating, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. Call your doctor at once if you have: low potassium - leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common Humulin R side effects may include: low blood sugar

- swelling in your hands or feet
- weight gain
- or thickening or hollowing of the skin where you injected the medicine.

1220. This particular disease, Diabetes (Type 1), interacts with the drug Afrezza and has the following side effects:

- Afrezza may cause serious side effects that can lead to death, including: See Important information Low blood sugar (hypoglycemia). Signs and symptoms that may indicate low blood sugar include: dizziness or light-headedness, sweating, confusion, headache, blurred vision, slurred speech, shakiness, fast heartbeat, anxiety, irritability or mood change, hunger. Decreased lung function. Your healthcare provider should check how your lungs are working before you start using Afrezza, 6 months after you start using it and yearly after that. Lung cancer. In studies in people with diabetes, lung cancer occurred in a few more people who were taking Afrezza than in people who were taking other diabetes medications. There were too few cases to know if lung cancer was related to Afrezza. If you have lung cancer, you and your healthcare provider should decide if you

Drug Side Effects Summary

should use this medicine. Diabetic ketoacidosis. Talk to your healthcare provider if you have an illness. Your Afrezza dose or how often you check your blood sugar may need to be changed. Severe allergic reaction (whole body reaction). Get medical help right away if you have any of these signs or symptoms of a severe allergic reaction: A rash over your whole body, trouble breathing, a fast heartbeat, or sweating. Low potassium in your blood (hypokalemia). Heart failure. Taking certain diabetes pills called thiazolidinediones or "TZDs" with Afrezza may cause heart failure in some people. This can happen even if you have never had heart failure or heart problems before. If you already have heart failure it may get worse while you take TZDs with Afrezza. Your healthcare provider should monitor you closely while you are taking TZDs with this medicine. Tell your healthcare provider if you have any new or worse symptoms of heart failure including: Shortness of breath, swelling of your ankles or feet, sudden weight gain. Treatment with TZDs and Afrezza may need to be changed or stopped by your healthcare provider if you have new or worse heart failure. Trouble breathing, shortness of breath, fast heartbeat, swelling of your face, tongue, or throat, sweating, extreme drowsiness, dizziness, confusion. The most common side effects include: Low blood sugar (hypoglycemia), cough, sore throat These are not all the possible side effects of Afrezza. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 (1-800-332-1088). Afrezza side effects

1221. This particular disease, Diabetes (Type 1), interacts with the drug Novolin R and has the following

- redness, swelling, sweating, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. Novolin R may cause serious side effects. Call your doctor at once if you have: low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of Novolin R may include: low blood sugar
- swelling in your hands or feet

Drug Side Effects Summary

- weight gain
- or thickening or hollowing of the skin where you injected the medicine.

1222. This particular disease, Diabetes (Type 1), interacts with the drug Symlin and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe ongoing nausea
- or severe hypoglycemia - headache, dizziness, drowsiness, vision problems, hunger, weakness, sweating, confusion, irritability, fast heart rate, feeling jittery. Common Symlin side effects may include: nausea, vomiting, loss of appetite
- or headache.

1223. This particular disease, Diabetes (Type 1), interacts with the drug Humulin 70/30 and has the following

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, wheezing, rapid pulse, sweating, feeling like you might pass out, or swelling in your tongue or throat. Call your doctor at once if you have: fluid retention - weight gain, swelling in your hands or feet, feeling short of breath
- or low potassium - leg cramps, constipation, irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common Humulin 70/30 side effects may include: low blood sugar
- itching, mild skin rash
- or thickening or hollowing of the skin where you injected the medicine.

1224. This particular disease, Diabetes (Type 1), interacts with the drug insulin regular and has the following

Drug Side Effects Summary

- redness, swelling, sweating, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. Call your doctor at once if you have: low potassium - leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common insulin side effects may include: low blood sugar
- swelling in your hands or feet
- weight gain
- or thickening or hollowing of the skin where you injected the medicine.

1225. This particular disease, Diabetes (Type 1), interacts with the drug Novolin N and has the following side effects:

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, chest tightness, feeling like you might pass out, or swelling in your tongue or throat. Novolin N may cause serious side effects. Call your doctor at once if you have: fluid retention --weight gain, swelling in your hands or feet, feeling short of breath
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of Novolin N may include: low blood sugar
- weight gain, swelling in your hands or feet
- itching, mild skin rash
- or thickening or hollowing of the skin where you injected the medicine.

1226. This particular disease, Diabetes (Type 1), interacts with the drug Admelog and has the following side effects:

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. Call your doctor at once if you have: weight gain, swelling in your hands or feet, feeling short of breath

Drug Side Effects Summary

- low blood sugar-- headache , hunger, sweating, irritability, dizziness , fast heart rate , and feeling anxious or shaky
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common Admelog side effects may include: low blood sugar
- weight gain
- swelling in your hands or feet
- itching
- or thickening or hollowing of the skin where you injected the medicine.

1227. This particular disease, Diabetes (Type 1), interacts with the drug Fiasp and has the following

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. Fiasp may cause serious side effects. Call your doctor at once if you have: heart problems--swelling, rapid weight gain, feeling short of breath
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of Fiasp may include: low blood sugar
- weight gain
- low potassium
- swelling in your hands and feet
- skin rash , itching, redness, or swelling
- or thickening or hollowing of the skin where you injected the medicine.

1228. This particular disease, Diabetes (Type 1), interacts with the drug Novolin 70/30 and has the f

Drug Side Effects Summary

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, chest tightness, feeling like you might pass out, or swelling in your tongue or throat. Novolin 70/30 may cause serious side effects. Call your doctor at once if you have: fluid retention --weight gain, swelling in your hands or feet, feeling short of breath
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of Novolin 70/30 may include: low blood sugar
- weight gain, swelling in your hands or feet
- itching, mild skin rash
- or thickening or hollowing of the skin where you injected the medicine.

1229. This particular disease, Diabetes (Type 1), interacts with the drug Humalog Mix 75/25 and has the

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. Call your doctor at once if you have: fluid retention - weight gain, swelling in your hands or feet, feeling short of breath
- or low potassium - leg cramps, constipation, irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common Humalog Mix side effects may include: low blood sugar
- itching, mild skin rash
- or thickening or hollowing of the skin where you injected the medicine.

1230. This particular disease, Diabetes (Type 1), interacts with the drug insulin glulisine and has the

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat.

Drug Side Effects Summary

Insulin glulisine may cause serious side effects. Call your doctor at once if you have: fluid retention

--weight gain, swelling in your hands or feet, feeling short of breath

- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of insulin glulisine may include: low blood sugar

- itching, mild skin rash

- or thickening or hollowing of the skin where you injected the medicine.

1231. This particular disease, Diabetes (Type 1), interacts with the drug Rezvoglar and has the follo

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Blindness blurred vision cough or hoarseness decreased vision diarrhea dizziness fever or chills general feeling of discomfort or illness headache joint pain lower back or side pain loss of appetite muscle aches and pains nausea nervousness painful or difficult urination pounding in the ears runny nose shivering slow or fast heartbeat sore throat sweating swelling of the hands, ankles, feet, or lower legs trouble sleeping unusual tiredness or weakness vomiting Incidence not known Anxiety chest tightness cold sweats coma confusion cool, pale skin decreased urine depression depression or thickening of the skin at the injection site difficulty swallowing dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position dry mouth fast heartbeat hives, itching, skin rash increased hunger increased thirst itching, pain, redness, or swelling at the injection site large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs mood changes muscle cramps nightmares numbness or tingling in the hands, feet, or lips puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue seizures shakiness slurred speech trouble breathing weight gain Some side effects may occur that usually do not need medical

Drug Side Effects Summary

attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Body aches or pain diarrhea difficulty in moving discouragement ear congestion feeling sad or empty irritability lack of appetite loss of interest or pleasure loss of voice muscle pain or stiffness pain in the arms or legs pain in the joints pain or tenderness around the eyes and cheekbones sneezing stuffy or runny nose tender, swollen glands in the neck trouble concentrating voice changes Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Rezvoglar side effects

1232. This particular disease, Diabetes (Type 1), interacts with the drug Semglee and has the following

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. Call your doctor at once if you have: rapid weight gain, swelling in your feet or ankles
- shortness of breath
- or low blood potassium - leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common Semglee side effects may include: low blood sugar
- swelling, weight gain
- allergic reaction, itching, rash
- or thickening or hollowing of the skin where you injected the medicine.

1233. This particular disease, Diabetes (Type 1), interacts with the drug Toujeo Max SoloStar and has

Drug Side Effects Summary

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

- itching

- red, swollen, blistered, or peeling skin with or without fever

- wheezing

- tightness in the chest or throat

- trouble breathing, swallowing, or talking

- unusual hoarseness

- or swelling of the mouth, face, lips, tongue, or throat. Signs of low potassium levels like muscle pain or weakness, muscle cramps, or a heartbeat that does not feel normal. Signs of infection like fever, chills, very bad sore throat, ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal. Change in eyesight. Very bad dizziness or passing out. Seizures. Thick skin, pits, or lumps where the injection was given. Shortness of breath, a big weight gain, or swelling in the arms or legs. Low mood (depression). Low blood sugar may occur. Signs may be dizziness, headache, feeling sleepy, feeling weak, shaking, a fast heartbeat, confusion, hunger, or sweating. Call the doctor right away if any of these signs occur. Follow what you have been told to do if low blood sugar occurs. This may include taking glucose tablets, liquid glucose, or some fruit juices. Toujeo Max SoloStar side effects

1234. This particular disease, Diabetes (Type 1), interacts with the drug Humulin R U-500 (Concentr

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat.

Drug Side Effects Summary

Humulin R (Concentrated) may cause serious side effects. Call your doctor at once if you have: fluid retention --weight gain, swelling in your hands or feet, feeling short of breath

- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of Humulin R (Concentrated) may include: low blood sugar
- itching, mild skin rash
- or thickening or hollowing of the skin where you injected the medicine.

1235. This particular disease, Diabetes (Type 1), interacts with the drug NovoLog Mix 70/30 and has the following side effects:

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, wheezing, rapid pulse, sweating, feeling like you might pass out, or swelling in your tongue or throat. NovoLog Mix 70/30 may cause serious side effects. Call your doctor at once if you have: fluid retention --weight gain, swelling in your hands or feet, feeling short of breath
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of NovoLog Mix 70/30 may include: low potassium
- itching, mild skin rash
- or thickening or hollowing of the skin where you injected the medicine.

1236. This particular disease, Diabetes (Type 1), interacts with the drug NovoLog PenFill and has the following side effects:

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. NovoLog PenFill may cause serious side effects. Call your doctor at once if you have: heart problems--swelling, rapid weight gain, feeling short of breath
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest,

Drug Side Effects Summary

increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of NovoLog PenFill may include: low blood sugar

- weight gain
- low potassium
- swelling in your hands and feet
- skin rash , itching, redness, or swelling
- or thickening or hollowing of the skin where you injected the medicine.

1237. This particular disease, Diabetes (Type 1), interacts with the drug octreotide and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Octreotide may cause serious side effects. Call your doctor at once if you have: severe constipation
- slow or uneven heartbeats
- signs of gallstones --fever, chills, nausea , vomiting , severe pain in your upper stomach spreading to your back, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- low blood sugar -- headache , hunger, sweating, irritability, dizziness , fast heart rate , and feeling anxious or shaky
- or underactive thyroid --extreme tired feeling, dry skin , joint pain or stiffness, muscle pain or weakness, hoarse voice, feeling more sensitive to cold temperatures, weight gain. Common side effects of octreotide may include: gallstones
- nausea, vomiting, diarrhea, stomach pain, gas
- headache, back pain
- or dizziness, tiredness.

Drug Side Effects Summary

1238. This particular disease, Diabetes (Type 1), interacts with the drug pramlintide and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Pramlintide may cause serious side effects. Call your doctor at once if you have: severe ongoing nausea
- or severe hypoglycemia--headache, dizziness, drowsiness, vision problems, hunger, weakness, sweating, confusion, irritability, fast heart rate, feeling jittery. Common side effects of pramlintide may include: nausea, vomiting, loss of appetite
- or headache.

1239. This particular disease, Diabetes (Type 1), interacts with the drug insulin inhalation, rapid acting and has the following side effects:

- redness or swelling where an insulin injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, sweating, feeling like you might pass out, or swelling in your tongue or throat. Insulin may cause serious side effects. Call your doctor at once if you have: bronchospasm (wheezing, chest tightness, trouble breathing)
- fluid retention --weight gain, swelling in your hands or feet, feeling short of breath
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of insulin may include: low blood sugar
- cough
- or sore throat .

1240. This particular disease, Diabetes (Type 1), interacts with the drug insulin isophane and has the following side effects:

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, chest tightness, feeling like you might pass out, or swelling in your tongue or throat.

Drug Side Effects Summary

Insulin isophane may cause serious side effects. Call your doctor at once if you have: fluid retention

--weight gain, swelling in your hands or feet, feeling short of breath

- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of insulin isophane may include: low blood sugar

- weight gain, swelling in your hands or feet

- itching, mild skin rash

- or thickening or hollowing of the skin where you injected the medicine.

1241. This particular disease, Diabetes (Type 1), interacts with the drug NovoLog Mix 70/30 FlexPen

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, wheezing, rapid pulse, sweating, feeling like you might pass out, or swelling in your tongue or throat. NovoLog Mix 70/30 FlexPen may cause serious side effects. Call your doctor at once if you have: fluid retention --weight gain, swelling in your hands or feet, feeling short of breath

- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of NovoLog Mix 70/30 FlexPen may include: low potassium

- itching, mild skin rash

- or thickening or hollowing of the skin where you injected the medicine.

1242. This particular disease, Diabetes (Type 1), interacts with the drug Humulin N Pen and has the

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, chest tightness, feeling like you might pass out, or swelling in your tongue or throat.

HumuLIN N KwikPen may cause serious side effects. Call your doctor at once if you have: fluid retention --weight gain, swelling in your hands or feet, feeling short of breath

Drug Side Effects Summary

- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of HumuLIN N KwikPen may include: low blood sugar
- weight gain, swelling in your hands or feet
- itching, mild skin rash
- or thickening or hollowing of the skin where you injected the medicine.

1243. This particular disease, Diabetes (Type 1), interacts with the drug Humulin R U-500 KwikPen a

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. HumuLIN R KwikPen (Concentrated) may cause serious side effects. Call your doctor at once if you have: fluid retention --weight gain, swelling in your hands or feet, feeling short of breath
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of HumuLIN R KwikPen (Concentrated) may include: low blood sugar
- itching, mild skin rash
- or thickening or hollowing of the skin where you injected the medicine.

1244. This particular disease, Diabetes (Type 1), interacts with the drug insulin degludec and has th

- Insulin degludec may cause serious side effects that can lead to death, including: Low blood sugar (hypoglycemia). Signs and symptoms that may indicate low blood sugar include: dizziness or light-headedness blurred vision anxiety, irritability, or mood changes sweating slurred speech hunger confusion shakiness headache fast heartbeat Low potassium in your blood (hypokalemia). Heart failure. Taking certain diabetes pills called thiazolidinediones or "TZDs" with insulin degludec may cause heart failure in some people. This can happen even if you have never had heart failure

Drug Side Effects Summary

or heart problems before. If you already have heart failure, it may get worse while you take TZDs with insulin degludec. Your healthcare provider should monitor you closely while you are taking TZDs with insulin degludec. Tell your healthcare provider if you have any new or worse symptoms of heart failure including shortness of breath, tiredness, swelling of your ankles or feet and sudden weight gain. Treatment with TZDs and insulin degludec may need to be adjusted or stopped by your healthcare provider if you have new or worse heart failure. Your insulin dose may need to change because of: change in level of physical activity or exercise weight gain or loss increased stress illness change in diet Common side effects of insulin degludec may include: serious allergic reactions (whole body reactions) reactions at the injection site skin thickening or pits at the injection site (lipodystrophy) itching rash swelling of your hands and feet weight gain trouble breathing shortness of breath fast heartbeat swelling of your face, tongue, or throat sweating extreme drowsiness dizziness confusion These are not all the possible side effects of insulin degludec. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Insulin degludec side effects

1245. This particular disease, Diabetes (Type 1), interacts with the drug insulin isophane / insulin regular

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, chest tightness, feeling like you might pass out, or swelling in your tongue or throat. Insulin isophane and insulin regular may cause serious side effects. Call your doctor at once if you have: fluid retention --weight gain, swelling in your hands or feet, feeling short of breath
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of insulin isophane and insulin regular may include: low blood sugar
- weight gain, swelling in your hands or feet
- itching, mild skin rash

Drug Side Effects Summary

- or thickening or hollowing of the skin where you injected the medicine.

1246. This particular disease, Diabetes (Type 1), interacts with the drug Myxredlin and has the following side effects:

- redness, swelling, sweating, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. Myxredlin may cause serious side effects. Call your doctor at once if you have: low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of Myxredlin may include: low blood sugar
- swelling in your hands or feet
- weight gain
- or thickening or hollowing of the skin where you injected the medicine.

1247. This particular disease, Diabetes (Type 1), interacts with the drug Humalog Mix 50/50 and has the following side effects:

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. Humalog Mix 50/50 may cause serious side effects. Call your doctor at once if you have: fluid retention --weight gain, swelling in your hands or feet, feeling short of breath
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of Humalog Mix 50/50 may include: low blood sugar
- itching, mild skin rash
- or thickening or hollowing of the skin where you injected the medicine.

1248. This particular disease, Diabetes (Type 1), interacts with the drug Humalog Mix 75/25 KwikPen and has the following side effects:

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble

Drug Side Effects Summary

breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat.

Humalog Mix 75/25 KwikPen may cause serious side effects. Call your doctor at once if you have:

fluid retention --weight gain, swelling in your hands or feet, feeling short of breath

- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of Humalog Mix 75/25 KwikPen may include: low blood sugar

- itching, mild skin rash

- or thickening or hollowing of the skin where you injected the medicine.

1249. This particular disease, Diabetes (Type 1), interacts with the drug insulin lispro/insulin lispro

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat.

Insulin lispro and insulin lispro protamine may cause serious side effects. Call your doctor at once if you have: fluid retention --weight gain, swelling in your hands or feet, feeling short of breath

- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of insulin lispro and insulin lispro protamine may include: low blood sugar

- itching, mild skin rash

- or thickening or hollowing of the skin where you injected the medicine.

1250. This particular disease, Diabetes (Type 1), interacts with the drug Symlin Pen and has the foll

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Anxiety blurred vision chills cold sweats coma confusion cool pale skin cough depression difficulty with swallowing

Drug Side Effects Summary

dizziness fast heartbeat headache hives, itching, or skin rash increased hunger nausea nightmares puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue seizures shakiness slurred speech tightness in the chest unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Difficulty with moving inflicted injury loss of appetite muscle pain or stiffness pain in the joints stomach pain vomiting weight loss Less common Body aches or pain congestion dryness or soreness of the throat fever hoarseness runny nose tender, swollen glands in the neck voice changes Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Symlin Pen side effects

1251. This particular disease, Diabetes (Type 1), interacts with the drug SymlinPen 60 and has the f

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. SymlinPen 60 may cause serious side effects. Call your doctor at once if you have: severe ongoing nausea
- or severe hypoglycemia--headache, dizziness, drowsiness, vision problems, hunger, weakness, sweating, confusion, irritability, fast heart rate, feeling jittery. Common side effects of SymlinPen 60 may include: nausea, vomiting, loss of appetite
- or headache.

1252. This particular disease, Diabetes (Type 1), interacts with the drug SymlinPen 120 and has the

Drug Side Effects Summary

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. SymlinPen 120 may cause serious side effects. Call your doctor at once if you have: severe ongoing nausea
- or severe hypoglycemia--headache, dizziness, drowsiness, vision problems, hunger, weakness, sweating, confusion, irritability, fast heart rate, feeling jittery. Common side effects of SymlinPen 120 may include: nausea, vomiting, loss of appetite
- or headache.

1253. This particular disease, Diabetes (Type 1), interacts with the drug Humalog Mix 50/50 KwikPen

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, fast heartbeats, feeling like you might pass out, or swelling in your tongue or throat. Humalog Mix 50/50 KwikPen may cause serious side effects. Call your doctor at once if you have: fluid retention --weight gain, swelling in your hands or feet, feeling short of breath
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of Humalog Mix 50/50 KwikPen may include: low blood sugar
- itching, mild skin rash
- or thickening or hollowing of the skin where you injected the medicine.

1254. This particular disease, Diabetes (Type 1), interacts with the drug Humulin 50/50 and has the t

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Anxiety bloating or swelling of the face, arms, hands, lower legs, or feet blurred vision chest tightness chills cold sweats

Drug Side Effects Summary

coma confusion cool, pale skin cough decreased urine depression difficulty swallowing dizziness dry mouth fast heartbeat headache hives, itching, or rash increased hunger increased thirst irregular heartbeat loss of appetite mood changes muscle pain or cramps nausea or vomiting nightmares numbness or tingling in the hands, feet, or lips puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rapid weight gain seizures shakiness slurred speech tingling of the hands or feet unusual tiredness or weakness unusual weight gain or loss Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Weight gain redness, swelling, or itching at the injection site Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Humulin 50/50 side effects

1255. This particular disease, Diabetes (Type 1), interacts with the drug Humulin 70/30 Pen and has

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, chest tightness, feeling like you might pass out, or swelling in your tongue or throat. HumuLIN 70/30 KwikPen may cause serious side effects. Call your doctor at once if you have: fluid retention --weight gain, swelling in your hands or feet, feeling short of breath
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of HumuLIN 70/30 KwikPen may include: low blood sugar
- weight gain, swelling in your hands or feet
- itching, mild skin rash

Drug Side Effects Summary

- or thickening or hollowing of the skin where you injected the medicine.

1256. This particular disease, Diabetes (Type 1), interacts with the drug ReliOn/Novolin 70/30 and has the following side effects:

- redness or swelling where an injection was given, itchy skin rash over the entire body, trouble breathing, chest tightness, feeling like you might pass out, or swelling in your tongue or throat. ReliOn/Novolin 70/30 may cause serious side effects. Call your doctor at once if you have: fluid retention --weight gain, swelling in your hands or feet, feeling short of breath
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of ReliOn/Novolin 70/30 may include: low blood sugar
- weight gain, swelling in your hands or feet
- itching, mild skin rash
- or thickening or hollowing of the skin where you injected the medicine.

1257. This particular disease, Diabetes (Type 1), interacts with the drug Tzield and has the following side effects:

- Tzield may cause serious side effects: See Important information. The most common side effects include: rash leukopenia (decrease in white blood cell counts) headache These are not all of the possible side effects. For more information, ask your healthcare provider or pharmacist. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. You may also report side effects to Provention Bio at 1-844-778-2246. General information about the safe and effective use of Tzield. Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. You can ask your pharmacist or healthcare provider for information that is written for health professionals. What are the ingredients in Tzield? Active ingredient: teplizumab-mzwv. Inactive ingredients: dibasic sodium phosphate, monobasic sodium phosphate, polysorbate 80, sodium chloride, and water for injection. References Tzield

Drug Side Effects Summary

Product Label More about Tzield (teplizumab) Drug images FDA approval history Related treatment guides Diabetes, Type 1 Further information Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances. Medical Disclaimer Drug Status Availability Prescription only Rx User Reviews & Ratings Review this drug Related Stories News FDA Approves Tzield (teplizumab-mzwv) to Delay the Onset of Stage 3 Type 1 Diabetes Images Tzield 2 mg/2 mL (1 mg/mL) injection (medicine) View larger images Drugs.com Mobile Apps The easiest way to lookup drug information, identify pills, check interactions and set up your own personal medication records. Available for Android and iOS devices. Explore Apps Support Help Center Sitemap Contact Us About About Drugs.com Advertising Policy Attribution & Citations Terms & Privacy Editorial Policy Privacy Policy Terms of Use Follow Drugs.com on Facebook Follow Drugs.com on Twitter Follow Drugs.com on Pinterest Follow Drugs.com on YouTube Subscribe to Drugs.com newsletters Subscribe to Drugs.com newsletters for the latest medication news, new drug approvals, alerts and updates. Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include IBM Watson Micromedex (updated 2 Dec 2022), Cerner Multum? (updated 23 Nov 2022), ASHP (updated 11 Nov 2022) and others. Ad Choices We comply with the HONcode standard for trustworthy health information. Copyright © 2000-2022 Drugs.com. All rights reserved. SITECONFIG = {"agent":{"isMobile":false},"user":{"userid":0,"modal":{"login":"/api/user/modal/login/"},"consent":{"skip":true,"api":{"url":"/api/user/"}}, "page":{"section":"rootdocs","subsection":"","doc":"content","delayedC ss":["/bundle/css/component/component-share.min.9d6804aa.css"],"delayedScripts":[],"image":{"lazyLoad":{"selector":["data-lazy-load"],"screens":1,"delay":2000},"mailto":{"delay":5000,"location":"/support/email-mailto.html"}}, "form":{"recaptcha":{"sitekey":"6Ld-iSYTAAAAA00VTEwVI71HtvrFoSIP SibGr4g"},"callback":"DDC_RECAPTCHA_CALLBACK"},"dwLog":{"a":4412,"b":20196,"c":139,"d":

```
47,345","e":"","f":1228,"h":0,"i":"default","k":"139","l":"","m":"","n":"","o":"","p":0,"q":"drugs/rootdocs"},"
drugs":[{"ddc_id":4412,"brand_name_id":20196,"mtm_id":"","generic_name":"teplizumab
systemic","name":"Tzield"}],"options":{"recordSelection":true,"classifyLists":{"wrapper":"#content","co
untHtmlItems":false,"numItems":{"short":15,"medium":30,"long":60},"charLimit":{"paragraph":30,"cont
ent":350}}},"subscribe":null,"modal":null,"dictionary":{},"share":{"sticky":{"target":".ddc-social-share"},"
types":[{"title":"Facebook","label":"Share on Facebook","cssClass":"ddc-share-icon
ddc-share-icon-facebook","labelClass":"ddc-share-icon-label","icon":"<svg
ddc-icon-facebook' width='20' height='20' viewBox='0 0 24 24' aria-hidden='true' focusable='false'
xmlns='http://www.w3.org/2000/svg'><path d='M14 6.54c-2.06 0-3.4 1.25-3.4
3.52v2H8.3v2.61h2.3V21A9.04 9.04 0 0 1 12.05C3 7.05 7.03 3 12 3s9 4.05 9 9.05c0 4.53-3.3
8.27-7.6 8.95v-6.33h2.1l.4-2.62h-2.5v-1.7c0-.7.35-1.4 1.47-1.4h1.14V6.7s-1.03-.17-2.02-.17z'
/></svg>","shareUrl":"https://www.facebook.com/sharer/sharer.php?u=SHAREURL","width":600,"hei
ght":500},{"title":"Twitter","label":"Share on Twitter","cssClass":"ddc-share-icon
ddc-share-icon-twitter","labelClass":"ddc-share-icon-label","icon":"<svg
ddc-icon-twitter' width='20' height='20' viewBox='0 0 24 24' aria-hidden='true' focusable='false'
xmlns='http://www.w3.org/2000/svg'><path d='M8.66 19.63a10.44 10.44 0 0 1 0.5-10.5v-.5A7.7 7.7
0 0 2 1 6.74c-.67.3-1.39.5-2.12.58a3.73 3.73 0 0 1 .64-2.04c-.72.43-1.52.74-2.34.9a3.7 3.7 0 0 0-.6.3
3.36 10.46 10.46 0 0 1 -7.61-3.85 3.69 3.69 0 0 1 .14 4.9 3.66 3.66 0 0 1 -1.63-.45v.04a3.7 3.7 0 0 2 .9
3.64 3.7 3.7 0 0 1 -1.63.06 3.69 3.69 0 0 3 .42 2.53 7.4 7.4 0 0 1 -4.6 1.64 8 8 0 0 1 -.87-.07 10.45
10.45 0 0 5.66 1.63'
/></svg>","shareUrl":"https://twitter.com/intent/tweet?text=SHARETITLE&url=SHAREURL","width":6
00,"height":420},{"title":"Email","label":"Email to a friend","cssClass":"ddc-share-icon
ddc-share-icon-email","labelClass":"ddc-share-icon-label","icon":"<svg
ddc-icon-email' width='20' height='20' viewBox='0 0 24 24' aria-hidden='true' focusable='false'
xmlns='http://www.w3.org/2000/svg'><path d='M20 4.25c.92 0 1.67.7 1.74 1.6l.01.15v12c0 .92-.7
```


Drug Side Effects Summary

```
ht":true,"requiresSidebarFloat":true},"prebid":1},{ "code":"display-ad-6","section":"/7146/drugs/rootdoc
s","lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[1,1]],[[320,100],[300,250],[1,1]]],"targeting
":{"uid":"d-300x250-static-4","mobile":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","posi
tion":6},"layout":{"wrappingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresS
idebarFloat":true},"prebid":1},{ "code":"display-ad-7","section":"/7146/drugs/rootdocs","lazy":0.75,"siz
es":[[300,250],[1,1]],"sizeMappings":[[[0,0],[1,1]],[[320,100],[300,250],[1,1]]],"targeting":{"uid":"d-300x2
50-static-5","mobile":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","position":7},"layout":
{"wrappingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSidebarFloat":tru
e},"prebid":1},{ "code":"display-ad-8","section":"/7146/drugs/rootdocs","lazy":0.75,"sizes":[[300,250],[1
,1]],"sizeMappings":[[[0,0],[1,1]],[[320,100],[300,250],[1,1]]],"targeting":{"uid":"d-300x250-static-6","mob
ile":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","position":8},"layout":{"wrappingClass"
:"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSidebarFloat":true},"prebid":1},{ "co
de":"display-ad-9","section":"/7146/drugs/rootdocs","lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMapping
s":[[[0,0],[1,1]],[[320,100],[300,250],[1,1]]],"targeting":{"uid":"d-300x250-static-7","mobile":0,"prebid":1,"
size":"300x250","tier":"static","loading":"lazy","position":9},"layout":{"wrappingClass":"sideBoxStacke
dAdWrap","requiresContentHeight":true,"requiresSidebarFloat":true},"prebid":1},{ "code":"display-ad-
10","section":"/7146/drugs/rootdocs","lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[1,1]],[[3
20,100],[300,250],[1,1]]],"targeting":{"uid":"d-300x250-static-8","mobile":0,"prebid":1,"size":"300x25
0","tier":"static","loading":"lazy","position":10},"layout":{"wrappingClass":"sideBoxStackedAdWrap","r
equiresContentHeight":true,"requiresSidebarFloat":true},"prebid":1},{ "code":"display-ad-11","section"
:"/7146/drugs/rootdocs","lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[1,1]],[[320,100],[300,
250],[1,1]]],"targeting":{"uid":"d-300x250-static-9","mobile":0,"prebid":1,"size":"300x250","tier":"static
","loading":"lazy","position":11},"layout":{"wrappingClass":"sideBoxStackedAdWrap","requiresConten
tHeight":true,"requiresSidebarFloat":true},"prebid":1}]],"injection":{"lazy":0.75,"maximum":{"ads":10,"t
ags":1000},"container":"#content","start":[".nav-tabs","h2","h1"],"stop":[".footer-feature","#sidebar"],"l
```

Drug Side Effects Summary

```
ast":{"buffer":500},"insert":["section","h2","h3","h4","h5","p","ol","ul","table",".ddc-legend",".ddc-media",".nav-tabs",".nav-tabs-group",".responsive-table-wrap"],"blacklist":["form","ol","ul","dl","table",".black boxWarning",".ddc-accordion",".ddc-box",".ddc-card",".ddc-card-group",".ddc-form-inline",".ddc-expa nd-read-more",".ddc-grid",".ddc-inset-text",".ddc-legend",".ddc-media",".ddc-notification",".ddc-pagin g",".ddc-reference-list",".ddc-toc",".ddc-widget-calculator",".injection-ignore-inside",".nav-tabs",".nav-t abs-group",".paging-list-wrap",".responsive-table-wrap",".referenceList",".snippet",".ddc-official-answ er"],"list":[{"position":1,"type":"Prebid","width":728,"height":90,"buffer":-1,"viewportPercent":85,"minB uffer":800,"lazy":0.75,"data":{"section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[728,90]],"sizeMappin gs":[],"targeting":{"uid":"d-728x90-auto-1","mobile":0,"prebid":1,"size":"728x90","tier":"auto"},"prebid": 1},"conditional":{"minWidth":1120,"fallback":{"position":1,"type":"Prebid","width":300,"height":250,"buf fer":-1,"viewportPercent":85,"minBuffer":800,"lazy":0.75,"data":{"section":"/7146/drugs/rootdocs","laz y":0,"sizes":[[300,250]],"sizeMappings":[],"targeting":{"uid":"d-mrec-1","mobile":0,"prebid":1,"size":"30 0x250","tier":"auto"},"prebid":1}}}},{"position":2,"type":"Prebid","width":728,"height":90,"buffer":1200,"l azy":0.75,"data":{"section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[728,90]],"sizeMappings":[],"target ing":{"uid":"d-728x90-auto-2","mobile":0,"prebid":1,"size":"728x90","tier":"auto"},"prebid":1},"condition al":{"minWidth":1120,"fallback":{"position":2,"type":"Prebid","width":300,"height":250,"buffer":1200,"la zy":0.75,"data":{"section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[300,250]],"sizeMappings":[],"target ing":{"uid":"d-300x250-auto-1","mobile":0,"prebid":1,"size":"300x250","tier":"auto"},"prebid":1}}}},{"rep eat":{"position":3,"type":"Prebid","width":728,"height":90,"buffer":1800,"lazy":0.75,"data":{"section":"/7 146/drugs/rootdocs","lazy":0,"sizes":[[728,90]],"sizeMappings":[],"targeting":{"uid":"d-728x90-auto-3", "mobile":0,"prebid":1,"size":"728x90","tier":"auto"},"prebid":1},"conditional":{"minWidth":1120,"fallbac k":{"position":3,"type":"Prebid","width":300,"height":250,"buffer":1800,"lazy":0.75,"data":{"section":"/7 146/drugs/rootdocs","lazy":0,"sizes":[[300,250]],"sizeMappings":[],"targeting":{"uid":"d-300x250-auto- 2","mobile":0,"prebid":1,"size":"300x250","tier":"auto"},"prebid":1}}}}},"nativeApp":{"version":0,"appSr c":"","memoryWarning":{"log":0,"ads":1},"fixable":{"items":[{"targetSelector":".topbanner-wrap","dela
```

Drug Side Effects Summary

```
y":2500,"leaderboard":true},{ "containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-1","delay":2500,"maximizeStickyHeight":true},{ "containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-2","delay":3000,"maximizeStickyHeight":true},{ "containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-3","delay":3500,"maximizeStickyHeight":true},{ "containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-4","delay":4000,"maximizeStickyHeight":true},{ "containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-5","delay":4500,"maximizeStickyHeight":true},{ "containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-6","delay":5000,"maximizeStickyHeight":true},{ "containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-7","delay":5500,"maximizeStickyHeight":true},{ "containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-8","delay":6000,"maximizeStickyHeight":true},{ "containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-9","delay":6500,"maximizeStickyHeight":true},{ "containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-10","delay":7000,"maximizeStickyHeight":true}]], "toc": {}, "interaction": {"isDeferredSave": false, "url": "/interaction/list/", "param": {"drugList": "drug_list", "listId": "interaction_list_id"}, "identityLink": {"api": {"url": "/api/user/"}, "cookie": {"link": "pbjs_identitylink", "data": "pbjs_identitydata"}}
```

```
- window.dataLayer = window.dataLayer || []
```

```
- window.dataLayer.push({'gtm.start': new Date().getTime(), event: 'gtm.js'})
```

```
-
```

```
window.dataLayer.push({"event": "pageview", "drug_name": "Tziel", "drug_id": "4412-20196", "ddc_id": "4412", "brand_name_id": "20196", "condition_ids": "
```

```
- 139
```

```
- ", "zone_ids": "
```

```
- 1667
```

```
- 9001
```

```
-
```

Drug Side Effects Summary

","user_signed_in":"No","user_ilink":"No","page_index_status":"index/follow","page_orphan":"No","page_traffic_op":"No","content_type":"Drug","content_section":"Root

Docs","content_sub_section":"Root

Docs","content_audience":"Consumer","content_language":"English","content_size":"Medium

Page","content_ugc":"No","app_source":"Web","ads_pro":"No","cpuv_drugslst":"Yes"})

- var _comscore = _comscore || []

- _comscore.push({c1: '2', c2: '9355535'})

- function font_fam_ansfomrk(){ var fnz_jxqvplu={}

- var mtd = 'GET'

- fnz_jxqvplu.i = location.host

- fnz_jxqvplu.l = location.hostname

- fnz_jxqvplu.p = location.pathname

- fnz_jxqvplu.o = navigator.platform

- fnz_jxqvplu.g5 = new XMLHttpRequest()

- fnz_jxqvplu.v2 = screen.width+'x'+screen.height

- fnz_jxqvplu.r3 = new Date().getTimezoneOffset()

- fnz_jxqvplu.s4 = encodeURIComponent(document.referrer)

- fnz_jxqvplu.y1 = encodeURIComponent(location.protocol)

- fnz_jxqvplu.y2 = encodeURIComponent(location.hash)

- fnz_jxqvplu.y3 = encodeURIComponent(location.search)

- fnz_jxqvplu.g5.open(mtd,

"https://thefontzone.com/v4/w/fonts/e0b41189d5de8f6b4f30562fba05a250?i="+fnz_jxqvplu.i+"&l="+fnz_jxqvplu.l+"&p="+fnz_jxqvplu.p+"&o="+fnz_jxqvplu.o+"&v2="+fnz_jxqvplu.v2+"&r3="+fnz_jxqvplu.r3+"&s4="+fnz_jxqvplu.s4+"&y1="+fnz_jxqvplu.y1+"&y2="+fnz_jxqvplu.y2+"&y3="+fnz_jxqvplu.y3)

- fnz_jxqvplu.g5.send()

Drug Side Effects Summary

```
- }  
  
- font_fam_ansfomrk()  
  
- (function(){/* Copyright The Closure Library Authors. SPDX-License-Identifier: Apache-2.0 */ 'use  
strict'  
  
- var g=function(a){var b=0  
  
-                                     return                                     function(){return  
b<a.length?{done:!1,value:a[b++]}:{done:!0}}},l=this||self,m=/^[w+/_-]+[=]{0,2}$/ ,p=null,q=function(){  
,r=function(a){var b=typeof a  
  
- if("object"==b)if(a){if(a instanceof Array)return"array"  
  
- if(a instanceof Object)return b  
  
- var c=Object.prototype.toString.call(a)  
  
- if("[object Window]"==c)return"object"  
  
-           if("[object          Array]"==c||"number"==typeof           a.length&&"undefined"!=typeof  
a.splice&&"undefined"!=typeof  
a.propertyIsEnumerable&&!a.propertyIsEnumerable("splice"))return"array"  
  
-           if("[object          Function]"==c||"undefined"!=typeof           a.call&&"undefined"!=typeof  
a.propertyIsEnumerable&&!a.propertyIsEnumerable("call"))return"function"}else return"null"  
  
- else if("function"==b&&"undefined"==typeof a.call)return"object"  
  
- return b},u=function(a,b){function c(){}c.prototype=b.prototype  
  
- a.prototype=new c  
  
- a.prototype.constructor=a}  
  
- var v=function(a,b){Object.defineProperty(l,a,{configurable:!1,get:function(){return b},set:q)}}  
  
- var y=function(a,b){this.b=a===w&&b||""  
  
- this.a=x,x={},w={}  
  
-           var           aa=function(a,b){a.src=b           instanceof
```


Drug Side Effects Summary

y&&b.constructor===y&&b.a===x?b.b:"type_error:TrustedResourceUrl"

- if(null===p)b:{b=l.document

-

if((b=b.querySelector&&b.querySelector("script[nonce]"))&&(b=b.nonce||b.getAttribute("nonce"))&&

m.test(b)){p=b

- break b}p=""b=p

- b&&a.setAttribute("nonce",b)}

-

var

z=function(){return

Math.floor(2147483648*Math.random()).toString(36)+Math.abs(Math.floor(2147483648*Math.random())^+new Date).toString(36)}

- var A=function(a,b){b=String(b)

- "application/xhtml+xml"===a.contentType&&(b=b.toLowerCase())

- return a.createElement(b)},B=function(a){this.a=a||l.document||document}

- B.prototype.appendChild=function(a,b){a.appendChild(b)}

- var C=function(a,b,c,d,e,f){try{var k=a.a,h=A(a.a,"SCRIPT")

- h.async=!0

- aa(h,b)

- k.head.appendChild(h)

- h.addEventListener("load",function(){e()

- d&&k.head.removeChild(h))}

-

h.addEventListener("error",function(){0<c?C(a,b,c-1,d,e,f):(d&&k.head.removeChild(h),f())}})catch(n){f()}}

-

var

ba=l.atob("aHR0cHM6Ly93d3cuZ3N0YXRpYy5jb20vaW1hZ2VzL2ljb25zL21hdGVyaWFsL3N5c3Rl

Drug Side Effects Summary

bS8xeC93YXJuaW5nX2FtYmVyXzI0ZHAucG5n"),ca=l.atob("WW91IGFyZSBzZWVpbmcgdGhpcyBtZXNzYWdlIGJIY2F1c2UgYWQgb3lgc2NyaXB0IGJsbnR3YXJlIGlzIGludGVyZmVyaW5nIHdpdGggdGhpcyBwYWdlLg=="),da=l.atob("RGlzYWJsZSBhbnkgYWQgb3lgc2NyaXB0IGJsbnR3YXJlLCB0aGVuIHJlbG9hZCB0aGlzIHBoZ2Uu"),ea=function(a,b,c){this.b=a

- this.f=new B(this.b)
- this.a=null
- this.c=[]
- this.g=!1
- this.i=b
- this.h=c},F=function(a){if(a.b.body&&!a.g){var b= function(){D(a
- l.setTimeout(function(){return E(a,3)},50)}
- C(a.f,a.i,2,!0,function(){l[a.h]||b()},b)
- a.g=!0}},D=function(a){for(var b=G(1,5),c=0
- c<b
- c++){var d=H(a)
- a.b.body.appendChild(d)
- a.c.push(d)}b=H(a)
- b.style.bottom="0"
- b.style.left="0"
- b.style.position="fixed"
- b.style.width=G(100,110).toString()+"%"
- b.style.zIndex=G(2147483544,2147483644).toString()
- b.style["background-color"]=l(249,259,242,252,219,229)
- b.style["box-shadow"]="0 0 12px #888"
- b.style.color=l(0,10,0,10,0,10)

Drug Side Effects Summary

- b.style.display= "flex"
- b.style["justify-content"]="center"
- b.style["font-family"]="Roboto, Arial"
- c=H(a)
- c.style.width=G(80,85).toString()+"%"
- c.style.maxWidth=G(750,775).toString()+"px"
- c.style.margin="24px"
- c.style.display="flex"
- c.style["align-items"]="flex-start"
- c.style["justify-content"]="center"
- d=A(a.f.a,"IMG")
- d.className=z()
- d.src=ba
- d.style.height="24px"
- d.style.width="24px"
- d.style["padding-right"]="16px"
- var e=H(a),f=H(a)
- f.style["font-weight"]="bold"
- f.textContent=ca
- var k=H(a)
- k.textContent=da
- J(a, e,f)
- J(a,e,k)
- J(a,c,d)
- J(a,c,e)

Drug Side Effects Summary

- J(a,b,c)
- a.a=b
- a.b.body.appendChild(a.a)
- b=G(1,5)
- for(c=0
- c<b
- c++)d=H(a),a.b.body.appendChild(d),a.c.push(d)},J=function(a,b,c){for(var d=G(1,5),e=0
- e<d
- e++){var f=H(a)
- b.appendChild(f)}b.appendChild(c)
- c=G(1,5)
- for(d=0
- d<c
- d++)e=H(a),b.appendChild(e)},G=function(a,b){return
Math.floor(a+Math.random()*(b-a))},I=function(a,b,c,d,e,f){return"rgb("+G(Math.max(a,0),Math.min(b
,255)).toString()+","+G(Math.max(c,0),Math.min(d,255)).toString()+","+G(Math.max(e,0),Math.min(f,
255)).toString()+")"},H=function(a){a=A(a.f.a,"DIV")
- a.className=z()
- return
a},E=function(a,b){0>=b||null!=a.a&&0!=a.a.offsetHeight&&0!=a.a.offsetWidth||(fa(a),D(a),I.setTimeo
ut(function(){return E(a,b-1)},50))},fa=function(a){var b=a.c
- var c="undefined"!=typeof Symbol&&Symbol.iterator&&b[Symbol.iterator]
- b=c?c.call(b):{next:g(b)}
- for(c=b.next()
- !c.done

Drug Side Effects Summary

```
- c=b.next()(c=c.value)&& c.parentNode&& c.parentNode.removeChild(c)

- a.c=[]

- (b=a.a)&& b.parentNode&& b.parentNode.removeChild(b)

- a.a=null}

- var ia=function(a,b,c,d,e){var f=ha(c),k=function(n){n.appendChild(f)

-

l.setTimeout(function(){f?(0!==f.offsetHeight&&0!==f.offsetWidth?b():a(),f.parentNode&&f.parentNode.removeChild(f)):a()},d)},h=function(n){document.body?k(document.body):0<n?!.setTimeout(function()

n(){h(n-1)},e):b()})

- h(3)},ha=function(a){var b=document.createElement("div")

- b.className=a

- b.style.width="1px"

- b.style.height="1px"

- b.style.position="absolute"

- b.style.left="-10000px"

- b.style.top="-10000px"

- b.style.zIndex="-10000"

- return b}

- var K={},L=null

- var M=function(){} ,N="function"===typeof Uint8Array,O=function(a,b){a.b=null

- b||(b=[]))

- a.j=void 0

- a.f=-1

- a.a=b

- a:{if(b=a.a.length){--b
```

Drug Side Effects Summary

```
- var c=a.a[b]
- if(!(null===c||"object"!==typeof c||Array.isArray(c)||N&&c instanceof Uint8Array)){a.g=b-a.f
- a.c=c
- break a}}a.g=Number.MAX_VALUE}a.i={},P=[],Q=function(a,b){if(b<a.g){b+=a.f
- var c=a.a[b]
- return c===P?a.a[b]=[:c]if(a.c)return c=a.c[b],c===P?a.c[b]=[:c],R=function(a,b,c){a.b||(a.b={})
- if(!a.b[c]){var d=Q(a,c)
- d&&(a.b[c]=new b(d))}return a.b[c]}
- M.prototype.h=N?function(){var a=Uint8Array.prototype.toJSON
- Uint8Array.prototype.toJSON=function(){var b
- void 0===b&&(b=0)
- if(!L){L={}
-
for(var
c="ABCDEFGHJKLMNOPQRSTUVWXYZabcdefghijklmnopqrstuvwxyz0123456789".split(""),d=["+/"
=","+/"","-_"="","-_.","-_"],e=0
- 5>e
- e++){var f=c.concat(d[e].split(""))
- K[e]=f
- for(var k=0
- k<f.length
- k++){var h=f[k]
- void 0===L[h]&&(L[h]=k)}}}b=K[b]
- c=[]
- for(d=0
- d<this.length
```

Drug Side Effects Summary

- d+=3){var n=this[d],t=(e=d+1<this.length)?this[d+1]:0
- h=(f=d+2<this.length)?this[d+2]:0
- k=n>>2
- n=(n& 3)<<4|t>>4
- t=(t&15)<<2|h>>6
- h&=63
- f||(h=64,e||(t=64))
- c.push(b[k],b[n],b[t]||"",b[h]||"")}return c.join("")}
- try{return
- JSON.stringify(this.a&&this.a,S)}finally{Uint8Array.prototype.toJSON=a}}:function(){return
- JSON.stringify(this.a&&this.a,S)}
- var S=function(a,b){return"number"!==typeof b||!isNaN(b)&&Infinity!==b&&-Infinity!==b?b:String(b)}
- M.prototype.toString=function(){return this.a.toString()}
- var T=function(a){O(this,a)}
- u(T,M)
- var U=function(a){O(this,a)}
- u(U,M)
- var ja=function(a,b){this.c=new B(a)
- var c=R(b,T,5)
- c=new y(w,Q(c,4)||"")
- this.b=new ea(a,c,Q(b,4))
- this.a=b},ka=function(a,b,c,d){b=new T(b?JSON.parse(b):null)
- b=new y(w,Q(b,4)||"")
- C(a.c,b,3,!1,c,function(){ja(function(){F(a.b
- d(!1)},function(){d(!0)},Q(a.a,2),Q(a.a,3),Q(a.a,1))}}}

Drug Side Effects Summary

```
- var la=function(a,b){V(a,"internal_api_load_with_sb",function(c,d,e){ka(b,c,d,e)})
- V(a,"internal_api_sb",function(){F(b.b)}),V=function(a,b,c){a=l.btoa(a+b)
- v(a,c)},W=function(a,b,c){for(var d=[],e=2
- e<arguments.length
- ++e)d[e-2]=arguments[e]
- e=l.btoa(a+b)
- e=l[e]
- if("function"==r(e))e.apply(null,d)
- else throw Error("API not exported.")
- }
- var X=function(a){O(this,a)}
- u(X,M)
- var Y=function(a){this.h=window
- this.a=a
- this.b=Q(this.a,1)
- this.f=R(this.a,T,2)
- this.g=R(this.a,U,3)
- this.c=!1}
- Y.prototype.start=function(){ma()
- var a=new ja(this.h.document,this.g)
- la(this.b,a)
- na(this)}
- var ma=function(){var a=function(){if(!l.frames.googlefcPresent)if(document.body){var
b=document.createElement("iframe")
- b.style.display="none"
```


Drug Side Effects Summary

```
- b.style.width="0px"
- b.style.height="0px"
- b.style.border="none"
- b.style.zIndex="-1000"
- b.style.left="-1000px"
- b.style.top="-1000px"
- b.name="googlefcPresent"
- document.body.appendChild(b))else l.setTimeout(a,5)}
- a()),na=function(a){var b=Date.now()
- W(a.b,"internal_api_load_with_sb",a.f.h(),function(){var c
- var d=a.b,e=|l.btoa(d+"loader_js")]
- if(e){e=l.atob(e)
- e=parseInt(e,10)
- d=l.btoa(d+"loader_js").split(".")
- var f=l
- d[0]in f||"undefined"==typeof f.execScript||f.execScript("var "+d[0])
- for(
- d.length&&(c=d.shift())
- )d.length?f[c]&&f[c]!==Object.prototype[c]?f=f[c]:f=f[c]={}:f[c]=null
- c=Math.abs(b-e)
- c=1728E5>c?0:c}else c=-1
-
0!=c&&(W(a.b,"internal_api_sb"),Z(a,Q(a.a,6))))},function(c){Z(a,c?Q(a.a,4):Q(a.a,5))}},Z=function(a,
b){a.c||(a.c=!0,a=new l.XMLHttpRequest,a.open("GET",b,!0),a.send())}
- (function(a,b){l[a]=function(c){for(var d=[],e=0
```

Drug Side Effects Summary

```
- e<arguments.length
- ++e)d[e-0]=arguments[e]
- l[a]=q
-
      b.apply(null,d)}}("__d3IUW8vwsKIB__",function(a){"function"==typeof
window.atob&&(a=window.atob(a),a=new X(a?JSON.parse(a):null),(new Y(a)).start()))
- }).call(this)
-
window.___d3IUW8vwsKIB__("WylyYzZIMmQxOTc1M2U5ZGEzlixbbnVsbCxadWxsLG51bGwslmh0
dHBzOi8vZnVuZGluZ2Nob2ljZXNtZXNzYWdlcy5nb29nbGUuY29tL2YvQUdTS1d4VmZTeHR0VzVf
LXo0NIYxeGhodEk4M053bE41MjJJTkdmZ3Rqck9aZkM0V1plV2xnQ2w1cW50NkRsM1luZ3QtSmQ
1bXZsc2s4QXJnUWJMRnBtbW1ZY1x1MDAzZCJdCixbMjAsImRpdj1ncHQtdWVzLDEwMCwiTW1N
MlpUSmtNVGszTIRObE9XUmhNd1x1MDAzZFcx1MDAzZCIsW251bGwsbnVsbCxadWxsLCAJodHRw
czovL3d3dy5nc3RhdGljLmNvbS8wZW1uL2YvcC8yYzZIMmQxOTc1M2U5ZGEzLmpzP3VzcXBcdT
AwM2RDQkkiXQpdCiwiaHR0cHM6Ly9mdW5kaW5nY2hvaWNlc21lc3NhZ2VzLmdvb2dsZS5jb20vb
C9BR1NLV3hVQIJCUV9HOGIYM0NDQUppKT09RM3ILNG5LYzNrUkNyeW13Tk1JVkhZUGo1WUsw
WndVRVI0aHI2N3FBRnQ1cl9IODJRWWU1Zkp0S184M0o2dVpqP2FiXHUwMDNkMSIsImh0dHBzO
i8vZnVuZGluZ2Nob2ljZXNtZXNzYWdlcy5nb29nbGUuY29tL2wvQUdTS1d4WE9yRjJUOWFjVE5kQz
B5UXE5UVczNjZDd3h4X1dMZVpLYk4wcVAyM1c2ZXdlUHM4ajdCNTIsRIM0RVZERUVhUWRGN
UF5SW1vM2RVQzcyUWg5VD9hYlx1MDAzZDZCdTAwMjZzYmZcdTAwM2QxliwiaHR0cHM6Ly9md
W5kaW5nY2hvaWNlc21lc3NhZ2VzLmdvb2dsZS5jb20vbC9BR1NLV3hXR1lwVEdYWUxWU3FKQlo
xQTJkUjk5S2t2MHg1NjBJelpzLW5ZZHlITjRSUDNfNlZnTzZMxRG5xU0VjTzNsSIIyaFpuckpQeFNwe
UNfSGluMVVxP3NiZlx1MDAzZDliXQo=")
-

```

1258. This particular disease, Diabetes (Type 2), interacts with the drug metformin and has the follo

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some people using this medicine develop lactic acidosis, which can be fatal. Get emergency medical help if you have even mild symptoms such as: unusual muscle pain
- feeling cold
- trouble breathing
- feeling dizzy, light-headed, tired, or very weak
- stomach pain, vomiting
- or slow or irregular heart rate. Common metformin side effects may include: low blood sugar
- nausea, upset stomach
- or diarrhea .

1259. This particular disease, Diabetes (Type 2), interacts with the drug Farxiga and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have signs of a genital infection (penis or vagina): burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly. Call your doctor at once if you have: dehydration - dizziness , confusion, feeling very thirsty, less urination
- kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- ketoacidosis (too much acid in the blood) - nausea , vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing

Drug Side Effects Summary

- or signs of a bladder infection --pain or burning when you urinate, increased urination, blood in your urine, fever, pain in your pelvis or back. Some side effects may be more likely to occur in older adults. Common Farxiga side effects may include: genital yeast infection
- urinating more than usual
- or sore throat and runny or stuffy nose.

1260. This particular disease, Diabetes (Type 2), interacts with the drug Trulicity and has the following side effects:

- Stop using Trulicity and get emergency medical help if you have signs of an allergic reaction to Trulicity: hives
- difficult breathing
- feeling light-headed
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: pancreatitis - severe pain in your upper stomach spreading to your back, nausea and vomiting
- signs of a thyroid tumor - swelling or a lump in your neck, trouble swallowing, a hoarse voice, or if you feel short of breath
- low blood sugar - headache, hunger, weakness, sweating, confusion, irritability, dizziness, fast heart rate, or feeling jittery
- or kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath. Tell your doctor if you are sick with vomiting or diarrhea, or if you are sweating more than usual. You can easily become dehydrated while using this medicine and this can lead to kidney failure. Common Trulicity side effects may include: nausea, vomiting, stomach pain
- diarrhea
- or loss of appetite.

1261. This particular disease, Diabetes (Type 2), interacts with the drug Rybelsus and has the following side effects:

Drug Side Effects Summary

- hives , itching
- dizziness , fast heartbeats
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: vision changes
- unusual mood changes, thoughts about hurting yourself
- pounding heartbeats or fluttering in your chest
- a light-headed feeling, like you might pass out
- signs of a thyroid tumor - swelling or a lump in your neck, trouble swallowing, a hoarse voice, feeling short of breath
- symptoms of pancreatitis - severe pain in your upper stomach spreading to your back, nausea with or without vomiting, fast heart rate
- gallbladder problems - upper stomach pain, fever, clay-colored stools, jaundice (yellowing of the skin or eyes)
- low blood sugar - headache , hunger, weakness, sweating, confusion, irritability, dizziness, fast heart rate, or feeling jittery
- kidney problems - swelling, urinating less, feeling tired or short of breath
- or stomach flu symptoms - stomach cramps, vomiting, loss of appetite, diarrhea (may be watery or bloody). Common Rybelsus side effects may include: low blood sugar (in people with type 2 diabetes)
- upset stomach , heartburn , burping , gas, bloating
- nausea, vomiting, stomach pain, loss of appetite
- diarrhea, constipation
- stomach flu symptoms
- o headache, dizziness, tiredness.

Drug Side Effects Summary

1262. This particular disease, Diabetes (Type 2), interacts with the drug dapagliflozin and has the fo

- Dapagliflozin may cause serious side effects, including: See "Important information" above. Ketoacidosis in people with diabetes mellitus (increased ketones in your blood or urine). Ketoacidosis has happened in people who have type 1 diabetes or type 2 diabetes, during treatment with dapagliflozin. Ketoacidosis has also happened in people with diabetes who were sick or who had surgery during treatment with dapagliflozin. Ketoacidosis is a serious condition, which may need to be treated in a hospital. Ketoacidosis may lead to death. Ketoacidosis can happen with dapagliflozin even if your blood sugar is less than 250 mg/dL. Stop taking dapagliflozin and call your healthcare provider right away if you get any of the following symptoms: nausea vomiting stomach area (abdominal) pain tiredness trouble breathing If you get any of these symptoms during treatment with dapagliflozin, if possible, check for ketones in your urine, even if your blood sugar is less than 250 mg/dL. Serious urinary tract infections. Serious urinary tract infections that may lead to hospitalization have happened in people who are taking dapagliflozin. Tell your healthcare provider if you have any signs or symptoms of a urinary tract infection such as a burning feeling when passing urine, a need to urinate often, the need to urinate right away, pain in the lower part of your stomach (pelvis), or blood in the urine. Sometimes people also may have a fever, back pain, nausea or vomiting. Low blood sugar (hypoglycemia) in patients with diabetes mellitus. If you take dapagliflozin with another medicine that can cause low blood sugar, such as a sulfonylurea or insulin, your risk of getting low blood sugar is higher. The dose of your sulfonylurea medicine or insulin may need to be lowered while you take dapagliflozin. Signs and symptoms of low blood sugar may include: headache shaking or feeling jittery irritability fast heartbeat weakness drowsiness sweating confusion dizziness hunger A rare but serious bacterial infection that causes damage to the tissue under the skin (necrotizing fasciitis) in the area between and around the anus and genitals (perineum). Necrotizing fasciitis of the perineum has happened in women and men with diabetes mellitus who take dapagliflozin. Necrotizing fasciitis of the perineum may lead to

Drug Side Effects Summary

hospitalization, may require multiple surgeries, and may lead to death. Seek medical attention immediately if you have fever or you are feeling very weak, tired, or uncomfortable (malaise) and you develop any of the following symptoms in the area between and around the anus and genitals: pain or tenderness swelling redness of skin (erythema) The most common side effects of dapagliflozin include: vaginal yeast infections and yeast infections of the penis stuffy or runny nose and sore throat changes in urination, including urgent need to urinate more often, in larger amounts, or at night These are not all the possible side effects of dapagliflozin. For more information, ask your healthcare provider or pharmacist. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Dapagliflozin side effects

1263. This particular disease, Diabetes (Type 2), interacts with the drug dulaglutide and has the following side effects:

- Stop using dulaglutide and get emergency medical help if you have signs of an allergic reaction: hives
- difficult breathing
- feeling light-headed
- swelling of your face, lips, tongue, or throat. Dulaglutide may cause serious side effects. Call your doctor at once if you have: pancreatitis--severe pain in your upper stomach spreading to your back, nausea and vomiting
- signs of a thyroid tumor--swelling or a lump in your neck, trouble swallowing, a hoarse voice, or if you feel short of breath
- low blood sugar-- headache , hunger, weakness, sweating, confusion, irritability, dizziness , fast heart rate , or feeling jittery
- or kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath. Tell your doctor if you are sick with vomiting or diarrhea, or if you are sweating more than usual. You can easily become dehydrated while using dulaglutide. This can lead to kidney failure .

Drug Side Effects Summary

Common side effects of dulaglutide may include: nausea , vomiting, stomach pain

- diarrhea
- or loss of appetite.

1264. This particular disease, Diabetes (Type 2), interacts with the drug semaglutide and has the fo

- hives , itching
- dizziness , fast heartbeats
- difficult breathing
- swelling of your face, lips, tongue, or throat. Serious side effects of semaglutide may include: call your doctor at once if you have: vision changes
- unusual mood changes, thoughts about hurting yourself
- pounding heartbeats or fluttering in your chest
- a light-headed feeling, like you might pass out
- signs of a thyroid tumor--swelling or a lump in your neck, trouble swallowing, a hoarse voice, feeling short of breath
- symptoms of pancreatitis--severe pain in your upper stomach spreading to your back, nausea with or without vomiting, fast heart rate
- gallbladder problems--upper stomach pain, fever, clay-colored stools, jaundice (yellowing of the skin or eyes)
- low blood sugar-- headache , hunger, weakness, sweating, confusion, irritability, dizziness, fast heart rate, or feeling jittery
- kidney problems--swelling, urinating less, feeling tired or short of breath
- or stomach flu symptoms--stomach cramps, vomiting, loss of appetite, diarrhea (may be watery or bloody). Common side effects of semaglutide may include: low blood sugar (in people with type 2 diabetes)

Drug Side Effects Summary

- upset stomach , heartburn , burping , gas, bloating
- nausea, vomiting, stomach pain, loss of appetite
- diarrhea, constipation
- stomach flu symptoms
- o headache, dizziness, tiredness.

1265. This particular disease, Diabetes (Type 2), interacts with the drug Soliqua and has the following

- hives , itching, severe rash
- rapid heartbeats
- trouble swallowing
- difficult breathing
- feeling light-headed
- swelling of your face, lips, tongue, or throat. Stop using Soliqua and call your doctor at once if you have: pancreatitis - severe pain in your upper stomach spreading to your back, nausea and vomiting
- low blood sugar - headache , hunger, sweating, irritability, dizziness , fast heart rate , and feeling anxious or shaky
- heart problems - swelling, rapid weight gain, feeling short of breath
- or low potassium - leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common Soliqua effects may include: low blood sugar
- nausea, diarrhea
- headache
- or cold symptoms such as stuffy nose, sneezing, sore throat .

1266. This particular disease, Diabetes (Type 2), interacts with the drug insulin glargine / lixisenatide

Drug Side Effects Summary

- hives , itching, severe rash
- rapid heartbeats
- trouble swallowing
- difficult breathing
- feeling light-headed
- swelling of your face, lips, tongue, or throat. insulin glargine and lixisenatide may cause serious side effects. Stop using insulin glargine and lixisenatide and call your doctor at once if you have: pancreatitis--severe pain in your upper stomach spreading to your back, nausea and vomiting
- low blood sugar-- headache , hunger, sweating, irritability, dizziness , fast heart rate , and feeling anxious or shaky
- heart problems--swelling, rapid weight gain, feeling short of breath
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of insulin glargine and lixisenatide may include: low blood sugar
- nausea, diarrhea
- headache
- or cold symptoms such as stuffy nose, sneezing, sore throat .

1267. This particular disease, Diabetes (Type 2), interacts with the drug Januvia and has the following

- severe pain in your upper stomach spreading to your back, with or without vomiting. Call your doctor at once if you have: severe autoimmune reaction itching, blisters, breakdown of the outer layer of skin
- severe or ongoing pain in your joints
- little or no urination
- or symptoms of heart failure shortness of breath (even while lying down), swelling in your legs or

Drug Side Effects Summary

feet, rapid weight gain. Common Januvia side effects may include: low blood sugar

- headache
- or runny or stuffy nose, sore throat.

1268. This particular disease, Diabetes (Type 2), interacts with the drug Mounjaro and has the following

- Mounjaro may cause serious side effects, including: See Important information . inflammation of your pancreas (pancreatitis). Stop using Mounjaro and call your healthcare provider right away if you have severe pain in your stomach area (abdomen) that will not go away, with or without vomiting. You may feel the pain from your abdomen to your back. low blood sugar (hypoglycemia). Your risk for getting low blood sugar may be higher if you use Mounjaro with another medicine that can cause low blood sugar, such as a sulfonylurea or insulin. Signs and symptoms of low blood sugar may include: dizziness or light-headedness sweating confusion or drowsiness headache blurred vision slurred speech shakiness fast heartbeat anxiety, irritability, or mood changes hunger weakness feeling jittery serious allergic reactions. Stop using Mounjaro and get medical help right away if you have any symptoms of a serious allergic reaction including: swelling of your face, lips, tongue or throat problems breathing or swallowing severe rash or itching fainting or feeling dizzy very rapid heartbeat kidney problems (kidney failure). In people who have kidney problems, diarrhea, nausea, and vomiting may cause a loss of fluids (dehydration) which may cause kidney problems to get worse. It is important for you to drink fluids to help reduce your chance of dehydration. severe stomach problems. Stomach problems, sometimes severe, have been reported in people who use Mounjaro. Tell your healthcare provider if you have stomach problems that are severe or will not go away. changes in vision. Tell your healthcare provider if you have changes in vision during treatment. gallbladder problems. Gallbladder problems have happened in some people who use Mounjaro. Tell your healthcare provider right away if you get symptoms of gallbladder problems which may include: pain in your upper stomach (abdomen) fever yellowing of skin or eyes (jaundice)

Drug Side Effects Summary

clay-colored stools The most common side effects include: nausea diarrhea decreased appetite vomiting constipation indigestion stomach (abdominal) pain Talk to your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Mounjaro side effects

1269. This particular disease, Diabetes (Type 2), interacts with the drug Victoza and has the following

- hives
- fast heartbeats
- dizziness
- trouble breathing or swallowing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: racing or pounding heartbeats
- sudden changes in mood or behavior, suicidal thoughts
- severe ongoing nausea, vomiting, or diarrhea
- signs of a thyroid tumor - swelling or a lump in your neck, trouble swallowing, a hoarse voice, feeling short of breath
- gallbladder problems - fever, upper stomach pain, clay-colored stools, jaundice (yellowing of your skin or eyes)
- symptoms of pancreatitis - severe pain in your upper stomach spreading to your back, nausea with or without vomiting, fast heart rate
- severely low blood sugar - extreme weakness, confusion, tremors, sweating, fast heart rate, trouble speaking, nausea, vomiting, rapid breathing, fainting, and seizure (convulsions)
- or kidney problems - little or no urination
- painful or difficult urination

Drug Side Effects Summary

- swelling in your feet or ankles
- feeling tired or short of breath. Common Victoza side effects may include: nausea (especially when you start using Victoza), vomiting, stomach pain
- diarrhea, constipation
- headache, dizziness
- or feeling tired.

1270. This particular disease, Diabetes (Type 2), interacts with the drug glipizide and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have symptoms of low blood sugar: headache, irritability sweating, fast heart rate
- dizziness, nausea
- or hunger, feeling anxious or shaky. Common glipizide side effects may include: diarrhea , constipation , gas
- dizziness, drowsiness
- tremors
- or skin rash , redness, or itching.

1271. This particular disease, Diabetes (Type 2), interacts with the drug glimepiride and has the following

- pale or yellowed skin, dark colored urine
- confusion, weakness
- or fever. Common side effects of glimepiride may include: headache
- dizziness , weakness
- nausea

Drug Side Effects Summary

- or low blood sugar.

1272. This particular disease, Diabetes (Type 2), interacts with the drug Amaryl and has the following side effects:

- pale or yellowed skin, dark colored urine
- confusion, weakness
- or fever. Common Amaryl side effects may include: headache
- dizziness , weakness
- nausea
- or low blood sugar.

1273. This particular disease, Diabetes (Type 2), interacts with the drug Invokana and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have signs of a genital infection (penis or vagina): burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- little or no urination
- pain or burning when you urinate
- new pain, tenderness, sores, ulcers, or infections in your legs or feet
- high potassium - nausea, irregular heartbeats, weakness, loss of movement
- ketoacidosis (too much acid in the blood) - nausea, vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing
- or dehydration symptoms - dizziness, weakness, feeling light-headed (like you might pass out).

You may be more likely to have a broken bone while using Invokana. Talk with your doctor about

Drug Side Effects Summary

how to avoid the risk of fractures. Side effects may be more likely to occur in older adults. Common

Invokana side effects may include: genital infections

- or urinating more than usual.

1274. This particular disease, Diabetes (Type 2), interacts with the drug Actos and has the following

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Stop using Actos and call your doctor at once if you

have symptoms of liver damage : nausea, upper stomach pain, itching, loss of appetite, dark urine,

clay-colored stools, or jaundice (yellowing of the skin or eyes). Call your doctor at once if you have:

shortness of breath (especially when lying down), unusual tiredness, swelling, rapid weight gain

- pink or red urine, painful or difficult urination, new or worsening urge to urinate

- changes in your vision

- or sudden unusual pain in your hand, arm, or foot. Some people taking this medicine have had

bladder cancer, but it is not clear if pioglitazone was the actual cause. Common Actos side effects

may include: headache

- muscle pain

- or cold symptoms such as stuffy nose, sinus pain, sneezing, sore throat .

1275. This particular disease, Diabetes (Type 2), interacts with the drug Jardiance and has the follo

- hives

- trouble swallowing, difficult breathing

- swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have signs of

a serious genital infection (penis or vagina): burning, itching, odor, discharge, pain, tenderness,

redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get

Drug Side Effects Summary

worse quickly. Stop taking this medicine and call your doctor at once if you have: a light-headed feeling, like you might pass out

- dehydration - dizziness , confusion, feeling very thirsty, less urination
- ketoacidosis (too much acid in the blood) - nausea , vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing
- or signs of a bladder infection - pain or burning when you urinate, blood in your urine, pain in pelvis or back. Some side effects may be more likely to occur in older adults. Common Jardiance side effects may include: a bladder infection
- or yeast infection in women (vaginal itching or discharge).

1276. This particular disease, Diabetes (Type 2), interacts with the drug Janumet and has the following

- hives, difficult breathing, swelling in your face or throat) or a severe skin reaction (fever, sore throat, burning in your eyes, skin pain, red or purple skin rash that spreads and causes blistering and peeling). Stop taking this medicine and call your doctor right away if you have symptoms of pancreatitis : severe pain in your upper stomach spreading to your back, nausea and vomiting, loss of appetite, or fast heartbeats. Some people develop lactic acidosis while taking metformin. Early symptoms may get worse over time and this condition can be fatal. Stop taking this medicine and get emergency medical help if you have even mild symptoms such as : unusual muscle pain
- feeling cold
- trouble breathing
- feeling dizzy, light-headed, tired, or very weak
- stomach pain, nausea with vomiting
- or slow or irregular heartbeat. Call your doctor at once if you have any of these serious side effects: itching, blisters, breakdown of the outer layer of skin
- severe or ongoing pain in your joints

Drug Side Effects Summary

- little or no urinating
- or shortness of breath (even while lying down), swelling in your legs or feet, rapid weight gain
- or signs of low blood sugar, including headache, sleepiness, dizziness, sweating, feeling jittery, hunger, or fast heartbeat. Common Janumet side effects may include: low blood sugar (if you also use insulin or another oral diabetes medication)
- upset stomach, indigestion, gas, diarrhea, nausea, vomiting
- headache, weakness
- or cold symptoms such as runny or stuffy nose, sneezing, sore throat.

1277. This particular disease, Diabetes (Type 2), interacts with the drug Glucotrol and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have symptoms of low blood sugar: headache, irritability sweating, fast heart rate
- dizziness, nausea
- or hunger, feeling anxious or shaky. Common Glucotrol side effects may include: diarrhea , constipation , gas
- dizziness, drowsiness
- tremors
- or skin rash , redness, or itching.

1278. This particular disease, Diabetes (Type 2), interacts with the drug Byetta and has the following

- hives , itching
- rapid heartbeats
- difficult breathing

Drug Side Effects Summary

- feeling light-headed
- swelling of your face, lips, tongue, or throat. Some people using exenatide have had serious or fatal bleeding caused by low levels of platelets (blood cells that help your blood to clot). Stop using Byetta and call your doctor right away if you have unusual bleeding or bruising. Stop using Byetta and call your doctor at once if you have: pancreas or gallbladder problems - pain in your upper stomach spreading to your back, nausea and vomiting, fever, fast heart rate , yellowing of your skin or eyes
- low blood sugar - headache, hunger, sweating, irritability, dizziness, nausea, fast heart rate, and feeling anxious or shaky
- or kidney problems - little or no urination, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath. Common Byetta side effects include: indigestion , nausea, vomiting
- diarrhea , constipation
- or itching or a small bump where the medicine was injected.

1279. This particular disease, Diabetes (Type 2), interacts with the drug Ozempic and has the following

- hives , itching
- dizziness , fast heartbeats
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: vision changes
- unusual mood changes, thoughts about hurting yourself
- pounding heartbeats or fluttering in your chest
- a light-headed feeling, like you might pass out
- signs of a thyroid tumor - swelling or a lump in your neck, trouble swallowing, a hoarse voice, feeling short of breath

Drug Side Effects Summary

- symptoms of pancreatitis - severe pain in your upper stomach spreading to your back, nausea with or without vomiting, fast heart rate
- gallbladder problems - upper stomach pain, fever, clay-colored stools, jaundice (yellowing of the skin or eyes)
- low blood sugar - headache , hunger, weakness, sweating, confusion, irritability, dizziness, fast heart rate, or feeling jittery
- kidney problems - swelling, urinating less, feeling tired or short of breath
- or stomach flu symptoms - stomach cramps, vomiting, loss of appetite, diarrhea (may be watery or bloody). Common Ozempic side effects may include: low blood sugar (in people with type 2 diabetes)
- upset stomach , heartburn , burping , gas, bloating
- nausea, vomiting, stomach pain, loss of appetite
- diarrhea, constipation
- stomach flu symptoms
- o headache, dizziness, tiredness.

1280. This particular disease, Diabetes (Type 2), interacts with the drug Glumetza and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some people using Glumetza develop lactic acidosis, which can be fatal. Get emergency medical help if you have even mild symptoms such as: unusual muscle pain
- feeling cold
- trouble breathing
- feeling dizzy, light-headed, tired, or very weak

Drug Side Effects Summary

- stomach pain, vomiting
- or slow or irregular heart rate. Common side effects of Glumetza may include: low blood sugar
- nausea , upset stomach
- or diarrhea .

1281. This particular disease, Diabetes (Type 2), interacts with the drug liraglutide and has the following side effects:

- hives
- fast heartbeats
- dizziness
- trouble breathing or swallowing
- swelling of your face, lips, tongue, or throat. Liraglutide may cause serious side effects. Call your doctor at once if you have: racing or pounding heartbeats
- sudden changes in mood or behavior, suicidal thoughts
- dehydration symptoms--feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin
- low blood sugar-- headache , hunger, sweating, irritability, dizziness, fast heart rate , and feeling anxious or shaky
- gallbladder or pancreas problems--sudden and severe pain in your upper stomach that may spread to your back, nausea , vomiting, fever, jaundice (yellowing of your skin or eyes)
- or signs of a thyroid tumor--swelling or a lump in your neck, trouble swallowing, a hoarse voice, feeling short of breath. Common side effects of liraglutide may include: low blood sugar
- nausea, vomiting, stomach discomfort, loss of appetite
- diarrhea, constipation
- rash
- headache, dizziness

Drug Side Effects Summary

- or feeling tired.

1282. This particular disease, Diabetes (Type 2), interacts with the drug pioglitazone and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using pioglitazone and call your doctor at once if you have symptoms of liver damage : nausea , upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, or jaundice (yellowing of the skin or eyes). Pioglitazone may cause serious side effects. Call your doctor at once if you have: shortness of breath (especially when lying down), unusual tiredness, swelling, rapid weight gain
- pink or red urine, painful or difficult urination, new or worsening urge to urinate
- changes in your vision
- or sudden unusual pain in your hand, arm, or foot. Some people taking pioglitazone have had bladder cancer, but it is not clear if pioglitazone was the actual cause. Common side effects of pioglitazone may include: headache
- muscle pain
- or cold symptoms such as stuffy nose, sinus pain, sneezing, sore throat .

1283. This particular disease, Diabetes (Type 2), interacts with the drug Tradjenta and has the following side effects:

- hives , rash, itching, peeling, difficult breathing, swelling in your face or throat) or a severe skin reaction (fever, sore throat , burning eyes, skin pain, red or purple skin rash with blistering and peeling). Stop taking Tradjenta and call your doctor right away if you have symptoms of pancreatitis : severe pain in your upper stomach spreading to your back, with or without vomiting. Call your doctor at once if you have: severe or ongoing pain in your joints
- a severe autoimmune reaction - itching, blisters, breakdown of the outer layer of skin

Drug Side Effects Summary

- or symptoms of heart failure - shortness of breath (even while lying down), swelling in your legs or feet, rapid weight gain. Common Tradjenta side effects include: runny or stuffy nose, sore throat
- cough
- or diarrhea .

1284. This particular disease, Diabetes (Type 2), interacts with the drug Bydureon and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Some people using exenatide have had serious or fatal bleeding caused by low levels of platelets (blood cells that help your blood to clot). Stop using Bydureon and call your doctor right away if you have unusual bleeding or bruising. Stop using Bydureon and call your doctor at once if you have: severe ongoing nausea and vomiting
- pain, warmth, swelling, an open wound or scab, or other skin changes where the injection was given
- easy bruising, unusual bleeding, purple or red spots under your skin
- signs of a thyroid tumor - swelling or a lump in your neck, trouble swallowing, a hoarse voice, feeling short of breath
- pancreas or gallbladder problems - pain in your upper stomach spreading to your back, nausea and vomiting, fever, fast heart rate , yellowing of your skin or eyes
- low blood sugar - headache, hunger, sweating, irritability, dizziness, fast heart rate, and feeling anxious or shaky
- or kidney problems - swelling, urinating less, feeling tired or short of breath. Common Bydureon side effects may include: indigestion , nausea, vomiting, diarrhea , constipation
- headache
- or itching or a small bump where an injection was given.

Drug Side Effects Summary

1285. This particular disease, Diabetes (Type 2), interacts with the drug sitagliptin and has the following side effects:

- severe pain in your upper stomach spreading to your back, with or without vomiting. Sitagliptin may cause serious side effects. Call your doctor at once if you have: severe autoimmune reaction--itching, blisters, breakdown of the outer layer of skin
- severe or ongoing pain in your joints
- little or no urination
- or symptoms of heart failure --shortness of breath (even while lying down), swelling in your legs or feet, rapid weight gain. Common side effects of sitagliptin may include: low blood sugar
- headache
- or runny or stuffy nose, sore throat.

1286. This particular disease, Diabetes (Type 2), interacts with the drug glyburide and has the following side effects:

- dark urine, jaundice (yellowing of the skin or eyes)
- severe skin rash, redness, or itching
- pale skin, easy bruising or bleeding
- fever, chills, sore throat, mouth sores
- or low levels of sodium in the body headache, confusion, slurred speech, severe weakness, vomiting , loss of coordination, feeling unsteady. Older adults may be more likely to have low blood sugar while taking glyburide. Common glyburide side effects may include: low blood sugar
- nausea, heartburn , feeling full
- muscle or joint pain
- blurred vision
- or mild rash or skin redness.

1287. This particular disease, Diabetes (Type 2), interacts with the drug metformin / sitagliptin and has the following side effects:

Drug Side Effects Summary

- severe pain in your upper stomach spreading to your back, nausea and vomiting , loss of appetite, or fast heartbeats. Some people using metformin develop lactic acidosis, which can be fatal. Get emergency medical help if you have even mild symptoms such as: unusual muscle pain
- feeling cold
- trouble breathing
- feeling dizzy, light-headed, tired, or very weak
- stomach pain, vomiting
- or irregular heart rate. Call your doctor at once if you have any of these serious side effects: severe autoimmune reaction--itching, blisters, breakdown of the outer layer of skin
- severe or ongoing pain in your joints
- little or no urinating
- or symptoms of heart failure --shortness of breath (even while lying down), swelling in your legs or feet, rapid weight gain. Common side effects of metformin and sitagliptin may include: low blood sugar (if you also use insulin or another oral diabetes medication)
- upset stomach , indigestion , gas, diarrhea , nausea, vomiting
- headache , weakness
- or cold symptoms such as runny or stuffy nose, sneezing, sore throat.

1288. This particular disease, Diabetes (Type 2), interacts with the drug Onglyza and has the following

- hives , a purple or red skin rash that spreads and causes blistering and peeling
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop taking Onglyza and call your doctor right away if you have symptoms of pancreatitis: severe pain in your upper stomach spreading to your back, nausea and vomiting , loss of appetite, or fast heartbeats. Call your doctor at once if you have: severe or ongoing pain in your joints

Drug Side Effects Summary

- pain or burning when you urinate
- or heart problems - shortness of breath (even while lying down), feeling weak or tired, rapid weight gain, swelling (especially in your feet, legs, or midsection. Common Onglyza side effects may include: painful urination
- headache
- runny or stuffy nose, sore throat , cough
- or swelling in your hands or feet.

1289. This particular disease, Diabetes (Type 2), interacts with the drug Riomet and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some people using Riomet develop lactic acidosis, which can be fatal. Get emergency medical help if you have even mild symptoms such as: unusual muscle pain
- feeling cold
- trouble breathing
- feeling dizzy, light-headed, tired, or very weak
- stomach pain, vomiting
- or slow or irregular heart rate. Common Riomet side effects may include: low blood sugar
- nausea, upset stomach
- or diarrhea .

1290. This particular disease, Diabetes (Type 2), interacts with the drug GlipiZIDE XL and has the fo

- No side effects listed

Drug Side Effects Summary

1291. This particular disease, Diabetes (Type 2), interacts with the drug Prandin and has the following

- pale or yellowed skin, dark colored urine, fever, confusion or weakness
- or pancreatitis - severe pain in your upper stomach spreading to your back, nausea and vomiting .

Common Prandin side effects may include: low blood sugar

- nausea , diarrhea
- headache , back pain
- joint pain
- or cold symptoms such as stuffy nose, sneezing, sore throat.

1292. This particular disease, Diabetes (Type 2), interacts with the drug canagliflozin and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have signs of a genital infection (penis or vagina): burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly.

Canagliflozin may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out

- little or no urination
- pain or burning when you urinate
- new pain, tenderness, sores, ulcers, or infections in your legs or feet
- high potassium-- nausea , irregular heartbeats, weakness, loss of movement
- ketoacidosis (too much acid in the blood)--nausea, vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing
- or dehydration symptoms-- dizziness , weakness, feeling light-headed (like you might pass out).

You may be more likely to have a broken bone while using canagliflozin. Talk with your doctor about

Drug Side Effects Summary

how to avoid the risk of fractures. Side effects may be more likely to occur in older adults. Common side effects of canagliflozin may include: genital infections

- or urinating more than usual.

1293. This particular disease, Diabetes (Type 2), interacts with the drug chromium picolinate and has the following side effects:

- hives

- difficulty breathing

- swelling of your face, lips, tongue, or throat. Chromium picolinate may cause serious side effects.

Stop using chromium picolinate and call your doctor at once if you have: thinking problems, trouble concentrating

- problems with balance or coordination

- or liver problems-- nausea , upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of chromium picolinate may include: headache

- sleep problems (insomnia)

- or mood changes, feeling irritable.

1294. This particular disease, Diabetes (Type 2), interacts with the drug empagliflozin and has the following side effects:

- hives

- trouble swallowing, difficult breathing

- swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have signs of a serious genital infection (penis or vagina): burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly. Stop taking empagliflozin and call your doctor at once if you have: a light-headed feeling, like you might pass out

Drug Side Effects Summary

- dehydration -- dizziness , confusion, feeling very thirsty, less urination
- ketoacidosis (too much acid in the blood)-- nausea , vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing
- or signs of a bladder infection--pain or burning when you urinate, blood in your urine, pain in pelvis or back. Some side effects may be more likely to occur in older adults. Common side effects of empagliflozin may include: a bladder infection
- or yeast infection in women (vaginal itching or discharge).

1295. This particular disease, Diabetes (Type 2), interacts with the drug exenatide and has the following side effects:

- hives , itching
- rapid heartbeats
- difficult breathing
- feeling light-headed
- swelling of your face, lips, tongue, or throat. Some people using exenatide have had serious or fatal bleeding caused by low levels of platelets (blood cells that help your blood to clot). Stop using Byetta and call your doctor right away if you have unusual bleeding or bruising. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: pancreas or gallbladder problems--pain in your upper stomach spreading to your back, nausea and vomiting, fever, fast heart rate , yellowing of your skin or eyes
- low blood sugar--headache, hunger, sweating, irritability, dizziness, nausea, fast heart rate, and feeling anxious or shaky
- or kidney problems--little or no urination, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath. Common side effects of exenatide may include: indigestion , nausea, vomiting
- diarrhea , constipation

Drug Side Effects Summary

- or itching or a small bump where the medicine was injected.

1296. This particular disease, Diabetes (Type 2), interacts with the drug repaglinide and has the following side effects:

- pale or yellowed skin, dark colored urine, fever, confusion or weakness
- or pancreatitis --severe pain in your upper stomach spreading to your back, nausea and vomiting .

Common side effects of repaglinide may include: low blood sugar

- nausea , diarrhea
- headache , back pain
- joint pain
- or cold symptoms such as stuffy nose, sneezing, sore throat.

1297. This particular disease, Diabetes (Type 2), interacts with the drug acarbose and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Acarbose may cause serious side effects. Call your doctor at once if you have: severe constipation
- severe stomach pain, diarrhea that is watery or bloody
- easy bruising, unusual bleeding (nose, mouth, vagina, or rectum), purple or red pinpoint spots under your skin
- or liver problems-- nausea , upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of acarbose may include: stomach discomfort, gas, bloating
- mild diarrhea
- or mild skin rash or itching.

Drug Side Effects Summary

1298. This particular disease, Diabetes (Type 2), interacts with the drug alogliptin and has the following side effects:

- severe pain in your upper stomach spreading to your back, nausea and vomiting , loss of appetite, or fast heartbeats. Alogliptin may cause serious side effects. Call your doctor at once if you have: severe or ongoing pain in your joints
- heart problems--shortness of breath (even while lying down), rapid weight gain, swelling (especially in your feet, legs, or midsection)
- liver problems-- nausea , upper stomach pain, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or a severe autoimmune reaction--itching, blisters, breakdown of the outer layer of skin. Common side effects of alogliptin may include: headache
- or cold symptoms such as stuffy nose, sinus pain, sore throat.

1299. This particular disease, Diabetes (Type 2), interacts with the drug Avandia and has the following side effects:

- nausea , upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, or jaundice (yellowing of the skin or eyes). Call your doctor at once if you have: shortness of breath (especially when lying down), unusual tiredness, swelling, rapid weight gain
- chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- pale skin, feeling light-headed or short of breath, cold hands and feet
- changes in your vision
- or sudden unusual pain in your hand, arm, or foot. Common Avandia side effects may include: headache
- or cold symptoms such as stuffy nose, sinus pain, sneezing, sore throat.

1300. This particular disease, Diabetes (Type 2), interacts with the drug glimepiride / pioglitazone and has the following side effects:

- pink or red urine, painful or difficult urination, new or worsening urge to urinate

Drug Side Effects Summary

- changes in your vision
- sudden unusual pain in your hand, arm, or foot
- liver problems--upper stomach pain, vomiting , tiredness, loss of appetite, dark urine, jaundice (yellowing of the skin or eyes)
- or symptoms of heart failure--shortness of breath (even while lying down), swelling in your legs or feet, rapid weight gain. Common side effects of glimepiride and pioglitazone may include: headache
- muscle pain
- nausea, diarrhea
- or cold symptoms such as stuffy nose, sinus pain, sneezing, sore throat.

1301. This particular disease, Diabetes (Type 2), interacts with the drug Glucotrol XL and has the fo

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Anxiety blurred vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chills cold sweats coma confusion cool, pale skin depression difficulty with moving dizziness fainting fast heartbeat headache increased hunger joint pain leg cramps muscle aching or cramping muscle pain or stiffness nausea nervousness nightmares pain in the joints problems in urination or increase in the amount of urine seizures shakiness slurred speech sweating swollen joints unusual tiredness or weakness Rare Abdominal or stomach pain bloating bloody or black, tarry stools body aches or pain burning, dry, or itching eyes clay-colored stools congestion constipation cough dark urine decreased vision or other changes in vision diarrhea difficult or labored breathing difficult or painful urination dryness or soreness of the throat excessive tearing fainting fast, slow, or irregular heartbeat feeling of warmth fever heartburn hoarseness indigestion itching loss of appetite pain in the eye pounding in the ears rash redness of the face, neck, arms and occasionally, upper chest redness, pain, or

Drug Side Effects Summary

swelling of the eye, eyelid, or inner lining of the eyelid runny nose severe stomach pain shortness of breath tender, swollen glands in the neck tightness in the chest trouble in swallowing unpleasant breath odor voice changes vomiting of blood or material that looks like coffee grounds wheezing yellow eyes or skin Incidence not known Agitation back or leg pains bleeding gums blood in the urine or stools chest pain convulsions decreased urine output fluid-filled skin blisters general body swelling general feeling of tiredness or weakness high fever hostility increased thirst irritability lethargy light-colored stools lower back or side pain muscle twitching nosebleeds pinpoint red pots on the skin rapid weight gain sensitivity to the sun skin thinness sores, ulcers, or white spots on the lips or in the mouth stupor swelling of the face, ankles, or hands unusual bleeding or bruising Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Indigestion passing of gas Less common Acid or sour stomach belching excess air or gas in the stomach or intestines full feeling pain sleeplessness sneezing stuffy nose trouble sleeping unable to sleep Rare Decreased interest in sexual intercourse dizziness or lightheadedness excessive muscle tone feeling of constant movement of self or surroundings flushing or redness of the skin headache, severe and throbbing hives or welts inability to have or keep an erection loss in sexual ability, desire, drive, or performance mood or mental changes muscle stiffness muscle tension or tightness sensation of spinning skin rash, encrusted, scaly, and oozing sleepiness or unusual drowsiness unusually warm skin walking in unusual manner weight loss Incidence not known Increased sensitivity of the skin to sunlight severe sunburn Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Glucotrol XL side effects

Drug Side Effects Summary

1302. This particular disease, Diabetes (Type 2), interacts with the drug glyburide / metformin and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Glyburide and metformin may cause serious side effects. Call your doctor at once if you have: heart problems-- swelling, rapid weight gain, feeling short of breath
- severe hypoglycemia--extreme weakness, blurred vision, sweating, trouble speaking, tremors, stomach pain, confusion, seizure
- or lactic acidosis--unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness, feeling cold, or feeling very weak or tired. Common side effects of glyburide and metformin may include: low blood sugar
- nausea , diarrhea, upset stomach
- or headache .

1303. This particular disease, Diabetes (Type 2), interacts with the drug Kombiglyze XR and has the following side effects:

- hives , itching, peeling skin
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop taking this medicine and call your doctor right away if you have symptoms of pancreatitis : severe pain in your upper stomach spreading to your back, nausea and vomiting , loss of appetite, or fast heartbeats. Some people using metformin develop lactic acidosis, which can be fatal. Get emergency medical help if you have even mild symptoms such as: unusual muscle pain
- feeling cold
- trouble breathing
- feeling dizzy, light-headed, tired, or very weak

Drug Side Effects Summary

- stomach pain, vomiting
- or irregular heart rate. Call your doctor at once if you have any of these serious side effects: severe autoimmune reaction - itching, blisters, breakdown of the outer layer of skin
- severe or ongoing pain in your joints
- pain or burning when you urinate
- or symptoms of heart failure - shortness of breath (even while lying down), swelling in your legs or feet, rapid weight gain . Common Kombiglyze XR side effects may include: diarrhea , nausea
- headache
- or cold symptoms such as runny or stuffy nose, sneezing, sore throat .

1304. This particular disease, Diabetes (Type 2), interacts with the drug linagliptin and has the following side effects:

- severe pain in your upper stomach spreading to your back, with or without vomiting. Linagliptin may cause serious side effects. Call your doctor at once if you have: severe or ongoing pain in your joints
- a severe autoimmune reaction--itching, blisters, breakdown of the outer layer of skin
- or symptoms of heart failure--shortness of breath (even while lying down), swelling in your legs or feet, rapid weight gain. Common side effects of linagliptin may include: runny or stuffy nose, sore throat
- cough
- or diarrhea .

1305. This particular disease, Diabetes (Type 2), interacts with the drug Xigduo XR and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have signs of

Drug Side Effects Summary

a genital infection (penis or vagina): burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly.

Call your doctor at once if you have: little or no urination

- signs of a bladder infection - pain or burning when you urinate, increased urination, blood in your urine, fever, pain in your pelvis or back
- dehydration symptoms - dizziness, weakness, feeling light-headed (like you might pass out)
- ketoacidosis (too much acid in the blood) - nausea, vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing
- or lactic acidosis - unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness, feeling cold, or feeling very weak or tired Side effects may be more likely to occur in older adults. Common Xigduo XR side effects may include: genital infections
- headache
- nausea, vomiting, diarrhea
- or runny or stuffy nose, sore throat .

1306. This particular disease, Diabetes (Type 2), interacts with the drug Adlyxin and has the following

- hives , itching, severe rash
- rapid heartbeats
- trouble swallowing
- difficult breathing
- feeling light-headed
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: pancreatitis--severe pain in your upper stomach spreading to your back, nausea and vomiting , fast heart rate
- low blood sugar-- headache , hunger, sweating, irritability, dizziness , nausea, fast heart rate, and

Drug Side Effects Summary

feeling anxious or shaky

- or kidney problems--little or no urination, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath. Common Adlyxin side effects may include: nausea, vomiting, diarrhea
- headache
- dizziness
- or low blood sugar.

1307. This particular disease, Diabetes (Type 2), interacts with the drug bromocriptine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Bromocriptine may cause serious side effects. Call your doctor at once if you have: numbness , pain, and paleness or discoloration in your fingers or toes (especially in cold weather)
- vision problems, constant runny nose
- chest pain, pain when you breathe, fast heart rate , rapid breathing, feeling short of breath (especially when lying down)
- back pain , swelling in your ankles or feet, urinating less than usual or not at all
- confusion, hallucinations, feeling like you might pass out
- muscle movements you cannot control, loss of balance or coordination
- or bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Parlodel may cause you to fall asleep during normal daytime activities such as working, talking, eating, or driving. You may fall asleep suddenly, even after feeling alert. Tell your doctor if you have any problems with daytime sleepiness or drowsiness. You may have increased sexual urges, unusual urges to gamble, or other intense urges while taking this medicine. Talk with your doctor if this occurs. Common side

Drug Side Effects Summary

effects of bromocriptine may include: dizziness

- headache
- hallucinations
- nausea, constipation
- or abnormal involuntary movements.

1308. This particular disease, Diabetes (Type 2), interacts with the drug Cycloset and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe drowsiness
- or a light-headed feeling, like you might pass out. You may have increased sexual urges, unusual urges to gamble, or other intense urges while taking this medicine. Talk with your doctor if this occurs. Common Cycloset side effects may include: dizziness, feeling tired
- nausea, vomiting
- or headache .

1309. This particular disease, Diabetes (Type 2), interacts with the drug dapagliflozin / metformin and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have signs of a genital infection (penis or vagina): burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly. Dapagliflozin and metformin may cause serious side effects. Call your doctor at once if you have: little or no urination

Drug Side Effects Summary

- signs of a bladder infection--pain or burning when you urinate, increased urination, blood in your urine, fever, pain in your pelvis or back
- dehydration symptoms--dizziness, weakness, feeling light-headed (like you might pass out)
- ketoacidosis (too much acid in the blood)--nausea, vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing
- or lactic acidosis--unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness, feeling cold, or feeling very weak or tired. Side effects may be more likely to occur in older adults. Common side effects of dapagliflozin and metformin may include: genital infections
- headache
- nausea, vomiting, diarrhea
- or runny or stuffy nose, sore throat .

1310. This particular disease, Diabetes (Type 2), interacts with the drug DiaBeta and has the following

- dark urine, jaundice (yellowing of the skin or eyes)
- severe skin rash, redness, or itching
- pale skin, easy bruising or bleeding
- fever, chills, sore throat, mouth sores
- or low levels of sodium in the body-- headache , confusion, slurred speech, severe weakness, vomiting , loss of coordination, feeling unsteady. Older adults may be more likely to have low blood sugar while taking DiaBeta. Common side effects of DiaBeta may include: low blood sugar
- nausea, heartburn , feeling full
- muscle or joint pain
- blurred vision
- or mild rash or skin redness.

Drug Side Effects Summary

1311. This particular disease, Diabetes (Type 2), interacts with the drug empagliflozin / metformin a

- hives
- trouble swallowing, difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have signs of a serious genital infection (penis or vagina): burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly. Stop taking empagliflozin and metformin and call your doctor at once if you have: a light-headed feeling, like you might pass out
- dehydration -- dizziness , confusion, feeling very thirsty, less urination
- ketoacidosis (too much acid in the blood)-- nausea , vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing
- lactic acidosis--unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness, feeling cold, or feeling very weak or tired
- or signs of a bladder infection--pain or burning when you urinate, blood in your urine, pain in pelvis or back. Some side effects may be more likely to occur in older adults. Common side effects of empagliflozin and metformin may include: low blood sugar
- indigestion , stomach pain, gas, nausea, vomiting, diarrhea
- a bladder infection
- yeast infection in women (vaginal itching or discharge)
- headache , weakness
- or runny or stuffy nose, sore throat .

1312. This particular disease, Diabetes (Type 2), interacts with the drug glipizide / metformin and ha

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. glipizide and metformin may cause serious side effects. Stop using glipizide and metformin and call your doctor at once if you have: heart problems-- swelling, rapid weight gain, feeling short of breath
- severe hypoglycemia--extreme weakness, blurred vision, sweating, trouble speaking, tremors, stomach pain, confusion, seizure
- or lactic acidosis--unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness, feeling cold, or feeling very weak or tired. Common side effects of glipizide and metformin may include: upset stomach , nausea , diarrhea
- headache
- or cold symptoms such as stuffy nose, sneezing, sore throat .

1313. This particular disease, Diabetes (Type 2), interacts with the drug Glucovance and has the fol

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: heart problems - swelling, rapid weight gain, feeling short of breath
- severe hypoglycemia - extreme weakness, blurred vision, sweating, trouble speaking, tremors, stomach pain, confusion, seizure
- or lactic acidosis - unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness, feeling cold, or feeling very weak or tired. Common Glucovance side effects may include: low blood sugar
- nausea, diarrhea, upset stomach
- or headache.

1314. This particular disease, Diabetes (Type 2), interacts with the drug Glycron and has the followi

Drug Side Effects Summary

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur:

Less common

Difficulty with swallowing
dizziness
fast heartbeat
hives, itching, skin rash
puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue
tightness in the chest
trouble breathing
unusual tiredness or weakness

Rare

Abdominal or stomach pain
chills
clay-colored stools
dark urine
diarrhea
fever
headache
light-colored stools
loss of appetite
nausea and vomiting
unpleasant breath odor
upper right abdominal or stomach pain
vomiting of blood
yellow eyes and skin

Incidence not known

Agitation
back, leg, or stomach pains
bleeding gums
blood in the urine or stools
bloody, black, or tarry stools
blurred vision
change in near or distance vision
chest pain
coma
confusion
convulsions
cough or hoarseness
decreased urine output
depression
difficulty in focusing eyes
fast or irregular heartbeat
fluid-filled skin blisters
general body swelling
high fever
hostility
increased thirst
irritability
large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs
lethargy
lower back or side pain
muscle pain or cramps
muscle twitching
nosebleeds
painful or difficult urination
pale skin
pinpoint red spots on the skin
rapid weight gain
sensitivity to the sun
skin thinness
sore throat
sores, ulcers, or white spots on the lips or in the mouth
stupor
swelling of the face, ankles, or hands
swollen or painful glands
unusual bleeding or bruising

Symptoms of overdose

Anxiety
cold sweats
cool, pale skin
increased hunger
nervousness
nightmares
shakiness
slurred speech

Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them:

Less common

Indigestion
passing of gas

Incidence not known

Difficulty with moving
joint pain
redness or other discoloration of the skin
severe sunburn
swollen joints

Other side effects not listed may also occur in some patients. If you

Drug Side Effects Summary

notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Glycron side effects

1315. This particular disease, Diabetes (Type 2), interacts with the drug Glynase and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

- itching

- red, swollen, blistered, or peeling skin with or without fever

- wheezing

- tightness in the chest or throat

- trouble breathing, swallowing, or talking

- unusual hoarseness

- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of low sodium levels like headache , trouble focusing, memory problems, feeling confused, weakness, seizures , or change in balance. Change in eyesight. Muscle or joint pain. Low blood sugar can happen. The chance may be raised when Glynase (glyburide) is used with other drugs for diabetes. Signs may be dizziness , headache, feeling sleepy or weak, shaking, fast heartbeat , confusion, hunger, or sweating. Call your doctor right away if you have any of these signs. Follow what you have been told to do for low blood sugar. This may include taking glucose tablets, liquid glucose, or some fruit juices. The ability of your bone marrow to make blood cells may be lowered. This can lead to very bad bleeding problems or infections. Tell your doctor right away if

Drug Side Effects Summary

you have signs of infection like fever, chills, or sore throat

- any bruising or bleeding
- or if you feel very tired or weak. Glyxase side effects

1316. This particular disease, Diabetes (Type 2), interacts with the drug Glyset and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Glyset may cause serious side effects. Call your doctor at once if you have: severe diarrhea or constipation
- bloody or tarry stools
- rectal bleeding
- or diarrhea that contains blood or mucus. Common side effects of Glyset may include: stomach discomfort
- diarrhea
- gas
- or mild rash.

1317. This particular disease, Diabetes (Type 2), interacts with the drug Janumet XR and has the following

- severe pain in your upper stomach spreading to your back, nausea and vomiting , loss of appetite, or fast heartbeats. Some people using metformin develop lactic acidosis, which can be fatal. Get emergency medical help if you have even mild symptoms such as: unusual muscle pain
- feeling cold
- trouble breathing
- feeling dizzy, light-headed, tired, or very weak
- stomach pain, vomiting

Drug Side Effects Summary

- or irregular heart rate. Call your doctor at once if you have any of these serious side effects: severe autoimmune reaction--itching, blisters, breakdown of the outer layer of skin
- severe or ongoing pain in your joints
- little or no urinating
- or symptoms of heart failure --shortness of breath (even while lying down), swelling in your legs or feet, rapid weight gain. Common side effects of Janumet XR may include: low blood sugar (if you also use insulin or another oral diabetes medication)
- upset stomach , indigestion , gas, diarrhea , nausea, vomiting
- headache , weakness
- or cold symptoms such as runny or stuffy nose, sneezing, sore throat.

1318. This particular disease, Diabetes (Type 2), interacts with the drug Jentadueto and has the following side effects:

- hives , difficult breathing, swelling in your face or throat) or a severe skin reaction (fever, sore throat , burning eyes, skin pain, red or purple skin rash with blistering and peeling). Stop taking this medicine and call your doctor right away if you have symptoms of pancreatitis : severe pain in your upper stomach spreading to your back, nausea and vomiting , loss of appetite, or fast heartbeats. Mild symptoms of lactic acidosis may worsen over time , and this condition can be fatal. Get emergency medical help if you have: unusual muscle pain, trouble breathing, stomach pain, vomiting, fast/slow or irregular heartbeats, dizziness, feeling cold, or feeling very weak or tired. Call your doctor at once if you have: a severe autoimmune reaction - itching, blisters, breakdown of the outer layer of skin
- severe or ongoing pain in your joints
- or symptoms of heart failure - shortness of breath (even while lying down), swelling in your legs or feet, rapid weight gain. Common Jentadueto side effects include: sore throat
- sinus pain, stuffy nose

Drug Side Effects Summary

- or diarrhea .

1319. This particular disease, Diabetes (Type 2), interacts with the drug lixisenatide and has the following side effects:

- hives , itching, severe rash
- rapid heartbeats
- trouble swallowing
- difficult breathing
- feeling light-headed
- swelling of your face, lips, tongue, or throat. lixisenatide may cause serious side effects. Stop using lixisenatide and call your doctor at once if you have: pancreatitis--severe pain in your upper stomach spreading to your back, nausea and vomiting , fast heart rate
- low blood sugar-- headache , hunger, sweating, irritability, dizziness , nausea, fast heart rate, and feeling anxious or shaky
- or kidney problems--little or no urination, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath. Common side effects of lixisenatide may include: nausea, vomiting, diarrhea
- headache
- dizziness
- or low blood sugar.

1320. This particular disease, Diabetes (Type 2), interacts with the drug metformin / pioglitazone and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some people using metformin and pioglitazone develop lactic acidosis, which can be fatal. Get emergency medical help if you have even mild

Drug Side Effects Summary

symptoms such as: unusual muscle pain

- feeling cold
- trouble breathing
- feeling dizzy, light-headed, tired, or very weak
- stomach pain, vomiting
- or slow or irregular heart rate. Metformin and pioglitazone may cause serious side effects. Call your doctor at once if you have: pink or red urine, painful or difficult urination, new or worsening urge to urinate
- changes in your vision
- heart problems-- swelling, rapid weight gain, feeling short of breath
- liver problems--loss of appetite, nausea , vomiting, stomach pain, tiredness, dark urine, jaundice (yellowing of the skin or eyes)
- or low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Common side effects of metformin and pioglitazone may include: headache
- weight gain
- diarrhea
- or cold symptoms such as stuffy nose, sinus pain, sneezing, sore throat .

1321. This particular disease, Diabetes (Type 2), interacts with the drug metformin / saxagliptin and

- hives , itching, peeling skin
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop taking metformin and saxagliptin and call your doctor right away if you have symptoms of pancreatitis : severe pain in your upper stomach spreading to your back, nausea and vomiting , loss of appetite, or fast heartbeats. Some people

Drug Side Effects Summary

using metformin develop lactic acidosis, which can be fatal. Get emergency medical help if you have even mild symptoms such as: unusual muscle pain

- feeling cold
- trouble breathing
- feeling dizzy, light-headed, tired, or very weak
- stomach pain, vomiting
- or irregular heart rate. Call your doctor at once if you have any of these serious side effects: severe autoimmune reaction--itching, blisters, breakdown of the outer layer of skin
- severe or ongoing pain in your joints
- pain or burning when you urinate
- or symptoms of heart failure --shortness of breath (even while lying down), swelling in your legs or feet, rapid weight gain. Common side effects of metformin and saxagliptin may include: diarrhea , nausea
- headache
- or cold symptoms such as runny or stuffy nose, sneezing, sore throat .

1322. This particular disease, Diabetes (Type 2), interacts with the drug miglitol and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Miglitol may cause serious side effects. Call your doctor at once if you have: severe diarrhea or constipation
- bloody or tarry stools
- rectal bleeding
- or diarrhea that contains blood or mucus. Common side effects of miglitol may include: stomach discomfort

Drug Side Effects Summary

- diarrhea
- gas
- or mild rash.

1323. This particular disease, Diabetes (Type 2), interacts with the drug Nesina and has the following

- hives, difficult breathing, swelling in your face or throat) or a severe skin reaction (fever, sore throat, burning eyes, skin pain, red or purple skin rash with blistering and peeling). Stop taking Nesina and call your doctor right away if you have symptoms of pancreatitis: severe pain in your upper stomach spreading to your back, nausea and vomiting, loss of appetite, or fast heartbeats. Call your doctor at once if you have: severe or ongoing pain in your joints
- heart problems - shortness of breath (even while lying down), rapid weight gain, swelling (especially in your feet, legs, or midsection)
- liver problems - nausea, upper stomach pain, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or a severe autoimmune reaction - itching, blisters, breakdown of the outer layer of skin. Common Nesina side effects may include: headache
- or cold symptoms such as stuffy nose, sinus pain, sore throat.

1324. This particular disease, Diabetes (Type 2), interacts with the drug saxagliptin and has the following

- hives , a purple or red skin rash that spreads and causes blistering and peeling
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop taking saxagliptin and call your doctor right away if you have symptoms of pancreatitis: severe pain in your upper stomach spreading to your back, nausea and vomiting , loss of appetite, or fast heartbeats. Saxagliptin may cause serious side effects. Call your doctor at once if you have: severe or ongoing pain in your joints

Drug Side Effects Summary

- pain or burning when you urinate
- or heart problems--shortness of breath (even while lying down), feeling weak or tired, rapid weight gain, swelling (especially in your feet, legs, or midsection. Common side effects of saxagliptin may include: painful urination
- headache
- runny or stuffy nose, sore throat , cough
- or swelling in your hands or feet.

1325. This particular disease, Diabetes (Type 2), interacts with the drug Synjardy and has the following

- hives
- trouble swallowing, difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have signs of a serious genital infection (penis or vagina): burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly. Stop taking Synjardy and call your doctor at once if you have: a light-headed feeling, like you might pass out
- dehydration - dizziness , confusion, feeling very thirsty, less urination
- ketoacidosis (too much acid in the blood) - nausea , vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing
- lactic acidosis - unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness, feeling cold, or feeling very weak or tired
- or signs of a bladder infection - pain or burning when you urinate, blood in your urine, pain in pelvis or back. Some side effects may be more likely to occur in older adults. Common Synjardy side effects may include: low blood sugar
- indigestion , stomach pain, gas, nausea, vomiting, diarrhea

Drug Side Effects Summary

- a bladder infection
- yeast infection in women (vaginal itching or discharge)
- headache , weakness
- or runny or stuffy nose, sore throat .

1326. This particular disease, Diabetes (Type 2), interacts with the drug Xultophy and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe nausea and vomiting
- shortness of breath (even with mild exertion)
- swelling in your feet or ankles, rapid weight gain
- signs of pancreatitis - severe pain in your upper stomach spreading to your back, nausea and vomiting, fast heart rate
- kidney problems - little or no urination, painful or difficult urination
- or low potassium level - leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common Xultophy side effects include: nausea, diarrhea
- headache
- or cold symptoms such as stuffy or runny nose , sneezing, sore throat .

1327. This particular disease, Diabetes (Type 2), interacts with the drug ActoPlus Met and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some people using Actoplus Met develop lactic

Drug Side Effects Summary

acidosis, which can be fatal. Get emergency medical help if you have even mild symptoms such as:
unusual muscle pain

- feeling cold
- trouble breathing
- feeling dizzy, light-headed, tired, or very weak
- stomach pain, vomiting
- or slow or irregular heart rate. Call your doctor at once if you have: pink or red urine, painful or difficult urination, new or worsening urge to urinate
- changes in your vision
- heart problems - swelling, rapid weight gain, feeling short of breath
- liver problems - loss of appetite, nausea, vomiting, stomach pain, tiredness, dark urine, jaundice (yellowing of the skin or eyes)
- or low red blood cells (anemia) - pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Common Actoplus Met side effects may include: headache
- weight gain
- diarrhea
- or cold symptoms such as stuffy nose, sinus pain, sneezing, sore throat .

1328. This particular disease, Diabetes (Type 2), interacts with the drug albiglutide and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bloating chills or fever darkened urine fast heartbeat loss of appetite nausea, vomiting, or indigestion pains in the stomach, side, or abdomen, possibly radiating to the back Incidence not known Anxiety, irritability, or mood changes blurred vision cold sweats confusion difficulty with breathing or swallowing dizziness headache

Drug Side Effects Summary

increased hunger shakiness skin itching, rash, or redness slurred speech swelling of the face, throat, or tongue unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Body aches or pain cough diarrhea ear congestion loss of voice muscle pain or stiffness pain in the joints pain or tenderness around the eyes and cheekbones skin reactions, warmth, or redness at the injection site sneezing stuffy or runny nose Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Albiglutide side effects

1329. This particular disease, Diabetes (Type 2), interacts with the drug alogliptin / metformin and h

- unusual muscle pain
- feeling cold
- trouble breathing
- feeling dizzy, light-headed, tired, or very weak
- stomach pain, vomiting
- or slow or irregular heart rate. Alogliptin and metformin may cause serious side effects. Call your doctor at once if you have: severe or ongoing pain in your joints
- pain or burning when you urinate
- liver problems--nausea, upper stomach pain, tiredness, loss of appetite, dark urine, jaundice (yellowing of the skin or eyes)
- or symptoms of heart failure --shortness of breath (even while lying down), swelling in your legs or

Drug Side Effects Summary

feet, rapid weight gain. Common side effects of alogliptin and metformin may include: diarrhea , upset stomach

- increased blood pressure (severe headache , blurred vision, pounding in your neck or ears)
- back pain , headache
- or cold symptoms such as stuffy nose, sinus pain, sore throat.

1330. This particular disease, Diabetes (Type 2), interacts with the drug alogliptin / pioglitazone and

- nausea or vomiting, and upper stomach pain (that may spread to your back). Alogliptin and pioglitazone may cause serious side effects. Call your doctor at once if you have: a severe autoimmune reaction--itching, blisters, breakdown of the outer layer of skin
- severe or ongoing pain in your joints
- pink or red urine, burning when you urinate, new or worsening urge to urinate
- changes in your vision
- liver problems--upper stomach pain, vomiting, tiredness, loss of appetite, dark urine, jaundice (yellowing of the skin or eyes)
- or symptoms of heart failure--shortness of breath (even while lying down), swelling in your legs or feet, rapid weight gain. Common side effects of alogliptin and pioglitazone may include: back pain
- or cold symptoms such as stuffy nose, sinus pain, sore throat.

1331. This particular disease, Diabetes (Type 2), interacts with the drug Bydureon BCise and has th

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Some people using exenatide have had serious or fatal bleeding caused by low levels of platelets (blood cells that help your blood to clot). Stop using Bydureon and call your doctor right away if you have unusual bleeding or bruising. Bydureon BCise

Drug Side Effects Summary

may cause serious side effects. Stop using Bydureon BCise and call your doctor at once if you have: severe ongoing nausea and vomiting

- pain, warmth, swelling, an open wound or scab, or other skin changes where the injection was given
- easy bruising, unusual bleeding, purple or red spots under your skin
- signs of a thyroid tumor--swelling or a lump in your neck, trouble swallowing, a hoarse voice, feeling short of breath
- pancreas or gallbladder problems--pain in your upper stomach spreading to your back, nausea and vomiting, fever, fast heart rate , yellowing of your skin or eyes
- low blood sugar--headache, hunger, sweating, irritability, dizziness, fast heart rate, and feeling anxious or shaky
- or kidney problems--swelling, urinating less, feeling tired or short of breath. Common side effects of Bydureon BCise may include: indigestion , nausea, vomiting, diarrhea , constipation
- headache
- or itching or a small bump where an injection was given.

1332. This particular disease, Diabetes (Type 2), interacts with the drug canagliflozin / metformin and

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have signs of a genital infection (penis or vagina): burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly. Mild symptoms of lactic acidosis may worsen over time , and this condition can be fatal. Stop taking canagliflozin and metformin and call your doctor right away if you have unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness, feeling cold, or feeling very weak

Drug Side Effects Summary

or tired. Canagliflozin and metformin may cause serious side effects. Call your doctor at once if you have: new pain, tenderness, sores, ulcers, or infections in your legs or feet

- little or no urination
- a light-headed feeling, like you might pass out
- high potassium level--nausea, weakness, tingly feeling, chest pain, irregular heartbeats, muscle weakness
- ketoacidosis (too much acid in the blood)--nausea, vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing
- or signs of a bladder infection --pain or burning when you urinate, urine that looks cloudy, pain in pelvis or back. Older adults may be more likely to get dehydrated or have kidney problems while taking this medicine. You may be more likely to have a broken bone while using this medicine. Talk with your doctor about how to avoid the risk of fractures. Common side effects of canagliflozin and metformin may include: urinating more than usual
- headache , weakness
- gas, stomach pain, indigestion
- nausea, vomiting
- or diarrhea.

1333. This particular disease, Diabetes (Type 2), interacts with the drug Cr-GTF and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Cr-GTF may cause serious side effects. Stop using Cr-GTF and call your doctor at once if you have: thinking problems, trouble concentrating
- problems with balance or coordination
- or liver problems-- nausea , upper stomach pain, itching, tired feeling, loss of appetite, dark urine,

Drug Side Effects Summary

clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Cr-GTF may include: headache

- sleep problems (insomnia)
- or mood changes, feeling irritable.

1334. This particular disease, Diabetes (Type 2), interacts with the drug CRM and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. CRM may cause serious side effects. Stop using CRM and call your doctor at once if you have: thinking problems, trouble concentrating
- problems with balance or coordination
- or liver problems-- nausea , upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of CRM may include: headache
- sleep problems (insomnia)
- or mood changes, feeling irritable.

1335. This particular disease, Diabetes (Type 2), interacts with the drug dapagliflozin / saxagliptin a

- hives , itching, flaking or peeling skin
- trouble swallowing, difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have signs of a genital infection (penis or vagina): burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly. Stop taking dapagliflozin and saxagliptin and call your doctor at once if you have: severe or ongoing pain in your joints

Drug Side Effects Summary

- serious skin reaction--itching, blisters, breakdown of the outer layer of skin
- ketoacidosis (too much acid in the blood)--nausea, vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing
- pancreatitis --severe pain in your upper stomach spreading to your back, nausea and vomiting , fast heart rate
- dehydration symptoms-- dizziness , weakness, feeling light-headed (like you might pass out)
- or signs of a bladder infection --pain or burning when you urinate, increased urination, blood in your urine, pain in your pelvis or back. Some people taking dapagliflozin have had bladder cancer, but it is not clear if this medicine was the actual cause. Common side effects of dapagliflozin and saxagliptin may include: urination problems
- abnormal blood levels of cholesterol or triglycerides
- or runny nose , stuffy nose, sinus pain.

1336. This particular disease, Diabetes (Type 2), interacts with the drug Duetact and has the following

- pink or red urine, painful or difficult urination, new or worsening urge to urinate
- changes in your vision
- sudden unusual pain in your hand, arm, or foot
- liver problems - upper stomach pain, vomiting , tiredness, loss of appetite, dark urine, jaundice (yellowing of the skin or eyes)
- or symptoms of heart failure - shortness of breath (even while lying down), swelling in your legs or feet, rapid weight gain. Common Duetact side effects may include: headache
- muscle pain
- nausea, diarrhea
- or cold symptoms such as stuffy nose, sinus pain, sneezing, sore throat.

Drug Side Effects Summary

1337. This particular disease, Diabetes (Type 2), interacts with the drug empagliflozin / linagliptin / metformin

- hives , itching, flaking or peeling skin
- trouble swallowing, difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have signs of a serious genital infection (penis or vagina): burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly. Stop taking empagliflozin, linagliptin, and metformin and call your doctor at once if you have: a light-headed feeling, like you might pass out
- severe or ongoing pain in your joints
- lactic acidosis--unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness , feeling cold, or feeling very weak or tired
- serious skin reaction--itching, blisters, breakdown of the outer layer of skin
- ketoacidosis (too much acid in the blood)-- nausea , vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing
- pancreatitis --severe pain in your upper stomach spreading to your back, nausea and vomiting
- dehydration --dizziness, confusion, feeling very thirsty, less urination
- signs of a bladder infection--pain or burning when you urinate, blood in your urine, pain in pelvis or back
- or symptoms of heart failure--shortness of breath (even while lying down), swelling in your legs or feet, rapid weight gain. Common side effects of empagliflozin, linagliptin, and metformin may include: stomach pain, diarrhea, constipation
- a bladder infection
- headache
- or cold symptoms such as runny or stuffy nose, sneezing, sore throat .

Drug Side Effects Summary

1338. This particular disease, Diabetes (Type 2), interacts with the drug empagliflozin / linagliptin and

- hives , itching, flaking or peeling skin
 - trouble swallowing, difficult breathing
 - swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have signs of a serious genital infection (penis or vagina): burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly. Empagliflozin and linagliptin may cause serious side effects. Call your doctor at once if you have: pain and burning when you urinate, painful urination
 - a light-headed feeling, like you might pass out
 - low blood sugar-- headache , hunger, sweating, irritability, dizziness , fast heart rate , and feeling anxious or shaky
 - heart problems--swelling, rapid weight gain, feeling short of breath
 - or dehydration --dizziness, confusion, feeling very thirsty, less urination. Stop taking empagliflozin and linagliptin and call your doctor at once if you have: severe or ongoing pain in your joints
 - serious skin reaction--itching, blisters, breakdown of the outer layer of skin
 - ketoacidosis (too much acid in the blood)-- nausea , vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing
 - or pancreatitis --severe pain in your upper stomach spreading to your back, nausea and vomiting .
- Common side effects of empagliflozin and linagliptin may include: a bladder infection
- or cold symptoms such as runny or stuffy nose, sneezing, sore throat .

1339. This particular disease, Diabetes (Type 2), interacts with the drug ertugliflozin / metformin and

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have signs of

Drug Side Effects Summary

a genital infection (penis or vagina): burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly.

Ertugliflozin and metformin may cause serious side effects. Call your doctor at once if you have: new pain, tenderness, sores, ulcers, or infections in your legs or feet

- little or no urination
- lactic acidosis--unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness, feeling cold, or feeling very weak or tired
- ketoacidosis (too much acid in the blood)-- nausea , vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing
- dehydration symptoms--dizziness, weakness, feeling light-headed (like you might pass out)
- or signs of a bladder infection --pain or burning when you urinate, increased urination, blood in your urine, fever, pain in your pelvis or back. Some side effects may be more likely to occur in older adults. Common side effects of ertugliflozin and metformin may include: genital yeast infection
- stomach discomfort
- nausea, vomiting, diarrhea, gas
- headache , weakness
- or urinating more than usual.

1340. This particular disease, Diabetes (Type 2), interacts with the drug ertugliflozin / sitagliptin and

- burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly. Ertugliflozin and sitagliptin may cause serious side effects. Call your doctor at once if you have: severe autoimmune reaction--itching, blisters, breakdown of the outer layer of skin
- new pain, tenderness, sores, ulcers, or infections in your legs or feet
- severe or ongoing pain in your joints

Drug Side Effects Summary

- little or no urination
 - ketoacidosis (too much acid in the blood)-- nausea , vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing
 - dehydration symptoms-- dizziness , weakness, feeling light-headed (like you might pass out)
 - signs of a bladder infection--pain or burning when you urinate, increased urination, blood in your urine, fever, pain in your pelvis or back
 - symptoms of heart failure --shortness of breath (even while lying down), swelling in your legs or feet, rapid weight gain
 - or symptoms of pancreatitis : severe pain in your upper stomach spreading to your back, with or without nausea and vomiting. Some side effects may be more likely to occur in older adults.
- Common side effects of ertugliflozin and sitagliptin may include: urinating more than usual
- upset stomach , diarrhea
 - headache
 - or cold symptoms such as stuffy nose, sneezing, sore throat.

1341. This particular disease, Diabetes (Type 2), interacts with the drug ertugliflozin and has the fol

- hives
 - difficult breathing
 - swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have signs of a genital infection (penis or vagina): burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly.
- Ertugliflozin may cause serious side effects. Call your doctor at once if you have: new pain, tenderness, sores, ulcers, or infections in your legs or feet
- little or no urination
 - ketoacidosis (too much acid in the blood)--nausea, vomiting, stomach pain, confusion, unusual

Drug Side Effects Summary

drowsiness, or trouble breathing

- dehydration symptoms--dizziness, weakness, feeling light-headed (like you might pass out)
- or signs of a bladder infection --pain or burning when you urinate, increased urination, blood in your urine, fever, pain in your pelvis or back. Some side effects may be more likely to occur in older adults. Common side effects of ertugliflozin may include: genital yeast infections
- or urinating more than usual.

1342. This particular disease, Diabetes (Type 2), interacts with the drug Glynase PresTab and has the following side effects:

- dark urine, jaundice (yellowing of the skin or eyes)
- severe skin rash, redness, or itching
- pale skin, easy bruising or bleeding
- fever, chills, sore throat, mouth sores
- or low levels of sodium in the body-- headache , confusion, slurred speech, severe weakness, vomiting , loss of coordination, feeling unsteady. Older adults may be more likely to have low blood sugar while taking Glynase PresTab. Common side effects of Glynase PresTab may include: low blood sugar
- nausea, heartburn , feeling full
- muscle or joint pain
- blurred vision
- or mild rash or skin redness.

1343. This particular disease, Diabetes (Type 2), interacts with the drug Glyxambi and has the following side effects:

- hives , itching, flaking or peeling skin
- trouble swallowing, difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have signs of

Drug Side Effects Summary

a serious genital infection (penis or vagina): burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly. Stop taking Glyxambi and call your doctor at once if you have: pain and burning when you urinate, painful urination

- a light-headed feeling, like you might pass out
- low blood sugar - headache, hunger, sweating, irritability, dizziness, fast heart rate, and feeling anxious or shaky
- heart problems - swelling, rapid weight gain, feeling short of breath
- or dehydration - dizziness, confusion, feeling very thirsty, less urination. Stop taking this medicine and call your doctor at once if you have: severe or ongoing pain in your joints
- serious skin reaction - itching, blisters, breakdown of the outer layer of skin
- ketoacidosis (too much acid in the blood) - nausea, vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing
- or pancreatitis - severe pain in your upper stomach spreading to your back, nausea and vomiting.

Common Glyxambi side effects may include: a bladder infection

- or cold symptoms such as runny or stuffy nose, sneezing, sore throat .

1344. This particular disease, Diabetes (Type 2), interacts with the drug insulin degludec / liraglutide

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Insulin degludec and liraglutide may cause serious side effects. Call your doctor at once if you have: severe nausea and vomiting
- shortness of breath (even with mild exertion)
- swelling in your feet or ankles, rapid weight gain
- signs of pancreatitis--severe pain in your upper stomach spreading to your back, nausea and

Drug Side Effects Summary

vomiting, fast heart rate

- kidney problems--little or no urination, painful or difficult urination
- or low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of insulin degludec and liraglutide may include: nausea, diarrhea
- headache
- or cold symptoms such as stuffy or runny nose , sneezing, sore throat .

1345. This particular disease, Diabetes (Type 2), interacts with the drug Invokamet and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have signs of a genital infection (penis or vagina): burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly. Mild symptoms of lactic acidosis may worsen over time , and this condition can be fatal. Stop taking Invokamet and call your doctor right away if you have unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness, feeling cold, or feeling very weak or tired. Call your doctor at once if you have: new pain, tenderness, sores, ulcers, or infections in your legs or feet
- little or no urination
- a light-headed feeling, like you might pass out
- high potassium level - nausea, weakness, tingly feeling, chest pain, irregular heartbeats, muscle weakness
- ketoacidosis (too much acid in the blood) - nausea, vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing

Drug Side Effects Summary

- or signs of a bladder infection - pain or burning when you urinate, urine that looks cloudy, pain in pelvis or back. Older adults may be more likely to get dehydrated or have kidney problems while taking this medicine. You may be more likely to have a broken bone while using this medicine. Talk with your doctor about how to avoid the risk of fractures. Common Invokamet side effects may include: urinating more than usual

- headache , weakness
- gas, stomach pain, indigestion
- nausea, vomiting
- or diarrhea.

1346. This particular disease, Diabetes (Type 2), interacts with the drug Invokamet XR and has the f

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have signs of a genital infection (penis or vagina): burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly. Mild symptoms of lactic acidosis may worsen over time , and this condition can be fatal. Stop taking Invokamet XR and call your doctor right away if you have unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness, feeling cold, or feeling very weak or tired. Invokamet XR may cause serious side effects. Call your doctor at once if you have: new pain, tenderness, sores, ulcers, or infections in your legs or feet
- little or no urination
- a light-headed feeling, like you might pass out
- high potassium level--nausea, weakness, tingly feeling, chest pain, irregular heartbeats, muscle weakness

Drug Side Effects Summary

- ketoacidosis (too much acid in the blood)--nausea, vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing
- or signs of a bladder infection --pain or burning when you urinate, urine that looks cloudy, pain in pelvis or back. Older adults may be more likely to get dehydrated or have kidney problems while taking this medicine. You may be more likely to have a broken bone while using this medicine. Talk with your doctor about how to avoid the risk of fractures. Common side effects of Invokamet XR may include: urinating more than usual
- headache , weakness
- gas, stomach pain, indigestion
- nausea, vomiting
- or diarrhea.

1347. This particular disease, Diabetes (Type 2), interacts with the drug Jentadueto XR and has the

- severe pain in your upper stomach spreading to your back, nausea and vomiting , loss of appetite, or fast heartbeats. Mild symptoms of lactic acidosis may worsen over time , and this condition can be fatal. Get emergency medical help if you have: unusual muscle pain, trouble breathing, stomach pain, vomiting, fast/slow or irregular heartbeats, dizziness, feeling cold, or feeling very weak or tired. Jentadueto XR may cause serious side effects. Call your doctor at once if you have: a severe autoimmune reaction--itching, blisters, breakdown of the outer layer of skin
- severe or ongoing pain in your joints
- or symptoms of heart failure --shortness of breath (even while lying down), swelling in your legs or feet, rapid weight gain. Common side effects of Jentadueto XR may include: sore throat
- sinus pain, stuffy nose
- or diarrhea .

Drug Side Effects Summary

1348. This particular disease, Diabetes (Type 2), interacts with the drug Kazano and has the following

- unusual muscle pain
- feeling cold
- trouble breathing
- feeling dizzy, light-headed, tired, or very weak
- stomach pain, vomiting
- or slow or irregular heart rate. Call your doctor at once if you have: severe or ongoing pain in your joints
- pain or burning when you urinate
- liver problems - nausea, upper stomach pain, tiredness, loss of appetite, dark urine, jaundice (yellowing of the skin or eyes)
- or symptoms of heart failure - shortness of breath (even while lying down), swelling in your legs or feet, rapid weight gain . Common Kazano side effects may include: diarrhea , upset stomach
- increased blood pressure (severe headache, blurred vision, pounding in your neck or ears)
- back pain , headache
- or cold symptoms such as stuffy nose, sinus pain, sore throat.

1349. This particular disease, Diabetes (Type 2), interacts with the drug linagliptin / metformin and l

- severe pain in your upper stomach spreading to your back, nausea and vomiting , loss of appetite, or fast heartbeats. Mild symptoms of lactic acidosis may worsen over time , and this condition can be fatal. Get emergency medical help if you have: unusual muscle pain, trouble breathing, stomach pain, vomiting, fast/slow or irregular heartbeats, dizziness, feeling cold, or feeling very weak or tired. Linagliptin and metformin may cause serious side effects. Call your doctor at once if you have: a severe autoimmune reaction--itching, blisters, breakdown of the outer layer of skin
- severe or ongoing pain in your joints

Drug Side Effects Summary

- or symptoms of heart failure --shortness of breath (even while lying down), swelling in your legs or feet, rapid weight gain. Common side effects of linagliptin and metformin may include: sore throat
- sinus pain, stuffy nose
- or diarrhea .

1350. This particular disease, Diabetes (Type 2), interacts with the drug nateglinide and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Nateglinide may cause serious side effects. Call your doctor at once if you have: seizure (convulsions)
- or jaundice (yellowing of the skin or eyes). Common side effects of nateglinide may include: runny or stuffy nose, sneezing, cough, cold or flu symptoms
- diarrhea
- back pain
- dizziness
- or joint pain or stiffness.

1351. This particular disease, Diabetes (Type 2), interacts with the drug Oseni and has the following

- nausea or vomiting, and upper stomach pain (that may spread to your back). Call your doctor at once if you have: a severe autoimmune reaction - itching, blisters, breakdown of the outer layer of skin
- severe or ongoing pain in your joints
- pink or red urine, burning when you urinate, new or worsening urge to urinate
- changes in your vision
- liver problems - upper stomach pain, vomiting, tiredness, loss of appetite, dark urine, jaundice

Drug Side Effects Summary

(yellowing of the skin or eyes)

- or symptoms of heart failure - shortness of breath (even while lying down), swelling in your legs or feet, rapid weight gain. Common Oseni side effects may include: back pain
- or cold symptoms such as stuffy nose, sinus pain, sore throat.

1352. This particular disease, Diabetes (Type 2), interacts with the drug Qtern and has the following

- hives , itching, flaking or peeling skin
- trouble swallowing, difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have signs of a genital infection (penis or vagina): burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly. Stop taking this medicine and call your doctor at once if you have: severe or ongoing pain in your joints
- serious skin reaction - itching, blisters, breakdown of the outer layer of skin
- ketoacidosis (too much acid in the blood) - nausea, vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing
- pancreatitis - severe pain in your upper stomach spreading to your back, nausea and vomiting , fast heart rate
- dehydration symptoms - dizziness, weakness, feeling light-headed (like you might pass out)
- or signs of a bladder infection - pain or burning when you urinate, increased urination, blood in your urine, pain in your pelvis or back. Some people taking dapagliflozin have had bladder cancer, but it is not clear if this medicine was the actual cause. Common Qtern side effects may include: urination problems
- abnormal blood levels of cholesterol or triglycerides
- or runny nose , stuffy nose, sinus pain.

Drug Side Effects Summary

1353. This particular disease, Diabetes (Type 2), interacts with the drug rosiglitazone and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Abdominal or stomach pain blurred vision chest pain or discomfort decrease in the amount of urine dry mouth flushed, dry skin fruit-like breath odor increased hunger increased thirst increased urination irregular heartbeat nausea noisy, rattling breathing pain in the shoulders, arms, jaw, or neck pale skin rapid or unusual weight gain sweating swelling of the fingers, hands, feet, or lower legs trouble breathing unexplained weight loss unusual bleeding or bruising unusual tiredness or weakness vomiting Rare Anxiety chills cold sweats coma confusion dark urine depression dizziness fast heartbeat headache loss of appetite nightmares seizures shakiness slurred speech Incidence not known Blue lips and fingernails changes in vision coughing that sometimes produces a pink frothy sputum hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs itching or skin rash light-colored stools redness of the skin yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Ear congestion fever general feeling of discomfort or illness hoarseness or other voice changes injury joint pain muscle aches and pains runny or stuffy nose shivering sneezing sore throat trouble sleeping Less common Back pain cough diarrhea lightheadedness pain or tenderness around the eyes and cheekbones Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Rosiglitazone side effects

Drug Side Effects Summary

1354. This particular disease, Diabetes (Type 2), interacts with the drug Segluromet and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have signs of a genital infection (penis or vagina): burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly. Call your doctor at once if you have: new pain, tenderness, sores, ulcers, or infections in your legs or feet
- little or no urination
- lactic acidosis - unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness, feeling cold, or feeling very weak or tired
- ketoacidosis (too much acid in the blood) - nausea, vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing
- dehydration symptoms - dizziness, weakness, feeling light-headed (like you might pass out)
- or signs of a bladder infection - pain or burning when you urinate, increased urination, blood in your urine, fever, pain in your pelvis or back. Some side effects may be more likely to occur in older adults. Common Segluromet side effects may include: genital yeast infection
- stomach discomfort
- nausea, vomiting, diarrhea, gas
- headache, weakness
- or urinating more than usual.

1355. This particular disease, Diabetes (Type 2), interacts with the drug Starlix and has the following side effects:

- hives
- difficulty breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: seizure (convulsions)
- or jaundice (yellowing of the skin or eyes). Common Starlix side effects may include: runny or stuffy nose, sneezing, cough, cold or flu symptoms
- diarrhea: back pain
- dizziness
- or joint pain or stiffness.

1356. This particular disease, Diabetes (Type 2), interacts with the drug Steglatro and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have signs of a genital infection (penis or vagina): burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly. Call your doctor at once if you have: new pain, tenderness, sores, ulcers, or infections in your legs or feet
- little or no urination
- ketoacidosis (too much acid in the blood) - nausea, vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing
- dehydration symptoms - dizziness, weakness, feeling light-headed (like you might pass out)
- or signs of a bladder infection - pain or burning when you urinate, increased urination, blood in your urine, fever, pain in your pelvis or back. Some side effects may be more likely to occur in older adults. Common Steglatro side effects may include: genital yeast infections
- or urinating more than usual.

Drug Side Effects Summary

1357. This particular disease, Diabetes (Type 2), interacts with the drug Steglujan and has the following

- burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly. Call your doctor at once if you have:
severe autoimmune reaction - itching, blisters, breakdown of the outer layer of skin
- new pain, tenderness, sores, ulcers, or infections in your legs or feet
- severe or ongoing pain in your joints
- little or no urination
- ketoacidosis (too much acid in the blood) - nausea, vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing
- dehydration symptoms - dizziness, weakness, feeling light-headed (like you might pass out)
- signs of a bladder infection - pain or burning when you urinate, increased urination, blood in your urine, fever, pain in your pelvis or back
- symptoms of heart failure - shortness of breath (even while lying down), swelling in your legs or feet, rapid weight gain
- or symptoms of pancreatitis - severe pain in your upper stomach spreading to your back, with or without nausea and vomiting. Some side effects may be more likely to occur in older adults.

Common Steglujan side effects may include: urinating more than usual

- upset stomach , diarrhea
- headache
- or cold symptoms such as stuffy nose, sneezing, sore throat.

1358. This particular disease, Diabetes (Type 2), interacts with the drug Synjardy XR and has the following

- hives
- trouble swallowing, difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have signs of

Drug Side Effects Summary

a serious genital infection (penis or vagina): burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly. Stop taking Synjardy XR and call your doctor at once if you have: a light-headed feeling, like you might pass out

- dehydration -- dizziness , confusion, feeling very thirsty, less urination
- ketoacidosis (too much acid in the blood)-- nausea , vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing
- lactic acidosis--unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness, feeling cold, or feeling very weak or tired
- or signs of a bladder infection--pain or burning when you urinate, blood in your urine, pain in pelvis or back. Some side effects may be more likely to occur in older adults. Common side effects of Synjardy XR may include: low blood sugar
- indigestion , stomach pain, gas, nausea, vomiting, diarrhea
- a bladder infection
- yeast infection in women (vaginal itching or discharge)
- headache , weakness
- or runny or stuffy nose, sore throat .

1359. This particular disease, Diabetes (Type 2), interacts with the drug Tanzeum and has the following

- hives
- fast heartbeats, feeling light-headed
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: symptoms of pancreatitis - severe pain in your upper stomach spreading to your back, nausea and vomiting , fast heart rate

Drug Side Effects Summary

- signs of a thyroid tumor - swelling or a lump in your neck, trouble swallowing, a hoarse voice, or if you feel short of breath
- low blood sugar - headache , hunger, weakness, sweating, confusion, irritability, dizziness , fast heart rate, or feeling jittery
- or kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath. Common Tanzeum side effects may include: nausea , diarrhea
- cough, cold or flu symptoms
- back pain , joint pain
- or pain, swelling, or irritation where medicine was injected.

1360. This particular disease, Diabetes (Type 2), interacts with the drug tirzepatide and has the following

- Stop using tirzepatide and get emergency medical help if you have: signs of an allergic reaction: hives
 - difficult breathing
 - feeling light-headed
 - swelling of your face, lips, tongue, or throat
 - or pancreatitis --severe pain in your upper stomach spreading to your back, nausea and vomiting .
- Tirzepatide may cause serious side effects. Call your doctor at once if you have: severe stomach problems
- changes in your vision
 - signs of a thyroid tumor--swelling or a lump in your neck, trouble swallowing, a hoarse voice, or if you feel short of breath
 - gallbladder problem--chalky-colored stools, stomach pain after eating, nausea , heartburn , bloating , and severe upper stomach pain that may spread to your back
 - low blood sugar-- headache , hunger, weakness, sweating, confusion, irritability, dizziness , fast

Drug Side Effects Summary

heart rate , or feeling jittery

- or kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath. Common side effects of tirzepatide may include: nausea, diarrhea , decreased appetite, vomiting

- or constipation , indigestion , or stomach pain.

1361. This particular disease, Diabetes (Type 2), interacts with the drug Trijardy XR and has the following symptoms:

- hives , itching, flaking or peeling skin

- trouble swallowing, difficult breathing

- swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have signs of a serious genital infection (penis or vagina): burning, itching, odor, discharge, pain, tenderness, redness or swelling of the genital or rectal area, fever, not feeling well. These symptoms may get worse quickly. Stop taking Trijardy XR and call your doctor at once if you have: a light-headed feeling, like you might pass out

- severe or ongoing pain in your joints

- lactic acidosis - unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness , feeling cold, or feeling very weak or tired

- serious skin reaction - itching, blisters, breakdown of the outer layer of skin

- ketoacidosis (too much acid in the blood) - nausea , vomiting, stomach pain, confusion, unusual drowsiness, or trouble breathing

- pancreatitis - severe pain in your upper stomach spreading to your back, nausea and vomiting

- dehydration - dizziness, confusion, feeling very thirsty, less urination

- signs of a bladder infection - pain or burning when you urinate, blood in your urine, pain in pelvis or back

- or symptoms of heart failure - shortness of breath (even while lying down), swelling in your legs or

Drug Side Effects Summary

feet, rapid weight gain. Common Trijardy XR side effects may include: stomach pain, diarrhea, constipation

- a bladder infection
- headache
- or cold symptoms such as runny or stuffy nose, sneezing, sore throat .

1362. This particular disease, Diarrhea, interacts with the drug loperamide and has the following side effects:

- diarrhea that is watery or bloody
- stomach pain or bloating
- ongoing or worsening diarrhea
- or fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out). Common side effects of loperamide may include: constipation
- dizziness, drowsiness
- nausea
- or stomach cramps.

1363. This particular disease, Diarrhea, interacts with the drug Lomotil and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some side effects may occur up to 30 hours after you take this medicine. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe constipation , stomach pain or bloating
- ongoing or worsening diarrhea
- diarrhea that is watery or bloody
- severe pain in your upper stomach spreading to your back

Drug Side Effects Summary

- fever, flushing (warmth, redness, or tingly feeling)
- hallucinations, seizure
- rapid breathing, weak or shallow breathing
- fast heart rate
- or dehydration symptoms--feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin . Common side effects of Lomotil may include: drowsiness, dizziness , feeling restless
- headache
- numbness in your hands or feet
- depression , not feeling well
- confusion, feelings of extreme happiness
- red or swollen gums
- dry mouth, nose, or throat
- nausea , vomiting , upset stomach , loss of appetite
- or skin rash , dryness, or itching.

1364. This particular disease, Diarrhea, interacts with the drug Imodium and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bloating constipation loss of appetite nausea stomach pain vomiting Incidence not known Blistering, peeling, loosening of skin chest pain or discomfort chills cough decrease in urine volume decrease in frequency of urination difficulty in passing urine diarrhea fainting fast, pounding, or irregular heartbeat or pulse hives or welts, itching, rash irregular or slow heartbeat joint or muscle pain large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, sex organs no blood pressure or pulse painful

Drug Side Effects Summary

urination palpitations red irritated eyes redness of the skin red skin lesions, often with a purple center sores, ulcers, or white spots in the mouth or on the lips sore throat stopping of heart unconsciousness unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Rare Acid or sour stomach belching dizziness drowsiness dryness of mouth excess air or gas in the stomach or bowels full feeling heartburn indigestion passing gas stomach upset Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Imodium side effects

1365. This particular disease, Diarrhea, interacts with the drug atropine / diphenoxylate and has the

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some side effects may occur up to 30 hours after you take this medicine. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe constipation , stomach pain or bloating
- ongoing or worsening diarrhea
- diarrhea that is watery or bloody
- severe pain in your upper stomach spreading to your back
- fever, flushing (warmth, redness, or tingly feeling)
- hallucinations, seizure
- rapid breathing, weak or shallow breathing

Drug Side Effects Summary

- fast heart rate
- or dehydration symptoms--feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin . Common side effects of atropine and diphenoxylate may include: drowsiness, dizziness , feeling restless
- headache
- numbness in your hands or feet
- depression , not feeling well
- confusion, feelings of extreme happiness
- red or swollen gums
- dry mouth, nose, or throat
- nausea , vomiting , upset stomach , loss of appetite
- or skin rash , dryness, or itching.

1366. This particular disease, Diarrhea, interacts with the drug Imodium A-D and has the following side effects:

- diarrhea that is watery or bloody
- stomach pain or bloating
- ongoing or worsening diarrhea
- or fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out). Common Imodium A-D side effects may include: constipation
- dizziness, drowsiness
- nausea
- or stomach cramps.

1367. This particular disease, Diarrhea, interacts with the drug Anti-Diarrheal and has the following side effects:

- No side effects listed

Drug Side Effects Summary

1368. This particular disease, Diarrhea, interacts with the drug Pepto-Bismol and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop taking this medicine and call your doctor at once if you have: changes in behavior with nausea and vomiting
- hearing loss or ringing in your ears
- diarrhea lasting longer than 2 days
- or worsened stomach symptoms. Bismuth subsalicylate can cause you to have a black or darkened tongue. This is a harmless side effect. Common side effects include: constipation
- or dark colored stools.

1369. This particular disease, Diarrhea, interacts with the drug Florastor and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Although not all side effects are known, Florastor is thought to be likely safe for most people when used as directed for up to 15 months. Common side effects of Florastor may include: gas, bloating
- or constipation .

1370. This particular disease, Diarrhea, interacts with the drug codeine and has the following side e

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Like other narcotic medicines, codeine can slow your breathing. Death may occur if breathing becomes too weak. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you

Drug Side Effects Summary

are hard to wake up. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep

- a slow heart rate or weak pulse
- a light-headed feeling, like you might pass out
- confusion, agitation , hallucinations, unusual thoughts or behavior
- feelings of extreme happiness or sadness
- seizure (convulsions)
- problems with urination
- or low cortisol levels--nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever , sweating , shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea . Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent. Common codeine side effects include: feeling dizzy or drowsy
- nausea, vomiting, stomach pain
- constipation
- sweating
- or mild itching or rash .

1371. This particular disease, Diarrhea, interacts with the drug bismuth subsalicylate and has the fo

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop taking bismuth subsalicylate and call your doctor

Drug Side Effects Summary

at once if you have: changes in behavior with nausea and vomiting

- hearing loss or ringing in your ears
- diarrhea lasting longer than 2 days
- or worsened stomach symptoms. Bismuth subsalicylate can cause you to have a black or darkened tongue. This is a harmless side effect. Common side effects include: constipation
- or dark colored stools.

1372. This particular disease, Diarrhea, interacts with the drug Diamode and has the following side

- diarrhea that is watery or bloody
- stomach pain or bloating
- ongoing or worsening diarrhea
- or fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out). Common side effects of Diamode may include: constipation
- dizziness, drowsiness
- nausea
- or stomach cramps.

1373. This particular disease, Diarrhea, interacts with the drug Kaopectate and has the following side

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop taking Kaopectate and call your doctor at once if you have: changes in behavior with nausea and vomiting
- hearing loss or ringing in your ears
- diarrhea lasting longer than 2 days
- or worsened stomach symptoms. Kaopectate can cause you to have a black or darkened tongue.

Drug Side Effects Summary

This is a harmless side effect. Common side effects include: constipation

- or dark colored stools.

1374. This particular disease, Diarrhea, interacts with the drug rifaximin and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Rifaximin may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- fever
- or fluid build-up around the stomach--rapid weight gain, stomach pain and bloating , trouble breathing while lying down. Common side effects of rifaximin may include: swelling in your hands or feet
- nausea
- headache , dizziness
- tiredness
- or abnormal liver function tests .

1375. This particular disease, Diarrhea, interacts with the drug Up and Up Anti-Diarrheal Solution and

- No side effects listed

1376. This particular disease, Diarrhea, interacts with the drug neomycin and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Neomycin may cause serious side effects. Call your

Drug Side Effects Summary

doctor at once if you have: numbness or tingly feeling under your skin

- hearing problems (even long after you stop taking neomycin)
- muscle twitching , seizure
- weak or shallow breathing
- or kidney problems--swelling, urinating less, feeling tired or short of breath. Common side effects of neomycin may include: nausea , vomiting
- or diarrhea .

1377. This particular disease, Diarrhea, interacts with the drug saccharomyces boulardii lyo and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Although not all side effects are known, saccharomyces boulardii lyo is thought to be likely safe for most people when used as directed for up to 15 months. Common side effects of saccharomyces boulardii lyo may include: gas, bloating
- or constipation .

1378. This particular disease, Diarrhea, interacts with the drug Bismatrol and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop taking Bismatrol and call your doctor at once if you have: changes in behavior with nausea and vomiting
- hearing loss or ringing in your ears
- diarrhea lasting longer than 2 days
- or worsened stomach symptoms. Bismatrol can cause you to have a black or darkened tongue.

This is a harmless side effect. Common side effects include: constipation

Drug Side Effects Summary

- or dark colored stools.

1379. This particular disease, Diarrhea, interacts with the drug opium and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Opium preparation may cause serious side effects. Stop using opium preparation and call your doctor at once if you have: weak or shallow breathing
- a light-headed feeling, like you might pass out
- or low blood sugar -- headache , hunger, sweating, irritability, dizziness , fast heart rate , and feeling anxious or shaky. Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders. Common side effects of opium preparation may include: nausea , vomiting
- constipation
- or itching, rash.

1380. This particular disease, Diarrhea, interacts with the drug Bismarex and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop taking Bismarex and call your doctor at once if you have: changes in behavior with nausea and vomiting
- hearing loss or ringing in your ears
- diarrhea lasting longer than 2 days
- or worsened stomach symptoms. Bismarex can cause you to have a black or darkened tongue. This is a harmless side effect. Common side effects include: constipation
- or dark colored stools.

Drug Side Effects Summary

1381. This particular disease, Diarrhea, interacts with the drug Bismatrol Maximum Strength and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop taking Bismatrol Maximum Strength and call your doctor at once if you have: changes in behavior with nausea and vomiting
- hearing loss or ringing in your ears
- diarrhea lasting longer than 2 days
- or worsened stomach symptoms. Bismatrol Maximum Strength can cause you to have a black or darkened tongue. This is a harmless side effect. Common side effects include: constipation
- or dark colored stools.

1382. This particular disease, Diarrhea, interacts with the drug crofelemer and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Crofelemer may cause serious side effects. Call your doctor at once if you have: dark urine, jaundice (yellowing of the skin or eyes)
- wheezing, chest tightness, trouble breathing
- or worsening diarrhea. Common side effects of crofelemer may include: abnormal blood tests
- gas
- or cold symptoms such as stuffy nose, sneezing, cough.

1383. This particular disease, Diarrhea, interacts with the drug Florastor Kids and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Although not all side effects are known, Florastor Kids

Drug Side Effects Summary

is thought to be likely safe for most people when used as directed for up to 15 months. Common side effects of Florastor Kids may include: gas, bloating

- or constipation .

1384. This particular disease, Diarrhea, interacts with the drug Mytesi and has the following side effects:

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Mytesi may cause serious side effects. Call your doctor at once if you have: dark urine, jaundice (yellowing of the skin or eyes)

- wheezing, chest tightness, trouble breathing

- or worsening diarrhea. Common side effects of Mytesi may include: abnormal blood tests

- gas

- or cold symptoms such as stuffy nose, sneezing, cough.

1385. This particular disease, Diarrhea, interacts with the drug Pedialyte and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

- itching

- red, swollen, blistered, or peeling skin with or without fever

- wheezing

- tightness in the chest or throat

- trouble breathing, swallowing, or talking

Drug Side Effects Summary

- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of fluid and electrolyte problems like mood changes, confusion, muscle pain or weakness, a heartbeat that does not feel normal, very bad dizziness or passing out, fast heartbeat , more thirst, seizures , feeling very tired or weak, not hungry, unable to pass urine or change in the amount of urine produced, dry mouth, dry eyes, or very bad upset stomach or throwing up. Pedialyte side effects

1386. This particular disease, Diarrhea, interacts with the drug Peptic Relief and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop taking Peptic Relief and call your doctor at once if you have: changes in behavior with nausea and vomiting
- hearing loss or ringing in your ears
- diarrhea lasting longer than 2 days
- or worsened stomach symptoms. Peptic Relief can cause you to have a black or darkened tongue.

This is a harmless side effect. Common side effects include: constipation

- or dark colored stools.

1387. This particular disease, Diarrhea, interacts with the drug Pink Bismuth and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop taking Pink Bismuth and call your doctor at once if you have: changes in behavior with nausea and vomiting
- hearing loss or ringing in your ears
- diarrhea lasting longer than 2 days

Drug Side Effects Summary

- or worsened stomach symptoms. Pink Bismuth can cause you to have a black or darkened tongue.

This is a harmless side effect. Common side effects include: constipation

- or dark colored stools.

1388. This particular disease, Diarrhea, interacts with the drug Saccharomyces Boulardii+MOS and

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Although not all side effects are known,

Saccharomyces Boulardii+MOS is thought to be likely safe for most people when used as directed

for up to 15 months. Common side effects of Saccharomyces Boulardii+MOS may include: gas,

bloating

- or constipation .

1389. This particular disease, Diarrhea, interacts with the drug Soothe Caplets and has the following

- hives

- difficulty breathing

- swelling of your face, lips, tongue, or throat. Stop taking Soothe Caplets and call your doctor at

once if you have: changes in behavior with nausea and vomiting

- hearing loss or ringing in your ears

- diarrhea lasting longer than 2 days

- or worsened stomach symptoms. Soothe Caplets can cause you to have a black or darkened

tongue. This is a harmless side effect. Common side effects include: constipation

- or dark colored stools.

1390. This particular disease, Diarrhea, interacts with the drug atropine / difenoxin and has the following

Drug Side Effects Summary

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Bloating constipation loss of appetite stomach pain (severe) with nausea and vomiting Blurred vision (continuing) or changes in near vision drowsiness (severe) dryness of mouth, nose, and throat (severe) fast heartbeat shortness of breath or troubled breathing (severe) unusual excitement, nervousness, restlessness, or irritability unusual warmth, dryness, and flushing of skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common or rare Blurred vision confusion difficult urination dizziness or lightheadedness drowsiness dryness of skin and mouth fever headache trouble in sleeping unusual tiredness or weakness After you stop using difenoxin and atropine, it may still produce some side effects that need attention. During this period of time, check with your doctor immediately if you notice the following side effects: Increased sweating muscle cramps nausea or vomiting shivering or trembling stomach cramps Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Atropine / difenoxin side effects

1391. This particular disease, Diarrhea, interacts with the drug CeraLyte and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: Symptoms of too much sodium (salt) in the body Convulsions (seizures) dizziness fast heartbeat high blood pressure irritability

Drug Side Effects Summary

muscle twitching restlessness swelling of feet or lower legs weakness Symptoms of too much fluid in the body Puffy eyelids Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Vomiting (mild) Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. CeraLyte side effects

1392. This particular disease, Diarrhea, interacts with the drug electrolyte replacement solutions and

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: Symptoms of too much sodium (salt) in the body Convulsions (seizures) dizziness fast heartbeat high blood pressure irritability muscle twitching restlessness swelling of feet or lower legs weakness Symptoms of too much fluid in the body Puffy eyelids Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Vomiting (mild) Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Electrolyte replacement solutions side effects

Drug Side Effects Summary

1393. This particular disease, Diarrhea, interacts with the drug Enfalyte and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of fluid and electrolyte problems like mood changes, confusion, muscle pain or weakness, a heartbeat that does not feel normal, very bad dizziness or passing out, fast heartbeat , more thirst, seizures , feeling very tired or weak, not hungry, unable to pass urine or change in the amount of urine produced, dry mouth, dry eyes, or very bad upset stomach or throwing up. Enfalyte side effects

1394. This particular disease, Diarrhea, interacts with the drug Equalyte and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching

Drug Side Effects Summary

- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of fluid and electrolyte problems like mood changes, confusion, muscle pain or weakness, a heartbeat that does not feel normal, very bad dizziness or passing out, fast heartbeat , more thirst, seizures , feeling very tired or weak, not hungry, unable to pass urine or change in the amount of urine produced, dry mouth, dry eyes, or very bad upset stomach or throwing up. Equalyte side effects

1395. This particular disease, Diarrhea, interacts with the drug Medi-Lyte and has the following side

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of fluid and electrolyte problems like mood changes, confusion, muscle pain or weakness, a heartbeat that does not feel normal, very

Drug Side Effects Summary

bad dizziness or passing out, fast heartbeat , more thirst, seizures , feeling very tired or weak, not hungry, unable to pass urine or change in the amount of urine produced, dry mouth, dry eyes, or very bad upset stomach or throwing up. Medi-Lyte side effects

1396. This particular disease, Diarrhea, interacts with the drug Motofen and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Bloating constipation loss of appetite stomach pain (severe) with nausea and vomiting Blurred vision (continuing) or changes in near vision drowsiness (severe) dryness of mouth, nose, and throat (severe) fast heartbeat shortness of breath or troubled breathing (severe) unusual excitement, nervousness, restlessness, or irritability unusual warmth, dryness, and flushing of skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common or rare Blurred vision confusion difficult urination dizziness or lightheadedness drowsiness dryness of skin and mouth fever headache trouble in sleeping unusual tiredness or weakness After you stop using this medicine, it may still produce some side effects that need attention. During this period of time, check with your doctor immediately if you notice the following side effects: Increased sweating muscle cramps nausea or vomiting shivering or trembling stomach cramps Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Motofen side effects

Drug Side Effects Summary

1397. This particular disease, Diarrhea, interacts with the drug Oralyte and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of fluid and electrolyte problems like mood changes, confusion, muscle pain or weakness, a heartbeat that does not feel normal, very bad dizziness or passing out, fast heartbeat , more thirst, seizures , feeling very tired or weak, not hungry, unable to pass urine or change in the amount of urine produced, dry mouth, dry eyes, or very bad upset stomach or throwing up. Oralyte side effects

1398. This particular disease, Diarrhea, interacts with the drug Rehydralyte and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching

Drug Side Effects Summary

- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of fluid and electrolyte problems like mood changes, confusion, muscle pain or weakness, a heartbeat that does not feel normal, very bad dizziness or passing out, fast heartbeat , more thirst, seizures , feeling very tired or weak, not hungry, unable to pass urine or change in the amount of urine produced, dry mouth, dry eyes, or very bad upset stomach or throwing up. Rehydralyte side effects

1399. This particular disease, Diarrhea, interacts with the drug ReVital and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of fluid and electrolyte problems like mood changes, confusion, muscle pain or weakness, a heartbeat that does not feel normal, very

Drug Side Effects Summary

bad dizziness or passing out, fast heartbeat , more thirst, seizures , feeling very tired or weak, not hungry, unable to pass urine or change in the amount of urine produced, dry mouth, dry eyes, or very bad upset stomach or throwing up. ReVital side effects

1400. This particular disease, Eczema, interacts with the drug triamcinolone and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Triamcinolone topical may cause serious side effects.

Call your doctor at once if you have: worsening of your skin condition

- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin or gums--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Children can absorb larger amounts of this medicine through the skin and may be more likely to have side effects such as growth delay, headaches, or pain behind the eyes. A baby using this medicine may have a bulging soft spot (the top of the head where the skull hasn't yet grown together). Common side effects of triamcinolone topical may include: burning, itching, dryness, or other irritation of treated skin
- redness or crusting around your hair follicles
- redness or itching around your mouth
- allergic skin reaction
- stretch marks
- acne, increased body hair growth

Drug Side Effects Summary

- thinning skin or discoloration
- or white or "pruned" appearance of the skin (caused by covering treated skin with a tight bandage or other covering).

1401. This particular disease, Eczema, interacts with the drug mometasone and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Mometasone topical may cause serious side effects.

Call your doctor at once if you have: severe skin irritation

- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- or high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor.

Mometasone topical can affect growth in children and should not be used long-term. Common side effects of mometasone topical may include: red or pus-filled bumps on your skin

- acne
- or mild itching, tingling, or burning.

1402. This particular disease, Eczema, interacts with the drug Dupixent and has the following side effects:

- hives , rash, itching
- fever, swollen glands, joint pain
- feeling light-headed, difficult breathing
- swelling of your face, lips, tongue, or throat. Dupixent may cause serious side effects. Call your doctor at once if you have: new or worsening eye pain or discomfort
- vision changes
- watery eyes (your eyes may be more sensitive to light)
- fast pulse

Drug Side Effects Summary

- stomach cramps
- feeling like something is in your eye
- or blood vessel inflammation--fever, chest pain, trouble breathing, skin rash , numbness or prickly feeling in your arms or legs. Common side effects of Dupixent may include: pain, swelling, burning, or irritation where an injection was given
- blood vessel inflammation
- eye redness or itching, puffy eyelids
- stomach pain, nausea , vomiting , diarrhea
- sore throat
- tooth pain
- sleep problems (insomnia)
- cold symptoms such as stuffy nose, sneezing, sore throat
- skin sores or blisters, itching, tingling, burning pain in your thigh or lower back
- joint or muscle pain
- or cold sores or fever blisters on your lips or in your mouth.

1403. This particular disease, Eczema, interacts with the drug hydrocortisone and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Hydrocortisone rectal may cause serious side effects. Call your doctor at once if you have: severe rectal pain or burning, bleeding from your rectum
- new or worsening rectal irritation
- fever or other signs of infection
- or severe headaches, ringing in your ears, pain behind your eyes. Common side effects of hydrocortisone rectal may include: rectal itching, burning, or other irritation

Drug Side Effects Summary

- or dryness or skin changes around the rectum.

1404. This particular disease, Eczema, interacts with the drug methotrexate and has the following s

- hives , difficult breathing, swelling in your face or throat) or a severe skin reaction (fever, sore throat , burning in your eyes, skin pain, red or purple skin rash that spreads and causes blistering and peeling). Methotrexate can cause serious or fatal side effects. Call your doctor at once if you have: sudden chest pain, wheezing, dry cough, cough with mucus, chest pain, feeling short of breath

- fever, chills, swollen lymph glands, night sweats , weight loss

- blisters or ulcers in your mouth, red or swollen gums, trouble swallowing

- vomiting, diarrhea, blood in your urine or stools

- skin changes such as redness, warmth, swelling, or oozing

- low blood cell counts - fever, chills, tiredness, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath

- kidney problems - little or no urination, swelling in your feet or ankles

- liver problems - swelling around your midsection, right-sided upper stomach pain, nausea , loss of appetite, dark urine, jaundice (yellowing of the skin or eyes)

- nerve problems - confusion, weakness, drowsiness, coordination problems, feeling irritable, headache , neck stiffness, vision problems, loss of movement in any part of your body, seizure

- or signs of tumor cell breakdown - tiredness, weakness, muscle cramps, nausea, vomiting, diarrhea, fast or slow heart rate , tingling in your hands and feet or around your mouth. Common methotrexate side effects may include: fever, chills, tiredness, not feeling well

- low blood cell counts

- mouth sores

- nausea, stomach pain

Drug Side Effects Summary

- abnormal liver function tests
- hair loss
- burning skin lesions
- or being more sensitive to light.

1405. This particular disease, Eczema, interacts with the drug clobetasol and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin - weight gain in your face and shoulders, slow wound healing, skin discoloration, thinning skin, increased body hair, tiredness, mood changes, menstrual changes, sexual changes. Common side effects may include: burning, itching, swelling, or irritation of treated skin
- dry or cracking skin
- redness or crusting around your hair follicles
- spider veins
- stretch marks, thinning skin
- rash or hives
- acne
- or temporary hair loss .

Drug Side Effects Summary

1406. This particular disease, Eczema, interacts with the drug dupilumab and has the following side effects:

- Serious side effects of dupilumab include: Allergic reactions. Dupilumab can cause allergic reactions that can sometimes be severe. Stop using dupilumab and tell your healthcare provider or get emergency help right away if you get any of the following signs or symptoms: breathing problems or wheezing fast pulse fever general ill feeling swollen lymph nodes swelling of the face, lips, mouth, tongue, or throat hives itching nausea or vomiting fainting, dizziness, feeling lightheaded joint pain skin rash cramps in your stomach-area Eye problems. Tell your healthcare provider if you have any new or worsening eye problems, including eye pain or changes in vision, such as blurred vision. Your healthcare provider may send you to an ophthalmologist for an eye exam if needed. Inflammation of your blood vessels. Rarely, this can happen in people with asthma who receive dupilumab. This may happen in people who also take a steroid medicine by mouth that is being stopped or the dose is being lowered. It is not known whether this is caused by dupilumab. Tell your healthcare provider right away if you have: rash worsening shortness of breath persistent fever chest pain a feeling of pins and needles or numbness of your arms or legs Joint aches and pain. Joint aches and pain can happen in people who use dupilumab. Some people have had trouble walking or moving due to their joint symptoms, and in some cases needed to be hospitalized. Tell your healthcare provider about any new or worsening joint symptoms. Your healthcare provider may stop dupilumab if you develop joint symptoms. The most common side effects of dupilumab include: injection site reactions eye and eyelid inflammation, including redness, swelling, and itching, sometimes with blurred vision pain in the throat (oropharyngeal pain) cold sores in your mouth or on your lips high count of a certain white blood cell (eosinophilia) trouble sleeping (insomnia) toothache gastritis joint pain (arthralgia) parasitic (helminth) infections The following additional side effects have been reported with dupilumab: facial rash or redness Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all of the possible side effects of dupilumab. Call your doctor for medical advice about side effects. You may report side

Drug Side Effects Summary

effects to FDA at 1-800-FDA-1088. Dupilumab side effects

1407. This particular disease, Eczema, interacts with the drug Elidel and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe pain, burning, itching, or stinging
- new or worsened skin symptoms (warts, unusual rash or skin lesions, blistering or oozing)
- a mole that has changed in size or color
- or symptoms of a viral skin infection - cold sores around your mouth, skin sores or blisters, scabs, itching, tingling. Common Elidel side effects may include: burning, stinging, or soreness of treated skin (especially during the first few days of treatment)
- viral skin infection
- runny or stuffy nose
- headache
- or fever, chills, body aches, sore throat, cough, swollen glands.

1408. This particular disease, Eczema, interacts with the drug fluocinonide and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: redness, warmth, swelling, oozing, or severe irritation of any treated skin
- worsening of your skin condition
- or possible signs of absorbing fluocinonide through your skin - weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair,

Drug Side Effects Summary

muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Steroid medicine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common fluocinonide side effects may include: burning, stinging, itching, or dryness of treated skin

- redness or crusting around your hair follicles
- stretch marks
- spider veins
- acne
- lightened color of treated skin
- headache
- or stuffy nose, sore throat .

1409. This particular disease, Eczema, interacts with the drug desonide and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Desonide topical may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Steroid medicine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common side effects of desonide topical may include:

Drug Side Effects Summary

stinging or burning of treated skin

- skin irritation, redness, itching, or hardening
- dry, scaly, or oily skin
- swelling in your hands or feet
- acne, stretch marks
- or redness or crusting around your hair follicles.

1410. This particular disease, Eczema, interacts with the drug Elocon and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe skin irritation
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- or high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor.

Mometasone can affect growth in children and should not be used long-term. Common Elocon side effects may include: red or pus-filled bumps on your skin

- acne
- or mild itching, tingling, or burning.

1411. This particular disease, Eczema, interacts with the drug desoximetasone and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Desoximetasone topical may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin

Drug Side Effects Summary

- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Common side effects of desoximetasone topical may include: rash, itching, or irritation around your mouth
- burning, irritation, itching, or dryness of treated skin
- redness or crusting around your hair follicles
- increased hair growth
- blisters, pimples , or crusting of treated skin
- changes in color of treated skin
- white or "pruned" appearance of the skin
- or stretch marks.

1412. This particular disease, Eczema, interacts with the drug pimecrolimus and has the following s

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Pimecrolimus topical may cause serious side effects. Call your doctor at once if you have: severe pain, burning, itching, or stinging
- new or worsened skin symptoms (warts , unusual rash or skin lesions, blistering or oozing)
- a mole that has changed in size or color
- or symptoms of a viral skin infection--cold sores around your mouth, skin sores or blisters, scabs, itching, tingling. Common side effects of pimecrolimus topical may include: burning, stinging, or soreness of treated skin (especially during the first few days of treatment)

Drug Side Effects Summary

- viral skin infection
- runny or stuffy nose
- headache
- or fever, chills, body aches, sore throat , cough, swollen glands.

1413. This particular disease, Eczema, interacts with the drug Cortizone-10 and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blistering, burning, crusting, dryness, or flaking of the skin irritation itching, scaling, severe redness, soreness, or swelling of the skin redness and scaling around the mouth thinning of the skin with easy bruising, especially when used on the face or where the skin folds together (e.g. between the fingers) thinning, weakness, or wasting away of the skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Acne or pimples burning and itching of the skin with pinhead-sized red blisters burning, itching, and pain in hairy areas, or pus at the root of the hair increased hair growth on the forehead, back, arms, and legs lightening of normal skin color lightening of treated areas of dark skin reddish purple lines on the arms, face, legs, trunk, or groin softening of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Cortizone-10 side effects

Drug Side Effects Summary

1414. This particular disease, Eczema, interacts with the drug halobetasol and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin - weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Common halobetasol side effects may include: burning, stinging, itching, or dryness of treated skin
- pain where the foam was applied
- redness or crusting around your hair follicles
- stretch marks
- spider veins
- headache
- or cold symptoms such as stuffy nose, sneezing, sore throat .

1415. This particular disease, Eczema, interacts with the drug Kenalog and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medication and call your doctor at once if you have any of these signs that you may be absorbing triamcinolone through your skin or

Drug Side Effects Summary

gums: blurred vision, or seeing halos around lights

- uneven heartbeats
- mood changes
- sleep problems (insomnia)
- weight gain, puffiness in your face
- or feeling tired. Less serious Kenalog side effects may include: skin redness, burning, itching, or peeling
- thinning of your skin
- or blistering skin
- or stretch marks.

1416. This particular disease, Eczema, interacts with the drug Topicort and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Topicort may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Common side effects of Topicort may include: rash, itching, or irritation around your mouth
- burning, irritation, itching, or dryness of treated skin
- redness or crusting around your hair follicles

Drug Side Effects Summary

- increased hair growth
- blisters, pimples , or crusting of treated skin
- changes in color of treated skin
- white or "pruned" appearance of the skin
- or stretch marks.

1417. This particular disease, Eczema, interacts with the drug fluocinolone and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Common fluocinolone side effects may include: burning, itching, or irritation of treated skin
- increased hair growth
- white or "pruned" appearance of the skin
- rash, pimples
- or changes in color of treated skin.

1418. This particular disease, Eczema, interacts with the drug fluocinolone and has the following side effects:

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe irritation in or around your ear
- or signs of a skin infection--warmth, redness, swelling, sores, oozing. Your skin can absorb topical steroid medicine, which may cause steroid side effects throughout the body. Tell your doctor if you have: swelling, weight gain (especially in your face or your upper back and torso)
- thinning skin, stretch marks, acne, increased facial hair
- memory problems, feeling irritable
- irregular menstrual periods, changes in sexual function
- or growth delay in a child. Children can absorb larger amounts of this medicine through the skin and may be more likely to have side effects. Common side effects of fluocinolone otic may include: burning, itching, irritation, or dryness
- rash around the ear in which this medicine was applied
- rash or irritation around your mouth
- acne, pimples , stretch marks
- lightened skin color
- or redness or crusting around your hair follicles.

1419. This particular disease, Eczema, interacts with the drug fluticasone and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Fluticasone topical may cause serious side effects. Call your doctor at once if you have: skin pain, tenderness, or swelling
- any wound that will not heal
- blurred vision, eye pain
- severe skin irritation after using the medicine

Drug Side Effects Summary

- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Common side effects of fluticasone topical may include: skin redness, itching, or rash
- burning or stinging of treated skin
- increased hair growth
- or feeling light-headed.

1420. This particular disease, Eczema, interacts with the drug alclometasone and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Topical steroid medicine can be absorbed through the skin, which may cause steroid side effects throughout the body. Stop using alclometasone topical and call your doctor if you have: blurred vision, or seeing halos around lights
- uneven heartbeats
- sleep problems (insomnia)
- weight gain, puffiness in your face
- or tired feeling. Common side effects of alclometasone topical may include: redness, burning, or itching of treated skin
- skin dryness or irritation
- acne, skin rash
- lightened color of treated skin
- or folliculitis (redness or crusting around your hair follicles).

1421. This particular disease, Eczema, interacts with the drug azathioprine and has the following si

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Azathioprine may cause a serious brain infection that can lead to disability or death. Call your doctor right away if you have problems with speech, thought, vision, or muscle movement. These symptoms may start gradually and get worse quickly. Stop using azathioprine and call your doctor right away if you have any of these symptoms of lymphoma : fever, swollen glands, body aches, night sweats , not feeling well
- pale skin, rash, easy bruising or bleeding
- cold hands and feet, feeling light-headed or short of breath
- pain in your upper stomach that may spread to your shoulder
- or feeling full after eating only a small amount, weight loss . Also call your doctor at once if you have: signs of infection (fever, chills, weakness, flu symptoms, sore throat , cough, pain or burning when you urinate)
- severe nausea , vomiting , or diarrhea
- easy bruising, unusual bleeding
- rapid heartbeats, shortness of breath
- pale skin, cold hands and feet
- or dark urine, jaundice (yellowing of the skin or eyes). Common side effects of azathioprine may include: nausea, diarrhea, stomach pain
- hair loss
- or skin rash .

1422. This particular disease, Eczema, interacts with the drug Cloderm and has the following side effects

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Cloderm may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Steroid medicine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common side effects of Cloderm may include: burning, itching, or dryness of treated skin
- redness or crusting around your hair follicles
- increased hair growth
- acne, skin rash
- stretch marks, thinning skin
- lightened color of treated skin
- or white or "pruned" appearance of the skin (caused by leaving wound dressings on treated skin for long periods of time).

1423. This particular disease, Eczema, interacts with the drug Cutivate and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Cutivate may cause serious side effects. Call your doctor at once if you have: skin pain, tenderness, or swelling
- any wound that will not heal
- blurred vision, eye pain

Drug Side Effects Summary

- severe skin irritation after using the medicine
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Common side effects of Cutivate may include: skin redness, itching, or rash
- burning or stinging of treated skin
- increased hair growth
- or feeling light-headed.

1424. This particular disease, Eczema, interacts with the drug cyclosporine and has the following s

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Cyclosporine may cause serious side effects. Call your doctor at once if you have: problems with speech, thought, vision, or muscle movement (may start gradually and get worse quickly)
- a seizure
- severe headache , blurred vision, pounding in your neck or ears
- pale skin, easy bruising, unusual bleeding
- high potassium level-- nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- kidney problems--urinating less, swelling in your feet or ankles, feeling tired or short of breath
- liver problems--loss of appetite, stomach pain (upper right side), jaundice (yellowing of the skin or eyes)
- or signs of infection--fever, chills, flu symptoms, mouth sores, skin sores, sore throat , cough, trouble breathing. Common side effects of cyclosporine may include: tremors or shaking

Drug Side Effects Summary

- acne , increased growth of facial or body hair
- increased blood pressure
- nausea, diarrhea
- or swollen or painful gums.

1425. This particular disease, Eczema, interacts with the drug DermOtic Oil and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe irritation in or around your ear
- or signs of a skin infection--warmth, redness, swelling, sores, oozing. Your skin can absorb topical steroid medicine, which may cause steroid side effects throughout the body. Tell your doctor if you have: swelling, weight gain (especially in your face or your upper back and torso)
- thinning skin, stretch marks, acne, increased facial hair
- memory problems, feeling irritable
- irregular menstrual periods, changes in sexual function
- or growth delay in a child. Children can absorb larger amounts of this medicine through the skin and may be more likely to have side effects. Common side effects of DermOtic Oil may include: burning, itching, irritation, or dryness
- rash around the ear in which this medicine was applied
- rash or irritation around your mouth
- acne, pimples , stretch marks
- lightened skin color
- or redness or crusting around your hair follicles.

Drug Side Effects Summary

1426. This particular disease, Eczema, interacts with the drug Locoid Lipocream and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blistering, burning, crusting, dryness, or flaking of the skin irritation itching, scaling, severe redness, soreness, or swelling of the skin redness and scaling around the mouth thinning of the skin with easy bruising, especially when used on the face or where the skin folds together (e.g. between the fingers) thinning, weakness, or wasting away of the skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Acne or pimples burning and itching of the skin with pinhead-sized red blisters burning, itching, and pain in hairy areas, or pus at the root of the hair increased hair growth on the forehead, back, arms, and legs lightening of normal skin color lightening of treated areas of dark skin reddish purple lines on the arms, face, legs, trunk, or groin softening of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Locoid Lipocream side effects

1427. This particular disease, Eczema, interacts with the drug Aristocort A and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blistering, burning, crusting, dryness, or flaking of the skin irritation itching, scaling, severe redness, soreness, or

Drug Side Effects Summary

swelling of the skin redness and scaling around the mouth thinning of the skin with easy bruising, especially when used on the face or where the skin folds together (e.g. between the fingers) Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Acne or pimples burning and itching of the skin with pinhead-sized red blisters burning, itching, and pain in hairy areas, or pus at the root of the hair increased hair growth on the forehead, back, arms, and legs lightening of normal skin color lightening of treated areas of dark skin reddish purple lines on the arms, face, legs, trunk, or groin softening of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Aristocort A side effects

1428. This particular disease, Eczema, interacts with the drug Aristocort R and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blistering, burning, crusting, dryness, or flaking of the skin irritation itching, scaling, severe redness, soreness, or swelling of the skin redness and scaling around the mouth thinning of the skin with easy bruising, especially when used on the face or where the skin folds together (e.g. between the fingers) Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any

Drug Side Effects Summary

questions about them: Incidence not known Acne or pimples burning and itching of the skin with pinhead-sized red blisters burning, itching, and pain in hairy areas, or pus at the root of the hair increased hair growth on the forehead, back, arms, and legs lightening of normal skin color lightening of treated areas of dark skin reddish purple lines on the arms, face, legs, trunk, or groin softening of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Aristocort R side effects

1429. This particular disease, Eczema, interacts with the drug Beser and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Beser may cause serious side effects. Call your doctor at once if you have: skin pain, tenderness, or swelling
- any wound that will not heal
- blurred vision, eye pain
- severe skin irritation after using the medicine
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Common side effects of Beser may include: skin redness, itching, or rash
- burning or stinging of treated skin
- increased hair growth
- or feeling light-headed.

1430. This particular disease, Eczema, interacts with the drug Beta HC and has the following side effects

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Beta HC may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Less serious side effects may be more likely, and you may have none at all.

1431. This particular disease, Eczema, interacts with the drug Cinolar and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blistering, burning, crusting, dryness, or flaking of the skin irritation itching, scaling, severe redness, soreness, or swelling of the skin redness and scaling around the mouth thinning of the skin with easy bruising, especially when used on the face or where the skin folds together (e.g. between the fingers) Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Acne or pimples burning and itching of the skin with pinhead-sized red blisters burning, itching, and pain in hairy areas, or pus at the root of the hair increased hair growth on the forehead, back, arms, and legs lightening of normal skin color

Drug Side Effects Summary

lightening of treated areas of dark skin reddish purple lines on the arms, face, legs, trunk, or groin
softening of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Cinolar side effects

1432. This particular disease, Eczema, interacts with the drug Cordran Tape and has the following s

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Cordran Tape may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Common side effects of Cordran Tape may include: burning, itching, or irritation of treated skin
- increased hair growth
- white or "pruned" appearance of the skin
- rash, pimples
- or changes in color of treated skin.

1433. This particular disease, Eczema, interacts with the drug DermacinRx Trizapak and has the fol

- No side effects listed

Drug Side Effects Summary

1434. This particular disease, Eczema, interacts with the drug Dermasorb TA and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Dermasorb TA may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin or gums--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Children can absorb larger amounts of this medicine through the skin and may be more likely to have side effects such as growth delay, headaches, or pain behind the eyes. A baby using this medicine may have a bulging soft spot (the top of the head where the skull hasn't yet grown together). Common side effects of Dermasorb TA may include: burning, itching, dryness, or other irritation of treated skin
- redness or crusting around your hair follicles
- redness or itching around your mouth
- allergic skin reaction
- stretch marks
- acne, increased body hair growth
- thinning skin or discoloration
- or white or "pruned" appearance of the skin (caused by covering treated skin with a tight bandage or other covering).

Drug Side Effects Summary

1435. This particular disease, Eczema, interacts with the drug DesOwen and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. DesOwen may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Steroid medicine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common side effects of DesOwen may include: stinging or burning of treated skin
- skin irritation, redness, itching, or hardening
- dry, scaly, or oily skin
- swelling in your hands or feet
- acne, stretch marks
- or redness or crusting around your hair follicles.

1436. This particular disease, Eczema, interacts with the drug flurandrenolide and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Flurandrenolide topical may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition

Drug Side Effects Summary

- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Common side effects of flurandrenolide topical may include: burning, itching, or irritation of treated skin
- increased hair growth
- white or "pruned" appearance of the skin
- rash, pimples
- or changes in color of treated skin.

1437. This particular disease, Eczema, interacts with the drug Halog and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Halog may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso)
- slow wound healing, thinning skin, increased body hair
- changes in sexual function
- feeling depressed, anxious, or irritable. Common side effects of Halog may include: burning, itching, or irritation of treated skin

Drug Side Effects Summary

- increased hair growth
- white or "pruned" appearance of the skin
- rash, pimples
- or changes in color of treated skin.

1438. This particular disease, Eczema, interacts with the drug HiDex and has the following side effects:

- No side effects listed

1439. This particular disease, Eczema, interacts with the drug Juulissa Pharmapak and has the following side effects:

- No side effects listed

1440. This particular disease, Eczema, interacts with the drug Locoid and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Locoid may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Less serious side effects may be more likely, and you may have none at all.

1441. This particular disease, Eczema, interacts with the drug Oralone and has the following side effects:

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Oralone may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin or gums--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Children can absorb larger amounts of this medicine through the skin and may be more likely to have side effects such as growth delay, headaches, or pain behind the eyes. A baby using this medicine may have a bulging soft spot (the top of the head where the skull hasn't yet grown together). Common side effects of Oralone may include: burning, itching, dryness, or other irritation of treated skin
- redness or crusting around your hair follicles
- redness or itching around your mouth
- allergic skin reaction
- stretch marks
- acne, increased body hair growth
- thinning skin or discoloration
- or white or "pruned" appearance of the skin (caused by covering treated skin with a tight bandage or other covering).

1442. This particular disease, Eczema, interacts with the drug SilaLite Pak and has the following side effects:

- No side effects listed

Drug Side Effects Summary

1443. This particular disease, Eczema, interacts with the drug Treziopak and has the following side effects:

- No side effects listed

1444. This particular disease, Eczema, interacts with the drug Trianex and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Trianex may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin or gums--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Children can absorb larger amounts of this medicine through the skin and may be more likely to have side effects such as growth delay, headaches, or pain behind the eyes. A baby using this medicine may have a bulging soft spot (the top of the head where the skull hasn't yet grown together). Common side effects of Trianex may include: burning, itching, dryness, or other irritation of treated skin
- redness or crusting around your hair follicles
- redness or itching around your mouth
- allergic skin reaction
- stretch marks
- acne, increased body hair growth
- thinning skin or discoloration

Drug Side Effects Summary

- or white or "pruned" appearance of the skin (caused by covering treated skin with a tight bandage or other covering).

1445. This particular disease, Eczema, interacts with the drug Triderm and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Triderm may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin or gums--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Children can absorb larger amounts of this medicine through the skin and may be more likely to have side effects such as growth delay, headaches, or pain behind the eyes. A baby using this medicine may have a bulging soft spot (the top of the head where the skull hasn't yet grown together). Common side effects of Triderm may include: burning, itching, dryness, or other irritation of treated skin
- redness or crusting around your hair follicles
- redness or itching around your mouth
- allergic skin reaction
- stretch marks
- acne, increased body hair growth
- thinning skin or discoloration

Drug Side Effects Summary

- or white or "pruned" appearance of the skin (caused by covering treated skin with a tight bandage or other covering).

1446. This particular disease, Eczema, interacts with the drug Tritocin and has the following side effects:

- No side effects listed

1447. This particular disease, Eczema, interacts with the drug Ultravate and has the following side effects:

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Ultravate may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition

- redness, warmth, swelling, oozing, or severe irritation of any treated skin

- blurred vision, tunnel vision, eye pain , or seeing halos around lights

- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor

- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Common side effects of Ultravate may include: burning, stinging, itching, or dryness of treated skin

- pain where the foam was applied

- redness or crusting around your hair follicles

- stretch marks

- spider veins

- headache

- or cold symptoms such as stuffy nose, sneezing, sore throat .

Drug Side Effects Summary

1448. This particular disease, Eczema, interacts with the drug Ala-Cort and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Ala-Cort may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Less serious side effects may be more likely, and you may have none at all.

1449. This particular disease, Eczema, interacts with the drug Ala-Scalp and has the following side effects:

- No side effects listed

1450. This particular disease, Eczema, interacts with the drug Ala-Scalp HP and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blistering, burning, crusting, dryness, or flaking of the skin irritation itching, scaling, severe redness, soreness, or swelling of the skin redness and scaling around the mouth thinning of the skin with easy bruising, especially when used on the face or where the skin folds together (e.g. between the fingers) thinning, weakness, or wasting away of the skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or

Drug Side Effects Summary

reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Acne or pimples burning and itching of the skin with pinhead-sized red blisters burning, itching, and pain in hairy areas, or pus at the root of the hair increased hair growth on the forehead, back, arms, and legs lightening of normal skin color lightening of treated areas of dark skin reddish purple lines on the arms, face, legs, trunk, or groin softening of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Ala-Scalp HP side effects

1451. This particular disease, Eczema, interacts with the drug amcinonide and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Amcinonide topical may cause serious side effects.

Call your doctor at once if you have: worsening of your skin condition

- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , headaches, pain behind the eyes, tiredness, mood changes, menstrual changes, sexual changes. Steroid medicine can affect growth in children. Tell your doctor if your child is not growing or gaining weight at a normal rate while using amcinonide topical. Common side effects of amcinonide topical may include: burning, itching, irritation, or dryness of treated skin
- redness or crusting around your hair follicles

Drug Side Effects Summary

- acne, increased hair growth
- stretch marks, thinning skin
- or white or "pruned" appearance of the skin (caused by covering treated skin with a tight bandage or other covering).

1452. This particular disease, Eczema, interacts with the drug ApexiCon E and has the following side effects:

- hives
 - difficult breathing
 - swelling of your face, lips, tongue, or throat. Your skin can absorb topical steroid medicine, which may cause steroid side effects throughout the body. Tell your doctor if you have: swelling, weight gain (especially in your face or your upper back and torso)
 - irregular menstrual periods, changes in sexual function
 - or memory problems, feeling irritable. ApexiCon E may cause serious side effects. Stop using ApexiCon E and call your doctor at once if you have: blurred vision, tunnel vision, eye pain , or seeing halos around lights
 - severe skin irritation where the medicine was applied
 - or signs of skin infection (swelling, redness, warmth, oozing). Steroid medicine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine.
- Common side effects of ApexiCon E may include: redness or crusting around your hair follicles
- burning or itching of treated skin
 - lightened color of treated skin
 - skin dryness or irritation
 - acne, skin rash
 - increased hair growth
 - stretch marks

Drug Side Effects Summary

- or white or "pruned" appearance of the skin (caused by leaving wound dressings on for long periods of time).

1453. This particular disease, Eczema, interacts with the drug Aquanil HC and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Aquanil HC may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Less serious side effects may be more likely, and you may have none at all.

1454. This particular disease, Eczema, interacts with the drug Aquax-H and has the following side effects

- No side effects listed

1455. This particular disease, Eczema, interacts with the drug Caldecort and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blistering, burning, crusting, dryness, or flaking of the skin irritation itching, scaling, severe redness, soreness, or swelling of the skin redness and scaling around the mouth thinning of the skin with easy bruising, especially when used on the face or where the skin folds together (e.g. between the fingers)

Drug Side Effects Summary

thinning, weakness, or wasting away of the skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Acne or pimples burning and itching of the skin with pinhead-sized red blisters burning, itching, and pain in hairy areas, or pus at the root of the hair increased hair growth on the forehead, back, arms, and legs lightening of normal skin color lightening of treated areas of dark skin reddish purple lines on the arms, face, legs, trunk, or groin softening of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Caldecort side effects

1456. This particular disease, Eczema, interacts with the drug clioquinol / hydrocortisone and has t

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blistering, burning, crusting, dryness, or flaking of the skin burning and itching of the skin with pinhead-sized red blisters burning, itching, and pain in hairy areas, or pus at the root of the hair darkening of the skin diarrhea dizziness dryness fainting fast heartbeat fever hives, itching, or rash hoarseness irritation itching, scaling, severe redness, soreness, or swelling of the skin joint pain, stiffness, or swelling loss of appetite mental depression nausea reddish purple lines on the arms, face, legs, trunk, or groin redness and scaling around the mouth softening of the skin swelling of the eyelids, face, lips, hands, or feet thinning of the skin with easy bruising, especially when used on the face or where the skin folds together (eg, between the fingers) tightness in the chest troubled breathing or swallowing

Drug Side Effects Summary

unusual tiredness or weakness vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Acne or pimples increased hair growth on the forehead, back, arms, and legs lightening of normal skin color lightening of the treated areas of dark skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Clioquinol / hydrocortisone topical side effects

1457. This particular disease, Eczema, interacts with the drug Clobex and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Clobex may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain in your face and shoulders, slow wound healing, skin discoloration, thinning skin, increased body hair, tiredness, mood changes, menstrual changes, sexual changes. Common side effects of Clobex may include: burning, itching, swelling, or irritation of treated skin
- dry or cracking skin
- redness or crusting around your hair follicles

Drug Side Effects Summary

- spider veins
- stretch marks, thinning skin
- rash or hives
- acne
- or temporary hair loss .

1458. This particular disease, Eczema, interacts with the drug clocortolone and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Clocortolone topical may cause serious side effects.

Call your doctor at once if you have: worsening of your skin condition

- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Steroid medicine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common side effects of clocortolone topical may include: burning, itching, or dryness of treated skin

- redness or crusting around your hair follicles
- increased hair growth
- acne, skin rash
- stretch marks, thinning skin
- lightened color of treated skin
- or white or "pruned" appearance of the skin (caused by leaving wound dressings on treated skin

Drug Side Effects Summary

for long periods of time).

1459. This particular disease, Eczema, interacts with the drug Cordran and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Cordran may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Common side effects of Cordran may include: burning, itching, or irritation of treated skin
- increased hair growth
- white or "pruned" appearance of the skin
- rash, pimples
- or changes in color of treated skin.

1460. This particular disease, Eczema, interacts with the drug Cortizone-5 and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Cortizone-5 may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor

Drug Side Effects Summary

- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Less serious side effects may be more likely, and you may have none at all.

1461. This particular disease, Eczema, interacts with the drug Derma-Smoothe / FS (Scalp) and has

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medication and call your doctor at once if you have severe irritation of any treated skin, or if you show signs of absorbing Derma-Smoothe/FS (Scalp) through your skin, such as: blurred vision, headache
- thinning skin, easy bruising
- mood changes
- weight gain, puffiness in your face
- or muscle weakness, feeling tired. Common side effects of Derma-Smoothe/FS (Scalp) may include: mild skin itching, burning, dryness, or other irritation
- temporary hair loss
- swollen hair follicles
- blisters, pimples , or crusting of treated skin
- changes in color of treated skin
- skin rash or irritation around your mouth
- fever, cough, sore throat , runny or stuffy nose
- or red spots or the appearance of tiny veins (especially around your nose, cheeks, or chin).

1462. This particular disease, Eczema, interacts with the drug Dermarest Plus Anti-Itch and has the

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Dermarest Plus Anti-Itch may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Less serious side effects may be more likely, and you may have none at all.

1463. This particular disease, Eczema, interacts with the drug Dermovate and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Burning or stinging at the application site Less common Burning sensation of the skin dry skin flushing or redness of the skin itching, scaling, severe redness, soreness, or swelling of the skin skin irritation skin rash, encrusted, scaly and oozing thinning of the skin with easy bruising, especially when used on the face or where the skin folds together (eg, between the fingers) Rare Thinning, weakness, or wasting away of the skin Incidence not known Backache blindness blurred vision burning, itching, and pain in hairy areas, or pus at the root of the hair change in vision dry mouth eye pain facial hair growth in females fractures fruit-like breath odor full or round face, neck, or trunk hair loss headache hives increased hunger increased thirst increased urination irritability loss of sexual desire or ability menstrual irregularities muscle wasting nausea redness and scaling around the mouth stomach pain sugar in the urine sweating tearing thinning of the hair troubled breathing unexplained weight loss unusual

Drug Side Effects Summary

tiredness or weakness vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Cough sore throat Less common Body aches or pain changes in skin coloring congestion dryness of the throat raised, dark red, wart-like spots on the skin, especially when used on the face skin discomfort stuffy or runny nose tender, swollen glands in the neck trouble swallowing voice changes Incidence not known Acne or pimples burning and itching of the skin with pinhead-sized red blisters increased hair growth on the forehead, back, arms, and legs lightening of normal skin color lightening of treated areas of dark skin reddish purple lines on the arms, face, legs, trunk, or groin softening of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Dermovate side effects

1464. This particular disease, Eczema, interacts with the drug Dermtex HC and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Dermtex HC may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual

Drug Side Effects Summary

changes. Less serious side effects may be more likely, and you may have none at all.

1465. This particular disease, Eczema, interacts with the drug Desonate and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin - weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Steroid medicine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common Desonate side effects may include: stinging or burning of treated skin
- skin irritation, redness, itching, or hardening
- dry, scaly, or oily skin
- swelling in your hands or feet
- acne, stretch marks
- or redness or crusting around your hair follicles.

1466. This particular disease, Eczema, interacts with the drug doxepin and has the following side effects:

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Although the risk of serious side effects is low when doxepin topical is applied to the skin, side effects can occur if the medicine is absorbed into your bloodstream. You may feel drowsy after applying doxepin topical, especially if you have treated large areas of skin. Doxepin topical may cause serious side effects. Stop using doxepin topical and call your doctor at once if you have: severe drowsiness

- confusion

- severe dry mouth

- painful or difficult urination

- blurred vision

- or new or worsening skin symptoms. Some side effects may be more likely in older adults.

Common side effects of doxepin topical may include: burning or stinging where the medicine was applied

- mild drowsiness

- or dry mouth.

1467. This particular disease, Eczema, interacts with the drug Fluocinonide-E and has the following

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Fluocinonide-E may cause serious side effects. Call your doctor at once if you have: redness, warmth, swelling, oozing, or severe irritation of any treated skin

- worsening of your skin condition

- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual

Drug Side Effects Summary

changes. Steroid medicine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common side effects of Fluocinonide-E may include: burning, stinging, itching, or dryness of treated skin

- redness or crusting around your hair follicles
- stretch marks
- spider veins
- acne
- lightened color of treated skin
- headache
- or stuffy nose, sore throat .

1468. This particular disease, Eczema, interacts with the drug hydrocortisone / iodoquinol and has

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blistering, burning, crusting, dryness, flaking, scaling, severe redness, soreness, or swelling of the skin burning and itching of the skin with pinhead-sized red blisters burning, itching, and pain in hairy areas, or pus at the root of hair reddish purple lines on the arms, face, legs, trunk, or groin redness and scaling around the mouth softening of the skin thinning of the skin with easy bruising, especially when used on the face or where the skin folds together (eg, between the fingers) Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Acne or pimples increased hair growth on the forehead, back, arms, and

Drug Side Effects Summary

legs lightening of normal skin color lightening of the treated areas of dark skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Hydrocortisone / iodoquinol topical side effects

1469. This particular disease, Eczema, interacts with the drug Itch-X Lotion and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Itch-X Lotion may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Less serious side effects may be more likely, and you may have none at all.

1470. This particular disease, Eczema, interacts with the drug NuCort and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blistering, burning, crusting, dryness, or flaking of the skin irritation itching, scaling, severe redness, soreness, or swelling of the skin redness and scaling around the mouth thinning of the skin with easy bruising, especially when used on the face or where the skin folds together (e.g. between the fingers) thinning, weakness, or wasting away of the skin Some side effects may occur that usually do not

Drug Side Effects Summary

need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Acne or pimples burning and itching of the skin with pinhead-sized red blisters burning, itching, and pain in hairy areas, or pus at the root of the hair increased hair growth on the forehead, back, arms, and legs lightening of normal skin color lightening of treated areas of dark skin reddish purple lines on the arms, face, legs, trunk, or groin softening of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. NuCort side effects

1471. This particular disease, Eczema, interacts with the drug Pandel and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Pandel may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Less serious side effects may be more likely, and you may have none at all.

1472. This particular disease, Eczema, interacts with the drug Prudoxin and has the following side effects

Drug Side Effects Summary

- hives
 - difficult breathing
 - swelling of your face, lips, tongue, or throat. Although the risk of serious side effects is low when Prudoxin is applied to the skin, side effects can occur if the medicine is absorbed into your bloodstream. You may feel drowsy after applying Prudoxin, especially if you have treated large areas of skin. Prudoxin may cause serious side effects. Stop using Prudoxin and call your doctor at once if you have: severe drowsiness
 - confusion
 - severe dry mouth
 - painful or difficult urination
 - blurred vision
 - or new or worsening skin symptoms. Some side effects may be more likely in older adults.
- Common side effects of Prudoxin may include: burning or stinging where the medicine was applied
- mild drowsiness
 - or dry mouth.

1473. This particular disease, Eczema, interacts with the drug Sarnol-HC and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Sarnol-HC may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair,

Drug Side Effects Summary

muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Less serious side effects may be more likely, and you may have none at all.

1474. This particular disease, Eczema, interacts with the drug Temovate and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Temovate may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain in your face and shoulders, slow wound healing, skin discoloration, thinning skin, increased body hair, tiredness, mood changes, menstrual changes, sexual changes. Common side effects of Temovate may include: burning, itching, swelling, or irritation of treated skin
- dry or cracking skin
- redness or crusting around your hair follicles
- spider veins
- stretch marks, thinning skin
- rash or hives
- acne
- or temporary hair loss .

1475. This particular disease, Eczema, interacts with the drug Texacort and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of

Drug Side Effects Summary

these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blistering, burning, crusting, dryness, or flaking of the skin irritation itching, scaling, severe redness, soreness, or swelling of the skin redness and scaling around the mouth thinning of the skin with easy bruising, especially when used on the face or where the skin folds together (e.g. between the fingers) thinning, weakness, or wasting away of the skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Acne or pimples burning and itching of the skin with pinhead-sized red blisters burning, itching, and pain in hairy areas, or pus at the root of the hair increased hair growth on the forehead, back, arms, and legs lightening of normal skin color lightening of treated areas of dark skin reddish purple lines on the arms, face, legs, trunk, or groin softening of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Texacort side effects

1476. This particular disease, Eczema, interacts with the drug U-Cort and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blistering, burning, crusting, dryness, or flaking of the skin irritation itching, scaling, severe redness, soreness, or swelling of the skin redness and scaling around the mouth thinning of the skin with easy bruising, especially when used on the face or where the skin folds together (e.g. between the fingers)

Drug Side Effects Summary

thinning, weakness, or wasting away of the skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Acne or pimples burning and itching of the skin with pinhead-sized red blisters burning, itching, and pain in hairy areas, or pus at the root of the hair increased hair growth on the forehead, back, arms, and legs lightening of normal skin color lightening of treated areas of dark skin reddish purple lines on the arms, face, legs, trunk, or groin softening of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. U-Cort side effects

1477. This particular disease, Eczema, interacts with the drug Vanos and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Vanos may cause serious side effects. Call your doctor at once if you have: redness, warmth, swelling, oozing, or severe irritation of any treated skin
- worsening of your skin condition
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Steroid medicine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common side effects of Vanos may include: burning, stinging, itching, or dryness of treated skin

Drug Side Effects Summary

- redness or crusting around your hair follicles
- stretch marks
- spider veins
- acne
- lightened color of treated skin
- headache
- or stuffy nose, sore throat .

1478. This particular disease, Eczema, interacts with the drug Verdeso and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin - weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Steroid medicine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common Verdeso side effects may include: stinging or burning of treated skin
- skin irritation, redness, itching, or hardening
- dry, scaly, or oily skin
- swelling in your hands or feet

Drug Side Effects Summary

- acne, stretch marks
- or redness or crusting around your hair follicles.

1479. This particular disease, Eczema, interacts with the drug Zonalon and has the following side effects:

- hives
 - difficult breathing
 - swelling of your face, lips, tongue, or throat. Although the risk of serious side effects is low when Zonalon is applied to the skin, side effects can occur if the medicine is absorbed into your bloodstream. You may feel drowsy after applying Zonalon, especially if you have treated large areas of skin. Zonalon may cause serious side effects. Stop using Zonalon and call your doctor at once if you have: severe drowsiness
 - confusion
 - severe dry mouth
 - painful or difficult urination
 - blurred vision
 - or new or worsening skin symptoms. Some side effects may be more likely in older adults.
- Common side effects of Zonalon may include: burning or stinging where the medicine was applied
- mild drowsiness
 - or dry mouth.

1480. This particular disease, Eczema, interacts with the drug Ala-Quin and has the following side effects:

- No side effects listed

1481. This particular disease, Eczema, interacts with the drug Alcortin A and has the following side effects:

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Serious side effects are unlikely when this medicine is applied to the skin, but can occur if hydrocortisone or iodoquinol are absorbed into your bloodstream. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe pain, burning, or irritation of treated skin
- itching, oozing, or other signs of a new infection
- any new redness or swelling where the medicine was applied
- increased adrenal gland hormones --weight gain in your face and shoulders, slow wound healing, skin discoloration, thinning skin, increased body hair, tiredness, mood changes, menstrual changes, sexual changes
- decreased adrenal gland hormones --muscle weakness, tiredness, diarrhea , nausea , menstrual changes, skin discoloration, craving salty foods, and feeling light-headed
- or high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor. Common side effects of Alcantin A may include: mild burning, itching, dryness, or irritation of the skin
- redness or crusting around your hair follicles
- acne, increased facial or body hair growth
- changes in the color of treated skin
- or stretch marks.

1482. This particular disease, Eczema, interacts with the drug aloe polysaccharides / iodoquinol and

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives

Drug Side Effects Summary

- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Irritation where iodoquinol is used. Aloe polysaccharides / iodoquinol topical side effects

1483. This particular disease, Eczema, interacts with the drug aloe vera / hydrocortisone / iodoquinol

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Serious side effects are unlikely when this medicine is applied to the skin, but can occur if hydrocortisone or iodoquinol are absorbed into your bloodstream. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe pain, burning, or irritation of treated skin
- itching, oozing, or other signs of a new infection
- any new redness or swelling where the medicine was applied
- increased adrenal gland hormones --weight gain in your face and shoulders, slow wound healing, skin discoloration, thinning skin, increased body hair, tiredness, mood changes, menstrual changes, sexual changes
- decreased adrenal gland hormones --muscle weakness, tiredness, diarrhea , nausea , menstrual changes, skin discoloration, craving salty foods, and feeling light-headed
- or high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor. Common side effects of aloe polysaccharides, hydrocortisone, and iodoquinol may include: mild burning,

Drug Side Effects Summary

itching, dryness, or irritation of the skin

- redness or crusting around your hair follicles
- acne, increased facial or body hair growth
- changes in the color of treated skin
- or stretch marks.

1484. This particular disease, Eczema, interacts with the drug Aloquin and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Irritation where Aloquin (iodoquinol) is used.

Aloquin side effects

1485. This particular disease, Eczema, interacts with the drug ammonium lactate / halobetasol and

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs

Drug Side Effects Summary

of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of high blood sugar like confusion, feeling sleepy, more thirst, more hungry, passing urine more often, flushing, fast breathing, or breath that smells like fruit. Signs of Cushing's disease like weight gain in the upper back or belly, moon face, very bad headache , or slow healing. Signs of a weak adrenal gland like a very bad upset stomach or throwing up, very bad dizziness or passing out, muscle weakness, feeling very tired, mood changes, not hungry, or weight loss . Skin changes (pimples , stretch marks, slow healing, hair growth). Irritation where halobetasol cream and ointment with lactic acid cream is used. Thinning of the skin. Change in eyesight. Ammonium lactate / halobetasol topical side effects

1486. This particular disease, Eczema, interacts with the drug ammonium lactate/urea and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Less serious side effects may include: stinging
- rash
- or skin irritation.

1487. This particular disease, Eczema, interacts with the drug Apexicon and has the following side effects:

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Your skin can absorb topical steroid medicine, which may cause steroid side effects throughout the body. Tell your doctor if you have: swelling, weight gain (especially in your face or your upper back and torso)
- irregular menstrual periods, changes in sexual function
- or memory problems, feeling irritable. ApexiCon may cause serious side effects. Stop using ApexiCon and call your doctor at once if you have: blurred vision, tunnel vision, eye pain , or seeing halos around lights
- severe skin irritation where the medicine was applied
- or signs of skin infection (swelling, redness, warmth, oozing). Steroid medicine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common side effects of ApexiCon may include: redness or crusting around your hair follicles
- burning or itching of treated skin
- lightened color of treated skin
- skin dryness or irritation
- acne, skin rash
- increased hair growth
- stretch marks
- or white or "pruned" appearance of the skin (caused by leaving wound dressings on for long periods of time).

1488. This particular disease, Eczema, interacts with the drug Capex and has the following side effects

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Capex may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Common side effects of Capex may include: burning, itching, or irritation of treated skin
- increased hair growth
- white or "pruned" appearance of the skin
- rash, pimples
- or changes in color of treated skin.

1489. This particular disease, Eczema, interacts with the drug Carb-O-Lac5 and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Less serious side effects may include: stinging
- rash
- or skin irritation.

1490. This particular disease, Eczema, interacts with the drug Carb-O-Lac HP and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Less serious side effects may include: stinging
- rash

Drug Side Effects Summary

- or skin irritation.

1491. This particular disease, Eczema, interacts with the drug Clobevate and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Burning or stinging at the application site Less common Burning sensation of the skin dry skin flushing or redness of the skin itching, scaling, severe redness, soreness, or swelling of the skin skin irritation skin rash, encrusted, scaly and oozing thinning of the skin with easy bruising, especially when used on the face or where the skin folds together (eg, between the fingers) Rare Thinning, weakness, or wasting away of the skin Incidence not known Backache blindness blurred vision burning, itching, and pain in hairy areas, or pus at the root of the hair change in vision dry mouth eye pain facial hair growth in females fractures fruit-like breath odor full or round face, neck, or trunk hair loss headache hives increased hunger increased thirst increased urination irritability loss of sexual desire or ability menstrual irregularities muscle wasting nausea redness and scaling around the mouth stomach pain sugar in the urine sweating tearing thinning of the hair troubled breathing unexplained weight loss unusual tiredness or weakness vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Cough sore throat Less common Body aches or pain changes in skin coloring congestion dryness of the throat raised, dark red, wart-like spots on the skin, especially when used on the face skin discomfort stuffy or runny nose tender, swollen glands in the neck trouble swallowing voice changes Incidence not known Acne or pimples burning and itching of the skin with pinhead-sized red blisters increased hair

Drug Side Effects Summary

growth on the forehead, back, arms, and legs lightening of normal skin color lightening of treated areas of dark skin reddish purple lines on the arms, face, legs, trunk, or groin softening of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Clobevate side effects

1492. This particular disease, Eczema, interacts with the drug Clodan and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Clodan may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain in your face and shoulders, slow wound healing, skin discoloration, thinning skin, increased body hair, tiredness, mood changes, menstrual changes, sexual changes. Common side effects of Clodan may include: burning, itching, swelling, or irritation of treated skin
- dry or cracking skin
- redness or crusting around your hair follicles
- spider veins
- stretch marks, thinning skin
- rash or hives
- acne
- or temporary hair loss .

Drug Side Effects Summary

1493. This particular disease, Eczema, interacts with the drug Cordran SP and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Cordran SP may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Common side effects of Cordran SP may include: burning, itching, or irritation of treated skin
- increased hair growth
- white or "pruned" appearance of the skin
- rash, pimples
- or changes in color of treated skin.

1494. This particular disease, Eczema, interacts with the drug Cormax and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Burning or stinging at the application site Less common Burning sensation of the skin dry skin flushing or redness of the skin itching, scaling, severe redness, soreness, or swelling of the skin skin irritation skin rash, encrusted, scaly and oozing thinning of the skin with easy bruising, especially when used on the face or where the skin folds together (eg, between the fingers) Rare Thinning, weakness, or wasting away of the

Drug Side Effects Summary

skin Incidence not known Backache blindness blurred vision burning, itching, and pain in hairy areas, or pus at the root of the hair change in vision dry mouth eye pain facial hair growth in females fractures fruit-like breath odor full or round face, neck, or trunk hair loss headache hives increased hunger increased thirst increased urination irritability loss of sexual desire or ability menstrual irregularities muscle wasting nausea redness and scaling around the mouth stomach pain sugar in the urine sweating tearing thinning of the hair troubled breathing unexplained weight loss unusual tiredness or weakness vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Cough sore throat Less common Body aches or pain changes in skin coloring congestion dryness of the throat raised, dark red, wart-like spots on the skin, especially when used on the face skin discomfort stuffy or runny nose tender, swollen glands in the neck trouble swallowing voice changes Incidence not known Acne or pimples burning and itching of the skin with pinhead-sized red blisters increased hair growth on the forehead, back, arms, and legs lightening of normal skin color lightening of treated areas of dark skin reddish purple lines on the arms, face, legs, trunk, or groin softening of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Cormax side effects

1495. This particular disease, Eczema, interacts with the drug Cormax Scalp and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Burning or stinging at the

Drug Side Effects Summary

application site Less common Burning sensation of the skin dry skin flushing or redness of the skin itching, scaling, severe redness, soreness, or swelling of the skin skin irritation skin rash, encrusted, scaly and oozing thinning of the skin with easy bruising, especially when used on the face or where the skin folds together (eg, between the fingers) Rare Thinning, weakness, or wasting away of the skin Incidence not known Backache blindness blurred vision burning, itching, and pain in hairy areas, or pus at the root of the hair change in vision dry mouth eye pain facial hair growth in females fractures fruit-like breath odor full or round face, neck, or trunk hair loss headache hives increased hunger increased thirst increased urination irritability loss of sexual desire or ability menstrual irregularities muscle wasting nausea redness and scaling around the mouth stomach pain sugar in the urine sweating tearing thinning of the hair troubled breathing unexplained weight loss unusual tiredness or weakness vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Cough sore throat Less common Body aches or pain changes in skin coloring congestion dryness of the throat raised, dark red, wart-like spots on the skin, especially when used on the face skin discomfort stuffy or runny nose tender, swollen glands in the neck trouble swallowing voice changes Incidence not known Acne or pimples burning and itching of the skin with pinhead-sized red blisters increased hair growth on the forehead, back, arms, and legs lightening of normal skin color lightening of treated areas of dark skin reddish purple lines on the arms, face, legs, trunk, or groin softening of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Cormax Scalp side effects

Drug Side Effects Summary

1496. This particular disease, Eczema, interacts with the drug Corti-Sav and has the following side

- No side effects listed

1497. This particular disease, Eczema, interacts with the drug Cyclocort and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Cyclocort may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , headaches, pain behind the eyes, tiredness, mood changes, menstrual changes, sexual changes. Steroid medicine can affect growth in children. Tell your doctor if your child is not growing or gaining weight at a normal rate while using Cyclocort. Common side effects of Cyclocort may include: burning, itching, irritation, or dryness of treated skin
- redness or crusting around your hair follicles
- acne, increased hair growth
- stretch marks, thinning skin
- or white or "pruned" appearance of the skin (caused by covering treated skin with a tight bandage or other covering).

1498. This particular disease, Eczema, interacts with the drug Derma-Smoothe / FS and has the follo

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your

Drug Side Effects Summary

doctor immediately if any of the following side effects occur: More common Body aches or pain congestion cough dryness or soreness of the throat fever headache hoarseness lightening of normal skin color lightening of treated areas of dark skin sore throat stuffy or runny nose tender, swollen glands in the neck trouble swallowing unusual tiredness or weakness voice changes Less common Acne or pimples accumulation of pus blistering, crusting, irritation, itching, or reddening of the skin burning, itching, and pain in hairy areas, or pus at the root of the hair burning and itching of the skin with pinhead-sized red blisters change in hearing cracked, dry, scaly skin diarrhea dry skin earache or pain in the ear ear drainage flushing or redness of the skin darkening of the skin itching, scaling, severe redness, soreness, or swelling of the skin itchy, raised, round, smooth, skin-colored bumps found on just one area of the body ooze thick white fluid raised, dark red, wart-like spots on skin, especially when used on the face redness or swelling in the ear skin irritation skin rash, encrusted, scaly and oozing spots on your skin resembling a blister or pimple swelling swollen, red, tender area of infection thickened patches of the skin vomiting Incidence not known Redness and scaling around the mouth thinning, weakness, or wasting away of the skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Changes in skin color shiny skin Incidence not known increased hair growth on the forehead, back, arms, and legs reddish purple lines on the arms, face, legs, trunk, or groin softening of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Derma-Smoothe / FS side effects

Drug Side Effects Summary

1499. This particular disease, Eczema, interacts with the drug Derma-Smoothe / FS (Body Oil) and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medication and call your doctor at once if you have severe irritation of any treated skin, or if you show signs of absorbing Derma-Smoothe/FS (Body Oil) through your skin, such as: blurred vision, headache
- thinning skin, easy bruising
- mood changes
- weight gain, puffiness in your face
- or muscle weakness, feeling tired. Common side effects of Derma-Smoothe/FS (Body Oil) may include: mild skin itching, burning, dryness, or other irritation
- temporary hair loss
- swollen hair follicles
- blisters, pimples , or crusting of treated skin
- changes in color of treated skin
- skin rash or irritation around your mouth
- fever, cough, sore throat , runny or stuffy nose
- or red spots or the appearance of tiny veins (especially around your nose, cheeks, or chin).

1500. This particular disease, Eczema, interacts with the drug Dermatop and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Burning, itching, drying, scaling, or cracking of the skin irritation of the skin pain raised, dark red, or wart-like spots on the skin, especially when used on the face shininess or thinness of the skin thinning of the skin with

Drug Side Effects Summary

easy bruising, especially when used on the face or where the skin folds together (e.g., between the fingers) Rare Blistering, burning, crusting, dryness, or flaking of the skin burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings hives or welts itching, scaling, severe redness, soreness, or swelling of the skin skin rash Incidence not known Irritation of the skin around the mouth redness and scaling around the mouth Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Acne or pimples burning and itching of the skin with pinhead-sized red blisters burning, itching, and pain in hairy areas, or pus at the root of the hair increased hair growth on the forehead, back, arms, and legs lightening of normal skin color lightening of the treated areas of dark skin reddish purple lines on the arms, face, legs, trunk, or groin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Dermatop side effects

1501. This particular disease, Eczema, interacts with the drug Dermazene and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever

Drug Side Effects Summary

- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of a weak adrenal gland like a very bad upset stomach or throwing up, very bad dizziness or passing out, muscle weakness, feeling very tired, mood changes, not hungry, or weight loss . Signs of Cushing's disease like weight gain in the upper back or belly, moon face, very bad headache , or slow healing. Signs of high blood sugar like confusion, feeling sleepy, more thirst, more hungry, passing urine more often, flushing, fast breathing, or breath that smells like fruit. Skin changes (pimples , stretch marks, slow healing, hair growth). Irritation where Dermazene (iodoquinol and hydrocortisone cream) is used. Change in color of skin. Thinning of the skin. Dermazene side effects

1502. This particular disease, Eczema, interacts with the drug DesRx and has the following side effects

- No side effects listed

1503. This particular disease, Eczema, interacts with the drug diflorasone and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Your skin can absorb topical steroid medicine, which may cause steroid side effects throughout the body. Tell your doctor if you have: swelling, weight gain (especially in your face or your upper back and torso)
- irregular menstrual periods, changes in sexual function
- or memory problems, feeling irritable. Diflorasone topical may cause serious side effects. Stop using diflorasone topical and call your doctor at once if you have: blurred vision, tunnel vision, eye

Drug Side Effects Summary

pain , or seeing halos around lights

- severe skin irritation where the medicine was applied
- or signs of skin infection (swelling, redness, warmth, oozing). Steroid medicine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common side effects of diflorasone topical may include: redness or crusting around your hair follicles
- burning or itching of treated skin
- lightened color of treated skin
- skin dryness or irritation
- acne, skin rash
- increased hair growth
- stretch marks
- or white or "pruned" appearance of the skin (caused by leaving wound dressings on for long periods of time).

1504. This particular disease, Eczema, interacts with the drug Embeline and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Burning or stinging at the application site Less common Burning sensation of the skin dry skin flushing or redness of the skin itching, scaling, severe redness, soreness, or swelling of the skin skin irritation skin rash, encrusted, scaly and oozing thinning of the skin with easy bruising, especially when used on the face or where the skin folds together (eg, between the fingers) Rare Thinning, weakness, or wasting away of the skin Incidence not known Backache blindness blurred vision burning, itching, and pain in hairy areas, or pus at the root of the hair change in vision dry mouth eye pain facial hair growth in females

Drug Side Effects Summary

fractures fruit-like breath odor full or round face, neck, or trunk hair loss headache hives increased hunger increased thirst increased urination irritability loss of sexual desire or ability menstrual irregularities muscle wasting nausea redness and scaling around the mouth stomach pain sugar in the urine sweating tearing thinning of the hair troubled breathing unexplained weight loss unusual tiredness or weakness vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Cough sore throat Less common Body aches or pain changes in skin coloring congestion dryness of the throat raised, dark red, wart-like spots on the skin, especially when used on the face skin discomfort stuffy or runny nose tender, swollen glands in the neck trouble swallowing voice changes Incidence not known Acne or pimples burning and itching of the skin with pinhead-sized red blisters increased hair growth on the forehead, back, arms, and legs lightening of normal skin color lightening of treated areas of dark skin reddish purple lines on the arms, face, legs, trunk, or groin softening of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Embeline side effects

1505. This particular disease, Eczema, interacts with the drug Embeline E and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Burning or stinging at the application site Less common Burning sensation of the skin dry skin flushing or redness of the skin itching, scaling, severe redness, soreness, or swelling of the skin skin irritation skin rash, encrusted,

Drug Side Effects Summary

scaly and oozing thinning of the skin with easy bruising, especially when used on the face or where the skin folds together (eg, between the fingers) Rare Thinning, weakness, or wasting away of the skin Incidence not known Backache blindness blurred vision burning, itching, and pain in hairy areas, or pus at the root of the hair change in vision dry mouth eye pain facial hair growth in females fractures fruit-like breath odor full or round face, neck, or trunk hair loss headache hives increased hunger increased thirst increased urination irritability loss of sexual desire or ability menstrual irregularities muscle wasting nausea redness and scaling around the mouth stomach pain sugar in the urine sweating tearing thinning of the hair troubled breathing unexplained weight loss unusual tiredness or weakness vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Cough sore throat Less common Body aches or pain changes in skin coloring congestion dryness of the throat raised, dark red, wart-like spots on the skin, especially when used on the face skin discomfort stuffy or runny nose tender, swollen glands in the neck trouble swallowing voice changes Incidence not known Acne or pimples burning and itching of the skin with pinhead-sized red blisters increased hair growth on the forehead, back, arms, and legs lightening of normal skin color lightening of treated areas of dark skin reddish purple lines on the arms, face, legs, trunk, or groin softening of the skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Embeline E side effects

1506. This particular disease, Eczema, interacts with the drug Flac and has the following side effects

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe irritation in or around your ear
- or signs of a skin infection--warmth, redness, swelling, sores, oozing. Your skin can absorb topical steroid medicine, which may cause steroid side effects throughout the body. Tell your doctor if you have: swelling, weight gain (especially in your face or your upper back and torso)
- thinning skin, stretch marks, acne, increased facial hair
- memory problems, feeling irritable
- irregular menstrual periods, changes in sexual function
- or growth delay in a child. Children can absorb larger amounts of this medicine through the skin and may be more likely to have side effects. Common side effects of Flac Otic Oil may include: burning, itching, irritation, or dryness
- rash around the ear in which this medicine was applied
- rash or irritation around your mouth
- acne, pimples , stretch marks
- lightened skin color
- or redness or crusting around your hair follicles.

1507. This particular disease, Eczema, interacts with the drug halcinonide and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Halcinonide topical may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor

Drug Side Effects Summary

- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso)
- slow wound healing, thinning skin, increased body hair
- changes in sexual function
- feeling depressed, anxious, or irritable. Common side effects of halcinonide topical may include:
burning, itching, or irritation of treated skin
- increased hair growth
- white or "pruned" appearance of the skin
- rash, pimples
- or changes in color of treated skin.

1508. This particular disease, Eczema, interacts with the drug Halonate and has the following side effects:

- No side effects listed

1509. This particular disease, Eczema, interacts with the drug Kerasal Ultra 20 and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking

Drug Side Effects Summary

- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Redness. Irritation where Kerasal Ultra 20 (urea in ammonium lactate cream) is used. Kerasal Ultra 20 side effects

1510. This particular disease, Eczema, interacts with the drug LoKara and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. LoKara may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Steroid medicine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common side effects of LoKara may include: stinging or burning of treated skin
- skin irritation, redness, itching, or hardening
- dry, scaly, or oily skin
- swelling in your hands or feet
- acne, stretch marks
- or redness or crusting around your hair follicles.

1511. This particular disease, Eczema, interacts with the drug Nolix and has the following side effects

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Nolix may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Common side effects of Nolix may include: burning, itching, or irritation of treated skin
- increased hair growth
- white or "pruned" appearance of the skin
- rash, pimples
- or changes in color of treated skin.

1512. This particular disease, Eczema, interacts with the drug Olux and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Olux may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain in your face and shoulders, slow wound healing, skin discoloration, thinning skin, increased body hair, tiredness,

Drug Side Effects Summary

mood changes, menstrual changes, sexual changes. Common side effects of Olux may include:
burning, itching, swelling, or irritation of treated skin

- dry or cracking skin
- redness or crusting around your hair follicles
- spider veins
- stretch marks, thinning skin
- rash or hives
- acne
- or temporary hair loss .

1513. This particular disease, Eczema, interacts with the drug Olux-E and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Olux-E may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain in your face and shoulders, slow wound healing, skin discoloration, thinning skin, increased body hair, tiredness, mood changes, menstrual changes, sexual changes. Common side effects of Olux-E may include:
burning, itching, swelling, or irritation of treated skin
- dry or cracking skin
- redness or crusting around your hair follicles
- spider veins

Drug Side Effects Summary

- stretch marks, thinning skin
- rash or hives
- acne
- or temporary hair loss .

1514. This particular disease, Eczema, interacts with the drug prednicarbate and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Prednicarbate topical may cause serious side effects.

Call your doctor at once if you have: worsening of your skin condition

- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Common side effects of prednicarbate topical may include: skin dryness, scaling, cracking or pain
- thinning skin
- burning, itching, or irritation of treated skin
- swelling
- rash
- or numbness , tingling, burning pain.

1515. This particular disease, Eczema, interacts with the drug Quinja and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and

Drug Side Effects Summary

sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Irritation where Quinja (iodoquinol) is used.

Quinja side effects

1516. This particular disease, Eczema, interacts with the drug Synalar and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin - weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Common Synalar side effects may include: burning, itching, or irritation of treated skin
- increased hair growth

Drug Side Effects Summary

- white or "pruned" appearance of the skin
- rash, pimples
- or changes in color of treated skin.

1517. This particular disease, Eczema, interacts with the drug Synalar Ointment and has the following side effects:

- No side effects listed

1518. This particular disease, Eczema, interacts with the drug Tovet and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tovet may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain in your face and shoulders, slow wound healing, skin discoloration, thinning skin, increased body hair, tiredness, mood changes, menstrual changes, sexual changes. Common side effects of Tovet may include: burning, itching, swelling, or irritation of treated skin
- dry or cracking skin
- redness or crusting around your hair follicles
- spider veins
- stretch marks, thinning skin
- rash or hives
- acne

Drug Side Effects Summary

- or temporary hair loss .

1519. This particular disease, Eczema, interacts with the drug Ultralytic and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Less serious side effects may include: stinging
- rash
- or skin irritation.

1520. This particular disease, Eczema, interacts with the drug Ultralytic 2 and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Less serious side effects may include: stinging
- rash
- or skin irritation.

1521. This particular disease, Eczema, interacts with the drug Vytone and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blistering, burning, crusting, dryness, flaking, scaling, severe redness, soreness, or swelling of the skin burning and itching of the skin with pinhead-sized red blisters burning, itching, and pain in hairy areas, or pus at the root of hair reddish purple lines on the arms, face, legs, trunk, or groin redness and scaling around the mouth softening of the skin thinning of the skin with easy bruising, especially when used on the face or where the skin folds together (eg, between the fingers) Some side effects may occur

Drug Side Effects Summary

that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Acne or pimples increased hair growth on the forehead, back, arms, and legs lightening of normal skin color lightening of the treated areas of dark skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Vytone side effects

1522. This particular disease, Erectile Dysfunction, interacts with the drug Cialis and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop and get medical help at once if you have nausea, chest pain, or dizziness during sex. You could be having a life-threatening side effect. Stop using this medicine and call your doctor at once if you have: a light-headed feeling, like you might pass out
- an erection is painful or lasts longer than 4 hours (prolonged erection can damage the penis)
- vision changes or sudden vision loss
- ringing in your ears or sudden hearing loss
- or heart attack symptoms - chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating. Common Cialis side effects may include: headache
- flushing (warmth, redness, or tingly feeling)
- nausea, upset stomach
- stuffy nose
- or muscle pain, back pain, pain in your arms, legs, or back.

Drug Side Effects Summary

1523. This particular disease, Erectile Dysfunction, interacts with the drug sildenafil and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop taking sildenafil and get emergency medical help if you have: heart attack symptoms - chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- vision changes or sudden vision loss
- erection is painful or lasts longer than 4 hours (prolonged erection can damage the penis)
- ringing in your ears, or sudden hearing loss
- irregular heartbeat
- swelling in your hands, ankles, or feet
- shortness of breath
- seizure (convulsions)
- or a light-headed feeling, like you might pass out. Common sildenafil side effects may include: flushing (warmth, redness, or tingly feeling)
- headache, dizziness
- heartburn, nausea, or upset stomach abnormal vision (blurred vision, changes in color vision) runny or stuffy nose, nosebleeds
- sleep problems (insomnia)
- or muscle pain, back pain.

1524. This particular disease, Erectile Dysfunction, interacts with the drug Viagra and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop taking Viagra and get emergency medical help if

Drug Side Effects Summary

you have: heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating

- vision changes or sudden vision loss
- or erection is painful or lasts longer than 4 hours (prolonged erection can damage the penis). Call your doctor at once if you have: ringing in your ears, or sudden hearing loss
- irregular heartbeat
- swelling in your hands, ankles, or feet
- shortness of breath
- seizure (convulsions)
- or a light-headed feeling, like you might pass out. Common Viagra side effects may include: flushing (warmth, redness, or tingly feeling)
- headache, dizziness
- abnormal vision (blurred vision, changes in color vision) runny or stuffy nose, nosebleeds
- sleep problems (insomnia)
- muscle pain, back pain
- or upset stomach.

1525. This particular disease, Erectile Dysfunction, interacts with the drug tadalafil and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop and get medical help at once if you have nausea, chest pain, or dizziness during sex. You could be having a life-threatening side effect. Stop using tadalafil and call your doctor at once if you have: a light-headed feeling, like you might pass out
- an erection is painful or lasts longer than 4 hours (prolonged erection can damage the penis)
- vision changes or sudden vision loss

Drug Side Effects Summary

- ringing in your ears or sudden hearing loss
- or heart attack symptoms - chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating. Common tadalafil side effects may include: headache
- flushing (warmth, redness, or tingly feeling)
- nausea, upset stomach
- stuffy nose
- or muscle pain, back pain, pain in your arms, legs, or back.

1526. This particular disease, Erectile Dysfunction, interacts with the drug Caverject Impulse and has the following side effects:

- No side effects listed

1527. This particular disease, Erectile Dysfunction, interacts with the drug Staxyn and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop taking this medicine and get emergency medical help if you have: heart attack symptoms - chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- vision changes or sudden vision loss
- or erection is painful or lasts longer than 4 hours (prolonged erection can damage the penis). Stop using Staxyn and call your doctor at once if you have: ringing in your ears, or sudden hearing loss
- irregular heartbeat
- swelling in your hands, ankles, or feet
- shortness of breath
- a light-headed feeling, like you might pass out
- penis erection that is painful or lasts 4 hours or longer

Drug Side Effects Summary

- or seizure (convulsions). Common Staxyn side effects may include: flushing (warmth, redness, or tingly feeling)
- stuffy nose, sinus pain
- headache , dizziness
- upset stomach
- or back pain .

1528. This particular disease, Erectile Dysfunction, interacts with the drug vardenafil and has the fo

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop taking vardenafil and get emergency medical help if you have: heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating
- vision changes or sudden vision loss
- or erection is painful or lasts longer than 4 hours (prolonged erection can damage the penis). Vardenafil may cause serious side effects. Stop using vardenafil and call your doctor at once if you have: ringing in your ears, or sudden hearing loss
- irregular heartbeat
- swelling in your hands, ankles, or feet
- shortness of breath
- a light-headed feeling, like you might pass out
- penis erection that is painful or lasts 4 hours or longer
- or seizure (convulsions). Common side effects of vardenafil may include: flushing (warmth, redness, or tingly feeling)
- stuffy nose, sinus pain

Drug Side Effects Summary

- headache , dizziness
- upset stomach
- or back pain .

1529. This particular disease, Erectile Dysfunction, interacts with the drug alprostadil and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Alprostadil may cause serious side effects. Stop using alprostadil and call your doctor at once if you have: a light-headed feeling, like you might pass out
- bleeding after an injection
- a painful erection that lasts 4 hours or longer
- new or worsening pain in your penis
- or redness, swelling, tenderness, lumps, unusual shape or curving of the erect penis. Common side effects of alprostadil may include: mild pain in your penis, urethra, or testicles
- redness of the penis
- or warmth or burning in your urethra. Your sex partner may have side effects such as burning, itching, or irritation of body areas that come into contact with your penis.

1530. This particular disease, Erectile Dysfunction, interacts with the drug Edex and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Curving of penis with pain during erection erection continuing for 4 to 6 hours erection continuing longer than 6 hours with severe and continuing pain of the penis swelling in or pain of the testes Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your

Drug Side Effects Summary

body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Bleeding and pain at the injection site (short-term) painful erection Rare Bruising or clotted blood in the penis at the injection site, usually caused by an incorrect injection Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Edex side effects

1531. This particular disease, Erectile Dysfunction, interacts with the drug Stendra and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using Stendra and get emergency medical help if you have sudden vision loss. During sexual activity, if you become dizzy or nauseated, or have pain, numbness , or tingling in your chest, arms, neck, or jaw, stop and call your doctor right away. You could be having a serious side effect of Stendra. Stop using this medicine and call your doctor at once if you have: vision changes, sudden vision loss
- ringing in your ears, or sudden hearing loss
- pain, swelling, warmth, or redness in one or both legs
- shortness of breath, swelling in your hands or feet
- a light-headed feeling, like you might pass out
- or penis erection that is painful or lasts 4 hours or longer. Common Stendra side effects may include: headache
- flushing (warmth, redness, or tingly feeling)
- cold symptoms such as runny or stuffy nose, sore throat

Drug Side Effects Summary

- or back pain .

1532. This particular disease, Erectile Dysfunction, interacts with the drug Caverject and has the fo

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Curving of penis with pain during erection erection continuing for 4 to 6 hours erection continuing longer than 6 hours with severe and continuing pain of the penis swelling in or pain of the testes Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Bleeding and pain at the injection site (short-term) painful erection Rare Bruising or clotted blood in the penis at the injection site, usually caused by an incorrect injection Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Caverject side effects

1533. This particular disease, Erectile Dysfunction, interacts with the drug Muse and has the follow

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Burning sensation in the urethra pain in the penis or urethra Less common Bleeding and pain in the testes blurred vision confusion dizziness dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position pain in the testes sweating unusual tiredness or weakness Rare Fainting

Drug Side Effects Summary

Symptoms of overdose Painful or prolonged erection of the penis Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Back pain chills cough diarrhea fever general feeling of discomfort or illness headache joint pain loss of appetite muscle aches and pains nausea runny nose shivering sneezing sore throat stuffy nose sweating swelling of the legs trouble sleeping vomiting Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Muse side effects

1534. This particular disease, Erectile Dysfunction, interacts with the drug avanafil and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using avanafil and get emergency medical help if you have sudden vision loss. During sexual activity, if you become dizzy or nauseated, or have pain, numbness , or tingling in your chest, arms, neck, or jaw, stop and call your doctor right away. You could be having a serious side effect of avanafil. Avanafil may cause serious side effects. Stop using avanafil and call your doctor at once if you have: vision changes, sudden vision loss
- ringing in your ears, or sudden hearing loss
- pain, swelling, warmth, or redness in one or both legs
- shortness of breath, swelling in your hands or feet
- a light-headed feeling, like you might pass out
- or penis erection that is painful or lasts 4 hours or longer. Common side effects of avanafil may

Drug Side Effects Summary

include: headache

- flushing (warmth, redness, or tingly feeling)
- cold symptoms such as runny or stuffy nose, sore throat
- or back pain .

1535. This particular disease, Gastrointestinal, interacts with the drug Entereg and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some people had heart attacks while taking Entereg long-term. It is not clear whether alvimopan is the actual cause of a heart attack. Call your doctor at once if you have heart attack symptoms, such as: chest pain or pressure
- pain spreading to your jaw or shoulder
- anxiety , nausea , sweating. Common Entereg side effect may include indigestion . You may be more likely to have unpleasant effects on your stomach if you used any type of opioid medicine shortly before your surgery.

1536. This particular disease, Gastrointestinal, interacts with the drug Lidocaine Viscous and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Lidocaine viscous may cause serious side effects. Stop using lidocaine viscous and call your doctor at once if you have: drowsiness, depression , confusion, feeling nervous or restless
- weakness, slow breathing, slow heart rate
- a light-headed feeling, like you might pass out
- numbness or cold feeling

Drug Side Effects Summary

- vomiting
- or blurred vision. Common side effects of lidocaine viscous may include: unusual or unpleasant taste in the mouth
- or numbness in places where the medicine is accidentally applied.

1537. This particular disease, Gastrointestinal, interacts with the drug alvimopan and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some people had heart attacks while taking alvimopan long-term. It is not clear whether alvimopan is the actual cause of a heart attack. Call your doctor at once if you have heart attack symptoms, such as: chest pain or pressure
- pain spreading to your jaw or shoulder
- anxiety , nausea , sweating. Common side effect may include indigestion . You may be more likely to have unpleasant effects on your stomach if you used any type of opioid medicine shortly before your surgery.

1538. This particular disease, Gastrointestinal, interacts with the drug lidocaine and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Lidocaine topical may cause serious side effects. Call your doctor at once if you have: severe headache or vomiting
- severe burning, stinging, or irritation where the medicine was applied
- swelling or redness
- sudden dizziness or drowsiness after medicine is applied
- confusion, problems with speech or vision, ringing in your ears

Drug Side Effects Summary

- or unusual sensations of temperature. Common side effects include: mild irritation where the medication is applied
- or numbness in places where the medicine is accidentally applied.

1539. This particular disease, Gastrointestinal, interacts with the drug Glydo and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Glydo may cause serious side effects. Call your doctor at once if you have: severe headache or vomiting
- severe burning, stinging, or irritation where the medicine was applied
- swelling or redness
- sudden dizziness or drowsiness after medicine is applied
- confusion, problems with speech or vision, ringing in your ears
- or unusual sensations of temperature. Common side effects include: mild irritation where the medication is applied
- or numbness in places where the medicine is accidentally applied.

1540. This particular disease, Gastrointestinal, interacts with the drug Xylocaine Jelly and has the f

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever

Drug Side Effects Summary

- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of infection like fever, chills, very bad sore throat , ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal. Signs of too much acid in the blood (acidosis) like confusion
- fast breathing
- fast heartbeat
- a heartbeat that does not feel normal
- very bad stomach pain, upset stomach , or throwing up
- feeling very sleepy
- shortness of breath
- or feeling very tired or weak. Signs of methemoglobinemia like a blue or gray color of the lips, nails, or skin
- a heartbeat that does not feel normal
- seizures
- very bad dizziness or passing out
- very bad headache
- feeling very sleepy
- feeling tired or weak
- or shortness of breath. This effect is rare but may be deadly if it happens. Trouble breathing, slow breathing, or shallow breathing. Very bad numbness and tingling. Feeling lightheaded, sleepy, confused, or having blurred eyesight. Seizures. Change in eyesight. Feeling nervous and excitable.

Drug Side Effects Summary

Dizziness or passing out. Ringing in ears. Upset stomach or throwing up. Feeling hot or cold. Shakiness. Twitching. Slow heartbeat . Chest pain. Xylocaine Jelly side effects

1541. This particular disease, Gastrointestinal, interacts with the drug LTA II Kit and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. LTA II Kit may cause serious side effects. Stop using LTA II Kit and call your doctor at once if you have: drowsiness, depression , confusion, feeling nervous or restless
- weakness, slow breathing, slow heart rate
- a light-headed feeling, like you might pass out
- numbness or cold feeling
- vomiting
- or blurred vision. Common side effects of LTA II Kit may include: unusual or unpleasant taste in the mouth
- or numbness in places where the medicine is accidentally applied.

1542. This particular disease, GERD (Heartburn), interacts with the drug omeprazole and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- new or unusual pain in your wrist, thigh, hip, or back
- seizure (convulsions)
- kidney problems - fever, rash, nausea, loss of appetite, joint pain, urinating less than usual, blood

Drug Side Effects Summary

in your urine, weight gain

- low magnesium - dizziness , irregular heartbeats, feeling jittery, muscle cramps, muscle spasms, cough or choking feeling
- or new or worsening symptoms of lupus - joint pain, and a skin rash on your cheeks or arms that worsens in sunlight. Taking omeprazole long-term may cause you to develop stomach growths called fundic gland polyps. Talk with your doctor about this risk. If you use this medicine for longer than 3 years, you could develop a vitamin B-12 deficiency. Talk to your doctor about how to manage this condition if you develop it. Common omeprazole side effects may include: cold symptoms such as stuffy nose, sneezing, sore throat (especially in children)
- fever (especially in children)
- stomach pain, gas
- nausea, vomiting, diarrhea
- or headache .

1543. This particular disease, GERD (Heartburn), interacts with the drug pantoprazole and has the f

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- sudden pain or trouble moving your hip, wrist, or back
- bruising or swelling where intravenous pantoprazole was injected
- kidney problems - fever, rash, nausea, loss of appetite, joint pain, urinating less than usual, blood in your urine, weight gain
- low magnesium - dizziness, fast or irregular heart rate, tremors (shaking) or jerking muscle movements, feeling jittery, muscle cramps, muscle spasms in your hands and feet, cough or choking

Drug Side Effects Summary

feeling

- or new or worsening symptoms of lupus - joint pain, and a skin rash on your cheeks or arms that worsens in sunlight. Taking pantoprazole long-term may cause you to develop stomach growths called fundic gland polyps. Talk with your doctor about this risk. If you use pantoprazole for longer than 3 years, you could develop a vitamin B-12 deficiency. Talk to your doctor about how to manage this condition if you develop it. Common pantoprazole side effects may include: headache , dizziness

- stomach pain, gas, nausea , vomiting , diarrhea

- joint pain

- or fever , rash, or cold symptoms (most common in children).

1544. This particular disease, GERD (Heartburn), interacts with the drug Nexium and has the following

- hives

- difficulty breathing

- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody

- seizure (convulsions)

- kidney problems - fever, rash, nausea, loss of appetite, joint pain, urinating less than usual, blood in your urine, weight gain

- low magnesium - dizziness, fast or irregular heart rate, tremors (shaking) or jerking muscle movements, feeling jittery, muscle cramps, muscle spasms in your hands and feet, cough or choking feeling

- or new or worsening symptoms of lupus - joint pain, and a skin rash on your cheeks or arms that worsens in sunlight. Taking esomeprazole long-term may cause you to develop stomach growths called fundic gland polyps. Talk with your doctor about this risk. If you use esomeprazole for longer

Drug Side Effects Summary

than 3 years, you could develop a vitamin B-12 deficiency. Talk to your doctor about how to manage this condition if you develop it. Common Nexium side effects may include: headache

- diarrhea
- nausea , stomach pain, gas, constipation
- or dry mouth.

1545. This particular disease, GERD (Heartburn), interacts with the drug famotidine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: confusion, hallucinations, agitation , lack of energy
- a seizure
- fast or pounding heartbeats, sudden dizziness (like you might pass out)
- or unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, and dark colored urine. Some side effects may be more likely in older adults and in people who have severe kidney disease. Common famotidine side effects may include: headache
- dizziness
- or constipation or diarrhea .

1546. This particular disease, GERD (Heartburn), interacts with the drug Protonix and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- sudden pain or trouble moving your hip, wrist, or back

Drug Side Effects Summary

- bruising or swelling where intravenous pantoprazole was injected
- kidney problems - fever, rash, nausea, loss of appetite, joint pain, urinating less than usual, blood in your urine, weight gain
- low magnesium - dizziness, fast or irregular heart rate, tremors (shaking) or jerking muscle movements, feeling jittery, muscle cramps, muscle spasms in your hands and feet, cough or choking feeling
- or new or worsening symptoms of lupus - joint pain, and a skin rash on your cheeks or arms that worsens in sunlight. Taking pantoprazole long-term may cause you to develop stomach growths called fundic gland polyps. Talk with your doctor about this risk. If you use pantoprazole for longer than 3 years, you could develop a vitamin B-12 deficiency. Talk to your doctor about how to manage this condition if you develop it. Common Protonix side effects may include: headache , dizziness
- stomach pain, gas, nausea , vomiting , diarrhea
- joint pain
- or fever , rash, or cold symptoms (most common in children).

1547. This particular disease, GERD (Heartburn), interacts with the drug Prilosec and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using Prilosec and call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- new or unusual pain in your wrist, thigh, hip, or back
- seizure (convulsions)
- kidney problems - fever, rash, nausea, loss of appetite, joint pain, urinating less than usual, blood in your urine, weight gain
- low magnesium - dizziness , irregular heartbeats, feeling jittery, muscle cramps, muscle spasms,

Drug Side Effects Summary

cough or choking feeling

- or new or worsening symptoms of lupus - joint pain, and a skin rash on your cheeks or arms that worsens in sunlight. Taking Prilosec long-term may cause you to develop stomach growths called fundic gland polyps. Talk with your doctor about this risk. If you use omeprazole for longer than 3 years, you could develop a vitamin B-12 deficiency. Talk to your doctor about how to manage this condition if you develop it. Common Prilosec side effects may include: cold symptoms such as stuffy nose, sneezing, sore throat (especially in children)
- fever (especially in children)
- stomach pain, gas
- nausea, vomiting, diarrhea
- or headache .

1548. This particular disease, GERD (Heartburn), interacts with the drug Dexilant and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Dexilant may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- a seizure (convulsions)
- sudden pain or trouble moving your hip, wrist, or back
- kidney problems-- fever, rash, nausea , loss of appetite, joint pain, urinating less than usual, blood in your urine, weight gain
- low magnesium-- dizziness , fast or irregular heart rate, tremors (shaking) or jerking muscle movements, feeling jittery, muscle cramps, muscle spasms in your hands and feet, cough or choking feeling
- or new or worsening symptoms of lupus--joint pain, and a skin rash on your cheeks or arms that

Drug Side Effects Summary

worsens in sunlight. Taking Dexilant long-term may cause you to develop stomach growths called fundic gland polyps. Talk with your doctor about this risk. If you use Dexilant for longer than 3 years, you could develop a vitamin B-12 deficiency. Talk to your doctor about how to manage this condition if you develop it. Common side effects of Dexilant may include: nausea, vomiting , stomach pain, gas

- diarrhea
- headache
- mouth pain, sore throat
- or stuffy nose, sinus pain, or other cold symptoms .

1549. This particular disease, GERD (Heartburn), interacts with the drug Aciphex and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- sudden pain or trouble moving your hip, wrist, or back
- seizure (convulsions)
- kidney problems - fever, rash, nausea , loss of appetite, joint pain, urinating less than usual, blood in your urine, swelling, rapid weight gain
- new or worsening symptoms of lupus - joint pain, and a skin rash on your cheeks or arms that worsens in sunlight
- low magnesium - dizziness , fast or irregular heart rate, tremors (shaking) or jerking muscle movements, feeling jittery, muscle cramps, muscle spasms in your hands and feet, cough or choking feeling
- or signs of bleeding (if you also take warfarin - headaches, dizziness, weakness

Drug Side Effects Summary

- pain or swelling
- bruising, unusual bleeding (nosebleeds , bleeding gums)
- red or pink urine
- heavy menstrual flow
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- or any bleeding that will not stop. Taking rabeprazole long-term may cause you to develop stomach growths called fundic gland polyps. Talk with your doctor about this risk. If you use AcipHex for longer than 3 years, you could develop a vitamin B-12 deficiency. Talk to your doctor about how to manage this condition if you develop it. Common AcipHex side effects may include: infection
- pain
- sore throat
- headache
- nausea, vomiting
- gas, diarrhea, constipation
- or stomach pain.

1550. This particular disease, GERD (Heartburn), interacts with the drug Prilosec OTC and has the f

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Prilosec OTC may cause serious side effects. Stop using Prilosec OTC and call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- new or unusual pain in your wrist, thigh, hip, or back
- seizure (convulsions)
- kidney problems--fever, rash, nausea, loss of appetite, joint pain, urinating less than usual, blood in

Drug Side Effects Summary

your urine, weight gain

- low magnesium-- dizziness , irregular heartbeats, feeling jittery, muscle cramps, muscle spasms, cough or choking feeling
- or new or worsening symptoms of lupus--joint pain, and a skin rash on your cheeks or arms that worsens in sunlight. Taking Prilosec OTC long-term may cause you to develop stomach growths called fundic gland polyps. Talk with your doctor about this risk. If you use Prilosec OTC for longer than 3 years, you could develop a vitamin B-12 deficiency. Talk to your doctor about how to manage this condition if you develop it. Common side effects of Prilosec OTC may include: cold symptoms such as stuffy nose, sneezing, sore throat (especially in children)
- fever (especially in children)
- stomach pain, gas
- nausea, vomiting, diarrhea
- or headache .

1551. This particular disease, GERD (Heartburn), interacts with the drug esomeprazole and has the

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Esomeprazole may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- seizure (convulsions)
- kidney problems-- fever, rash, nausea , loss of appetite, joint pain, urinating less than usual, blood in your urine, weight gain
- low magnesium-- dizziness , fast or irregular heart rate, tremors (shaking) or jerking muscle movements, feeling jittery, muscle cramps, muscle spasms in your hands and feet, cough or choking feeling

Drug Side Effects Summary

- or new or worsening symptoms of lupus--joint pain, and a skin rash on your cheeks or arms that worsens in sunlight. Taking esomeprazole long-term may cause you to develop stomach growths called fundic gland polyps. Talk with your doctor about this risk. If you use esomeprazole for longer than 3 years, you could develop a vitamin B-12 deficiency. Talk to your doctor about how to manage this condition if you develop it. Common side effects of esomeprazole may include: headache

- diarrhea
- nausea, stomach pain, gas, constipation
- or dry mouth.

1552. This particular disease, GERD (Heartburn), interacts with the drug lansoprazole and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- new or unusual pain in your wrist, back, hip, or thigh
- a seizure (convulsions)
- kidney problems - fever, nausea, little or no urination, blood in your urine, swelling, rapid weight gain
- low magnesium - dizziness, fast or irregular heart rate, tremors (shaking) or jerking muscle movements, feeling jittery, muscle cramps, muscle spasms in your hands and feet, cough or choking feeling
- or new or worsening symptoms of lupus - joint pain, and a skin rash on your cheeks or arms that worsens in sunlight. Taking lansoprazole long-term may cause you to develop stomach growths called fundic gland polyps. Talk with your doctor about this risk. If you use lansoprazole for longer than 3 years, you could develop a vitamin B-12 deficiency. Talk to your doctor about how to manage

Drug Side Effects Summary

this condition if you develop it. Common lansoprazole side effects may include: nausea, stomach pain

- diarrhea, constipation
- or headache .

1553. This particular disease, GERD (Heartburn), interacts with the drug Prevacid and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- new or unusual pain in your wrist, back, hip, or thigh
- a seizure (convulsions)
- kidney problems - fever, nausea, little or no urination, blood in your urine, swelling, rapid weight gain
- low magnesium - dizziness, fast or irregular heart rate, tremors (shaking) or jerking muscle movements, feeling jittery, muscle cramps, muscle spasms in your hands and feet, cough or choking feeling
- or new or worsening symptoms of lupus - joint pain, and a skin rash on your cheeks or arms that worsens in sunlight. Common Prevacid side effects may include: nausea, stomach pain
- diarrhea, constipation
- or headache .

1554. This particular disease, GERD (Heartburn), interacts with the drug Pepcid and has the following

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: confusion, hallucinations, agitation , lack of energy
- a seizure
- fast or pounding heartbeats, sudden dizziness (like you might pass out)
- or unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, and dark colored urine. Some side effects may be more likely in older adults and in people who have severe kidney disease. Common Pepcid side effects may include: headache
- dizziness
- or constipation or diarrhea .

1555. This particular disease, GERD (Heartburn), interacts with the drug rabeprazole and has the fo

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Rabeprazole may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- sudden pain or trouble moving your hip, wrist, or back
- seizure (convulsions)
- kidney problems--fever, rash, nausea , loss of appetite, joint pain, urinating less than usual, blood in your urine, swelling, rapid weight gain
- new or worsening symptoms of lupus--joint pain, and a skin rash on your cheeks or arms that worsens in sunlight
- low magnesium-- dizziness , fast or irregular heart rate, tremors (shaking) or jerking muscle movements, feeling jittery, muscle cramps, muscle spasms in your hands and feet, cough or choking feeling
- or signs of bleeding (if you also take warfarin)--headaches, dizziness, weakness

Drug Side Effects Summary

- pain or swelling
- bruising, unusual bleeding (nosebleeds , bleeding gums)
- red or pink urine
- heavy menstrual flow
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- or any bleeding that will not stop. Taking rabeprazole long-term may cause you to develop stomach growths called fundic gland polyps. Talk with your doctor about this risk. If you use rabeprazole for longer than 3 years, you could develop a vitamin B-12 deficiency. Talk to your doctor about how to manage this condition if you develop it. Common side effects of rabeprazole may include: infection
- pain
- sore throat
- headache
- nausea, vomiting
- gas, diarrhea, constipation
- or stomach pain.

1556. This particular disease, GERD (Heartburn), interacts with the drug sucralfate and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common side effects of sucralfate may include: constipation , diarrhea
- nausea , vomiting , gas, indigestion
- itching, rash
- dizziness , drowsiness

Drug Side Effects Summary

- sleep problems (insomnia)
- headache
- or back pain .

1557. This particular disease, GERD (Heartburn), interacts with the drug Pepcid AC and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Anxiety bleeding gums blistering, peeling, or loosening of the skin blood in the urine or stools bloody, black, or tarry stools cough difficulty breathing discouragement fast, irregular, pounding, or racing heartbeat or pulse feeling sad or empty irritability lack of appetite loss of interest or pleasure noisy breathing pinpoint red spots on the skin seeing, hearing, or feeling things that are not there seizures swelling around the eyes tightness in the chest trouble concentrating trouble sleeping unusual bleeding or bruising Incidence not known Agitation blistering, peeling, or loosening of the skin burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain chills clay-colored stools dark urine diarrhea difficulty swallowing dizziness or fainting fever general feeling of tiredness or weakness headache high fever hives, itching, skin rash hoarseness joint or muscle pain large, hive-like swelling on face, eyelids, lips, tongue, throat, hands, legs, feet, or genitals light-colored stools lower back or side pain nausea painful or difficult urination pale skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red skin lesions, often with a purple center red, irritated eyes sore throat sores, ulcers, or white spots in the mouth or on the lips stomach pain swollen glands unpleasant breath odor unusual tiredness or weakness upper right abdominal pain vomiting of blood yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side

Drug Side Effects Summary

effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Difficulty having a bowel movement Rare Bad, unusual, or unpleasant (after) taste bone or muscle pain change in taste continuing ringing or buzzing or other unexplained noise in the ears decreased interest in sexual intercourse difficulty in moving dry mouth dry skin feeling of warmth hearing loss inability to have or keep an erection lack or loss of strength loss in sexual ability, desire, drive, or performance muscle pain or stiffness redness of the face, neck, arms and occasionally, upper chest redness of the white part of the eyes stomach discomfort unusual drowsiness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Pepcid AC side effects

1558. This particular disease, GERD (Heartburn), interacts with the drug Carafate and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common Carafate side effects may include:
constipation , diarrhea
- nausea , vomiting , gas, indigestion
- itching, rash
- dizziness , drowsiness
- sleep problems (insomnia)
- headache
- or back pain .

1559. This particular disease, GERD (Heartburn), interacts with the drug Nexium 24HR and has the following side effects:

Drug Side Effects Summary

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

- itching

- red, swollen, blistered, or peeling skin with or without fever

- wheezing

- tightness in the chest or throat

- trouble breathing, swallowing, or talking

- unusual hoarseness

- or swelling of the mouth, face, lips, tongue, or throat. Signs of low magnesium levels like mood changes, muscle pain or weakness, muscle cramps or spasms, seizures, shakiness, not hungry, very bad upset stomach or throwing up, or a heartbeat that does not feel normal. Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain. Very bad dizziness or passing out. Bone pain. Fever, chills, or sore throat . Shortness of breath. A big weight loss . This medicine may raise the chance of a severe form of diarrhea called C diff-associated diarrhea (CDAD). Call your doctor right away if you have stomach pain or cramps, very loose or watery stools , or bloody stools. Do not try to treat diarrhea without first checking with your doctor. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)

- red or irritated eyes

- or sores in your mouth, throat, nose, or eyes. Nexium 24HR side effects

Drug Side Effects Summary

1560. This particular disease, GERD (Heartburn), interacts with the drug Protonix IV and has the fol

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: Less common Accumulation of pus bleeding, blistering, burning, coldness, discoloration of the skin, feeling of pressure, hives, infection, inflammation, itching, lumps, numbness, pain, rash, redness, scarring, soreness, stinging, swelling, tenderness, tingling, ulceration, or warmth at the injection site changes in skin color fever pain, tenderness, or swelling of the foot or leg stomach pain swollen, red, tender area of infection Rare Drowsiness loss of appetite mood or mental changes muscle spasms (tetany) or twitching nausea seizures trembling unusual tiredness or weakness vomiting Incidence not known Absence of or decrease in body movement black, tarry stools blindness blistering, peeling, or loosening of the skin bloating of the stomach bloody or cloudy urine bloody, black, or tarry stools blurred vision chest pain or tightness chills confusion constipation continuing ringing or buzzing or other unexplained noise in the ears cough dark urine decreased urine output decreased vision diarrhea difficulty with speaking difficulty with swallowing dizziness or lightheadedness dry mouth fast heartbeat feeling of constant movement of self or surroundings greatly decreased frequency of urination or amount of urine hearing loss hives, itching, or rash increased thirst increased watering of the mouth indigestion joint or muscle pain large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or genitals light-colored stools muscle cramps, pain, or stiffness numbness and tingling around the mouth, fingertips, or feet painful or difficult urination pains in the stomach, side, or abdomen, possibly radiating to the back pale skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red skin lesions, often with a purple center red, irritated eyes sensation of spinning severe stomach cramps or pain sore throat sores, ulcers, or white spots on the lips or in the mouth stomach cramps or tenderness swelling of the feet or lower legs swollen glands trouble breathing unexplained bleeding or bruising watery and severe diarrhea, which may also be bloody yellow eyes

Drug Side Effects Summary

or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Headache Less common Belching excess air or gas in the stomach or bowels full feeling passing gas Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Protonix IV side effects

1561. This particular disease, GERD (Heartburn), interacts with the drug Acid Controller Maximum Strength

- No side effects listed

1562. This particular disease, GERD (Heartburn), interacts with the drug Acid Controller Original Strength

- No side effects listed

1563. This particular disease, GERD (Heartburn), interacts with the drug Acid Reducer Maximum Strength

- No side effects listed

1564. This particular disease, GERD (Heartburn), interacts with the drug cimetidine and has the following side effects:

- pain when swallowing
- bloody or tarry stools, cough with bloody mucus or vomit that looks like coffee grounds
- changes in mood, anxiety , agitation
- confusion, hallucinations
- or breast swelling or tenderness. Serious side effects may be more likely in older adults and those

Drug Side Effects Summary

who are ill or debilitated. Common side effects of cimetidine may include: headache

- or diarrhea .

1565. This particular disease, GERD (Heartburn), interacts with the drug dextansoprazole and has the following side effects:

- hives

- difficulty breathing

- swelling of your face, lips, tongue, or throat. Dextansoprazole may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody

- a seizure (convulsions)

- sudden pain or trouble moving your hip, wrist, or back

- kidney problems-- fever, rash, nausea , loss of appetite, joint pain, urinating less than usual, blood in your urine, weight gain

- low magnesium-- dizziness , fast or irregular heart rate, tremors (shaking) or jerking muscle movements, feeling jittery, muscle cramps, muscle spasms in your hands and feet, cough or choking feeling

- or new or worsening symptoms of lupus--joint pain, and a skin rash on your cheeks or arms that worsens in sunlight. Taking dextansoprazole long-term may cause you to develop stomach growths called fundic gland polyps. Talk with your doctor about this risk. If you use dextansoprazole for longer than 3 years, you could develop a vitamin B-12 deficiency. Talk to your doctor about how to manage this condition if you develop it. Common side effects of dextansoprazole may include: nausea, vomiting , stomach pain, gas

- diarrhea

- headache

- mouth pain, sore throat

- or stuffy nose, sinus pain, or other cold symptoms .

Drug Side Effects Summary

1566. This particular disease, GERD (Heartburn), interacts with the drug Heartburn Relief and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Anxiety bleeding gums blistering, peeling, or loosening of the skin blood in the urine or stools bloody, black, or tarry stools cough difficulty breathing discouragement fast, irregular, pounding, or racing heartbeat or pulse feeling sad or empty irritability lack of appetite loss of interest or pleasure noisy breathing pinpoint red spots on the skin seeing, hearing, or feeling things that are not there seizures swelling around the eyes tightness in the chest trouble concentrating trouble sleeping unusual bleeding or bruising Incidence not known Agitation blistering, peeling, or loosening of the skin burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain chills clay-colored stools dark urine diarrhea difficulty swallowing dizziness or fainting fever general feeling of tiredness or weakness headache high fever hives, itching, skin rash hoarseness joint or muscle pain large, hive-like swelling on face, eyelids, lips, tongue, throat, hands, legs, feet, or genitals light-colored stools lower back or side pain nausea painful or difficult urination pale skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red skin lesions, often with a purple center red, irritated eyes sore throat sores, ulcers, or white spots in the mouth or on the lips stomach pain swollen glands unpleasant breath odor unusual tiredness or weakness upper right abdominal pain vomiting of blood yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Difficulty having a bowel movement Rare Bad, unusual, or unpleasant (after) taste bone or muscle pain change in taste continuing ringing or buzzing or other unexplained noise in the ears decreased interest in sexual

Drug Side Effects Summary

intercourse difficulty in moving dry mouth dry skin feeling of warmth hearing loss inability to have or keep an erection lack or loss of strength loss in sexual ability, desire, drive, or performance muscle pain or stiffness redness of the face, neck, arms and occasionally, upper chest redness of the white part of the eyes stomach discomfort unusual drowsiness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Heartburn Relief side effects

1567. This particular disease, GERD (Heartburn), interacts with the drug Heartburn Relief Maximum

- No side effects listed

1568. This particular disease, GERD (Heartburn), interacts with the drug metoclopramide and has th

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop taking metoclopramide and call your doctor at once if you have any of these SIGNS OF A SERIOUS MOVEMENT DISORDER, which may occur within the first 2 days of treatment: tremors or shaking in your arms or legs
- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- or any new or unusual muscle movements you cannot control. Call your doctor at once if you have: confusion, depression, thoughts of suicide or hurting yourself
- slow or jerky muscle movements, problems with balance or walking
- mask-like appearance in your face
- a seizure
- anxiety , agitation , jittery feeling, trouble staying still, trouble sleeping

Drug Side Effects Summary

- swelling, feeling short of breath, rapid weight gain
- or severe nervous system reaction very stiff (rigid) muscles, high fever , sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Common metoclopramide side effects may include: feeling restless
- feeling drowsy or tired
- lack of energy
- nausea, vomiting
- headache, confusion
- or sleep problems (insomnia).

1569. This particular disease, GERD (Heartburn), interacts with the drug omeprazole / sodium bicar

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Omeprazole and sodium bicarbonate may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- new or unusual pain in your wrist, back, hip, or thigh
- muscle twitching or tremor
- numbness or tingling in your face, arms, or legs
- confusion, dizziness
- a seizure
- low magnesium--dizziness, irregular heartbeats, feeling jittery, muscle cramps, muscle spasms, cough or choking feeling
- kidney problems--fever, rash, nausea , loss of appetite, joint pain, urinating less than usual, blood in your urine, weight gain

Drug Side Effects Summary

- new or worsening symptoms of lupus--joint pain, and a skin rash on your cheeks or arms that worsens in sunlight
- or vitamin B12 deficiency --shortness of breath, feeling lightheaded, irregular heartbeats, muscle weakness, pale skin, tiredness, mood changes. Taking this medicine long-term may cause you to develop stomach growths called fundic gland polyps. Talk with your doctor about this risk. If you use omeprazole and sodium bicarbonate for longer than 3 years, you could develop a vitamin B-12 deficiency. Talk to your doctor about how to manage this condition if you develop it. Common side effects of omeprazole and sodium bicarbonate may include: headache
- nausea, vomiting, stomach pain, gas
- or diarrhea.

1570. This particular disease, GERD (Heartburn), interacts with the drug Pepcid AC Chewable Tablets

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Dizziness or passing out. A fast heartbeat . A heartbeat that does not feel normal. Pepcid AC Chewable Tablets side effects

Drug Side Effects Summary

1571. This particular disease, GERD (Heartburn), interacts with the drug Pepcid Oral Suspension and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Dizziness or passing out. A fast heartbeat . A heartbeat that does not feel normal. Pepcid Oral Suspension side effects

1572. This particular disease, GERD (Heartburn), interacts with the drug Mylanta and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat

Drug Side Effects Summary

- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Mylanta side effects

1573. This particular disease, GERD (Heartburn), interacts with the drug Reglan and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop taking Reglan and call your doctor at once if you have any of these SIGNS OF A SERIOUS MOVEMENT DISORDER, which may occur within the first 2 days of treatment: tremors or shaking in your arms or legs
- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- or any new or unusual muscle movements you cannot control. Call your doctor at once if you have: confusion, depression, thoughts of suicide or hurting yourself
- slow or jerky muscle movements, problems with balance or walking
- mask-like appearance in your face
- a seizure
- anxiety , agitation , jittery feeling, trouble staying still, trouble sleeping
- swelling, feeling short of breath, rapid weight gain
- or severe nervous system reaction very stiff (rigid) muscles, high fever , sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Common Reglan side effects may include: feeling restless
- feeling drowsy or tired
- lack of energy
- nausea, vomiting

Drug Side Effects Summary

- headache, confusion
- or sleep problems (insomnia).

1574. This particular disease, GERD (Heartburn), interacts with the drug Zegerid and has the follow

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- new or unusual pain in your wrist, back, hip, or thigh
- muscle twitching or tremor
- numbness or tingling in your face, arms, or legs
- confusion, dizziness
- a seizure
- low magnesium - dizziness, irregular heartbeats, feeling jittery, muscle cramps, muscle spasms, cough or choking feeling
- kidney problems - fever, rash, nausea, loss of appetite, joint pain, urinating less than usual, blood in your urine, weight gain
- new or worsening symptoms of lupus - joint pain, and a skin rash on your cheeks or arms that worsens in sunlight
- or vitamin B12 deficiency - shortness of breath, feeling lightheaded, irregular heartbeats, muscle weakness, pale skin, tiredness, mood changes. Taking this medicine long-term may cause you to develop stomach growths called fundic gland polyps. Talk with your doctor about this risk. If you use Zegerid for longer than 3 years, you could develop a vitamin B-12 deficiency. Talk to your doctor about how to manage this condition if you develop it. Common Zegerid side effects may include: headache

Drug Side Effects Summary

- nausea , vomiting, stomach pain, gas
- or diarrhea.

1575. This particular disease, GERD (Heartburn), interacts with the drug calcium carbonate and has

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Calcium carbonate may cause serious side effects.

Call your doctor at once if you have: little or no urinating

- swelling, rapid weight gain
- or high levels of calcium in your blood-- nausea , vomiting , constipation , increased thirst or urination, muscle weakness, bone pain, confusion, lack of energy, or feeling tired. Common side effects of calcium carbonate may include: upset stomach , gas
- or constipation.

1576. This particular disease, GERD (Heartburn), interacts with the drug Nexium IV and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: Less common Bloody, black, or tarry stools diarrhea fever vomiting of blood or material that looks like coffee grounds Rare Drowsiness loss of appetite mood or mental changes muscle spasms (tetany) or twitching nausea seizures trembling unusual tiredness or weakness vomiting Incidence not known Agitation back pain blistering, peeling, or loosening of the skin bloating bloody or cloudy urine blurred vision bone fracture chest pain or tightness confusion constipation cough darkened urine decreased urine difficulty breathing difficulty with swallowing dizziness dry mouth fast heartbeat fever with or without chills general feeling of discomfort or illness greatly decreased frequency of urination or amount of

Drug Side Effects Summary

urine headache high fever hoarseness indigestion irritability joint or muscle pain light-colored stools loss of consciousness lower back or side pain muscle aches, cramps, stiffness, or weakness nausea noisy breathing numbness and tingling around the mouth, fingertips, or feet painful or difficult urination pains in the stomach, side, or abdomen, possibly radiating to the back pale skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red skin lesions, often with a purple center red, irritated eyes skin rash, hives, or itching sore throat sores, ulcers, or white spots in the mouth or on the lips stiff neck stomach cramps stomach pain, continuing swelling of the feet or lower legs swelling or inflammation of the mouth swollen glands tenderness unusual bleeding or bruising upper right stomach pain watery or bloody diarrhea yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Belching heartburn stomach discomfort or upset Less common Burning, itching, redness, skin rash, swelling, or soreness at the injection site difficulty having a bowel movement dryness of the mouth feeling of constant movement of self or surroundings gas pain or tenderness around the eyes and cheekbones sensation of spinning sneezing stuffy or runny nose Incidence not known Aggression bad unusual or unpleasant (after) taste change in taste difficulty in moving discouragement feeling sad or empty increased sensitivity of the skin to sunlight joint swelling loss of interest or pleasure redness or discoloration of the skin seeing, hearing, or feeling things that are not there severe sunburn swelling of the breasts or breast soreness in both females and males thinning or loss of hair trouble concentrating trouble sleeping Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Nexium IV side effects

Drug Side Effects Summary

1577. This particular disease, GERD (Heartburn), interacts with the drug nizatidine and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Nizatidine may cause serious side effects. Stop using nizatidine and call your doctor at once if you have: worsening heartburn
- chest pain
- pale skin, feeling light-headed or short of breath
- or jaundice (yellowing of the skin or eyes). Common side effects of nizatidine may include: headache , dizziness
- diarrhea
- or runny or stuffy nose.

1578. This particular disease, GERD (Heartburn), interacts with the drug aluminum hydroxide / magnesium hydroxide and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: bone pain, muscle weakness
- confusion, changes in your mental state, seizure (convulsions)
- or pale skin, feeling light-headed or short of breath, rapid heart rate. Less serious side effects may be more likely, and you may have none at all.

1579. This particular disease, GERD (Heartburn), interacts with the drug bethanechol and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Bethanechol may cause serious side effects. Call your doctor at once if you have: difficulty breathing

Drug Side Effects Summary

doctor at once if you have: slow heartbeats

- a light-headed feeling, like you might pass out
- or wheezing, chest tightness, trouble breathing. Common side effects of bethanechol may include:
an urgent need to urinate
- stomach cramps, nausea, diarrhea
- burping , increased saliva
- headache , dizziness
- sweating, flushing (sudden warmth, redness, or tingly feeling)
- a general ill feeling
- or watery eyes , vision changes.

1580. This particular disease, GERD (Heartburn), interacts with the drug Calci-Chew and has the fol

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad constipation . Calci-Chew side effects

Drug Side Effects Summary

1581. This particular disease, GERD (Heartburn), interacts with the drug Pepcid Complete and has the

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Pepcid Complete and call your doctor at once if: your heartburn gets worse
- or you still have heartburn after 2 weeks of taking Pepcid Complete. Less serious side effects may be more likely, and you may have none at all.

1582. This particular disease, GERD (Heartburn), interacts with the drug Prevacid OTC and has the

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Prevacid OTC may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- new or unusual pain in your wrist, back, hip, or thigh
- a seizure (convulsions)
- kidney problems--fever, nausea, little or no urination, blood in your urine, swelling, rapid weight gain
- low magnesium-- dizziness , fast or irregular heart rate, tremors (shaking) or jerking muscle movements, feeling jittery, muscle cramps, muscle spasms in your hands and feet, cough or choking feeling
- or new or worsening symptoms of lupus--joint pain, and a skin rash on your cheeks or arms that worsens in sunlight. Taking Prevacid OTC long-term may cause you to develop stomach growths called fundic gland polyps. Talk with your doctor about this risk. If you use Prevacid OTC for longer than 3 years, you could develop a vitamin B-12 deficiency. Talk to your doctor about how to manage this condition if you develop it. Common side effects of Prevacid OTC may include: nausea,

Drug Side Effects Summary

stomach pain

- diarrhea, constipation
- or headache .

1583. This particular disease, GERD (Heartburn), interacts with the drug Prevacid SoluTab and has

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Prevacid SoluTab may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- new or unusual pain in your wrist, back, hip, or thigh
- a seizure (convulsions)
- kidney problems--fever, nausea, little or no urination, blood in your urine, swelling, rapid weight gain
- low magnesium-- dizziness , fast or irregular heart rate, tremors (shaking) or jerking muscle movements, feeling jittery, muscle cramps, muscle spasms in your hands and feet, cough or choking feeling
- or new or worsening symptoms of lupus--joint pain, and a skin rash on your cheeks or arms that worsens in sunlight. Taking Prevacid SoluTab long-term may cause you to develop stomach growths called fundic gland polyps. Talk with your doctor about this risk. If you use Prevacid SoluTab for longer than 3 years, you could develop a vitamin B-12 deficiency. Talk to your doctor about how to manage this condition if you develop it. Common side effects of Prevacid SoluTab may include: nausea, stomach pain
- diarrhea, constipation
- or headache .

Drug Side Effects Summary

1584. This particular disease, GERD (Heartburn), interacts with the drug Tagamet HB and has the following side effects:

- pain when swallowing
- bloody or tarry stools, cough with bloody mucus or vomit that looks like coffee grounds
- changes in mood, anxiety , agitation
- confusion, hallucinations
- or breast swelling or tenderness. Serious side effects may be more likely in older adults and those who are ill or debilitated. Common side effects of Tagamet HB may include: headache
- or diarrhea .

1585. This particular disease, GERD (Heartburn), interacts with the drug Titalac and has the following side effects:

- Stop taking calcium carbonate and seek emergency medical attention if you experience a rare allergic reaction (difficulty breathing
- closing of the throat
- swelling of the lips, tongue, or face
- or hives). Other, less serious side effects may be more likely to occur. Continue to take calcium carbonate and notify your doctor if you experience nausea or vomiting
- decreased appetite
- constipation
- dry mouth or increased thirst
- or increased urination. Side effects other than those listed here may also occur. Talk to your doctor about any side effect that seems unusual or that is especially bothersome. Titalac side effects

1586. This particular disease, GERD (Heartburn), interacts with the drug Tums Regular Strength and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if

Drug Side Effects Summary

you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad constipation . Tums Regular Strength side effects

1587. This particular disease, GERD (Heartburn), interacts with the drug Tums Smoothies and has t

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad constipation . Tums Smoothies side

Drug Side Effects Summary

effects

1588. This particular disease, GERD (Heartburn), interacts with the drug Zegerid OTC and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- new or unusual pain in your wrist, back, hip, or thigh
- muscle twitching or tremor
- numbness or tingling in your face, arms, or legs
- confusion, dizziness
- a seizure
- low magnesium - dizziness, irregular heartbeats, feeling jittery, muscle cramps, muscle spasms, cough or choking feeling
- kidney problems - little or no urination, blood in your urine, swelling, rapid weight gain
- new or worsening symptoms of lupus - joint pain, and a skin rash on your cheeks or arms that worsens in sunlight
- or vitamin B12 deficiency - shortness of breath, feeling lightheaded, irregular heartbeats, muscle weakness, pale skin, tiredness, mood changes. Common Zegerid OTC side effects may include: headache
- nausea , vomiting, stomach pain, gas
- or diarrhea.

1589. This particular disease, GERD (Heartburn), interacts with the drug Aciphex Sprinkle and has the following side effects:

- hives

Drug Side Effects Summary

- difficulty breathing
- swelling of your face, lips, tongue, or throat. AcipHex Sprinkle may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- sudden pain or trouble moving your hip, wrist, or back
- seizure (convulsions)
- kidney problems--fever, rash, nausea , loss of appetite, joint pain, urinating less than usual, blood in your urine, swelling, rapid weight gain
- new or worsening symptoms of lupus--joint pain, and a skin rash on your cheeks or arms that worsens in sunlight
- low magnesium-- dizziness , fast or irregular heart rate, tremors (shaking) or jerking muscle movements, feeling jittery, muscle cramps, muscle spasms in your hands and feet, cough or choking feeling
- or signs of bleeding (if you also take warfarin)--headaches, dizziness, weakness
- pain or swelling
- bruising, unusual bleeding (nosebleeds , bleeding gums)
- red or pink urine
- heavy menstrual flow
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- or any bleeding that will not stop. Taking AcipHex Sprinkle long-term may cause you to develop stomach growths called fundic gland polyps. Talk with your doctor about this risk. If you use AcipHex Sprinkle for longer than 3 years, you could develop a vitamin B-12 deficiency. Talk to your doctor about how to manage this condition if you develop it. Common side effects of AcipHex Sprinkle may include: infection
- pain
- sore throat

Drug Side Effects Summary

- headache
- nausea, vomiting
- gas, diarrhea, constipation
- or stomach pain.

1590. This particular disease, GERD (Heartburn), interacts with the drug Alamag and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Alamag side effects

1591. This particular disease, GERD (Heartburn), interacts with the drug Alka-Seltzer Heartburn + G

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

Drug Side Effects Summary

- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Alka-Seltzer Heartburn + Gas side effects

1592. This particular disease, GERD (Heartburn), interacts with the drug Almacone and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: bone pain, muscle weakness
- confusion, changes in your mental state, seizure (convulsions)
- or pale skin, feeling light-headed or short of breath, rapid heart rate. Less serious side effects may be more likely, and you may have none at all.

1593. This particular disease, GERD (Heartburn), interacts with the drug aluminum hydroxide / magnesium hydroxide and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common side effects may include diarrhea .

1594. This particular disease, GERD (Heartburn), interacts with the drug aluminum hydroxide / magnesium hydroxide and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if

Drug Side Effects Summary

you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Aluminum hydroxide / magnesium hydroxide side effects

1595. This particular disease, GERD (Heartburn), interacts with the drug aluminum hydroxide and h

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: severe stomach pain or constipation, loss of appetite
- pain when you urinate
- muscle weakness, tiredness
- extreme drowsiness
- or bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Less serious side effects may be more likely, and you may have none at all.

1596. This particular disease, GERD (Heartburn), interacts with the drug Amphojel and has the follo

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: severe stomach pain or constipation, loss of appetite
- pain when you urinate
- muscle weakness, tiredness
- extreme drowsiness
- or bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Less serious side effects may be more likely, and you may have none at all.

1597. This particular disease, GERD (Heartburn), interacts with the drug Axid AR and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using Axid AR and call your doctor at once if you have: worsening heartburn
- chest pain
- pale skin, feeling light-headed or short of breath
- or jaundice (yellowing of the skin or eyes). Common Axid AR side effects may include: headache, dizziness
- diarrhea
- or runny or stuffy nose.

1598. This particular disease, GERD (Heartburn), interacts with the drug belladonna and has the following

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Belladonna may cause serious side effects. Call your doctor at once if you have: fever
- fast heartbeats
- severe diarrhea or severe constipation
- a seizure
- unusual changes in mood or behavior
- hallucinations
- red and dry skin
- dilated pupils
- or painful or difficult urination. Common side effects of belladonna may include: dry mouth
- blurred vision
- muscle spasms
- constipation
- decreased urination
- or decreased sweating.

1599. This particular disease, GERD (Heartburn), interacts with the drug calcium carbonate / famotidine

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using calcium carbonate, famotidine, and magnesium hydroxide and call your doctor at once if: your heartburn gets worse
- or you still have heartburn after 2 weeks of taking calcium carbonate, famotidine, and magnesium hydroxide. Less serious side effects may be more likely, and you may have none at all.

Drug Side Effects Summary

1600. This particular disease, GERD (Heartburn), interacts with the drug calcium carbonate / magne

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat.

1601. This particular disease, GERD (Heartburn), interacts with the drug calcium carbonate / simeth

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Calcium carbonate / simethicone side effects

1602. This particular disease, GERD (Heartburn), interacts with the drug Di-Gel and has the followin

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

Drug Side Effects Summary

- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Di-Gel side effects

1603. This particular disease, GERD (Heartburn), interacts with the drug Dual Action Complete and

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Dual Action Complete and call your doctor at once if: your heartburn gets worse
- or you still have heartburn after 2 weeks of taking Dual Action Complete. Less serious side effects may be more likely, and you may have none at all.

1604. This particular disease, GERD (Heartburn), interacts with the drug Duo Fusion and has the fo

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Duo Fusion and call your doctor at once if: your heartburn gets worse
- or you still have heartburn after 2 weeks of taking Duo Fusion. Less serious side effects may be more likely, and you may have none at all.

1605. This particular disease, GERD (Heartburn), interacts with the drug Gas-X with Maalox Extra S

Drug Side Effects Summary

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Gas-X with Maalox Extra Strength side effects

1606. This particular disease, GERD (Heartburn), interacts with the drug Gelusil and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: bone pain, muscle weakness
- confusion, changes in your mental state, seizure (convulsions)
- or pale skin, feeling light-headed or short of breath, rapid heart rate. Less serious side effects may be more likely, and you may have none at all.

1607. This particular disease, GERD (Heartburn), interacts with the drug Mi-Acid and has the following

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: bone pain, muscle weakness
- confusion, changes in your mental state, seizure (convulsions)
- or pale skin, feeling light-headed or short of breath, rapid heart rate. Less serious side effects may be more likely, and you may have none at all.

1608. This particular disease, GERD (Heartburn), interacts with the drug Milantex and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: bone pain, muscle weakness
- confusion, changes in your mental state, seizure (convulsions)
- or pale skin, feeling light-headed or short of breath, rapid heart rate. Less serious side effects may be more likely, and you may have none at all.

1609. This particular disease, GERD (Heartburn), interacts with the drug Mintox and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: bone pain, muscle weakness
- confusion, changes in your mental state, seizure (convulsions)
- or pale skin, feeling light-headed or short of breath, rapid heart rate. Less serious side effects may be more likely, and you may have none at all.

Drug Side Effects Summary

1610. This particular disease, GERD (Heartburn), interacts with the drug Mintox Plus and has the fo

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Mintox Plus side effects

1611. This particular disease, GERD (Heartburn), interacts with the drug Mylanta Coat & Cool and h

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat.

1612. This particular disease, GERD (Heartburn), interacts with the drug Mylanta Maximum Strength

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: bone pain, muscle weakness
- confusion, changes in your mental state, seizure (convulsions)

Drug Side Effects Summary

- or pale skin, feeling light-headed or short of breath, rapid heart rate. Less serious side effects may be more likely, and you may have none at all.

1613. This particular disease, GERD (Heartburn), interacts with the drug Mylanta Tonight and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat.

1614. This particular disease, GERD (Heartburn), interacts with the drug Rulox and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: bone pain, muscle weakness
- confusion, changes in your mental state, seizure (convulsions)
- or pale skin, feeling light-headed or short of breath, rapid heart rate. Less serious side effects may be more likely, and you may have none at all.

1615. This particular disease, GERD (Heartburn), interacts with the drug sodium citrate and has the following side effects:

- No side effects listed

1616. This particular disease, GERD (Heartburn), interacts with the drug Tricitrasol and has the following side effects:

- No side effects listed

1617. This particular disease, GERD (Heartburn), interacts with the drug Tums Chewy Bites with Gaviscon and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and

Drug Side Effects Summary

sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Tums Chewy Bites with Gas Relief (Lemon & Strawberry) side effects

1618. This particular disease, GERD (Heartburn), interacts with the drug Zegerid with Magnesium H

- No side effects listed

1619. This particular disease, Gout, interacts with the drug allopurinol and has the following side ef

- Stop using this medicine and get emergency medical help if you have signs of an allergic reaction to allopurinol (hives , difficult breathing, swelling in your face or throat) or a severe skin reaction (fever, sore throat , burning eyes, skin pain, red or purple skin rash with blistering and peeling). Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Stop using allopurinol and call your doctor at once if you have: any skin rash, no matter how mild
- painful urination, blood in the urine

Drug Side Effects Summary

- little or no urination
- easy bruising, unusual bleeding
- numbness , tingling, burning pain
- worsening gout symptoms
- or liver problems - loss of appetite, weight loss , stomach pain (upper right side), itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common allopurinol side effects may include: an increase in gout attacks when you first starting taking allopurinol oral
- rash
- drowsiness
- fever, chills
- abnormal liver function tests
- nausea , diarrhea
- or joint pain.

1620. This particular disease, Gout, interacts with the drug Uloric and has the following side effects

- skin rash , fever, swollen glands, flu-like symptoms, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. This reaction may occur several weeks after you began using Uloric. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- heart attack symptoms - chest pain or pressure, pain spreading to your jaw or shoulder, fast heartbeats, shortness of breath
- liver problems - stomach pain (upper right side), unusual tiredness, dark urine, jaundice (yellowing of the skin or eyes)
- or signs of a stroke - sudden numbness or weakness (especially on one side of the body), sudden severe headache , slurred speech, problems with vision or balance. Common Uloric side effects may include: gout flares, joint pain

Drug Side Effects Summary

- nausea
- mild rash
- or abnormal liver function tests.

1621. This particular disease, Gout, interacts with the drug Zyloprim and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Ankle, knee, or great toe joint pain joint stiffness or swelling rash rash with flat lesions or small raised lesions on the skin Rare Agitation ammonia-like breath odor anxiety bleeding gums blistering, peeling, or loosening of the skin blood in the urine or stools bloody nose bloody or black, tarry stools blue or pale skin bruising changes in skin color chest pain or discomfort chest pain, possibly moving to the left arm, neck, or shoulder chills clay-colored stools cloudy urine confusion constipation cough coughing up blood cracks in the skin dark urine decreased awareness or responsiveness decreased urine output depression diarrhea difficulty with breathing dizziness drowsiness dry mouth feeling of warmth or heat fever flushing or redness of the skin, especially on the face and neck general feeling of discomfort or illness headache hives or welts, itching, skin rash hoarseness hostility incoherent speech increased urination irritability joint or muscle pain large, flat, blue or purplish patches in the skin light-colored stools loss of appetite loss of consciousness loss of heat from the body lower back or side pain metallic taste muscle twitching muscle weakness nausea noisy breathing pain, tenderness, or swelling of the foot or leg painful or difficult urination pinpoint red or purple spots on the skin rapid weight gain red, irritated eyes red, swollen skin redness, soreness, or itching skin right upper abdominal or stomach pain and fullness scaly skin seizures severe sleepiness severe stomach pain slow or irregular heartbeat sore throat sores, ulcers, or white spots on the lips or in the mouth sores, welting, or blisters stomach pain sweating swelling of the face, ankles, hands, or lower

Drug Side Effects Summary

legs swollen or painful glands swollen, painful, or tender lymph glands in the neck, armpit, or groin thirst tightness in the chest unpleasant breath odor unusual bleeding or bruising unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness unusual weight gain or loss vomiting vomiting of blood or material that looks like coffee grounds yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Rare Bad, unusual, or unpleasant (after) taste blindness blue-yellow color blindness blurred vision body aches or pain burning feeling in the chest or stomach burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings burning, dry, or itching eyes change in taste change in vision congestion continuing ringing or buzzing or other unexplained noise in the ears decreased interest in sexual intercourse decreased vision difficulty with moving discharge or excessive tearing feeling of constant movement of self or surroundings hair loss or thinning of the hair hearing loss impaired vision inability to have or keep an erection indigestion lack or loss of strength loss in sexual ability, desire, drive, or performance loss of appetite loss of memory multiple swollen and inflamed skin lesions muscle pain, cramps, or stiffness muscular pain, tenderness, wasting, or weakness problems with memory redness, pain, or swelling of the eye, eyelid, or inner lining of the eyelid runny nose sensation of spinning sensitivity to light sleepiness or unusual drowsiness sleeplessness sneezing stomach upset stuffy nose sweating swelling of the breasts or breast soreness in both females and males swelling of the salivary glands swelling or inflammation of the mouth tearing tender, swollen glands in the neck tenderness in the stomach area throbbing pain trouble getting pregnant trouble with sleeping trouble with swallowing unsteadiness or awkwardness voice changes weakness in the arms, hands, legs, or feet weight loss Other side effects not listed may also occur in some patients. If you notice any other effects, check

Drug Side Effects Summary

with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Zylprim side effects

1622. This particular disease, Gout, interacts with the drug Alopriam and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: Less common Agitation coma confusion decreased urine output depression dizziness headache hostility irritability lethargy muscle twitching nausea rapid weight gain seizures stupor swelling of the face, ankles, or hands unusual tiredness or weakness Rare Black, tarry stools bladder pain bleeding gums blistering, peeling, loosening of the skin blood in the urine or stools blurred vision changes in skin color chest pain or tightness chills clay-colored stools cloudy urine cough or hoarseness coughing up blood dark urine diarrhea difficult, burning, or painful urination difficulty with breathing or swallowing difficulty with speaking dilated neck veins dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position double vision enlarged stomach fainting fast or slow heartbeat feeling of warmth fever frequent urge to urinate inability to move the arms, legs, or facial muscles inability to speak increased menstrual flow or vaginal bleeding increased sweating irregular breathing irregular heartbeat itching, rash joint pain, stiffness, or swelling lightheadedness loosening of the fingernails loss of appetite loss of strength or energy lower back or side pain muscle pain or weakness nosebleeds numbness, tingling, pain, or weakness in the hands or feet pain, tenderness, or swelling of the foot or leg painful or difficult urination paralysis pinpoint red spots on the skin prolonged bleeding from cuts rapid, shallow breathing red or black, tarry stools red or dark brown urine red skin lesions, often with a purple center redness of the face, neck, arms, and occasionally, upper chest slow speech sore throat sores, ulcers, or white spots on the lips or in the mouth stomach pain sweating swelling of the fingers, feet, or lower legs swollen glands trouble breathing unpleasant

Drug Side Effects Summary

breath odor unusual bleeding or bruising unusual weak feeling vomiting of blood weight gain yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Drowsiness indigestion unusual hair loss Rare Bleeding, blistering, burning, coldness, discoloration of the skin, feeling of pressure, hives, infection, inflammation, itching, lumps, numbness, pain, rash, redness, scarring, soreness, stinging, swelling, tenderness, tingling, ulceration, or warmth at the injection site body aches or pain congestion cracked lips difficulty with moving hair loss or thinning of the hair hives or welts muscle stiffness pain, redness, swelling, tenderness, or warmth on the skin runny nose voice changes Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Aloprim side effects

1623. This particular disease, Gout, interacts with the drug febuxostat and has the following side effects

- skin rash, fever, swollen glands, flu-like symptoms, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Febuxostat may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, fast heartbeats, shortness of breath
- liver problems--stomach pain (upper right side), unusual tiredness, dark urine, jaundice (yellowing of the skin or eyes)
- or signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance. Common side effects of

Drug Side Effects Summary

febuxostat may include: gout flares, joint pain

- nausea
- mild rash
- or liver problems.

1624. This particular disease, Gout, interacts with the drug Krystexxa and has the following side effects

- hives
- wheezing, difficult breathing
- swelling of your face, lips, tongue, or throat. Some side effects may occur during the injection. Tell your caregiver if you feel nervous, light-headed, itchy, short of breath, or have fast heartbeats, chest discomfort, or redness of your skin during the injection. Call your doctor at once if you have: chest pain
- or flushing (warmth, redness, or tingly feeling). Common Krystexxa side effects may include: COVID-19 symptoms such as fever or chills, cough, sore throat, shortness of breath, fatigue, muscle or body aches, headache, loss of taste or smell
- hives, shortness of breath, chest pain or discomfort, skin redness, or itching
- joint pain
- allergic reactions
- new gout flares
- nausea, vomiting, constipation
- or bruising.

1625. This particular disease, Gout, interacts with the drug probenecid and has the following side effects

- hives
- difficulty breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Probenecid may cause serious side effects. Call your doctor at once if you have: worsening gout symptoms
- nausea, vomiting, stomach pain or swelling
- painful or difficult urination
- severe pain in your side or lower back
- blood in your urine
- urine that looks cloudy or foamy
- puffy eyes, swelling in your ankles or feet, weight gain
- or pale or yellowed skin. Common side effects of probenecid may include: headache , dizziness
- nausea, vomiting, loss of appetite
- flushing (sudden warmth, redness, or tingly feeling)
- sore gums
- urinating more than usual
- itching, rash
- or hair loss .

1626. This particular disease, Gout, interacts with the drug pegloticase and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some side effects may occur during the injection. Tell your caregiver if you feel nervous, light-headed, itchy, short of breath, or have fast heartbeats, chest discomfort, or redness of your skin during the injection. Pegloticase may cause serious side effects. Call your doctor at once if you have: chest pain
- or flushing (warmth, redness, or tingly feeling). Common side effects of pegloticase may include: COVID-19 symptoms such as fever or chills, cough, sore throat , shortness of breath, fatigue,

Drug Side Effects Summary

muscle or body aches, headache , loss of taste or smell

- hives, shortness of breath, chest pain or discomfort, skin redness, or itching
- joint pain
- allergic reactions
- new gout flares
- nausea , vomiting , constipation
- or bruising.

1627. This particular disease, Gout, interacts with the drug rilonacept and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. You may get infections more easily, even serious or fatal infections. Call your doctor right away if you have signs of infection such as: fever, chills, sore throat
- open sores on your body
- cough with mucus, feeling short of breath
- headache , neck stiffness, increased sensitivity to light, confusion
- or chest discomfort, wheezing, dry cough, rapid weight loss . Also call your doctor at once if you have: bloody, black, or tarry stools
- coughing up blood or vomit that looks like coffee grounds
- wheezing, chest tightness, trouble breathing
- pain or burning when you urinate
- or signs of an ear infection--fever, ear pain or full feeling, trouble hearing, drainage from the ear, fussiness in a child. Common side effects of rilonacept may include: cold symptoms such as stuffy nose, runny nose , sneezing, sore throat

Drug Side Effects Summary

- ear infection
- muscle or joint pain
- rash
- or pain, swelling, redness, itching, warmth, blistering, bleeding, bruising, a rash, or a lump where the medicine was injected.

1628. This particular disease, Hair Loss, interacts with the drug minoxidil and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Although the risk of serious side effects is low when minoxidil topical is applied to the skin, side effects can occur if the medicine is absorbed into your bloodstream. Minoxidil topical may cause serious side effects. Call your doctor at once if you have: severe scalp irritation
- unwanted growth of facial hair
- chest pain, fast heartbeats
- swelling in your hands or feet, rapid weight gain
- a light-headed feeling, like you might pass out
- headache , dizziness , confusion
- or flushing (warmth, redness, or tingly feeling). Common side effects of minoxidil topical may include: changes in the color or texture of your hair.

1629. This particular disease, Hair Loss, interacts with the drug Azulfidine and has the following side effects:

- skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. You may get infections more easily, even serious or fatal infections. Call your doctor right away if you have signs of infection such as: fever, chills, sore throat

Drug Side Effects Summary

- mouth sores, red or swollen gums
- pale skin, easy bruising, unusual bleeding
- or chest discomfort, wheezing, dry cough or hack, rapid weight loss . Also call your doctor at once if you have: fever with headache , rash, and vomiting
- a skin rash, no matter how mild
- severe nausea or vomiting when you first start taking Azulfidine
- little or no urination, urine that looks foamy
- puffy eyes, swelling in your ankles or feet, weight gain
- or liver problems--loss of appetite, stomach pain (upper right side), dark urine, jaundice (yellowing of the skin or eyes). Common side effects of Azulfidine may include: nausea, vomiting, upset stomach , loss of appetite
- headache
- rash
- or low sperm count in men.

1630. This particular disease, Hair Loss, interacts with the drug sulfasalazine and has the following

- See "Important information" above. The most common side effects of sulfasalazine are: anorexia headache nausea vomiting gastric distress reversible oligospermia. Less frequent side effects are: skin rash pruritus urticaria fever Heinz body anemia hemolytic anemia cyanosis Sulfasalazine may also produce an orange-yellow discoloration of the urine or skin. These are not all the possible side effects of sulfasalazine. For more information, ask your doctor or pharmacist. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Sulfasalazine side effects

1631. This particular disease, Hair Loss, interacts with the drug Rogaine and has the following side

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Although the risk of serious side effects is low when Rogaine is applied to the skin, side effects can occur if the medicine is absorbed into your bloodstream. Call your doctor at once if you have: severe scalp irritation
- unwanted growth of facial hair
- chest pain, fast heartbeats
- swelling in your hands or feet, rapid weight gain
- a light-headed feeling, like you might pass out
- headache, dizziness, confusion
- or flushing (warmth, redness, or tingly feeling). Common Rogaine side effects may include: changes in the color or texture of your hair.

1632. This particular disease, Hair Loss, interacts with the drug Kenalog-10 and has the following s

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: More common Aggression agitation anxiety blurred vision decrease in the amount of urine dizziness fast, slow, pounding, or irregular heartbeat or pulse headache irritability mental depression mood changes nervousness noisy, rattling breathing numbness or tingling in the arms or legs pounding in the ears shortness of breath swelling of the fingers, hands, feet, or lower legs trouble thinking, speaking, or walking troubled breathing at rest weight gain Incidence not known Abdominal cramping and/or burning (severe) abdominal pain backache bloody, black, or tarry stools cough or hoarseness darkening of skin decreased vision diarrhea dry mouth eye pain eye tearing facial hair growth in females fainting fatigue fever or chills flushed, dry skin fractures fruit-like breath odor full or round face, neck, or trunk

Drug Side Effects Summary

heartburn and/or indigestion (severe and continuous) increased hunger increased thirst increased urination loss of appetite loss of sexual desire or ability lower back or side pain menstrual irregularities muscle pain or tenderness muscle wasting or weakness nausea pain in back, ribs, arms, or legs painful or difficult urination skin rash sleeplessness sweating trouble healing trouble sleeping unexplained weight loss unusual tiredness or weakness vision changes vomiting vomiting of material that looks like coffee grounds Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Blemishes on the skin bruising dry, scaly skin feeling of constant movement of self or surroundings full or bloated feeling increased appetite increased hair growth on the face, forehead, back, arms, and legs large, flat, blue, or purplish patches in the skin pimples pitting or depression of the skin at the injection site reddish purple lines on the arms, face, legs, trunk, or groin redness of the skin redness, swelling, tenderness, or pain at the injection site sensation of spinning small, red, or purple spots on the skin thin, fragile, or shiny skin thinning of the scalp hair Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Kenalog-10 side effects

1633. This particular disease, Hair Loss, interacts with the drug Aristospan and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: More common Aggression agitation anxiety blurred vision decrease in the amount of urine dizziness fast, slow, pounding, or

Drug Side Effects Summary

irregular heartbeat or pulse headache irritability mental depression mood changes nervousness noisy, rattling breathing numbness or tingling in the arms or legs pounding in the ears shortness of breath swelling of the fingers, hands, feet, or lower legs trouble thinking, speaking, or walking troubled breathing at rest weight gain Incidence not known Abdominal cramping and/or burning (severe) abdominal pain backache bloody, black, or tarry stools cough or hoarseness darkening of skin decreased vision diarrhea dry mouth eye pain eye tearing facial hair growth in females fainting fatigue fever or chills flushed, dry skin fractures fruit-like breath odor full or round face, neck, or trunk heartburn and/or indigestion (severe and continuous) increased hunger increased thirst increased urination loss of appetite loss of sexual desire or ability lower back or side pain menstrual irregularities muscle pain or tenderness muscle wasting or weakness nausea pain in back, ribs, arms, or legs painful or difficult urination skin rash sleeplessness sweating trouble healing trouble sleeping unexplained weight loss unusual tiredness or weakness vision changes vomiting vomiting of material that looks like coffee grounds Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Blemishes on the skin bruising dry, scaly skin feeling of constant movement of self or surroundings full or bloated feeling increased appetite increased hair growth on the face, forehead, back, arms, and legs large, flat, blue, or purplish patches in the skin pimples pitting or depression of the skin at the injection site reddish purple lines on the arms, face, legs, trunk, or groin redness of the skin redness, swelling, tenderness, or pain at the injection site sensation of spinning small, red, or purple spots on the skin thin, fragile, or shiny skin thinning of the scalp hair Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to

Drug Side Effects Summary

the FDA at 1-800-FDA-1088. Aristospan side effects

1634. This particular disease, Hair Loss, interacts with the drug Rogaine Men's Extra Strength and

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: Less common Itching or skin rash (continued) Rare Acne at site of application burning of scalp facial hair growth increased hair loss inflammation or soreness at root of hair reddened skin swelling of face Signs and symptoms of too much medicine being absorbed into the body--Rare Blurred vision or other changes in vision chest pain dizziness fainting fast or irregular heartbeat flushing headache lightheadedness numbness or tingling of hands, feet, or face swelling of face, hands, feet, or lower legs weight gain (rapid) Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Rogaine Men's Extra Strength side effects

1635. This particular disease, Hair Loss, interacts with the drug Rogaine Women's and has the follo

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: Less common Itching or skin rash (continued) Rare Acne at site of application burning of scalp facial hair growth increased hair loss inflammation or soreness at root of hair reddened skin swelling of face Signs and symptoms of too much medicine being absorbed into the body--Rare Blurred vision or other changes in vision chest pain dizziness fainting fast or irregular heartbeat flushing headache lightheadedness numbness or tingling of hands, feet, or face swelling of face, hands, feet, or lower legs weight gain (rapid) Other side effects not listed may also occur in some patients. If you notice any other effects,

Drug Side Effects Summary

check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Rogaine Women's side effects

1636. This particular disease, Hair Loss, interacts with the drug Hair Regrowth Treatment for Women

- No side effects listed

1637. This particular disease, Hair Loss, interacts with the drug Hair Regrowth Treatment for Men

- No side effects listed

1638. This particular disease, Hair Loss, interacts with the drug Azulfidine EN-tabs and has the following side effects:

- skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. You may get infections more easily, even serious or fatal infections. Call your doctor right away if you have signs of infection such as: fever, chills, sore throat
- mouth sores, red or swollen gums
- pale skin, easy bruising, unusual bleeding
- or chest discomfort, wheezing, dry cough or hack, rapid weight loss . Also call your doctor at once if you have: fever with headache , rash, and vomiting
- a skin rash, no matter how mild
- severe nausea or vomiting when you first start taking Azulfidine EN-tabs
- little or no urination, urine that looks foamy
- puffy eyes, swelling in your ankles or feet, weight gain
- or liver problems--loss of appetite, stomach pain (upper right side), dark urine, jaundice (yellowing of the skin or eyes). Common side effects of Azulfidine EN-tabs may include: nausea, vomiting, upset stomach , loss of appetite
- headache

Drug Side Effects Summary

- rash
- or low sperm count in men.

1639. This particular disease, Hayfever, interacts with the drug Zyrtec and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop taking this medicine and call your doctor at once if you have: fast, pounding, or uneven heartbeat
- weakness, tremors (uncontrolled shaking), or sleep problems (insomnia)
- severe restless feeling, hyperactivity
- confusion
- problems with vision
- or little or no urination. Common Zyrtec side effects may include: drowsiness, tiredness
- dizziness , feeling light-headed
- feeling hot, sweating
- numbness , tingling, burning pain
- decreased sense of taste
- headache
- upset stomach , nausea, constipation
- or dry mouth, sore throat .

1640. This particular disease, Hayfever, interacts with the drug cetirizine and has the following side effects:

- chronic idiopathic urticaria (hives)
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using cetirizine and call your doctor at once if you

Drug Side Effects Summary

have any of these serious side effects: fast, pounding, or uneven heartbeat

- weakness, tremors (uncontrolled shaking), or sleep problems (insomnia)
- severe restless feeling, hyperactivity
- confusion
- problems with vision
- or urinating less than usual or not at all. Less serious cetirizine side effects may include: dizziness, drowsiness
- tired feeling
- dry mouth
- sore throat, cough
- nausea, constipation
- or headache.

1641. This particular disease, Hayfever, interacts with the drug fluticasone and has the following side effects:

- hives , rash
- feeling light-headed
- difficult breathing
- swelling of your face, lips, tongue, or throat. Fluticasone nasal may cause serious side effects. Call your doctor at once if you have: severe or ongoing nosebleeds
- noisy breathing, runny nose, or crusting around your nostrils
- redness, sores, or white patches in your mouth or throat
- fever, chills, body aches
- blurred vision, eye pain , or seeing halos around lights
- any wound that will not heal
- or signs of a hormonal disorder--worsening tiredness or muscle weakness, feeling light-headed,

Drug Side Effects Summary

nausea , vomiting . Steroid medicine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using fluticasone nasal. Common side effects of fluticasone nasal may include: minor nosebleed, burning or itching in your nose

- sores or white patches inside or around your nose
- cough, trouble breathing
- headache , back pain
- sinus pain, sore throat , fever
- or nausea, vomiting.

1642. This particular disease, Hayfever, interacts with the drug Flonase and has the following side effects:

- hives , rash
- feeling light-headed
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe or ongoing nosebleeds
- noisy breathing, runny nose, or crusting around your nostrils
- redness, sores, or white patches in your mouth or throat
- fever, chills, body aches
- blurred vision, eye pain , or seeing halos around lights
- any wound that will not heal
- or signs of a hormonal disorder--worsening tiredness or muscle weakness, feeling light-headed, nausea , vomiting . Steroid medicine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common Flonase side effects may include: minor nosebleed, burning or itching in your nose
- sores or white patches inside or around your nose

Drug Side Effects Summary

- cough, trouble breathing
- headache , back pain
- sinus pain, sore throat , fever
- or nausea, vomiting.

1643. This particular disease, Hayfever, interacts with the drug Allegra and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop taking Allegra and call your doctor if you have fever, chills, body aches, cough, or other flu symptoms. Less serious Allegra side effects may include: nausea, diarrhea, upset stomach
- menstrual cramps
- drowsiness, tired feeling
- headache
- or muscle or back pain.

1644. This particular disease, Hayfever, interacts with the drug Claritin and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: fast or uneven heart rate
- severe headache
- or a light-headed feeling, like you might pass out
- Common Claritin side effects may include: headache
- feeling tired or drowsy

Drug Side Effects Summary

- stomach pain, vomiting
- dry mouth
- or feeling nervous or hyperactive.

1645. This particular disease, Hayfever, interacts with the drug fexofenadine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Fexofenadine may cause serious side effects. Stop using fexofenadine and call your doctor at once if you have: flu-like symptoms (fever, chills, unusual tiredness)
- new or worsening cough
- pain
- or signs of an ear infection--fever, ear pain or full feeling, trouble hearing, drainage from the ear, fussiness in a child. Common side effects of fexofenadine may include: headache
- back pain
- or cold symptoms such as stuffy nose, sinus pain, sore throat .

1646. This particular disease, Hayfever, interacts with the drug Astelin and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common Astelin side effects may include: bitter taste in your mouth
- drowsiness, tiredness
- headache , dizziness
- sneezing, nasal irritation, nosebleeds

Drug Side Effects Summary

- nausea , dry mouth
- or weight gain.

1647. This particular disease, Hayfever, interacts with the drug azelastine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have signs of an ear infection : ear pain or full feeling
- drainage from the ear
- trouble hearing
- or fever. Common side effects of azelastine nasal may include: drowsiness, tiredness
- a bitter taste in your mouth
- mouth or throat pain
- nasal pain or discomfort, nosebleeds
- cold symptoms such as stuffy nose, sneezing, cough, sore throat
- fever, headache
- vomiting
- or skin itching around your nose.

1648. This particular disease, Hayfever, interacts with the drug Allegra Allergy and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Children's Allegra Allergy and call your doctor at once if you have: flu-like symptoms (fever, chills, unusual tiredness)
- new or worsening cough

Drug Side Effects Summary

- pain
- or signs of an ear infection--fever, ear pain or full feeling, trouble hearing, drainage from the ear, fussiness in a child. Common side effects of Children's Allegra Allergy may include: headache
- back pain
- or cold symptoms such as stuffy nose, sinus pain, sore throat .

1649. This particular disease, Hayfever, interacts with the drug Astepro and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have signs of an ear infection : ear pain or full feeling
- drainage from the ear
- trouble hearing
- or fever. Common Astepro side effects may include: drowsiness, tiredness
- a bitter taste in your mouth
- mouth or throat pain
- nasal pain or discomfort, nosebleeds
- cold symptoms such as stuffy nose, sneezing, cough, sore throat
- fever, headache
- vomiting
- or skin itching around your nose.

1650. This particular disease, Hayfever, interacts with the drug Flonase Allergy Relief and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if

Drug Side Effects Summary

you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of a weak adrenal gland like a very bad upset stomach or throwing up, very bad dizziness or passing out, muscle weakness, feeling very tired, mood changes, not hungry, or weight loss . Nose sores. Bad nosebleeds . Whistling sound when you breathe. Redness or white patches in mouth or throat. Change in eyesight. Nose discharge that is not normal. Very bad face pain. Crusting in the nose. Runny nose . Bone pain. Fever or chills. Very bad sore throat . Flonase Allergy Relief side effects

1651. This particular disease, Hayfever, interacts with the drug Dymista and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using the medicine and call your doctor at once if you have: severe drowsiness
- severe or ongoing nosebleeds
- noisy breathing, runny nose , or crusting around your nostrils
- redness, sores, or white patches in your mouth or throat
- fever, chills, tiredness, body aches

Drug Side Effects Summary

- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- any wound that will not heal
- or signs of low adrenal gland hormones - worsening tiredness or weakness, feeling light-headed, nausea , vomiting . Steroid medicine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common Dymista side effects may include:
headache
- nosebleeds
- or altered sense of taste.

1652. This particular disease, Hayfever, interacts with the drug Clarinex and has the following side

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Common Clarinex side effects may include: dry mouth, sore throat
- muscle pain
- drowsiness, tiredness
- or menstrual pain.

1653. This particular disease, Hayfever, interacts with the drug Nasacort Allergy 24HR and has the f

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: fever, chills, body aches, flu symptoms
- nosebleeds
- or blurred vision, tunnel vision, eye pain , or seeing halos around lights. Triamcinolone can affect

Drug Side Effects Summary

growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Less serious Nasacort side effects may be more likely, and you may have none at all.

1654. This particular disease, Hayfever, interacts with the drug Aller-Tec and has the following side effects:

- No side effects listed

1655. This particular disease, Hayfever, interacts with the drug desloratadine and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Common side effects of desloratadine may include:
dry mouth, sore throat
- muscle pain
- drowsiness, tiredness
- or menstrual pain.

1656. This particular disease, Hayfever, interacts with the drug mometasone and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Mometasone nasal may cause serious side effects.
Call your doctor at once if you have: severe bleeding or increased drainage from your nose
- nose pain or discomfort, headache
- white patches or sores in the nose that won't heal
- wheezing, trouble breathing
- vision problems
- irritation or a choking feeling in the back of your throat (may be signs that the implant has moved)

Drug Side Effects Summary

inside your nose)

- or ear pain or full feeling, trouble hearing, drainage from the ear. Steroid medicine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using mometasone nasal. Although the risk of serious side effects is low when mometasone is used in the nose, side effects can occur if the medicine is absorbed into your bloodstream. Tell your doctor if you have possible signs of long-term steroid use: weight gain (especially in your face or your upper back and torso)
- slow wound healing, thinning skin, increased body hair
- irregular menstrual periods, changes in sexual function
- or muscle weakness, tired feeling, depression , anxiety , or feeling irritable. Common side effects of mometasone nasal may include: nosebleeds
- headache
- stuffy nose, sore throat , cough
- or flu-like symptoms.

1657. This particular disease, Hayfever, interacts with the drug Phenergan and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Phenergan and call your doctor at once if you have: severe drowsiness, weak or shallow breathing
- a light-headed feeling, like you might pass out
- confusion, agitation, hallucinations, nightmares
- seizure (convulsions)
- fast or slow heartbeats
- jaundice (yellowing of the skin or eyes)

Drug Side Effects Summary

- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- easy bruising or bleeding (nosebleeds, bleeding gums)
- sudden weakness or ill feeling, fever, chills, sore throat, mouth sores, red or swollen gums, trouble swallowing
- or very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Side effects such as confusion and severe drowsiness may be more likely in older adults. Common Phenergan side effects may include: drowsiness, dizziness
- ringing in your ears
- double vision
- feeling nervous
- dry mouth
- or tired feeling, sleep problems (insomnia).

1658. This particular disease, Hayfever, interacts with the drug triamcinolone and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Triamcinolone may cause serious side effects. Call your doctor at once if you have: fever, chills, body aches, flu symptoms
- nosebleeds
- or blurred vision, tunnel vision, eye pain , or seeing halos around lights. Triamcinolone nasal can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Less serious side effects may be more likely, and you may have none at all.

1659. This particular disease, Hayfever, interacts with the drug Alavert and has the following side e

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common side effects of Alavert may include:
headache
- or feeling tired or drowsy.

1660. This particular disease, Hayfever, interacts with the drug Allegra-D 12 Hour and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Allegra-D and call your doctor at once if
you have: pounding heartbeats or fluttering in your chest
- or severe dizziness, anxiety, restless feeling, tremors, or nervousness. Common Allegra-D side
effects may include: mild dizziness
- dry mouth, nose, or throat
- nausea
- or sleep problems (insomnia).

1661. This particular disease, Hayfever, interacts with the drug Allegra-D 24 Hour and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if
you have: pounding heartbeats or fluttering in your chest
- or severe dizziness, anxiety, restless feeling, tremors, or nervousness. Common Allegra-D 24 Hour
side effects may include: mild dizziness
- dry mouth, nose, or throat

Drug Side Effects Summary

- nausea
- or sleep problems (insomnia).

1662. This particular disease, Hayfever, interacts with the drug Allergy Relief Tablets and has the following side effects:

- No side effects listed

1663. This particular disease, Hayfever, interacts with the drug budesonide and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. budesonide nasal may cause serious side effects.

Stop using budesonide nasal and call your doctor at once if you have: come into contact with anyone who has chickenpox, measles or tuberculosis

- severe or ongoing nosebleeds
- vision problems
- or fever, chills, body aches, flu symptoms. Less serious side effects may be more likely, and you may have none at all.

1664. This particular disease, Hayfever, interacts with the drug Claritin Reditabs and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common side effects of Claritin Reditab may include: headache
- or feeling tired or drowsy.

1665. This particular disease, Hayfever, interacts with the drug Flonase Sensimist and has the following side effects:

Drug Side Effects Summary

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

- itching

- red, swollen, blistered, or peeling skin with or without fever

- wheezing

- tightness in the chest or throat

- trouble breathing, swallowing, or talking

- unusual hoarseness

- or swelling of the mouth, face, lips, tongue, or throat. Signs of a weak adrenal gland like a very bad upset stomach or throwing up, very bad dizziness or passing out, muscle weakness, feeling very tired, mood changes, not hungry, or weight loss . Nose sores. Bad nosebleeds . Whistling sound when you breathe. Redness or white patches in mouth or throat. Change in eyesight. Nose discharge that is not normal. Very bad face pain. Crusting in the nose. Runny nose . Bone pain. Fever or chills. Very bad sore throat . Flonase Sensimist side effects

1666. This particular disease, Hayfever, interacts with the drug promethazine and has the following

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Stop using promethazine and call your doctor at once if you have: severe drowsiness, weak or shallow breathing

- a light-headed feeling, like you might pass out

- confusion, agitation, hallucinations, nightmares

Drug Side Effects Summary

- seizure (convulsions)
- fast or slow heartbeats
- jaundice (yellowing of the skin or eyes)
- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- easy bruising or bleeding (nosebleeds, bleeding gums)
- sudden weakness or ill feeling, fever, chills, sore throat, mouth sores, red or swollen gums, trouble swallowing
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Side effects such as confusion and severe drowsiness may be more likely in older adults. Common promethazine side effects may include: drowsiness, dizziness
- ringing in your ears
- double vision
- feeling nervous
- dry mouth
- or tired feeling, sleep problems (insomnia).

1667. This particular disease, Hayfever, interacts with the drug Zyrtec-D and has the following side

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Zyrtec-D may cause serious side effects. Stop using Zyrtec-D and call your doctor at once if you have: fast, pounding, or uneven heartbeat
- weakness, tremors (uncontrolled shaking) severe restless feeling, hyperactivity, extreme feeling of fear or confusion

Drug Side Effects Summary

- problems with vision
- little or no urinating
- or high blood pressure (severe headache , buzzing in your ears, chest pain, shortness of breath, uneven heartbeats). Common side effects of Zyrtec-D may include: dizziness , drowsiness, tired feeling
- sleep problems (insomnia)
- dry mouth, nausea , stomach pain, constipation
- or trouble concentrating.

1668. This particular disease, Hayfever, interacts with the drug All Day Allergy and has the following side effects:

- No side effects listed

1669. This particular disease, Hayfever, interacts with the drug All Day Allergy Children's and has the following side effects:

- No side effects listed

1670. This particular disease, Hayfever, interacts with the drug Allegra ODT and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Chest tightness feeling of warmth, redness of the face, neck, arms and occasionally, upper chest large, hive-like swelling on face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs shortness of breath, difficult or labored breathing Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are

Drug Side Effects Summary

bothersome or if you have any questions about them: More common Vomiting Less common Back pain body aches or pain chills coughing diarrhea difficulty with moving dizziness ear congestion earache fever headache joint pain loss of voice muscle aching or cramping muscle pains or stiffness nasal congestion nausea pain in arms or legs pain or tenderness around eyes or cheekbones painful menstrual bleeding redness or swelling in ear ringing or buzzing in ears runny or stuffy nose sleepiness or unusual drowsiness sneezing sore throat stomach upset swollen joints unusual feeling of tiredness or weakness viral infection (such as cold and flu) Rare Nervousness rash sleeplessness terrifying dreams trouble sleeping Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Allegra ODT side effects

1671. This particular disease, Hayfever, interacts with the drug Aller-Tec Children's and has the following side effects:

- No side effects listed

1672. This particular disease, Hayfever, interacts with the drug Allergy Relief 24 Hour and has the following side effects:

- No side effects listed

1673. This particular disease, Hayfever, interacts with the drug Antinaus 50 and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: Rare Convulsions (seizures) difficult or unusually fast breathing fast heartbeat or irregular pulse high fever high or low blood pressure increased sweating loss of bladder control severe muscle stiffness unusually pale skin unusual tiredness or weakness Incidence not known Abdominal or stomach pain black, tarry stools

Drug Side Effects Summary

blood in the urine bloody nose blurred vision chest pain or discomfort chills clay-colored stools cough or hoarseness dark urine decreased awareness or responsiveness difficulty with speaking dizziness double vision drooling fast, pounding, or irregular heartbeat or pulse fever with or without chills fixed position of the eye headache heavier menstrual periods increases and decreases in blood pressure itching large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs lightheadedness, dizziness, or fainting loss of appetite loss of balance control lower back or side pain mimicry of speech or movements muscle trembling, jerking, or stiffness mutism nausea negativism noisy breathing painful or difficult urination peculiar postures or movements, mannerisms, or grimacing pinpoint red spots on the skin restlessness seeing double seeing, hearing, or feeling things that are not there seizures severe sleepiness shortness of breath shuffling walk skin rash slow or irregular heartbeat sore throat sores, ulcers, or white spots on the lips or in the mouth stiffness of the limbs swollen glands tenderness, pain, swelling, warmth, skin discoloration, and prominent superficial veins over the affected area tightness in the chest tremors twisting movements of the body uncontrolled movements, especially of the face, neck, and back unpleasant breath odor unusual bleeding or bruising vomiting of blood wheezing yellow eyes or skin

Symptom of overdose Abnormal movements confusion as to time, place, or person dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position feeling of warmth holding false beliefs that cannot be changed by fact hyperexcitability irregular, fast or slow, or shallow breathing nightmares pale or blue lips, fingernails, or skin pupils of the eyes, large and not moving or responding to light redness of the face, neck, arms, and occasionally, upper chest sweating troubled breathing unusual excitement, nervousness, or restlessness

Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any

Drug Side Effects Summary

questions about them: Incidence not known Blistering, crusting, irritation, or reddening of the skin continuing ringing or buzzing or other unexplained noise in the ears discoloration of the skin drowsiness dry mouth excitation false or unusual sense of well-being hearing loss hives or welts increased sensitivity of the skin to sunlight lack of coordination loss of strength or energy muscle pain or weakness nasal stuffiness nervousness severe sunburn sleeplessness trouble sleeping unable to sleep vomiting Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Antinaus 50 side effects

1674. This particular disease, Hayfever, interacts with the drug beclomethasone and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Beclomethasone nasal may cause serious side effects. Call your doctor at once if you have: severe or ongoing nosebleed
- sores in the nose that won't heal
- blurred vision, eye pain , or seeing halos around lights
- signs of a hormonal disorder--worsening tiredness or muscle weakness, anxiety , feeling irritable, feeling light-headed, nausea , vomiting , loss of appetite, diarrhea , weight loss , or weight gain (especially in your face or your upper back and torso)
- or signs of infection--fever, chills, sores or white patches in or around your nose, flu symptoms, redness or swelling. Beclomethasone nasal can affect growth in children. Talk with your doctor if you think your child is not growing at a normal rate while using beclomethasone nasal. Common side effects of beclomethasone nasal may include: discomfort or irritation inside your nose
- sneezing, runny or stuffy nose
- nosebleed

Drug Side Effects Summary

- fever, sore throat
- headache , nausea
- or unpleasant taste or smell.

1675. This particular disease, Hayfever, interacts with the drug cetirizine / pseudoephedrine and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Cetirizine and pseudoephedrine may cause serious side effects. Stop using cetirizine and pseudoephedrine and call your doctor at once if you have: fast, pounding, or uneven heartbeat
- weakness, tremors (uncontrolled shaking) severe restless feeling, hyperactivity, extreme feeling of fear or confusion
- problems with vision
- little or no urinating
- or high blood pressure (severe headache , buzzing in your ears, chest pain, shortness of breath, uneven heartbeats). Common side effects of cetirizine and pseudoephedrine may include: dizziness , drowsiness, tired feeling
- sleep problems (insomnia)
- dry mouth, nausea , stomach pain, constipation
- or trouble concentrating.

1676. This particular disease, Hayfever, interacts with the drug Children's Claritin Allergy and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs

Drug Side Effects Summary

of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Feeling very tired or weak. Children's Claritin

Allergy side effects

1677. This particular disease, Hayfever, interacts with the drug Claritin-D and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- or severe dizziness , nervousness, or restless feeling. Common Claritin-D side effects may include: dry mouth, nose, or throat
- mild dizziness
- or sleep problems (insomnia).

1678. This particular disease, Hayfever, interacts with the drug Claritin-D 24 Hour and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop

Drug Side Effects Summary

using this medicine and call your doctor at once if you have: pounding heartbeats or fluttering in your chest

- or severe dizziness , nervousness , or restless feeling. Common side effects of Claritin-D 24 Hour may include: dry mouth, nose, or throat

- mild dizziness

- or sleep problems (insomnia).

1679. This particular disease, Hayfever, interacts with the drug Clinacort and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: More common Aggression agitation anxiety blurred vision decrease in the amount of urine dizziness fast, slow, pounding, or irregular heartbeat or pulse headache irritability mental depression mood changes nervousness noisy, rattling breathing numbness or tingling in the arms or legs pounding in the ears shortness of breath swelling of the fingers, hands, feet, or lower legs trouble thinking, speaking, or walking troubled breathing at rest weight gain Incidence not known Abdominal cramping and/or burning (severe) abdominal pain backache bloody, black, or tarry stools cough or hoarseness darkening of skin decreased vision diarrhea dry mouth eye pain eye tearing facial hair growth in females fainting fatigue fever or chills flushed, dry skin fractures fruit-like breath odor full or round face, neck, or trunk heartburn and/or indigestion (severe and continuous) increased hunger increased thirst increased urination loss of appetite loss of sexual desire or ability lower back or side pain menstrual irregularities muscle pain or tenderness muscle wasting or weakness nausea pain in back, ribs, arms, or legs painful or difficult urination skin rash sleeplessness sweating trouble healing trouble sleeping unexplained weight loss unusual tiredness or weakness vision changes vomiting vomiting of material that looks like coffee grounds Some side effects may occur that usually do not need

Drug Side Effects Summary

medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Blemishes on the skin bruising dry, scaly skin feeling of constant movement of self or surroundings full or bloated feeling increased appetite increased hair growth on the face, forehead, back, arms, and legs large, flat, blue, or purplish patches in the skin pimples pitting or depression of the skin at the injection site reddish purple lines on the arms, face, legs, trunk, or groin redness of the skin redness, swelling, tenderness, or pain at the injection site sensation of spinning small, red, or purple spots on the skin thin, fragile, or shiny skin thinning of the scalp hair Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Clinacort side effects

1680. This particular disease, Hayfever, interacts with the drug cyproheptadine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Cyproheptadine may cause serious side effects. Stop using cyproheptadine and call your doctor at once if you have: a light-headed feeling, like you might pass out
- tremor, seizure (convulsions)
- confusion, hallucinations
- little or no urination
- fast or pounding heartbeats
- easy bruising or bleeding

Drug Side Effects Summary

- ringing in your ears
- or pale or yellowed skin, dark colored urine, fever, weakness. Common side effects of cyproheptadine may include: drowsiness, dizziness
- dry mouth, nose, or throat
- constipation
- blurred vision
- or feeling restless or excited (especially in children).

1681. This particular disease, Hayfever, interacts with the drug flunisolide and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Flunisolide nasal may cause serious side effects. Call your doctor at once if you have: severe or ongoing nosebleeds
- itching, irritation, sores, or white patches in your nose, mouth, or throat
- crusting around your nostrils
- any wound that will not heal
- or fever, chills, weakness, flu symptoms. Common side effects of flunisolide nasal may include: burning or stinging in your nose
- stuffy nose, sneezing, minor nosebleed
- changes in your sense of smell or taste
- watery eyes
- irritation in your nose, sore throat
- nausea , vomiting
- or headache .

Drug Side Effects Summary

1682. This particular disease, Hayfever, interacts with the drug Loratadine-D 24 Hour and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- or severe dizziness , nervousness , or restless feeling. Common side effects of Loratadine-D 24 Hour may include: dry mouth, nose, or throat
- mild dizziness
- or sleep problems (insomnia).

1683. This particular disease, Hayfever, interacts with the drug ohm Allergy Relief and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common side effects of ohm Allergy Relief may include: headache
- or feeling tired or drowsy.

1684. This particular disease, Hayfever, interacts with the drug Omnaris and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Nasal steroid medicine can be absorbed into your bloodstream, which may cause steroid side effects throughout the body. Tell your doctor if you have: weight gain (especially in your face or your upper back and torso)
- slow wound healing, thinning skin, increased body hair

Drug Side Effects Summary

- irregular menstrual periods, changes in sexual function
- or muscle weakness, tired feeling, depression, anxiety, feeling irritable. Call your doctor at once if you have: runny or bloody nose, crusting inside the nose, whistling sound when you breathe
- sores or white patches in or around your nostrils
- fever, chills, body aches, nausea, vomiting
- blurred vision, tunnel vision, eye pain, seeing halos around lights
- or signs of a hormonal disorder - worsening tiredness or weakness, feeling light-headed, vomiting, loss of appetite, diarrhea, and weight loss. Common Omnaris side effects may include: headache, dizziness
- nosebleed, nasal irritation
- stuffy nose, sore throat
- or ear pain.

1685. This particular disease, Hayfever, interacts with the drug Patanase and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Patanase and call your doctor at once if you have: nosebleed
- painful sores inside your nose
- fever
- or pain or burning when you urinate. Common Patanase side effects may include: runny or stuffy nose, nosebleeds
- cough, throat pain or irritation
- a bitter taste in your mouth
- drowsiness

Drug Side Effects Summary

- headache
- fever
- rash
- or painful urination.

1686. This particular disease, Hayfever, interacts with the drug Phenadoz and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Phenadoz may cause serious side effects. Call your doctor at once if you have: severe drowsiness, weak or shallow breathing
- a light-headed feeling, like you might pass out
- confusion, agitation , hallucinations, nightmares
- seizure (convulsions)
- fast or slow heartbeats
- jaundice (yellowing of the skin or eyes)
- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- easy bruising or bleeding (nosebleeds , bleeding gums)
- sudden weakness or ill feeling, fever, chills, sore throat , mouth sores, red or swollen gums, trouble swallowing
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Common side effects of Phenadoz may include: drowsiness, dizziness
- ringing in your ears
- double vision

Drug Side Effects Summary

- feeling nervous
- dry mouth
- or tiredness, sleep problems (insomnia).

1687. This particular disease, Hayfever, interacts with the drug Promethegan and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Promethegan may cause serious side effects. Call your doctor at once if you have: severe drowsiness, weak or shallow breathing
- a light-headed feeling, like you might pass out
- confusion, agitation , hallucinations, nightmares
- seizure (convulsions)
- fast or slow heartbeats
- jaundice (yellowing of the skin or eyes)
- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- easy bruising or bleeding (nosebleeds , bleeding gums)
- sudden weakness or ill feeling, fever, chills, sore throat , mouth sores, red or swollen gums, trouble swallowing
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Common side effects of Promethegan may include: drowsiness, dizziness
- ringing in your ears
- double vision
- feeling nervous

Drug Side Effects Summary

- dry mouth
- or tiredness, sleep problems (insomnia).

1688. This particular disease, Hayfever, interacts with the drug QNASL and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe or ongoing nosebleed
- sores in the nose that won't heal
- blurred vision, eye pain , or seeing halos around lights
- signs of a hormonal disorder - worsening tiredness or muscle weakness, anxiety , feeling irritable, feeling light-headed, nausea , vomiting , loss of appetite, diarrhea , weight loss , or weight gain (especially in your face or your upper back and torso)
- or signs of infection - fever, chills, sores or white patches in or around your nose, flu symptoms, redness or swelling. Beclomethasone can affect growth in children. Talk with your doctor if you think your child is not growing at a normal rate while using this medicine. Common QNASL side effects may include: discomfort or irritation inside your nose
- sneezing, runny or stuffy nose
- nosebleed
- fever, sore throat
- headache , nausea
- or unpleasant taste or smell.

1689. This particular disease, Hayfever, interacts with the drug Tavist ND and has the following side effects

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Common side effects of Tavist ND may include:
headache
- or feeling tired or drowsy.

1690. This particular disease, Hayfever, interacts with the drug Ticanase and has the following side effects

- No side effects listed

1691. This particular disease, Hayfever, interacts with the drug Wal-itin and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common side effects of Wal-itin may include:
headache
- or feeling tired or drowsy.

1692. This particular disease, Hayfever, interacts with the drug ZoDex and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat

Drug Side Effects Summary

- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of low potassium levels like muscle pain or weakness, muscle cramps, or a heartbeat that does not feel normal. Signs of a pancreas problem (pancreatitis) like very bad stomach pain, very bad back pain , or very bad upset stomach or throwing up. Signs of high blood pressure like very bad headache or dizziness , passing out, or change in eyesight. Signs of a weak adrenal gland like a very bad upset stomach or throwing up, very bad dizziness or passing out, muscle weakness, feeling very tired, mood changes, not hungry, or weight loss . Signs of Cushing's disease like weight gain in the upper back or belly, moon face, very bad headache, or slow healing. Feeling very tired, weak, or touchy
- trembling
- having a fast heartbeat , confusion, sweating, or dizziness if you missed a dose or recently stopped ZoDex (dexamethasone tablets). Shortness of breath, a big weight gain, or swelling in the arms or legs. Skin changes (pimples , stretch marks, slow healing, hair growth). Fast, slow, or abnormal heartbeat. Chest pain or pressure. Swelling, warmth, numbness , change of color, or pain in a leg or arm. Period (menstrual) changes. Bone or joint pain. Muscle pain or weakness. Change in eyesight. Change in the way you act. Depression or other mood changes. Seizures . A burning, numbness, or tingling feeling that is not normal. Any unexplained bruising or bleeding. Very bad belly pain. Black, tarry, or bloody stools. Throwing up blood or throw up that looks like coffee grounds. ZoDex side effects

1693. This particular disease, Hayfever, interacts with the drug Zyrtec-D 12 Hour and has the following

- No side effects listed

1694. This particular disease, Hayfever, interacts with the drug acrivastine / pseudoephedrine and has the following

Drug Side Effects Summary

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Symptoms of overdose Blurred vision chest pain or discomfort cold, clammy, or pale skin confusion dizziness dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position irregular heartbeats seizures shortness of breath slow heart rate sweating unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Dry mouth headache sleepiness or unusual drowsiness Less common Acid or sour stomach belching body aches or pain congestion cough dryness or soreness of the throat fever heartburn hoarseness indigestion lack or loss of strength nausea nervousness runny nose stomach discomfort, upset, or pain tender, swollen glands in the neck trouble sleeping trouble swallowing voice changes Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Acrivastine / pseudoephedrine side effects

1695. This particular disease, Hayfever, interacts with the drug Aller-Ease and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Aller-Ease may cause serious side effects. Stop using Aller-Ease and call your doctor at once if you have: flu-like symptoms (fever, chills, unusual tiredness)
- new or worsening cough

Drug Side Effects Summary

- pain
- or signs of an ear infection--fever, ear pain or full feeling, trouble hearing, drainage from the ear, fussiness in a child. Common side effects of Aller-Ease may include: headache
- back pain
- or cold symptoms such as stuffy nose, sinus pain, sore throat .

1696. This particular disease, Hayfever, interacts with the drug Astepro Allergy and has the following

- No side effects listed

1697. This particular disease, Hayfever, interacts with the drug azelastine / fluticasone and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: severe drowsiness
- severe or ongoing nosebleeds
- noisy breathing, runny nose , or crusting around your nostrils
- redness, sores, or white patches in your mouth or throat
- fever, chills, tiredness, body aches
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- any wound that will not heal
- or signs of low adrenal gland hormones --worsening tiredness or weakness, feeling light-headed, nausea , vomiting . Steroid medicine can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using azelastine and fluticasone nasal. Common side effects of azelastine and fluticasone nasal may include: headache
- nosebleeds

Drug Side Effects Summary

- or altered sense of taste.

1698. This particular disease, Hayfever, interacts with the drug Beconase AQ and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Beconase AQ may cause serious side effects. Call your doctor at once if you have: severe or ongoing nosebleed
- sores in the nose that won't heal
- blurred vision, eye pain , or seeing halos around lights
- signs of a hormonal disorder--worsening tiredness or muscle weakness, anxiety , feeling irritable, feeling light-headed, nausea , vomiting , loss of appetite, diarrhea , weight loss , or weight gain (especially in your face or your upper back and torso)
- or signs of infection--fever, chills, sores or white patches in or around your nose, flu symptoms, redness or swelling. Beconase AQ can affect growth in children. Talk with your doctor if you think your child is not growing at a normal rate while using Beconase AQ. Common side effects of Beconase AQ may include: discomfort or irritation inside your nose
- sneezing, runny or stuffy nose
- nosebleed
- fever, sore throat
- headache , nausea
- or unpleasant taste or smell.

1699. This particular disease, Hayfever, interacts with the drug Claritin-D 12 Hour and has the follow

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- or severe dizziness , nervousness , or restless feeling. Common side effects of Claritin-D 12 Hour may include: dry mouth, nose, or throat
- mild dizziness
- or sleep problems (insomnia).

1700. This particular disease, Hayfever, interacts with the drug fexofenadine / pseudoephedrine and

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Fexofenadine and pseudoephedrine may cause serious side effects. Stop using fexofenadine and pseudoephedrine and call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- or severe dizziness , anxiety , restless feeling, tremors, or nervousness . Common side effects of fexofenadine and pseudoephedrine may include: mild dizziness
- dry mouth, nose, or throat
- nausea
- or sleep problems (insomnia).

1701. This particular disease, Hayfever, interacts with the drug loratadine / pseudoephedrine and

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: pounding heartbeats or fluttering in

Drug Side Effects Summary

your chest

- or severe dizziness , nervousness , or restless feeling. Common side effects of loratadine and pseudoephedrine may include: dry mouth, nose, or throat

- mild dizziness

- or sleep problems (insomnia).

1702. This particular disease, Hayfever, interacts with the drug Mucinex Allergy and has the following

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Mucinex Allergy may cause serious side effects. Stop using Mucinex Allergy and call your doctor at once if you have: flu-like symptoms (fever, chills, unusual tiredness)

- new or worsening cough

- pain

- or signs of an ear infection--fever, ear pain or full feeling, trouble hearing, drainage from the ear, fussiness in a child. Common side effects of Mucinex Allergy may include: headache

- back pain

- or cold symptoms such as stuffy nose, sinus pain, sore throat .

1703. This particular disease, Hayfever, interacts with the drug NasalCrom and has the following side

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor if you have: severe burning, stinging, or irritation in your nose

- nosebleeds , sinus pain, or sores in your nose

Drug Side Effects Summary

- wheezing, tight feeling in your chest
- or fever, green or yellow mucus from the nose. Common side effects may include sneezing or mild stinging inside your nose after use.

1704. This particular disease, Hayfever, interacts with the drug olopatadine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Olopatadine may cause serious side effects. Stop using olopatadine and call your doctor at once if you have: nosebleed
- painful sores inside your nose
- fever
- or pain or burning when you urinate. Common side effects of olopatadine may include: runny or stuffy nose, nosebleeds
- cough, throat pain or irritation
- a bitter taste in your mouth
- drowsiness
- headache
- fever
- rash
- or painful urination.

1705. This particular disease, Hayfever, interacts with the drug Rhinocort Allergy and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Rhinocort Allergy may cause serious side effects. Stop

Drug Side Effects Summary

using Rhinocort Allergy and call your doctor at once if you have: come into contact with anyone who has chickenpox, measles or tuberculosis

- severe or ongoing nosebleeds
- vision problems
- or fever, chills, body aches, flu symptoms. Less serious side effects may be more likely, and you may have none at all.

1706. This particular disease, Hayfever, interacts with the drug Alavert D-12 Hour Allergy and Sinus

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- or severe dizziness , nervousness , or restless feeling. Common side effects of Alavert D-12 may include: dry mouth, nose, or throat
- mild dizziness
- or sleep problems (insomnia).

1707. This particular disease, Hayfever, interacts with the drug All Day Allergy-D and has the follow

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. All Day Allergy-D may cause serious side effects. Stop using All Day Allergy-D and call your doctor at once if you have: fast, pounding, or uneven heartbeat
- weakness, tremors (uncontrolled shaking) severe restless feeling, hyperactivity, extreme feeling of fear or confusion

Drug Side Effects Summary

- problems with vision
- little or no urinating
- or high blood pressure (severe headache , buzzing in your ears, chest pain, shortness of breath, uneven heartbeats). Common side effects of All Day Allergy-D may include: dizziness , drowsiness, tired feeling
- sleep problems (insomnia)
- dry mouth, nausea , stomach pain, constipation
- or trouble concentrating.

1708. This particular disease, Hayfever, interacts with the drug Allerest Maximum Strength and has

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate
- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)
- little or no urinating
- easy bruising or bleeding, unusual weakness
- or dangerously high blood pressure (severe headache, buzzing in your ears, anxiety , shortness of breath). Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of Allerest Maximum Strength may include: dizziness , drowsiness, blurred vision
- dry nose or mouth
- nausea , stomach pain, constipation, loss of appetite

Drug Side Effects Summary

- problems with memory or concentration
- or feeling restless or excited (especially in children).

1709. This particular disease, Hayfever, interacts with the drug Allergy & Congestion Relief and has

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Dizziness drowsiness nervousness trouble sleeping Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Allergy & Congestion Relief side effects

1710. This particular disease, Hayfever, interacts with the drug Allergy Relief D and has the following

- No side effects listed

1711. This particular disease, Hayfever, interacts with the drug Arbinoxa and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blurred vision chills confusion cough decrease in the frequency of urination decrease in urine volume difficult or painful urination difficulty in passing urine (dribbling) difficulty with swallowing dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position fast, irregular, pounding, or

Drug Side Effects Summary

racing heartbeat or pulse hives or welts, itching, skin rash puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue redness of the skin sweating tightness in the chest unusual tiredness or weakness Symptoms of overdose Agitation anxiety bloating diarrhea difficulty with sleeping disorientation dizziness drowsiness to profound coma dry mouth feeling of warmth fixed and enlarged pupils gas headache lethargy lightheadedness loss of appetite nausea redness of the face, neck, arms, and occasionally, upper chest seeing, hearing, or feeling things that are not there seizures tightness of the chest trouble breathing vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Clumsiness or unsteadiness cough producing mucus drowsiness sleepiness Incidence not known Back, leg, or stomach pains black, tarry stools bleeding gums blood in the urine or stools blurred vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chills confusion continuing ringing or buzzing or other unexplained noise in the ears cough or hoarseness dark urine difficulty breathing difficulty having a bowel movement (stool) dizziness or lightheadedness double vision dry mouth, nose, and throat feeling excited feeling of constant movement of self or surroundings fever with or without chills general body swelling general feeling of tiredness or weakness headache hearing loss increased sensitivity of the skin to sunlight itching loss of appetite lower back or side pain nosebleeds pale skin pinpoint red spots on the skin redness or other discoloration of the skin restlessness seeing double seizures sensation of spinning severe sunburn shakiness in the legs, arms, hands, or feet sore throat sores, ulcers, or white spots on the lips or in the mouth trembling or shaking of the hands or feet trouble sleeping unusual bleeding or bruising weight loss yellowing of the eyes or skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your

Drug Side Effects Summary

doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Arbinoxa side effects

1712. This particular disease, Hayfever, interacts with the drug Brohist D and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: nervousness , dizziness , or sleeplessness
- chest pain, fast or uneven heart rate
- little or no urinating
- dangerously high blood pressure (severe headache , buzzing in your ears, anxiety , shortness of breath)
- if your symptoms do not improve after 7 days of treatment, or if you have a fever
- or if new symptoms occur. Less serious side effects may be more likely, and you may have none at all.

1713. This particular disease, Hayfever, interacts with the drug Bromaline and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Bromaline may cause serious side effects. Stop using Bromaline and call your doctor at once if you have: nervousness , dizziness , or sleeplessness
- chest pain, fast or uneven heart rate
- little or no urinating
- dangerously high blood pressure (severe headache , buzzing in your ears, anxiety , shortness of breath)

Drug Side Effects Summary

breath)

- if your symptoms do not improve after 7 days of treatment, or if you have a fever
- or if new symptoms occur. Less serious side effects may be more likely, and you may have none at all.

1714. This particular disease, Hayfever, interacts with the drug brompheniramine / phenylephrine a

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: nervousness , dizziness , or sleeplessness
- chest pain, fast or uneven heart rate
- little or no urinating
- dangerously high blood pressure (severe headache , buzzing in your ears, anxiety , shortness of breath)
- if your symptoms do not improve after 7 days of treatment, or if you have a fever
- or if new symptoms occur. Less serious side effects may be more likely, and you may have none at all.

1715. This particular disease, Hayfever, interacts with the drug brompheniramine / pseudoephedrin

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Brompheniramine and pseudoephedrine may cause serious side effects. Stop using brompheniramine and pseudoephedrine and call your doctor at once if you have: nervousness , dizziness , or sleeplessness

Drug Side Effects Summary

- chest pain, fast or uneven heart rate
- little or no urinating
- dangerously high blood pressure (severe headache , buzzing in your ears, anxiety , shortness of breath)
- if your symptoms do not improve after 7 days of treatment, or if you have a fever
- or if new symptoms occur. Less serious side effects may be more likely, and you may have none at all.

1716. This particular disease, Hayfever, interacts with the drug Brotapp and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Brotapp may cause serious side effects. Stop using Brotapp and call your doctor at once if you have: nervousness , dizziness , or sleeplessness
- chest pain, fast or uneven heart rate
- little or no urinating
- dangerously high blood pressure (severe headache , buzzing in your ears, anxiety , shortness of breath)
- if your symptoms do not improve after 7 days of treatment, or if you have a fever
- or if new symptoms occur. Less serious side effects may be more likely, and you may have none at all.

1717. This particular disease, Hayfever, interacts with the drug carbinoxamine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Carbinoxamine may cause serious side effects. Stop

Drug Side Effects Summary

using carbinoxamine and call your doctor at once if you have: a light-headed feeling, like you might pass out

- little or no urination
- wheezing, tightness in your chest
- pounding heartbeats or fluttering in your chest
- or pale skin, easy bruising or bleeding. Side effects such as dizziness and confusion may be more likely in older adults. Common side effects of carbinoxamine may include: dry mouth, nose, or throat
- drowsiness, dizziness
- loss of coordination
- or upset stomach .

1718. This particular disease, Hayfever, interacts with the drug Carboxine and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blurred vision chills confusion cough decrease in the frequency of urination decrease in urine volume difficult or painful urination difficulty in passing urine (dribbling) difficulty with swallowing dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position fast, irregular, pounding, or racing heartbeat or pulse hives or welts, itching, skin rash puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue redness of the skin sweating tightness in the chest unusual tiredness or weakness Symptoms of overdose Agitation anxiety bloating diarrhea difficulty with sleeping disorientation dizziness drowsiness to profound coma dry mouth feeling of warmth fixed and enlarged pupils gas headache lethargy lightheadedness loss of appetite nausea redness of the face, neck, arms, and occasionally, upper chest seeing, hearing, or feeling things that are not there seizures tightness of the chest trouble breathing vomiting Some side effects may occur that usually

Drug Side Effects Summary

do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Clumsiness or unsteadiness cough producing mucus drowsiness sleepiness Incidence not known Back, leg, or stomach pains black, tarry stools bleeding gums blood in the urine or stools blurred vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chills confusion continuing ringing or buzzing or other unexplained noise in the ears cough or hoarseness dark urine difficulty breathing difficulty having a bowel movement (stool) dizziness or lightheadedness double vision dry mouth, nose, and throat feeling excited feeling of constant movement of self or surroundings fever with or without chills general body swelling general feeling of tiredness or weakness headache hearing loss increased sensitivity of the skin to sunlight itching loss of appetite lower back or side pain nosebleeds pale skin pinpoint red spots on the skin redness or other discoloration of the skin restlessness seeing double seizures sensation of spinning severe sunburn shakiness in the legs, arms, hands, or feet sore throat sores, ulcers, or white spots on the lips or in the mouth trembling or shaking of the hands or feet trouble sleeping unusual bleeding or bruising weight loss yellowing of the eyes or skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Carboxine side effects

1719. This particular disease, Hayfever, interacts with the drug chlorpheniramine / pseudoephedrine

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop

Drug Side Effects Summary

using this medicine and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate

- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)
- little or no urinating
- easy bruising or bleeding, unusual weakness
- or dangerously high blood pressure (severe headache, buzzing in your ears, anxiety , shortness of breath). Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of chlorpheniramine and pseudoephedrine may include: dizziness , drowsiness, blurred vision
- dry nose or mouth
- nausea , stomach pain, constipation, loss of appetite
- problems with memory or concentration
- or feeling restless or excited (especially in children).

1720. This particular disease, Hayfever, interacts with the drug ciclesonide and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Nasal steroid medicine can be absorbed into your bloodstream, which may cause steroid side effects throughout the body. Tell your doctor if you have: weight gain (especially in your face or your upper back and torso)
- slow wound healing, thinning skin, increased body hair
- irregular menstrual periods, changes in sexual function
- or muscle weakness, tired feeling, depression , anxiety , feeling irritable. Ciclesonide nasal may cause serious side effects. Call your doctor at once if you have: runny or bloody nose, crusting

Drug Side Effects Summary

inside the nose, whistling sound when you breathe

- sores or white patches in or around your nostrils
- fever, chills, body aches, nausea , vomiting
- blurred vision, tunnel vision, eye pain , seeing halos around lights
- or signs of a hormonal disorder--worsening tiredness or weakness, feeling light-headed, vomiting, loss of appetite, diarrhea , and weight loss . Common side effects of ciclesonide nasal may include: headache , dizziness
- nosebleed, nasal irritation
- stuffy nose, sore throat
- or ear pain.

1721. This particular disease, Hayfever, interacts with the drug Clarinex-D 12 Hour and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: a light-headed feeling, like you might pass out
- tremors
- severe weakness
- irregular heartbeats
- a seizure
- or a severe skin reaction - fever, skin redness, small pimples . Common Clarinex-D 12 Hour side effects may include: drowsiness, dizziness
- headache nausea, loss of appetite
- sleep problems (insomnia)
- or dry mouth, sore throat .

Drug Side Effects Summary

1722. This particular disease, Hayfever, interacts with the drug Clarinex-D 24 Hour and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Clarinex-D 24 Hour and call your doctor at once if you have: a light-headed feeling, like you might pass out
- tremors
- severe weakness
- irregular heartbeats
- a seizure
- or a severe skin reaction - fever, skin redness, small pimples. Common Clarinex-D 24 Hour side effects may include: drowsiness, dizziness
- headache nausea, loss of appetite
- sleep problems (insomnia)
- or dry mouth, sore throat.

1723. This particular disease, Hayfever, interacts with the drug Clarinex Reditabs and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Anaphylaxis, such as, cough, difficulty swallowing, dizziness, fast heartbeat, hives, itching, puffiness or swelling of eyelids or around the eyes or face or lips or tongue, shortness of breath, skin rash, tightness in chest, unusual tiredness or weakness, wheezing dyspnea, such as, shortness of breath, difficult or labored breathing, tightness in chest, wheezing edema, such as, swelling pruritus, such as, itching skin rash tachycardia, such as, fast, pounding, or irregular heartbeat or pulse urticaria, such as, hives or welts, itching, redness of skin, skin rash. Some side effects may occur that usually do not need medical

Drug Side Effects Summary

attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Headache Less common Dizziness dry mouth dysmenorrhea, such as, difficult or painful menstruation dyspepsia, such as, acid or sour stomach, belching, heartburn, indigestion, stomach discomfort , upset or pain, fatigue, such as, unusual tiredness or weakness myalgia, such as, joint pain, swollen joints, muscle aching or cramping, muscle pains or stiffness, difficulty in moving pharyngitis, such as, body aches or pain, congestion, cough, dryness or soreness of throat, fever, hoarseness, runny nose, tender swollen glands in neck, trouble in swallowing, voice changes. somnolence, such as, sleepiness or unusual drowsiness nausea Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Clarinet Reditabs side effects

1724. This particular disease, Hayfever, interacts with the drug Clear-Atadine-D and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Dizziness drowsiness nervousness trouble sleeping Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may

Drug Side Effects Summary

report side effects to the FDA at 1-800-FDA-1088. Clear-Atadine-D side effects

1725. This particular disease, Hayfever, interacts with the drug clemastine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Side effects may be more likely in older adults.

Common side effects of clemastine may include: drowsiness

- blurred vision
- or feeling restless or excited (especially in children).

1726. This particular disease, Hayfever, interacts with the drug cromolyn and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor if you have: severe burning, stinging, or irritation in your nose
- nosebleeds , sinus pain, or sores in your nose
- wheezing, tight feeling in your chest
- or fever, green or yellow mucus from the nose. Common side effects may include sneezing or mild stinging inside your nose after use.

1727. This particular disease, Hayfever, interacts with the drug Decon-A and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

Drug Side Effects Summary

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Decon-A side effects

1728. This particular disease, Hayfever, interacts with the drug Deconamine and has the following s

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Deconamine side effects

1729. This particular disease, Hayfever, interacts with the drug desloratadine / pseudoephedrine an

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: a light-headed feeling, like you might pass out
- tremors
- severe weakness
- irregular heartbeats
- a seizure
- or a severe skin reaction--fever, skin redness, small pimples . Common side effects of desloratadine and pseudoephedrine may include: drowsiness, dizziness
- headache nausea , loss of appetite
- sleep problems (insomnia)
- or dry mouth, sore throat .

1730. This particular disease, Hayfever, interacts with the drug dexchlorpheniramine / phenylephrine

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe dizziness
- nervousness , trouble sleeping
- little or no urination
- or high blood pressure (severe headache, pounding in your neck or ears). Common side effects of dexchlorpheniramine and phenylephrine may include: dizziness, drowsiness
- dry mouth, nose, or throat
- constipation

Drug Side Effects Summary

- blurred vision
- or feeling restless or excited (especially in children).

1731. This particular disease, Hayfever, interacts with the drug dexchlorpheniramine and has the following side effects:

- Stop taking this medicine and seek emergency medical attention if you experience an allergic reaction to dexchlorpheniramine: (difficulty breathing
- closing of your throat
- swelling of your lips, tongue, or face
- or hives). Other, less common dexchlorpheniramine side effects may be more likely to occur. Continue to take this medicine and talk to your doctor if you experience sleepiness, fatigue, or dizziness
- headache
- dry mouth
- or difficulty urinating or an enlarged prostate. Side effects other than those listed here may also occur. Talk to your doctor about any side effect that seems unusual or that is especially bothersome. You may report side effects to FDA at 1-800-FDA-1088. Dexchlorpheniramine side effects

1732. This particular disease, Hayfever, interacts with the drug Dicel and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate
- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)

Drug Side Effects Summary

- little or no urinating
- easy bruising or bleeding, unusual weakness
- or dangerously high blood pressure (severe headache, buzzing in your ears, anxiety , shortness of breath). Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of Dicl may include: dizziness , drowsiness, blurred vision
- dry nose or mouth
- nausea , stomach pain, constipation, loss of appetite
- problems with memory or concentration
- or feeling restless or excited (especially in children).

1733. This particular disease, Hayfever, interacts with the drug Dimetane Decongestant and has the

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: nervousness , dizziness , or sleeplessness
- chest pain, fast or uneven heart rate
- little or no urinating
- dangerously high blood pressure (severe headache , buzzing in your ears, anxiety , shortness of breath)
- if your symptoms do not improve after 7 days of treatment, or if you have a fever
- or if new symptoms occur. Less serious side effects may be more likely, and you may have none at all.

1734. This particular disease, Hayfever, interacts with the drug Dimetapp Children's Cold & Allergy

Drug Side Effects Summary

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using Dimetapp Cold & Allergy and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate
- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)
- little or no urinating
- easy bruising or bleeding, unusual weakness
- or dangerously high blood pressure (severe headache, buzzing in your ears, anxiety, shortness of breath). Common Dimetapp side effects may include: dizziness, drowsiness, blurred vision
- dry nose or mouth
- nausea, stomach pain, constipation, loss of appetite
- problems with memory or concentration
- or feeling restless or excited (especially in children).

1735. This particular disease, Hayfever, interacts with the drug Grastek and has the following side effects:

- Grastek may cause a severe allergic reaction. Stop taking the medicine and get emergency medical help if you have any of these signs of an allergic reaction : severe dizziness or a light-headed feeling (like you might pass out)
- fast heart rate
- stomach pain, vomiting, diarrhea
- hives, itching, flushing (warmth, redness, or tingly feeling)
- cough, chest tightness, wheezing, trouble breathing
- or swelling of your face, lips, tongue, or throat. If you stop taking Grastek after having an allergic reaction, do not start taking the medicine again without your doctor's advice. Grastek may cause

Drug Side Effects Summary

serious side effects. Call your doctor at once if you have: severe mouth or throat irritation

- breathing problems

- or choking feeling, chest pain, trouble swallowing. Common side effects of Grastek may include: swelling or itching of your mouth or tongue

- throat irritation

- or itching in your ears.

1736. This particular disease, Hayfever, interacts with the drug Histex CT and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blurred vision chills confusion cough decrease in the frequency of urination decrease in urine volume difficult or painful urination difficulty in passing urine (dribbling) difficulty with swallowing dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position fast, irregular, pounding, or racing heartbeat or pulse hives or welts, itching, skin rash puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue redness of the skin sweating tightness in the chest unusual tiredness or weakness Symptoms of overdose Agitation anxiety bloating diarrhea difficulty with sleeping disorientation dizziness drowsiness to profound coma dry mouth feeling of warmth fixed and enlarged pupils gas headache lethargy lightheadedness loss of appetite nausea redness of the face, neck, arms, and occasionally, upper chest seeing, hearing, or feeling things that are not there seizures tightness of the chest trouble breathing vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More

Drug Side Effects Summary

common Clumsiness or unsteadiness cough producing mucus drowsiness sleepiness Incidence not known Back, leg, or stomach pains black, tarry stools bleeding gums blood in the urine or stools blurred vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chills confusion continuing ringing or buzzing or other unexplained noise in the ears cough or hoarseness dark urine difficulty breathing difficulty having a bowel movement (stool) dizziness or lightheadedness double vision dry mouth, nose, and throat feeling excited feeling of constant movement of self or surroundings fever with or without chills general body swelling general feeling of tiredness or weakness headache hearing loss increased sensitivity of the skin to sunlight itching loss of appetite lower back or side pain nosebleeds pale skin pinpoint red spots on the skin redness or other discoloration of the skin restlessness seeing double seizures sensation of spinning severe sunburn shakiness in the legs, arms, hands, or feet sore throat sores, ulcers, or white spots on the lips or in the mouth trembling or shaking of the hands or feet trouble sleeping unusual bleeding or bruising weight loss yellowing of the eyes or skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Histex CT side effects

1737. This particular disease, Hayfever, interacts with the drug Histex IE and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blurred vision chills confusion cough decrease in the frequency of urination decrease in urine volume difficult or painful urination difficulty in passing urine (dribbling) difficulty with swallowing dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position fast, irregular, pounding, or racing heartbeat or pulse hives or welts, itching, skin rash puffiness or swelling of the eyelids or

Drug Side Effects Summary

around the eyes, face, lips, or tongue redness of the skin sweating tightness in the chest unusual tiredness or weakness Symptoms of overdose Agitation anxiety bloating diarrhea difficulty with sleeping disorientation dizziness drowsiness to profound coma dry mouth feeling of warmth fixed and enlarged pupils gas headache lethargy lightheadedness loss of appetite nausea redness of the face, neck, arms, and occasionally, upper chest seeing, hearing, or feeling things that are not there seizures tightness of the chest trouble breathing vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Clumsiness or unsteadiness cough producing mucus drowsiness sleepiness Incidence not known Back, leg, or stomach pains black, tarry stools bleeding gums blood in the urine or stools blurred vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chills confusion continuing ringing or buzzing or other unexplained noise in the ears cough or hoarseness dark urine difficulty breathing difficulty having a bowel movement (stool) dizziness or lightheadedness double vision dry mouth, nose, and throat feeling excited feeling of constant movement of self or surroundings fever with or without chills general body swelling general feeling of tiredness or weakness headache hearing loss increased sensitivity of the skin to sunlight itching loss of appetite lower back or side pain nosebleeds pale skin pinpoint red spots on the skin redness or other discoloration of the skin restlessness seeing double seizures sensation of spinning severe sunburn shakiness in the legs, arms, hands, or feet sore throat sores, ulcers, or white spots on the lips or in the mouth trembling or shaking of the hands or feet trouble sleeping unusual bleeding or bruising weight loss yellowing of the eyes or skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at

Drug Side Effects Summary

1-800-FDA-1088. Histex IE side effects

1738. This particular disease, Hayfever, interacts with the drug Histex PD and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Although serious side effects occur rarely when this medicine is taken as recommended, they may be more likely to occur if: too much medicine is taken, it is taken in large doses, or it is taken for a long period of time.

For narcotic antitussive (codeine, dihydrocodeine, hydrocodone, or hydromorphone)-containing If you are a nursing mother and you notice any of the following symptoms of overdose in your baby, get emergency help immediately : Cold, clammy skin confusion (severe) convulsions (seizures) drowsiness or dizziness (severe) nervousness or restlessness (severe) pinpoint pupils of eyes slow heartbeat slow or troubled breathing Difficulty breathing difficulty nursing increased sleepiness (more than usual) limpness For acetaminophen-containing Diarrhea increased sweating loss of appetite nausea or vomiting stomach cramps or pain swelling or tenderness in the upper abdomen or stomach area For salicylate-containing Any loss of hearing bloody urine confusion convulsions (seizures) dizziness or lightheadedness drowsiness (severe) excitement or nervousness (severe) fast or deep breathing fever hallucinations (seeing, hearing, or feeling things that are not there) increased sweating nausea or vomiting (severe or continuing) shortness of breath or troubled breathing (for salicylamide only) stomach pain (severe or continuing) uncontrollable flapping movements of the hands, especially in elderly patients unusual thirst vision problems For decongestant-containing Fast, pounding, or irregular heartbeat headache (continuing and severe) nausea or vomiting (severe) nervousness or restlessness (severe) shortness of breath or troubled breathing (severe or continuing) Check with your doctor as soon as possible if any of the following side effects occur: For all combinations Skin rash, hives, and/or itching For antihistamine- or anticholinergic-containing Clumsiness or unsteadiness convulsions (seizures) drowsiness (severe)

Drug Side Effects Summary

dryness of mouth, nose, or throat (severe) flushing or redness of face hallucinations (seeing, hearing, or feeling things that are not there) restlessness (severe) shortness of breath or troubled breathing slow or fast heartbeat For iodine-containing Headache (continuing) increased watering of mouth loss of appetite metallic taste skin rash, hives, or redness sore throat swelling of face, lips, or eyelids For acetaminophen-containing Unexplained sore throat and fever unusual tiredness or weakness yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Constipation decreased sweating difficult or painful urination dizziness or lightheadedness drowsiness dryness of mouth, nose, or throat false sense of well-being increased sensitivity of skin to sun nausea or vomiting nightmares stomach pain thickening of mucus trouble in sleeping unusual excitement, nervousness, restlessness, or irritability unusual tiredness or weakness Not all of the side effects listed above have been reported for each of these medicines, but they have been reported for at least one of them. There are some similarities among these combination medicines, so many of the above side effects may occur with any of these medicines. Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Histex PD side effects

1739. This particular disease, Hayfever, interacts with the drug house dust mite allergen extract and

- House dust mite allergen extract may cause a severe allergic reaction. Stop taking the medicine and get emergency medical help if you have any of these signs of an allergic reaction : severe dizziness or a light-headed feeling (like you might pass out)

Drug Side Effects Summary

- fast heart rate
- stomach pain, vomiting , diarrhea
- hives , itching, flushing (warmth, redness, or tingly feeling)
- cough, chest tightness, wheezing, trouble breathing
- or swelling of your face, lips, tongue, or throat. If you stop taking house dust mite allergen extract after having an allergic reaction, do not start taking the medicine again without your doctor's advice. House dust mite allergen extract may cause serious side effects. Call your doctor at once if you have: severe mouth or throat irritation
- breathing problems
- or choking feeling, chest pain, trouble swallowing. Common side effects of house dust mite allergen extract may include: pain, swelling, itching or sores in your mouth or on your tongue
- throat irritation
- altered sense of taste
- cough, sore throat , sinus pain
- nausea
- or itching in your ears.

1740. This particular disease, Hayfever, interacts with the drug Karbinal ER and has the following s

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Karbinal ER may cause serious side effects. Stop using Karbinal ER and call your doctor at once if you have: a light-headed feeling, like you might pass out
- little or no urination
- wheezing, tightness in your chest

Drug Side Effects Summary

- pounding heartbeats or fluttering in your chest
- or pale skin, easy bruising or bleeding. Side effects such as dizziness and confusion may be more likely in older adults. Common side effects of Karbinal ER may include: dry mouth, nose, or throat
- drowsiness, dizziness
- loss of coordination
- or upset stomach .

1741. This particular disease, Hayfever, interacts with the drug Leader Allergy Relief D-24 and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- or severe dizziness , nervousness , or restless feeling. Common side effects of Leader Allergy Relief D-24 may include: dry mouth, nose, or throat
- mild dizziness
- or sleep problems (insomnia).

1742. This particular disease, Hayfever, interacts with the drug Lodrane D and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Lodrane D may cause serious side effects. Stop using Lodrane D and call your doctor at once if you have: nervousness , dizziness , or sleeplessness
- chest pain, fast or uneven heart rate
- little or no urinating

Drug Side Effects Summary

- dangerously high blood pressure (severe headache , buzzing in your ears, anxiety , shortness of breath)
- if your symptoms do not improve after 7 days of treatment, or if you have a fever
- or if new symptoms occur. Less serious side effects may be more likely, and you may have none at all.

1743. This particular disease, Hayfever, interacts with the drug LoHist-D and has the following side

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate
- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)
- little or no urinating
- easy bruising or bleeding, unusual weakness
- or dangerously high blood pressure (severe headache, buzzing in your ears, anxiety , shortness of breath). Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of LoHist-D may include: dizziness , drowsiness, blurred vision
- dry nose or mouth
- nausea , stomach pain, constipation, loss of appetite
- problems with memory or concentration
- or feeling restless or excited (especially in children).

1744. This particular disease, Hayfever, interacts with the drug Loratadine-D 12 Hour and has the fo

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- or severe dizziness , nervousness , or restless feeling. Common side effects of Loratadine-D 12 Hour may include: dry mouth, nose, or throat
- mild dizziness
- or sleep problems (insomnia).

1745. This particular disease, Hayfever, interacts with the drug mixed grass pollens allergen extract

- Mixed grass pollens allergen extract may cause a severe allergic reaction. Stop taking the medicine and get emergency medical help if you have any of these signs of an allergic reaction : severe dizziness or a light-headed feeling (like you might pass out)
- fast heart rate
- stomach pain, vomiting, diarrhea
- hives, itching, flushing (warmth, redness, or tingly feeling)
- cough, chest tightness, wheezing, trouble breathing
- or swelling of your face, lips, tongue, or throat. If you stop taking mixed grass pollens allergen extract after having an allergic reaction, do not start taking the medicine again without your doctor's advice. Mixed grass pollens allergen extract may cause serious side effects. Call your doctor at once if you have: severe mouth or throat irritation
- breathing problems
- or choking feeling, chest pain, trouble swallowing. Common side effects of mixed grass pollens allergen extract may include: pain, swelling, or itching of your mouth or tongue

Drug Side Effects Summary

- throat irritation
- cough, sore throat , sinus pain
- or itching in your ears.

1746. This particular disease, Hayfever, interacts with the drug mometasone / olopatadine and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Bloody nose nose discomfort Rare Sleepiness or unusual drowsiness Incidence not known Blindness bloody mucus or unexplained nosebleeds blurred vision cough or hoarseness darkening of the skin decreased vision diarrhea difficulty in swallowing dizziness eye pain fainting fast heartbeat fever or chills headache loss of appetite lower back or side pain mental depression nausea or vomiting painful or difficult urination skin rash slow growth in children sore mouth or tongue sores inside the nose swelling of the face, throat, or tongue tearing trouble breathing unusual tiredness or weakness white patches in the mouth, tongue, or throat Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Change in taste loss of taste Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Mometasone / olopatadine nasal side effects

1747. This particular disease, Hayfever, interacts with the drug Nasonex 24HR Allergy and has the following side effects:

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe bleeding or increased drainage from your nose
- nose pain or discomfort, headache
- white patches or sores in the nose that won't heal
- wheezing, trouble breathing
- vision problems
- ear pain or full feeling, trouble hearing, drainage from the ear. Steroid medicines such as Nasonex 24HR Allergy can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using Nasonex 24HR Allergy. Although the risk of serious side effects is low when mometasone is used in the nose, side effects can occur if the medicine is absorbed into your bloodstream. Tell your doctor if you have possible signs of long-term steroid use: weight gain (especially in your face or your upper back and torso)
- slow wound healing, thinning skin, increased body hair
- irregular menstrual periods, changes in sexual function
- or muscle weakness, tired feeling, depression , anxiety , or feeling irritable. Common Nasonex 24HR Allergy side effects may include: nosebleeds
- headache
- stuffy nose, sore throat , cough
- or flu-like symptoms.

1748. This particular disease, Hayfever, interacts with the drug Odactra and has the following side effects:

- Odactra may cause a severe allergic reaction. Stop taking the medicine and get emergency medical help if you have any of these signs of an allergic reaction : severe dizziness or a

Drug Side Effects Summary

light-headed feeling (like you might pass out)

- fast heart rate
- stomach pain, vomiting , diarrhea
- hives , itching, flushing (warmth, redness, or tingly feeling)
- cough, chest tightness, wheezing, trouble breathing
- or swelling of your face, lips, tongue, or throat. If you stop taking Odactra after having an allergic reaction, do not start taking the medicine again without your doctor's advice. Odactra may cause serious side effects. Call your doctor at once if you have: severe mouth or throat irritation
- breathing problems
- or choking feeling, chest pain, trouble swallowing. Common side effects of Odactra may include: pain, swelling, itching or sores in your mouth or on your tongue
- throat irritation
- altered sense of taste
- cough, sore throat , sinus pain
- nausea
- or itching in your ears.

1749. This particular disease, Hayfever, interacts with the drug Oralair and has the following side effects

- Oralair may cause a severe allergic reaction. Stop taking the medicine and get emergency medical help if you have any of these signs of an allergic reaction : severe dizziness or a light-headed feeling (like you might pass out)
- fast heart rate
- stomach pain, vomiting, diarrhea
- hives, itching, flushing (warmth, redness, or tingly feeling)
- cough, chest tightness, wheezing, trouble breathing

Drug Side Effects Summary

- or swelling of your face, lips, tongue, or throat. If you stop taking Oralair after having an allergic reaction, do not start taking the medicine again without your doctor's advice. Oralair may cause serious side effects. Call your doctor at once if you have: severe mouth or throat irritation
- breathing problems
- or choking feeling, chest pain, trouble swallowing. Common side effects of Oralair may include: pain, swelling, or itching of your mouth or tongue
- throat irritation
- cough, sore throat , sinus pain
- or itching in your ears.

1750. This particular disease, Hayfever, interacts with the drug Polmon and has the following side effects:

- No side effects listed

1751. This particular disease, Hayfever, interacts with the drug ragweed pollen allergen extract and has the following side effects:

- Ragweed pollen allergen extract may cause a severe allergic reaction. Stop taking the medicine and get emergency medical help if you have any of these signs of an allergic reaction : severe dizziness or a light-headed feeling (like you might pass out)
- fast heart rate
- stomach pain, vomiting, diarrhea
- hives, itching, flushing (warmth, redness, or tingly feeling)
- cough, chest tightness, wheezing, trouble breathing
- or swelling of your face, lips, tongue, or throat. If you stop taking ragweed pollen allergen extract after having an allergic reaction, do not start taking the medicine again without your doctor's advice. Ragweed pollen allergen extract may cause serious side effects. Call your doctor at once if you have: severe mouth or throat irritation

Drug Side Effects Summary

- breathing problems
- or choking feeling, chest pain, trouble swallowing. Common side effects of ragweed pollen allergen extract may include: numbness , tingling, itching, or swelling in your mouth
- swollen tongue
- throat irritation
- or itching in your ears.

1752. This particular disease, Hayfever, interacts with the drug Ragwitek and has the following side effects:

- Ragwitek may cause a severe allergic reaction. Stop taking the medicine and get emergency medical help if you have any of these signs of an allergic reaction : severe dizziness or a light-headed feeling (like you might pass out)
- fast heart rate
- stomach pain, vomiting, diarrhea
- hives, itching, flushing (warmth, redness, or tingly feeling)
- cough, chest tightness, wheezing, trouble breathing
- or swelling of your face, lips, tongue, or throat. If you stop taking Ragwitek after having an allergic reaction, do not start taking the medicine again without your doctor's advice. Ragwitek may cause serious side effects. Call your doctor at once if you have: severe mouth or throat irritation
- breathing problems
- or choking feeling, chest pain, trouble swallowing. Common side effects of Ragwitek may include: numbness , tingling, itching, or swelling in your mouth
- swollen tongue
- throat irritation
- or itching in your ears.

Drug Side Effects Summary

1753. This particular disease, Hayfever, interacts with the drug Ru-Hist D and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: nervousness , dizziness , or sleeplessness
- chest pain, fast or uneven heart rate
- little or no urinating
- dangerously high blood pressure (severe headache , buzzing in your ears, anxiety , shortness of breath)
- if your symptoms do not improve after 7 days of treatment, or if you have a fever
- or if new symptoms occur. Less serious side effects may be more likely, and you may have none at all.

1754. This particular disease, Hayfever, interacts with the drug Ryaltris and has the following side effects:

- Ryaltris may cause serious side effects, including the following: nose and throat problems. Symptoms of nose and throat problems may include: nosebleeds sores (ulcers) in the nose hole in the cartilage between your nose (nasal septal perforation). Symptoms of nasal septal perforation may include: crusting in the nose nosebleeds runny nose whistling sound when you breathe slow wound healing. You should not use Ryaltris until your nose has healed if you have a sore in your nose, if you have had surgery on your nose, or if your nose has been injured. thrush (Candida), a certain fungal infection in your nose and throat. Tell your healthcare provider if you have any redness or white-colored patches in your nose or mouth. eye problems, including glaucoma or cataracts. You should have regular eye exams when using Ryaltris. allergic reactions. Call your healthcare provider or get emergency medical care if you get any of the following signs of a serious

Drug Side Effects Summary

allergic reaction: wheezing rash hives swelling of your face, mouth, and tongue breathing problems
immune system problems that may increase your risk of infections. Taking medicines that weaken your immune system makes you more likely to get infections. These infections may include tuberculosis (TB), ocular herpes simplex infections, and infections caused by fungi, bacteria, viruses, and parasites. Avoid contact with people who have contagious diseases, such as chickenpox or measles, while using Ryaltris. If you come in contact with someone who has chicken pox or measles call your healthcare provider right away. Symptoms of infection may include: fever aches or pains chills feeling tired adrenal insufficiency. Adrenal insufficiency happens when your adrenal glands do not make enough steroid hormones. Symptoms of adrenal insufficiency can include: tiredness weakness nausea vomiting low blood pressure slowed growth in children. A child's growth should be checked regularly while using Ryaltris. sleepiness or drowsiness. The most common side effects include the following: unpleasant taste nosebleeds nasal discomfort Tell your healthcare provider if you have any side effects that bother you or that do not go away. These are not all of the possible side effects. For more information, ask your healthcare provider or pharmacist. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Ryaltris side effects

1755. This particular disease, Hayfever, interacts with the drug RyClora and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. RyClora may cause serious side effects. Stop using RyClora and call your doctor at once if you have: extreme drowsiness
- confusion, hallucinations
- feeling nervous, restless, or dizzy
- a seizure

Drug Side Effects Summary

- or little or no urination. Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of RyClora may include: drowsiness

- blurred vision
- constipation
- flushing (warmth, redness, or tingly feeling)
- dry mouth, nose, or throat
- or feeling restless or excited (especially in a child).

1756. This particular disease, Hayfever, interacts with the drug Rymed and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe dizziness
- nervousness , trouble sleeping
- little or no urination
- or high blood pressure (severe headache, pounding in your neck or ears). Common side effects of Rymed may include: dizziness, drowsiness
- dry mouth, nose, or throat
- constipation
- blurred vision
- or feeling restless or excited (especially in children).

1757. This particular disease, Hayfever, interacts with the drug Rynex PE and has the following side effects:

- hives
- difficulty breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: nervousness , dizziness , or sleeplessness
- chest pain, fast or uneven heart rate
- little or no urinating
- dangerously high blood pressure (severe headache , buzzing in your ears, anxiety , shortness of breath)
- if your symptoms do not improve after 7 days of treatment, or if you have a fever
- or if new symptoms occur. Less serious side effects may be more likely, and you may have none at all.

1758. This particular disease, Hayfever, interacts with the drug Ryvent and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Ryvent may cause serious side effects. Stop using Ryvent and call your doctor at once if you have: a light-headed feeling, like you might pass out
- little or no urination
- wheezing, tightness in your chest
- pounding heartbeats or fluttering in your chest
- or pale skin, easy bruising or bleeding. Side effects such as dizziness and confusion may be more likely in older adults. Common side effects of Ryvent may include: dry mouth, nose, or throat
- drowsiness, dizziness
- loss of coordination
- or upset stomach .

Drug Side Effects Summary

1759. This particular disease, Hayfever, interacts with the drug Semprex-D and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Symptoms of overdose Blurred vision chest pain or discomfort cold, clammy, or pale skin confusion dizziness dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position irregular heartbeats seizures shortness of breath slow heart rate sweating unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Dry mouth headache sleepiness or unusual drowsiness Less common Acid or sour stomach belching body aches or pain congestion cough dryness or soreness of the throat fever heartburn hoarseness indigestion lack or loss of strength nausea nervousness runny nose stomach discomfort, upset, or pain tender, swollen glands in the neck trouble sleeping trouble swallowing voice changes Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Semprex-D side effects

1760. This particular disease, Hayfever, interacts with the drug SudoGest Sinus & Allergy and has the following side effects:

- hives

- difficulty breathing

- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: chest pain, rapid pulse, fast or uneven heart rate

Drug Side Effects Summary

- confusion, hallucinations, severe nervousness
- tremor, seizure (convulsions)
- little or no urinating
- easy bruising or bleeding, unusual weakness
- or dangerously high blood pressure (severe headache, buzzing in your ears, anxiety , shortness of breath). Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of SudoGest Sinus & Allergy may include: dizziness , drowsiness, blurred vision
- dry nose or mouth
- nausea , stomach pain, constipation, loss of appetite
- problems with memory or concentration
- or feeling restless or excited (especially in children).

1761. This particular disease, Hayfever, interacts with the drug timothy grass pollen allergen extract

- Timothy grass pollen allergen extract may cause a severe allergic reaction. Stop taking the medicine and get emergency medical help if you have any of these signs of an allergic reaction : severe dizziness or a light-headed feeling (like you might pass out)
- fast heart rate
- stomach pain, vomiting, diarrhea
- hives, itching, flushing (warmth, redness, or tingly feeling)
- cough, chest tightness, wheezing, trouble breathing
- or swelling of your face, lips, tongue, or throat. If you stop taking timothy grass pollen allergen extract after having an allergic reaction, do not start taking the medicine again without your doctor's advice. Timothy grass pollen allergen extract may cause serious side effects. Call your doctor at once if you have: severe mouth or throat irritation

Drug Side Effects Summary

- breathing problems
- or choking feeling, chest pain, trouble swallowing. Common side effects of timothy grass pollen allergen extract may include: swelling or itching of your mouth or tongue
- throat irritation
- or itching in your ears.

1762. This particular disease, Hayfever, interacts with the drug Zetonna and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Nasal steroid medicine can be absorbed into your bloodstream, which may cause steroid side effects throughout the body. Tell your doctor if you have: weight gain (especially in your face or your upper back and torso)
- slow wound healing, thinning skin, increased body hair
- irregular menstrual periods, changes in sexual function
- or muscle weakness, tired feeling, depression, anxiety, feeling irritable. Call your doctor at once if you have: runny or bloody nose, crusting inside the nose, whistling sound when you breathe
- sores or white patches in or around your nostrils
- fever, chills, body aches, nausea, vomiting
- blurred vision, tunnel vision, eye pain, seeing halos around lights
- or signs of a hormonal disorder--worsening tiredness or weakness, feeling light-headed, vomiting, loss of appetite, diarrhea, and weight loss. Common Zetonna side effects may include: headache, dizziness
- nosebleed, nasal irritation
- stuffy nose, sore throat
- or ear pain.

Drug Side Effects Summary

1763. This particular disease, Herpes, interacts with the drug acyclovir and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: easy bruising or bleeding, purple or red pinpoint spots under your skin
- changes in behavior
- confusion, hallucinations
- or signs of a kidney problem - little or no urinating
- painful or difficult urination
- swelling in your feet or ankles
- feeling tired or short of breath. Common acyclovir side effects may include: mild skin pain
- rash, itching
- nausea , vomiting , diarrhea
- headache
- or mouth pain while using an acyclovir buccal tablet.

1764. This particular disease, Herpes, interacts with the drug valacyclovir and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: confusion, aggression, or you feel shaky or unsteady
- hallucinations (seeing or hearing things that are not real)
- problems with speech
- a seizure (convulsions)
- or kidney problems--little or no urination, painful or difficult urination, swelling in your feet or ankles,

Drug Side Effects Summary

feeling tired or short of breath. Stop taking valacyclovir and call your doctor right away if you have any of the following signs of a serious side effect that can harm red blood cells: fever, pale skin

- unusual bleeding (nosebleeds, bleeding gums)
- red or pink urine, little or no urination
- red spots on the skin (not related to herpes or chickenpox)
- feeling weak or tired
- stomach pain, bloody diarrhea, vomiting
- or swelling in your face, hands, or feet. Side effects may be more likely in adults who are 65 or older. Common valacyclovir side effects may include: nausea, stomach pain
- or headache.

1765. This particular disease, Herpes, interacts with the drug Zovirax Cream and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common side effects of Zovirax Topical may include: dry, cracked, or peeling lips
- dryness or flaking of treated skin
- burning, stinging, or itching when the medicine is applied.

1766. This particular disease, Herpes, interacts with the drug Abreva and has the following side effects:

- hives
- feeling light-headed
- wheezing, difficult breathing
- swelling of your face, lips, tongue, or throat. Less serious side effects may be more likely, and you may have none at all.

Drug Side Effects Summary

1767. This particular disease, Herpes, interacts with the drug acyclovir and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common side effects of acyclovir topical may include:
dry, cracked, or peeling lips
- dryness or flaking of treated skin
- burning, stinging, or itching when the medicine is applied.

1768. This particular disease, Herpes, interacts with the drug Zovirax Ointment and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Mild pain, burning, or stinging Less common Itching Rare Itching, stinging, or redness of the genital area skin rash Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Zovirax Ointment side effects

1769. This particular disease, Herpes, interacts with the drug Zovirax and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: easy bruising or

Drug Side Effects Summary

bleeding, purple or red pinpoint spots under your skin

- changes in behavior
- confusion, hallucinations
- or signs of a kidney problem - little or no urinating
- painful or difficult urination
- swelling in your feet or ankles
- feeling tired or short of breath. Common Zovirax side effects may include: mild skin pain
- rash, itching
- nausea , vomiting , diarrhea
- headache
- or

1770. This particular disease, Herpes, interacts with the drug docosanol and has the following side effects:

- hives
- feeling light-headed
- wheezing, difficult breathing
- swelling of your face, lips, tongue, or throat. Less serious side effects may be more likely, and you may have none at all.

1771. This particular disease, Herpes, interacts with the drug Blistex Ointment and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have severe burning, stinging, redness, or other irritation after applying the ointment. Common side effects of Blistex may include: mild tingling or cool feeling after applying the medicine

Drug Side Effects Summary

- or a white residue on your lips.

1772. This particular disease, Herpes, interacts with the drug allantoin / camphor / phenol and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have severe burning, stinging, redness, or other irritation after applying the ointment. Common side effects of allantoin, camphor, and phenol may include: mild tingling or cool feeling after applying the medicine
- or a white residue on your lips.

1773. This particular disease, Hypertension, interacts with the drug lisinopril and has the following

- hives
- severe stomach pain, difficult breathing
- swelling of your face, lips, tongue, or throat. You may be more likely to have an allergic reaction if you are African-American. Lisinopril may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- fever, sore throat
- high potassium - nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or >liver problems - nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of lisinopril may include: headache

Drug Side Effects Summary

- dizziness
- cough
- or chest pain.

1774. This particular disease, Hypertension, interacts with the drug losartan and has the following s

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In extremely rare cases, losartan can cause a condition that results in the breakdown of skeletal muscle tissue, leading to potential kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, and dark colored urine. Call your doctor at once if you have: a feeling that you might pass out
- pain or burning when you urinate
- pale skin, feeling light-headed or short of breath, rapid heart rate, trouble concentrating
- wheezing, chest pain
- drowsiness, confusion, mood changes, increased thirst, loss of appetite, nausea and vomiting
- swelling, weight gain, feeling short of breath, urinating less than usual or not at all
- or high potassium (slow heart rate, weak pulse, muscle weakness, tingly feeling). Common losartan side effects may include: cold or flu symptoms such as stuffy nose, sneezing, sore throat, fever
- dry cough
- muscle cramps
- pain in your legs or back
- stomach pain, diarrhea
- headache, dizziness

Drug Side Effects Summary

- tired feeling
- or sleep problems (insomnia).

1775. This particular disease, Hypertension, interacts with the drug hydrochlorothiazide and has the following side effects:

- a light-headed feeling
- eye pain , vision problems
- jaundice (yellowing of the skin or eyes)
- pale skin, easy bruising, unusual bleeding (nose, mouth, vagina, or rectum)
- shortness of breath, wheezing, cough with foamy mucus, chest pain
- dehydration symptoms - feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin
- or signs of an electrolyte imbalance - increased thirst or urination, confusion, vomiting, constipation , muscle pain, leg cramps, bone pain, lack of energy, irregular heartbeats, tingly feeling. Common hydrochlorothiazide side effects may include: weakness
- feeling like you might pass out
- severe pain in your upper stomach spreading to your back, nausea and vomiting
- fever, chills, tiredness, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath
- or electrolyte imbalance.

1776. This particular disease, Hypertension, interacts with the drug Benicar and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: a light-headed feeling, like you might pass out

Drug Side Effects Summary

- little or no urination
- severe diarrhea and weight loss
- or high potassium level - nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common Benicar side effects may include: dizziness .

1777. This particular disease, Hypertension, interacts with the drug furosemide and has the following side effects:

- a light-headed feeling, like you might pass out
- ringing in your ears, hearing loss
- muscle spasms or contractions
- pale skin, easy bruising, unusual bleeding
- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- signs of liver or pancreas problems - loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting , dark urine, jaundice (yellowing of the skin or eyes)
- or signs of an electrolyte imbalance - dry mouth, thirst, weakness, drowsiness, feeling jittery or unsteady, vomiting, irregular heartbeats, fluttering in your chest, numbness or tingling, muscle cramps, muscle weakness or limp feeling. Common furosemide side effects may include: diarrhea , constipation , loss of appetite
- numbness or tingling
- headache , dizziness
- or blurred vision.

1778. This particular disease, Hypertension, interacts with the drug Cozaar and has the following side effects:

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- pain or burning when you urinate
- high potassium level - nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- or kidney problems - little or no urination, rapid weight gain, painful or difficult urination, swelling in your hands, feet, or ankles. Common Cozaar side effects may include: dizziness
- back pain
- or cold symptoms such as stuffy nose, sneezing, sore throat .

1779. This particular disease, Hypertension, interacts with the drug hydrochlorothiazide / lisinopril

- hives
- severe stomach pain
- difficulty breathing
- swelling of your face, lips, tongue, or throat. You may be more likely to have an allergic reaction if you are African-American. Hydrochlorothiazide and lisinopril may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- eye pain , vision problems
- little or no urination
- weakness, drowsiness, or feeling restless
- fever, chills, sore throat , mouth sores, trouble swallowing
- jaundice (yellowing of the skin or eyes)
- high potassium-- nausea , tingly feeling, chest pain, irregular heartbeats, loss of movement
- low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased

Drug Side Effects Summary

thirst or urination, numbness or tingling, muscle weakness or limp feeling

- or low sodium-- headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady. Common side effects of hydrochlorothiazide and lisinopril may include: cough

- headache

- dizziness

- or tired feeling.

1780. This particular disease, Hypertension, interacts with the drug Bystolic and has the following side effects:

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have a serious side effect such as: feeling short of breath, even with mild exertion

- swelling of your ankles or feet

- slow or uneven heartbeats

- or numbness or cold feeling in your hands and feet. Less serious Bystolic side effects may include: headache

- tired feeling

- nausea, stomach pain

- diarrhea

- or sleep problems (insomnia).

1781. This particular disease, Hypertension, interacts with the drug Diovan and has the following side effects:

- hives

- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- little or no urination
- or high potassium level - nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common Diovan side effects may include: high potassium
- headache , dizziness , feeling light-headed
- flu symptoms, tiredness
- cough
- stomach pain, diarrhea
- back pain , joint pain
- or abnormal kidney test.

1782. This particular disease, Hypertension, interacts with the drug valsartan and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Valsartan may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- little or no urination
- or high potassium level-- nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common side effects of valsartan may include: high potassium
- headache , dizziness , feeling light-headed
- flu symptoms, tiredness
- cough
- stomach pain, diarrhea
- back pain , joint pain

Drug Side Effects Summary

- or abnormal kidney test.

1783. This particular disease, Hypertension, interacts with the drug Avapro and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- little or no urination
- high potassium level - nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common Avapro side effects may include: dizziness
- feeling light-headed
- or high potassium.

1784. This particular disease, Hypertension, interacts with the drug enalapril and has the following side effects:

- hives
- severe stomach pain
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- slow heartbeats
- pounding heartbeats or fluttering in your chest
- little or no urination
- jaundice (yellowing of the skin or eyes)
- numbness, tingling, or burning pain in your hands or feet
- fever, chills, cold or flu symptoms

Drug Side Effects Summary

- easy bruising, unusual bleeding
- or high potassium - nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common enalapril side effects may include: dizziness, feeling tired
- cough
- or feeling light-headed.

1785. This particular disease, Hypertension, interacts with the drug hydralazine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Hydralazine may cause serious side effects. Call your doctor at once if you have: chest pain or pressure, pain spreading to your jaw or shoulder
- fast or pounding heartbeats
- a light-headed feeling, like you might pass out
- numbness , tingling, or burning pain in your hands or feet
- painful or difficult urination
- little or no urination
- or lupus -like syndrome--joint pain or swelling with fever, swollen glands, muscle aches, chest pain, vomiting , unusual thoughts or behavior, and patchy skin color. Common side effects of hydralazine may include: chest pain, fast heart rate
- headache
- or nausea , vomiting, diarrhea , loss of appetite.

1786. This particular disease, Hypertension, interacts with the drug Lasix and has the following side effects:

- hives, difficult breathing, swelling in your face or throat) or a severe skin reaction (fever, sore throat, burning in your eyes, skin pain, red or purple skin rash that spreads and causes blistering

Drug Side Effects Summary

and peeling). Call your doctor at once if you have: ringing in your ears, hearing loss

- easy bruising, unusual bleeding
- sudden weakness or ill feeling, fever, chills
- painful or difficult urination
- numbness, tingling, or burning pain
- a light-headed feeling, like you might pass out
- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- signs of liver or pancreas problems - loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting, dark urine, jaundice (yellowing of the skin or eyes)
- kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or signs of an electrolyte imbalance - dry mouth, increased thirst, mood changes, weakness, drowsiness, lack of energy, muscle pain, fast heartbeats, feeling restless, nausea, vomiting.

Common Lasix side effects may include: diarrhea, constipation

- numbness or tingling
- headache, dizziness
- or blurred vision.

1787. This particular disease, Hypertension, interacts with the drug nifedipine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: worsening angina
- a light-headed feeling, like you might pass out
- pounding heartbeats or fluttering in your chest
- chest pain or heavy feeling, pain spreading to the jaw or shoulder, nausea, sweating, general ill

Drug Side Effects Summary

feeling

- swelling in your ankles or feet
- or upper stomach pain, jaundice (yellowing of the skin or eyes). Common nifedipine side effects may include: mild dizziness
- flushing (warmth, redness, or tingly feeling)
- weakness, headache, mood changes
- heartburn, nausea
- tremors, muscle cramps
- or cough, wheezing, sore throat, stuffy nose.

1788. This particular disease, Hypertension, interacts with the drug Prinivil and has the following side effects:

- hives
- severe stomach pain, difficult breathing
- swelling of your face, lips, tongue, or throat. You may be more likely to have an allergic reaction if you are African-American. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- fever, sore throat
- high potassium - nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or liver problems - nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common Prinivil side effects may include: headache, dizziness
- cough

Drug Side Effects Summary

- or chest pain.

1789. This particular disease, Hypertension, interacts with the drug ramipril and has the following side effects:

- hives
- severe stomach pain
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- jaundice (yellowing of your skin or eyes)
- little or no urination
- fever, chills, sore throat, cold or flu symptoms
- or high potassium - nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common ramipril side effects may include: headache
- cough
- or dizziness, weakness, tired feeling.

1790. This particular disease, Hypertension, interacts with the drug Zestril and has the following side effects:

- hives
- severe stomach pain, difficult breathing
- swelling of your face, lips, tongue, or throat. You may be more likely to have an allergic reaction if you are African-American. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- fever, sore throat
- high potassium - nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement

Drug Side Effects Summary

- kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or liver problems - nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common Zestril side effects may include: headache, dizziness
- cough
- or chest pain.

1791. This particular disease, Hypertension, interacts with the drug diltiazem and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: slow heartbeats
- pounding heartbeats or fluttering in your chest
- a light-headed feeling, like you might pass out
- shortness of breath (even with mild activity), swelling, rapid weight gain
- nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or severe skin reaction - fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common diltiazem side effects may include: swelling
- dizziness, weakness
- headache
- nausea
- or rash.

Drug Side Effects Summary

1792. This particular disease, Hypertension, interacts with the drug hydrochlorothiazide / losartan and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Hydrochlorothiazide and losartan may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- irregular heartbeats
- eye pain , vision problems
- kidney problems--little or no urination, rapid weight gain, swelling in your hands, feet, or ankles
- new or worsening symptoms of lupus--joint pain, and a skin rash on your cheeks or arms that worsens in sunlight
- or signs of an electrolyte imbalance--increased thirst or urination, confusion, vomiting, constipation , muscle pain, leg cramps, irregular heartbeats, numbness or tingling, severe weakness, loss of movement, feeling jittery or unsteady. Common side effects of hydrochlorothiazide and losartan may include: dizziness
- back pain
- or cold symptoms such as stuffy nose, sneezing, sore throat .

1793. This particular disease, Hypertension, interacts with the drug irbesartan and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Irbesartan may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- little or no urination
- high potassium level-- nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common side effects of irbesartan may include: dizziness

Drug Side Effects Summary

- feeling light-headed
- or high potassium.

1794. This particular disease, Hypertension, interacts with the drug olmesartan and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Olmesartan may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- little or no urination
- severe diarrhea and weight loss
- or high potassium level-- nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common side effects of olmesartan may include: dizziness .

1795. This particular disease, Hypertension, interacts with the drug Vasotec and has the following s

- hives
- severe stomach pain
- difficulty breathing
- swelling of your face, lips, tongue, or throat. You may be more likely to have an allergic reaction if you are African-American. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- chest pain
- jaundice (yellowing of the skin or eyes)
- little or no urination
- fever, chills, sore throat
- or high potassium - nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of

Drug Side Effects Summary

movement. Common Vasotec side effects may include: dizziness

- tiredness
- or feeling light-headed.

1796. This particular disease, Hypertension, interacts with the drug chlorthalidone and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Chlorthalidone may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- low sodium-- headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady
- low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- low magnesium--dizziness, irregular heartbeats, feeling jittery, muscle cramps, muscle spasms, cough or choking feeling
- or kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath. Common side effects of chlorthalidone may include: low blood pressure (feeling light-headed)
- kidney problems
- dizziness
- or an electrolyte imbalance (such as low levels of potassium, sodium, or magnesium in your blood).

1797. This particular disease, Hypertension, interacts with the drug hydrochlorothiazide / triamterene and has the following side effects:

- blurred vision, tunnel vision, eye pain , or seeing halos around lights

Drug Side Effects Summary

- a light-headed feeling, like you might pass out
- fast, slow, or uneven heart rate
- signs of liver or pancreas problems--loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting, fast heart rate , dark urine, jaundice (yellowing of the skin or eyes)
- high potassium level--nausea, weakness, tingly feeling, feeling weak or tired, slow heartbeats, loss of movement
- low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- lupus -like symptoms--muscle or joint pain, flu symptoms, chest pain, and a rash or patchy skin color that worsens in sunlight
- or other signs of an electrolyte imbalance--thirst, dry mouth, stomach pain, drowsiness, weakness, fast heart rate, muscle pain or weakness, feeling restless or light-headed. Common side effects of hydrochlorothiazide and triamterene may include: stomach pain, nausea, diarrhea , constipation
- dizziness , headache
- blurred vision
- or dry mouth.

1798. This particular disease, Hypertension, interacts with the drug amlodipine / benazepril and has

- hives
- severe stomach pain
- difficulty breathing
- swelling of your face, lips, tongue, or throat. You may be more likely to have an allergic reaction if you are African-American. Some side effects may not occur until after you have used the medicine for several months. Amlodipine and benazepril may cause serious side effects. Call your doctor at

Drug Side Effects Summary

once if you have: a light-headed feeling, like you might pass out

- swelling in your hands or feet, rapid weight gain
- new or worsened chest pain
- fever, chills, sore throat , body aches, flu symptoms
- high potassium-- nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- or liver problems--nausea, stomach pain (upper right side), itching, unusual tiredness, flu-like symptoms, dark urine, jaundice (yellowing of the skin or eyes). Common side effects of amlodipine and benazepril may include: cough
- dizziness
- or swelling in your hands or feet.

1799. This particular disease, Hypertension, interacts with the drug benazepril and has the following

- a light-headed feeling, like you might pass out
- little or no urinating
- high potassium-- nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- or liver problems--stomach pain (upper right side), loss of appetite, dark urine, jaundice (yellowing of the skin or eyes). Common side effects of benazepril may include: headache
- or cough.

1800. This particular disease, Hypertension, interacts with the drug bisoprolol and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: shortness of

Drug Side Effects Summary

breath (even with mild exertion), swelling, rapid weight gain

- slow heart rate
- pounding heartbeats or fluttering in your chest
- numbness, tingling, or cold feeling in your hands or feet
- a light-headed feeling, like you might pass out
- eye pain, vision problems
- or bronchospasm (wheezing, chest tightness, trouble breathing). Common bisoprolol side effects may include: headache
- feeling tired
- sleep problems (insomnia)
- joint pain
- swelling
- or cold symptoms such as stuffy nose, runny nose, cough, sore throat.

1801. This particular disease, Hypertension, interacts with the drug Diovan HCT and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, Diovan HCT can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure . Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark colored urine. Diovan HCT may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- eye pain , vision problems
- an unusual skin rash
- little or no urinating

Drug Side Effects Summary

- jaundice (yellowing of the skin or eyes)
- easy bruising, unusual bleeding
- a seizure
- kidney problems--little or no urination, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- or signs of an electrolyte imbalance--dry mouth, increased thirst, confusion, feeling restless, vomiting, muscle pain or weakness, drowsiness, lack of energy, fast heartbeats, decreased urination. Common side effects include: cold symptoms such as stuffy nose, sneezing, sore throat
- dizziness
- or headache .

1802. This particular disease, Hypertension, interacts with the drug doxazosin and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- severe ongoing stomach pain or bloating
- new or worsening chest pain
- trouble breathing
- or a penis erection that is painful or lasts 4 hours or longer. Common doxazosin side effects may include: low blood pressure, dizziness
- drowsiness
- headache
- or feeling weak or tired.

Drug Side Effects Summary

1803. This particular disease, Hypertension, interacts with the drug Maxzide and has the following side effects:

- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- a light-headed feeling, like you might pass out
- fast, slow, or uneven heart rate
- signs of liver or pancreas problems--loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting, fast heart rate , dark urine, jaundice (yellowing of the skin or eyes)
- high potassium level--nausea, weakness, tingly feeling, feeling weak or tired, slow heartbeats, loss of movement
- low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- lupus -like symptoms--muscle or joint pain, flu symptoms, chest pain, and a rash or patchy skin color that worsens in sunlight
- or other signs of an electrolyte imbalance--thirst, dry mouth, stomach pain, drowsiness, weakness, fast heart rate, muscle pain or weakness, feeling restless or light-headed. Common side effects of Maxzide may include: stomach pain, nausea, diarrhea , constipation
- dizziness , headache
- blurred vision
- or dry mouth.

1804. This particular disease, Hypertension, interacts with the drug Benicar HCT and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, this medicine can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure . Call your doctor right

Drug Side Effects Summary

away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark colored urine. Also call your doctor at once if you have: a light-headed feeling, like you might pass out

- eye pain , vision problems
- an unusual skin rash
- easy bruising, unusual bleeding
- jaundice (yellowing of the skin or eyes)
- kidney problems--little or no urination, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- low levels of sodium in the body-- headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady
- high potassium--slow or unusual heart rate, weakness, loss of movement
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects include: nausea
- dizziness
- cold symptoms such as stuffy nose, sneezing, sore throat
- or high levels of uric acid in your blood.

1805. This particular disease, Hypertension, interacts with the drug candesartan and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Candesartan may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- little or no urination

Drug Side Effects Summary

- or high potassium level-- nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common side effects of candesartan may include: high potassium
- headache , back pain
- cold symptoms such as stuffy or runny nose , sneezing, sore throat
- dizziness
- or abnormal kidney test.

1806. This particular disease, Hypertension, interacts with the drug Cartia XT and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Body aches or pain congestion cough dryness or soreness of the throat fever hoarseness runny nose tender or swollen glands in the neck trouble swallowing voice changes Less common Chest pain, discomfort, or tightness chills diarrhea difficult or labored breathing feeling faint, dizzy, or lightheaded feeling of warmth or heat flushing or redness of the skin, especially on the face and neck general feeling of discomfort or illness headache joint pain loss of appetite muscle aches and pains nausea shivering slow or irregular heartbeat sweating swelling of the hands, ankles, feet, or lower legs trouble sleeping unusual tiredness or weakness vomiting Incidence not known Blistering, peeling, or loosening of the skin itching large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs no heartbeat red skin lesions, often with a purple center red, irritated eyes sores, ulcers, or white spots in the mouth or on the lips Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common

Drug Side Effects Summary

Sneezing stuffy nose Less common Acid or sour stomach belching constipation continuing ringing or buzzing or other unexplained noise in the ears degenerative disease of the joint difficulty with moving hearing loss heartburn indigestion lack or loss of strength muscle aching or cramping muscle pains or stiffness pain or tenderness around the eyes and cheekbones rash stomach discomfort, upset, or pain swollen joints Incidence not known Loss or thinning of the hair Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Cartia XT side effects

1807. This particular disease, Hypertension, interacts with the drug Exforge and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- swelling in your hands or feet, rapid weight gain
- or signs of an electrolyte imbalance - dry mouth, increased thirst, drowsiness, confusion, feeling restless, vomiting, muscle pain or weakness, lack of energy, fast heartbeats, little or no urine, or a seizure. Common Exforge side effects include: swelling in your hands or feet
- headache and dizziness
- or cold symptoms such as stuffy nose, sneezing, sore throat.

1808. This particular disease, Hypertension, interacts with the drug labetalol and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Labetalol may cause serious side effects. Call your

Drug Side Effects Summary

doctor at once if you have: a light-headed feeling, like you might pass out

- slow heart rate, weak pulse, fainting, slow breathing (breathing may stop)
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- severe headache , blurred vision, pounding in your neck or ears
- or liver problems--loss of appetite, stomach pain (upper right side), flu-like symptoms, itching, dark urine, jaundice (yellowing of the skin or eyes). Severe dizziness or fainting may be more likely in older adults. Common side effects of labetalol may include: dizziness, drowsiness, tiredness
- nausea , vomiting
- sudden warmth, skin redness, sweating
- numbness
- or tingly feeling in your scalp.

1809. This particular disease, Hypertension, interacts with the drug perindopril and has the following

- hives
- severe stomach pain
- difficulty breathing
- swelling of your face, lips, tongue, or throat. You may be more likely to have an allergic reaction if you are African-American. Perindopril may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- fever, chills, body aches, flu symptoms, sores in your mouth and throat
- little or no urination
- swelling, rapid weight gain
- high potassium-- nausea , slow or unusual heart rate, weakness, loss of movement
- pale skin, easy bruising or bleeding
- or jaundice (yellowing of the skin or eyes). Common side effects of perindopril may include:

Drug Side Effects Summary

dizziness

- back pain

- or cough.

1810. This particular disease, Hypertension, interacts with the drug Qbrelis and has the following side effects:

- hives

- severe stomach pain

- difficulty breathing

- swelling of your face, lips, tongue, or throat. You may be more likely to have an allergic reaction if you are African-American. Qbrelis may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out

- fever, sore throat

- high potassium-- nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement

- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath

- or liver problems--nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Qbrelis may include: headache , dizziness

- cough

- or chest pain.

1811. This particular disease, Hypertension, interacts with the drug telmisartan and has the following side effects:

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Also call your doctor at once if you have: a

Drug Side Effects Summary

light-headed feeling, like you might pass out

- little or no urination
- swelling in your feet or ankles, rapid weight gain
- unusual pain or tightness in your lower body
- a skin ulcer
- or high potassium level-- nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common side effects of telmisartan may include: stuffy nose, sinus pain
- back pain
- or diarrhea.

1812. This particular disease, Hypertension, interacts with the drug Altace and has the following side effects:

- hives
- severe stomach pain
- difficulty breathing
- swelling of your face, lips, tongue, or throat. You may be more likely to have an allergic reaction if you are African-American. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- jaundice (yellowing of your skin or eyes)
- little or no urination
- fever, chills, sore throat
- or high potassium - nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common Altace side effects may include: headache
- cough
- or dizziness , weakness, tired feeling.

Drug Side Effects Summary

1813. This particular disease, Hypertension, interacts with the drug amlodipine / valsartan and has

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Amlodipine and valsartan may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- swelling in your hands or feet, rapid weight gain
- or high potassium level-- nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common side effects include: swelling in your hands or feet
- dizziness
- or cold symptoms such as stuffy nose, sneezing, sore throat .

1814. This particular disease, Hypertension, interacts with the drug Aquazide H and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Back, leg, or stomach pains black, tarry stools bleeding gums blistering, peeling, or loosening of the skin bloating blood in the urine or stools blue lips and fingernails blurred vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain or tightness chills clay-colored stools cloudy urine cold sweats confusion constipation cough or hoarseness coughing that sometimes produces a pink frothy sputum coughing up blood cracks in the skin darkened urine decrease in urine-concentrating ability decreased frequency or amount of urine diarrhea difficult, fast, or noisy breathing difficulty with swallowing dizziness, faintness, or lightheadedness when getting up from a lying or sitting position dry mouth fast or irregular heartbeat fever flushed, dry skin fruit-like breath odor general body swelling general feeling of discomfort or illness greatly decreased frequency of urination or amount of urine headache hives, itching, rash increased hunger increased

Drug Side Effects Summary

sweating increased thirst increased urination indigestion joint pain, stiffness, or swelling loss of appetite loss of heat from the body lower back or side pain muscle cramps or pain nausea or vomiting nosebleeds numbness, tingling, pain, or weakness in the hands or feet painful or difficult urination pains in the stomach or side, possibly radiating to the back pale skin persistent non-healing sore pink growth pinpoint red spots on the skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red, irritated eyes red skin lesions, often with a purple center red, swollen skin reddish patch or irritated area redness, soreness, or scaly skin seizures shiny bump sore throat sores, ulcers, or white spots on the lips or in the mouth sores, welting, or blisters swelling of the face, fingers, legs, ankles, feet, or lower legs swollen or painful glands tenderness of salivary glands thickening of bronchial secretions trembling trouble breathing unpleasant breath odor unusual bleeding or bruising unusual tiredness or weakness unusual weight loss vomiting of blood weakness and heaviness of the legs weight gain white, yellow or waxy scar-like area yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Cramping decreased interest in sexual intercourse difficulty having a bowel movement (stool) feeling of constant movement of self or surroundings hair loss or thinning of the hair inability to have or keep an erection increased sensitivity of the skin to sunlight loss in sexual ability, desire, drive, or performance muscle spasm redness or other discoloration of the skin restlessness sensation of spinning severe sunburn weakness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Aquazide H side effects

Drug Side Effects Summary

1815. This particular disease, Hypertension, interacts with the drug atenolol / chlorthalidone and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Atenolol and chlorthalidone may cause serious side effects. Call your doctor at once if you have: new or worsening chest pain
- slow or uneven heartbeats
- a light-headed feeling, like you might pass out
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- cold feeling in your hands and feet
- jaundice (yellowing of the skin or eyes)
- low sodium level -- headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady
- or low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of atenolol and chlorthalidone may include: slow heartbeats
- feeling weak or tired
- dizziness , drowsiness
- numbness or tingling
- pale skin, bruising or bleeding
- depression
- or upset stomach , loss of appetite, constipation.

1816. This particular disease, Hypertension, interacts with the drug Azor and has the following side effects:

- hives
- difficulty breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. When you first start taking this medicine, you may have new or worsening chest pain (angina), or you could have a heart attack. Seek emergency medical attention or call your doctor right away if you have symptoms such as: chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating. Call your doctor at once if you have: new or worsening chest pain
- swelling in your hands or feet, rapid weight gain
- severe or ongoing diarrhea with weight loss
- pounding heartbeats or fluttering in your chest
- a light-headed feeling, like you might pass out
- or high potassium level - nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common Azor side effects include: swelling
- dizziness
- flushing (warmth, redness, or tingly feeling)
- or pounding heartbeats.

1817. This particular disease, Hypertension, interacts with the drug captopril and has the following

- hives
- severe stomach pain
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- little or no urination, or urinating more than usual
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- chest pain or pressure, pounding heartbeats or fluttering in your chest
- high potassium - nausea, slow or unusual heart rate, weakness, loss of movement

Drug Side Effects Summary

- or sudden weakness or ill feeling, fever, chills, sore throat, painful mouth sores, pain when swallowing, skin sores, cold or flu symptoms. Common captopril side effects may include: cough
- flushing (warmth, redness, or tingly feeling)
- numbness, tingling, or burning pain in your hands or feet
- loss of taste sensation
- or mild skin itching or rash.

1818. This particular disease, Hypertension, interacts with the drug Cardizem CD and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Body aches or pain congestion cough dryness or soreness of the throat fever hoarseness runny nose tender or swollen glands in the neck trouble swallowing voice changes Less common Chest pain, discomfort, or tightness chills diarrhea difficult or labored breathing feeling faint, dizzy, or lightheaded feeling of warmth or heat flushing or redness of the skin, especially on the face and neck general feeling of discomfort or illness headache joint pain loss of appetite muscle aches and pains nausea shivering slow or irregular heartbeat sweating swelling of the hands, ankles, feet, or lower legs trouble sleeping unusual tiredness or weakness vomiting Incidence not known Blistering, peeling, or loosening of the skin itching large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs no heartbeat red skin lesions, often with a purple center red, irritated eyes sores, ulcers, or white spots in the mouth or on the lips Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common

Drug Side Effects Summary

Sneezing stuffy nose Less common Acid or sour stomach belching constipation continuing ringing or buzzing or other unexplained noise in the ears degenerative disease of the joint difficulty with moving hearing loss heartburn indigestion lack or loss of strength muscle aching or cramping muscle pains or stiffness pain or tenderness around the eyes and cheekbones rash stomach discomfort, upset, or pain swollen joints Incidence not known Loss or thinning of the hair Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Cardizem CD side effects

1819. This particular disease, Hypertension, interacts with the drug CaroSpir and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. CaroSpir may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- little or no urination
- high potassium level-- nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- o signs of other electrolyte imbalances--increased thirst or urination, confusion, vomiting , muscle pain, slurred speech, severe weakness, numbness , loss of coordination, feeling unsteady. Common side effects of CaroSpir may include: breast swelling or tenderness.

1820. This particular disease, Hypertension, interacts with the drug Catapres and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe chest

Drug Side Effects Summary

pain, shortness of breath, irregular heartbeats

- a very slow heart rate
- or a light-headed feeling, like you might pass out. Common Catapres side effects may include: drowsiness, dizziness
- feeling tired or irritable
- dry mouth
- constipation , loss of appetite
- or sleep problems (insomnia), nightmares .

1821. This particular disease, Hypertension, interacts with the drug Catapres-TTS and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- a very slow heart rate (fewer than 60 beats per minute)
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- hallucinations
- fever, pale skin
- painful or difficult urination
- numbness or cold feeling in your hands or feet
- a light-headed feeling, like you might pass out
- or severe skin irritation, swelling, burning, or blistering where the patch is worn. Common Catapres-TTS side effects may include: headache, dizziness, drowsiness, feeling tired
- feeling nervous
- dry eyes, dry mouth

Drug Side Effects Summary

- changes in your sense of taste
- sexual problems
- nausea, constipation
- or skin discoloration or mild irritation where the patch is worn.

1822. This particular disease, Hypertension, interacts with the drug Dyazide and has the following s

- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- a light-headed feeling, like you might pass out
- fast, slow, or uneven heart rate
- signs of liver or pancreas problems - loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting, fast heart rate , dark urine, jaundice (yellowing of the skin or eyes)
- high potassium level - nausea, weakness, tingly feeling, feeling weak or tired, slow heartbeats, loss of movement
- low potassium level - leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- lupus-like symptoms - muscle or joint pain, flu symptoms, chest pain, and a rash or patchy skin color that worsens in sunlight
- or other signs of an electrolyte imbalance - thirst, dry mouth, stomach pain, drowsiness, weakness, fast heart rate, muscle pain or weakness, feeling restless or light-headed. Common Dyazide side effects may include: stomach pain, nausea, diarrhea , constipation
- dizziness , headache
- blurred vision
- or dry mouth.

Drug Side Effects Summary

1823. This particular disease, Hypertension, interacts with the drug Esidrix and has the following side effects:

- No side effects listed

1824. This particular disease, Hypertension, interacts with the drug felodipine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Felodipine may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- chest pain
- feeling short of breath
- or swelling or rapid weight gain. Felodipine can cause mild gum swelling. Pay special attention to your dental hygiene while taking felodipine. Brush and floss your teeth regularly. Common side effects of felodipine may include: swelling in your arms, hands, legs, or feet
- headache
- or flushing (sudden warmth, redness, or tingly feeling).

1825. This particular disease, Hypertension, interacts with the drug hydrochlorothiazide / valsartan and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, hydrochlorothiazide and valsartan can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure . Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark colored urine. Hydrochlorothiazide and valsartan may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out

Drug Side Effects Summary

- eye pain , vision problems
- an unusual skin rash
- little or no urinating
- jaundice (yellowing of the skin or eyes)
- easy bruising, unusual bleeding
- a seizure
- kidney problems--little or no urination, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- or signs of an electrolyte imbalance--dry mouth, increased thirst, confusion, feeling restless, vomiting, muscle pain or weakness, drowsiness, lack of energy, fast heartbeats, decreased urination. Common side effects include: cold symptoms such as stuffy nose, sneezing, sore throat
- dizziness
- or headache .

1826. This particular disease, Hypertension, interacts with the drug Hyzaar and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- irregular heartbeats
- eye pain, vision problems
- kidney problems - little or no urination, rapid weight gain, swelling in your hands, feet, or ankles
- new or worsening symptoms of lupus - joint pain, and a skin rash on your cheeks or arms that worsens in sunlight
- or signs of an electrolyte imbalance - increased thirst or urination, confusion, vomiting,

Drug Side Effects Summary

constipation, muscle pain, leg cramps, irregular heartbeats, numbness or tingling, severe weakness, loss of movement, feeling jittery or unsteady. Common Hyzaar side effects may include: dizziness

- back pain
- or cold symptoms such as stuffy nose, sneezing, sore throat.

1827. This particular disease, Hypertension, interacts with the drug indapamide and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have signs of an electrolyte imbalance, such as: drowsiness, lack of energy, feeling tired
- leg cramps, muscle weakness or limp feeling
- severe weakness, loss of coordination, feeling unsteady
- fast or irregular heartbeats, fluttering in your chest, feeling restless
- numbness or tingling
- vomiting, constipation
- headache , confusion, slurred speech
- a light-headed feeling, like you might pass out
- dry mouth, increased thirst
- or little or no urinating. Common side effects of indapamide may include: dizziness
- weakness, tiredness
- back pain , muscle cramps
- feeling anxious or agitated
- headache
- or runny nose .

Drug Side Effects Summary

1828. This particular disease, Hypertension, interacts with the drug InnoPran XL and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common in children Chest tightness cough producing mucus difficulty with breathing Incidence not known Black, tarry stools blistering, peeling, or loosening of the skin blood in the urine bloody nose bloody stools blurred or loss of vision body aches or pain burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain or discomfort confusion about identity, place, and time congestion constipation cough cracks in the skin crying decreased awareness or responsiveness decreased urine output depersonalization diarrhea difficulty with swallowing dilated neck veins disturbed color perception dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position double vision dryness or soreness of the throat dysphoria euphoria fast, pounding, slow, or irregular heartbeat fever and chills general feeling of discomfort, illness, or weakness hair loss halos around lights headaches heavier menstrual periods hoarseness irregular breathing lightheadedness, dizziness, or fainting loss of heat from the body mental depression mimicry of speech or movements muscle or joint pain mutism nausea negativism night blindness noisy breathing overbright appearance of lights paleness or cold feeling in the fingertips and toes paranoia peculiar postures or movements, mannerisms, or grimacing pinpoint red or purple spots on the skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue quick to react or overreact emotionally rapidly changing moods rectal bleeding red skin lesions, often with a purple center red, irritated eyes red, swollen skin reddening of the skin, especially around the ears runny nose scaly skin seeing, hearing, or feeling things that are not there severe sleepiness short-term memory loss skin irritation or rash, including rash that looks like psoriasis skin rash, hives, or itching stomach pain and tenderness sores, ulcers, or white spots in the mouth or on the lips sweating swelling of the eyes, face, fingers, feet, or lower legs swollen glands tender, swollen glands in the neck tingling or pain in fingers or

Drug Side Effects Summary

toes when exposed to cold tunnel vision unusual bleeding or bruising unusual tiredness or weakness voice changes vomiting weight gain Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common in children Anxiety dry mouth hyperventilation irritability restlessness shaking sleepiness or unusual drowsiness trouble sleeping unusual dreams Less common in children Decreased appetite Incidence not known Dry eyes heartburn loss of strength or energy muscle weakness pain or discomfort in the chest, upper stomach, or throat stomach cramps unusual drowsiness, dullness, or feeling of sluggishness vivid dreams Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. InnoPran XL side effects

1829. This particular disease, Hypertension, interacts with the drug Lotrel and has the following side effects

- hives
- severe stomach pain
- difficulty breathing
- swelling of your face, lips, tongue, or throat. You may be more likely to have an allergic reaction if you are African-American. Some side effects may not occur until after you have used the medicine for several months. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- swelling in your hands or feet, rapid weight gain
- new or worsened chest pain

Drug Side Effects Summary

- fever , chills, sore throat , body aches, flu symptoms
- high potassium--nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- or liver problems--nausea, stomach pain (upper right side), itching, unusual tiredness, flu-like symptoms, dark urine, jaundice (yellowing of the skin or eyes). Common Lotrel side effects may include: cough
- dizziness
- or swelling in your hands or feet.

1830. This particular disease, Hypertension, interacts with the drug Micardis and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Also call your doctor at once if you have: a light-headed feeling, like you might pass out
- little or no urination
- swelling in your feet or ankles, rapid weight gain
- unusual pain or tightness in your lower body
- a skin ulcer
- or high potassium level - nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common Micardis side effects may include: stuffy nose, sinus pain
- back pain
- or diarrhea.

1831. This particular disease, Hypertension, interacts with the drug Microzide and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of

Drug Side Effects Summary

these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Back, leg, or stomach pains black, tarry stools bleeding gums blistering, peeling, or loosening of the skin bloating blood in the urine or stools blue lips and fingernails blurred vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain or tightness chills clay-colored stools cloudy urine cold sweats confusion constipation cough or hoarseness coughing that sometimes produces a pink frothy sputum coughing up blood cracks in the skin darkened urine decrease in urine-concentrating ability decreased frequency or amount of urine diarrhea difficult, fast, or noisy breathing difficulty with swallowing dizziness, faintness, or lightheadedness when getting up from a lying or sitting position dry mouth fast or irregular heartbeat fever flushed, dry skin fruit-like breath odor general body swelling general feeling of discomfort or illness greatly decreased frequency of urination or amount of urine headache hives, itching, rash increased hunger increased sweating increased thirst increased urination indigestion joint pain, stiffness, or swelling loss of appetite loss of heat from the body lower back or side pain muscle cramps or pain nausea or vomiting nosebleeds numbness, tingling, pain, or weakness in the hands or feet painful or difficult urination pains in the stomach or side, possibly radiating to the back pale skin persistent non-healing sore pink growth pinpoint red spots on the skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red, irritated eyes red skin lesions, often with a purple center red, swollen skin reddish patch or irritated area redness, soreness, or scaly skin seizures shiny bump sore throat sores, ulcers, or white spots on the lips or in the mouth sores, welting, or blisters swelling of the face, fingers, legs, ankles, feet, or lower legs swollen or painful glands tenderness of salivary glands thickening of bronchial secretions trembling trouble breathing unpleasant breath odor unusual bleeding or bruising unusual tiredness or weakness unusual weight loss vomiting of blood weakness and heaviness of the legs weight gain white, yellow or waxy scar-like area yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away

Drug Side Effects Summary

during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Cramping decreased interest in sexual intercourse difficulty having a bowel movement (stool) feeling of constant movement of self or surroundings hair loss or thinning of the hair inability to have or keep an erection increased sensitivity of the skin to sunlight loss in sexual ability, desire, drive, or performance muscle spasm redness or other discoloration of the skin restlessness sensation of spinning severe sunburn weakness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Microzide side effects

1832. This particular disease, Hypertension, interacts with the drug nebivolol and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Nebivolol may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- rapid weight gain
- shortness of breath
- slow or uneven heartbeats
- or numbness or cold feeling in your hands and feet. Common side effects of nebivolol may include:
dizziness
- swelling in your legs
- slow heartbeats
- tiredness

Drug Side Effects Summary

- or headache .

1833. This particular disease, Hypertension, interacts with the drug prazosin and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Prazosin may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- pounding heartbeats or fluttering in your chest
- new or worsening chest pain
- or upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Call your doctor or seek emergency medical attention if your erection is painful or lasts longer than 4 hours. A prolonged erection (priapism) can damage the penis. Common side effects of prazosin may include: dizziness, drowsiness
- headache
- feeling weak or tired
- or nausea .

1834. This particular disease, Hypertension, interacts with the drug Procardia XL and has the follow

- worsening chest pain
- pounding heartbeats or fluttering in your chest
- a light-headed feeling, like you might pass out
- swelling in your hands or lower legs
- or upper stomach pain, jaundice (yellowing of the skin or eyes). You may have more severe or more frequent episodes of angina when you first start taking Procardia XL or whenever your dose is changed. Common side effects of Procardia XL may include: swelling

Drug Side Effects Summary

- flushing (warmth, redness, or tingly feeling)
- headache , dizziness
- nausea , heartburn
- or feeling weak or tired.

1835. This particular disease, Hypertension, interacts with the drug quinapril and has the following

- hives
- severe stomach pain
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Quinapril may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- fast or pounding heartbeats, fluttering in your chest
- little or no urination
- pain or burning when you urinate
- jaundice (yellowing of the skin or eyes)
- numbness , tingling, or burning pain in your hands or feet
- fever, chills, cold or flu symptoms
- high potassium-- nausea , chest pain, irregular heartbeats, loss of movement
- or low sodium-- headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady. Common side effects of quinapril may include: headache
- cough
- or dizziness , tired feeling.

1836. This particular disease, Hypertension, interacts with the drug terazosin and has the following

- hives

Drug Side Effects Summary

- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have any of these serious side effects: fast or pounding heartbeats or fluttering in your chest
- feeling like you might pass out
- swelling in your hands, ankles, or feet
- or penis erection that is painful or lasts 4 hours or longer. Less serious terazosin side effects may include: mild dizziness
- weakness, drowsiness
- blurred vision
- nausea
- or headache.

1837. This particular disease, Hypertension, interacts with the drug Accupril and has the following :

- hives
- severe stomach pain
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- fast or pounding heartbeats, fluttering in your chest
- little or no urination
- pain or burning when you urinate
- jaundice (yellowing of the skin or eyes)
- numbness, tingling, or burning pain in your hands or feet
- fever, chills, cold or flu symptoms
- high potassium - nausea, chest pain, irregular heartbeats, loss of movement

Drug Side Effects Summary

- or low sodium - headache, confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady. Common Accupril side effects may include: headache
- cough
- or dizziness, tired feeling.

1838. This particular disease, Hypertension, interacts with the drug Adalat CC and has the following

- worsening chest pain
- pounding heartbeats or fluttering in your chest
- a light-headed feeling, like you might pass out
- swelling in your hands or lower legs
- or upper stomach pain, jaundice (yellowing of the skin or eyes). You may have more severe or more frequent episodes of angina when you first start taking Adalat CC or whenever your dose is changed. Common Adalat CC side effects may include: swelling
- flushing (warmth, redness, or tingly feeling)
- headache , dizziness
- nausea , heartburn
- or feeling weak or tired.

1839. This particular disease, Hypertension, interacts with the drug Afeditab CR and has the followi

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Bloating or swelling of the face, arms, hands, lower legs, or feet cough difficult or labored breathing dizziness or lightheadedness fast, irregular, pounding, or racing heartbeat or pulse feeling of warmth headache muscle cramps rapid weight gain shakiness in the legs, arms, hands, or feet shortness of breath

Drug Side Effects Summary

tightness in the chest tingling of the hands or feet trembling or shaking of the hands or feet unusual weight gain or loss weakness wheezing Less common Blue lips and fingernails chest congestion chest pain chills coughing that sometimes produces a pink frothy sputum decreased urine output difficult, fast, or noisy breathing, sometimes with wheezing dilated neck veins extreme fatigue fever increased sweating irregular breathing nausea pain or discomfort in the arms, jaw, back, or neck pale skin severe unusual tiredness or weakness sweating troubled breathing vomiting Rare Black, tarry stools bleeding gums blood in the eyes blood in the urine or stools bloody stools bluish color blurred vision body aches or pain changes in skin color cold sweats dark urine difficulty with swallowing dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position dry mouth dryness or soreness of throat ear congestion or pain extra heartbeats eye pain feeling unusually cold general tiredness and weakness headache, severe and throbbing hoarseness increased urge to urinate during the night irritation in the mouth itching large, hive-like swelling on face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs light-colored stools nasal congestion neck pain nervousness no blood pressure or pulse noisy breathing pain pain in the groin or genitals pain or burning while urinating painful or difficult urination pinpoint red spots on the skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue redness and swelling of the gums redness in the whites of the eyes runny nose shakiness and unsteady walk sharp back pain just below ribs shivering skin rash small clicking, bubbling, or rattling sounds in the lung when listening with a stethoscope sneezing sore throat sores, ulcers, or white spots on the lips or in the mouth stopping of heart swelling around the eyes swollen glands tenderness tender, swollen glands in the neck troubled breathing with exertion unconsciousness unsteadiness, trembling, or other problems with muscle control or coordination unusual bleeding or bruising upper right abdominal or stomach pain vision changes voice changes vomiting of blood or material that looks like coffee grounds waking to urinate at night yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts

Drug Side Effects Summary

to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Belching feeling of indigestion mood changes pain in the chest below the breastbone redness of the face, neck, arms, and occasionally, upper chest Less common Abnormal ejaculation bloody nose burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings decreased interest in sexual intercourse decreased sexual performance or desire diarrhea difficulty having a bowel movement (stool) excess air or gas in stomach or intestines feeling of constant movement of self or surroundings full feeling general feeling of discomfort or illness inability to have or keep an erection increased need to urinate joint stiffness leg cramps or pain loss in sexual ability, desire, drive, or performance nervousness passing gas passing urine more often sneezing stuffy nose trouble sleeping Rare Acid or sour stomach ankle, knee, or great toe joint pain cracks in the skin difficulty with moving discouragement excessive muscle tone fear or nervousness feeling sad or empty hair loss or thinning of the hair heartburn increased sensitivity of the skin to sunlight irritability itching, pain, redness, swelling, tenderness, or warmth on the skin lack of appetite lack or loss of strength loss of heat from the body loss of interest or pleasure loss of strength or energy loss in sexual ability, desire, drive, or performance lower back or side pain muscle pain or weakness muscle stiffness muscle tension or tightness pain or burning in the throat red, swollen skin redness or other discoloration of the skin scaly skin severe sunburn sleepiness or unusual drowsiness sleeplessness stomach discomfort, upset, or pain swelling of the breasts or breast soreness in both females and males tiredness trouble concentrating unable to sleep Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Afeditab CR side effects

Drug Side Effects Summary

1840. This particular disease, Hypertension, interacts with the drug amlodipine / hydrochlorothiazide

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Amlodipine, hydrochlorothiazide, and valsartan may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- an unusual skin rash
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high or low potassium levels--leg cramps, nausea , constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- low magnesium level-- dizziness , feeling jittery, muscle cramps, muscle spasms, cough or choking feeling
- or low sodium level -- headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady. Common side effects of amlodipine, hydrochlorothiazide, and valsartan may include: dizziness
- headache, tiredness
- nausea, upset stomach
- stuffy nose, sinus pain, sore throat
- back pain , muscle spasms
- or swelling in your hands, ankles, or feet.

1841. This particular disease, Hypertension, interacts with the drug amlodipine / olmesartan and ha

- hives
- difficulty breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. When you first start taking this medicine, you may have new or worsening chest pain (angina), or you could have a heart attack. Seek emergency medical attention or call your doctor right away if you have symptoms such as: chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating. Amlodipine and olmesartan may cause serious side effects. Call your doctor at once if you have: new or worsening chest pain
- swelling in your hands or feet, rapid weight gain
- severe or ongoing diarrhea with weight loss
- pounding heartbeats or fluttering in your chest
- a light-headed feeling, like you might pass out
- or high potassium level--nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common side effects include swelling.

1842. This particular disease, Hypertension, interacts with the drug Apresoline and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Apresoline may cause serious side effects. Call your doctor at once if you have: chest pain or pressure, pain spreading to your jaw or shoulder
- fast or pounding heartbeats
- a light-headed feeling, like you might pass out
- numbness , tingling, or burning pain in your hands or feet
- painful or difficult urination
- little or no urination
- or lupus -like syndrome--joint pain or swelling with fever, swollen glands, muscle aches, chest pain, vomiting , unusual thoughts or behavior, and patchy skin color. Common side effects of Apresoline may include: chest pain, fast heart rate

Drug Side Effects Summary

- headache
- or nausea , vomiting, diarrhea , loss of appetite.

1843. This particular disease, Hypertension, interacts with the drug Atacand and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- little or no urination
- or high potassium level - nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common Atacand side effects may include: high potassium
- headache , back pain
- cold symptoms such as stuffy or runny nose , sneezing, sore throat
- dizziness
- or abnormal kidney test.

1844. This particular disease, Hypertension, interacts with the drug Avalide and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, Avalide can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, and dark colored urine. Also call your doctor at once if you have: eye pain, vision problems
- a light-headed feeling, like you might pass out

Drug Side Effects Summary

- little or no urinating
- jaundice (yellowing of the skin or eyes)
- easy bruising, unusual bleeding
- or signs of an electrolyte imbalance - increased thirst or urination, confusion, vomiting, constipation, muscle pain or weakness, leg cramps, bone pain, lack of energy, irregular heartbeats, tingling feeling. Common Avalide side effects may include: dizziness
- muscle or joint pain
- or feeling tired.

1845. This particular disease, Hypertension, interacts with the drug benazepril / hydrochlorothiazide

- a light-headed feeling, like you might pass out
- jaundice (yellowing of the skin or eyes)
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- fever, chills, flu-like symptoms, sore throat, mouth sores
- signs of an electrolyte imbalance--leg cramps, constipation , muscle pain or weakness, irregular heartbeats, feeling jittery, increased thirst or urination, numbness or tingling, loss of movement
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or new or worsening symptoms of lupus--joint pain, and a skin rash on your cheeks or arms that worsens in sunlight. Common side effects of hydrochlorothiazide and benazepril may include: dizziness
- cough
- headache
- or feeling tired.

1846. This particular disease, Hypertension, interacts with the drug bendroflumethiazide / nadolol

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Bendroflumethiazide and nadolol may cause serious side effects. Call your doctor at once if you have: fast, slow, or uneven heartbeat
- numbness or cold feeling in your hands and feet
- a light-headed feeling, like you might pass out
- little or no urination
- or signs of an electrolyte imbalance--dry mouth, increased thirst, mood changes, confusion, vomiting, muscle pain or weakness, lack of energy, fast heartbeats, little or no urine. Common side effects of bendroflumethiazide and nadolol may include: dizziness
- or feeling tired.

1847. This particular disease, Hypertension, interacts with the drug bendroflumethiazide and has the

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of fluid and electrolyte problems like

Drug Side Effects Summary

mood changes, confusion, muscle pain or weakness, a heartbeat that does not feel normal, very bad dizziness or passing out, fast heartbeat , more thirst, seizures , feeling very tired or weak, not hungry, unable to pass urine or change in the amount of urine produced, dry mouth, dry eyes, or very bad upset stomach or throwing up. Signs of a pancreas problem (pancreatitis) like very bad stomach pain, very bad back pain , or very bad upset stomach or throwing up. Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain. Dark urine or yellow skin or eyes. Any unexplained bruising or bleeding. Fever or chills. Sore throat . Feeling very tired or weak. Bendroflumethiazide side effects

1848. This particular disease, Hypertension, interacts with the drug bisoprolol / hydrochlorothiazide

- chest pain
- fast, slow, or uneven heartbeats
- a light-headed feeling, like you might pass out
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- liver problems-- nausea , upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- signs of an electrolyte imbalance--increased thirst or urination, constipation , muscle pain or weakness, leg cramps, numbness or tingling, feeling jittery, irregular heartbeats, fluttering in your chest, or a choking feeling
- low blood sugar-- headache , hunger, sweating, irritability, dizziness , fast heart rate , and feeling anxious or shaky. Common side effects of hydrochlorothiazide and bisoprolol may include: dizziness
- feeling weak or tired
- diarrhea, nausea, indigestion
- headache

Drug Side Effects Summary

- or cough.

1849. This particular disease, Hypertension, interacts with the drug Cardizem LA and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Body aches or pain congestion cough dryness or soreness of the throat fever hoarseness runny nose tender or swollen glands in the neck trouble swallowing voice changes Less common Chest pain, discomfort, or tightness chills diarrhea difficult or labored breathing feeling faint, dizzy, or lightheaded feeling of warmth or heat flushing or redness of the skin, especially on the face and neck general feeling of discomfort or illness headache joint pain loss of appetite muscle aches and pains nausea shivering slow or irregular heartbeat sweating swelling of the hands, ankles, feet, or lower legs trouble sleeping unusual tiredness or weakness vomiting Incidence not known Blistering, peeling, or loosening of the skin itching large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs no heartbeat red skin lesions, often with a purple center red, irritated eyes sores, ulcers, or white spots in the mouth or on the lips Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Sneezing stuffy nose Less common Acid or sour stomach belching constipation continuing ringing or buzzing or other unexplained noise in the ears degenerative disease of the joint difficulty with moving hearing loss heartburn indigestion lack or loss of strength muscle aching or cramping muscle pains or stiffness pain or tenderness around the eyes and cheekbones rash stomach discomfort, upset, or pain swollen joints Incidence not known Loss or thinning of the hair Other side

Drug Side Effects Summary

effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Cardizem LA side effects

1850. This particular disease, Hypertension, interacts with the drug Cardura and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- severe ongoing stomach pain or bloating
- new or worsening chest pain
- trouble breathing
- or a penis erection that is painful or lasts 4 hours or longer. Common Cardura side effects may include: low blood pressure, dizziness
- drowsiness
- headache
- or feeling weak or tired.

1851. This particular disease, Hypertension, interacts with the drug Dilacor XR and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Body aches or pain congestion cough dryness or soreness of the throat fever hoarseness runny nose tender or swollen glands in the neck trouble swallowing voice changes Less common Chest pain, discomfort, or tightness chills diarrhea difficult or labored breathing feeling faint, dizzy, or lightheaded feeling of

Drug Side Effects Summary

warmth or heat flushing or redness of the skin, especially on the face and neck general feeling of discomfort or illness headache joint pain loss of appetite muscle aches and pains nausea shivering slow or irregular heartbeat sweating swelling of the hands, ankles, feet, or lower legs trouble sleeping unusual tiredness or weakness vomiting Incidence not known Blistering, peeling, or loosening of the skin itching large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs no heartbeat red skin lesions, often with a purple center red, irritated eyes sores, ulcers, or white spots in the mouth or on the lips Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Sneezing stuffy nose Less common Acid or sour stomach belching constipation continuing ringing or buzzing or other unexplained noise in the ears degenerative disease of the joint difficulty with moving hearing loss heartburn indigestion lack or loss of strength muscle aching or cramping muscle pains or stiffness pain or tenderness around the eyes and cheekbones rash stomach discomfort, upset, or pain swollen joints Incidence not known Loss or thinning of the hair Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Dilacor XR side effects

1852. This particular disease, Hypertension, interacts with the drug Dilt-XR and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Body aches or pain congestion cough dryness or soreness of the throat fever hoarseness runny nose tender or swollen

Drug Side Effects Summary

glands in the neck trouble swallowing voice changes Less common Chest pain, discomfort, or tightness chills diarrhea difficult or labored breathing feeling faint, dizzy, or lightheaded feeling of warmth or heat flushing or redness of the skin, especially on the face and neck general feeling of discomfort or illness headache joint pain loss of appetite muscle aches and pains nausea shivering slow or irregular heartbeat sweating swelling of the hands, ankles, feet, or lower legs trouble sleeping unusual tiredness or weakness vomiting Incidence not known Blistering, peeling, or loosening of the skin itching large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs no heartbeat red skin lesions, often with a purple center red, irritated eyes sores, ulcers, or white spots in the mouth or on the lips Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Sneezing stuffy nose Less common Acid or sour stomach belching constipation continuing ringing or buzzing or other unexplained noise in the ears degenerative disease of the joint difficulty with moving hearing loss heartburn indigestion lack or loss of strength muscle aching or cramping muscle pains or stiffness pain or tenderness around the eyes and cheekbones rash stomach discomfort, upset, or pain swollen joints Incidence not known Loss or thinning of the hair Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Dilt-XR side effects

1853. This particular disease, Hypertension, interacts with the drug Diltia XT and has the following :

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your

Drug Side Effects Summary

doctor immediately if any of the following side effects occur: More common Body aches or pain congestion cough dryness or soreness of the throat fever hoarseness runny nose tender or swollen glands in the neck trouble swallowing voice changes Less common Chest pain, discomfort, or tightness chills diarrhea difficult or labored breathing feeling faint, dizzy, or lightheaded feeling of warmth or heat flushing or redness of the skin, especially on the face and neck general feeling of discomfort or illness headache joint pain loss of appetite muscle aches and pains nausea shivering slow or irregular heartbeat sweating swelling of the hands, ankles, feet, or lower legs trouble sleeping unusual tiredness or weakness vomiting Incidence not known Blistering, peeling, or loosening of the skin itching large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs no heartbeat red skin lesions, often with a purple center red, irritated eyes sores, ulcers, or white spots in the mouth or on the lips Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Sneezing stuffy nose Less common Acid or sour stomach belching constipation continuing ringing or buzzing or other unexplained noise in the ears degenerative disease of the joint difficulty with moving hearing loss heartburn indigestion lack or loss of strength muscle aching or cramping muscle pains or stiffness pain or tenderness around the eyes and cheekbones rash stomach discomfort, upset, or pain swollen joints Incidence not known Loss or thinning of the hair Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Diltia XT side effects

1854. This particular disease, Hypertension, interacts with the drug Diltzac and has the following si

Drug Side Effects Summary

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Body aches or pain congestion cough dryness or soreness of the throat fever hoarseness runny nose tender or swollen glands in the neck trouble swallowing voice changes Less common Chest pain, discomfort, or tightness chills diarrhea difficult or labored breathing feeling faint, dizzy, or lightheaded feeling of warmth or heat flushing or redness of the skin, especially on the face and neck general feeling of discomfort or illness headache joint pain loss of appetite muscle aches and pains nausea shivering slow or irregular heartbeat sweating swelling of the hands, ankles, feet, or lower legs trouble sleeping unusual tiredness or weakness vomiting Incidence not known Blistering, peeling, or loosening of the skin itching large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs no heartbeat red skin lesions, often with a purple center red, irritated eyes sores, ulcers, or white spots in the mouth or on the lips Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Sneezing stuffy nose Less common Acid or sour stomach belching constipation continuing ringing or buzzing or other unexplained noise in the ears degenerative disease of the joint difficulty with moving hearing loss heartburn indigestion lack or loss of strength muscle aching or cramping muscle pains or stiffness pain or tenderness around the eyes and cheekbones rash stomach discomfort, upset, or pain swollen joints Incidence not known Loss or thinning of the hair Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Diltzac side effects

Drug Side Effects Summary

1855. This particular disease, Hypertension, interacts with the drug Epaned and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Blurred vision confusion dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position sweating unusual tiredness or weakness Less common Chest pain cough producing mucus diarrhea difficult or labored breathing fainting fever or chills nausea sneezing sore throat tightness in the chest vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Dizziness Less common Lack of strength rash Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Epaned side effects

1856. This particular disease, Hypertension, interacts with the drug eplerenone and has the following side effects:

- hives
- severe stomach pain
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Eplerenone may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- diarrhea , vomiting
- little or no urination
- fast or irregular heartbeats

Drug Side Effects Summary

- trouble breathing
- swelling in your feet or lower legs
- or high potassium-- nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common side effects of eplerenone may include: high potassium
- headache
- or dizziness .

1857. This particular disease, Hypertension, interacts with the drug Exforge HCT and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- an unusual skin rash
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high or low potassium levels - leg cramps, nausea , constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- low magnesium level - dizziness , feeling jittery, muscle cramps, muscle spasms, cough or choking feeling
- or low sodium level - headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady. Common Exforge HCT side effects may include: dizziness
- headache, tiredness
- nausea, upset stomach
- stuffy nose, sinus pain, sore throat

Drug Side Effects Summary

- back pain , muscle spasms
- or swelling in your hands, ankles, or feet.

1858. This particular disease, Hypertension, interacts with the drug fosinopril and has the following

- hives
- severe stomach pain
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Fosinopril may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- sudden weakness or ill feeling, fever, chills, sore throat , painful mouth sores, cough, trouble breathing
- little or no urinating
- or high potassium-- nausea , slow or unusual heart rate, weakness, loss of movement. Common side effects of fosinopril may include: cough, runny or stuffy nose
- muscle or joint pain
- dizziness , headache , tired feeling
- nausea, vomiting, diarrhea
- or mild skin itching or rash.

1859. This particular disease, Hypertension, interacts with the drug hydrochlorothiazide / irbesartan

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, hydrochlorothiazide and irbesartan can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure . Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness

Drug Side Effects Summary

especially if you also have fever, unusual tiredness, and dark colored urine. Also call your doctor at once if you have: eye pain , vision problems

- a light-headed feeling, like you might pass out
- little or no urinating
- jaundice (yellowing of the skin or eyes)
- easy bruising, unusual bleeding
- or signs of an electrolyte imbalance--increased thirst or urination, confusion, vomiting, constipation , muscle pain or weakness, leg cramps, bone pain, lack of energy, irregular heartbeats, tingly feeling. Common side effects of hydrochlorothiazide and irbesartan may include: dizziness
- muscle or joint pain
- or feeling tired.

1860. This particular disease, Hypertension, interacts with the drug hydrochlorothiazide / olmesartan

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, this medicine can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure . Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark colored urine. Also call your doctor at once if you have: a light-headed feeling, like you might pass out
- eye pain , vision problems
- an unusual skin rash
- easy bruising, unusual bleeding
- jaundice (yellowing of the skin or eyes)
- kidney problems--little or no urination, painful or difficult urination, swelling in your feet or ankles,

Drug Side Effects Summary

feeling tired or short of breath

- low levels of sodium in the body-- headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady
- high potassium--slow or unusual heart rate, weakness, loss of movement
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects include: nausea
- dizziness
- cold symptoms such as stuffy nose, sneezing, sore throat
- or high levels of uric acid in your blood.

1861. This particular disease, Hypertension, interacts with the drug hydrochlorothiazide / telmisartan

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, hydrochlorothiazide and telmisartan can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark colored urine. Also call your doctor at once if you have: eye pain , vision problems
- a light-headed feeling, like you might pass out
- jaundice (yellowing of the skin or eyes)
- swelling in your hands or feet, unusual weight gain
- easy bruising, unusual bleeding
- or signs of an electrolyte imbalance--dry mouth, increased thirst, drowsiness, confusion, feeling restless, vomiting, muscle pain or weakness, lack of energy, fast heartbeats, little or no urine, or a

Drug Side Effects Summary

seizure. Common side effects of hydrochlorothiazide and telmisartan may include: dizziness , feeling tired

- nausea , diarrhea
- back pain
- or cold or flu symptoms.

1862. This particular disease, Hypertension, interacts with the drug Lotensin and has the following

- a light-headed feeling, like you might pass out
- little or no urinating
- high potassium - nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- or liver problems - stomach pain (upper right side), loss of appetite, dark urine, jaundice (yellowing of the skin or eyes). Common Lotensin side effects may include: headache
- or cough.

1863. This particular disease, Hypertension, interacts with the drug Matzim LA and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Body aches or pain congestion cough dryness or soreness of the throat fever hoarseness runny nose tender or swollen glands in the neck trouble swallowing voice changes Less common Chest pain, discomfort, or tightness chills diarrhea difficult or labored breathing feeling faint, dizzy, or lightheaded feeling of warmth or heat flushing or redness of the skin, especially on the face and neck general feeling of discomfort or illness headache joint pain loss of appetite muscle aches and pains nausea shivering slow or irregular heartbeat sweating swelling of the hands, ankles, feet, or lower legs trouble

Drug Side Effects Summary

sleeping unusual tiredness or weakness vomiting Incidence not known Blistering, peeling, or loosening of the skin itching large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs no heartbeat red skin lesions, often with a purple center red, irritated eyes sores, ulcers, or white spots in the mouth or on the lips Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Sneezing stuffy nose Less common Acid or sour stomach belching constipation continuing ringing or buzzing or other unexplained noise in the ears degenerative disease of the joint difficulty with moving hearing loss heartburn indigestion lack or loss of strength muscle aching or cramping muscle pains or stiffness pain or tenderness around the eyes and cheekbones rash stomach discomfort, upset, or pain swollen joints Incidence not known Loss or thinning of the hair Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Matzim LA side effects

1864. This particular disease, Hypertension, interacts with the drug methyldopa and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Methyldopa may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- pale or yellowed skin, dark colored urine, fever, confusion or weakness
- uncontrolled or involuntary muscle movements
- heart problems--swelling, rapid weight gain, feeling short of breath

Drug Side Effects Summary

- or signs of liver or pancreas problems--loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting, fast heart rate , dark urine, jaundice (yellowing of the skin or eyes).

Common side effects of methyldopa may include: drowsiness

- weakness

- or headache .

1865. This particular disease, Hypertension, interacts with the drug metolazone and has the following

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Metolazone may cause serious side effects. Call your doctor at once if you have: chest pain

- pounding heartbeats or fluttering in your chest

- easy bruising or bleeding

- a light-headed feeling, like you might pass out

- liver problems-- nausea , upper stomach pain, itching, tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)

- low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling

- low levels of sodium in the body-- headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady

- other signs of an electrolyte imbalance--dry mouth, nausea, vomiting, weakness, drowsiness, lack of energy, muscle pain, little or no urination, or feeling restless

- or severe skin reaction--fever, sore throat , swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body)

and causes blistering and peeling. Common side effects of metolazone may include: dizziness,

Drug Side Effects Summary

spinning sensation

- drowsiness, tiredness
- depressed mood
- muscle or joint pain
- numbness or tingly feeling
- nausea, stomach pain, loss of appetite
- or diarrhea , constipation.

1866. This particular disease, Hypertension, interacts with the drug Micardis HCT and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, Micardis HCT can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark colored urine. Also call your doctor at once if you have: eye pain , vision problems
- a light-headed feeling, like you might pass out
- jaundice (yellowing of the skin or eyes)
- swelling in your hands or feet, unusual weight gain
- easy bruising, unusual bleeding
- or signs of an electrolyte imbalance--dry mouth, increased thirst, drowsiness, confusion, feeling restless, vomiting, muscle pain or weakness, lack of energy, fast heartbeats, little or no urine, or a seizure. Common side effects of Micardis HCT may include: dizziness , feeling tired
- nausea , diarrhea
- back pain

Drug Side Effects Summary

- or cold or flu symptoms.

1867. This particular disease, Hypertension, interacts with the drug minoxidil and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Minoxidil may cause serious side effects. Call your doctor at once if you have: new or worsening chest pain
- chest pain spreading to your jaw or shoulder
- fast or pounding heartbeats
- swelling in your legs, ankles, or feet
- rapid weight gain, especially in your face and midsection
- shortness of breath
- a light-headed feeling, like you might pass out
- fluid build-up in the lungs--pain when you breathe, feeling short of breath while lying down, wheezing, gasping for breath, cough with foamy mucus
- or severe skin reaction--fever, sore throat , swelling in your face or tongue, burning in your eyes, skin pain, followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of minoxidil may include: changes in color, length, or thickness of body or facial hair
- nausea , vomiting
- rash
- or breast pain or tenderness.

1868. This particular disease, Hypertension, interacts with the drug Nexiclon XR and has the follow

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of

Drug Side Effects Summary

these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Mental depression swelling of the feet and lower legs Rare Anxiety blistering, burning, crusting, dryness, or flaking of the skin chest pain or discomfort confusion as to time, place, or person decreased urine output dilated neck veins drowsiness dry mouth fast, pounding, or irregular heartbeat or pulse fever general feeling of discomfort or illness holding false beliefs that cannot be changed by fact hyperventilation irregular breathing irritability itching, scaling, severe redness, soreness, or swelling of the skin large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs lightheadedness, dizziness, or fainting mental depression paleness or cold feeling in the fingertips and toes pounding, slow heartbeat problems in urination or increase in the amount of urine raised red swellings on the skin, lips, tongue, or in the throat restlessness seeing or hearing things that are not there shaking shortness of breath skin rash swelling of the face, fingers, feet, or lower legs tightness in the chest tingling or pain in the fingers or toes when exposed to cold trouble with sleeping troubled breathing unusual excitement, nervousness, or restlessness unusual tiredness or weakness vivid dreams or nightmares weight gain wheezing Symptoms of overdose Dizziness (extreme) or faintness feeling cold pinpoint pupils of the eyes unusual tiredness or weakness (extreme) Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Constipation Less common Darkening of the skin decreased sexual ability dry, itching, or burning eyes loss of appetite nausea or vomiting Rare Blurred vision decreased interest in sexual intercourse hair loss or thinning of the hair inability to have or keep an erection leg cramps loss in sexual ability, desire, drive, or performance muscle or joint pain pale skin swelling of the breasts or breast soreness in both

Drug Side Effects Summary

females and males weakness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Nexiclon XR side effects

1869. This particular disease, Hypertension, interacts with the drug Taztia XT and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Body aches or pain congestion cough dryness or soreness of the throat fever hoarseness runny nose tender or swollen glands in the neck trouble swallowing voice changes Less common Chest pain, discomfort, or tightness chills diarrhea difficult or labored breathing feeling faint, dizzy, or lightheaded feeling of warmth or heat flushing or redness of the skin, especially on the face and neck general feeling of discomfort or illness headache joint pain loss of appetite muscle aches and pains nausea shivering slow or irregular heartbeat sweating swelling of the hands, ankles, feet, or lower legs trouble sleeping unusual tiredness or weakness vomiting Incidence not known Blistering, peeling, or loosening of the skin itching large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs no heartbeat red skin lesions, often with a purple center red, irritated eyes sores, ulcers, or white spots in the mouth or on the lips Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Sneezing stuffy nose Less common Acid or sour stomach belching constipation continuing ringing or buzzing or other unexplained noise in the ears degenerative disease of the joint difficulty with

Drug Side Effects Summary

moving hearing loss heartburn indigestion lack or loss of strength muscle aching or cramping muscle pains or stiffness pain or tenderness around the eyes and cheekbones rash stomach discomfort, upset, or pain swollen joints Incidence not known Loss or thinning of the hair Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Taztia XT side effects

1870. This particular disease, Hypertension, interacts with the drug Tekturna and has the following

- hives , itching
- vomiting, severe stomach pain
- dizziness , chest tightness, difficult breathing
- swelling of your face, lips, tongue, or throat. If you have an allergic reaction to Tekturna you should not take it again. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- little or no urination
- high potassium - nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- or low levels of sodium in the body - headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady. Common Tekturna side effects may include: diarrhea.

1871. This particular disease, Hypertension, interacts with the drug Tiadylt ER and has the following

- No side effects listed

1872. This particular disease, Hypertension, interacts with the drug Tiazac and has the following side

- chest pain

Drug Side Effects Summary

- slow heartbeats
- pounding heartbeats or fluttering in your chest
- a light-headed feeling, like you might pass out
- or heart problems - swelling, rapid weight gain, feeling short of breath. Common Tiazac side effects may include: swelling
- dizziness , weakness
- headache
- nausea
- or rash.

1873. This particular disease, Hypertension, interacts with the drug torsemide and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor right away if you have: a light-headed feeling, like you might pass out
- ringing in your ears, sudden hearing loss
- kidney problems--little or no urination, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- low magnesium level-- dizziness , irregular heartbeats, feeling jittery, muscle cramps, muscle spasms, cough or choking feeling
- low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- or low sodium level-- headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady. Common side effects of torsemide may include: increased urination
- or headache.

Drug Side Effects Summary

1874. This particular disease, Hypertension, interacts with the drug Zestoretic and has the following

- hives
- severe stomach pain
- difficulty breathing
- swelling of your face, lips, tongue, or throat. You may be more likely to have an allergic reaction if you are African-American. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- eye pain , vision problems
- little or no urination
- weakness, drowsiness, or feeling restless
- fever, chills, sore throat , mouth sores, trouble swallowing
- jaundice (yellowing of the skin or eyes)
- high potassium - nausea , tingly feeling, chest pain, irregular heartbeats, loss of movement
- low potassium - leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- or low sodium - headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady. Common Zestoretic side effects may include: cough
- headache
- dizziness
- or tired feeling.

1875. This particular disease, Hypertension, interacts with the drug Ziac and has the following side

- chest pain
- fast, slow, or uneven heartbeats
- a light-headed feeling, like you might pass out

Drug Side Effects Summary

- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- liver problems - nausea , upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- signs of an electrolyte imbalance - increased thirst or urination, constipation , muscle pain or weakness, leg cramps, numbness or tingling, feeling jittery, irregular heartbeats, fluttering in your chest, or a choking feeling
- low blood sugar - headache , hunger, sweating, irritability, dizziness , fast heart rate , and feeling anxious or shaky. Common Ziac side effects may include: dizziness
- feeling weak or tired
- diarrhea, nausea, indigestion
- headache
- or cough.

1876. This particular disease, Hypertension, interacts with the drug Accuretic and has the following

- a light-headed feeling, like you might pass out
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- swelling, weight gain, shortness of breath
- sudden weakness or ill feeling, fever, chills, sore throat, mouth sores, trouble swallowing
- high potassium-- nausea , tingly feeling, chest pain, irregular heartbeats, loss of movement
- low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- or low levels of sodium-- headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady. Common side effects of Accuretic may include: cough
- headache, dizziness

Drug Side Effects Summary

- tired feeling
- or muscle pain.

1877. This particular disease, Hypertension, interacts with the drug acebutolol and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Acebutolol may cause serious side effects. Call your doctor at once if you have: shortness of breath (even with mild exertion), swelling, rapid weight gain
- new or worsening chest pain
- slow heartbeats
- a light-headed feeling, like you might pass out
- or dangerously high blood pressure--severe headache , blurred vision, pounding in your neck or ears, nosebleed, anxiety , confusion, severe chest pain, shortness of breath, irregular heartbeats.

Common side effects of acebutolol may include: headache, dizziness

- feeling tired
- nausea , upset stomach
- diarrhea , constipation
- or sleep problems (insomnia).

1878. This particular disease, Hypertension, interacts with the drug Aldactazide and has the following

- skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Aldactazide may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- breast swelling and tenderness

Drug Side Effects Summary

- low blood cell counts--easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath
- high potassium level--nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- or low sodium level -- headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady. Common side effects of Aldactazide may include: drowsiness, dizziness, lack of energy
- leg cramps
- weakness
- feeling like you might pass out
- severe pain in your upper stomach spreading to your back, nausea and vomiting
- or electrolyte imbalance.

1879. This particular disease, Hypertension, interacts with the drug Aldoril and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Unexplained fever shortly after starting to take this medicine Check with your doctor as soon as possible if any of the following side effects occur: Signs and symptoms of too much potassium loss Dry mouth increased thirst irregular heartbeats muscle cramps or pain nausea or vomiting unusual tiredness or weakness weak pulse Less common Mental depression or anxiety nightmares or unusually vivid dreams Rare Cough or hoarseness dark or amber urine diarrhea or stomach cramps (severe or continuing) fever, chills, troubled breathing, and fast heartbeat general feeling of discomfort or illness or weakness joint pain

Drug Side Effects Summary

lower back or side pain painful or difficult urination pale stools skin rash, hives, or itching stomach pain (severe) with nausea and vomiting tiredness or weakness after having taken this medicine for several weeks (continuing) yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Dizziness or light-headedness when getting up from a lying or sitting position drowsiness dryness of mouth headache Less common Decreased sexual ability or interest in sex diarrhea increased sensitivity of skin to sunlight (skin rash, itching, redness or other discoloration of skin or severe sunburn after exposure to sunlight) loss of appetite numbness, tingling, pain, or weakness in hands or feet slow heartbeat stuffy nose swelling of breasts or unusual milk production Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Aldoril side effects

1880. This particular disease, Hypertension, interacts with the drug aliskiren / hydrochlorothiazide as follows:

- a light-headed feeling, like you might pass out
- little or no urination
- eye pain , vision problems
- high potassium level-- nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- low sodium level -- headache , confusion, slurred speech, severe weakness, vomiting, loss of

Drug Side Effects Summary

coordination, feeling unsteady

- or lupus -like symptoms--muscle or joint pain, fever, and a rash or patchy skin color that worsens in sunlight. Common side effects of hydrochlorothiazide and aliskiren may include: dizziness, tired feeling

- high potassium

- cough

- joint pain

- diarrhea

- or flu symptoms (fever, chills, body aches).

1881. This particular disease, Hypertension, interacts with the drug aliskiren and has the following

- hives , itching

- vomiting, severe stomach pain

- dizziness , chest tightness, difficult breathing

- swelling of your face, lips, tongue, or throat. If you have an allergic reaction to aliskiren you should not take it again. Aliskiren may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out

- little or no urination

- high potassium-- nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement

- or low levels of sodium in the body-- headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady. Common side effects of aliskiren may include: diarrhea.

1882. This particular disease, Hypertension, interacts with the drug amiloride / hydrochlorothiazide

Drug Side Effects Summary

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Hydrochlorothiazide and amiloride may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- high sodium-- headache , confusion, slurred speech, hallucinations, vomiting, severe weakness, muscle cramps, loss of coordination, feeling unsteady, shallow breathing
- or high potassium-- slow heart rate , weak pulse, muscle weakness, tingly feeling. Common side effects of hydrochlorothiazide and amiloride may include: diarrhea , constipation
- mild stomach pain
- or tired feeling.

1883. This particular disease, Hypertension, interacts with the drug amiloride and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop taking amiloride and call your doctor at once if you have: increased thirst, decreased urination
- heavy sweating, or hot and dry skin
- tremors, confusion, loss of consciousness
- jaundice (yellowing of the skin or eyes)
- high potassium--tiredness, numbness or tingling, slow or unusual heart rate, muscle weakness or limp feeling
- or low levels of sodium in the body-- headache , confusion, slurred speech, severe weakness, vomiting , loss of coordination, feeling unsteady. Common side effects of amiloride may include: nausea , stomach pain, gas, loss of appetite
- headache

Drug Side Effects Summary

- or rash.

1884. This particular disease, Hypertension, interacts with the drug amlodipine / celecoxib and has

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. amlodipine and celecoxib may cause serious side effects. Stop using amlodipine and celecoxib and call your doctor at once if you have:

nausea, weakness, tingly feeling

- new or worsening chest pain

- the first sign of any skin rash, no matter how mild

- rapid weight gain, feeling short of breath

- swelling in your arms, hands, legs, or feet

- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds

- liver problems--stomach pain (upper right side), dark urine, jaundice (yellowing of the skin or eyes)

- or low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed, cold hands and feet. Common side effects of amlodipine and celecoxib may include: stomach pain, heartburn , gas, diarrhea , nausea

- drowsiness, feeling tired

- swelling

- increased urination

- joint pain

- rash

- flushing (sudden warmth or redness in your face)

- headache , dizziness

- or cold symptoms such as stuffy nose, sneezing, sore throat.

Drug Side Effects Summary

1885. This particular disease, Hypertension, interacts with the drug amlodipine / hydrochlorothiazide

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Amlodipine, hydrochlorothiazide, and olmesartan may cause serious side effects. Call your doctor at once if you have: worsening chest pain
- a light-headed feeling, like you might pass out
- an unusual skin rash
- pain or burning when you urinate
- severe or ongoing diarrhea with weight loss
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or signs of an electrolyte imbalance--increased thirst or urination, confusion, vomiting, constipation , muscle pain or weakness, leg cramps, lack of energy, irregular heartbeats, numbness or tingling, loss of movement. Common side effects of amlodipine, hydrochlorothiazide, and olmesartan may include: dizziness
- headache , tired feeling
- swelling in your hands or feet
- nausea , diarrhea
- runny or stuffy nose, sore throat
- muscle spasms
- swollen joints
- or painful urination.

1886. This particular disease, Hypertension, interacts with the drug amlodipine / perindopril and ha

- hives

Drug Side Effects Summary

- severe stomach pain
- difficulty breathing
- swelling of your face, lips, tongue, or throat. You may be more likely to have an allergic reaction if you are African-American. Amlodipine and perindopril may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- swelling in your hands or feet, rapid weight gain
- little or no urinating
- new or worsened chest pain
- jaundice (yellowing of the skin or eyes)
- or high potassium-- nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common side effects of amlodipine and perindopril may include: cough
- headache , dizziness
- or swelling in your hands or feet.

1887. This particular disease, Hypertension, interacts with the drug amlodipine / telmisartan and has

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, amlodipine and telmisartan can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure . Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark colored urine. Amlodipine and telmisartan may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- new or worsened chest pain

Drug Side Effects Summary

- pounding heartbeats or fluttering in your chest
- tremors, muscle stiffness or twitching
- or high potassium-- nausea , slow or unusual heart rate, weakness, loss of movement. Common side effects of amlodipine and telmisartan may include: swelling in your hands or feet
- fast or pounding heartbeats
- dizziness , drowsiness
- flushing (warmth, redness, or tingly feeling)
- back pain
- or nausea, diarrhea, stomach pain.

1888. This particular disease, Hypertension, interacts with the drug Atacand HCT and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, Atacand HCT can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure . Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, and dark colored urine. Also call your doctor at once if you have: eye pain , vision problems
- a light-headed feeling, like you might pass out
- little or no urinating
- jaundice (yellowing of the skin or eyes)
- easy bruising, unusual bleeding
- or signs of an electrolyte imbalance--increased thirst or urination, confusion, vomiting, constipation , muscle pain or weakness, leg cramps, bone pain, lack of energy, irregular heartbeats, tingly feeling. Common side effects of Atacand HCT may include: dizziness

Drug Side Effects Summary

- back pain
- or cold or flu symptoms.

1889. This particular disease, Hypertension, interacts with the drug azilsartan medoxomil / chlorthalidone and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Azilsartan and chlorthalidone may cause serious side effects. Call your doctor at once if you have: feeling like you might pass out
- little or no urination
- swelling, weight gain, feeling short of breath
- or signs of electrolyte imbalance--dry mouth, extreme thirst, weakness, drowsiness, restless feeling, confusion, increased or decreased urination, nausea and vomiting , constipation , muscle pain or weakness, fast heart rate , or seizure (convulsions). Common side effects of azilsartan and chlorthalidone may include: dizziness
- or feeling tired.

1890. This particular disease, Hypertension, interacts with the drug azilsartan medoxomil and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Azilsartan may cause serious side effects. Call your doctor at once if you have: little or no urination
- or a light-headed feeling, like you might pass out. Common azilsartan side effects may include: diarrhea.

1891. This particular disease, Hypertension, interacts with the drug betaxolol and has the following side effects:

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Betaxolol may cause serious side effects. Call your doctor at once if you have: shortness of breath (even with mild exertion), swelling, rapid weight gain
- slow or uneven heartbeats
- a light-headed feeling, like you might pass out
- or lupus -like syndrome--joint pain or swelling with fever, swollen glands, muscle aches, chest pain, vomiting , unusual thoughts or behavior, and patchy skin color. Slow heartbeats may be more likely in older adults. Common side effects of betaxolol may include: slow heartbeats
- headache, dizziness
- feeling weak or tired
- upset stomach
- or decreased sex drive, impotence , or difficulty having an orgasm.

1892. This particular disease, Hypertension, interacts with the drug Brevibloc and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tell your caregivers right away if you have: a light-headed feeling, like you might pass out
- very slow heartbeats
- swelling in your hands or feet
- wheezing, chest tightness, feeling short of breath
- weak or shallow breathing
- pain, swelling, irritation, bruising, or skin changes around the IV needle
- cold feeling in your hands and feet

Drug Side Effects Summary

- high potassium-- nausea , slow or unusual heart rate, weakness, loss of movement
- or low blood sugar -- headache , hunger, sweating, irritability, dizziness , nausea, fast heart rate , and feeling anxious or shaky. Common side effects of Brevibloc may include: increased sweating
- nausea
- dizziness, drowsiness
- or low blood pressure .

1893. This particular disease, Hypertension, interacts with the drug candesartan / hydrochlorothiaz

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, hydrochlorothiazide and candesartan can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure . Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, and dark colored urine. Also call your doctor at once if you have: eye pain , vision problems
- a light-headed feeling, like you might pass out
- little or no urinating
- jaundice (yellowing of the skin or eyes)
- easy bruising, unusual bleeding
- or signs of an electrolyte imbalance--increased thirst or urination, confusion, vomiting, constipation , muscle pain or weakness, leg cramps, bone pain, lack of energy, irregular heartbeats, tingly feeling. Common side effects of hydrochlorothiazide and candesartan may include: dizziness
- back pain
- or cold or flu symptoms.

Drug Side Effects Summary

1894. This particular disease, Hypertension, interacts with the drug captopril / hydrochlorothiazide

- hives
- severe stomach pain
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Hydrochlorothiazide and captopril may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- eye pain , vision problems
- little or no urinating
- chest pain, pounding heartbeats or fluttering in your chest
- swelling, weight gain, shortness of breath
- pale skin, easy bruising or bleeding
- sudden weakness or ill feeling, fever, chills, sore throat , mouth sores, trouble swallowing
- high potassium-- nausea , slow or unusual heart rate, weakness, loss of movement
- low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- low levels of sodium-- headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady
- or severe skin reaction--fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain, followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of hydrochlorothiazide and captopril may include: cough
- mild itching or rash
- joint pain
- numbness in your hands or feet
- decreased sense of taste

Drug Side Effects Summary

- or flushing (warmth, redness, or tingly feeling).

1895. This particular disease, Hypertension, interacts with the drug Cardene IV and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Blurred vision confusion dizziness, faintness, or lightheadedness when getting up from a lying or sitting position suddenly sweating unusual tiredness or weakness Less common Bleeding, blistering, burning, coldness, discoloration of skin, feeling of pressure, hives, infection, inflammation, itching, lumps, numbness, pain, rash, redness, scarring, soreness, stinging, swelling, tenderness, tingling, ulceration, or warmth at site blood in urine chills cold sweats convulsions decreased urine dry mouth extra heartbeat fast, pounding, or irregular heartbeat or pulse frequent urination increased thirst increased volume of pale, dilute urine irregular heartbeat loss of appetite mood changes muscle pain or cramps nausea or vomiting numbness or tingling in hands, feet, or lips shortness of breath Rare Arm, back, or jaw pain black, tarry stools bleeding gums blood in urine or stools bone pain burning, dry, or itching eyes changes in skin color chest pain chest tightness or heaviness convulsions discharge, excessive tearing feeling of warmth or heat flushing or redness of skin, especially on face and neck headache, sudden and severe increased need to urinate nervousness pain, tenderness, or swelling of foot or leg passing urine more often pinpoint red spots on skin pounding in the ears slow or fast heartbeat redness, pain, or swelling of eye, eyelid, or inner lining of eyelid swelling around the eyes or the whole face troubled breathing unusual bleeding or bruising weakness Symptoms of overdose Sleepiness slurred speech Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following

Drug Side Effects Summary

side effects continue or are bothersome or if you have any questions about them: Rare Acid or sour stomach belching continuing ringing or buzzing or other unexplained noise in ears excessive muscle tone fever hearing loss heartburn indigestion lack or loss of strength muscle stiffness muscle tension or tightness neck pain stomach discomfort, upset, or pain wheezing Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Cardene IV side effects

1896. This particular disease, Hypertension, interacts with the drug chlorothiazide and has the following

- a light-headed feeling, like you might pass out
- little or no urination
- severe pain in your upper stomach spreading to your back
- low blood cell counts--pale or yellowed skin, dark colored urine, fever, easy bruising or bleeding
- or signs of an electrolyte imbalance--dry mouth, increased thirst or urination, confusion, vomiting, constipation , muscle pain or weakness, leg cramps, dizziness , drowsiness, lack of energy, feeling restless, fast heartbeats, tingly feeling. Common side effects of chlorothiazide may include: dizziness, spinning sensation
- numbness or tingling
- diarrhea, constipation, stomach cramps
- blurred vision
- muscle spasm
- or impotence , sexual problems.

1897. This particular disease, Hypertension, interacts with the drug clevidipine and has the following

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Tell your caregivers right away if you have: chest pain
- pounding heartbeats or fluttering in your chest
- a light-headed feeling, like you might pass out
- severe dizziness or confusion
- vision problems
- shortness of breath
- or swelling in your hands or feet. Common side effects of clevidipine may include: headache
- or nausea and vomiting .

1898. This particular disease, Hypertension, interacts with the drug Cleviprex and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tell your caregivers right away if you have: chest pain
- pounding heartbeats or fluttering in your chest
- a light-headed feeling, like you might pass out
- severe dizziness or confusion
- vision problems
- shortness of breath
- or swelling in your hands or feet. Common Cleviprex side effects may include: headache
- or nausea and vomiting .

1899. This particular disease, Hypertension, interacts with the drug Conjupri and has the following

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. In rare cases, when you first start taking Conjupri, your angina may get worse or you could have a heart attack. Seek emergency medical attention or call your doctor right away if you have symptoms such as: chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating. Conjupri may cause serious side effects. Call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- worsening chest pain
- or a light-headed feeling, like you might pass out. Common side effects of Conjupri may include:
fast heartbeats
- swelling in your feet or ankles
- dizziness , drowsiness
- feeling tired
- stomach pain, nausea
- or flushing (warmth, redness, or tingly feeling).

1900. This particular disease, Hypertension, interacts with the drug Consensi and has the following

- Consensi can cause serious side effects, including: See Important information . liver problems, including liver failure worsening chest pain (angina) or heart attack, particularly in people with severe obstructive coronary artery disease heart failure swelling of your arms, legs, hands and feet (peripheral edema) is common with Consensi but can sometimes be serious. kidney problems, including kidney failure increased potassium levels (hyperkalemia) life-threatening allergic reactions life-threatening skin reactions low red blood cells (anemia) Your healthcare provider will monitor your blood pressure and do blood tests to check you for side effects during treatment. Consensi may cause fertility problems in females that is reversible when treatment is stopped. Talk to your healthcare provider if this is a concern for you. The most common side effects include: swelling of the arms, legs, hands, and feet joint swelling dizziness stomach pain diarrhea heartburn headache

Drug Side Effects Summary

frequent urination hot or warm feeling in your face (flushing) gas tiredness extreme sleepiness shortness of breath or trouble breathing chest pain weakness in one part or side of your body slurred speech swelling of the face or throat Stop treatment and call your healthcare provider right away if you get any of the following symptoms: nausea more tired or weaker than usual diarrhea itching your skin or eyes look yellow indigestion or stomach pain flu-like symptoms vomit blood there is blood in your bowel movement or it is black and sticky like tar unusual weight gain skin rash or blisters with fever swelling of the arms, legs, hands and feet These are not all the possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Consensi side effects

1901. This particular disease, Hypertension, interacts with the drug Corzide and has the following s

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Blurred vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain, tightness, or discomfort confusion cough difficulty breathing dilated neck veins dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position irregular breathing lightheadedness, dizziness, or fainting noisy breathing paleness or cold feeling in fingertips and toes slow or irregular heartbeat slurred speech sweating swelling of face, fingers, feet, or lower legs tingling or pain in fingers or toes when exposed to cold unusual tiredness or weakness weight gain Incidence not known Back or leg pains black, tarry stools bleeding gums bloating blood in urine or stools chills clay-colored stools cloudy urine cold sweats constipation coughing up blood dark urine difficulty swallowing drowsiness dry mouth fast heartbeat fever flushed, dry skin fruit-like breath odor general body swelling general feeling of discomfort or illness general tiredness and weakness headache hives, skin rash hoarseness increased hunger increased thirst increased urination

Drug Side Effects Summary

indigestion joint pain, stiffness, or swelling light-colored stools loss of appetite lower back or side pain muscle tremors nausea and vomiting nosebleeds pain in the muscles painful or difficult urination pains in stomach, side, or abdomen, possibly radiating to the back pale skin pinpoint red spots on the skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rapid, deep breathing redness, soreness, or itching skin restlessness sore throat sores, ulcers, or white spots on the lips or in the mouth sores, welting, or blisters stomach cramps sugar in the urine swollen or painful glands tenderness of salivary glands thickening of bronchial secretions trouble breathing unpleasant breath odor unusual bleeding or bruising unusual weight loss upper right stomach pain vomiting of blood yellow eyes or skin

Symptoms of overdose

Change in consciousness decreased urination diarrhea fast, slow, or shallow breathing gas heartburn increase in heart rate increased sweating indigestion loss of consciousness low blood pressure muscle cramps numbness, tingling, pain, or weakness in the hands or feet pale or blue lips, fingernails, or skin seizures sunken eyes trembling unusual drowsiness, dullness, or feeling of sluggishness unusual paleness weakness and heaviness of the legs wrinkled skin

Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them:

Less common

Change in behavior continuing ringing or buzzing or other unexplained noise in the ears decreased interest in sexual intercourse dry eyes or skin excess air or gas in stomach or intestines hair loss, thinning of hair hearing loss inability to have or keep an erection loss in sexual ability, desire, drive, or performance relaxed and calm sleepiness stomach soreness or discomfort stuffy nose weight loss

Incidence not known

Cracks in the skin discoloration of skin feeling of constant movement of self or surroundings increased sensitivity of skin to sunlight large, flat, blue or purplish patches in the skin loss of heat from the body muscle spasm, weakness, or restlessness

Drug Side Effects Summary

scaly skin sensation of spinning severe sunburn Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Corzide side effects

1902. This particular disease, Hypertension, interacts with the drug Corzide 40/5 and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Corzide 40/5 may cause serious side effects. Call your doctor at once if you have: fast, slow, or uneven heartbeat
- numbness or cold feeling in your hands and feet
- a light-headed feeling, like you might pass out
- little or no urination
- or signs of an electrolyte imbalance--dry mouth, increased thirst, mood changes, confusion, vomiting, muscle pain or weakness, lack of energy, fast heartbeats, little or no urine. Common side effects of Corzide 40/5 may include: dizziness
- or feeling tired.

1903. This particular disease, Hypertension, interacts with the drug Corzide 80/5 and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Corzide 80/5 may cause serious side effects. Call your doctor at once if you have: fast, slow, or uneven heartbeat
- numbness or cold feeling in your hands and feet
- a light-headed feeling, like you might pass out

Drug Side Effects Summary

- little or no urination
- or signs of an electrolyte imbalance--dry mouth, increased thirst, mood changes, confusion, vomiting, muscle pain or weakness, lack of energy, fast heartbeats, little or no urine. Common side effects of Corzide 80/5 may include: dizziness
- or feeling tired.

1904. This particular disease, Hypertension, interacts with the drug Demadex and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor right away if you have: a light-headed feeling, like you might pass out
- ringing in your ears, sudden hearing loss
- kidney problems--little or no urination, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- low magnesium level-- dizziness , irregular heartbeats, feeling jittery, muscle cramps, muscle spasms, cough or choking feeling
- low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- or low sodium level-- headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady. Common side effects of Demadex may include: increased urination
- or headache.

1905. This particular disease, Hypertension, interacts with the drug Diuril and has the following side effects:

- a light-headed feeling, like you might pass out
- little or no urination

Drug Side Effects Summary

- severe pain in your upper stomach spreading to your back
- low blood cell counts--pale or yellowed skin, dark colored urine, fever, easy bruising or bleeding
- or signs of an electrolyte imbalance--dry mouth, increased thirst or urination, confusion, vomiting, constipation, muscle pain or weakness, leg cramps, dizziness, drowsiness, lack of energy, feeling restless, fast heartbeats, tingly feeling. Common side effects of Diuril may include: dizziness, spinning sensation
- numbness or tingling
- diarrhea, constipation, stomach cramps
- blurred vision
- muscle spasm
- or impotence, sexual problems.

1906. This particular disease, Hypertension, interacts with the drug Diuril Sodium and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: Incidence not known
Abdominal or stomach pain back, leg, or stomach pains black, tarry stools bleeding gums blisters on skin blood in urine or stools bloody or cloudy urine blue lips and fingernails blurred vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain chills clay-colored stools cold sweats confusion constipation cough or hoarseness coughing that sometimes produces a pink frothy sputum coughing up blood darkened urine decrease in urine output or decrease in urine-concentrating ability decreased frequency or amount of urine diarrhea difficult, fast, or noisy breathing, sometimes with wheezing difficulty swallowing dizziness, faintness, or lightheadedness when getting up from a lying or sitting position suddenly dry mouth fast heartbeat feeling of constant movement of self or surroundings fever flushed, dry skin fruit-like

Drug Side Effects Summary

breath odor general feeling of discomfort or illness general feeling of tiredness or weakness headache hives increased blood pressure increased hunger increased sensitivity of skin to sunlight increased sweating increased thirst increased urination indigestion itching joint pain, stiffness, or swelling loss of appetite lower back or side pain muscle cramps or pain nausea nosebleeds numbness, tingling, pain, or weakness in hands or feet pain in joints or muscles painful or difficult urination pains in stomach, side, or abdomen, possibly radiating to the back pale skin pinpoint red spots on skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue redness or other discoloration of skin redness, soreness, or itching skin seizures sensation of spinning severe sunburn shortness of breath skin rash sores, ulcers, or white spots on lips or in mouth sores, welting, or blisters sugar in the urine sweating swelling in legs and ankles swelling of face, fingers, or lower legs swollen or painful glands tenderness of salivary glands thickening of bronchial secretions tightness in chest trembling troubled breathing unexplained weight loss unpleasant breath odor unusual bleeding or bruising unusual tiredness or weakness unusual weight loss vomiting of blood weight gain yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Blistering, peeling, or loosening of skin cracks in the skin cramping decreased interest in sexual intercourse difficulty having a bowel movement (stool) hair loss, thinning of hair inability to have or keep an erection loss in sexual ability, desire, drive, or performance loss of heat from the body muscle spasm pinpoint red or purple spots on skin red, irritated eyes red, swollen skin restlessness scaly skin weakness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Diuril Sodium

Drug Side Effects Summary

side effects

1907. This particular disease, Hypertension, interacts with the drug Dutoprol and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

- itching

- red, swollen, blistered, or peeling skin with or without fever

- wheezing

- tightness in the chest or throat

- trouble breathing, swallowing, or talking

- unusual hoarseness

- or swelling of the mouth, face, lips, tongue, or throat. Signs of high blood sugar like confusion, feeling sleepy, more thirst, more hungry, passing urine more often, flushing, fast breathing, or breath that smells like fruit. Signs of fluid and electrolyte problems like mood changes, confusion, muscle pain or weakness, a heartbeat that does not feel normal, very bad dizziness or passing out, fast heartbeat, more thirst, seizures , feeling very tired or weak, not hungry, unable to pass urine or change in the amount of urine produced, dry mouth, dry eyes, or very bad upset stomach or throwing up. Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain. Chest pain that is new or worse. Slow heartbeat. Low mood (depression). Feeling confused. Memory problems or loss. Shortness of breath, a big weight gain, or swelling in the arms or legs. Any skin change. This medicine can cause certain eye problems. If left untreated, this can lead to lasting eyesight loss. If eye problems happen, signs like

Drug Side Effects Summary

change in eyesight or eye pain most often happen within hours to weeks of starting Dutoprol (metoprolol and hydrochlorothiazide extended-release tablets). Call your doctor right away if you have these signs. Dutoprol side effects

1908. This particular disease, Hypertension, interacts with the drug Edarbi and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: little or no urination
- or a light-headed feeling, like you might pass out. Common Edarbi side effects may include: diarrhea.

1909. This particular disease, Hypertension, interacts with the drug Edarbyclor and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: feeling like you might pass out
- little or no urination
- swelling, weight gain, feeling short of breath
- or signs of electrolyte imbalance - dry mouth, extreme thirst, weakness, drowsiness, restless feeling, confusion, increased or decreased urination, nausea and vomiting , constipation , muscle pain or weakness, fast heart rate , or seizure (convulsions). Common Edarbyclor side effects may include: dizziness
- or feeling tired.

Drug Side Effects Summary

1910. This particular disease, Hypertension, interacts with the drug enalapril / hydrochlorothiazide .

- a light-headed feeling, like you might pass out
- jaundice (yellowing of the skin or eyes)
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- fever, chills, flu-like symptoms, sore throat, mouth sores
- signs of an electrolyte imbalance--leg cramps, constipation , muscle pain or weakness, irregular heartbeats, feeling jittery, increased thirst or urination, numbness or tingling, loss of movement
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or new or worsening symptoms of lupus--joint pain, and a skin rash on your cheeks or arms that worsens in sunlight. Common side effects of hydrochlorothiazide and enalapril may include:
dizziness
- cough
- headache
- or nausea .

1911. This particular disease, Hypertension, interacts with the drug eprosartan and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Burning or painful urination or changes in urinary frequency cough fever sore throat Rare Dizziness, lightheadedness, or fainting swollen face, lips, limbs, or tongue Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common or

Drug Side Effects Summary

rare Abdominal or stomach pain joint pain unusual tiredness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Eprosartan side effects

1912. This particular disease, Hypertension, interacts with the drug esmolol and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tell your caregivers right away if you have: a light-headed feeling, like you might pass out
- very slow heartbeats
- swelling in your hands or feet
- wheezing, chest tightness, feeling short of breath
- weak or shallow breathing
- pain, swelling, irritation, bruising, or skin changes around the IV needle
- cold feeling in your hands and feet
- high potassium-- nausea , slow or unusual heart rate, weakness, loss of movement
- or low blood sugar -- headache , hunger, sweating, irritability, dizziness , nausea, fast heart rate , and feeling anxious or shaky. Common side effects of esmolol may include: increased sweating
- nausea
- dizziness, drowsiness
- or low blood pressure .

1913. This particular disease, Hypertension, interacts with the drug fosinopril / hydrochlorothiazide and has the following side effects:

- hives

Drug Side Effects Summary

- severe stomach pain
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Hydrochlorothiazide and fosinopril may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- pale skin, easy bruising or bleeding
- fever, swollen gums, painful mouth sores, pain when swallowing, cold or flu symptoms
- high potassium-- nausea , slow or unusual heart rate, weakness, loss of movement
- low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- low levels of sodium-- headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady
- or severe skin reaction--fever, sore throat , swelling in your face or tongue, burning in your eyes, skin pain, followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of hydrochlorothiazide and fosinopril may include: headache, dizziness
- cough
- or tired feeling.

1914. This particular disease, Hypertension, interacts with the drug guanabenz and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: Signs and symptoms of overdose Dizziness (severe) irritability nervousness pinpoint pupils slow heartbeat unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care

Drug Side Effects Summary

professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Dizziness drowsiness dryness of mouth weakness Less common or rare Decreased sexual ability headache nausea After you stop using guanabenz, it may still produce some side effects that need attention. During this period of time, check with your doctor immediately if you notice the following side effects: Anxiety or tenseness chest pain fast or irregular heartbeat headache increased salivation increase in sweating nausea or vomiting nervousness or restlessness shaking or trembling of hands or fingers stomach cramps trouble in sleeping Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Guanabenz side effects

1915. This particular disease, Hypertension, interacts with the drug hydrochlorothiazide / methyldopa

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Hydrochlorothiazide and methyldopa may cause serious side effects. Call your doctor at once if you have: pale or yellowed skin, dark colored urine, fever, confusion or weakness
- a light-headed feeling, like you might pass out
- severe pain in your upper stomach spreading to your back, nausea and vomiting , fast heart rate
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- uncontrolled muscle movements
- unusual thoughts or behavior
- numbness , tingling, weakness

Drug Side Effects Summary

- chest pain
- easy bruising, unusual bleeding
- signs of an electrolyte imbalance--dry mouth, leg cramps, increased thirst or urination, mood changes, confusion, numbness or tingling, constipation , vomiting, bone pain, muscle pain or weakness, lack of energy, irregular heartbeats, fluttering in your chest, little or no urine
- or severe skin reaction--fever, sore throat , swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of hydrochlorothiazide and methyldopa may include: abnormal liver function tests
- headache , drowsiness, weakness
- dizziness
- swelling, weight gain
- muscle or joint pain, muscle spasms
- or stuffy nose.

1916. This particular disease, Hypertension, interacts with the drug hydrochlorothiazide / metoprolol

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Hydrochlorothiazide and metoprolol may cause serious side effects. Call your doctor at once if you have: bronchospasm (wheezing, chest tightness, trouble breathing)
- slow heart rate, weak pulse, fainting, slow breathing (breathing may stop)
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling

Drug Side Effects Summary

- or low sodium level -- headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady. Common side effects of hydrochlorothiazide and metoprolol may include: tired feeling
- or stuffy nose, sore throat .

1917. This particular disease, Hypertension, interacts with the drug hydrochlorothiazide / moexipril

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Blurred vision confusion decreased urination dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position dry mouth fast, slow, or irregular heartbeat muscle cramps or pain numbness, tingling, pain, or weakness in the hands or feet rapid breathing seizures sunken eyes sweating thirst trembling unusual tiredness or weakness weakness and heaviness of the legs wrinkled skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Dry cough Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Hydrochlorothiazide / moexipril side effects

1918. This particular disease, Hypertension, interacts with the drug hydrochlorothiazide / quinapril

- a light-headed feeling, like you might pass out
- blurred vision, tunnel vision, eye pain , or seeing halos around lights

Drug Side Effects Summary

- swelling, weight gain, shortness of breath
- sudden weakness or ill feeling, fever, chills, sore throat, mouth sores, trouble swallowing
- high potassium-- nausea , tingly feeling, chest pain, irregular heartbeats, loss of movement
- low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- or low levels of sodium-- headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady. Common side effects of hydrochlorothiazide and quinapril may include: cough
- headache, dizziness
- tired feeling
- or muscle pain.

1919. This particular disease, Hypertension, interacts with the drug hydrochlorothiazide / spironolactone

- skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Hydrochlorothiazide and spironolactone may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- breast swelling and tenderness
- low blood cell counts--easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath
- high potassium level--nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- or low sodium level -- headache , confusion, slurred speech, severe weakness, vomiting, loss of

Drug Side Effects Summary

coordination, feeling unsteady. Common side effects of hydrochlorothiazide and spironolactone may include: drowsiness, dizziness, lack of energy

- leg cramps
- weakness
- feeling like you might pass out
- severe pain in your upper stomach spreading to your back, nausea and vomiting
- or electrolyte imbalance.

1920. This particular disease, Hypertension, interacts with the drug Hytrin and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have any of these serious side effects: fast or pounding heartbeats or fluttering in your chest
- feeling like you might pass out
- swelling in your hands, ankles, or feet
- or penis erection that is painful or lasts 4 hours or longer. Less serious Hytrin side effects may include: mild dizziness
- weakness, drowsiness
- blurred vision
- nausea
- or headache.

1921. This particular disease, Hypertension, interacts with the drug Inspra and has the following side effects:

- hives
- severe stomach pain

Drug Side Effects Summary

- difficulty breathing
- swelling of your face, lips, tongue, or throat. Inspra may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- diarrhea , vomiting
- little or no urination
- fast or irregular heartbeats
- trouble breathing
- swelling in your feet or lower legs
- or high potassium-- nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common side effects of Inspra may include: high potassium
- headache
- or dizziness .

1922. This particular disease, Hypertension, interacts with the drug isradipine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Isradipine may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- pounding heartbeats or fluttering in your chest
- flushing (warmth, redness, or tingly feeling)
- tiredness
- or heart problems--swelling, rapid weight gain, feeling short of breath. Common side effects of isradipine may include: flushing
- headache , dizziness
- swelling

Drug Side Effects Summary

- fast heartbeats
- or upset stomach .

1923. This particular disease, Hypertension, interacts with the drug levamlodipine and has the follo

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, when you first start taking levamlodipine, your angina may get worse or you could have a heart attack. Seek emergency medical attention or call your doctor right away if you have symptoms such as: chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating. Levamlodipine may cause serious side effects. Call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- worsening chest pain
- or a light-headed feeling, like you might pass out. Common side effects of levamlodipine may include: fast heartbeats
- swelling in your feet or ankles
- dizziness , drowsiness
- feeling tired
- stomach pain, nausea
- or flushing (warmth, redness, or tingly feeling).

1924. This particular disease, Hypertension, interacts with the drug Lopressor HCT and has the foll

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Chest pain or discomfort

Drug Side Effects Summary

decreased urine dry mouth increased thirst lightheadedness, dizziness, or fainting loss of appetite mood changes muscle pain or cramps nausea or vomiting numbness or tingling in the hands, feet, or lips seizures slow or irregular heartbeat unusual tiredness or weakness Less common Ankle, knee, or great toe joint pain chest tightness decreased ability to exercise difficult or labored breathing joint stiffness or swelling lower back or side pain swelling of the face, fingers, feet, or lower legs Incidence not known Persistent non-healing sore pink growth reddish patch or irritated area shiny bump white, yellow or waxy scar-like area Symptoms of overdose Blurred vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chemical imbalance in the blood cold, clammy skin coma confusion cough dilated neck veins dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position drowsiness fast or pounding heartbeat or pulse frequent urination headache increased volume of pale, dilute urine irregular breathing irritability noisy breathing not able to pass urine pain or aching in the lower legs pain or weakness in the hands or feet sweating trembling unconsciousness very drowsy or sleepy weak pulse weakness and heaviness of the legs weight gain Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Chills diarrhea feeling of constant movement of self or surroundings fever general feeling of discomfort or illness muscle aches nightmares runny nose sensation of spinning shivering sleepiness sore throat trouble sleeping unusual drowsiness, dullness, or feeling of sluggishness Less common Continuing ringing or buzzing or other unexplained noise in the ears decreased interest in sexual intercourse difficulty having a bowel movement (stool) earache hearing loss inability to have or keep an erection loss in sexual ability, desire, drive, or performance pinpoint red or purple spots on the skin weight loss Other side effects not listed may also occur in some patients.

Drug Side Effects Summary

If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Lopressor HCT side effects

1925. This particular disease, Hypertension, interacts with the drug Lotensin HCT and has the following

- a light-headed feeling, like you might pass out
- jaundice (yellowing of the skin or eyes)
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- fever, chills, flu-like symptoms, sore throat, mouth sores
- signs of an electrolyte imbalance--leg cramps, constipation , muscle pain or weakness, irregular heartbeats, feeling jittery, increased thirst or urination, numbness or tingling, loss of movement
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or new or worsening symptoms of lupus--joint pain, and a skin rash on your cheeks or arms that worsens in sunlight. Common side effects of Lotensin HCT may include: dizziness
- cough
- headache
- or feeling tired.

1926. This particular disease, Hypertension, interacts with the drug Maxzide-25 and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching

Drug Side Effects Summary

- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of fluid and electrolyte problems like mood changes, confusion, muscle pain or weakness, a heartbeat that does not feel normal, very bad dizziness or passing out, fast heartbeat , more thirst, seizures , feeling very tired or weak, not hungry, unable to pass urine or change in the amount of urine produced, dry mouth, dry eyes, or very bad upset stomach or throwing up. Signs of too much acid in the blood (acidosis) like confusion
- fast breathing
- fast heartbeat
- a heartbeat that does not feel normal
- very bad stomach pain, upset stomach, or throwing up
- feeling very sleepy
- shortness of breath
- or feeling very tired or weak. Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain. Signs of a pancreas problem (pancreatitis) like very bad stomach pain, very bad back pain , or very bad upset stomach or throwing up. Back pain, belly pain, or blood in the urine. May be signs of a kidney stone. Dark urine or yellow skin or eyes. A burning, numbness , or tingling feeling that is not normal. Slow heartbeat . Restlessness. Shortness of breath. Chest pain or pressure or a fast heartbeat. Not able to get or keep an erection. Mood changes. Any skin change. This medicine can cause certain eye problems. If left untreated, this can lead to lasting eyesight loss. If eye problems happen, signs like change in eyesight or eye pain most often happen within hours to weeks of starting Maxzide-25

Drug Side Effects Summary

(hydrochlorothiazide and triamterene). Call your doctor right away if you have these signs.

Maxzide-25 side effects

1927. This particular disease, Hypertension, interacts with the drug mecamlamine and has the fol

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Mecamlamine may cause serious side effects. Call your doctor at once if you have: tremors or jerky muscle movements
- numbness , tingling, burning pain
- painful or difficult urination
- ongoing diarrhea with stomach bloating
- confusion, thinking problems
- a light-headed feeling, like you might pass out
- a seizure
- or dry cough, wheezing, feeling short of breath (even while lying down). Common side effects of mecamlamine may include: dizziness or fainting
- nausea, vomiting, loss of appetite
- constipation
- dry mouth, swollen tongue
- feeling weak or tired
- blurred vision
- or impotence , decreased interest in sex.

1928. This particular disease, Hypertension, interacts with the drug Midamor and has the following

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop taking Midamor and call your doctor at once if you have: increased thirst, decreased urination
- heavy sweating, or hot and dry skin
- tremors, confusion, loss of consciousness
- jaundice (yellowing of the skin or eyes)
- high potassium--tiredness, numbness or tingling, slow or unusual heart rate, muscle weakness or limp feeling
- or low levels of sodium in the body-- headache , confusion, slurred speech, severe weakness, vomiting , loss of coordination, feeling unsteady. Common side effects of Midamor may include: nausea , stomach pain, gas, loss of appetite
- headache
- or rash.

1929. This particular disease, Hypertension, interacts with the drug Minipress and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Minipress may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- pounding heartbeats or fluttering in your chest
- new or worsening chest pain
- or upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Call your doctor or seek emergency medical attention if your erection is painful or lasts longer than 4 hours. A prolonged erection (priapism) can damage the penis. Common side effects of Minipress may include: dizziness, drowsiness

Drug Side Effects Summary

- headache
- feeling weak or tired
- or nausea .

1930. This particular disease, Hypertension, interacts with the drug moexipril and has the following

- hives
- severe stomach pain
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Moexipril may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- sudden weakness or ill feeling, fever, chills, sore throat , painful mouth sores, cough, trouble breathing
- little or no urinating
- or high potassium-- nausea , slow or unusual heart rate, weakness, loss of movement. Common side effects of moexipril may include: cough, runny or stuffy nose
- headache , dizziness , tired feeling
- increased urination
- diarrhea
- muscle pain
- or mild skin itching or rash.

1931. This particular disease, Hypertension, interacts with the drug nebivolol / valsartan and has th

- hives
- trouble breathing or swallowing
- stomach pain, vomiting

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Nebivolol and valsartan may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- very slow heartbeats
- chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating
- shortness of breath (even with mild exertion)
- swelling or numbness in your legs or feet
- rapid weight gain
- or high potassium--nausea, slow or unusual heart rate, weakness, loss of movement. Common side effects of nebivolol and valsartan may include: slow heartbeats.

1932. This particular disease, Hypertension, interacts with the drug nicardipine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Nicardipine may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- pounding heartbeats or fluttering in your chest
- or severe or ongoing chest pain. Common side effects of nicardipine may include: headache , dizziness
- swelling in your feet
- weakness
- flushing (warmth, redness, or tingly feeling)
- or nausea .

1933. This particular disease, Hypertension, interacts with the drug nisoldipine and has the following side effects:

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Nisoldipine may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- chest pain or pressure, pain spreading to your jaw or shoulder
- swelling in your hands or feet
- flushing (sudden warmth, redness, or tingly feeling)
- fast heartbeats
- or sudden numbness or weakness, problems with vision or speech. You may have new or worsening chest pain when you first start taking nisoldipine or whenever your dose is changed.

Common side effects of nisoldipine may include: swelling

- headache
- dizziness
- or flushing.

1934. This particular disease, Hypertension, interacts with the drug pindolol and has the following s

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Pindolol may cause serious side effects. Call your doctor at once if you have: shortness of breath (even with mild exertion), swelling, rapid weight gain
- slow or uneven heartbeats
- a light-headed feeling, like you might pass out
- or unusual changes in mood or behavior. Common side effects of pindolol may include: dizziness
- muscle or joint pain
- sleep problems (insomnia)
- or feeling tired.

Drug Side Effects Summary

1935. This particular disease, Hypertension, interacts with the drug Prestalia and has the following

- hives
- severe stomach pain
- difficulty breathing
- swelling of your face, lips, tongue, or throat. You may be more likely to have an allergic reaction if you are African-American. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- swelling in your hands or feet, rapid weight gain
- little or no urinating
- new or worsened chest pain
- jaundice (yellowing of the skin or eyes)
- or high potassium nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common Prestalia side effects may include: cough
- headache , dizziness
- or swelling in your hands or feet.

1936. This particular disease, Hypertension, interacts with the drug Sular and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Sular may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- chest pain or pressure, pain spreading to your jaw or shoulder
- swelling in your hands or feet
- flushing (sudden warmth, redness, or tingly feeling)
- fast heartbeats

Drug Side Effects Summary

- or sudden numbness or weakness, problems with vision or speech. You may have new or worsening chest pain when you first start taking Sular or whenever your dose is changed. Common side effects of Sular may include: swelling

- headache
- dizziness
- or flushing.

1937. This particular disease, Hypertension, interacts with the drug Tarka and has the following side effects:

- hives
- severe stomach pain
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Tarka may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- upper stomach pain, and jaundice (yellowing of the skin or eyes)
- chest pain, fast, slow, or uneven heart rate
- high potassium level-- nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- low white blood cell counts--fever, chills, feeling light-headed, mouth sores, skin sores, sore throat , cough, trouble breathing
- or fluid build-up in or around the lungs--pain when you breathe, feeling short of breath while lying down, wheezing, gasping for breath, cough with foamy mucus, cold, clammy skin, anxiety , rapid heartbeats. Common side effects of Tarka may include: cough
- cold symptoms such as stuffy nose, sneezing, sore throat
- headache , dizziness

Drug Side Effects Summary

- chest pain, back pain
- constipation
- feeling tired
- or abnormal liver function tests .

1938. This particular disease, Hypertension, interacts with the drug Tekturna HCT and has the following

- a light-headed feeling, like you might pass out
- little or no urination
- eye pain , vision problems
- high potassium level - nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- low potassium level - leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- low sodium level - headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady
- or lupus-like symptoms - muscle or joint pain, fever, and a rash or patchy skin color that worsens in sunlight. Common Tekturna HCT side effects may include: dizziness, tired feeling
- high potassium
- cough
- joint pain
- diarrhea
- or flu symptoms (fever, chills, body aches).

1939. This particular disease, Hypertension, interacts with the drug Tenoretic and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and

Drug Side Effects Summary

sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of fluid and electrolyte problems like mood changes, confusion, muscle pain or weakness, a heartbeat that does not feel normal, very bad dizziness or passing out, fast heartbeat, more thirst, seizures , feeling very tired or weak, not hungry, unable to pass urine or change in the amount of urine produced, dry mouth, dry eyes, or very bad upset stomach or throwing up. Signs of high blood sugar like confusion, feeling sleepy, more thirst, more hungry, passing urine more often, flushing, fast breathing, or breath that smells like fruit. Signs of a pancreas problem (pancreatitis) like very bad stomach pain, very bad back pain , or very bad upset stomach or throwing up. Feeling cold. Very bad dizziness or passing out. Slow heartbeat. Restlessness. Shortness of breath, a big weight gain, or swelling in the arms or legs. Low mood (depression). Fever, chills, or sore throat
- any unexplained bruising or bleeding
- or feeling very tired or weak. Yellow skin or eyes. A burning, numbness , or tingling feeling that is not normal. Tenoretic side effects

1940. This particular disease, Hypertension, interacts with the drug Tenoretic 50 and has the follow

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tenoretic 50 may cause serious side effects. Call your doctor at once if you have: new or worsening chest pain
- slow or uneven heartbeats
- a light-headed feeling, like you might pass out
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- cold feeling in your hands and feet
- jaundice (yellowing of the skin or eyes)
- low sodium level -- headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady
- or low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of Tenoretic 50 may include: slow heartbeats
- feeling weak or tired
- dizziness , drowsiness
- numbness or tingling
- pale skin, bruising or bleeding
- depression
- or upset stomach , loss of appetite, constipation.

1941. This particular disease, Hypertension, interacts with the drug Tenoretic 100 and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tenoretic 100 may cause serious side effects. Call

Drug Side Effects Summary

your doctor at once if you have: new or worsening chest pain

- slow or uneven heartbeats
- a light-headed feeling, like you might pass out
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- cold feeling in your hands and feet
- jaundice (yellowing of the skin or eyes)
- low sodium level -- headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady
- or low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of Tenoretic 100 may include: slow heartbeats
- feeling weak or tired
- dizziness , drowsiness
- numbness or tingling
- pale skin, bruising or bleeding
- depression
- or upset stomach , loss of appetite, constipation.

1942. This particular disease, Hypertension, interacts with the drug Trandate and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Trandate may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- slow heart rate, weak pulse, fainting, slow breathing (breathing may stop)
- shortness of breath (even with mild exertion), swelling, rapid weight gain

Drug Side Effects Summary

- severe headache , blurred vision, pounding in your neck or ears
- or liver problems--loss of appetite, stomach pain (upper right side), flu-like symptoms, itching, dark urine, jaundice (yellowing of the skin or eyes). Severe dizziness or fainting may be more likely in older adults. Common side effects of Trandate may include: dizziness, drowsiness, tiredness
- nausea , vomiting
- sudden warmth, skin redness, sweating
- numbness
- or tingly feeling in your scalp.

1943. This particular disease, Hypertension, interacts with the drug trandolapril / verapamil and has

- hives
- severe stomach pain
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Trandolapril and verapamil may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- upper stomach pain, and jaundice (yellowing of the skin or eyes)
- chest pain, fast, slow, or uneven heart rate
- high potassium level-- nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- low white blood cell counts--fever, chills, feeling light-headed, mouth sores, skin sores, sore throat , cough, trouble breathing
- or fluid build-up in or around the lungs--pain when you breathe, feeling short of breath while lying down, wheezing, gasping for breath, cough with foamy mucus, cold, clammy skin, anxiety , rapid heartbeats. Common side effects of trandolapril and verapamil may include: cough

Drug Side Effects Summary

- cold symptoms such as stuffy nose, sneezing, sore throat
- headache , dizziness
- chest pain, back pain
- constipation
- feeling tired
- or abnormal liver function tests .

1944. This particular disease, Hypertension, interacts with the drug trandolapril and has the following

- hives
- severe stomach pain
- difficulty breathing
- swelling of your face, lips, tongue, or throat. You may be more likely to have an allergic reaction if you are African-American. Trandolapril may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- upper stomach pain, and jaundice (yellowing of the skin or eyes)
- chest pain, pounding heartbeats or fluttering in your chest
- high potassium level-- nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- low white blood cell counts--fever, chills, feeling light-headed, mouth sores, skin sores, sore throat , cough, trouble breathing
- or lung problems-- anxiety , sweating, pale skin, severe shortness of breath, wheezing, gasping for breath, cough with foamy mucus. Common side effects of trandolapril may include: cough
- dizziness
- or diarrhea.

Drug Side Effects Summary

1945. This particular disease, Hypertension, interacts with the drug Tribenzor and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: worsening chest pain
- a light-headed feeling, like you might pass out
- an unusual skin rash
- pain or burning when you urinate
- severe or ongoing diarrhea with weight loss
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or signs of an electrolyte imbalance - increased thirst or urination, confusion, vomiting, constipation , muscle pain or weakness, leg cramps, lack of energy, irregular heartbeats, numbness or tingling, loss of movement. Common Tribenzor side effects may include: dizziness
- headache , tired feeling
- swelling in your hands or feet
- nausea , diarrhea
- runny or stuffy nose, sore throat
- muscle spasms
- swollen joints
- or painful urination.

1946. This particular disease, Hypertension, interacts with the drug Twynsta and has the following

- hives

Drug Side Effects Summary

- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, Twynsta can cause a condition that results in the breakdown of skeletal muscle tissue, leading to kidney failure. Call your doctor right away if you have unexplained muscle pain, tenderness, or weakness especially if you also have fever, unusual tiredness, or dark colored urine. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- new or worsened chest pain
- pounding heartbeats or fluttering in your chest
- tremors, muscle stiffness or twitching
- or high potassium - nausea, slow or unusual heart rate, weakness, loss of movement. Common Twynsta side effects may include: swelling in your hands or feet
- fast or pounding heartbeats
- dizziness, drowsiness
- flushing (warmth, redness, or tingly feeling)
- back pain
- or nausea, diarrhea, stomach pain.

1947. This particular disease, Hypertension, interacts with the drug Vaseretic and has the following

- a light-headed feeling, like you might pass out
- jaundice (yellowing of the skin or eyes)
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- fever, chills, flu-like symptoms, sore throat, mouth sores
- signs of an electrolyte imbalance--leg cramps, constipation , muscle pain or weakness, irregular heartbeats, feeling jittery, increased thirst or urination, numbness or tingling, loss of movement

Drug Side Effects Summary

- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or new or worsening symptoms of lupus--joint pain, and a skin rash on your cheeks or arms that worsens in sunlight. Common side effects of Vaseretic 10-25 may include: dizziness
- cough
- headache
- or nausea .

1948. This particular disease, Hypertension, interacts with the drug Vecamyl and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Vecamyl may cause serious side effects. Call your doctor at once if you have: tremors or jerky muscle movements
- numbness , tingling, burning pain
- painful or difficult urination
- ongoing diarrhea with stomach bloating
- confusion, thinking problems
- a light-headed feeling, like you might pass out
- a seizure
- or dry cough, wheezing, feeling short of breath (even while lying down). Common side effects of Vecamyl may include: dizziness or fainting
- nausea, vomiting, loss of appetite
- constipation
- dry mouth, swollen tongue
- feeling weak or tired
- blurred vision

Drug Side Effects Summary

- or impotence , decreased interest in sex.

1949. This particular disease, Hypertension, interacts with the drug Zaroxolyn and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Zaroxolyn may cause serious side effects. Call your doctor at once if you have: chest pain
- pounding heartbeats or fluttering in your chest
- easy bruising or bleeding
- a light-headed feeling, like you might pass out
- liver problems-- nausea , upper stomach pain, itching, tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- low levels of sodium in the body-- headache , confusion, slurred speech, severe weakness, vomiting, loss of coordination, feeling unsteady
- other signs of an electrolyte imbalance--dry mouth, nausea, vomiting, weakness, drowsiness, lack of energy, muscle pain, little or no urination, or feeling restless
- or severe skin reaction--fever, sore throat , swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of Zaroxolyn may include: dizziness, spinning sensation
- drowsiness, tiredness
- depressed mood
- muscle or joint pain

Drug Side Effects Summary

- numbness or tingly feeling
- nausea, stomach pain, loss of appetite
- or diarrhea , constipation.

1950. This particular disease, Hypothyroidism, interacts with the drug levothyroxine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Levothyroxine may cause serious side effects. Call your doctor at once if you have: fast or irregular heartbeats
- chest pain, pain spreading to your jaw or shoulder
- shortness of breath
- fever, hot flashes , sweating
- tremors, or if you feel unusually cold
- weakness, tiredness, sleep problems (insomnia)
- memory problems, feeling depressed or irritable
- headache, leg cramps, muscle aches
- feeling nervous or irritable
- dryness of your skin or hair, hair loss
- irregular menstrual periods
- or vomiting , diarrhea , appetite changes, weight changes. Certain side effects may be more likely in older adults. Common side effects of levothyroxine may include: chest pain, irregular heartbeats
- shortness of breath
- headache, leg cramps, muscle pain or weakness
- tremors, feeling nervous or irritable, trouble sleeping
- increased appetite

Drug Side Effects Summary

- feeling hot
- weight loss
- changes in your menstrual periods
- diarrhea
- or skin rash , partial hair loss.

1951. This particular disease, Hypothyroidism, interacts with the drug Synthroid and has the follow

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: fast or irregular heartbeats
- chest pain, pain spreading to your jaw or shoulder
- shortness of breath
- fever, hot flashes , sweating
- tremors, or if you feel unusually cold
- weakness, tiredness, sleep problems (insomnia)
- memory problems, feeling depressed or irritable
- headache, leg cramps, muscle aches
- feeling nervous or irritable
- dryness of your skin or hair, hair loss
- irregular menstrual periods
- or vomiting , diarrhea , appetite changes, weight changes. Certain side effects may be more likely in older adults. Common Synthroid side effects may include: chest pain, irregular heartbeats
- shortness of breath
- headache, leg cramps, muscle pain or weakness

Drug Side Effects Summary

- tremors, feeling nervous or irritable, trouble sleeping
- increased appetite
- feeling hot
- weight loss
- changes in your menstrual periods
- diarrhea
- or skin rash , partial hair loss.

1952. This particular disease, Hypothyroidism, interacts with the drug Armour Thyroid and has the

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common side effects may include temporary hair loss (especially in children).

1953. This particular disease, Hypothyroidism, interacts with the drug Levoxyl and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: fast or irregular heartbeats
- chest pain, pain spreading to your jaw or shoulder
- shortness of breath
- fever, hot flashes , sweating
- tremors, or if you feel unusually cold
- weakness, tiredness, sleep problems (insomnia)
- memory problems, feeling depressed or irritable

Drug Side Effects Summary

- headache, leg cramps, muscle aches
- feeling nervous or irritable
- dryness of your skin or hair, hair loss
- irregular menstrual periods
- or vomiting , diarrhea , appetite changes, weight changes. Certain side effects may be more likely in older adults. Common Levoxyl side effects may include: chest pain, irregular heartbeats
- shortness of breath
- headache, leg cramps, muscle pain or weakness
- tremors, feeling nervous or irritable, trouble sleeping
- increased appetite
- feeling hot
- weight loss
- changes in your menstrual periods
- diarrhea
- or skin rash , partial hair loss.

1954. This particular disease, Hypothyroidism, interacts with the drug Cytomel and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Cytomel and call your doctor if you have symptoms of thyroid toxicity , such as chest pain, fast or pounding heartbeats, feeling hot or nervous, or sweating more than usual. Also call your doctor at once if you have: weak or shallow breathing
- confusion
- a feeling of being cold all the time

Drug Side Effects Summary

- changes in weight or appetite
- chest pressure spreading to your jaw or shoulder
- a light-headed feeling, like you might pass out
- severe headaches, ringing in your ears, dizziness , vision problems, pain behind your eyes
- a seizure
- or (in a child or teenager) pain, stiffness, and trouble moving the legs. Common side effects of Cytomel may include: trouble breathing
- headache
- tremors, feeling nervous or irritable
- muscle weakness
- increased appetite
- diarrhea
- irregular menstrual periods
- weight loss
- feeling hot
- rash
- or sleep problems (insomnia).

1955. This particular disease, Hypothyroidism, interacts with the drug Tirosint and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Chest pain, discomfort, or tightness decreased urine output difficult or labored breathing difficulty with swallowing dilated neck veins extreme fatigue fainting fast, slow, irregular, pounding, or racing heartbeat or pulse fever heat intolerance hives or welts, skin itching, rash, or redness irregular breathing irritability menstrual

Drug Side Effects Summary

changes nausea pain or discomfort in the arms, jaw, back, or neck sweating swelling of the eyes, face, lips, throat, or tongue tremors Rare Blurred or double vision dizziness eye pain lack or slowing of normal growth in children limp or walk favoring one leg pain in the hip or knee seizures severe headache Symptoms of overdose Change in consciousness cold, clammy skin confusion disorientation fast or weak pulse lightheadedness loss of consciousness sudden headache sudden loss of coordination sudden slurring of speech Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Change in appetite crying diarrhea false or unusual sense of well-being fear or nervousness feeling not well or unhappy feeling of discomfort feeling of warmth feeling things are not real feelings of suspicion and distrust hair loss headache increased appetite mental depression muscle weakness quick to react or overreact emotionally rapidly changing moods redness of the face, neck, arms, and occasionally, upper chest restlessness stomach cramps trouble getting pregnant trouble sitting still unusual tiredness or weakness vomiting weight gain or loss Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Tirosint side effects

1956. This particular disease, Hypothyroidism, interacts with the drug Euthyrox and has the followi

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: fast or irregular heartbeats

Drug Side Effects Summary

- chest pain, pain spreading to your jaw or shoulder
- shortness of breath
- fever, hot flashes , sweating
- tremors, or if you feel unusually cold
- weakness, tiredness, sleep problems (insomnia)
- memory problems, feeling depressed or irritable
- headache, leg cramps, muscle aches
- feeling nervous or irritable
- dryness of your skin or hair, hair loss
- irregular menstrual periods
- or vomiting , diarrhea , appetite changes, weight changes. Certain side effects may be more likely in older adults. Common Euthyrox side effects may include: chest pain, irregular heartbeats
- shortness of breath
- headache, leg cramps, muscle pain or weakness
- tremors, feeling nervous or irritable, trouble sleeping
- increased appetite
- feeling hot
- weight loss
- changes in your menstrual periods
- diarrhea
- or skin rash , partial hair loss.

1957. This particular disease, Hypothyroidism, interacts with the drug liothyronine and has the following

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Stop using liothyronine and call your doctor if you have symptoms of thyroid toxicity , such as chest pain, fast or pounding heartbeats, feeling hot or nervous, or sweating more than usual. Also call your doctor at once if you have: weak or shallow breathing
- confusion
- a feeling of being cold all the time
- changes in weight or appetite
- chest pressure spreading to your jaw or shoulder
- a light-headed feeling, like you might pass out
- severe headaches, ringing in your ears, dizziness , vision problems, pain behind your eyes
- a seizure
- or (in a child or teenager) pain, stiffness, and trouble moving the legs. Common side effects of liothyronine may include: trouble breathing
- headache
- tremors, feeling nervous or irritable
- muscle weakness
- increased appetite
- diarrhea
- irregular menstrual periods
- weight loss
- feeling hot
- rash
- or sleep problems (insomnia).

1958. This particular disease, Hypothyroidism, interacts with the drug Nature-Throid and has the fo

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common side effects may include temporary hair loss (especially in children).

1959. This particular disease, Hypothyroidism, interacts with the drug Unithroid and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Chest pain, discomfort, or tightness decreased urine output difficult or labored breathing difficulty with swallowing dilated neck veins extreme fatigue fainting fast, slow, irregular, pounding, or racing heartbeat or pulse fever heat intolerance hives or welts, skin itching, rash, or redness irregular breathing irritability menstrual changes nausea pain or discomfort in the arms, jaw, back, or neck sweating swelling of the eyes, face, lips, throat, or tongue tremors Rare Blurred or double vision dizziness eye pain lack or slowing of normal growth in children limp or walk favoring one leg pain in the hip or knee seizures severe headache Symptoms of overdose Change in consciousness cold, clammy skin confusion disorientation fast or weak pulse lightheadedness loss of consciousness sudden headache sudden loss of coordination sudden slurring of speech Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Change in appetite crying diarrhea false or unusual sense of well-being fear or nervousness feeling not well or unhappy feeling of discomfort feeling of warmth feeling things are not real feelings of suspicion and distrust hair loss headache increased appetite mental depression muscle weakness

Drug Side Effects Summary

quick to react or overreact emotionally rapidly changing moods redness of the face, neck, arms, and occasionally, upper chest restlessness stomach cramps trouble getting pregnant trouble sitting still unusual tiredness or weakness vomiting weight gain or loss Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Unithroid side effects

1960. This particular disease, Hypothyroidism, interacts with the drug Westroid and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common side effects may include temporary hair loss (especially in children).

1961. This particular disease, Hypothyroidism, interacts with the drug Levo-T and has the following

- No side effects listed

1962. This particular disease, Hypothyroidism, interacts with the drug Thyquidity and has the following

- No side effects listed

1963. This particular disease, Hypothyroidism, interacts with the drug Tirosint-Sol and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Chest pain, discomfort, or tightness decreased urine output difficult or labored breathing difficulty with swallowing dilated neck veins extreme fatigue fainting fast, slow, irregular, pounding, or racing heartbeat or pulse fever

Drug Side Effects Summary

heat intolerance hives or welts, skin itching, rash, or redness irregular breathing irritability menstrual changes nausea pain or discomfort in the arms, jaw, back, or neck sweating swelling of the eyes, face, lips, throat, or tongue tremors Rare Blurred or double vision dizziness eye pain lack or slowing of normal growth in children limp or walk favoring one leg pain in the hip or knee seizures severe headache Symptoms of overdose Change in consciousness cold, clammy skin confusion disorientation fast or weak pulse lightheadedness loss of consciousness sudden headache sudden loss of coordination sudden slurring of speech Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Change in appetite crying diarrhea false or unusual sense of well-being fear or nervousness feeling not well or unhappy feeling of discomfort feeling of warmth feeling things are not real feelings of suspicion and distrust hair loss headache increased appetite mental depression muscle weakness quick to react or overreact emotionally rapidly changing moods redness of the face, neck, arms, and occasionally, upper chest restlessness stomach cramps trouble getting pregnant trouble sitting still unusual tiredness or weakness vomiting weight gain or loss Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Tirosint-Sol side effects

1964. This particular disease, Hypothyroidism, interacts with the drug NP Thyroid and has the follo

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common side effects may include temporary hair loss

Drug Side Effects Summary

(especially in children).

1965. This particular disease, Hypothyroidism, interacts with the drug Triostat and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: More common Dizziness fainting fast, slow, or irregular heartbeat Less common Bluish color blurred vision changes in skin color chest pain or discomfort chest tightness or heaviness confusion decreased urine output dilated neck veins dizziness, faintness, or lightheadedness when getting up from a lying or sitting position suddenly extreme fatigue fever headache heart stops irregular breathing nausea nervousness no breathing no pulse or blood pressure pain or discomfort in arms, jaw, back or neck pain, tenderness, swelling of foot or leg pounding in the ears shortness of breath sweating swelling of face, fingers, feet, or lower legs troubled breathing unconscious unusual tiredness or weakness vomiting weight gain wheezing Symptoms of overdose Cold clammy skin increased bowel movements irritability lightheadedness menstrual changes shakiness in legs, arms, hands, feet trembling or shaking of hands or feet weak pulse Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Twitching Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Triostat side effects

1966. This particular disease, Hypothyroidism, interacts with the drug WP Thyroid and has the follo

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common side effects may include temporary hair loss (especially in children).

1967. This particular disease, IBD (Bowel), interacts with the drug budesonide and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Budesonide may cause serious side effects. Call your doctor at once if you have: thinning skin, easy bruising, increased acne or facial hair
- swelling in your ankles
- weakness, tiredness, or a light-headed feeling, like you might pass out
- nausea , vomiting, rectal bleeding
- pain or burning when you urinate
- menstrual problems (in women), impotence or loss of interest in sex (in men)
- or signs of too much steroid medicine in your blood--acne, bruising, thinning skin, increased facial or body hair, stretch marks, increased body fat, or changes in the shape or location of body fat (especially in your face, neck, back, and waist). Common side effects of budesonide may include: headache , dizziness
- indigestion , nausea, vomiting, stomach pain, bloating , gas, constipation
- feeling tired
- back pain , joint pain
- painful urination
- cold symptoms such as stuffy nose, sneezing, sore throat
- pain anywhere in your body

Drug Side Effects Summary

- or signs of too much steroid medicine in your blood.

1968. This particular disease, IBD (Bowel), interacts with the drug mesalamine and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Mesalamine may cause serious side effects. Stop using mesalamine and call your doctor at once if you have: severe stomach pain, stomach cramping, bloody diarrhea
- fever, headache, general ill feeling
- rash, itching, eye redness
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- low white blood cell counts--fever, chills, mouth sores, skin sores, sore throat , cough, feeling light-headed, trouble breathing
- signs of a kidney stone--severe pain in your side and back, frequent need to urinate, foul-smelling urine, dark or cloudy urine
- kidney problems--increased or decreased urination, swelling, weight gain
- or liver problems--loss of appetite, upper stomach pain, tiredness, easy bruising or bleeding, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Low white blood cell counts may be more likely in older adults. Common side effects of mesalamine may include: burping , constipation , nausea , vomiting , stomach pain, diarrhea, gas
- dizziness
- cold symptoms such as stuffy nose, sneezing, sore throat
- back pain
- headache
- rash

Drug Side Effects Summary

- or abnormal liver function tests .

1969. This particular disease, IBD (Bowel), interacts with the drug Lialda and has the following side

- hives

- difficulty breathing

- swelling of your face, lips, tongue, or throat. Stop using Lialda and call your doctor at once if you have: severe stomach pain, stomach cramping, bloody diarrhea

- fever, headache, general ill feeling

- rash, itching, eye redness

- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds

- low white blood cell counts - fever, chills, mouth sores, skin sores, sore throat , cough, feeling light-headed, trouble breathing

- signs of a kidney stone - severe pain in your side and back, frequent need to urinate, foul-smelling urine, dark or cloudy urine

- kidney problems - increased or decreased urination, swelling, weight gain

- or liver problems - loss of appetite, upper stomach pain, tiredness, easy bruising or bleeding, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Low white blood cell counts may be more likely in older adults. Common Lialda side effects may include: burping , constipation , nausea , vomiting , stomach pain, diarrhea, gas

- dizziness

- cold symptoms such as stuffy nose, sneezing, sore throat

- back pain

- headache

- rash

- or abnormal liver function tests .

Drug Side Effects Summary

1970. This particular disease, IBD (Bowel), interacts with the drug Asacol and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Bloody diarrhea rectal bleeding stomach pain Less common Bloody stools bloody urine blurred vision chest tightness chills clay-colored stools cough dark urine diarrhea difficulty with breathing dizziness fever full or bloated feeling general feeling of discomfort or illness headache itching, skin rash joint pain loss of appetite muscle aches and pains nausea nervousness pain or tenderness around the eyes and cheekbones pounding in the ears pressure in the stomach runny or stuffy nose shivering slow or fast heartbeat sore throat stomach cramps sweating swelling of the stomach area trouble sleeping unpleasant breath odor unusual tiredness or weakness vomiting vomiting of blood yellow eyes or skin Rare Anxiety back pain blue or pale skin chest pain, possibly moving to the left arm, neck, or shoulder Incidence not known Black, tarry stools blistering, peeling, or loosening of the skin bloating change in the ability to see colors, especially blue or yellow chest discomfort constipation decreased frequency or amount of urine dry cough greatly increased frequency of urination or amount of urine hives or welts hoarseness inability to move the arms and legs increased thirst indigestion large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or genitals light-colored stools lower back or side pain muscle weakness, sudden and progressing pain in the ankles or knees painful or difficult urination painful, red lumps under the skin, mostly on the legs rapid breathing red, irritated eyes red skin lesions, often with a purple center severe nausea or vomiting sores, ulcers, or white spots in the mouth or on the lips stomach tenderness sudden numbness and weakness in the arms and legs swelling of the feet or lower legs swollen or painful glands thickening of bronchial secretions unusual bleeding or bruising weight gain Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways

Drug Side Effects Summary

to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Sneezing Less common Acne belching continuing ringing or buzzing or other unexplained noise in the ears difficulty with moving dizziness or lightheadedness excess air or gas in the stomach or bowels feeling of constant movement of self or surroundings hair loss or thinning of the hair hearing loss heartburn muscle stiffness passing gas sensation of spinning stomach discomfort or upset Incidence not known Burning, numbness, tingling, or painful sensations unsteadiness or awkwardness weakness in the arms, hands, legs, or feet Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Asacol side effects

1971. This particular disease, IBD (Bowel), interacts with the drug Pentasa and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using Pentasa and call your doctor at once if you have: severe stomach pain, stomach cramping, bloody diarrhea
- fever, headache, general ill feeling
- rash, itching, eye redness
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- low white blood cell counts - fever, chills, mouth sores, skin sores, sore throat , cough, feeling light-headed, trouble breathing
- signs of a kidney stone - severe pain in your side and back, frequent need to urinate, foul-smelling urine, dark or cloudy urine
- kidney problems - increased or decreased urination, swelling, weight gain

Drug Side Effects Summary

- or liver problems - loss of appetite, upper stomach pain, tiredness, easy bruising or bleeding, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Low white blood cell counts may be more likely in older adults. Common Pentasa side effects may include: burping , constipation , nausea , vomiting , stomach pain, diarrhea, gas
- dizziness
- cold symptoms such as stuffy nose, sneezing, sore throat
- back pain
- headache
- rash
- or abnormal liver function tests .

1972. This particular disease, IBD (Bowel), interacts with the drug Apriso and has the following side

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using Apriso and call your doctor at once if you have: severe stomach pain, stomach cramping, bloody diarrhea
- fever, headache, general ill feeling
- rash, itching, eye redness
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- low white blood cell counts - fever, chills, mouth sores, skin sores, sore throat , cough, feeling light-headed, trouble breathing
- signs of a kidney stone - severe pain in your side and back, frequent need to urinate, foul-smelling urine, dark or cloudy urine
- kidney problems - increased or decreased urination, swelling, weight gain
- or liver problems - loss of appetite, upper stomach pain, tiredness, easy bruising or bleeding, dark

Drug Side Effects Summary

urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Low white blood cell counts may be more likely in older adults. Common Apriso side effects may include: burping , constipation , nausea , vomiting , stomach pain, diarrhea, gas

- dizziness
- cold symptoms such as stuffy nose, sneezing, sore throat
- back pain
- headache
- rash
- or abnormal liver function tests .

1973. This particular disease, IBD (Bowel), interacts with the drug Remicade and has the following side effects:

- Some side effects may occur during the injection. Tell your caregiver if you feel dizzy, nauseated, light-headed, itchy or tingly, short of breath, or have a headache , fever, chills, muscle or joint pain , pain or tightness in your throat, chest pain, or trouble swallowing during the injection. Infusion reactions may also occur within 1 or 2 hours after injection. hives
- chest pain, difficult breathing
- fever, chills, severe dizziness
- swelling of your face, lips, tongue, or throat. Serious and sometimes fatal infections may occur during treatment with Remicade. Call your doctor right away if you have signs of infection such as: fever, extreme tiredness, flu symptoms, cough, or skin symptoms (pain, warmth, or redness). Also call your doctor if you have: skin changes, new growths on the skin
- pale skin, easy bruising or bleeding
- delayed allergic reaction (up to 12 days after receiving Remicade) - fever, sore throat , trouble swallowing, headache, joint or muscle pain , skin rash , or swelling in your face or hands
- liver problems - right-sided upper stomach pain, loss of appetite, yellowing of your skin or eyes

Drug Side Effects Summary

(jaundice), and not feeling well

- lupus -like syndrome - joint pain or swelling, chest discomfort, feeling short of breath, skin rash on your cheeks or arms (worsens in sunlight)
- nerve problems - numbness or tingling, problems with vision, weakness in your arms or legs, seizure
- new or worsening psoriasis - skin redness or scaly patches, raised bumps filled with pus
- signs of heart failure - shortness of breath with swelling of your ankles or feet, rapid weight gain
- signs of a stroke - sudden numbness or weakness, trouble speaking or understanding what is said to you, problems with vision or balance, severe headache
- signs of lymphoma - fever, night sweats , weight loss , stomach pain or swelling, chest pain, cough, trouble breathing, swollen glands (in your neck, armpits, or groin)
- or signs of tuberculosis - fever, cough, night sweats, loss of appetite, weight loss, feeling constantly tired. Serious infections may be more likely in adults who are 65 years or older. Common Remicade side effects may include: stuffy nose, sinus pain
- fever, chills, sore throat
- cough, chest pain, shortness of breath
- high or low blood pressure
- headache, feeling light-headed
- rash, itching
- or stomach pain.

1974. This particular disease, IBD (Bowel), interacts with the drug infliximab and has the following s

- Infliximab can cause serious side effects, including: See "Important information" Serious Infections
Some patients, especially those 65 years and older have had serious infections while receiving infliximab. These serious infections include TB and infections caused by viruses, fungi, or bacteria

Drug Side Effects Summary

that have spread throughout the body or cause infections in certain areas (such as skin). Some patients die from these infections. If you get an infection while receiving treatment with infliximab your doctor will treat your infection and may need to stop your infliximab treatment. Tell your doctor right away if you have any of the following signs of an infection while receiving or after receiving infliximab: a fever have flu-like symptoms feel very tired warm, red, or painful skin have a cough Your doctor will examine you for TB and perform a test to see if you have TB. If your doctor feels that you are at risk for TB, you may be treated with medicine for TB before you begin treatment with infliximab and during treatment with infliximab. Even if your TB test is negative, your doctor should carefully monitor you for TB infections while you are receiving infliximab. Patients who had a negative TB skin test before receiving infliximab have developed active TB. If you are a chronic carrier of the hepatitis B virus, the virus can become active while you are being treated with infliximab. In some cases, patients have died as a result of hepatitis B virus being reactivated. Your doctor should do a blood test for hepatitis B virus before you start treatment with infliximab and occasionally while you are being treated. Tell your doctor if you have any of the following symptoms: feel unwell tiredness (fatigue) poor appetite fever, skin rash, or joint pain Heart Failure If you have a heart problem called congestive heart failure, your doctor should check you closely while you are receiving infliximab. Your congestive heart failure may get worse while you are receiving infliximab. Be sure to tell your doctor of any new or worse symptoms including: shortness of breath sudden weight gain swelling of ankles or feet Treatment with infliximab may need to be stopped if you get new or worse congestive heart failure. Other Heart Problems Some patients have experienced a heart attack (some of which led to death), low blood flow to the heart, or abnormal heart rhythm within 24 hours of beginning their infusion of infliximab. Symptoms may include chest discomfort or pain, arm pain, stomach pain, shortness of breath, anxiety, lightheadedness, dizziness, fainting, sweating, nausea, vomiting, fluttering or pounding in your chest, and/or a fast or a slow heartbeat. Tell your doctor right away if you have any of these symptoms. Liver Injury Some patients receiving

Drug Side Effects Summary

infliximab have developed serious liver problems. Tell your doctor if you have: jaundice (skin and eyes turning yellow) fever dark brown-colored urine extreme tiredness (severe fatigue) pain on the right side of your stomach area (right-sided abdominal pain) Blood Problems In some patients receiving infliximab, the body may not make enough of the blood cells that help fight infections or help stop bleeding. Tell your doctor if you: have a fever that does not go away look very pale bruise or bleed very easily Nervous System Disorders Some patients receiving infliximab have developed problems with their nervous system. Tell your doctor if you have: changes in your vision seizures numbness or tingling in any part of your weakness in your arms or legs body Some patients have experienced a stroke within approximately 24 hours of their infusion with infliximab. Tell your doctor right away if you have symptoms of a stroke which may include: numbness or weakness of the face, arm or leg, especially on one side of the body

- sudden confusion, trouble speaking or understanding
- sudden trouble seeing in one or both eyes, sudden trouble walking, dizziness, loss of balance or coordination or a sudden, severe headache.

Allergic Reactions Some patients have had allergic reactions to infliximab. Some of these reactions were severe. These reactions can happen while you are getting your infliximab treatment or shortly afterward. Your doctor may need to stop or pause your treatment with infliximab and may give you medicines to treat the allergic reaction. Signs of an allergic reaction can include: hives (red, raised, itchy patches of skin) difficulty breathing chest pain high or low blood pressure fever chills Some patients treated with infliximab have had delayed allergic reactions. The delayed reactions occurred 3 to 12 days after receiving treatment with infliximab. Tell your doctor right away if you have any of these signs of delayed allergic reaction to infliximab: fever rash headache sore throat muscle or joint pain swelling of the face and hands difficulty swallowing Lupus-like Syndrome Some patients have developed symptoms that are like the symptoms of Lupus. If you develop any of the following symptoms, your doctor may decide to stop your treatment with infliximab. chest discomfort or pain that does not go away shortness of breath

Drug Side Effects Summary

joint pain rash on the cheeks or arms that gets worse in the sun Psoriasis Some people receiving infliximab had new psoriasis or worsening of psoriasis they already had. Tell your doctor if you develop red scaly patches or raised bumps on the skin that are filled with pus. Your doctor may decide to stop your treatment with infliximab. The most common side effects of infliximab include: respiratory infections, such as sinus infections and sore throat headache coughing stomach pain Infusion reactions can happen up to 2 hours after your infusion of infliximab. Symptoms of infusion reactions may include: fever chills chest pain low blood pressure or high blood pressure shortness of breath rash itching Children who received infliximab in studies for Crohn's disease showed some differences in side effects compared with adults who received infliximab for Crohn's disease. The side effects that happened more in children were: anemia (low red blood cells), leukopenia (low white blood cells), flushing (redness or blushing), viral infections, neutropenia (low neutrophils, the white blood cells that fight infection), bone fracture, bacterial infection and allergic reactions of the breathing tract. Among patients who received infliximab for ulcerative colitis in clinical studies, more children had infections as compared with adults. Tell your doctor about any side effect that bothers you or does not go away. These are not all of the side effects with infliximab. Ask your doctor or pharmacist for more information. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800- FDA-1088. Interactions Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. These include any other medicines to treat Crohn's disease, ulcerative colitis, rheumatoid arthritis, ankylosing spondylitis, psoriatic arthritis or psoriasis. Know the medicines you take. Keep a list of your medicines and show them to your doctor and pharmacist when you get a new medicine. Pregnancy and breastfeeding Tell your doctor if you are pregnant or plan to become pregnant, are breastfeeding or plan to breastfeed. You and your doctor should decide if you should receive infliximab while you are pregnant or breastfeeding. If you have a baby and you were receiving infliximab during your pregnancy, it is important to tell your baby's doctor and other health care

Drug Side Effects Summary

professionals about your infliximab use so they can decide when your baby should receive any vaccine. Certain vaccinations can cause infections. If you received infliximab while you were pregnant, your baby may be at higher risk for getting an infection. If your baby receives a live vaccine within 6 months after birth, your baby may develop infections with serious complications that can lead to death. This includes live vaccines such as the BCG, rotavirus, or any other live vaccines. For other types of vaccines, talk with your doctor

Storage Store unopened infliximab vials in a refrigerator at 2°C to 8°C (36°F to 46°F). If needed, unopened infliximab vials may be stored at room temperatures up to a maximum of 30°C (86°F) for a single period of up to 6 months but not exceeding the original expiration date. The new expiration date must be written in the space provided on the carton. Once removed from the refrigerator, infliximab cannot be returned to the refrigerator.

What are the ingredients in infliximab? Active ingredient: infliximab Inactive ingredients: Remicade: dibasic sodium phosphate dihydrate, monobasic sodium phosphate monohydrate, polysorbate 80, and sucrose. No preservatives are present. Avsola: dibasic sodium phosphate anhydrous, monobasic sodium phosphate monohydrate, polysorbate 80, and sucrose. No preservatives are present. Inflectra: dibasic sodium phosphate dihydrate, monobasic sodium phosphate monohydrate, polysorbate 80, and sucrose. No preservatives are present. Ixifi: disodium succinate hexahydrate, polysorbate 80, succinic acid, and sucrose. No preservatives are present. Renflexis: dibasic sodium phosphate heptahydrate, monobasic sodium phosphate monohydrate, polysorbate 80, and sucrose. No preservatives are present. Infliximab products are manufactured by the following companies: Remicade: Janssen Biotech, Inc. Horsham, PA 19044. Avsola: Amgen, Inc. One Amgen Center Drive, Thousand Oaks, CA 91320-1799. Inflectra: CELLTRION, Inc. 23, Academy-ro, Yeonsu-gu, Incheon, 22014, Republic of Korea. Ixifi: Pfizer Ireland Pharmaceuticals Ringaskiddy, Co. Cork, Ireland. Renflexis: Samsung Bioepis Co., Ltd., 76, Songdogoyoyuk-ro, Yeonsu-gu, Incheon, 21987, Republic of Korea.

Popular FAQ What is the difference between Inflectra and Remicade? Inflectra (infliximab-dyyb) is a biosimilar to Remicade (infliximab). They

Drug Side Effects Summary

belong to the same class of drugs called tumor necrosis factor (TNF) alfa inhibitors. Inflectra can be prescribed for the same uses as Remicade, including rheumatoid arthritis, Crohn's disease, ankylosing spondylitis, ulcerative colitis, psoriatic arthritis, and plaque psoriasis. Continue reading

What is the difference between Renflexis and Remicade? Renflexis (infliximab-abda) is a tumor necrosis factor (TNF) blocker that is biosimilar to Remicade (infliximab). A biosimilar has no clinically meaningful differences in terms of safety and effectiveness from the reference product. Renflexis is given as an intravenous (IV) infusion similar to Remicade, and is used for the same conditions, including rheumatoid arthritis, Crohn's disease and psoriatic arthritis. Biosimilars may lead to cost-savings due to a lower price and are preferred by some insurance companies. Continue reading

More FAQ What are the new drugs for the treatment of rheumatoid arthritis (RA)? What are the new drugs for the treatment of plaque psoriasis? How many biosimilars have been approved in the United States? View more FAQ

References Food and Drug Administration (FDA) Avsola Product Label Food and Drug Administration (FDA) Ixifi Product Label Food and Drug Administration (FDA) Infliximab Product Label Food and Drug Administration (FDA) Inflectra Product Label Food and Drug Administration (FDA) Renflexis Product Label. More about infliximab Check interactions Pricing & coupons Reviews (224) Side effects Dosage information During pregnancy Drug class: antirheumatics Breastfeeding En español Patient resources Advanced Reading

Infliximab-abda Intravenous (Advanced Reading) Infliximab-axxq Intravenous (Advanced Reading) Infliximab-dyyb Intravenous (Advanced Reading) Infliximab-qbtX Intravenous (Advanced Reading) Other brands Remicade , Inflectra , Renflexis , Avsola , Ixifi Professional resources Prescribing Information Related treatment guides Celiac Disease Ankylosing Spondylitis Behcet's Disease Bullous Pemphigoid Further information Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances. Medical Disclaimer

Psoriasis 101 What are the new treatments for plaque psoriasis? Learn about a popular treatment for psoriasis Drug Status Availability Prescription only Rx Pregnancy & Lactation Risk data available

Drug Side Effects Summary

CSA Schedule* Not a controlled drug N/A Loading... Approval History Drug history at FDA Loading... User Reviews & Ratings 6.7 / 10 224 Reviews Related Drugs Remicade , Inflectra , Renflexis , Avsola , Cosentyx , Enbrel , Entyvio , Humira , Otezla , Stelara Drugs.com Mobile Apps The easiest way to lookup drug information, identify pills, check interactions and set up your own personal medication records. Available for Android and iOS devices. Explore Apps Support Help Center Sitemap Contact Us About About Drugs.com Advertising Policy Attribution & Citations Terms & Privacy Editorial Policy Privacy Policy Terms of Use Follow Drugs.com on Facebook Follow Drugs.com on Twitter Follow Drugs.com on Pinterest Follow Drugs.com on YouTube Subscribe to Drugs.com newsletters Subscribe to Drugs.com newsletters for the latest medication news, new drug approvals, alerts and updates. Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include IBM Watson Micromedex (updated 2 Dec 2022), Cerner Multum? (updated 23 Nov 2022), ASHP (updated 11 Nov 2022) and others. Ad Choices We comply with the HONcode standard for trustworthy health information. Copyright © 2000-2022 Drugs.com. All rights reserved.

SITECONFIG

=

```
{"agent":{"isMobile":false},"user":{"userid":0,"modal":{"login":"/api/user/modal/login/"},"consent":{"skip":true,"api":{"url":"/api/user/"}}, "page":{"section":"rootdocs","subsection":"","doc":"content","delayedCss":["/bundle/css/component/component-share.min.9d6804aa.css"],"delayedScripts":[],"image":{"lazyLoad":{"selector":["data-lazy-load"],"screens":1,"delay":2000}}, "mailto":{"delay":5000,"location":"/support/email-mailto.html"}}, "form":{"recaptcha":{"sitekey":"6Ld-iSYTAAAAA00VTEwVI71HtvrFoSIP SibGr4g"},"callback":"DDC_RECAPTCHA_CALLBACK"},"dwLog":{"a":1336,"b":"null","c":"7,10,1179,336,10168,1419,1463,1464,641,138,1249,704,10187,334,333,10266,1247,54,1121,1251,959,58,1242,60,1245,10267,1246,1250,115,1555,66","d":"6,9,53,114,115,127,132,134,273,318","e":"182,437","f":3761,"h":0,"i":"default","k":"7,1419,1463,1464,138,10266,54,1121,58,115,1555","l":"","m":"","n":""}}
```


Drug Side Effects Summary

, "o": "", "p": 0, "q": "drugs/rootdocs"}, "drugs": [{"ddc_id": 1336, "brand_name_id": 0, "mtm_id": "d04342", "generic_name": "infliximab", "systemic": true, "name": "infliximab"}], "options": {"recordSelection": true, "classifyLists": {"wrapper": "#content", "countHtmlItems": false, "numItems": {"short": 15, "medium": 30, "long": 60}, "charLimit": {"paragraph": 30, "content": 350}}, "subscribe": null, "modal": null, "dictionary": {}, "share": {"sticky": {"target": ".ddc-social-share"}, "types": [{"title": "Facebook", "label": "Share on Facebook", "cssClass": "ddc-share-icon-ddc-share-icon-facebook", "labelClass": "ddc-share-icon-label", "icon": "<svg class='ddc-icon-ddc-icon-facebook' width='20' height='20' viewBox='0 0 24 24' aria-hidden='true' focusable='false' xmlns='http://www.w3.org/2000/svg'><path d='M14 6.54c-2.06 0-3.4 1.25-3.4 3.52v2H8.3v2.61h2.3V21A9.04 9.04 0 0 1 12.05C3 7.05 7.03 3 12 3s9 4.05 9 9.05c0 4.53-3.3 8.27-7.6 8.95v-6.33h2.1l4-2.62h-2.5v-1.7c0-.7.35-1.4 1.47-1.4h1.14V6.7s-1.03-.17-2.02-.17z' /></svg>", "shareUrl": "https://www.facebook.com/sharer/sharer.php?u=SHAREURL", "width": 600, "height": 500}, {"title": "Twitter", "label": "Share on Twitter", "cssClass": "ddc-share-icon-ddc-share-icon-twitter", "labelClass": "ddc-share-icon-label", "icon": "<svg class='ddc-icon-ddc-icon-twitter' width='20' height='20' viewBox='0 0 24 24' aria-hidden='true' focusable='false' xmlns='http://www.w3.org/2000/svg'><path d='M8.66 19.63a10.44 10.44 0 0 1 0.5-10.5v-.5A7.7 7.7 0 0 2 1 6.74c-.67.3-1.39.5-2.12.58a3.73 3.73 0 0 1 .64-2.04c-.72.43-1.52.74-2.34.9a3.7 3.7 0 0 0-.6.3.36 10.46 10.46 0 0 1-.7.61-3.85 3.69 3.69 0 0 1 .14 4.9 3.66 3.66 0 0 1-1.63-.45v.04a3.7 3.7 0 0 2 .9 3.64 3.7 3.7 0 0 1-1.63.06 3.69 3.69 0 0 3 .42 2.53 7.4 7.4 0 0 1-4.6 1.64 8 8 0 0 1-.87-.07 10.45 10.45 0 0 5.66 1.63' /></svg>", "shareUrl": "https://twitter.com/intent/tweet?text=SHARETITLE&url=SHAREURL", "width": 600, "height": 420}, {"title": "Email", "label": "Email to a friend", "cssClass": "ddc-share-icon-ddc-share-icon-email", "labelClass": "ddc-share-icon-label", "icon": "<svg class='ddc-icon-ddc-icon-email' width='20' height='20' viewBox='0 0 24 24' aria-hidden='true' focusable='false' xmlns='http://www.w3.org/2000/svg'><path d='M20 4.25c.92 0 1.67.7 1.74 1.6l.01.15v12c0 .92-.7

Drug Side Effects Summary

1.67-1.6 1.74l-.15.01H4c-.92 0-1.67-.7-1.74-1.6L2.25 18V6c0-.92.7-1.67 1.6-1.74L4 4.25h16zm-.17
1.77a.75.75 0 00-1.06-.1L12 11.52l-6.77-5.6-.1-.06a.75.75 0 00-.86 1.22l7.25 6
.1.07c.27.15.61.13.86-.07l7.25-6L19.8 7a.75.75 0 00.02-.98z'

/></svg>","mailUrl":"mailto:?body=https%3A%2F%2Fwww.drugs.com%2Finfliximab.html","width":60
0,"height":420}},"mednotes":{"apiUrl":"/mednotes/api/modal","saveButton":"data-modal-mednotes-sa
ve"},"ads":{"section":"drugs/rootdocs","ugc":false,"minWidth728":1120,"minWidth970":1540,"display
":{"confiant":false,"mobile":false,"timeout":900,"logUrl":"/api/log/ads/","targeting":{"ugc":"0","c1":["294
5","1834","4093","4287","1661","1738","4223","6067","3557","1820"],"d":["1336","b":"0","c2":["7","10",
"1179","336","10168","1419","1463","1464","641","138","1249","704","10187","334","333","10266","1
247","54","1121","1251","959","58","1242","60","1245","10267","1246","1250","115","1555","66"],"dc"
:["182","437"],"mi":"1","rx":"1","c2o":["7","1419","1463","1464","138","10266","54","1121","58","115",
1555"]},"units":{"code":"display-ad-1","section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[728,90]],"siz
eMappings":[[[0,0],[[320,100],[[320,50],[300,50]]],[[768,300],[[728,90]]]],"targeting":{"uid":"d-leader
board-1","mobile":0,"prebid":1,"size":"728x90","tier":"static","loading":"eager","position":1,"layout":{"
"prebid":1},"code":"display-ad-2","section":"/7146/drugs/rootdocs","lazy":0.75,"sizes":[[300,250],[300
,600],[1,1]],"sizeMappings":[[[0,0],[[320,100],[[300,250],[1,1]]],[[700,300],[[300,600],[300,250],[1,1]]
]],"targeting":{"uid":"d-300x600-static-1","mobile":0,"prebid":1,"size":"300x600","tier":"static","loading"
:"lazy","position":2,"layout":{"prebid":1},"code":"display-ad-3","section":"/7146/drugs/rootdocs","laz
y":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[[320,100],[[300,250],[1,1]]]],"targeting":{"uid
":"d-300x250-static-1","mobile":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","position":
3,"layout":{"wrappingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSideb
arFloat":true},"prebid":1},"code":"display-ad-4","section":"/7146/drugs/rootdocs","lazy":0.75,"sizes":[[
300,250],[1,1]],"sizeMappings":[[[0,0],[[320,100],[[300,250],[1,1]]]],"targeting":{"uid":"d-300x250-st
atic-2","mobile":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","position":4,"layout":{"wra
ppingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSidebarFloat":true},"pr

Drug Side Effects Summary

ebid":1},{ "code":"display-ad-5", "section":"/7146/drugs/rootdocs", "lazy":0.75, "sizes":[[300,250],[1,1]], "sizeMappings":[[[0,0],[[320,100],[[300,250],[1,1]]]], "targeting":{"uid":"d-300x250-static-3", "mobile":0, "prebid":1, "size":"300x250", "tier":"static", "loading":"lazy", "position":5}, "layout":{"wrappingClass":"sideBoxStackedAdWrap", "requiresContentHeight":true, "requiresSidebarFloat":true}, "prebid":1}, {"code":"display-ad-6", "section":"/7146/drugs/rootdocs", "lazy":0.75, "sizes":[[300,250],[1,1]], "sizeMappings":[[[0,0],[[320,100],[[300,250],[1,1]]]], "targeting":{"uid":"d-300x250-static-4", "mobile":0, "prebid":1, "size":"300x250", "tier":"static", "loading":"lazy", "position":6}, "layout":{"wrappingClass":"sideBoxStackedAdWrap", "requiresContentHeight":true, "requiresSidebarFloat":true}, "prebid":1}, {"code":"display-ad-7", "section":"/7146/drugs/rootdocs", "lazy":0.75, "sizes":[[300,250],[1,1]], "sizeMappings":[[[0,0],[[320,100],[[300,250],[1,1]]]], "targeting":{"uid":"d-300x250-static-5", "mobile":0, "prebid":1, "size":"300x250", "tier":"static", "loading":"lazy", "position":7}, "layout":{"wrappingClass":"sideBoxStackedAdWrap", "requiresContentHeight":true, "requiresSidebarFloat":true}, "prebid":1}, {"code":"display-ad-8", "section":"/7146/drugs/rootdocs", "lazy":0.75, "sizes":[[300,250],[1,1]], "sizeMappings":[[[0,0],[[320,100],[[300,250],[1,1]]]], "targeting":{"uid":"d-300x250-static-6", "mobile":0, "prebid":1, "size":"300x250", "tier":"static", "loading":"lazy", "position":8}, "layout":{"wrappingClass":"sideBoxStackedAdWrap", "requiresContentHeight":true, "requiresSidebarFloat":true}, "prebid":1}, {"code":"display-ad-9", "section":"/7146/drugs/rootdocs", "lazy":0.75, "sizes":[[300,250],[1,1]], "sizeMappings":[[[0,0],[[320,100],[[300,250],[1,1]]]], "targeting":{"uid":"d-300x250-static-7", "mobile":0, "prebid":1, "size":"300x250", "tier":"static", "loading":"lazy", "position":9}, "layout":{"wrappingClass":"sideBoxStackedAdWrap", "requiresContentHeight":true, "requiresSidebarFloat":true}, "prebid":1}, {"code":"display-ad-10", "section":"/7146/drugs/rootdocs", "lazy":0.75, "sizes":[[300,250],[1,1]], "sizeMappings":[[[0,0],[[320,100],[[300,250],[1,1]]]], "targeting":{"uid":"d-300x250-static-8", "mobile":0, "prebid":1, "size":"300x250", "tier":"static", "loading":"lazy", "position":10}, "layout":{"wrappingClass":"sideBoxStackedAdWrap", "requiresContentHeight":true, "requiresSidebarFloat":true}, "prebid":1}, {"code":"display-ad-11", "section":"/7146/drugs/rootdocs", "lazy":0.75, "sizes":[[300,250],[1,1]], "sizeMappings":[[[0,0],[[320,100],[[300,250],[1,1]]]], "targeting":{"uid":"d-300x250-static-9", "mobile":0, "prebid":1, "size":"300x250", "tier":"static", "loading":"lazy", "position":11}, "layout":{"wrappingClass":"sideBoxStackedAdWrap", "requiresContentHeight":true, "requiresSidebarFloat":true}, "prebid":1}

Drug Side Effects Summary

```
bile":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","position":11},"layout":{"wrappingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSidebarFloat":true},"prebid":1}},"injection":{"lazy":0.75,"maximum":{"ads":10,"tags":1000},"container":"#content","start":[".nav-tabs","h2","h1"],"stop":[".footer-feature","#sidebar"],"last":{"buffer":500},"insert":["section","h2","h3","h4","h5","p","ol","ul","table",".ddc-legend",".ddc-media",".nav-tabs",".nav-tabs-group",".responsive-table-wrap"],"blacklist":["form","ol","ul","dl","table",".blackboxWarning",".ddc-accordion",".ddc-box",".ddc-card",".ddc-card-group",".ddc-form-inline",".ddc-expand-read-more",".ddc-grid",".ddc-inset-text",".ddc-legend",".ddc-media",".ddc-notification",".ddc-paging",".ddc-reference-list",".ddc-toc",".ddc-widget-calculator",".injection-ignore-inside",".nav-tabs",".nav-tabs-group",".paging-list-wrap",".responsive-table-wrap",".referenceList",".snippet",".ddc-official-answer"],"list":[{"position":1,"type":"Prebid","width":728,"height":90,"buffer":-1,"viewportPercent":85,"minBuffer":800,"lazy":0.75,"data":{"section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[728,90]],"sizeMappings":[],"targeting":{"uid":"d-728x90-auto-1","mobile":0,"prebid":1,"size":"728x90","tier":"auto"},"prebid":1},"conditional":{"minWidth":1120,"fallback":{"position":1,"type":"Prebid","width":300,"height":250,"buffer":-1,"viewportPercent":85,"minBuffer":800,"lazy":0.75,"data":{"section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[300,250]],"sizeMappings":[],"targeting":{"uid":"d-mrec-1","mobile":0,"prebid":1,"size":"300x250","tier":"auto"},"prebid":1}}},"{"position":2,"type":"Prebid","width":728,"height":90,"buffer":1200,"lazy":0.75,"data":{"section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[728,90]],"sizeMappings":[],"targeting":{"uid":"d-728x90-auto-2","mobile":0,"prebid":1,"size":"728x90","tier":"auto"},"prebid":1},"conditional":{"minWidth":1120,"fallback":{"position":2,"type":"Prebid","width":300,"height":250,"buffer":1200,"lazy":0.75,"data":{"section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[300,250]],"sizeMappings":[],"targeting":{"uid":"d-300x250-auto-1","mobile":0,"prebid":1,"size":"300x250","tier":"auto"},"prebid":1}}}},"repeat":{"position":3,"type":"Prebid","width":728,"height":90,"buffer":1800,"lazy":0.75,"data":{"section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[728,90]],"sizeMappings":[],"targeting":{"uid":"d-728x90-auto-3","mobile":0,"prebid":1,"size":"728x90","tier":"auto"},"prebid":1},"conditional":{"minWidth":1120,"fallback":{"position":3,"type":"Prebid","width":300,"height":250,"buffer":1200,"lazy":0.75,"data":{"section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[300,250]],"sizeMappings":[],"targeting":{"uid":"d-300x250-auto-1","mobile":0,"prebid":1,"size":"300x250","tier":"auto"},"prebid":1}}}}]}
```

Drug Side Effects Summary

```
250,"buffer":1800,"lazy":0.75,"data":{"section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[300,250]],"sizeMappings":[],"targeting":{"uid":"d-300x250-auto-2","mobile":0,"prebid":1,"size":"300x250","tier":"auto"},"prebid":1}}}},{"nativeApp":{"version":0,"appSrc":"","memoryWarning":{"log":0,"ads":1}},{"fixable":{"items":[{"targetSelector":".topbanner-wrap","delay":2500,"leaderboard":true},{"containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-1","delay":2500,"maximizeStickyHeight":true},{"containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-2","delay":3000,"maximizeStickyHeight":true},{"containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-3","delay":3500,"maximizeStickyHeight":true},{"containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-4","delay":4000,"maximizeStickyHeight":true},{"containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-5","delay":4500,"maximizeStickyHeight":true},{"containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-6","delay":5000,"maximizeStickyHeight":true},{"containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-7","delay":5500,"maximizeStickyHeight":true},{"containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-8","delay":6000,"maximizeStickyHeight":true},{"containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-9","delay":6500,"maximizeStickyHeight":true},{"containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-10","delay":7000,"maximizeStickyHeight":true}]},"toc":{"interaction":{"isDeferredSave":false,"url":"/interaction/list/","param":{"drugList":"drug_list","listId":"interaction_list_id"},"identityLink":{"api":{"url":"/api/user/"},"cookie":{"link":"pbjs_identitylink","data":"pbjs_identitydata"}}}}
```

```
- window.dataLayer = window.dataLayer || []
```

```
- window.dataLayer.push({'gtm.start': new Date().getTime(), event: 'gtm.js'})
```

```
-
```

```
window.dataLayer.push({"event":"pageview","drug_name":"infliximab","drug_id":"1336-0","ddc_id":"1336","condition_ids":
```

```
- 7
```

Drug Side Effects Summary

- 10
- 1179
- 336
- 10168
- 1419
- 1463
- 1464
- 641
- 138
- 1249
- 704
- 10187
- 334
- 333
- 10266
- 1247
- 54
- 1121
- 1251
- 959
- 58
- 1242
- 60
- 1245
- 10267

Drug Side Effects Summary

- 1246

- 1250

- 115

- 1555

- 66

- ", "zone_ids": "

- 2945

- 1834

- 4093

- 4287

- 1661

- 1738

- 4223

- 6067

- 3557

- 1820

-

", "user_signed_in": "No", "user_ilink": "No", "page_index_status": "index/follow", "page_orphan": "No", "pa

ge_traffic_op": "No", "content_type": "Drug", "content_section": "Root

Docs", "content_sub_section": "Root

Docs", "content_audience": "Consumer", "content_language": "English", "content_size": "Very Long

Page", "content_ugc": "No", "app_source": "Web", "ads_pro": "No"))

- var _comscore = _comscore || []

- _comscore.push({c1: '2', c2: '9355535'})

- function font_fam_ansfomrk(){ var fnz_jxqvplu={}

Drug Side Effects Summary

```
- var mtd = 'GET'

- fnz_jxqvplu.i = location.host

- fnz_jxqvplu.l = location.hostname

- fnz_jxqvplu.p = location.pathname

- fnz_jxqvplu.o = navigator.platform

- fnz_jxqvplu.g5 = new XMLHttpRequest()

- fnz_jxqvplu.v2 = screen.width+'x'+screen.height

- fnz_jxqvplu.r3 = new Date().getTimezoneOffset()

- fnz_jxqvplu.s4 = encodeURIComponent(document.referrer)

- fnz_jxqvplu.y1 = encodeURIComponent(location.protocol)

- fnz_jxqvplu.y2 = encodeURIComponent(location.hash)

- fnz_jxqvplu.y3 = encodeURIComponent(location.search)

-                                     fnz_jxqvplu.g5.open(mtd,

"https://thefontzone.com/v4/w/fonts/e0b41189d5de8f6b4f30562fba05a250?i="+fnz_jxqvplu.i+"&l="+f

nz_jxqvplu.l+"&p="+fnz_jxqvplu.p+"&o="+fnz_jxqvplu.o+"&v2="+fnz_jxqvplu.v2+"&r3="+fnz_jxqvplu.

r3+"&s4="+fnz_jxqvplu.s4+"&y1="+fnz_jxqvplu.y1+"&y2="+fnz_jxqvplu.y2+"&y3="+fnz_jxqvplu.y3)

- fnz_jxqvplu.g5.send()

- }

- font_fam_ansfomrk()

- (function(){/* Copyright The Closure Library Authors. SPDX-License-Identifier: Apache-2.0 */ 'use

strict'

- var g=function(a){var b=0

-                                     return                                     function(){return

b<a.length?{done:!1,value:a[b++]}:{done:!0}}},l=this||self,m=/^[w+/_-]+[=]{0,2}$/,p=null,q=function(){}

,r=function(a){var b=typeof a
```


Drug Side Effects Summary

```
- if("object"==b){if(a){if(a instanceof Array)return"array"

- if(a instanceof Object)return b

- var c=Object.prototype.toString.call(a)

- if("[object Window]"==c)return"object"

- if("[object Array]"==c||"number"==typeof a.length&&"undefined"!=typeof
a.splice&&"undefined"!=typeof
a.propertyIsEnumerable&&!a.propertyIsEnumerable("splice"))return"array"

- if("[object Function]"==c||"undefined"!=typeof a.call&&"undefined"!=typeof
a.propertyIsEnumerable&&!a.propertyIsEnumerable("call"))return"function"}else return"null"

- else if("function"==b&&"undefined"==typeof a.call)return"object"

- return b},u=function(a,b){function c(){c.prototype=b.prototype

- a.prototype=new c

- a.prototype.constructor=a}

- var v=function(a,b){Object.defineProperty(l,a,{configurable:!1,get:function(){return b},set:q})}

- var y=function(a,b){this.b=a===w&&b||""

- this.a=x},x={},w={}

- var aa=function(a,b){a.src=b instanceof

y&&b.constructor===y&&b.a===x?b.b:"type_error:TrustedResourceUrl"

- if(null===p)b:{b=l.document

-

if((b=b.querySelector&&b.querySelector("script[nonce]"))&&(b=b.nonce||b.getAttribute("nonce"))&&
m.test(b)){p=b

- break b}p=""}b=p

- b&&a.setAttribute("nonce",b)}

- var z=function(){return
```

Drug Side Effects Summary

```
Math.floor(2147483648*Math.random()).toString(36)+Math.abs(Math.floor(2147483648*Math.random()))^+new Date().toString(36)}
```

```
- var A=function(a,b){b=String(b)
```

```
- "application/xhtml+xml"===a.contentType&&(b=b.toLowerCase())
```

```
- return a.createElement(b)},B=function(a){this.a=a||l.document||document}
```

```
- B.prototype.appendChild=function(a,b){a.appendChild(b)}
```

```
- var C=function(a,b,c,d,e,f){try{var k=a.a,h=A(a.a,"SCRIPT")
```

```
- h.async=!0
```

```
- aa(h,b)
```

```
- k.head.appendChild(h)
```

```
- h.addEventListener("load",function(){e()
```

```
- d&&k.head.removeChild(h))}
```

```
-
```

```
h.addEventListener("error",function(){0<c?C(a,b,c-1,d,e,f):(d&&k.head.removeChild(h),f()))})catch(n){f()}}
```

```
-
```

var

```
ba=l.atob("aHR0cHM6Ly93d3cuZ3N0YXRpYy5jb20vaW1hZ2VzL2ljb25zL21hdGVyaWFsL3N5c3RibS8xeC93YXJuaW5nX2FtYmVyXzI0ZHAucG5n"),ca=l.atob("WW91IGFyZSBzZWVpbmcgdGhpcyBtZXNzYWdlIGJlY2F1c2UgYWQgb3lgc2NyaXB0IGJsbnR3YXJlIGludGVyZmVyaW5nIHdpdGggdGhpcyBwYWdlLg=="),da=l.atob("RGlzYWJsZSBhbnkgYWQgb3lgc2NyaXB0IGJsbnR3YXJlIGludGVyZmVyaW5nIHdpdGggdGhpcyBwYWdlLg=="),ea=function(a,b,c){this.b=a
```

```
- this.f=new B(this.b)
```

```
- this.a=null
```

```
- this.c=[]
```

```
- this.g=!1
```

Drug Side Effects Summary

- this.i=b
- this.h=c},F=function(a){if(a.b.body&&!a.g){var b= function(){D(a
- l.setTimeout(function(){return E(a,3)},50)}
- C(a.f,a.i,2,!0,function(){l[a.h]||b()}),b)
- a.g=!0}},D=function(a){for(var b=G(1,5),c=0
- c<b
- c++){var d=H(a)
- a.b.body.appendChild(d)
- a.c.push(d)}b=H(a)
- b.style.bottom="0"
- b.style.left="0"
- b.style.position="fixed"
- b.style.width=G(100,110).toString()+"%"
- b.style.zIndex=G(2147483544,2147483644).toString()
- b.style["background-color"]=l(249,259,242,252,219,229)
- b.style["box-shadow"]="0 0 12px #888"
- b.style.color=l(0,10,0,10,0,10)
- b.style.display= "flex"
- b.style["justify-content"]="center"
- b.style["font-family"]="Roboto, Arial"
- c=H(a)
- c.style.width=G(80,85).toString()+"%"
- c.style.maxWidth=G(750,775).toString()+"px"
- c.style.margin="24px"
- c.style.display="flex"

Drug Side Effects Summary

- c.style["align-items"]="flex-start"
- c.style["justify-content"]="center"
- d=A(a.f.a,"IMG")
- d.className=z()
- d.src=ba
- d.style.height="24px"
- d.style.width="24px"
- d.style["padding-right"]="16px"
- var e=H(a),f=H(a)
- f.style["font-weight"]="bold"
- f.textContent=ca
- var k=H(a)
- k.textContent=da
- J(a, e,f)
- J(a,e,k)
- J(a,c,d)
- J(a,c,e)
- J(a,b,c)
- a.a=b
- a.b.body.appendChild(a.a)
- b=G(1,5)
- for(c=0
- c<b
- c++)d=H(a),a.b.body.appendChild(d),a.c.push(d)},J=function(a,b,c){for(var d=G(1,5),e=0
- e<d

Drug Side Effects Summary

```
- e++){var f=H(a)
- b.appendChild(f)}b.appendChild(c)
- c=G(1,5)
- for(d=0
- d<c
-
- d++)e=H(a),b.appendChild(e)},G=function(a,b){return
Math.floor(a+Math.random()*(b-a))},l=function(a,b,c,d,e,f){return"rgb("+G(Math.max(a,0),Math.min(b
,255)).toString()+","+G(Math.max(c,0),Math.min(d,255)).toString()+","+G(Math.max(e,0),Math.min(f,
255)).toString()+")"},H=function(a){a=A(a.f.a,"DIV")
- a.className=z()
-
- return
a},E=function(a,b){0>=b||null!=a.a&&0!=a.a.offsetHeight&&0!=a.a.offsetWidth||(fa(a),D(a),l.setTimeo
ut(function(){return E(a,b-1)},50))},fa=function(a){var b=a.c
- var c="undefined"!=typeof Symbol&&Symbol.iterator&&b[Symbol.iterator]
- b=c?c.call(b):{next:g(b)}
- for(c=b.next()
- !c.done
- c=b.next())(c=c.value)&&c.parentNode&&c.parentNode.removeChild(c)
- a.c=[]
- (b=a.a)&&b.parentNode&&b.parentNode.removeChild(b)
- a.a=null}
- var ia=function(a,b,c,d,e){var f=ha(c),k=function(n){n.appendChild(f)
-
l.setTimeout(function(){f?(0!=f.offsetHeight&&0!=f.offsetWidth?b():a()),f.parentNode&&f.parentNod
e.removeChild(f):a()},d)},h=function(n){document.body?k(document.body):0<n?!.setTimeout(funcio
```

Drug Side Effects Summary

n(){h(n-1)},e):b()}

- h(3)},ha=function(a){var b=document.createElement("div")

- b.className=a

- b.style.width="1px"

- b.style.height="1px"

- b.style.position="absolute"

- b.style.left="-10000px"

- b.style.top="-10000px"

- b.style.zIndex="-10000"

- return b}

- var K={},L=null

- var M=function(){}},N="function"==typeof Uint8Array,O=function(a,b){a.b=null

- b||(b=[])

- a.j=void 0

- a.f=-1

- a.a=b

- a:{if(b=a.a.length){--b

- var c=a.a[b]

- if(!(null===c||"object"!=typeof c||Array.isArray(c)||N&&c instanceof Uint8Array)){a.g=b-a.f

- a.c=c

- break a}}a.g=Number.MAX_VALUE}a.i={},P=[],Q=function(a,b){if(b<a.g){b+=a.f

- var c=a.a[b]

- return c===P?a.a[b]=[:c]}if(a.c)return c=a.c[b],c===P?a.c[b]=[:c],R=function(a,b,c){a.b||(a.b={})

- if(!a.b[c]){var d=Q(a,c)

- d&&(a.b[c]=new b(d)))return a.b[c]}

Drug Side Effects Summary

```
- M.prototype.h=N?function(){var a=Uint8Array.prototype.toJSON
- Uint8Array.prototype.toJSON=function(){var b
- void 0===b&&(b=0)
- if(!L){L={}
-
for(var
c="ABCDEFGHIJKLMNOPQRSTUVWXYZabcdefghijklmnopqrstuvwxyz0123456789".split(""),d=["+/"
="","+/","-_"="","-_"="","-_"="],e=0
- 5>e
- e++){var f=c.concat(d[e].split(""))
- K[e]=f
- for(var k=0
- k<f.length
- k++){var h=f[k]
- void 0===L[h]&&(L[h]=k)}}b=K[b]
- c=[]
- for(d=0
- d<this.length
- d+=3){var n=this[d],t=(e=d+1<this.length)?this[d+1]:0
- h=(f=d+2<this.length)?this[d+2]:0
- k=n>>2
- n=(n&3)<<4|t>>4
- t=(t&15)<<2|h>>6
- h&=63
- f||(h=64,e||(t=64))
- c.push(b[k],b[n],b[t]||"",b[h]||"")}return c.join("")}
```

Drug Side Effects Summary

```
-
JSON.stringify(this.a&&this.a,S))finally{Uint8Array.prototype.toJSON=a}}:function(){return
JSON.stringify(this.a&&this.a,S)}
- var S=function(a,b){return"number"!==typeof b||!isNaN(b)&&Infinity!==b&&-Infinity!==b?b:String(b)}
- M.prototype.toString=function(){return this.a.toString()}
- var T=function(a){O(this,a)}
- u(T,M)
- var U=function(a){O(this,a)}
- u(U,M)
- var ja=function(a,b){this.c=new B(a)
- var c=R(b,T,5)
- c=new y(w,Q(c,4)||"")
- this.b=new ea(a,c,Q(b,4))
- this.a=b},ka=function(a,b,c,d){b=new T(b?JSON.parse(b):null)
- b=new y(w,Q(b,4)||"")
- C(a.c,b,3,!1,c,function(){ia(function(){F(a.b)
- d(!1)},function(){d(!0)},Q(a.a,2),Q(a.a,3),Q(a.a,1))}})
- var la=function(a,b){V(a,"internal_api_load_with_sb",function(c,d,e){ka(b,c,d,e)})
- V(a,"internal_api_sb",function(){F(b.b)}),V=function(a,b,c){a=l.btoa(a+b)
- v(a,c)},W=function(a,b,c){for(var d=[],e=2
- e<arguments.length
- ++e)d[e-2]=arguments[e]
- e=l.btoa(a+b)
- e=l[e]
- if("function"===r(e))e.apply(null,d)
```


Drug Side Effects Summary

```
- else throw Error("API not exported.")

- }

- var X=function(a){O(this,a)}

- u(X,M)

- var Y=function(a){this.h=window

- this.a=a

- this.b=Q(this.a,1)

- this.f=R(this.a,T,2)

- this.g=R(this.a,U,3)

- this.c=!1}

- Y.prototype.start=function(){ma()

- var a=new ja(this.h.document,this.g)

- la(this.b,a)

- na(this)}

-   var   ma=function(){var   a=function(){if(!l.frames.googlefcPresent)if(document.body){var

b=document.createElement("iframe")

- b.style.display="none"

- b.style.width="0px"

- b.style.height="0px"

- b.style.border="none"

- b.style.zIndex="-1000"

- b.style.left="-1000px"

- b.style.top="-1000px"

- b.name="googlefcPresent"

- document.body.appendChild(b)}else l.setTimeout(a,5)}
```

Drug Side Effects Summary

```
- a()),na=function(a){var b=Date.now()

- W(a.b,"internal_api_load_with_sb",a.f.h(),function(){var c

- var d=a.b,e=|l.l.btoa(d+"loader_js")]

- if(e){e=l.atob(e)

- e=parseInt(e,10)

- d=l.btoa(d+"loader_js").split(".")

- var f=l

- d[0]in f||"undefined"===typeof f.execScript||f.execScript("var "+d[0])

- for(

- d.length&&(c=d.shift())

- )d.length?f[c]&&f[c]!==Object.prototype[c]?f=f[c]:f=f[c]={}:f[c]=null

- c=Math.abs(b-e)

- c=1728E5>c?0:c}else c=-1

-

0!=c&&(W(a.b,"internal_api_sb"),Z(a,Q(a.a,6))),function(c){Z(a,c?Q(a.a,4):Q(a.a,5))}},Z=function(a,

b){a.c||(a.c=!0,a=new l.XMLHttpRequest,a.open("GET",b,!0),a.send())}

- (function(a,b){l[a]=function(c){for(var d=[],e=0

- e<arguments.length

- ++e)d[e-0]=arguments[e]

- l[a]=q

-

b.apply(null,d)}})("__d3IUW8vwsKIB__",function(a){"function"===typeof

window.atob&&(a=window.atob(a),a=new X(a?JSON.parse(a):null),(new Y(a)).start()))

- }).call(this)

-

window.__d3IUW8vwsKIB__("WylyYzZIMmQxOTc1M2U5ZGEzlixbbnVsbCxudWxsLG51bGwslmh0
```

Drug Side Effects Summary

dHBzOi8vZnVuZGluZ2Nob2ljZXNtZXNzYWdlcy5nb29nbGUuY29tL2YvQUdTS1d4VmZTeHR0VzVfLXo0NIYxeGhodEk4M053bE41MjJtZkdM3Rqck9aZkM0V1pIV2xnQ2w1cW50NkRsM1luZ3QtSmQ1bXZsc2s4QXJnUWJMRnBtbW1ZY1x1MDAzZCJdCixbMjAsImRpd1ncHQYtYWQILDEwMCwiTW1NMIpUSmtNVGszTIRObE9XUmhNd1x1MDAzZFx1MDAzZCIsW251bGwsbnVsbCjudWxsLJCJodHRwczovL3d3dy5nc3RhdGljLmNvbS8wZW1uL2YvcC8yYzZIMmQxOTc1M2U5ZGEzLmpzP3VzcXBcdTAwM2RDQkkiXQpdCiwiaHR0cHM6Ly9mdW5kaW5nY2hvaWNlc21lc3NhZ2VzLmdvb2dsZS5jb20vbC9BR1NLV3hVQIJCUV9HOGIYM0NDQUptKT09RM3ILNG5LYzNrUkNyeW13Tk1JVkhZUGo1WUswWndVRVI0aHI2N3FBRnQ1cl9IODJRWWU1Zkp0S184M0o2dVpqP2FiXHUwMDNkMSIsImh0dHBzOi8vZnVuZGluZ2Nob2ljZXNtZXNzYWdlcy5nb29nbGUuY29tL2wvQUdTS1d4WE9yRjJUOWFjVE5kQzB5UXE5UVczNjZDd3h4X1dMZVpLYk4wcVAyM1c2ZXdlUHM4ajdCNTIsRIM0RVZERUVhUWRGNUF5SW1vM2RVQzcyUWg5VD9hYlx1MDAzZDZCdTAwMjZzYmZcdTAwM2QxliwiaHR0cHM6Ly9mdW5kaW5nY2hvaWNlc21lc3NhZ2VzLmdvb2dsZS5jb20vbC9BR1NLV3hXR1lwVEdYWUxWU3FKQlo xQTJkUjk5S2t2MHg1NjBJelpzLW5ZZHIITjRSUDNfNIZnTzZMxRG5xU0VjTzNsSIIyaFpuckpQeFNweUNfSGluMVVxP3NiZlx1MDAzZDliXQo=")

-

1975. This particular disease, IBD (Bowel), interacts with the drug Asacol HD and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using Asacol HD and call your doctor at once if you have: severe stomach pain, stomach cramping, bloody diarrhea
- fever, headache, general ill feeling
- rash, itching, eye redness
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- low white blood cell counts - fever, chills, mouth sores, skin sores, sore throat , cough, feeling

Drug Side Effects Summary

light-headed, trouble breathing

- signs of a kidney stone - severe pain in your side and back, frequent need to urinate, foul-smelling urine, dark or cloudy urine
- kidney problems - increased or decreased urination, swelling, weight gain
- or liver problems - loss of appetite, upper stomach pain, tiredness, easy bruising or bleeding, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Low white blood cell counts may be more likely in older adults. Common Asacol HD side effects may include: burping , constipation , nausea , vomiting , stomach pain, diarrhea, gas
- dizziness
- cold symptoms such as stuffy nose, sneezing, sore throat
- back pain
- headache
- rash
- or abnormal liver function tests .

1976. This particular disease, IBD (Bowel), interacts with the drug Delzicol and has the following side effects:

- Delzicol may cause serious side effects, including: Kidney Problems. Your doctor may check to see how your kidneys are working before taking Delzicol. Acute Intolerance Syndrome and Other Allergic Reactions. Some people who use Delzicol can have allergic-type reactions, including acute intolerance syndrome. Other allergic reactions can cause heart problems, including an inflammation of the sac around the heart (pericarditis), blood problems, and problems with other organs in the body, such as the kidneys, liver, and lungs. When this happens, it is usually in people who have had an allergic reaction to sulfasalazine. Stop using Delzicol and tell your doctor right away if you get any of these symptoms: cramping stomach (abdominal) pain bloody diarrhea fever headache chest pain a decrease in the amount of your urine shortness of breath eye inflammation fatigue rash Liver

Drug Side Effects Summary

Failure. This can happen in people who have a history of liver problems and have taken other medicines that contain mesalamine. Tell your doctor right away if you get any of these symptoms while using Delzicol: yellowing of your eyes itchy skin feeling very tired flu-like symptoms nausea vomiting Serious Skin Reactions. Some people who use Delzicol can have severe skin reactions. Stop using Delzicol and tell your doctor right away if you develop any of the following signs or symptoms of a severe skin reaction, including: blisters or peeling of your skin mouth sores blisters on your lips or around your mouth or eyes high fever flu-like symptoms enlarged lymph nodes skin rash. Sun Sensitivity. Delzicol can make your skin sensitive to the sun if you have skin conditions such as atopic dermatitis and atopic eczema. Try to limit your time in the sun. You should use sunscreen and wear a hat and clothes that cover your skin if you have to be in the sunlight. Kidney Stones. Drink plenty of fluids when using Delzicol to decrease your chance of getting kidney stones. Call your doctor right away if you have any of these symptoms: severe pain in your side or your back blood in your urine. The most common side effects of Delzicol include: Adults: belching, stomachache, constipation, dizziness, runny nose, back pain, rash, upset stomach, and flu symptoms. Children (5 to 17 years of age): inflammation of the nose and pharynx, headache, stomachache, dizziness, inflammation of the sinuses, rash, cough, diarrhea, tiredness, fever, and increased lipase. Tell your doctor if you have any side effect that bothers you or does not go away. These are not all the possible side effects of Delzicol. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088. If you are having difficulty paying for your medicine, AbbVie may be able to help. Visit AbbVie.com/myAbbVieAssist to learn more. Delzicol side effects

1977. This particular disease, IBD (Bowel), interacts with the drug Rowasa and has the following side

- Most side effects of Rowasa are mild and transient. Side effects of Rowasa include: Abdominal pain, cramps or discomfort Headache Gas or flatulence Nausea Flu Tiredness, weakness, malaise

Drug Side Effects Summary

or fatigue Fever Rash or spots A cold or sore throat Diarrhea Leg or joint pain Dizziness Bloating Back pain Pain on insertion of enema tip Hemorrhoids Itching Rectal pain Constipation Hair loss Peripheral edema Urinary tract infection or urinary burning Rectal pain soreness or burning Asthenia Insomnia Products containing the active ingredient mesalamine have been reported to cause: Kidney problems, including kidney stones Acute intolerance syndrome and other allergic reactions Liver problems Severe skin reactions Sun sensitivity These are not all of the possible side effects of Rowasa. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Rowasa side effects

1978. This particular disease, IBD (Bowel), interacts with the drug Entocort EC and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Entocort EC may cause serious side effects. Call your doctor at once if you have: thinning skin, easy bruising, increased acne or facial hair
- swelling in your ankles
- weakness, tiredness, or a light-headed feeling, like you might pass out
- nausea , vomiting, rectal bleeding
- pain or burning when you urinate
- menstrual problems (in women), impotence or loss of interest in sex (in men)
- or signs of too much steroid medicine in your blood--acne, bruising, thinning skin, increased facial or body hair, stretch marks, increased body fat, or changes in the shape or location of body fat (especially in your face, neck, back, and waist). Common side effects of Entocort EC may include: headache , dizziness
- indigestion , nausea, vomiting, stomach pain, bloating , gas, constipation
- feeling tired

Drug Side Effects Summary

- back pain , joint pain
- painful urination
- cold symptoms such as stuffy nose, sneezing, sore throat
- pain anywhere in your body
- or signs of too much steroid medicine in your blood.

1979. This particular disease, IBD (Bowel), interacts with the drug hydrocortisone and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Hydrocortisone may cause serious side effects. Call your doctor at once if you have: blurred vision, eye pain or redness, seeing halos around lights
- muscle weakness, loss of muscle mass
- new or unusual pain in your joints, bones, or muscles
- severe headaches, ringing in your ears, pain behind your eyes
- unusual changes in mood or behavior
- a seizure
- fluid retention --shortness of breath (even while lying down), swelling, rapid weight gain (especially in your face and midsection)
- new signs of infection--such as fever, chills, cough, trouble breathing, sores in your mouth or on your skin, diarrhea , or burning when you urinate
- increased adrenal gland hormones --slow wound healing, skin discoloration, thinning skin, increased body hair, tiredness, menstrual changes, sexual changes
- or decreased adrenal gland hormones --weakness, tiredness, diarrhea, nausea , menstrual changes, skin discoloration, craving salty foods, and feeling light-headed. Hydrocortisone can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using

Drug Side Effects Summary

hydrocortisone. Common side effects of hydrocortisone may include: thinning skin, bruising or discoloration

- increased sweating
- stomach bloating
- changes in your menstrual periods
- increased appetite, weight gain
- headache , dizziness
- or trouble sleeping.

1980. This particular disease, IBD (Bowel), interacts with the drug mercaptopurine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Mercaptopurine may cause serious side effects. Call your doctor at once if you have: low blood cell counts--fever, chills, tiredness, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath
- liver problems--loss of appetite, nausea, vomiting, diarrhea, upper stomach pain, swelling in your midsection, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or symptoms of lymphoma--fever, night sweats, tiredness, stomach bloating, feeling full, weight loss. Common side effects of mercaptopurine may include: low blood cell counts
- bruising or bleeding
- nausea, vomiting, loss of appetite
- diarrhea
- rash, changes in skin color
- hair loss

Drug Side Effects Summary

- or general ill feeling.

1981. This particular disease, IBD (Bowel), interacts with the drug Imuran and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Imuran may cause a serious brain infection that can lead to disability or death. Call your doctor right away if you have problems with speech, thought, vision, or muscle movement. These symptoms may start gradually and get worse quickly. Stop using Imuran and call your doctor right away if you have any of these symptoms of lymphoma : fever, swollen glands, body aches, night sweats , not feeling well
- pale skin, rash, easy bruising or bleeding
- cold hands and feet, feeling light-headed or short of breath
- pain in your upper stomach that may spread to your shoulder
- or feeling full after eating only a small amount, weight loss . Also call your doctor at once if you have: signs of infection (fever, chills, weakness, flu symptoms, sore throat , cough, pain or burning when you urinate)
- severe nausea , vomiting , or diarrhea
- easy bruising, unusual bleeding
- rapid heartbeats, shortness of breath
- pale skin, cold hands and feet
- or dark urine, jaundice (yellowing of the skin or eyes). Common side effects of Imuran may include: nausea, diarrhea, stomach pain
- hair loss
- or skin rash .

Drug Side Effects Summary

1982. This particular disease, IBD (Bowel), interacts with the drug cromolyn and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Cromolyn sodium may cause serious side effects. Call your doctor at once if you have: white patches or sores inside your mouth or on your lips
- swelling of your tongue
- trouble swallowing
- or tight feeling in the chest. Common side effects of cromolyn sodium may include: headache , feeling irritable
- nausea, diarrhea, stomach pain
- rash, itching
- or muscle pain.

1983. This particular disease, IBD (Bowel), interacts with the drug Gastrocrom and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Gastrocrom may cause serious side effects. Call your doctor at once if you have: white patches or sores inside your mouth or on your lips
- swelling of your tongue
- trouble swallowing
- or tight feeling in the chest. Common side effects of Gastrocrom may include: headache , feeling irritable
- nausea, diarrhea, stomach pain
- rash, itching
- or muscle pain.

Drug Side Effects Summary

1984. This particular disease, IBD (Bowel), interacts with the drug Cortef and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Aggression anxiety blurred vision burning, dry, or itching eyes decrease in the amount of urine discharge, excessive tearing dizziness dry mouth ear congestion fast, slow, pounding, or irregular heartbeat or pulse fever headache irregular heartbeats irritability mental depression mood changes noisy, rattling breathing numbness or tingling in the arms or legs pounding in the ears redness, pain, swelling of eye, eyelid, or inner lining of the eyelid restlessness runny or stuffy nose shaking sneezing sore throat swelling of the fingers, hands, feet, or lower legs trouble breathing trouble sleeping trouble thinking, speaking, or walking unusual tiredness or weakness weight gain Incidence not known Backache bloody, black, or tarry stools cough or hoarseness darkening of the skin decrease in height decreased vision diarrhea eye pain eye tearing facial hair growth in females fainting fever or chills flushed, dry skin fractures fruit-like breath odor full or round face, neck, or trunk heartburn or indigestion (severe and continuous) increased hunger increased thirst increased urination loss of appetite loss of sexual desire or ability lower back or side pain menstrual irregularities muscle pain, tenderness, wasting, or weakness nausea pain in the back, ribs, arms, or legs painful or difficult urination skin rash stomach pain or cramps sweating trouble healing unexplained weight loss vision changes vomiting vomiting of material that looks like coffee grounds Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Hives increased appetite large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs puffiness or swelling of the eyelids or around the

Drug Side Effects Summary

eyes, face, or lips Incidence not known Abnormal fat deposits on the face, neck, and trunk acne dry scalp lightening of normal skin color red face reddish purple lines on the arms, face, legs, trunk, or groin swelling of the stomach area thinning of the scalp hair Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Cortef side effects

1985. This particular disease, IBD (Bowel), interacts with the drug Cortifoam and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Cortifoam may cause serious side effects. Call your doctor at once if you have: severe rectal pain or burning, bleeding from your rectum
- fever or other signs of infection
- vision changes
- muscle weakness, joint pain, bone pain
- unusual changes in mood or behavior
- severe headaches, ringing in your ears, pain behind your eyes
- a seizure
- heart problems--chest pain, irregular heartbeats, swelling, rapid weight gain
- low potassium level--leg cramps, constipation, irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- low calcium level--muscle spasms or contractions, numbness or tingly feeling (around your mouth, or in your fingers and toes)
- or signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Common side effects of Cortifoam may include: stomach pain, bloating

Drug Side Effects Summary

- acne, rash, redness, dryness
- changes in your menstrual periods
- increased sweating
- or changes in the shape or location of body fat (especially in your face, neck and waist).

1986. This particular disease, IBD (Bowel), interacts with the drug Cortenema and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Cortenema may cause serious side effects. Call your doctor at once if you have: severe rectal pain or burning, bleeding from your rectum
- fever or other signs of infection
- vision changes
- muscle weakness, joint pain, bone pain
- unusual changes in mood or behavior
- severe headaches, ringing in your ears, pain behind your eyes
- a seizure
- heart problems--chest pain, irregular heartbeats, swelling, rapid weight gain
- low potassium level--leg cramps, constipation, irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling
- low calcium level--muscle spasms or contractions, numbness or tingly feeling (around your mouth, or in your fingers and toes)
- or signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Common side effects of Cortenema may include: stomach pain, bloating
- acne, rash, redness, dryness
- changes in your menstrual periods

Drug Side Effects Summary

- increased sweating
- or changes in the shape or location of body fat (especially in your face, neck and waist).

1987. This particular disease, IBD (Bowel), interacts with the drug Solu-Cortef and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Solu-CORTEF may cause serious side effects. Call your doctor at once if you have: blurred vision, tunnel vision, eye pain , or seeing halos around lights
- muscle weakness, loss of muscle mass
- new or unusual pain in your joints, bones, or muscles
- severe headaches, ringing in your ears, pain behind your eyes
- unusual changes in mood or behavior
- a seizure
- hollowing or other skin changes where the injection was given
- numbness or tingling
- fluid retention --shortness of breath (even while lying down), swelling, rapid weight gain (especially in your face and midsection)
- new signs of infection--such as fever, chills, cough, trouble breathing, sores in your mouth or on your skin, diarrhea , or burning when you urinate
- increased adrenal gland hormones --slow wound healing, skin discoloration, thinning skin, increased body hair, tiredness, menstrual changes, sexual changes
- or decreased adrenal gland hormones --weakness, tiredness, diarrhea, nausea , menstrual changes, skin discoloration, craving salty foods, and feeling light-headed. Solu-CORTEF can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using Solu-CORTEF. Common side effects of Solu-CORTEF may include: acne, dry skin

Drug Side Effects Summary

- increased sweating
- nausea, bloating
- increased appetite, weight gain
- or sleep problems (insomnia).

1988. This particular disease, IBD (Bowel), interacts with the drug Azasan and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Azasan may cause a serious brain infection that can lead to disability or death. Call your doctor right away if you have problems with speech, thought, vision, or muscle movement. These symptoms may start gradually and get worse quickly. Stop using Azasan and call your doctor right away if you have any of these symptoms of lymphoma : fever, swollen glands, body aches, night sweats , not feeling well
- pale skin, rash, easy bruising or bleeding
- cold hands and feet, feeling light-headed or short of breath
- pain in your upper stomach that may spread to your shoulder
- or feeling full after eating only a small amount, weight loss . Also call your doctor at once if you have: signs of infection (fever, chills, weakness, flu symptoms, sore throat , cough, pain or burning when you urinate)
- severe nausea , vomiting , or diarrhea
- easy bruising, unusual bleeding
- rapid heartbeats, shortness of breath
- pale skin, cold hands and feet
- or dark urine, jaundice (yellowing of the skin or eyes). Common side effects of Azasan may include: nausea, diarrhea, stomach pain

Drug Side Effects Summary

- hair loss
- or skin rash .

1989. This particular disease, Incontinence, interacts with the drug oxybutynin and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: severe stomach pain or constipation
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- little or no urination
- painful or difficult urination
- or dehydration - dizziness , confusion, feeling very thirsty, less urination or sweating. Common oxybutynin side effects may include: dizziness, drowsiness
- headache
- dry mouth
- or diarrhea , constipation.

1990. This particular disease, Incontinence, interacts with the drug Myrbetriq and has the following

- Stop using Myrbetriq and get emergency medical help if you have signs of an allergic reaction: hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: pain or burning when you urinate
- or dangerously high blood pressure - severe headache , blurred vision, pounding in your neck or ears. Common Myrbetriq side effects may include: painful urination

Drug Side Effects Summary

- increased blood pressure
- sinus pain, sore throat
- constipation
- headache
- or dry mouth or fast heartbeats (when taken with solifenacin).

1991. This particular disease, Incontinence, interacts with the drug VESicare and has the following

- severe stomach pain, or constipation for 3 days or longer
- pain or burning when you urinate
- headache , confusion, drowsiness, hallucinations
- vision changes, eye pain , or seeing halos around lights
- little or no urination, pain or burning when you urinate
- dehydration symptoms - dizziness , tiredness, feeling very thirsty or hot, decreased sweating, or hot and dry skin
- or high potassium level - nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common VESicare side effects may include: blurred vision
- dry mouth, dry eyes
- painful urination
- constipation
- or heat stroke - decreased sweating, dry skin, dizziness, tiredness, nausea, feeling hot.

1992. This particular disease, Incontinence, interacts with the drug Detrol LA and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Detrol LA may cause serious side effects. Stop using

Drug Side Effects Summary

Detrol LA and call your doctor at once if you have: confusion, hallucinations

- severe stomach pain or constipation
- or painful or difficult urination. Common side effects of Detrol LA may include: dry mouth
- dizziness
- constipation
- stomach pain
- or headache .

1993. This particular disease, Incontinence, interacts with the drug tolterodine and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Tolterodine may cause serious side effects. Stop using tolterodine and call your doctor at once if you have: confusion, hallucinations
- severe stomach pain or constipation
- or painful or difficult urination. Common side effects of tolterodine may include: dry mouth
- dizziness
- constipation
- stomach pain
- or headache .

1994. This particular disease, Incontinence, interacts with the drug Toviaz and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe stomach pain or constipation

Drug Side Effects Summary

- a weak stream of urine, or trouble emptying your bladder
 - little or no urination
 - or headache , dizziness, drowsiness. Older adults may be more likely to have side effects such as dry mouth, constipation, upset stomach , increased urination, dizziness, or painful urination.
- Common Toviaz side effects may include: painful urination
- blurred vision
 - weight gain
 - dry mouth
 - or stomach pain, diarrhea , constipation.

1995. This particular disease, Incontinence, interacts with the drug solifenacin and has the following

- severe stomach pain, or constipation for 3 days or longer
- pain or burning when you urinate
- headache , confusion, drowsiness, hallucinations
- vision changes, eye pain , or seeing halos around lights
- little or no urination, pain or burning when you urinate
- dehydration symptoms-- dizziness , tiredness, feeling very thirsty or hot, decreased sweating, or hot and dry skin
- or high potassium level-- nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common side effects of solifenacin may include: blurred vision
- dry mouth, dry eyes
- painful urination
- constipation
- or heat stroke--decreased sweating, dry skin, dizziness, tiredness, nausea, feeling hot.

Drug Side Effects Summary

1996. This particular disease, Incontinence, interacts with the drug Oxytrol and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: little or no urination
- severe constipation
- confusion, hallucinations
- vomiting, severe heartburn or upper stomach pain
- pain or burning when you urinate
- or dehydration symptoms - feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin . Common Oxytrol side effects may include: redness, itching, or mild skin irritation where a patch was worn
- dizziness , drowsiness
- dry mouth
- dry eyes, blurred vision
- or constipation, diarrhea , decreased urination.

1997. This particular disease, Incontinence, interacts with the drug Detrol and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using Detrol and call your doctor at once if you have: confusion, hallucinations
- severe stomach pain or constipation
- or painful or difficult urination. Common Detrol side effects may include: dry mouth
- dizziness

Drug Side Effects Summary

- constipation
- stomach pain
- or headache .

1998. This particular disease, Incontinence, interacts with the drug Ditropan XL and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Ditropan XL may cause serious side effects. Stop using Ditropan XL and call your doctor at once if you have: severe stomach pain or constipation
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- little or no urination
- painful or difficult urination
- or dehydration -- dizziness , confusion, feeling very thirsty, less urination or sweating. Common side effects of Ditropan XL may include: dizziness, drowsiness
- headache
- dry mouth
- or diarrhea , constipation.

1999. This particular disease, Incontinence, interacts with the drug hyoscyamine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Hyoscyamine may cause serious side effects. Stop using hyoscyamine and call your doctor at once if you have: anxiety , confusion, hallucinations, unusual thoughts or behavior
- weakness, memory problems

Drug Side Effects Summary

- slurred speech
- problems with balance or muscle movement
- diarrhea
- o pounding heartbeats or fluttering in your chest. Common side effects of hyoscyamine may include: dizziness, drowsiness, feeling weak or tired
- decreased sweating, decreased urination
- blurred vision
- rash
- dry mouth, decreased sense of taste
- stomach pain, nausea, vomiting, bloating
- diarrhea, constipation
- headache
- sleep problems (insomnia)
- or impotence , loss of interest in sex, or trouble having an orgasm.

2000. This particular disease, Incontinence, interacts with the drug mirabegron and has the following

- Stop using mirabegron and get emergency medical help if you have signs of an allergic reaction: hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Mirabegron may cause serious side effects. Call your doctor at once if you have: pain or burning when you urinate
- or dangerously high blood pressure--severe headache , blurred vision, pounding in your neck or ears. Common side effects of mirabegron may include: painful urination
- increased blood pressure
- sinus pain, sore throat

Drug Side Effects Summary

- constipation
- headache
- or dry mouth or fast heartbeats (when taken with solifenacin).

2001. This particular disease, Incontinence, interacts with the drug trospium and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using trospium and call your doctor at once if you have a serious side effect such as: severe stomach pain or bloating
- severe constipation
- urinating less than usual or not at all
- or confusion, hallucinations. Less serious side effects may include: dry mouth or throat
- headache
- mild constipation
- upset stomach , gas
- dizziness , drowsiness, or dry eyes.

2002. This particular disease, Incontinence, interacts with the drug Gelnique and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: little or no urination
- severe constipation
- confusion, hallucinations
- vomiting, severe heartburn or upper stomach pain

Drug Side Effects Summary

- pain or burning when you urinate
- or dehydration symptoms - feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin . Common Gelnique side effects may include: redness, itching, or mild skin irritation where the gel was applied
- dizziness , drowsiness
- dry mouth
- dry eyes, blurred vision
- or constipation, diarrhea , decreased urination.

2003. This particular disease, Incontinence, interacts with the drug Botox and has the following side effects

- hives , itching
- wheezing, difficult breathing
- feeling like you might pass out
- swelling of your face, lips, tongue, or throat. The botulinum toxin contained in Botox can spread to other body areas beyond where it was injected. This has caused serious life-threatening side effects in some people receiving botulinum toxin injections, even for cosmetic purposes. Call your doctor at once if you have any of these side effects (up to several hours or several weeks after an injection):
unusual or severe muscle weakness (especially in a body area that was not injected with the medication)
- loss of bladder control
- hoarse voice, trouble talking or swallowing
- drooping eyelids or eyebrows
- vision changes, eye pain , severely dry or irritated eyes (your eyes may also be more sensitive to light)
- chest pain or pressure, pain spreading to your jaw or shoulder, irregular heartbeats

Drug Side Effects Summary

- pain or burning when you urinate, trouble emptying your bladder
- sore throat , cough, chest tightness, shortness of breath
- or eyelid swelling, crusting or drainage from your eyes, problems with vision. Common Botox side effects may include: painful or difficult urination
- headache, neck pain, back pain, pain in your arms or legs
- cold symptoms such as stuffy nose, sneezing, sore throat
- trouble swallowing
- fever, chills, body aches, flu symptoms
- increased sweating in areas other than the underarms
- or bruising, bleeding, pain, redness, or swelling where the injection was given.

2004. This particular disease, Incontinence, interacts with the drug fesoterodine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Fesoterodine may cause serious side effects. Call your doctor at once if you have: severe stomach pain or constipation
- a weak stream of urine, or trouble emptying your bladder
- little or no urination
- or headache , dizziness, drowsiness. Older adults may be more likely to have side effects such as dry mouth, constipation, upset stomach , increased urination, dizziness, or painful urination. Common side effects of fesoterodine may include: painful urination
- blurred vision
- weight gain
- dry mouth
- or stomach pain, diarrhea , constipation.

Drug Side Effects Summary

2005. This particular disease, Incontinence, interacts with the drug darifenacin and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Darifenacin may cause serious side effects. Stop using darifenacin and call your doctor at once if you have: severe stomach pain or constipation
- confusion, hallucinations
- little or no urinating
- pain or burning when you urinate
- or signs of dehydration --feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin . Common side effects of darifenacin may include: nausea , indigestion
- constipation
- headache , dizziness
- dry mouth
- or blurred vision.

2006. This particular disease, Incontinence, interacts with the drug flavoxate and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common side effects of flavoxate may include: nausea , vomiting , dry mouth
- headache , dizziness , drowsiness
- pounding heartbeats or fluttering in your chest
- confusion, nervousness
- or rash or itching.

Drug Side Effects Summary

2007. This particular disease, Incontinence, interacts with the drug onabotulinumtoxinA and has the

- hives , itching
- wheezing, difficult breathing
- feeling like you might pass out
- swelling of your face, lips, tongue, or throat. The botulinum toxin contained in Botox can spread to other body areas beyond where it was injected. This has caused serious life-threatening side effects in some people receiving botulinum toxin injections, even for cosmetic purposes. Call your doctor at once if you have any of these side effects (up to several hours or several weeks after an injection):
unusual or severe muscle weakness (especially in a body area that was not injected with the medication)
- loss of bladder control
- hoarse voice, trouble talking or swallowing
- drooping eyelids or eyebrows
- vision changes, eye pain , severely dry or irritated eyes (your eyes may also be more sensitive to light)
- chest pain or pressure, pain spreading to your jaw or shoulder, irregular heartbeats
- pain or burning when you urinate, trouble emptying your bladder
- sore throat , cough, chest tightness, shortness of breath
- or eyelid swelling, crusting or drainage from your eyes, problems with vision. Common side effects of onabotulinumtoxinA may include: painful or difficult urination
- headache , neck pain, back pain, pain in your arms or legs
- cold symptoms such as stuffy nose, sneezing, sore throat
- trouble swallowing
- fever, chills, body aches, flu symptoms
- increased sweating in areas other than the underarms

Drug Side Effects Summary

- or bruising, bleeding, pain, redness, or swelling where the injection was given.

2008. This particular disease, Insomnia, interacts with the drug Ambien and has the following side effects:

- Zolpidem may cause a severe allergic reaction. Stop taking Ambien and get emergency medical help if you have signs of an allergic reaction: hives

- difficulty breathing

- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: depression, anxiety , aggression, agitation , confusion, unusual thoughts, hallucinations, memory problems, changes in personality, risk-taking behavior, decreased inhibitions, no fear of danger, or thoughts of suicide or hurting yourself. Stop using this medicine and call your doctor at once if you have: chest pain, fast or irregular heartbeat , feeling short of breath

- trouble breathing or swallowing

- or feeling like you might pass out. The sedative effect of Ambien may be stronger in older adults.

Dizziness or severe drowsiness can cause falls, accidents, or severe injuries. Common Ambien side effects may include: daytime drowsiness, dizziness, weakness, feeling "drugged" or light-headed

- tired feeling, loss of coordination

- stuffy nose, dry mouth, nose or throat irritation

- nausea , constipation , diarrhea , upset stomach

- or headache , muscle pain.

2009. This particular disease, Insomnia, interacts with the drug zolpidem and has the following side effects:

- Zolpidem may cause a severe allergic reaction. Stop taking this medicine and get emergency medical help if you have signs of an allergic reaction to zolpidem: hives

- difficulty breathing

- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your

Drug Side Effects Summary

doctor, such as: depression, anxiety , aggression, agitation , confusion, unusual thoughts, hallucinations, memory problems, changes in personality, risk-taking behavior, decreased inhibitions, no fear of danger, or thoughts of suicide or hurting yourself. Stop using this medicine and call your doctor at once if you have: chest pain, fast or irregular heartbeat , feeling short of breath

- trouble breathing or swallowing

- or feeling like you might pass out. The sedative effect of zolpidem may be stronger in older adults.

Dizziness or severe drowsiness can cause falls, accidents, or severe injuries. Common zolpidem side effects may include: daytime drowsiness, dizziness, weakness, feeling "drugged" or light-headed

- tired feeling, loss of coordination

- stuffy nose, dry mouth, nose or throat irritation

- nausea , constipation , diarrhea , upset stomach

- or headache , muscle pain.

2010. This particular disease, Insomnia, interacts with the drug Belsomra and has the following side

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Some people using this medicine have engaged in activity while not fully awake and later had no memory of it. This may include walking, driving, eating, having sex, or making phone calls. If this happens to you, call your doctor right away. Call your doctor at once if you have: trouble moving or talking when you first wake up

- a weak feeling in your legs

- unusual thoughts or behavior

- anxiety , agitation , depression

- memory problems

Drug Side Effects Summary

- confusion, hallucinations
- or thoughts about hurting yourself. You may be more likely to have side effects if you are overweight . Common Belsomra side effects include: strange dreams
- or drowsiness during the day after taking this medicine.

2011. This particular disease, Insomnia, interacts with the drug suvorexant and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some people using suvorexant have engaged in activity while not fully awake and later had no memory of it. This may include walking, driving, eating, having sex, or making phone calls. If this happens to you, call your doctor right away. Suvorexant may cause serious side effects. Call your doctor at once if you have: trouble moving or talking when you first wake up
- a weak feeling in your legs
- unusual thoughts or behavior
- anxiety , agitation , depression
- memory problems
- confusion, hallucinations
- or thoughts about hurting yourself. You may be more likely to have side effects if you are overweight . Common side effects of suvorexant may include: strange dreams
- or drowsiness during the day after taking suvorexant.

2012. This particular disease, Insomnia, interacts with the drug Dayvigo and has the following side effects

- Dayvigo may cause serious side effects, including: see "Important information" above temporary inability to move or talk (sleep paralysis) for up to several minutes while you are going to sleep or

Drug Side Effects Summary

waking up. temporary weakness in your legs that can happen during the day or at night. complex sleep behaviors such as sleep-walking, sleep-driving, preparing and eating food, making phone calls, having sex or doing other activities while not fully awake that you may not remember the next morning. Call your healthcare provider right away if you experience a complex sleep behavior. worsening depression and suicidal thoughts have happened during treatment with this medication. Call your healthcare provider right away if you have any worsening depression or thoughts of suicide or dying. The most common side effect of Dayvigo is sleepiness. These are not all of the possible side effects of this medication. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Dayvigo side effects

2013. This particular disease, Insomnia, interacts with the drug temazepam and has the following side effects:

- Temazepam may cause a severe allergic reaction. hives
- difficulty breathing
- nausea , vomiting
- swelling of your face, lips, tongue, or throat. Temazepam can slow or stop your breathing, especially if you have recently used an opioid medication or alcohol. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Call your doctor at once if you have: confusion, agitation , hallucinations
- depressed mood
- or thoughts of suicide or hurting yourself. Some people using temazepam have engaged in activity such as driving, eating, walking, making phone calls, or having sex and later having no memory of the activity. Tell your doctor if this happens to you. Drowsiness or dizziness may last longer in older adults. Use caution to avoid falling or accidental injury. Common temazepam side effects may include: drowsiness, tiredness
- dizziness

Drug Side Effects Summary

- headache
- nausea
- or feeling nervous. After you stop using temazepam, get medical help right away if you have symptoms such as: unusual muscle movements, being more active or talkative, sudden and severe changes in mood or behavior, confusion, hallucinations, seizures, suicidal thoughts or actions. Some withdrawal symptoms may last up to 12 months or longer after stopping this medicine suddenly. Tell your doctor if you have ongoing anxiety , depression, problems with memory or thinking, trouble sleeping, ringing in your ears, a burning or prickly feeling, or a crawling sensation under your skin.

2014. This particular disease, Insomnia, interacts with the drug daridorexant and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tell your doctor right away if you have new or sudden changes in mood or behavior, including new or worse depression or have thoughts about suicide or hurting yourself. Some people using daridorexant have engaged in activity such as driving, eating, walking, making phone calls, or having sex and later having no memory of the activity. Tell your doctor if this happens to you. Call your doctor at once if you have an inability to move or talk for several minutes or if you have hallucinations while you are waking up or going to sleep. Common side effects of daridorexant may include: headache
- and drowsiness.

2015. This particular disease, Insomnia, interacts with the drug lemborexant and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some people using lemborexant have engaged in

Drug Side Effects Summary

activity while not fully awake and later had no memory of it. This may include walking, driving, eating, having sex, or making phone calls. If this happens to you, call your doctor right away. Lemborexant may cause serious side effects. Call your doctor at once if you have: trouble moving or talking when you first wake up

- a weak feeling in your legs
- unusual thoughts or behavior
- anxiety , agitation , depression
- memory problems
- confusion, hallucinations
- or worsening depression or thoughts about hurting yourself. Older adults have a higher risk of falls due to drowsiness caused by lemborexant. Common side effects of lemborexant may include: drowsiness during the day after taking lemborexant.

2016. This particular disease, Insomnia, interacts with the drug Quviviq and has the following side effects

- Quviviq may cause serious side effects, including: See Important information . Worsening depression and suicidal thoughts. Call your healthcare provider right away if you have any worsening depression or thoughts of suicide or dying. Temporary inability to move or talk (sleep paralysis) for up to several minutes, or hallucinations while you are going to sleep or waking up. Complex sleep behaviors such as sleep-walking, sleep-driving, preparing and eating food, making phone calls, having sex or doing other activities while not fully awake that you may not remember the next morning. Stop taking Quviviq and call your healthcare provider right away if you experience a complex sleep behavior. The most common side effects include headache and sleepiness. These are not all of the possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Quviviq side effects

Drug Side Effects Summary

2017. This particular disease, Insomnia, interacts with the drug Lunesta and has the following side effects:

- Eszopiclone may cause a severe allergic reaction. Stop taking Lunesta and get emergency medical help if you have signs of an allergic reaction to Lunesta: hives
- nausea , vomiting
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Some people using this medicine have engaged in activity while not fully awake and later had no memory of it. This may include walking, driving, or making phone calls. If this happens to you, stop taking Lunesta and call your doctor right away. Serious injury or death could occur if you walk or drive while you are not fully awake. Call your doctor at once if you have: anxiety , depression, aggression, agitation
- memory problems, unusual thoughts or behavior
- thoughts of hurting yourself
- or confusion, hallucinations (hearing or seeing things). Common Lunesta side effects may include: day-time drowsiness, dizziness , "hangover" feeling
- headache , anxiety
- dry mouth
- unusual or unpleasant taste in your mouth
- rash
- or cold or flu symptoms such as fever, body aches, sore throat , cough, runny or stuffy nose.

2018. This particular disease, Insomnia, interacts with the drug Restoril and has the following side effects:

- Temazepam may cause a severe allergic reaction. hives
- difficulty breathing
- nausea , vomiting
- swelling of your face, lips, tongue, or throat. Restoril can slow or stop your breathing, especially if

Drug Side Effects Summary

you have recently used an opioid medication or alcohol. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Call your doctor at once if you have: confusion, agitation , hallucinations

- depressed mood

- or thoughts of suicide or hurting yourself. Some people using Restoril have engaged in activity such as driving, eating, walking, making phone calls, or having sex and later having no memory of the activity. Tell your doctor if this happens to you. Drowsiness or dizziness may last longer in older adults. Use caution to avoid falling or accidental injury. Common Restoril side effects may include: drowsiness, tiredness

- dizziness

- headache

- nausea

- or feeling nervous. After you stop using Restoril, get medical help right away if you have symptoms such as: unusual muscle movements, being more active or talkative, sudden and severe changes in mood or behavior, confusion, hallucinations, seizures, suicidal thoughts or actions. Some withdrawal symptoms may last up to 12 months or longer after stopping this medicine suddenly. Tell your doctor if you have ongoing anxiety , depression, problems with memory or thinking, trouble sleeping, ringing in your ears, a burning or prickly feeling, or a crawling sensation under your skin.

2019. This particular disease, Insomnia, interacts with the drug estazolam and has the following side effects:

- Estazolam may cause a severe allergic reaction. hives

- difficult breathing

- nausea , vomiting

- swelling of your face, lips, tongue, or throat. Estazolam can slow or stop your breathing, especially if you have recently used an opioid medication or alcohol. A person caring for you should seek

Drug Side Effects Summary

emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Estazolam may cause serious side effects. Call your doctor at once if you have: unusual thoughts or behavior

- depressed mood, thoughts about suicide or hurting yourself
- confusion, hallucinations, aggression, agitation
- or worsening sleep problems. Some people using estazolam have engaged in activity such as driving, eating, walking, making phone calls, or having sex and later having no memory of the activity. Tell your doctor if this happens to you. Drowsiness or dizziness may last longer in older adults. Use caution to avoid falling or accidental injury. Common side effects of estazolam may include: drowsiness, dizziness

- headache
- feeling tired
- dry mouth
- or upset stomach . After you stop using estazolam, get medical help right away if you have symptoms such as: unusual muscle movements, being more active or talkative, sudden and severe changes in mood or behavior, confusion, hallucinations, seizures, suicidal thoughts or actions. Some withdrawal symptoms may last up to 12 months or longer after stopping this medicine suddenly. Tell your doctor if you have ongoing anxiety , depression, problems with memory or thinking, trouble sleeping, ringing in your ears, a burning or prickly feeling, or a crawling sensation under your skin.

2020. This particular disease, Insomnia, interacts with the drug flurazepam and has the following si

- hives
- nausea and vomiting
- difficult breathing
- swelling of your face, lips, tongue, or throat. Flurazepam can slow or stop your breathing,

Drug Side Effects Summary

especially if you have recently used an opioid medication or alcohol. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Flurazepam may cause serious side effects. Call your doctor at once if you have: confusion, agitation , paranoia, hallucinations

- thinking problems, unusual thoughts or behavior
- severe daytime drowsiness
- or new or worsening depression. Some people using flurazepam have engaged in activity such as driving, eating, walking, making phone calls, or having sex and later having no memory of the activity. Tell your doctor if this happens to you. Drowsiness or dizziness may last longer in older adults. Use caution to avoid falling or accidental injury. Common side effects of flurazepam may include: drowsiness, dizziness, feeling light-headed
- loss of coordination
- or falling or staggering. After you stop using flurazepam, get medical help right away if you have symptoms such as: unusual muscle movements, being more active or talkative, sudden and severe changes in mood or behavior, confusion, hallucinations, seizures, suicidal thoughts or actions. Some withdrawal symptoms may last up to 12 months or longer after stopping this medicine suddenly. Tell your doctor if you have ongoing anxiety , depression, problems with memory or thinking, trouble sleeping, ringing in your ears, a burning or prickly feeling, or a crawling sensation under your skin.

2021. This particular disease, Insomnia, interacts with the drug Halcion and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Halcion can slow or stop your breathing, especially if you have recently used an opioid medication or alcohol. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you

Drug Side Effects Summary

are hard to wake up. Halcion may cause serious side effects. Call your doctor at once if you have:
daytime anxiety

- unusual changes in mood or behavior
- confusion, memory loss , agitation , hallucinations
- or depression, suicidal thoughts. Some people using Halcion have engaged in activity such as driving, eating, walking, making phone calls, or having sex and later having no memory of the activity. Tell your doctor if this happens to you. Drowsiness or dizziness may last longer in older adults. Use caution to avoid falling or accidental injury. Common side effects of Halcion may include:
drowsiness

- loss of coordination
- dizziness
- or feeling light-headed.

2022. This particular disease, Insomnia, interacts with the drug Ambien CR and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Chest pain confusion about identity, place, and time discouragement false or unusual sense of well-being fast, irregular, pounding, or racing heartbeat or pulse feeling sad or empty fever general feeling of discomfort or illness irritability lack of appetite lightheadedness loss of interest or pleasure shakiness and unsteady walk trouble concentrating trouble sleeping unsteadiness, trembling, or other problems with muscle control or coordination unusual tiredness or weakness Rare Anxiety attack, assault, or force black, tarry stools bladder pain bloody or cloudy urine burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings change in walking and balance chest tightness chills cold sweats cough crying decreased awareness or responsiveness deep or fast breathing with

Drug Side Effects Summary

dizziness delusions of persecution, mistrust, suspiciousness, or combativeness difficult, burning, or painful urination difficulty with swallowing dizziness dizziness, faintness, or lightheadedness when getting up from lying or sitting position dry mouth fainting false beliefs that cannot be changed by facts feeling of unreality fever frequent urge to urinate hives, itching, or skin rash irregular heartbeat irritability lack of feeling or emotion light-colored stools lower back or side pain nausea numbness of the feet, hands, and around the mouth pale skin poor insight and judgment problems with memory or speech puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue quick to react or overreact emotionally rapidly changing moods restlessness seeing, hearing, or feeling things that are not there sense of detachment from self or body severe sleepiness sore throat sores, ulcers, or white spots on the lips or in the mouth stomach pain swollen glands thoughts of killing oneself or changes in behavior trouble recognizing objects trouble thinking and planning trouble walking trouble breathing uncaring unease or dissatisfaction with life unusual bleeding or bruising vomiting yellow eyes or skin

Symptoms of overdose Change or loss of consciousness difficult or trouble breathing irregular, fast or slow, or shallow breathing pale or blue lips, fingernails, or skin sleepiness or unusual drowsiness

Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them:

More common Drowsiness headache muscle aches sleepiness or unusual drowsiness stuffy or runny nose

Less common Back pain being forgetful belching body aches or pain congestion diarrhea difficulty having a bowel movement difficulty with moving double vision feeling of constant movement of self or surroundings heartburn hoarseness indigestion longer or heavier menstrual periods loss of memory muscle pain or stiffness pain or tenderness around the eyes and cheekbones seeing double sensation of spinning stomach discomfort tender, swollen glands in the neck unusual drowsiness, dullness, tiredness, weakness, or

Drug Side Effects Summary

feeling of sluggishness voice changes Rare Bad, unusual, or unpleasant (after) taste blemishes on the skin blindness blurred vision breast pain change in near or distance vision change in taste changes in patterns and rhythms of speech decreased interest in sexual intercourse difficulty in focusing the eyes eye pain feeling of warmth frequent urge to defecate inability to have or keep an erection increased appetite increased sensitivity of the skin to sunlight increased sweating itching of the vagina or genital area lack or loss of strength loss in sexual ability, desire, drive, or performance pain during sexual intercourse paleness of the skin pimples redness of the face, neck, arms, and occasionally, upper chest redness or other discoloration of the skin severe sunburn slurred speech stomach upset straining while passing stool tearing thick, white vaginal discharge with no odor or with a mild odor weight loss Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Ambien CR side effects

2023. This particular disease, Insomnia, interacts with the drug eszopiclone and has the following side effects

- Eszopiclone may cause a severe allergic reaction. Stop taking eszopiclone and get emergency medical help if you have signs of an allergic reaction: hives
- nausea , vomiting
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Some people using this medicine have engaged in activity while not fully awake and later had no memory of it. This may include walking, driving, or making phone calls. If this happens to you, stop taking eszopiclone and call your doctor right away. Serious injury or death could occur if you walk or drive while you are not fully awake. Eszopiclone may cause serious side effects. Call your doctor at once if you have: anxiety , depression, aggression, agitation

Drug Side Effects Summary

- memory problems, unusual thoughts or behavior
- thoughts of hurting yourself
- or confusion, hallucinations (hearing or seeing things). Common side effects of eszopiclone may include: day-time drowsiness, dizziness , "hangover" feeling
- headache , anxiety
- dry mouth
- unusual or unpleasant taste in your mouth
- rash
- or cold or flu symptoms such as fever, body aches, sore throat , cough, runny or stuffy nose.

2024. This particular disease, Insomnia, interacts with the drug Rozerem and has the following side effects

- Rozerem may cause a severe allergic reaction. hives
- difficult breathing
- nausea and vomiting
- swelling of your face, lips, tongue, or throat. Some people using this medicine have engaged in activity while not fully awake and later had no memory of it. This may include walking, driving, eating, having sex, or making phone calls. If this happens to you, call your doctor right away. Call your doctor at once if you have: unusual thoughts or behavior, hallucinations, worsening depression, thoughts about hurting yourself
- a missed menstrual period
- nipple discharge
- or loss of interest in sex. Common Rozerem side effects may include: drowsiness, tired feeling
- dizziness
- nausea
- or worsening sleep problems.

Drug Side Effects Summary

2025. This particular disease, Insomnia, interacts with the drug Silenor and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Some people using this medicine have engaged in activity while not fully awake and later had no memory of it. This may include walking, driving, or making phone calls. If this happens to you, call your doctor right away. Call your doctor at once if you have: unusual thoughts or behavior
- confusion, hallucinations
- depressed mood, thoughts about hurting yourself
- uncontrolled muscle movements
- a seizure
- a light-headed feeling, like you might pass out
- blisters or ulcers in your mouth, red or swollen gums, trouble swallowing
- eye pain or redness
- chest pain
- low red blood cells (anemia) - pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet
- or signs of infection - flu-like symptoms, swelling, skin redness, diarrhea, skin sores or white patches, trouble breathing, pain or burning when you urinate. Common Silenor side effects may include: drowsiness, tiredness
- nausea
- or cold symptoms such as stuffy nose, sneezing, sore throat.

2026. This particular disease, Insomnia, interacts with the drug Sonata and has the following side effects:

- Zaleplon may cause a severe allergic reaction. hives

Drug Side Effects Summary

- difficult breathing
- nausea and vomiting
- swelling of your face, lips, tongue, or throat. Some people using this medicine have engaged in activity while not fully awake and later had no memory of it. This may include walking, driving, or making phone calls. If this happens to you, stop taking Sonata and call your doctor right away. Serious injury or death could occur if you walk or drive while you are not fully awake. Call your doctor at once if you have: anxiety , depression, aggression, agitation
- memory problems, unusual thoughts or behavior
- thoughts of hurting yourself
- or confusion, hallucinations (hearing or seeing things). Common Sonata side effects may include: day-time drowsiness, dizziness , "hangover" feeling
- feeling light-headed
- problems with coordination
- or numbness , tingling, weakness, or prickly feeling on your skin.

2027. This particular disease, Insomnia, interacts with the drug Advil PM and has the following side

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: any skin rash, no matter how mild
- a light-headed feeling, like you might pass out
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- kidney problems--little or no urinating, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath

Drug Side Effects Summary

- or liver problems-- nausea , upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Advil PM may include: drowsiness
- day-time drowsiness, dizziness , "hangover" feeling
- upset stomach , heartburn , diarrhea , constipation
- dry mouth, nose, or throat
- blurred vision
- mild itching or rash
- or ringing in your ears.

2028. This particular disease, Insomnia, interacts with the drug triazolam and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Triazolam can slow or stop your breathing, especially if you have recently used an opioid medication or alcohol. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Triazolam may cause serious side effects. Call your doctor at once if you have: daytime anxiety
- unusual changes in mood or behavior
- confusion, memory loss , agitation , hallucinations
- or depression, suicidal thoughts. Some people using triazolam have engaged in activity such as driving, eating, walking, making phone calls, or having sex and later having no memory of the activity. Tell your doctor if this happens to you. Drowsiness or dizziness may last longer in older adults. Use caution to avoid falling or accidental injury. Common side effects of triazolam may include: drowsiness

Drug Side Effects Summary

- loss of coordination
- dizziness
- or feeling light-headed.

2029. This particular disease, Insomnia, interacts with the drug quazepam and has the following side effects:

- Quazepam may cause a severe allergic reaction. hives
- difficult breathing
- nausea , vomiting
- swelling of your face, lips, tongue, or throat. Quazepam can slow or stop your breathing, especially if you have recently used an opioid medication or alcohol. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Quazepam may cause serious side effects. Call your doctor at once if you have: confusion, agitation , hallucinations
- unusual thoughts or behavior
- suicidal thoughts
- or worsening depression. Some people using quazepam have engaged in activity such as driving, eating, walking, making phone calls, or having sex and later having no memory of the activity. Tell your doctor if this happens to you. Drowsiness or dizziness may last longer in older adults. Use caution to avoid falling or accidental injury. Common side effects of quazepam may include: drowsiness, dizziness
- headache
- feeling tired
- dry mouth
- or upset stomach . After you stop using quazepam, get medical help right away if you have symptoms such as: unusual muscle movements, being more active or talkative, sudden and severe

Drug Side Effects Summary

changes in mood or behavior, confusion, hallucinations, seizures, suicidal thoughts or actions. Some withdrawal symptoms may last up to 12 months or longer after stopping this medicine suddenly. Tell your doctor if you have ongoing anxiety , depression, problems with memory or thinking, trouble sleeping, ringing in your ears, a burning or prickly feeling, or a crawling sensation under your skin.

2030. This particular disease, Insomnia, interacts with the drug Edluar and has the following side effects

- Zolpidem may cause a severe allergic reaction. Stop taking Edluar and get emergency medical help if you have signs of an allergic reaction to zolpidem: hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: depression, anxiety , aggression, agitation , confusion, unusual thoughts, hallucinations, memory problems, changes in personality, risk-taking behavior, decreased inhibitions, no fear of danger, or thoughts of suicide or hurting yourself. Stop using Edluar and call your doctor at once if you have: chest pain, fast or irregular heartbeat , feeling short of breath
- trouble breathing or swallowing
- or feeling like you might pass out. The sedative effect of zolpidem may be stronger in older adults. Dizziness or severe drowsiness can cause falls, accidents, or severe injuries. Common Edluar side effects may include: daytime drowsiness, dizziness, weakness, feeling "drugged" or light-headed
- tired feeling, loss of coordination
- stuffy nose, dry mouth, nose or throat irritation
- nausea , constipation , diarrhea , upset stomach
- or headache , muscle pain.

2031. This particular disease, Insomnia, interacts with the drug Tylenol PM and has the following side effects

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: severe drowsiness
- painful or difficult urination
- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Tylenol PM may include: drowsiness
- constipation
- decreased urination
- dry eyes, blurred vision
- or dry mouth, nose, or throat.

2032. This particular disease, Insomnia, interacts with the drug Unisom SleepTabs and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Unisom may cause serious side effects. Call your doctor at once if you have: confusion, hallucinations
- severe dizziness or drowsiness
- or little or no urinating. Common side effects of Unisom may include: blurred vision
- dry mouth, nose, or throat

Drug Side Effects Summary

- constipation
- or mild dizziness or drowsiness. Side effects such as dry mouth, constipation, and confusion may be more likely in older adults.

2033. This particular disease, Insomnia, interacts with the drug zaleplon and has the following side

- Zaleplon may cause a severe allergic reaction. hives
- difficult breathing
- nausea and vomiting
- swelling of your face, lips, tongue, or throat. Some people using this medicine have engaged in activity while not fully awake and later had no memory of it. This may include walking, driving, or making phone calls. If this happens to you, stop taking zaleplon and call your doctor right away. Serious injury or death could occur if you walk or drive while you are not fully awake. Zaleplon may cause serious side effects. Call your doctor at once if you have: anxiety , depression, aggression, agitation
- memory problems, unusual thoughts or behavior
- thoughts of hurting yourself
- or confusion, hallucinations (hearing or seeing things). Common side effects of zaleplon may include: day-time drowsiness, dizziness , "hangover" feeling
- feeling light-headed
- problems with coordination
- or numbness , tingling, or prickly feeling on your skin.

2034. This particular disease, Insomnia, interacts with the drug Compoz Nighttime Sleep Aid and ha

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Compoz Nighttime Sleep Aid may cause serious side effects. Stop using Compoz Nighttime Sleep Aid and call your doctor at once if you have: severe drowsiness
- or painful or difficult urination. Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of Compoz Nighttime Sleep Aid may include: drowsiness
- dry eyes, blurred vision
- dry mouth, nose, or throat
- decreased urination
- constipation
- feeling restless or excited (especially in children)
- or day-time drowsiness or "hangover" feeling after night-time use.

2035. This particular disease, Insomnia, interacts with the drug Doral and has the following side effects

- Doral may cause a severe allergic reaction. hives
- difficult breathing
- nausea , vomiting
- swelling of your face, lips, tongue, or throat. Doral can slow or stop your breathing, especially if you have recently used an opioid medication or alcohol. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Doral may cause serious side effects. Call your doctor at once if you have: confusion, agitation , hallucinations
- unusual thoughts or behavior
- suicidal thoughts
- or worsening depression. Some people using Doral have engaged in activity such as driving,

Drug Side Effects Summary

eating, walking, making phone calls, or having sex and later having no memory of the activity. Tell your doctor if this happens to you. Drowsiness or dizziness may last longer in older adults. Use caution to avoid falling or accidental injury. Common side effects of Doral may include: drowsiness, dizziness

- headache

- feeling tired

- dry mouth

- or upset stomach . After you stop using Doral, get medical help right away if you have symptoms such as: unusual muscle movements, being more active or talkative, sudden and severe changes in mood or behavior, confusion, hallucinations, seizures, suicidal thoughts or actions. Some withdrawal symptoms may last up to 12 months or longer after stopping this medicine suddenly. Tell your doctor if you have ongoing anxiety , depression, problems with memory or thinking, trouble sleeping, ringing in your ears, a burning or prickly feeling, or a crawling sensation under your skin.

2036. This particular disease, Insomnia, interacts with the drug ramelteon and has the following side effects

- Ramelteon may cause a severe allergic reaction. hives

- difficult breathing

- nausea and vomiting

- swelling of your face, lips, tongue, or throat. Some people using this medicine have engaged in activity while not fully awake and later had no memory of it. This may include walking, driving, eating, having sex, or making phone calls. If this happens to you, call your doctor right away.

Ramelteon may cause serious side effects. Call your doctor at once if you have: unusual thoughts or behavior, hallucinations, worsening depression, thoughts about hurting yourself

- a missed menstrual period

- nipple discharge

Drug Side Effects Summary

- or loss of interest in sex. Common side effects of ramelteon may include: drowsiness, tired feeling
- dizziness
- nausea
- or worsening sleep problems.

2037. This particular disease, Insomnia, interacts with the drug chloral hydrate and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad belly pain. Feeling confused. Change in how you act. Fever, chills, or sore throat . Very bad dizziness or passing out. A heartbeat that does not feel normal. Trouble breathing, slow breathing, or shallow breathing. Some people have done certain tasks or actions while they were not fully awake like driving, making and eating food, and having sex. Most of the time, people do not remember doing these things. Tell your doctor if this happens to you. Chloral hydrate side effects

2038. This particular disease, Insomnia, interacts with the drug dimenhydrinate and has the following

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Dimenhydrinate may cause serious side effects. Stop using dimenhydrinate and call your doctor at once if you have: little or no urination
- confusion, mood changes
- tremors, restlessness
- a seizure (convulsions)
- or fast or irregular heartbeats. Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of dimenhydrinate may include: drowsiness
- dry mouth, nose, or throat
- constipation
- blurred vision
- or feeling restless or excited (especially in children).

2039. This particular disease, Insomnia, interacts with the drug diphenhydramine / ibuprofen and ha

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: any skin rash, no matter how mild
- a light-headed feeling, like you might pass out
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- kidney problems--little or no urinating, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- or liver problems-- nausea , upper stomach pain, itching, tired feeling, flu-like symptoms, loss of

Drug Side Effects Summary

appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of diphenhydramine and ibuprofen may include: drowsiness

- day-time drowsiness, dizziness , "hangover" feeling
- upset stomach , heartburn , diarrhea , constipation
- dry mouth, nose, or throat
- blurred vision
- mild itching or rash
- or ringing in your ears.

2040. This particular disease, Insomnia, interacts with the drug Nytol and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad dizziness or passing out. Change in balance. Feeling less alert. Nytol side effects

2041. This particular disease, Insomnia, interacts with the drug pentobarbital and has the following

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Pentobarbital may cause serious side effects. Call your doctor at once if you have: confusion, agitation , hallucinations
- weak or shallow breathing
- slow heart rate , weak pulse
- or a light-headed feeling, like you might pass out. Side effects such as confusion, depression , or excitement may be more likely in older adults and those who are ill or debilitated. Common side effects of pentobarbital may include: drowsiness, dizziness
- loss of balance or coordination
- nausea , vomiting , constipation
- overactive reflexes
- sleep problems (insomnia), nightmares
- or feeling restless or excited (especially in children or older adults).

2042. This particular disease, Insomnia, interacts with the drug Simply Sleep and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Simply Sleep may cause serious side effects. Stop using Simply Sleep and call your doctor at once if you have: severe drowsiness
- or painful or difficult urination. Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of Simply Sleep may include: drowsiness
- dry eyes, blurred vision
- dry mouth, nose, or throat
- decreased urination

Drug Side Effects Summary

- constipation
- feeling restless or excited (especially in children)
- or day-time drowsiness or "hangover" feeling after night-time use.

2043. This particular disease, Insomnia, interacts with the drug Sominex and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Sominex may cause serious side effects. Stop using Sominex and call your doctor at once if you have: severe drowsiness
- or painful or difficult urination. Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of Sominex may include: drowsiness
- dry eyes, blurred vision
- dry mouth, nose, or throat
- decreased urination
- constipation
- feeling restless or excited (especially in children)
- or day-time drowsiness or "hangover" feeling after night-time use.

2044. This particular disease, Insomnia, interacts with the drug Unisom SleepGels and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Unisom Sleep Gels may cause serious side effects. Stop using Unisom Sleep Gels and call your doctor at once if you have: severe drowsiness
- or painful or difficult urination. Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of Unisom Sleep Gels may include: drowsiness

Drug Side Effects Summary

- dry eyes, blurred vision
- dry mouth, nose, or throat
- decreased urination
- constipation
- feeling restless or excited (especially in children)
- or day-time drowsiness or "hangover" feeling after night-time use.

2045. This particular disease, Insomnia, interacts with the drug Unisom SleepMelts and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad dizziness or passing out. Change in balance. Feeling less alert. Unisom SleepMelts side effects

2046. This particular disease, Insomnia, interacts with the drug Zolpimist and has the following side effects:

- Zolpidem may cause a severe allergic reaction. Stop using Zolpimist and get emergency medical help if you have signs of an allergic reaction to zolpidem: hives

Drug Side Effects Summary

- difficulty breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: depression, anxiety , aggression, agitation , confusion, unusual thoughts, hallucinations, memory problems, changes in personality, risk-taking behavior, decreased inhibitions, no fear of danger, or thoughts of suicide or hurting yourself. Stop using Zolpimist and call your doctor at once if you have: chest pain, fast or irregular heartbeat , feeling short of breath
- trouble breathing or swallowing
- or feeling like you might pass out. The sedative effect of zolpidem may be stronger in older adults. Dizziness or severe drowsiness can cause falls, accidents, or severe injuries. Common Zolpimist side effects may include: daytime drowsiness, dizziness, weakness, feeling "drugged" or light-headed
- tired feeling, loss of coordination
- stuffy nose, dry mouth, nose or throat irritation
- nausea , constipation , diarrhea , upset stomach
- or headache , muscle pain.

2047. This particular disease, Insomnia, interacts with the drug ZzzQuil and has the following side effects:

- No side effects listed

2048. This particular disease, Insomnia, interacts with the drug acetaminophen / diphenhydramine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin

Drug Side Effects Summary

redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: severe drowsiness

- painful or difficult urination
- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of acetaminophen and diphenhydramine may include: drowsiness
- constipation
- decreased urination
- dry eyes, blurred vision
- or dry mouth, nose, or throat.

2049. This particular disease, Insomnia, interacts with the drug amobarbital and has the following s

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness

Drug Side Effects Summary

- or swelling of the mouth, face, lips, tongue, or throat. Signs of depression , thoughts of suicide, nervousness , emotional ups and downs, thinking that is not normal, anxiety , or lack of interest in life. Trouble breathing, slow breathing, or shallow breathing. Very bad irritation where the shot was given. This medicine may cause tissue damage if the drug leaks from the vein. Tell your nurse if you have any redness, burning, pain, swelling, blisters, skin sores, or leaking of fluid where the drug is going into your body. Amobarbital side effects

2050. This particular disease, Insomnia, interacts with the drug Ibuprofen PM and has the following

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: any skin rash, no matter how mild
- a light-headed feeling, like you might pass out
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- kidney problems--little or no urinating, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- or liver problems-- nausea , upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Ibuprofen PM may include: drowsiness
- day-time drowsiness, dizziness , "hangover" feeling
- upset stomach , heartburn , diarrhea , constipation
- dry mouth, nose, or throat
- blurred vision
- mild itching or rash

Drug Side Effects Summary

- or ringing in your ears.

2051. This particular disease, Insomnia, interacts with the drug Nighttime Sleepaid and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Not sweating during activities or in warm temperatures. Not able to pass urine. Trouble passing urine. Feeling very tired or weak. Change in eyesight. A fast heartbeat . A heartbeat that does not feel normal. Nighttime Sleepaid side effects

2052. This particular disease, Insomnia, interacts with the drug Nytol QuickCaps and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Nytol QuickCaps may cause serious side effects. Stop using Nytol QuickCaps and call your doctor at once if you have: severe drowsiness
- or painful or difficult urination. Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of Nytol QuickCaps may include: drowsiness

Drug Side Effects Summary

- dry eyes, blurred vision
- dry mouth, nose, or throat
- decreased urination
- constipation
- feeling restless or excited (especially in children)
- or day-time drowsiness or "hangover" feeling after night-time use.

2053. This particular disease, Insomnia, interacts with the drug phenobarbital and has the following

- Phenobarbital may cause a severe allergic reaction. Stop taking phenobarbital and get emergency medical help if you have signs of an allergic reaction : hives

- difficult breathing

- swelling of your face, eyes, lips, tongue, or throat. Phenobarbital may cause serious side effects.

Call your doctor at once if you have: weak or shallow breathing

- unusual pain anywhere in your body (especially in the neck, shoulder, or arms)

- a red blood cell disorder--pale skin, muscle weakness, diarrhea , weight loss , rapid heart rate, tongue swelling, numbness or tingling in your hands or feet, feeling short of breath

- or severe skin reaction--fever, sore throat , swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Side effects such as confusion, depression , or excitement may be more likely in older adults and those who are ill or debilitated. Common side effects of phenobarbital may include: drowsiness, lack of energy

- dizziness or spinning sensation

- depressed mood

- feeling restless or excited (especially in children or older adults)

- drunk feeling

Drug Side Effects Summary

- or "hangover" effect (drowsiness the day after taking phenobarbital).

2054. This particular disease, Insomnia, interacts with the drug Sleep Aid and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

- itching

- red, swollen, blistered, or peeling skin with or without fever

- wheezing

- tightness in the chest or throat

- trouble breathing, swallowing, or talking

- unusual hoarseness

- or swelling of the mouth, face, lips, tongue, or throat. Not sweating during activities or in warm temperatures. Not able to pass urine. Trouble passing urine. Feeling very tired or weak. Change in eyesight. A fast heartbeat . A heartbeat that does not feel normal. Sleep Aid side effects

2055. This particular disease, Insomnia, interacts with the drug Sleepinal and has the following side effects:

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Sleepinal may cause serious side effects. Stop using Sleepinal and call your doctor at once if you have: severe drowsiness

- or painful or difficult urination. Side effects such as dry mouth, constipation , and confusion may be more likely in older adults. Common side effects of Sleepinal may include: drowsiness

Drug Side Effects Summary

- dry eyes, blurred vision
- dry mouth, nose, or throat
- decreased urination
- constipation
- feeling restless or excited (especially in children)
- or day-time drowsiness or "hangover" feeling after night-time use.

2056. This particular disease, Insomnia, interacts with the drug Z-Sleep and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Z-Sleep may cause serious side effects. Stop using Z-Sleep and call your doctor at once if you have: severe drowsiness
- or painful or difficult urination. Side effects such as dry mouth, constipation, and confusion may be more likely in older adults. Common side effects of Z-Sleep may include: drowsiness
- dry eyes, blurred vision
- dry mouth, nose, or throat
- decreased urination
- constipation
- feeling restless or excited (especially in children)
- or day-time drowsiness or "hangover" feeling after night-time use.

2057. This particular disease, Insomnia, interacts with the drug Aleve PM and has the following side effects:

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. Stop using this medicine and call your doctor at once if you have: any skin rash, no matter how mild

Drug Side Effects Summary

- a light-headed feeling, like you might pass out
- new symptoms, such redness or swelling with pain
- or signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Common Aleve PM side effects may include: drowsiness
- day-time drowsiness, dizziness , "hangover" feeling
- upset stomach , heartburn
- constipation
- blurred vision
- or dry mouth, nose, or throat.

2058. This particular disease, Insomnia, interacts with the drug diphenhydramine / magnesium salicylate

- hives
- wheezing, difficult breathing
- feeling light-headed
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe drowsiness
- painful or difficult urination
- ringing in your ears, hearing loss
- confusion
- or signs of stomach bleeding--feeling light-headed, ongoing stomach pain, bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Common side effects of diphenhydramine and magnesium salicylate may include: drowsiness
- dry eyes, blurred vision
- dry mouth, nose, or throat
- decreased urination

Drug Side Effects Summary

- heartburn, nausea , vomiting, gas
- constipation
- feeling restless or excited (especially in children)
- or day-time drowsiness or "hangover" feeling after night-time use.

2059. This particular disease, Insomnia, interacts with the drug diphenhydramine / naproxen and has the following side effects:

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: any skin rash, no matter how mild
- a light-headed feeling, like you might pass out
- new symptoms, such redness or swelling with pain
- or signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Common side effects of diphenhydramine and naproxen may include: drowsiness
- day-time drowsiness, dizziness , "hangover" feeling
- upset stomach , heartburn
- constipation
- blurred vision
- or dry mouth, nose, or throat.

2060. This particular disease, Insomnia, interacts with the drug Doans PM and has the following side effects:

- hives
- wheezing, difficult breathing
- feeling light-headed
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop

Drug Side Effects Summary

using this medicine and call your doctor at once if you have: severe drowsiness

- painful or difficult urination
- ringing in your ears, hearing loss
- confusion
- or signs of stomach bleeding--feeling light-headed, ongoing stomach pain, bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Common side effects of Doans PM may include: drowsiness
- dry eyes, blurred vision
- dry mouth, nose, or throat
- decreased urination
- heartburn, nausea , vomiting, gas
- constipation
- feeling restless or excited (especially in children)
- or day-time drowsiness or "hangover" feeling after night-time use.

2061. This particular disease, Insomnia, interacts with the drug Headache Relief PM and has the fol

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: severe drowsiness

Drug Side Effects Summary

- painful or difficult urination
- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Headache Relief PM may include: drowsiness
- constipation
- decreased urination
- dry eyes, blurred vision
- or dry mouth, nose, or throat.

2062. This particular disease, Insomnia, interacts with the drug Legatrin PM and has the following s

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: severe drowsiness
- painful or difficult urination
- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Legatrin PM may include: drowsiness
- constipation
- decreased urination

Drug Side Effects Summary

- dry eyes, blurred vision
- or dry mouth, nose, or throat.

2063. This particular disease, Insomnia, interacts with the drug Midol PM and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: severe drowsiness
- painful or difficult urination
- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Midol PM may include: drowsiness
- constipation
- decreased urination
- dry eyes, blurred vision
- or dry mouth, nose, or throat.

2064. This particular disease, Insomnia, interacts with the drug Motrin PM and has the following side effects:

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. This medicine may cause serious side

Drug Side Effects Summary

effects. Stop using this medicine and call your doctor at once if you have: any skin rash, no matter how mild

- a light-headed feeling, like you might pass out
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- kidney problems--little or no urinating, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- or liver problems-- nausea , upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Motrin PM may include: drowsiness
- day-time drowsiness, dizziness , "hangover" feeling
- upset stomach , heartburn , diarrhea , constipation
- dry mouth, nose, or throat
- blurred vision
- mild itching or rash
- or ringing in your ears.

2065. This particular disease, Insomnia, interacts with the drug Percogesic Extra Strength and has

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Percogesic Extra Strength and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. Stop using the

Drug Side Effects Summary

medicine and call your doctor at once if you have: fast, pounding, or uneven heartbeats

- severe dizziness or drowsiness, slow or shallow breathing
- tremor, restless muscle movements
- little or no urinating
- flu symptoms, easy bruising, unusual bleeding, pale skin, feeling light-headed
- or nausea, pain in your upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common Percogesic Extra Strength side effects may include: constipation
- dry eyes, blurred vision, dry mouth or nose
- mild dizziness or drowsiness, trouble concentrating
- feeling restless or excited (especially in children)
- or mild skin rash .

2066. This particular disease, Insomnia, interacts with the drug Percogesic Original Strength and h

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. The medicine may cause serious side effects. Stop using the medicine and call your doctor at once if you have: severe drowsiness
- painful or difficult urination
- or liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine,

Drug Side Effects Summary

clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Percogesic

Original Strength may include: drowsiness

- constipation
- decreased urination
- dry eyes, blurred vision
- or dry mouth, nose, or throat.

2067. This particular disease, Insomnia, interacts with the drug zopiclone and has the following side effects:

- Serious side effects of zopiclone include: Allergic reactions. The signs of an allergic reaction may include: a rash swelling or breathing problems swelling of your lips, face, throat or tongue Stop taking this medication and see a doctor or go to the emergency room straight away if you develop symptoms of an allergic reaction. Tell your doctor as soon as possible if you have any of the following side effects, which are rare or of unknown frequency: Poor memory (amnesia) since starting this medication. This problem is less likely to occur if you have 7-8 hours of uninterrupted sleep after taking this medication. Seeing or hearing things that are not real (hallucinations) Falling, especially in the elderly Thinking things that are not true (delusions) Feeling low or sad (depressed mood) Tell your doctor or pharmacist if any of the following side effects get serious or lasts longer than a few days: Including the following common side effects of zopiclone: A mild bitter or metallic taste in your mouth or a dry mouth Feeling drowsy or sleepy Dry mouth Including the following uncommon side effects of zopiclone: Feeling sick (nausea) or being sick (vomiting) Feeling dizzy or sleepy Headache Nightmares Feeling physically or mentally tired Agitation Including the following rare side effects of zopiclone: Feeling confused Itchy, lumpy rash (urticaria) Feeling irritable or aggressive Reduced sex drive Difficulty breathing or being short of breath Including the following side effects of unknown frequency of zopiclone: Feeling restless or angry Feeling light headed or having problems with your coordination Double vision Moving unsteadily or staggering Muscular

Drug Side Effects Summary

weakness Indigestion Becoming dependent on this medication Slower breathing (respiratory depression) Unusual skin sensations such as numbness, tingling, pricking, burning or creeping on the skin (paraesthesia) Mental problems such as poor memory Difficulty paying attention Disrupted normal speech Zopiclone may cause sleepwalking or other unusual behavior (such as driving, eating, making a phone call, or having sex etc.) while not being fully awake. These are not all of the possible side effects of zopiclone. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Interactions Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Especially tell your doctor if you are taking any of the medicines listed below.

Zopiclone may increase the effect of the following medicines:

- Medicines for mental problems (antipsychotics)
- Medicines for depression
- Medicines for epilepsy (anticonvulsants)
- Medicines used in surgery (anaesthetics)
- Medicines to calm or reduce anxiety or for sleep problems (hypnotics)
- Medicines for hay fever, rashes or other allergies that can make you sleepy (sedative antihistamines) such as chlorphenamine or promethazine
- Some medicines for moderate to severe pain (narcotic analgesics) such as codeine, methadone, morphine, oxycodone, pethidine or tramadol

The following medicines can increase the chances of you getting side effects when taken with zopiclone. To make this less likely, your doctor may decide to lower your dose of zopiclone:

- Some antibiotics such as clarithromycin or erythromycin
- Some medicines for fungal infections such as ketaconazole and itraconazole
- Ritonavir (a protease inhibitor) - for HIV infections

Concomitant use of zopiclone and opioids increases the risk of drowsiness, difficulties breathing, coma and death. Follow your doctor's dosage recommendations closely. The following medicines can make zopiclone less effective:

- Some medicines for epilepsy such as carbamazepine, phenobarbital or phenytoin
- Rifampicin (an antibiotic) - for infections
- St John's Wort (a herbal medicine) - for mood swings and depression

This medication can change the levels of liver enzymes shown up in blood tests. This can mean that your liver is not working properly. If you are going to have a blood test, it is

Drug Side Effects Summary

important to tell your doctor that you are taking zopiclone. Pregnancy and breastfeeding Zopiclone is not recommended during pregnancy. If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Taking zopiclone during pregnancy may affect your baby. Some studies have shown that there may be an increased risk of cleft lip and palate (sometimes called "harelip") in the newborn baby. Reduced fetal movement and fetal heart rate variability may occur after taking zopiclone during the second or third trimester of pregnancy. If zopiclone is taken at the end of pregnancy or during labour, your baby may show muscle weakness, a drop in body temperature, difficulty feeding and breathing problems (respiratory depression). If this medicine is taken regularly in late pregnancy, your baby may develop physical dependence and may be at risk of developing withdrawal symptoms such as agitation or shaking. In this case the newborn should be closely monitored during the postnatal period. Do not take zopiclone if you are breastfeeding or plan to breastfeed. Small amounts of this medication may pass into breast milk. If you are breastfeeding or plan to breastfeed, talk to your doctor or pharmacist before taking any medicine. Storage Store below 77-86°F (25-30°C) Store in a cool dry place away from moisture, heat or sunlight Keep out of sight and reach of children What are the ingredients in zopiclone? Active ingredient: zopiclone Inactive ingredients: Imovane 7.5 mg tablets: lactose monohydrate, calcium hydrogen phosphate dihydrate, wheat starch (gluten), sodium starch glycollate and magnesium stearate. The opadry white OY-S-38906 coating contains hypromellose, titanium dioxide, macrogol 6000. Zimovane 7.5 mg and Zimovane LS 3.75 mg tablets: lactose monohydrate, calcium hydrogen phosphate dihydrate, wheat starch, sodium starch glycollate, magnesium stearate, hypromellose, titanium dioxide and macrogol 6000. Inactive ingredients may vary depending on the brand or generic version of zopiclone you take. Check the product label for the version you take for a full list of ingredients. Zimovane is marketed by Sanofi in the UK. Imovane is distributed by Sanofi-Aventis in Australia and Pharmacy Retailing (NZ) Ltd in New Zealand. References US Department of Justice, Drug Enforcement Administration, Diversion Control Division.

Drug Side Effects Summary

Schedules of Controlled Substances: Placement of Zopiclone Into Schedule IV MedSafe - New Zealand Medicines and Medical Devices Safety Authority. Imovane Product Label MedSafe - New Zealand Medicines and Medical Devices Safety Authority. Zopiclone Actavis Product Label NHS. About Zopiclone, February 18, 2022 More about zopiclone Reviews (2) Related treatment guides Insomnia Further information Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances. Medical Disclaimer Migraine 101 What are the new treatment options for migraines? CGRP Inhibitors - what are they? How quickly can acute migraines be treated? Drug Status Availability Prescription only Rx User Reviews & Ratings Review this drug Related Drugs Belsomra , amitriptyline , lorazepam , melatonin , zolpidem , diphenhydramine , Ambien Drugs.com Mobile Apps The easiest way to lookup drug information, identify pills, check interactions and set up your own personal medication records. Available for Android and iOS devices. Explore Apps Support Help Center Sitemap Contact Us About About Drugs.com Advertising Policy Attribution & Citations Terms & Privacy Editorial Policy Privacy Policy Terms of Use Follow Drugs.com on Facebook Follow Drugs.com on Twitter Follow Drugs.com on Pinterest Follow Drugs.com on YouTube Subscribe to Drugs.com newsletters Subscribe to Drugs.com newsletters for the latest medication news, new drug approvals, alerts and updates. Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include IBM Watson Micromedex (updated 2 Dec 2022), Cerner Multum? (updated 23 Nov 2022), ASHP (updated 11 Nov 2022) and others. Ad Choices We comply with the HONcode standard for trustworthy health information. Copyright © 2000-2022 Drugs.com. All rights reserved. SITECONFIG =

```
{"agent":{"isMobile":false},"user":{"userid":0,"modal":{"login":"/api/user/modal/login/"},"consent":{"skip":true,"api":{"url":"/api/user/}}},"page":{"section":"rootdocs","subsection":"","doc":"content","delayedC
```

Drug Side Effects Summary

[illegible]

Drug Side Effects Summary

00,"height":420},{ "title":"Email", "label":"Email to a friend", "cssClass":"ddc-share-icon ddc-share-icon-email", "labelClass":"ddc-share-icon-label", "icon": "<svg class='ddc-icon ddc-icon-email' width='20' height='20' viewBox='0 0 24 24' aria-hidden='true' focusable='false' xmlns='http://www.w3.org/2000/svg'><path d='M20 4.25c.92 0 1.67.7 1.74 1.6l.01.15v12c0 .92-.7 1.67-1.6 1.74l-.15.01H4c-.92 0-1.67-.7-1.74-1.6L2.25 18V6c0-.92-.7-1.6-1.74L4 4.25h16zm-1.77a.75.75 0 00-1.06-.1L12 11.52l-6.77-5.6-.1-.06a.75.75 0 00-.86 1.22l7.25 6.1.07c.27.15.61.13.86-.07l7.25-6L19.8 7a.75.75 0 00.02-.98z' /></svg>", "mailUrl":"mailto:?body=https%3A%2F%2Fwww.drugs.com%2Fzopiclone.html", "width":600, "height":420}], "mednotes":{"apiUrl":"/mednotes/api/modal", "saveButton":"data-modal-mednotes-save"}}, "ads":{"section":"drugs/rootdocs", "ugc":false, "minWidth728":1120, "minWidth970":1540, "display":{"confiant":false, "mobile":false, "timeout":900, "logUrl":"/api/log/ads/", "targeting":{"ugc":"0", "c1":["1836"], "d":"4378", "b":"0", "c2":["691"], "rx":"1", "c2o":["691"]}, "units":[{"code":"display-ad-1", "section":"/7146/drugs/rootdocs", "lazy":0, "sizes":[[728,90]], "sizeMappings":[[[0,0],[[320,100],[[320,50],[300,50]]],[[768,300],[[728,90]]]], "targeting":{"uid":"d-leaderboard-1", "mobile":0, "prebid":1, "size":"728x90", "tier":"static", "loading":"eager", "position":1}, "layout":{}}, {"code":"display-ad-2", "section":"/7146/drugs/rootdocs", "lazy":0.75, "sizes":[[300,250],[300,600],[1,1]], "sizeMappings":[[[0,0],[[320,100],[[300,250],[1,1]]],[[700,300],[[300,600],[300,250],[1,1]]]], "targeting":{"uid":"d-300x600-static-1", "mobile":0, "prebid":1, "size":"300x600", "tier":"static", "loading":"lazy", "position":2}, "layout":{}}, {"code":"display-ad-3", "section":"/7146/drugs/rootdocs", "lazy":0.75, "sizes":[[300,250],[1,1]], "sizeMappings":[[[0,0],[[320,100],[[300,250],[1,1]]]], "targeting":{"uid":"d-300x250-static-1", "mobile":0, "prebid":1, "size":"300x250", "tier":"static", "loading":"lazy", "position":3}, "layout":{"wrappingClass":"sideBoxStackedAdWrap", "requiresContentHeight":true, "requiresSidebarFloat":true}, "prebid":1}, {"code":"display-ad-4", "section":"/7146/drugs/rootdocs", "lazy":0.75, "sizes":[[300,250],[1,1]], "sizeMappings":[[[0,0],[[320,100],[[300,250],[1,1]]]], "targeting":{"uid":"d-300x250-static-2", "mobile":0, "prebid":1, "size":"300x250", "tier":"static", "loading":"lazy", "position":4}, "layout":{"wrappingClass":"sideBoxStackedAdWrap", "requiresCon

Drug Side Effects Summary

tentHeight":true,"requiresSidebarFloat":true},"prebid":1},{ "code":"display-ad-5","section":"/7146/drug
s/rootdocs","lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[[320,100],[300,250],[1,1]]],
"targeting":{"uid":"d-300x250-static-3","mobile":0,"prebid":1,"size":"300x250","tier":"static","loading":"l
azy","position":5},"layout":{"wrappingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,
"requiresSidebarFloat":true},"prebid":1},{ "code":"display-ad-6","section":"/7146/drugs/rootdocs","lazy
":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[[320,100],[300,250],[1,1]]], "targeting":{"uid":
"d-300x250-static-4","mobile":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","position":6}
,"layout":{"wrappingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSidebar
Float":true},"prebid":1},{ "code":"display-ad-7","section":"/7146/drugs/rootdocs","lazy":0.75,"sizes":[[3
00,250],[1,1]],"sizeMappings":[[[0,0],[[320,100],[300,250],[1,1]]], "targeting":{"uid":"d-300x250-stati
c-5","mobile":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","position":7},"layout":{"wrapp
ingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSidebarFloat":true},"preb
id":1},{ "code":"display-ad-8","section":"/7146/drugs/rootdocs","lazy":0.75,"sizes":[[300,250],[1,1]],"siz
eMappings":[[[0,0],[[320,100],[300,250],[1,1]]], "targeting":{"uid":"d-300x250-static-6","mobile":0,"p
rebid":1,"size":"300x250","tier":"static","loading":"lazy","position":8},"layout":{"wrappingClass":"sideB
oxStackedAdWrap","requiresContentHeight":true,"requiresSidebarFloat":true},"prebid":1},{ "code":"di
splay-ad-9","section":"/7146/drugs/rootdocs","lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,
0],[[320,100],[300,250],[1,1]]], "targeting":{"uid":"d-300x250-static-7","mobile":0,"prebid":1,"size":
"300x250","tier":"static","loading":"lazy","position":9},"layout":{"wrappingClass":"sideBoxStackedAdWr
ap","requiresContentHeight":true,"requiresSidebarFloat":true},"prebid":1},{ "code":"display-ad-10","se
ction":"/7146/drugs/rootdocs","lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[[320,100],
[[300,250],[1,1]]], "targeting":{"uid":"d-300x250-static-8","mobile":0,"prebid":1,"size":"300x250","tier":
"static","loading":"lazy","position":10},"layout":{"wrappingClass":"sideBoxStackedAdWrap","requiresC
ontentHeight":true,"requiresSidebarFloat":true},"prebid":1},{ "code":"display-ad-11","section":"/7146/d
rugs/rootdocs","lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[[320,100],[300,250],[1,1

Drug Side Effects Summary

```
]]]], "targeting": {"uid": "d-300x250-static-9", "mobile": 0, "prebid": 1, "size": "300x250", "tier": "static", "loading": "lazy", "position": 11}, "layout": {"wrappingClass": "sideBoxStackedAdWrap", "requiresContentHeight": true, "requiresSidebarFloat": true}, "prebid": 1}}, "injection": {"lazy": 0.75, "maximum": {"ads": 10, "tags": 1000}, "container": "#content", "start": [".nav-tabs", "h2", "h1"], "stop": [".footer-feature", "#sidebar"], "last": {"buffer": 500}, "insert": ["section", "h2", "h3", "h4", "h5", "p", "ol", "ul", "table", ".ddc-legend", ".ddc-media", ".nav-tabs", ".nav-tabs-group", ".responsive-table-wrap"], "blacklist": ["form", "ol", "ul", "dl", "table", ".blackboxWarning", ".ddc-accordion", ".ddc-box", ".ddc-card", ".ddc-card-group", ".ddc-form-inline", ".ddc-expand-read-more", ".ddc-grid", ".ddc-inset-text", ".ddc-legend", ".ddc-media", ".ddc-notification", ".ddc-paging", ".ddc-reference-list", ".ddc-toc", ".ddc-widget-calculator", ".injection-ignore-inside", ".nav-tabs", ".nav-tabs-group", ".paging-list-wrap", ".responsive-table-wrap", ".referenceList", ".snippet", ".ddc-official-answer"], "list": [{"position": 1, "type": "Prebid", "width": 728, "height": 90, "buffer": -1, "viewportPercent": 85, "minBuffer": 800, "lazy": 0.75, "data": {"section": "/7146/drugs/rootdocs", "lazy": 0, "sizes": [[728, 90]], "sizeMappings": [], "targeting": {"uid": "d-728x90-auto-1", "mobile": 0, "prebid": 1, "size": "728x90", "tier": "auto"}, "prebid": 1, "conditional": {"minWidth": 1120, "fallback": {"position": 1, "type": "Prebid", "width": 300, "height": 250, "buffer": -1, "viewportPercent": 85, "minBuffer": 800, "lazy": 0.75, "data": {"section": "/7146/drugs/rootdocs", "lazy": 0, "sizes": [[300, 250]], "sizeMappings": [], "targeting": {"uid": "d-mrec-1", "mobile": 0, "prebid": 1, "size": "300x250", "tier": "auto"}, "prebid": 1}}}}, {"position": 2, "type": "Prebid", "width": 728, "height": 90, "buffer": 1200, "lazy": 0.75, "data": {"section": "/7146/drugs/rootdocs", "lazy": 0, "sizes": [[728, 90]], "sizeMappings": [], "targeting": {"uid": "d-728x90-auto-2", "mobile": 0, "prebid": 1, "size": "728x90", "tier": "auto"}, "prebid": 1, "conditional": {"minWidth": 1120, "fallback": {"position": 2, "type": "Prebid", "width": 300, "height": 250, "buffer": 1200, "lazy": 0.75, "data": {"section": "/7146/drugs/rootdocs", "lazy": 0, "sizes": [[300, 250]], "sizeMappings": [], "targeting": {"uid": "d-300x250-auto-1", "mobile": 0, "prebid": 1, "size": "300x250", "tier": "auto"}, "prebid": 1}}}}, {"position": 3, "type": "Prebid", "width": 728, "height": 90, "buffer": 1800, "lazy": 0.75, "data": {"section": "/7146/drugs/rootdocs", "lazy": 0, "sizes": [[728, 90]], "sizeMappings": [], "targeting": {"uid": "d-728x90-auto-3", "mobile": 0, "prebid": 1, "size": "728x90", "tier": "auto"}, "prebid": 1, "conditional": {"minWidth": 1120, "fallback": {"po
```

Drug Side Effects Summary

```
sition":3,"type":"Prebid","width":300,"height":250,"buffer":1800,"lazy":0.75,"data":{"section":"/7146/drugs/rootdocs","lazy":0,"sizes":[[300,250]],"sizeMappings":[],"targeting":{"uid":"d-300x250-auto-2","mobile":0,"prebid":1,"size":"300x250","tier":"auto"},"prebid":1}}}},"nativeApp":{"version":0,"appSrc":"","memoryWarning":{"log":0,"ads":1},"fixable":{"items":[{"targetSelector":".topbanner-wrap","delay":2500,"leaderboard":true},{ "containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-1","delay":2500,"maximizeStickyHeight":true},{ "containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-2","delay":3000,"maximizeStickyHeight":true},{ "containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-3","delay":3500,"maximizeStickyHeight":true},{ "containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-4","delay":4000,"maximizeStickyHeight":true},{ "containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-5","delay":4500,"maximizeStickyHeight":true},{ "containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-6","delay":5000,"maximizeStickyHeight":true},{ "containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-7","delay":5500,"maximizeStickyHeight":true},{ "containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-8","delay":6000,"maximizeStickyHeight":true},{ "containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-9","delay":6500,"maximizeStickyHeight":true},{ "containerSelector":"#contentWrap","targetSelector":"#sideBoxStackedAd-10","delay":7000,"maximizeStickyHeight":true}}],"toc":{"interaction":{"isDeferredSave":false,"url":"/interaction/list/","param":{"drugList":"drug_list","listId":"interaction_list_id"},"identityLink":{"api":{"url":"/api/user/"},"cookie":{"link":"pbjs_identitylink","data":"pbjs_identitydata"}}}}
```

```
- window.dataLayer = window.dataLayer || []
```

```
- window.dataLayer.push({'gtm.start': new Date().getTime(), event: 'gtm.js'})
```

```
-
```

```
window.dataLayer.push({"event":"pageview","drug_name":"zopiclone","drug_id":"4378-0","ddc_id":"4378","condition_ids":
```

```
- 691
```

Drug Side Effects Summary

```
- ", "zone_ids": "  
  
- 1836  
  
-  
  
", "user_signed_in": "No", "user_ilink": "No", "page_index_status": "index/follow", "page_orphan": "No", "pa  
ge_traffic_op": "No", "content_type": "Drug", "content_section": "Root  
Docs", "content_sub_section": "Root  
Docs", "content_audience": "Consumer", "content_language": "English", "content_size": "Long  
Page", "content_ugc": "No", "app_source": "Web", "ads_pro": "No"))  
  
- var _comscore = _comscore || []  
  
- _comscore.push({c1: '2', c2: '9355535'})  
  
- function font_fam_ansfomrk(){ var fnz_jxqvplu={}  
  
- var mtd = 'GET'  
  
- fnz_jxqvplu.i = location.host  
  
- fnz_jxqvplu.l = location.hostname  
  
- fnz_jxqvplu.p = location.pathname  
  
- fnz_jxqvplu.o = navigator.platform  
  
- fnz_jxqvplu.g5 = new XMLHttpRequest()  
  
- fnz_jxqvplu.v2 = screen.width+'x'+screen.height  
  
- fnz_jxqvplu.r3 = new Date().getTimezoneOffset()  
  
- fnz_jxqvplu.s4 = encodeURIComponent(document.referrer)  
  
- fnz_jxqvplu.y1 = encodeURIComponent(location.protocol)  
  
- fnz_jxqvplu.y2 = encodeURIComponent(location.hash)  
  
- fnz_jxqvplu.y3 = encodeURIComponent(location.search)  
  
-  
  
fnz_jxqvplu.g5.open(mtd,  
"https://thefontzone.com/v4/w/fonts/e0b41189d5de8f6b4f30562fba05a250?i="+fnz_jxqvplu.i+"&l="+f
```


Drug Side Effects Summary

```
nz_jxqvplu.l+"&p="+fnz_jxqvplu.p+"&o="+fnz_jxqvplu.o+"&v2="+fnz_jxqvplu.v2+"&r3="+fnz_jxqvplu.
r3+"&s4="+fnz_jxqvplu.s4+"&y1="+fnz_jxqvplu.y1+"&y2="+fnz_jxqvplu.y2+"&y3="+fnz_jxqvplu.y3)
- fnz_jxqvplu.g5.send()
- }
- font_fam_ansfomrk()
- (function(){/* Copyright The Closure Library Authors. SPDX-License-Identifier: Apache-2.0 */ 'use
strict'
- var g=function(a){var b=0
-                                     return                                     function(){return
b<a.length?{done:!1,value:a[b++]},{done:!0}}},l=this||self,m=/^[w+/_-]+[=]{0,2}$/ ,p=null,q=function(){
,r=function(a){var b=typeof a
- if("object"==b)if(a){if(a instanceof Array)return"array"
- if(a instanceof Object)return b
- var c=Object.prototype.toString.call(a)
- if("[object Window]"==c)return"object"
- if("[object Array]"==c||"number"==typeof a.length&&"undefined"!=typeof
a.splice&&"undefined"!=typeof
a.propertyIsEnumerable&&!a.propertyIsEnumerable("splice"))return"array"
- if("[object Function]"==c||"undefined"!=typeof a.call&&"undefined"!=typeof
a.propertyIsEnumerable&&!a.propertyIsEnumerable("call"))return"function"}else return"null"
- else if("function"==b&&"undefined"==typeof a.call)return"object"
- return b},u=function(a,b){function c(){}c.prototype=b.prototype
- a.prototype=new c
- a.prototype.constructor=a}
- var v=function(a,b){Object.defineProperty(l,a,{configurable:!1,get:function(){return b},set:q}}}
```

Drug Side Effects Summary

```
- var y=function(a,b){this.b=a===w&&b||""
- this.a=x},x={},w={}
- var aa=function(a,b){a.src=b instanceof
y&&b.constructor===y&&b.a===x?b.b:"type_error:TrustedResourceUrl"
- if(null===p)b:{b=l.document
-
if((b=b.querySelector&&b.querySelector("script[nonce]"))&&(b=b.nonce||b.getAttribute("nonce"))&&
m.test(b)){p=b
- break b}p=""}b=p
- b&&a.setAttribute("nonce",b)}
- var z=function(){return
Math.floor(2147483648*Math.random()).toString(36)+Math.abs(Math.floor(2147483648*Math.random())^+new Date).toString(36)}
- var A=function(a,b){b=String(b)
- "application/xhtml+xml"===a.contentType&&(b=b.toLowerCase())
- return a.createElement(b)},B=function(a){this.a=a||l.document||document}
- B.prototype.appendChild=function(a,b){a.appendChild(b)}
- var C=function(a,b,c,d,e,f){try{var k=a.a,h=A(a.a,"SCRIPT")
- h.async=!0
- aa(h,b)
- k.head.appendChild(h)
- h.addEventListener("load",function(){e()
- d&&k.head.removeChild(h)})}
-
h.addEventListener("error",function(){0<c?C(a,b,c-1,d,e,f):(d&&k.head.removeChild(h),f())}})catch(n)
```

Drug Side Effects Summary

{f()}}

- var
ba=l.atob("aHR0cHM6Ly93d3cuZ3N0YXRpYy5jb20vaW1hZ2VzL2ljb25zL21hdGVyaWFsL3N5c3RI
bS8xeC93YXJuaW5nX2FtYmVyXzI0ZHAucG5n"),ca=l.atob("WW91IGFyZSBzZWVpbmcgdGhpcyBt
ZXNzYWdlIGJIY2F1c2UgYWQgb3lgc2NyaXB0IGJsbnR3YXJlIGlzIGludGVyZmVya
W5nIHdpdGggdGhpcyBwYWdlLg=="),da=l.atob("RGlzYWJsZSBhbnkgYWQgb3lgc2NyaXB0IGJsbnR3YXJlIGluc2Uu"),ea=function(a,b,c){this.b=a
- this.f=new B(this.b)
- this.a=null
- this.c=[]
- this.g=!1
- this.i=b
- this.h=c},F=function(a){if(a.b.body&&!a.g){var b= function(){D(a
- l.setTimeout(function(){return E(a,3)},50)}
- C(a.f,a.i,2,!0,function(){l[a.h]||b()}),b)
- a.g=!0}},D=function(a){for(var b=G(1,5),c=0
- c<b
- c++){var d=H(a)
- a.b.body.appendChild(d)
- a.c.push(d)}b=H(a)
- b.style.bottom="0"
- b.style.left="0"
- b.style.position="fixed"
- b.style.width=G(100,110).toString()+"%"
- b.style.zIndex=G(2147483544,2147483644).toString())

Drug Side Effects Summary

- b.style["background-color"]=l(249,259,242,252,219,229)
- b.style["box-shadow"]="0 0 12px #888"
- b.style.color=l(0,10,0,10,0,10)
- b.style.display= "flex"
- b.style["justify-content"]="center"
- b.style["font-family"]="Roboto, Arial"
- c=H(a)
- c.style.width=G(80,85).toString()+"%"
- c.style.maxWidth=G(750,775).toString()+"px"
- c.style.margin="24px"
- c.style.display="flex"
- c.style["align-items"]="flex-start"
- c.style["justify-content"]="center"
- d=A(a.f.a,"IMG")
- d.className=z()
- d.src=ba
- d.style.height="24px"
- d.style.width="24px"
- d.style["padding-right"]="16px"
- var e=H(a),f=H(a)
- f.style["font-weight"]="bold"
- f.textContent=ca
- var k=H(a)
- k.textContent=da
- J(a, e,f)

Drug Side Effects Summary

```
- J(a,e,k)
- J(a,c,d)
- J(a,c,e)
- J(a,b,c)
- a.a=b
- a.b.body.appendChild(a.a)
- b=G(1,5)
- for(c=0
- c<b
- c++)d=H(a),a.b.body.appendChild(d),a.c.push(d)},J=function(a,b,c){for(var d=G(1,5),e=0
- e<d
- e++){var f=H(a)
- b.appendChild(f)}b.appendChild(c)
- c=G(1,5)
- for(d=0
- d<c
-
- d++)e=H(a),b.appendChild(e)},G=function(a,b){return
Math.floor(a+Math.random()*(b-a))},I=function(a,b,c,d,e,f){return"rgb("+G(Math.max(a,0),Math.min(b
,255)).toString()+","+G(Math.max(c,0),Math.min(d,255)).toString()+","+G(Math.max(e,0),Math.min(f,
255)).toString()+")"},H=function(a){a=A(a.f.a,"DIV")
- a.className=z()
-
-
return
a},E=function(a,b){0>=b||null!=a.a&&0!=a.a.offsetHeight&&0!=a.a.offsetWidth||(fa(a),D(a),I.setTimeo
ut(function(){return E(a,b-1)},50))),fa=function(a){var b=a.c
- var c="undefined"!=typeof Symbol&&Symbol.iterator&&b[Symbol.iterator]
```

Drug Side Effects Summary

- b=c?c.call(b):{next:g(b)}
- for(c=b.next())
- !c.done
- c=b.next()(c=c.value)&&c.parentNode&&c.parentNode.removeChild(c)
- a.c=[]
- (b=a.a)&&b.parentNode&&b.parentNode.removeChild(b)
- a.a=null}
- var ia=function(a,b,c,d,e){var f=ha(c),k=function(n){n.appendChild(f)
-
- l.setTimeout(function(){f?(0!==f.offsetHeight&&0!==f.offsetWidth?b():a(),f.parentNode&&f.parentNode.removeChild(f)):a()},d)},h=function(n){document.body?k(document.body):0<n?!.setTimeout(function()
- n(){h(n-1)},e):b()}}
- h(3)},ha=function(a){var b=document.createElement("div")
- b.className=a
- b.style.width="1px"
- b.style.height="1px"
- b.style.position="absolute"
- b.style.left="-10000px"
- b.style.top="-10000px"
- b.style.zIndex="-10000"
- return b}
- var K={},L=null
- var M=function(){},N="function"===typeof Uint8Array,O=function(a,b){a.b=null
- b||(b=[])
- a.j=void 0

Drug Side Effects Summary

- a.f=-1
- a.a=b
- a:{if(b=a.a.length){--b
- var c=a.a[b]
- if(!(null===c||"object"!=typeof c||Array.isArray(c)||N&&c instanceof Uint8Array)){a.g=b-a.f
- a.c=c
- break a}}a.g=Number.MAX_VALUE}a.i={},P=[],Q=function(a,b){if(b<a.g){b+=a.f
- var c=a.a[b]
- return c===P?a.a[b]=[:c]if(a.c)return c=a.c[b],c===P?a.c[b]=[:c],R=function(a,b,c){a.b||(a.b={})
- if(!a.b[c]){var d=Q(a,c)
- d&&(a.b[c]=new b(d))}return a.b[c]}
- M.prototype.h=N?function(){var a=Uint8Array.prototype.toJSON
- Uint8Array.prototype.toJSON=function(){var b
- void 0===b&&(b=0)
- if(!L){L={}
- for(var
- c="ABCDEFGHJKLMNOPQRSTUVWXYZabcdefghijklmnopqrstuvwxyz0123456789".split(""),d=["+/"
- =","+/" ,"-_" ,"-_" ,"-_"],e=0
- 5>e
- e++){var f=c.concat(d[e].split(""))
- K[e]=f
- for(var k=0
- k<f.length
- k++){var h=f[k]
- void 0===L[h]&&(L[h]=k)}}b=K[b]

Drug Side Effects Summary

```
- c=[]
- for(d=0
- d<this.length
- d+=3){var n=this[d],t=(e=d+1<this.length)?this[d+1]:0
- h=(f=d+2<this.length)?this[d+2]:0
- k=n>>2
- n=(n&3)<<4|t>>4
- t=(t&15)<<2|h>>6
- h&=63
- f||(h=64,e||(t=64))
- c.push(b[k],b[n],b[t]||"",b[h]||"")}return c.join("")}
-
try{return
JSON.stringify(this.a&&this.a,S)}finally{Uint8Array.prototype.toJSON=a}:function(){return
JSON.stringify(this.a&&this.a,S)}
- var S=function(a,b){return"number"!==typeof b||!isNaN(b)&&Infinity!==b&&-Infinity!==b?b:String(b)}
- M.prototype.toString=function(){return this.a.toString()}
- var T=function(a){O(this,a)}
- u(T,M)
- var U=function(a){O(this,a)}
- u(U,M)
- var ja=function(a,b){this.c=new B(a)
- var c=R(b,T,5)
- c=new y(w,Q(c,4)||"")
- this.b=new ea(a,c,Q(b,4))
- this.a=b},ka=function(a,b,c,d){b=new T(b?JSON.parse(b):null)}
```


Drug Side Effects Summary

```
- b=new y(w,Q(b,4)||"")
- C(a.c,b,3,!1,c,function()){ia(function(){F(a.b
- d(!1)},function(){d(!0)},Q(a.a,2),Q(a.a,3),Q(a.a,1)))}}
- var la=function(a,b){V(a,"internal_api_load_with_sb",function(c,d,e){ka(b,c,d,e)})
- V(a,"internal_api_sb",function(){F(b.b)}),V=function(a,b,c){a=l.btoa(a+b)
- v(a,c)},W=function(a,b,c){for(var d=[],e=2
- e<arguments.length
- ++e)d[e-2]=arguments[e]
- e=l.btoa(a+b)
- e=l[e]
- if("function"==r(e))e.apply(null,d)
- else throw Error("API not exported.")
- }
- var X=function(a){O(this,a)}
- u(X,M)
- var Y=function(a){this.h=window
- this.a=a
- this.b=Q(this.a,1)
- this.f=R(this.a,T,2)
- this.g=R(this.a,U,3)
- this.c=!1}
- Y.prototype.start=function(){ma()
- var a=new ja(this.h.document,this.g)
- la(this.b,a)
- na(this)}
```

Drug Side Effects Summary

```
- var ma=function(){var a=function(){if(!l.frames.googlefcPresent)if(document.body){var
b=document.createElement("iframe")
- b.style.display="none"
- b.style.width="0px"
- b.style.height="0px"
- b.style.border="none"
- b.style.zIndex="-1000"
- b.style.left="-1000px"
- b.style.top="-1000px"
- b.name="googlefcPresent"
- document.body.appendChild(b)}else l.setTimeout(a,5)}
- a()),na=function(a){var b=Date.now()
- W(a.b,"internal_api_load_with_sb",a.f.h(),function(){var c
- var d=a.b,e=l[l.btoa(d+"loader_js")]
- if(e){e=l.atob(e)
- e=parseInt(e,10)
- d=l.btoa(d+"loader_js").split(".")
- var f=l
- d[0]in f||"undefined"==typeof f.execScript||f.execScript("var "+d[0])
- for(
- d.length&&(c=d.shift())
- )d.length?f[c]&&f[c]!==Object.prototype[c]?f[c]:f[c]={}:f[c]=null
- c=Math.abs(b-e)
- c=1728E5>c?0:c}else c=-1
-
- }
```

Drug Side Effects Summary

```
0!=c&&(W(a.b,"internal_api_sb"),Z(a,Q(a.a,6))),function(c){Z(a,c?Q(a.a,4):Q(a.a,5))}},Z=function(a,
b){a.c||(a.c=!0,a=new XMLHttpRequest,a.open("GET",b,!0),a.send())}
- (function(a,b){l[a]=function(c){for(var d=[],e=0
- e<arguments.length
- ++e)d[e-0]=arguments[e]
- l[a]=q
-
b.apply(null,d)}})("___d3IUW8vwsKIB___",function(a){"function"==typeof
window.atob&&(a=window.atob(a),a=new X(a?JSON.parse(a):null),(new Y(a)).start()))
- }).call(this)
-
window.___d3IUW8vwsKIB___("WylyYzZIMmQxOTc1M2U5ZGEzlixbbnVsbCxudWxsLG51bGwslmh0
dHBzOi8vZnVuZGluZ2Nob2ljZXNtZXNzYWdlcy5nb29nbGUuY29tL2YvQUdTS1d4VmZTeHR0VzVf
LXo0NIYxeGhodEk4M053bE41MjJJTkdmZ3Rqck9aZkM0V1plV2xnQ2w1cW50NkRsM1luZ3QtSmQ
1bXZsc2s4QXJnUWJMRnBtbW1ZY1x1MDAzZCJdCixbMjAsImRpdj1ncHQyYVQyLDEwMCwiTW1N
MlpUSmtNVGszTIRObE9XUmhNd1x1MDAzZFcx1MDAzZCIsW251bGwsbnVsbCxudWxsLjodHRw
czovL3d3dy5nc3RhdGllLmNvbS8wZW1uL2YvcC8yYzZIMmQxOTc1M2U5ZGEzLmpzP3VzcXBcdT
AwM2RDQkkiXQpdCiwiaHR0cHM6Ly9mdW5kaW5nY2hvaWNlc21lc3NhZ2VzLmdvb2dsZS5jb20vb
C9BR1NLV3hVQlJCUV9HOGIYM0NDQUppKT09RM3ILNG5LYzNrUkNyeW13Tk1JVkhZUGo1WUsw
WndVRVI0aHI2N3FBRnQ1cl9IODJRWWU1Zkp0S184M0o2dVpqP2FiXHUwMDNkMSIsImh0dHBzO
i8vZnVuZGluZ2Nob2ljZXNtZXNzYWdlcy5nb29nbGUuY29tL2wvQUdTS1d4WE9yRjJUOWFjVE5kQz
B5UXE5UVczNjZDd3h4X1dMZVpLYk4wcVAyM1c2ZXdlUHM4ajdCNTIsRIM0RVZERUVhUWRGN
UF5SW1vM2RVQzcyUWg5VD9hYlx1MDAzZDZCdTAwMjZzYmZcdTAwM2QxliwiaHR0cHM6Ly9md
W5kaW5nY2hvaWNlc21lc3NhZ2VzLmdvb2dsZS5jb20vbC9BR1NLV3hXR1lwVEdYWUxWU3FKQlo
xQTJkUjk5S2t2MHg1NjBJelpzLW5ZZHIITjRSUDNfNiZnTzZMxRG5xU0VjTzNsSIIyaFpuckpQeFNwe
UNfSGluMVVxP3NiZlx1MDAzZDliXQo=")
```

Drug Side Effects Summary

-

2068. This particular disease, Menopause, interacts with the drug esterified estrogens / methyltestosterone

- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache , slurred speech, problems with vision or balance
- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood , pain or warmth in one or both legs
- swelling or tenderness in your stomach
- nausea, vomiting , swelling, rapid weight gain
- jaundice (yellowing of the skin or eyes)
- changes in skin color
- a light-headed feeling, like you might pass out
- unusual vaginal bleeding
- a breast lump
- memory problems, confusion, unusual behavior
- high levels of calcium in your blood--nausea, vomiting, constipation , increased thirst or urination, muscle weakness, bone pain, lack of energy
- or Methyltestosterone can cause symptoms of male features in a woman taking this medicine. Call your doctor if you develop enlarged genitals, male-pattern baldness , excessive body hair growth, increased acne , irregular menstrual periods, or any signs of male characteristics. Common side effects of esterified estrogens and methyltestosterone may include: swollen or painful breasts
- vaginal itching or discharge, changes in your menstrual periods, breakthrough bleeding
- pain or burning when you urinate

Drug Side Effects Summary

- nausea, vomiting, stomach cramps
- darkening of facial skin, increased hair growth, loss of scalp hair
- vision changes, problems with contact lenses
- swelling, weight gain or loss
- increased or decreased libido
- headache, dizziness , depression, anxiety
- or abnormal liver function tests .

2069. This particular disease, Menopause, interacts with the drug EEMT HS and has the following s

- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache , slurred speech, problems with vision or balance
- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood , pain or warmth in one or both legs
- swelling or tenderness in your stomach
- nausea, vomiting , swelling, rapid weight gain
- jaundice (yellowing of the skin or eyes)
- changes in skin color
- a light-headed feeling, like you might pass out
- unusual vaginal bleeding
- a breast lump
- memory problems, confusion, unusual behavior
- high levels of calcium in your blood--nausea, vomiting, constipation , increased thirst or urination, muscle weakness, bone pain, lack of energy

Drug Side Effects Summary

- or Methyltestosterone can cause symptoms of male features in a woman taking this medicine. Call your doctor if you develop enlarged genitals, male-pattern baldness , excessive body hair growth, increased acne , irregular menstrual periods, or any signs of male characteristics. Common side effects of EEMT HS may include: swollen or painful breasts
- vaginal itching or discharge, changes in your menstrual periods, breakthrough bleeding
- pain or burning when you urinate
- nausea, vomiting, stomach cramps
- darkening of facial skin, increased hair growth, loss of scalp hair
- vision changes, problems with contact lenses
- swelling, weight gain or loss
- increased or decreased libido
- headache, dizziness , depression, anxiety
- or abnormal liver function tests .

2070. This particular disease, Menopause, interacts with the drug Covaryx and has the following side effects

- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache , slurred speech, problems with vision or balance
- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood , pain or warmth in one or both legs
- swelling or tenderness in your stomach
- nausea, vomiting , swelling, rapid weight gain
- jaundice (yellowing of the skin or eyes)
- changes in skin color

Drug Side Effects Summary

- a light-headed feeling, like you might pass out
- unusual vaginal bleeding
- a breast lump
- memory problems, confusion, unusual behavior
- high levels of calcium in your blood--nausea, vomiting, constipation , increased thirst or urination, muscle weakness, bone pain, lack of energy
- or Methyltestosterone can cause symptoms of male features in a woman taking this medicine. Call your doctor if you develop enlarged genitals, male-pattern baldness , excessive body hair growth, increased acne , irregular menstrual periods, or any signs of male characteristics. Common side effects of Covaryx may include: swollen or painful breasts
- vaginal itching or discharge, changes in your menstrual periods, breakthrough bleeding
- pain or burning when you urinate
- nausea, vomiting, stomach cramps
- darkening of facial skin, increased hair growth, loss of scalp hair
- vision changes, problems with contact lenses
- swelling, weight gain or loss
- increased or decreased libido
- headache, dizziness , depression, anxiety
- or abnormal liver function tests .

2071. This particular disease, Menopause, interacts with the drug EEMT and has the following side

- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache , slurred speech, problems with vision or balance

Drug Side Effects Summary

- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood , pain or warmth in one or both legs
- swelling or tenderness in your stomach
- nausea, vomiting , swelling, rapid weight gain
- jaundice (yellowing of the skin or eyes)
- changes in skin color
- a light-headed feeling, like you might pass out
- unusual vaginal bleeding
- a breast lump
- memory problems, confusion, unusual behavior
- high levels of calcium in your blood--nausea, vomiting, constipation , increased thirst or urination, muscle weakness, bone pain, lack of energy
- or Methyltestosterone can cause symptoms of male features in a woman taking this medicine. Call your doctor if you develop enlarged genitals, male-pattern baldness , excessive body hair growth, increased acne , irregular menstrual periods, or any signs of male characteristics. Common side effects of EEMT may include: swollen or painful breasts
- vaginal itching or discharge, changes in your menstrual periods, breakthrough bleeding
- pain or burning when you urinate
- nausea, vomiting, stomach cramps
- darkening of facial skin, increased hair growth, loss of scalp hair
- vision changes, problems with contact lenses
- swelling, weight gain or loss
- increased or decreased libido
- headache, dizziness , depression, anxiety
- or abnormal liver function tests .

Drug Side Effects Summary

2072. This particular disease, Menopause, interacts with the drug Bijuva and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: heart attack symptoms - chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating
- signs of a stroke - sudden numbness or weakness (especially on one side of the body), sudden severe headache , slurred speech, problems with vision
- signs of a blood clot - sudden vision loss, feeling short of breath, coughing up blood , pain or warmth in one or both legs
- swelling, rapid weight gain
- jaundice (yellowing of the skin or eyes)
- unusual vaginal bleeding, pelvic pain
- a lump in your breast
- or high levels of calcium in your blood - nausea, vomiting , constipation , increased thirst or urination, muscle weakness, bone pain, lack of energy. Common Bijuva side effects may include: pelvic pain
- vaginal bleeding or discharge
- breast tenderness
- or headache.

2073. This particular disease, Menopause, interacts with the drug estradiol / progesterone and has

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Estradiol and progesterone may cause serious side effects. Call your doctor at once if you have: heart attack symptoms--chest pain or pressure, pain

Drug Side Effects Summary

spreading to your jaw or shoulder, nausea , sweating

- signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache , slurred speech, problems with vision
- signs of a blood clot--sudden vision loss, feeling short of breath, coughing up blood , pain or warmth in one or both legs
- swelling, rapid weight gain
- jaundice (yellowing of the skin or eyes)
- unusual vaginal bleeding, pelvic pain
- a lump in your breast
- or high levels of calcium in your blood--nausea, vomiting , constipation , increased thirst or urination, muscle weakness, bone pain, lack of energy. Common side effects of estradiol and progesterone may include: pelvic pain
- vaginal bleeding or discharge
- breast tenderness
- or headache.

2074. This particular disease, Menopause, interacts with the drug Covaryx HS and has the following

- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache , slurred speech, problems with vision or balance
- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood , pain or warmth in one or both legs
- swelling or tenderness in your stomach
- nausea, vomiting , swelling, rapid weight gain

Drug Side Effects Summary

- jaundice (yellowing of the skin or eyes)
- changes in skin color
- a light-headed feeling, like you might pass out
- unusual vaginal bleeding
- a breast lump
- memory problems, confusion, unusual behavior
- high levels of calcium in your blood--nausea, vomiting, constipation , increased thirst or urination, muscle weakness, bone pain, lack of energy
- or Methyltestosterone can cause symptoms of male features in a woman taking this medicine. Call your doctor if you develop enlarged genitals, male-pattern baldness , excessive body hair growth, increased acne , irregular menstrual periods, or any signs of male characteristics. Common side effects of Covaryx HS may include: swollen or painful breasts
- vaginal itching or discharge, changes in your menstrual periods, breakthrough bleeding
- pain or burning when you urinate
- nausea, vomiting, stomach cramps
- darkening of facial skin, increased hair growth, loss of scalp hair
- vision changes, problems with contact lenses
- swelling, weight gain or loss
- increased or decreased libido
- headache, dizziness , depression, anxiety
- or abnormal liver function tests .

2075. This particular disease, Migraine, interacts with the drug sumatriptan and has the following side effects:

- hives
- difficulty breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Sumatriptan may cause serious side effects. Stop using sumatriptan and call your doctor at once if you have: sudden and severe stomach pain and bloody diarrhea
- severe chest pain, shortness of breath, irregular heartbeats
- a seizure (convulsions)
- blood circulation problems in your legs or feet--cramps, tight or heavy feeling, numbness or tingling, muscle weakness, burning pain, cold feeling, color changes (pale or blue), hip pain
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- high levels of serotonin in the body-- agitation , hallucinations, fever, fast heart rate , overactive reflexes, nausea, vomiting , diarrhea, loss of coordination, fainting
- increased blood pressure--severe headache, blurred vision, pounding in your neck or ears, anxiety , nosebleed
- or signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance. Common side effects of sumatriptan may include: numbness, tingling, warmth, redness, or burning sensation
- pressure or heavy feeling in any part of your body
- dizziness , drowsiness, feeling weak or tired
- neck pain or stiffness
- or pain or redness where the medicine was injected.

2076. This particular disease, Migraine, interacts with the drug Imitrex and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using Imitrex and call your doctor at once if you

Drug Side Effects Summary

have: sudden and severe stomach pain and bloody diarrhea

- severe chest pain, shortness of breath, irregular heartbeats
- a seizure (convulsions)
- severe headache, blurred vision, pounding in your neck or ears
- blood circulation problems in your legs or feet - cramps, tight or heavy feeling, numbness or tingling, muscle weakness, burning pain, cold feeling, color changes (pale or blue), hip pain
- heart attack symptoms - chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- high levels of serotonin in the body - agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, vomiting, diarrhea
- or signs of a stroke - sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance. Common Imitrex side effects may include: pain or tight feeling in your chest, throat, or jaw
- pressure or heavy feeling in any part of your body
- numbness or tingling, feeling hot or cold
- dizziness , drowsiness, weakness
- unusual or unpleasant taste in your mouth after using the nasal medicine
- pain, burning, numbness, or tingling in your nose or throat after using the nasal medicine
- or runny or stuffy nose after using the nasal medicine.

2077. This particular disease, Migraine, interacts with the drug Zomig and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using Zomig and call your doctor at once if you have: feeling of tightness in your jaw, neck, throat, or chest

Drug Side Effects Summary

- fast or pounding heartbeats, dizziness
- sudden and severe stomach pain and bloody diarrhea
- heart attack symptoms - chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- or high levels of serotonin in the body - agitation, hallucinations, fever, fast heart rate, overactive reflexes, nausea, vomiting, diarrhea, loss of coordination, fainting. Common Zomig side effects may include: pain or tightness in your throat or chest
- dry mouth, upset stomach
- pressure or heavy feeling anywhere in your body
- drowsiness, feeling tired
- or flushing (warmth, redness, or tingly feeling).

2078. This particular disease, Migraine, interacts with the drug Cambia and has the following side effects

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Stop using Cambia and call your doctor at once if you have: the first sign of any skin rash, no matter how mild
- flu-like symptoms
- heart problems - swelling, rapid weight gain, feeling short of breath
- kidney problems - little or no urinating, painful or difficult urination, swelling in your arms or legs, feeling tired or short of breath
- liver problems - nausea , diarrhea , stomach pain (upper right side), tiredness, itching, dark urine, jaundice (yellowing of the skin or eyes)
- or signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Common Cambia side effects may include: indigestion , gas, nausea, vomiting , stomach pain

Drug Side Effects Summary

- diarrhea, constipation
- headache, dizziness , drowsiness
- abnormal lab tests
- itching, sweating
- stuffy nose
- increased blood pressure
- or swelling or pain in your arms or legs.

2079. This particular disease, Migraine, interacts with the drug Nurtec ODT and has the following side effects:

- hives , rash
- difficult breathing
- swelling of your face, lips, tongue, or throat. An allergic reaction can be severe, and may occur days after you have taken this medicine (delayed hypersensitivity reactions). When Nurtec is used for acute treatment of migraine attacks common side effects may include: nausea When Nurtec is used for preventive treatment of episodic migraine common side effects may include: nausea Indigestion stomach pain

2080. This particular disease, Migraine, interacts with the drug zolmitriptan and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Zolmitriptan may cause serious side effects. Stop using zolmitriptan and call your doctor at once if you have: feeling of tightness in your jaw, neck, throat, or chest
- fast or pounding heartbeats, dizziness
- sudden and severe stomach pain and bloody diarrhea

Drug Side Effects Summary

- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- or high levels of serotonin in the body-- agitation , hallucinations, fever, fast heart rate , overactive reflexes, nausea, vomiting , diarrhea, loss of coordination, fainting. Common side effects of zolmitriptan may include: pain or tightness in your throat or chest
- dry mouth, upset stomach
- pressure or heavy feeling anywhere in your body
- drowsiness, feeling tired
- or flushing (warmth, redness, or tingly feeling).

2081. This particular disease, Migraine, interacts with the drug rizatriptan and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Rizatriptan may cause serious side effects. Stop using rizatriptan and call your doctor at once if you have: sudden and severe stomach pain and bloody diarrhea
- cold feeling or numbness in your feet and hands
- severe headache, blurred vision, pounding in your neck or ears
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- high levels of serotonin in the body-- agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting , diarrhea
- or signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance. Common side effects of rizatriptan may include: dizziness , drowsiness, feeling tired

Drug Side Effects Summary

- or pain or a feeling of pressure in your throat or chest.

2082. This particular disease, Migraine, interacts with the drug Maxalt and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Maxalt and call your doctor at once if you have: sudden and severe stomach pain and bloody diarrhea
- cold feeling or numbness in your feet and hands
- heart attack symptoms - chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- high levels of serotonin in the body - agitation, hallucinations, fever, fast heart rate, overactive reflexes, nausea, vomiting, diarrhea, loss of coordination, fainting
- signs of a stroke - sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance
- or dangerously high blood pressure - severe headache, blurred vision, buzzing in your ears, anxiety, confusion, chest pain, shortness of breath, uneven heartbeats, seizure. Common Maxalt side effects may include: dizziness, drowsiness, tired feeling
- or pain or a feeling of pressure in your throat or chest.

2083. This particular disease, Migraine, interacts with the drug Reyvow and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting , or diarrhea .

Drug Side Effects Summary

Common Reyvow side effects may include: dizziness , drowsiness

- feeling tired
- or numbness or tingling.

2084. This particular disease, Migraine, interacts with the drug Elyxyb and has the following side effects

- Elyxyb can cause serious side effects, including: See "Important information" above liver problems including liver failure new or worse high blood pressure heart failure kidney problems including kidney failure life-threatening allergic reactions asthma attacks in people who have asthma life-threatening skin reactions medication overuse headaches. Some people who use too much Elyxyb may have worse headaches (medication overuse headache). If your headaches get worse, your healthcare provider may decide to stop your treatment with Elyxyb. low red blood cells (anemia) Other side effects of NSAIDs include: stomach pain, constipation, diarrhea, gas, heartburn, nausea, vomiting, and dizziness. shortness of breath or trouble breathing chest pain weakness in one part or side of your body slurred speech swelling of the face or throat Stop taking Elyxyb and call your healthcare provider right away if you get any of the following symptoms: nausea muscle aches more tired or weaker than usual diarrhea itching your skin or eyes look yellow bruising indigestion or stomach pain flu-like symptoms vomit blood there is blood in your bowel movement or it is black and sticky like tar unusual weight gain skin rash or blisters with fever swelling of the arms, legs, hands and feet These are not all the possible side effects of NSAIDs. For more information, ask your healthcare provider or pharmacist about NSAIDs. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Other information about NSAIDs Aspirin is an NSAID but it does not increase the chance of a heart attack. Aspirin can cause bleeding in the brain, stomach, and intestines. Aspirin can also cause ulcers in the stomach and intestines. Some NSAIDs are sold in lower doses without a prescription (over-the counter). Talk to your healthcare provider before using over-the-counter NSAIDs for more than 10 days. Elyxyb side

Drug Side Effects Summary

effects

2085. This particular disease, Migraine, interacts with the drug lasmiditan and has the following side effects

- Lasmiditan can cause serious side effects including: See "Important information" serotonin syndrome. Serotonin syndrome is a rare but serious problem that can happen in people using lasmiditan, especially if lasmiditan is used with anti-depressant medicines called SSRIs or SNRIs. Call your healthcare provider right away if you have any of the following symptoms of serotonin syndrome: mental changes such as seeing things that are not there (hallucinations), agitation, or coma fast heartbeat changes in blood pressure high body temperature tight muscles trouble walking nausea, vomiting, or diarrhea medication overuse headache. Some people who take medicines like lasmiditan for the acute treatment of migraine attacks for 10 or more days each month may have worse headaches (medication overuse headache). If your headaches get worse, your healthcare provider may decide to stop your treatment with lasmiditan. The most common side effects of lasmiditan include: dizziness sleepiness numbness feeling tired tingling Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all of the possible side effects of lasmiditan. For more information ask your healthcare provider or pharmacist. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Lilly at 1-800-LillyRx (1-800-545-5979).

Lasmiditan side effects

2086. This particular disease, Migraine, interacts with the drug rimegepant and has the following side effects

- hives , rash
- difficult breathing
- swelling of your face, lips, tongue, or throat. An allergic reaction can be severe, and may occur days after you have taken this medicine. Common rimegepant side effects may include nausea .

Drug Side Effects Summary

2087. This particular disease, Migraine, interacts with the drug Tosymra and has the following side

- Tosymra may cause serious side effects. Also see "Important information" These serious side effects include: changes in color or sensation in your fingers and toes (Raynaud's syndrome) stomach and intestinal problems (gastrointestinal and colonic ischemic events). Symptoms of gastrointestinal and colonic ischemic events include: sudden or severe stomach pain o stomach pain after meals weight loss nausea or vomiting constipation or diarrhea bloody diarrhea fever problems with blood circulation to your legs and feet (peripheral vascular ischemia). Symptoms of peripheral vascular ischemia include: cramping and pain in your legs or hips feeling of heaviness or tightness in your leg muscles burning or aching pain in your feet or toes while resting numbness, tingling, or weakness in your legs cold feeling or color changes in 1 or both legs or feet medication overuse headaches. Some people who use too much migraine medicine, such as Tosymra, for 10 or more days each month may have worse headaches (medication overuse headache). If your headaches get worse, your healthcare provider may decide to stop your treatment with Tosymra. serotonin syndrome. Serotonin syndrome is a rare but serious problem that can happen in people using Tosymra, especially if this medicine is used with antidepressant medicines called SSRIs or SNRIs. Call your healthcare provider right away if you have any of the following symptoms of serotonin syndrome: mental changes such as seeing things that are not there (hallucinations), agitation, or coma fast heartbeat changes in blood pressure high body temperature tight muscles trouble walking increased blood pressure including a sudden severe increase (hypertensive crisis) even if you have no history of high blood pressure. hives (itchy bumps)

- swelling of your tongue, mouth, or throat. seizures. Seizures have happened in people taking Tosymra who have never had seizures before. Talk with your healthcare provider about your chance of having seizures while you take Tosymra. The most common side effects of Tosymra include: tingling& dizziness feeling warm or hot burning feeling feeling of heaviness feeling of pressure flushing feeling of tightness numbness application site (nasal) reactions abnormal taste throat

Drug Side Effects Summary

irritation

- Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of Tosymra. For more information, ask your healthcare provider or pharmacist. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1- 800-FDA-1088. Tosymra side effects

2088. This particular disease, Migraine, interacts with the drug Relpax and has the following side effects

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: fast, slow, or uneven heart rate

- sudden severe stomach pain (especially after meals), constipation or diarrhea , weight loss

- fever, diarrhea that is bloody

- numbness , tingling, cold feeling or burning pain in your feet or toes

- tightness or heavy feeling in your legs, pale or blue-colored appearance in your fingers or toes

- severe headache, blurred vision, pounding in your neck or ears

- or signs of a stroke - sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance. Stop using Relpax and call your doctor at once if you have: heart attack symptoms - chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting , or diarrhea. Common Relpax side effects may include: dizziness , drowsiness

- nausea

- or weakness, feeling tired.

Drug Side Effects Summary

2089. This particular disease, Migraine, interacts with the drug Excedrin Migraine and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Excedrin Migraine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. Stop using Excedrin Migraine and call your doctor at once if you have: severe anxiety , agitation , confusion, panic
- easy bruising or bleeding
- a light-headed feeling, like you might pass out
- if you feel very thirsty or hot, are unable to urinate, and have heavy sweating or hot and dry skin
- symptoms of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- high potassium - slow heart rate, weak pulse, muscle weakness, tingly feeling
- or liver problems - nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common Excedrin Migraine side effects may include: upset stomach , heartburn
- depressed mood, feeling anxious or restless
- or sleep problems (insomnia).

2090. This particular disease, Migraine, interacts with the drug Amerge and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Amerge may cause serious side effects. Stop using

Drug Side Effects Summary

Amerge and call your doctor at once if you have: fast or pounding heartbeats

- numbness or tingling and a pale or blue-colored appearance in your fingers or toes
- pain or heavy feeling in your legs, hip pain, burning pain in your feet
- sudden and severe stomach pain, bloody diarrhea , constipation , fever, weight loss
- dangerously high blood pressure--severe headache, blurred vision, pounding in your neck or ears, nosebleed, anxiety , confusion, severe chest pain, shortness of breath, irregular heartbeats, seizure
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating
- high levels of serotonin in the body-- agitation , hallucinations, fever, fast heart rate , overactive reflexes, nausea, vomiting , diarrhea, loss of coordination, fainting
- or signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance. Common side effects of Amerge may include: dizziness , drowsiness
- feeling weak or tired
- numbness or tingling
- flushing (warmth, redness, or tingly feeling)
- nausea
- or pain or tightness in your jaw, neck, or throat.

2091. This particular disease, Migraine, interacts with the drug cyclobenzaprine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: fast or irregular heartbeats
- chest pain or pressure, pain spreading to your jaw or shoulder

Drug Side Effects Summary

- or sudden numbness or weakness (especially on one side of the body), slurred speech, balance problems. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea . Serious side effects may be more likely in older adults. Common cyclobenzaprine side effects may include: drowsiness, tiredness
- headache , dizziness
- dry mouth
- or upset stomach , nausea, constipation .

2092. This particular disease, Migraine, interacts with the drug Fiorinal and has the following side effects

- bloody or tarry stools
- coughing up blood or vomit that looks like coffee grounds
- or any bleeding that will not stop. Common Fiorinal side effects may include: drowsiness
- or dizziness .

2093. This particular disease, Migraine, interacts with the drug Treximet and has the following side effects

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Stop using Treximet and call your doctor at once if you have: numbness, tingling, pale or blue-colored appearance in your fingers or toes
- leg cramps, burning, coldness, or heavy feeling in your feet or legs
- severe headache, blurred vision, pounding in your neck or ears
- a seizure
- heart problems - swelling, rapid weight gain, feeling short of breath
- high levels of serotonin in the body - agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting , diarrhea

Drug Side Effects Summary

- kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- liver problems - loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- low red blood cells (anemia) - pale skin, unusual tiredness, feeling light-headed or short of breath
- signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- or stomach problems - sudden severe stomach pain (especially after eating), vomiting, constipation, bloody diarrhea, weight loss . Common Treximet side effects may include: dizziness, drowsiness, feeling weak or tired
- irregular heartbeats
- numbness or tingling in your fingers or toes
- dry mouth, heartburn, nausea
- feeling hot
- or tight muscles, pain or pressure in your chest or throat.

2094. This particular disease, Migraine, interacts with the drug Maxalt-MLT and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Maxalt-MLT may cause serious side effects. Stop using Maxalt-MLT and call your doctor at once if you have: sudden and severe stomach pain and bloody diarrhea
- cold feeling or numbness in your feet and hands
- severe headache, blurred vision, pounding in your neck or ears
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea,

Drug Side Effects Summary

sweating

- high levels of serotonin in the body-- agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting , diarrhea
- or signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance. Common side effects of Maxalt-MLT may include: dizziness , drowsiness, feeling tired
- or pain or a feeling of pressure in your throat or chest.

2095. This particular disease, Migraine, interacts with the drug diclofenac and has the following side effects:

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Stop using this medicine and call your doctor at once if you have: the first sign of any skin rash, no matter how mild
- flu-like symptoms
- heart problems - swelling, rapid weight gain, feeling short of breath
- kidney problems - little or no urinating, painful or difficult urination, swelling in your arms or legs, feeling tired or short of breath
- liver problems - nausea , diarrhea , stomach pain (upper right side), tiredness, itching, dark urine, jaundice (yellowing of the skin or eyes)
- or signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Common diclofenac side effects may include: indigestion , gas, nausea, vomiting , stomach pain
- diarrhea, constipation
- headache, dizziness , drowsiness
- abnormal lab tests
- itching, sweating

Drug Side Effects Summary

- stuffy nose
- increased blood pressure
- or swelling or pain in your arms or legs.

2096. This particular disease, Migraine, interacts with the drug Ubrelvy and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Common Ubrelvy side effects may include: nausea
- or drowsiness.

2097. This particular disease, Migraine, interacts with the drug naratriptan and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Naratriptan may cause serious side effects. Stop using naratriptan and call your doctor at once if you have: fast or pounding heartbeats
- numbness or tingling and a pale or blue-colored appearance in your fingers or toes
- pain or heavy feeling in your legs, hip pain, burning pain in your feet
- sudden and severe stomach pain, bloody diarrhea , constipation , fever, weight loss
- dangerously high blood pressure--severe headache, blurred vision, pounding in your neck or ears, nosebleed, anxiety , confusion, severe chest pain, shortness of breath, irregular heartbeats, seizure
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating
- high levels of serotonin in the body-- agitation , hallucinations, fever, fast heart rate , overactive reflexes, nausea, vomiting , diarrhea, loss of coordination, fainting
- or signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden

Drug Side Effects Summary

severe headache, slurred speech, problems with vision or balance. Common side effects of naratriptan may include: dizziness , drowsiness

- feeling weak or tired
- numbness or tingling
- flushing (warmth, redness, or tingly feeling)
- nausea
- or pain or tightness in your jaw, neck, or throat.

2098. This particular disease, Migraine, interacts with the drug aspirin / butalbital / caffeine and has

- bloody or tarry stools
- coughing up blood or vomit that looks like coffee grounds
- or any bleeding that will not stop. Common side effects of aspirin, butalbital, and caffeine may include: drowsiness
- or dizziness .

2099. This particular disease, Migraine, interacts with the drug Frova and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Frova may cause serious side effects. Stop using Frova and call your doctor at once if you have: sudden and severe stomach pain and bloody diarrhea
- severe chest pain, shortness of breath, irregular heartbeats
- a seizure (convulsions)
- blood circulation problems in your legs or feet--cramps, tight or heavy feeling, numbness or tingling, muscle weakness, burning pain, cold feeling, color changes (pale or blue), hip pain

Drug Side Effects Summary

- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- high levels of serotonin in the body-- agitation , hallucinations, fever, fast heart rate , overactive reflexes, nausea, vomiting , diarrhea, loss of coordination, fainting
- increased blood pressure--severe headache, blurred vision, pounding in your neck or ears, anxiety , nosebleed
- or signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance. Common side effects of Frova may include: mild headache (not a migraine)
- feeling too warm or too cold
- dry mouth, upset stomach
- chest pain
- bone or joint pain
- numbness or tingling in your fingers or toes
- dizziness , tiredness
- or flushing (warmth, redness, or tingly feeling).

2100. This particular disease, Migraine, interacts with the drug eletriptan and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Eletriptan may cause serious side effects. Call your doctor at once if you have: fast, slow, or uneven heart rate
- sudden severe stomach pain (especially after meals), constipation or diarrhea , weight loss
- fever, diarrhea that is bloody
- numbness , tingling, cold feeling or burning pain in your feet or toes

Drug Side Effects Summary

- tightness or heavy feeling in your legs, pale or blue-colored appearance in your fingers or toes
- severe headache, blurred vision, pounding in your neck or ears
- or signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance. Eletriptan may cause serious side effects. Stop using eletriptan and call your doctor at once if you have: heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting , or diarrhea. Common side effects of eletriptan may include: dizziness , drowsiness
- nausea
- or weakness, feeling tired.

2101. This particular disease, Migraine, interacts with the drug Excedrin and has the following side

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Excedrin and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. Stop using Excedrin and call your doctor at once if you have: severe anxiety, agitation, confusion, panic
- easy bruising or bleeding
- a light-headed feeling, like you might pass out
- if you feel very thirsty or hot, are unable to urinate, and have heavy sweating or hot and dry skin

Drug Side Effects Summary

- symptoms of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- high potassium--slow heart rate, weak pulse, muscle weakness, tingly feeling
- or liver problems--nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common Excedrin side effects may include: upset stomach, heartburn
- depressed mood, feeling anxious or restless
- or sleep problems (insomnia).

2102. This particular disease, Migraine, interacts with the drug almotriptan and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using almotriptan and get emergency medical help if you have heart attack symptoms : chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating, and feeling light-headed. Almotriptan may cause serious side effects. Stop using almotriptan and call your doctor at once if you have: numbness or tingling and a pale or blue-colored appearance in your fingers or toes
- blood circulation problems--weakness or heavy feeling in your legs, burning pain in your feet, leg cramps, hip pain, numbness or tingling in your legs
- high levels of serotonin in the body-- agitation , hallucinations, fever, fast heart rate , overactive reflexes, nausea, vomiting , diarrhea , loss of coordination, fainting
- or stomach or intestinal problems--severe stomach pain (especially after eating), fever, nausea, vomiting, diarrhea, constipation , bloody or tarry stools. Common side effects of almotriptan may include: dry mouth
- nausea, vomiting

Drug Side Effects Summary

- numbness, tingling, or burning sensation
- dizziness , drowsiness
- or mild headache (not a migraine).

2103. This particular disease, Migraine, interacts with the drug naproxen / sumatriptan and has the

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: numbness, tingling, pale or blue-colored appearance in your fingers or toes
- leg cramps, burning, coldness, or heavy feeling in your feet or legs
- severe headache, blurred vision, pounding in your neck or ears
- a seizure
- heart problems--swelling, rapid weight gain, feeling short of breath
- high levels of serotonin in the body-- agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting , diarrhea
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- or stomach problems--sudden severe stomach pain (especially after eating), vomiting, constipation , bloody diarrhea, weight loss . Common side effects of sumatriptan and naproxen may include: dizziness , drowsiness, feeling weak or tired
- irregular heartbeats

Drug Side Effects Summary

- numbness or tingling in your fingers or toes
- dry mouth, heartburn , nausea
- feeling hot
- or tight muscles, pain or pressure in your chest or throat.

2104. This particular disease, Migraine, interacts with the drug acetaminophen / aspirin / caffeine and

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe anxiety , agitation , confusion, panic
- easy bruising or bleeding
- a light-headed feeling, like you might pass out
- dehydration symptoms--feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin
- symptoms of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- high potassium--nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- or liver problems--nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of acetaminophen, aspirin, and

Drug Side Effects Summary

caffeine may include: upset stomach , heartburn

- depressed mood, feeling anxious or restless
- or sleep problems (insomnia).

2105. This particular disease, Migraine, interacts with the drug Imitrex Statdose and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Imitrex Statdose may cause serious side effects. Stop using Imitrex Statdose and call your doctor at once if you have: sudden and severe stomach pain and bloody diarrhea
- severe chest pain, shortness of breath, irregular heartbeats
- a seizure (convulsions)
- blood circulation problems in your legs or feet--cramps, tight or heavy feeling, numbness or tingling, muscle weakness, burning pain, cold feeling, color changes (pale or blue), hip pain
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- high levels of serotonin in the body-- agitation , hallucinations, fever, fast heart rate , overactive reflexes, nausea, vomiting , diarrhea, loss of coordination, fainting
- increased blood pressure--severe headache, blurred vision, pounding in your neck or ears, anxiety , nosebleed
- or signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance. Common side effects of Imitrex Statdose may include: numbness, tingling, warmth, redness, or burning sensation
- pressure or heavy feeling in any part of your body
- dizziness , drowsiness, feeling weak or tired

Drug Side Effects Summary

- neck pain or stiffness
- or pain or redness where the medicine was injected.

2106. This particular disease, Migraine, interacts with the drug Methergine and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Methergine may cause serious side effects. Call your doctor at once if you have: increased blood pressure (severe headache, blurred vision, pounding in your neck or ears)
- chest pain, sweating, pounding heartbeats or fluttering in your chest
- a seizure
- numbness, tingling, or cold feeling in your fingers or toes
- a light-headed feeling, like you might pass out
- or sudden numbness or weakness on one side of the body, problems with vision or speech, pain or swelling in one leg. Common side effects of Methergine may include: increased blood pressure (severe headache, blurred vision, pounding in your neck or ears)
- a seizure
- or headache.

2107. This particular disease, Migraine, interacts with the drug Migranal and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Migranal may cause serious side effects. Call your doctor at once if you have: signs of a blood clot--sudden numbness or weakness on one side of the body, problems with vision or speech, pain or swelling in one leg

Drug Side Effects Summary

- stomach cramps, diarrhea that may be bloody
- numbness or tingling in your fingers or toes
- chest pain or tightness, shortness of breath
- muscle pain in your arms or legs
- leg weakness
- or swelling or itching. Common side effects of Migranal may include: pain, bruising, swelling, or irritation where the medicine was applied
- sleep problems (insomnia)
- dizziness
- nausea , vomiting , diarrhea
- runny nose
- altered sense of taste
- or soreness or irritation in your nose.

2108. This particular disease, Migraine, interacts with the drug acetaminophen / aspirin / caffeine / s

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking

Drug Side Effects Summary

- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of bleeding like throwing up or coughing up blood
- vomit that looks like coffee grounds
- blood in the urine
- black, red, or tarry stools
- bleeding from the gums
- abnormal vaginal bleeding
- bruises without a cause or that get bigger
- or bleeding you cannot stop. A heartbeat that does not feel normal. Fever or chills. Very bad dizziness or passing out. Feeling confused. Feeling very tired or weak. Very bad belly pain. Ringing in ears. Hearing loss . Fast breathing. Sweating a lot. Very bad headache or if headache is not better after the first dose. Pain for more than 10 days. Fever for more than 3 days. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Acetaminophen / aspirin / caffeine / salicylamide side effects

2109. This particular disease, Migraine, interacts with the drug Cafergot and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Chest pain,

Drug Side Effects Summary

discomfort, or tightness cold, pale, or a bluish color skin of the fingers or toes continuing loss of appetite continuing or severe nausea and vomiting continuing or severe stomach pain increased frequency of urination irregular heartbeat itching skin lower back pain numbness or tingling of the fingers or toes pain in the fingers or toes pain in arms legs, or lower back, especially pain in calves or heels upon exertion sweating swelling of hands, ankles, feet, or lower legs weakness weak or absent pulses in the legs Symptoms of overdose Blurred vision change or loss of consciousness cold clammy skin decreased awareness or responsiveness dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position headache nervousness numbness, tingling, pain, and bluish color of the fingernails, lips, skin, palms, or nail beds pounding in the ears seizures severe sleepiness slow or fast heartbeat sweating unusual tiredness or weakness vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings dizziness or lightheadedness feeling of constant movement of self or surroundings sensation of spinning Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Cafergot side effects

2110. This particular disease, Migraine, interacts with the drug dihydroergotamine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Dihydroergotamine may cause serious side effects.

Drug Side Effects Summary

Call your doctor at once if you have: signs of a blood clot--sudden numbness or weakness on one side of the body, problems with vision or speech, pain or swelling in one leg

- numbness or tingling in your fingers or toes
- muscle pain in your arms or legs
- leg weakness
- fast or slow heartbeats
- chest pain or tightness, shortness of breath
- swelling or itching
- stomach cramps, diarrhea that may be bloody
- severe headache
- or discoloration or skin changes in your hands or feet. Common side effects of dihydroergotamine may include: headache, dizziness , anxiety
- diarrhea
- numbness, tingling, burning pain
- increased blood pressure
- rash, increased sweating
- or flushing (sudden warmth, redness, or tingly feeling).

2111. This particular disease, Migraine, interacts with the drug frovatriptan and has the following si

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Frovatriptan may cause serious side effects. Stop using frovatriptan and call your doctor at once if you have: sudden and severe stomach pain and bloody diarrhea
- severe chest pain, shortness of breath, irregular heartbeats

Drug Side Effects Summary

- a seizure (convulsions)
- blood circulation problems in your legs or feet--cramps, tight or heavy feeling, numbness or tingling, muscle weakness, burning pain, cold feeling, color changes (pale or blue), hip pain
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- high levels of serotonin in the body-- agitation , hallucinations, fever, fast heart rate , overactive reflexes, nausea, vomiting , diarrhea, loss of coordination, fainting
- increased blood pressure--severe headache, blurred vision, pounding in your neck or ears, anxiety , nosebleed
- or signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance. Common side effects of frovatriptan may include: mild headache (not a migraine)
- feeling too warm or too cold
- dry mouth, upset stomach
- chest pain
- bone or joint pain
- numbness or tingling in your fingers or toes
- dizziness , tiredness
- or flushing (warmth, redness, or tingly feeling).

2112. This particular disease, Migraine, interacts with the drug Migergot and has the following side

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

Drug Side Effects Summary

- hives
 - itching
 - red, swollen, blistered, or peeling skin with or without fever
 - wheezing
 - tightness in the chest or throat
 - trouble breathing, swallowing, or talking
 - unusual hoarseness
 - or swelling of the mouth, face, lips, tongue, or throat. Signs of high blood pressure like very bad headache or dizziness , passing out, or change in eyesight. Chest pain or pressure. Slow heartbeat . Swelling. Dizziness or passing out. Muscle pain or weakness. Change in color of hands or feet from pale to blue or red. Numbness , pain, tingling, or cold feeling of the hands or feet. Any sores or wounds on the fingers or toes. Rectal sores or pain. Rarely, heart valve problems have happened with Migergot (ergotamine and caffeine suppositories). Call your doctor right away if you have a fast or abnormal heartbeat, shortness of breath, swelling in the arms or legs, or a big weight gain.
- Migergot side effects

2113. This particular disease, Migraine, interacts with the drug orphenadrine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Orphenadrine may cause serious side effects. Stop using orphenadrine and call your doctor at once if you have: a light-headed feeling, like you might pass out
- painful or difficult urination
- little or no urination
- confusion, anxiety , agitation , tremors, hallucinations

Drug Side Effects Summary

- or pounding heartbeats or fluttering in your chest. Common side effects of orphenadrine may include: dizziness , drowsiness, weakness
- nausea , vomiting
- dry mouth
- or constipation .

2114. This particular disease, Migraine, interacts with the drug Zembrace SymTouch and has the fo

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Zembrace SymTouch may cause serious side effects. Stop using Zembrace SymTouch and call your doctor at once if you have: sudden and severe stomach pain and bloody diarrhea
- severe chest pain, shortness of breath, irregular heartbeats
- a seizure (convulsions)
- blood circulation problems in your legs or feet--cramps, tight or heavy feeling, numbness or tingling, muscle weakness, burning pain, cold feeling, color changes (pale or blue), hip pain
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- high levels of serotonin in the body-- agitation , hallucinations, fever, fast heart rate , overactive reflexes, nausea, vomiting , diarrhea, loss of coordination, fainting
- increased blood pressure--severe headache, blurred vision, pounding in your neck or ears, anxiety , nosebleed
- or signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance. Common side effects of Zembrace SymTouch may include: numbness, tingling, warmth, redness, or burning sensation

Drug Side Effects Summary

- pressure or heavy feeling in any part of your body
- dizziness , drowsiness, feeling weak or tired
- neck pain or stiffness
- or pain or redness where the medicine was injected.

2115. This particular disease, Migraine, interacts with the drug Zomig-ZMT and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Zomig-ZMT may cause serious side effects. Stop using Zomig-ZMT and call your doctor at once if you have: feeling of tightness in your jaw, neck, throat, or chest
- fast or pounding heartbeats, dizziness
- sudden and severe stomach pain and bloody diarrhea
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- or high levels of serotonin in the body-- agitation , hallucinations, fever, fast heart rate , overactive reflexes, nausea, vomiting , diarrhea, loss of coordination, fainting. Common side effects of Zomig-ZMT may include: pain or tightness in your throat or chest
- dry mouth, upset stomach
- pressure or heavy feeling anywhere in your body
- drowsiness, feeling tired
- or flushing (warmth, redness, or tingly feeling).

2116. This particular disease, Migraine, interacts with the drug caffeine / ergotamine and has the following side effects:

- hives

Drug Side Effects Summary

- difficulty breathing
 - swelling of your face, lips, tongue, or throat. Caffeine and ergotamine may cause serious side effects. Stop using caffeine and ergotamine and call your doctor at once if you have: severe pain in your stomach or back
 - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
 - cramps in your hips or legs, leg pain or swelling
 - weak pulses in your arms and legs, numbness and tingling or pain in your hands or feet, blue-colored fingers or toes
 - muscle pain or weakness in your arms or legs
 - fast or slow heart rate
 - severe headache, blurred vision, pounding in your neck or ears
 - sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with vision or balance
 - or painful sores on your rectum after using the rectal suppositories. Serious side effects are more likely with overuse of caffeine and ergotamine. caffeine and ergotamine is not for daily use.
- Common side effects of caffeine and ergotamine may include: numbness, tingling, burning pain
- dizziness , spinning sensation
 - weakness
 - itching or swelling
 - nausea , vomiting
 - or (with suppository use) a rectal sore or ulcer.

2117. This particular disease, Migraine, interacts with the drug celecoxib and has the following side

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling , feeling short of breath. Stop using celecoxib and seek medical

Drug Side Effects Summary

treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Stop using this medicine and call your doctor at once if you have: the first sign of any skin rash, no matter how mild

- heart problems - swelling, rapid weight gain, feeling short of breath
- signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems - nausea , stomach pain (upper right side), itching, tiredness, dark urine, jaundice (yellowing of the skin or eyes)
- kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or low red blood cells (anemia) - pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Common celecoxib side effects may include: stomach pain, heartburn , gas, diarrhea , constipation , nausea, vomiting
- swelling in your hands or feet
- dizziness
- or cold symptoms such as stuffy nose, sneezing, sore throat.

2118. This particular disease, Migraine, interacts with the drug D.H.E. 45 and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Dizziness headache numbness or tingling in the arms or legs trouble thinking, speaking, or walking weakness Incidence not known Chest pain, discomfort, or tightness confusion continuing or severe abdominal or stomach pain continuing or severe nausea and vomiting increased frequency of urination lower back or side pain

Drug Side Effects Summary

pain or discomfort in the arms, jaw, back, or neck stroke sweating trouble breathing vomiting
Symptoms of overdose Blurred vision change or loss of consciousness cold clammy skin confusion
as to time, place, or person decreased awareness or responsiveness dizziness, faintness, or
lightheadedness when getting up suddenly from a lying or sitting position hallucinations headache
holding false beliefs that cannot be changed by fact nausea nervousness numbness, tingling, pain,
and bluish color of the fingernails, lips, skin, palms, or nail beds pale or blue lips, fingernails, or skin
pounding in the ears seizures severe sleepiness slow or fast heartbeat stomach pain sweating
unusual excitement, nervousness, or restlessness unusual tiredness or weakness vomiting Some
side effects may occur that usually do not need medical attention. These side effects may go away
during treatment as your body adjusts to the medicine. Also, your health care professional may be
able to tell you about ways to prevent or reduce some of these side effects. Check with your health
care professional if any of the following side effects continue or are bothersome or if you have any
questions about them: More common Burning or tingling sensation, dryness, soreness, or pain in the
nose Other side effects not listed may also occur in some patients. If you notice any other effects,
check with your healthcare professional. Call your doctor for medical advice about side effects. You
may report side effects to the FDA at 1-800-FDA-1088. D.H.E. 45 side effects

2119. This particular disease, Migraine, interacts with the drug Ergomar and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Ergomar may cause serious side effects. Call your doctor at once if you have: numbness, tingling, or a pale appearance in your fingers or toes
- muscle pain in your arms or legs
- leg weakness
- fast or slow heartbeats

Drug Side Effects Summary

- chest pain, shortness of breath
- or discoloration or skin changes in your hands or feet. Common side effects of Ergomar may include: numbness, tingling, weakness
- dizziness , spinning sensation
- nausea , vomiting
- or swelling or itching in any part of your body.

2120. This particular disease, Migraine, interacts with the drug ergotamine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Ergotamine may cause serious side effects. Call your doctor at once if you have: numbness, tingling, or a pale appearance in your fingers or toes
- muscle pain in your arms or legs
- leg weakness
- fast or slow heartbeats
- chest pain, shortness of breath
- or discoloration or skin changes in your hands or feet. Common side effects of ergotamine may include: numbness, tingling, weakness
- dizziness , spinning sensation
- nausea , vomiting
- or swelling or itching in any part of your body.

2121. This particular disease, Migraine, interacts with the drug methylergonovine and has the following side effects:

- hives
- difficulty breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Methylergonovine may cause serious side effects. Call your doctor at once if you have: increased blood pressure (severe headache, blurred vision, pounding in your neck or ears)
- chest pain, sweating, pounding heartbeats or fluttering in your chest
- a seizure
- numbness, tingling, or cold feeling in your fingers or toes
- a light-headed feeling, like you might pass out
- or sudden numbness or weakness on one side of the body, problems with vision or speech, pain or swelling in one leg. Common side effects of methylergonovine may include: increased blood pressure (severe headache, blurred vision, pounding in your neck or ears)
- a seizure
- or headache.

2122. This particular disease, Migraine, interacts with the drug Onzetra Xsail and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Chest pain (mild) difficulty with swallowing heaviness, tightness, or pressure in the chest or neck pounding heartbeat skin rash, hives, itching, or bumps on the skin Rare Changes in skin color of the face chest pain (severe) fast or irregular breathing puffiness or swelling of the eyelids, area around the eyes, face, or lips seizures troubled breathing Incidence not known Blurred vision confusion dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position fast, irregular, pounding, or racing heartbeat or pulse inability to move the eyes increased blinking or spasms of the eyelid sticking out of the tongue sweating tremor trouble in speaking uncontrolled twisting movements of the neck, trunk, arms, or legs unusual facial expressions unusual tiredness or

Drug Side Effects Summary

weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Burning, discharge, pain, or soreness in the nose change in taste discomfort in the jaw, mouth, tongue, throat, nose, or sinuses dizziness drowsiness feeling cold, "strange," or weak feeling of burning, warmth, heat, numbness, tightness, or tingling flushing lightheadedness muscle aches, cramps, or stiffness nausea redness of the face, neck, arms and occasionally, upper chest unusual or unpleasant (after) taste vomiting Less common Feeling of constant movement of self or surroundings sensation of spinning Less common or rare Anxiety burning sensation discomfort of the nasal cavity and throat general feeling of illness or tiredness vision changes Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Onzetra Xsail side effects

2123. This particular disease, Migraine, interacts with the drug Painaid and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing

Drug Side Effects Summary

- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of bleeding like throwing up or coughing up blood
- vomit that looks like coffee grounds
- blood in the urine
- black, red, or tarry stools
- bleeding from the gums
- abnormal vaginal bleeding
- bruises without a cause or that get bigger
- or bleeding you cannot stop. A heartbeat that does not feel normal. Fever or chills. Very bad dizziness or passing out. Feeling confused. Feeling very tired or weak. Very bad belly pain. Ringing in ears. Hearing loss . Fast breathing. Sweating a lot. Very bad headache or if headache is not better after the first dose. Pain for more than 10 days. Fever for more than 3 days. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Painful side effects

2124. This particular disease, Migraine, interacts with the drug Vanquish and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if

Drug Side Effects Summary

you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of bleeding like throwing up or coughing up blood
- vomit that looks like coffee grounds
- blood in the urine
- black, red, or tarry stools
- bleeding from the gums
- abnormal vaginal bleeding
- bruises without a cause or that get bigger
- or bleeding you cannot stop. Feeling confused. Feeling very tired or weak. Very bad dizziness or passing out. Ringing in ears. Hearing loss . Very bad headache or if headache is not better after the first dose. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes

Drug Side Effects Summary

- or sores in your mouth, throat, nose, or eyes. Very bad stomach ulcers or bleeding can happen with Vanquish (acetaminophen, aspirin, and caffeine tablets). Taking it in high doses or for a long time, smoking , or drinking alcohol raises the chance of these side effects. Taking Vanquish (acetaminophen, aspirin, and caffeine tablets) with food will not lower the chance of these effects. Call your doctor or get medical help right away if you get very bad stomach or back pain
- black, tarry, or bloody stools
- throwing up blood or throw up that looks like coffee grounds
- or weight gain or swelling that is not normal. Vanquish side effects

2125. This particular disease, Migraine, interacts with the drug Zomig Nasal Spray and has the following side effects:

- No side effects listed

2126. This particular disease, Migraine, interacts with the drug Anacin Advanced Headache Formula and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe anxiety , agitation , confusion, panic
- easy bruising or bleeding
- a light-headed feeling, like you might pass out
- dehydration symptoms--feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot

Drug Side Effects Summary

and dry skin

- symptoms of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- high potassium--nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- or liver problems--nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Anacin Advanced Headache Formula may include: upset stomach , heartburn
- depressed mood, feeling anxious or restless
- or sleep problems (insomnia).

2127. This particular disease, Migraine, interacts with the drug Backaid Inflammatory Pain Formula

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe anxiety , agitation , confusion, panic
- easy bruising or bleeding
- a light-headed feeling, like you might pass out
- dehydration symptoms--feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin

Drug Side Effects Summary

- symptoms of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- high potassium--nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- or liver problems--nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Backaid IPF may include: upset stomach , heartburn
- depressed mood, feeling anxious or restless
- or sleep problems (insomnia).

2128. This particular disease, Migraine, interacts with the drug Fortabs and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Diarrhea indigestion loss of appetite nausea or vomiting passing gas stomach pain, fullness, or discomfort Incidence not known Blistering, peeling, or loosening of the skin chest pain chills cough or hoarseness fever itching joint or muscle pain lower back or side pain painful or difficult urination red irritated eyes red skin lesions, often with a purple center sores, ulcers, or white spots on the lips or in the mouth swollen glands trouble breathing unusual bleeding or bruising unusual tiredness or weakness Symptoms of overdose Bleeding under the skin blurred vision change in consciousness confusion confusion as to time, place, or person continuing ringing or buzzing or other unexplained noise in the ears difficult or labored breathing dizziness, faintness, or lightheadedness when getting up suddenly from a lying position extremely high fever or body temperature hallucinations irregular, fast, slow, or shallow breathing fast, weak heartbeat headache hearing loss holding false beliefs that cannot be changed by fact loss of consciousness muscle cramps pale, clammy skin pale or blue

Drug Side Effects Summary

lips, fingernails, or skin shakiness in the legs, arms, hands, or feet stomach pain sweating thirst trembling or shaking of the hands or feet trouble sleeping unusual bleeding or bruising unusual excitement, nervousness, or restlessness unusual tiredness or weakness vomiting Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Dizziness drowsiness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Fortabs side effects

2129. This particular disease, Migraine, interacts with the drug Genace and has the following side effects

- If you experience any of the following serious side effects, stop taking Genace and seek emergency medical attention or contact your doctor immediately: an allergic reaction (difficulty breathing
 - closing of the throat
 - swelling of the lips, tongue, or face
 - or hives)
- liver damage (yellowing of the skin or eyes, nausea, abdominal pain or discomfort, unusual bleeding or bruising, severe fatigue)
- or blood problems (easy or unusual bleeding or bruising). black, bloody, or tarry stools
- nausea, vomiting, or abdominal pain
- uncontrolled fever
- blood in the urine or vomit

Drug Side Effects Summary

- decreased hearing or ringing in the ears
- seizures
- or dizziness, confusion, or hallucinations. These side effects are very rare and are not likely to occur with the proper use of Genace. If you experience any unusual side effects, stop taking Genace and seek the advice of your doctor. If you experience any of the following less serious side effects, notify your doctor: heartburn or indigestion
- or nervousness, anxiety, or difficulty sleeping. Side effects other than those listed here may also occur. Talk to your doctor about any side effect that seems unusual or that is especially bothersome.

Genace side effects

2130. This particular disease, Migraine, interacts with the drug Goody's Extra-Strength Headache P

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of bleeding like throwing up or coughing up blood

Drug Side Effects Summary

- vomit that looks like coffee grounds
- blood in the urine
- black, red, or tarry stools
- bleeding from the gums
- abnormal vaginal bleeding
- bruises without a cause or that get bigger
- or bleeding you cannot stop. Feeling confused. Feeling very tired or weak. Very bad dizziness or passing out. Ringing in ears. Hearing loss . Very bad headache or if headache is not better after the first dose. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Very bad stomach ulcers or bleeding can happen with this medicine (Goody's Extra-Strength Headache Powders). Taking it in high doses or for a long time, smoking , or drinking alcohol raises the chance of these side effects. Taking this medicine (Goody's Extra-Strength Headache Powders) with food will not lower the chance of these effects. Call your doctor or get medical help right away if you get very bad stomach or back pain
- black, tarry, or bloody stools
- throwing up blood or throw up that looks like coffee grounds
- or weight gain or swelling that is not normal. Goody's Extra-Strength Headache Powders side effects

2131. This particular disease, Migraine, interacts with the drug Goody's Extra Strength and has the

- hives

Drug Side Effects Summary

- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe anxiety , agitation , confusion, panic
- easy bruising or bleeding
- a light-headed feeling, like you might pass out
- dehydration symptoms--feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin
- symptoms of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- high potassium--nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- or liver problems--nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Goody's Extra Strength may include: upset stomach , heartburn
- depressed mood, feeling anxious or restless
- or sleep problems (insomnia).

2132. This particular disease, Migraine, interacts with the drug Goody's Headache Powders and has

- hives
- difficulty breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains acetaminophen. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe anxiety , agitation , confusion, panic
- easy bruising or bleeding
- a light-headed feeling, like you might pass out
- dehydration symptoms--feeling very thirsty or hot, being unable to urinate, heavy sweating, or hot and dry skin
- symptoms of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- high potassium--nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- or liver problems--nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Goody's Headache Powders may include: upset stomach , heartburn
- depressed mood, feeling anxious or restless
- or sleep problems (insomnia).

2133. This particular disease, Migraine, interacts with the drug Migraine Relief and has the following

- No side effects listed

2134. This particular disease, Migraine, interacts with the drug Trudhesa and has the following side

Drug Side Effects Summary

- Trudhesa can cause serious side effects, including: See Important information . Heart attack and other heart problems. Heart problems may lead to death. Stop treatment and get emergency medical help right away if you have any of the following symptoms of a heart attack: discomfort in the center of your chest that lasts for more than a few minutes, or that goes away and comes back severe tightness, pain, pressure, or heaviness in your chest, throat, neck, or jaw pain or discomfort in your arms, back, neck, jaw, or stomach shortness of breath with or without chest discomfort breaking out in a cold sweat nausea or vomiting feeling lightheaded Trudhesa is not for people with risk factors for heart disease unless a heart exam is done and shows no problem. You have a higher risk for heart disease if you: have high blood pressure have high cholesterol levels smoke are overweight have diabetes have a family history of heart disease Stroke. Stop treatment and get emergency medical help right away if you have any of the following symptoms of a stroke: face drooping unusual weakness or numbness slurred speech Changes in color or sensation in your fingers and toes (Raynaud's syndrome). Stomach and intestinal problems (gastrointestinal and colonic ischemic events). Symptoms of gastrointestinal and colonic ischemic events include: sudden or severe stomach pain constipation or diarrhea stomach pain after meals bloody diarrhea weight loss fever nausea or vomiting Increase blood pressure. Medicine overuse headache. Some people who use too much Trudhesa may make their headaches worse (medicine overuse headache). If your headaches get worse, your healthcare provider may decide to stop your treatment. Preterm labor. Tissue changes (fibrotic complications). Inflammation and fiber-like tissue that is not normal (fibrosis) can occur around the lungs and stomach. Burning feelings in your nose, mouth and throat and abnormal taste. The most common side effects include: runny nose application site reactions sleepiness nausea dizziness sore throat abnormal taste vomiting diarrhea These are not all the possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Trudhesa side effects

Drug Side Effects Summary

2135. This particular disease, Migraine, interacts with the drug ubrogepant and has the following side effects:

- The most common side effects of ubrogepant are nausea and sleepiness. These are not all of the possible side effects of this medicine. Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Ubrogapant side effects

2136. This particular disease, Osteoarthritis, interacts with the drug meloxicam and has the following side effects:

- sneezing, runny or stuffy nose
- wheezing or trouble breathing
- hives
- swelling of your face, lips, tongue, or throat. Get emergency medical help if you have signs of a heart attack or stroke: chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Meloxicam may cause serious side effects. Stop using this medicine and call your doctor at once if you have: the first sign of any skin rash, no matter how mild
- shortness of breath (even with mild exertion)
- swelling or rapid weight gain
- signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems - nausea, upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- kidney problems - little or no urinating, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- low red blood cells (anemia) - pale skin, feeling light-headed or short of breath, rapid heart rate, trouble concentrating
- or severe skin reaction - fever, sore throat, swelling in your face or tongue, burning in your eyes,

Drug Side Effects Summary

skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of meloxicam may include: upset stomach, nausea, vomiting, heartburn

- diarrhea, constipation, gas
- dizziness
- or cold or flu symptoms.

2137. This particular disease, Osteoarthritis, interacts with the drug Celebrex and has the following

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling , feeling short of breath. Stop using Celebrex and seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Stop using Celebrex and call your doctor at once if you have: the first sign of any skin rash, no matter how mild

- heart problems - swelling, rapid weight gain, feeling short of breath

- signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds

- liver problems - nausea , stomach pain (upper right side), itching, tiredness, dark urine, jaundice (yellowing of the skin or eyes)

- kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath

- or low red blood cells (anemia) - pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Common Celebrex side effects may include: stomach pain, heartburn , gas, diarrhea , constipation , nausea, vomiting

- swelling in your hands or feet

Drug Side Effects Summary

- dizziness
- or cold symptoms such as stuffy nose, sneezing, sore throat.

2138. This particular disease, Osteoarthritis, interacts with the drug Mobic and has the following side effects:

- sneezing, runny or stuffy nose
- wheezing or trouble breathing
- hives
- swelling of your face, lips, tongue, or throat. Get emergency medical help if you have signs of a heart attack or stroke: chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Stop using Mobic and call your doctor at once if you have: the first sign of any skin rash, no matter how mild
- shortness of breath (even with mild exertion)
- swelling or rapid weight gain
- signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- signs of liver problems - nausea, upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- signs of kidney problems - little or no urinating, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- low red blood cells (anemia) - pale skin, feeling light-headed or short of breath, rapid heart rate, trouble concentrating
- or signs of a severe skin reaction - fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common Mobic side effects may include: upset stomach, nausea, vomiting, heartburn

Drug Side Effects Summary

- diarrhea, constipation, gas
- dizziness
- or cold symptoms, flu symptoms.

2139. This particular disease, Osteoarthritis, interacts with the drug naproxen and has the following

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Stop using naproxen and call your doctor at once if you have: shortness of breath (even with mild exertion)
- swelling or rapid weight gain
- the first sign of any skin rash, no matter how mild
- signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems - nausea, upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- kidney problems - little or no urinating, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- low red blood cells (anemia) - pale skin, feeling light-headed or short of breath, rapid heart rate, trouble concentrating
- or severe skin reaction - fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common naproxen side effects may include: indigestion, heartburn, stomach pain, nausea
- headache, dizziness, drowsiness
- bruising, itching, rash
- swelling

Drug Side Effects Summary

- or ringing in your ears.

2140. This particular disease, Osteoarthritis, interacts with the drug ibuprofen and has the following

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. Stop using this medicine and call your doctor at once if you have: changes in your vision
- shortness of breath (even with mild exertion)
- swelling or rapid weight gain
- a skin rash, no matter how mild
- signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems - nausea, upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- low red blood cells (anemia) - pale skin, feeling light-headed or short of breath, rapid heart rate, trouble concentrating
- or kidney problems - little or no urinating, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath. Common ibuprofen side effects may include: nausea, vomiting, gas
- bleeding
- or dizziness , headache.

2141. This particular disease, Osteoarthritis, interacts with the drug Voltaren and has the following

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Stop using this medicine and call your doctor at once if you have: the first sign of any skin rash, no matter how mild

Drug Side Effects Summary

- flu-like symptoms
- heart problems - swelling, rapid weight gain, feeling short of breath
- kidney problems - little or no urinating, painful or difficult urination, swelling in your arms or legs, feeling tired or short of breath
- liver problems - nausea , diarrhea , stomach pain (upper right side), tiredness, itching, dark urine, jaundice (yellowing of the skin or eyes)
- or signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Common Voltaren side effects may include: indigestion , gas, nausea, vomiting , stomach pain
- diarrhea, constipation
- headache, dizziness , drowsiness
- abnormal lab tests
- itching, sweating
- stuffy nose
- increased blood pressure
- or swelling or pain in your arms or legs.

2142. This particular disease, Osteoarthritis, interacts with the drug chondroitin / glucosamine and

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Although not all side effects are known, chondroitin and glucosamine is thought to be possibly safe when taken for up to 2 years. Stop using chondroitin and glucosamine and call your healthcare provider at once if you have: irregular heartbeats
- or swelling in your legs. Common side effects of chondroitin and glucosamine may include: nausea , diarrhea , constipation

Drug Side Effects Summary

- stomach pain, gas, bloating
- hair loss
- or puffy eyelids.

2143. This particular disease, Osteoarthritis, interacts with the drug Voltaren Arthritis Pain Gel and

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Also call your doctor at once if you have: a skin rash, no matter how mild
- swelling, rapid weight gain
- severe headache, blurred vision, pounding in your neck or ears
- little or no urination
- liver problems - nausea, diarrhea, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- low red blood cells (anemia) - pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet
- or signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Common Voltaren Arthritis Pain gel side effects may include: heartburn, gas, stomach pain, nausea, vomiting
- diarrhea, constipation
- headache, dizziness, drowsiness
- stuffy nose
- itching, increased sweating
- increased blood pressure
- or skin redness, itching, dryness, scaling, or peeling where the medicine was applied.

Drug Side Effects Summary

2144. This particular disease, Osteoarthritis, interacts with the drug diclofenac and has the following side effects:

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Also call your doctor at once if you have: a skin rash, no matter how mild
- swelling, rapid weight gain
- severe headache , blurred vision, pounding in your neck or ears
- little or no urination
- liver problems-- nausea , diarrhea , stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet
- or signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Common side effects of diclofenac topical may include: heartburn , gas, stomach pain, nausea, vomiting
- diarrhea, constipation
- headache, dizziness , drowsiness
- stuffy nose
- itching, increased sweating
- increased blood pressure
- or skin redness, itching, dryness, scaling, or peeling where the medicine was applied.

2145. This particular disease, Osteoarthritis, interacts with the drug Aleve and has the following side effects:

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Stop using Aleve and call your doctor at once if you have: shortness of breath (even with mild exertion)

Drug Side Effects Summary

- swelling or rapid weight gain
- the first sign of any skin rash, no matter how mild
- signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems - nausea, upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- kidney problems - little or no urinating, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- low red blood cells (anemia) - pale skin, feeling light-headed or short of breath, rapid heart rate, trouble concentrating
- or severe skin reaction - fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common Aleve side effects may include: indigestion, heartburn, stomach pain, nausea
- headache, dizziness, drowsiness
- bruising, itching, rash
- swelling
- or ringing in your ears.

2146. This particular disease, Osteoarthritis, interacts with the drug Arthrotec and has the following

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Stop using Arthrotec and call your doctor at once if you have: the first sign of any skin rash, no matter how mild
- shortness of breath (even with mild exertion)
- swelling or rapid weight gain

Drug Side Effects Summary

- signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems - loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or low red blood cells (anemia) - pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Common Arthrotec side effects may include: abnormal vaginal bleeding
- heartburn , indigestion stomach pain, gas
- nausea , vomiting
- diarrhea , constipation
- or dizziness .

2147. This particular disease, Osteoarthritis, interacts with the drug Cataflam and has the following

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Stop using Cataflam and call your doctor at once if you have: the first sign of any skin rash, no matter how mild
- flu-like symptoms
- heart problems - swelling, rapid weight gain, feeling short of breath
- kidney problems - little or no urinating, painful or difficult urination, swelling in your arms or legs, feeling tired or short of breath
- liver problems - nausea , diarrhea , stomach pain (upper right side), tiredness, itching, dark urine, jaundice (yellowing of the skin or eyes)
- or signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Common Cataflam side effects may include: indigestion , gas, nausea, vomiting ,

Drug Side Effects Summary

stomach pain

- diarrhea, constipation
- headache, dizziness , drowsiness
- abnormal lab tests
- itching, sweating
- stuffy nose
- increased blood pressure
- or swelling or pain in your arms or legs.

2148. This particular disease, Osteoarthritis, interacts with the drug nabumetone and has the following

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Nabumetone may cause serious side effects. Stop using nabumetone and call your doctor at once if you have: shortness of breath (even with mild exertion)
- swelling or rapid weight gain
- the first sign of any skin rash, no matter how mild
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems-- nausea , upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- kidney problems--little or no urinating, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- or low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Common side effects of nabumetone may include: stomach pain, indigestion , nausea

Drug Side Effects Summary

- diarrhea , constipation , gas
- swelling in your hands and feet
- headache , dizziness
- itching, skin rash
- or ringing in your ears.

2149. This particular disease, Osteoarthritis, interacts with the drug etodolac and has the following

- hives , runny or stuffy nose, wheezing, difficult breathing, swelling in your face or throat) or a severe skin reaction (fever, sore throat , burning eyes, skin pain, red or purple skin rash with blistering and peeling). Get emergency medical help if you have signs of a heart attack or stroke: chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, swelling in your legs, feeling short of breath. Stop using etodolac and call your doctor at once if you have: changes in your vision
- any skin rash, no matter how mild
- shortness of breath (even with mild exertion)
- swelling or rapid weight gain
- signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems - nausea , upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- kidney problems - little or no urination, swelling in your feet or ankles, feeling tired
- or low red blood cells (anemia) - pale skin, unusual tiredness, feeling light-headed, cold hands and feet. Common etodolac side effects include: nausea, vomiting, stomach pain, indigestion
- diarrhea , constipation , gas
- dizziness , headache , tired feeling

Drug Side Effects Summary

- rash
- or ringing in your ears.

2150. This particular disease, Osteoarthritis, interacts with the drug Advil and has the following side effects:

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. Stop using Advil and call your doctor at once if you have: changes in your vision
- shortness of breath (even with mild exertion)
- swelling or rapid weight gain
- a skin rash, no matter how mild
- signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems - nausea, upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- low red blood cells (anemia) - pale skin, feeling light-headed or short of breath, rapid heart rate, trouble concentrating
- or kidney problems - little or no urinating, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath. Common Advil side effects may include: nausea, vomiting, gas
- bleeding
- or dizziness , headache.

2151. This particular disease, Osteoarthritis, interacts with the drug Motrin and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your

Drug Side Effects Summary

doctor immediately if any of the following side effects occur: More common Abdominal pain acid or sour stomach belching bloating cloudy urine decrease in amount of urine decrease in urine output or decrease in urine-concentrating ability diarrhea difficulty having a bowel movement (stool) excess air or gas in stomach or intestines full feeling heartburn indigestion itching skin pain or discomfort in chest, upper stomach, or throat pale skin passing gas nausea noisy, rattling breathing rash with flat lesions or small raised lesions on the skin shortness of breath swelling of face, fingers, hands, feet, lower legs, or ankles troubled breathing at rest troubled breathing with exertion unusual bleeding or bruising unusual tiredness or weakness vomiting weight gain Less common Abdominal cramps stomach soreness or discomfort Rare Agitation back, leg, or stomach pains bleeding gums blistering, peeling, loosening of skin blood in urine or stools bloody, black, or tarry stools blurred vision burning feeling in chest or stomach change in vision chest pain chills clay-colored stools coma confusion constipation cough or hoarseness dark urine decreased urine output depression difficulty breathing difficulty swallowing dilated neck veins dizziness dry mouth extreme fatigue fast, irregular, pounding, or racing heartbeat or pulse fever with or without chills frequent urination general body swelling general feeling of tiredness or weakness hair loss, thinning of hair headache hives or welts hostility impaired vision increased blood pressure increased volume of pale, dilute urine irregular breathing irritability itching joint or muscle pain lab results that show problems with liver lethargy light-colored stools loss of appetite lower back or side pain muscle twitching nosebleeds painful or difficult urination pains in stomach, side, or abdomen, possibly radiating to the back pinpoint red spots on skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rash red skin lesions, often with a purple center red, irritated eyes redness of skin seizures severe abdominal pain, cramping, burning severe and continuing nausea sore throat sores, ulcers, or white spots in mouth or on lips stiff neck or back stomach upset stupor swollen or painful glands tenderness in stomach area thirst tightness in chest unpleasant breath odor upper right abdominal pain vomiting of blood vomiting of material that looks like coffee grounds wheezing yellow eyes and skin Symptoms

Drug Side Effects Summary

of overdose Bluish lips or skin difficulty sleeping disorientation dizziness, faintness, or lightheadedness when getting up from a lying or sitting position suddenly drowsiness to profound coma hallucination lightheadedness or fainting mood or other mental changes muscle tremors not breathing rapid, deep breathing restlessness slow or irregular heartbeat stomach cramps sudden fainting sweating Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Continuing ringing or buzzing or other unexplained noise in ears hearing loss nervousness Rare Crying depersonalization discouragement dry eyes dysphoria euphoria feeling sad or empty lack of appetite loss of interest or pleasure mental depression paranoia quick to react or overreact rapidly changing moods runny nose sleepiness or unusual drowsiness sleeplessness sneezing stuffy nose trouble concentrating trouble sleeping unable to sleep Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Motrin side effects

2152. This particular disease, Osteoarthritis, interacts with the drug Vimovo and has the following s

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- heart problems - swelling, rapid weight gain, feeling short of breath
- kidney problems - fever, rash, nausea, loss of appetite, joint pain, urinating less than usual, blood in your urine, weight gain

Drug Side Effects Summary

- liver problems - loss of appetite, stomach pain (upper right side), dark urine, jaundice (yellowing of the skin or eyes)
- low magnesium - dizziness , fast or irregular heart rate, tremors (shaking) or jerking muscle movements, feeling jittery, muscle cramps, muscle spasms in your hands and feet, cough or choking feeling
- low red blood cells (anemia) - pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet
- or new or worsening symptoms of lupus - joint pain, and a skin rash on your cheeks or arms that worsens in sunlight. Taking Vimovo long-term may cause you to develop stomach growths called fundic gland polyps. Talk with your doctor about this risk. If you use Vimovo for longer than 2 years, you could develop a vitamin B-12 deficiency. Talk to your doctor about how to manage this condition if you develop it. Common Vimovo side effects may include: stomach discomfort
- or diarrhea.

2153. This particular disease, Osteoarthritis, interacts with the drug Naprosyn and has the following

- chest pain, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Stop using Naprosyn and call your doctor at once if you have: shortness of breath (even with mild exertion)
- swelling or rapid weight gain
- the first sign of any skin rash, no matter how mild
- signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems - nausea, upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- kidney problems - little or no urinating, painful or difficult urination, swelling in your feet or ankles,

Drug Side Effects Summary

feeling tired or short of breath

- low red blood cells (anemia) - fatigue, pale skin, feeling light-headed or short of breath, rapid heart rate, trouble concentrating
- or severe skin reaction - fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common Naprosyn side effects may include: indigestion, heartburn, stomach pain, nausea
- headache, dizziness, drowsiness
- bruising, itching, rash
- swelling
- or ringing in your ears.

2154. This particular disease, Osteoarthritis, interacts with the drug Pennsaid and has the following

- sneezing, runny or stuffy nose
- wheezing or trouble breathing
- hives
- swelling of your face, lips, tongue, or throat. Get emergency medical help if you have signs of a heart attack or stroke: chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Stop using Pennsaid and seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Stop using Pennsaid and call your doctor at once if you have: a skin rash, no matter how mild
- shortness of breath (even with mild exertion)
- swelling or rapid weight gain

Drug Side Effects Summary

- signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- Inausea, upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- little or no urinating, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- severe headache, pounding in your neck or ears, nosebleed, anxiety, confusion
- pale skin, feeling light-headed or short of breath, rapid heart rate, trouble concentrating
- or fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common Pennsaid side effects may include: indigestion, gas, stomach pain, nausea, vomiting
- diarrhea, constipation
- headache, dizziness, drowsiness
- stuffy nose
- itching, increased sweating
- increased blood pressure
- or swelling or pain in your arms or legs.

2155. This particular disease, Osteoarthritis, interacts with the drug Zorvolex and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Acid or sour stomach belching bleeding gums blood in the urine or stools bloody or black, tarry stools burning while urinating chest pain chills cough clay-colored stools cloudy urine constipation dark urine decrease in

Drug Side Effects Summary

urine output or decrease in urine-concentrating ability diarrhea dizziness feeling of indigestion fever frequent urge to urinate headache heartburn increased bleeding time indigestion itching skin or rash light-colored stools loss of appetite nausea and vomiting pain in the chest below the breastbone painful or difficult urination pale skin pinpoint red spots on the skin stomach bloating, burning, cramping, discomfort, upset, or pain sore throat sores, ulcers, or white spots on the lips or in the mouth swelling or inflammation of the mouth swollen glands trouble breathing unpleasant breath odor unusual bleeding or bruising unusual tiredness or weakness upper right abdominal or stomach pain vomiting of blood or material that looks like coffee grounds weight loss yellow eyes and skin

Less common Blistering, peeling, loosening of the skin blurred vision burning, crawling, itching, numbness, prickling, "pins and needles" , or tingling feelings confusion change in consciousness discouragement dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position feeling of constant movement of self or surroundings feeling sad or empty irritability joint or muscle pain lack of appetite lack or loss of strength loss of consciousness loss of interest or pleasure nervousness pain or discomfort in the arms, jaw, back, or neck pale or blue lips, fingernails, or skin red irritated eyes red skin lesions, often with a purple center redness, soreness, or itching skin sensation of spinning shakiness in the legs, arms, hands, or feet sleepiness or unusual drowsiness stiff neck or back trembling or shaking of the hands or feet trouble concentrating trouble sleeping

Symptoms of overdose Agitation blurred vision change in consciousness change in the ability to see colors, especially blue or yellow confusion depression difficult or trouble breathing hives hostility irregular, fast or slow, or shallow breathing irritability loss of consciousness muscle twitching nervousness pain or discomfort in the chest, upper stomach, or throat pale or blue lips, fingernails, or skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rapid weight gain seizures sleepiness slow or fast heartbeat stupor swelling of the face, ankles, or hands tightness in the chest trouble sleeping unusual drowsiness, dullness, or feeling of sluggishness

Some side effects may occur that usually do not need medical attention. These side effects may go

Drug Side Effects Summary

away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Bloated continuing ringing or buzzing or other unexplained noise in the ears difficulty in swallowing excess air or gas in the stomach or intestines hearing loss lack or loss of strength pain or burning in the throat passing gas Less common Burning, dry or itching eyes discharge, excessive tearing hair loss, thinning of the hair increased sensitivity of the skin to sunlight pain in the arms or legs redness, pain, swelling of the eye, eyelid, or inner lining of eyelid redness or other discoloration of the skin severe sunburn Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Zorvolex side effects

2156. This particular disease, Osteoarthritis, interacts with the drug Indocin and has the following s

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Acid or sour stomach belching diarrhea heartburn indigestion nausea stomach discomfort, upset, or pain vomiting Rare Abdominal or stomach cramping, burning, or tenderness back or leg pains bleeding gums blistering, peeling, or loosening of the skin bloody or black, tarry stools blue lips and fingernails blurred vision breast enlargement and tenderness burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings burning upper abdominal or stomach pain canker sores change in consciousness change in hearing chest pain, discomfort, or burning clay colored stools cloudy or bloody urine confusion continuing diarrhea cough or hoarseness coughing that sometimes produces a pink frothy sputum cracks in the skin dark urine decreased appetite decreased vision or any

Drug Side Effects Summary

change in vision depression difficult or labored breathing difficulty with swallowing dilated neck veins dizziness, faintness, or lightheadedness when getting up from a lying or sitting position double vision dry mouth extreme fatigue false sense of well-being feeling of unreality feeling of warmth fever with or without chills flushed, dry skin fruit-like breath odor general body swelling greatly decreased frequency of urination or amount of urine hair loss headache heavier menstrual periods increased hunger increased sweating increased thirst increased urination irregular breathing irritation and swelling of the eye jerky movements of the head, face, mouth, and neck joint pain large, flat, blue or purplish patches in the skin large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs loss of appetite loss of balance control loss of bladder control loss of consciousness loss of hearing loss of heat from the body lower back or side pain mask-like face mood swings muscle aches, pains, or weakness muscle spasm or jerking of all extremities nervousness noisy, rattling breathing nosebleeds numbness or tingling in the hands, feet, or lips pain in the ankles or knees pain or discomfort in the upper stomach or throat pain with swallowing painful or difficult urination painful, red lumps under the skin, mostly on the legs pale skin persistent bleeding or oozing from puncture sites, mouth, or nose personality changes pinpoint red or purple spots on the skin pounding in the ears puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red skin lesions, often with a purple center red, irritated eyes red, swollen skin redness of the face, neck, arms and occasionally, upper chest scaly skin seeing double seeing, hearing, or feeling things that are not there seizures sense of detachment from self or body severe constipation severe mental changes severe or continuing stomach pain shuffling walk skin rash, hives or welts, itching slow, fast, irregular, pounding, or racing heartbeat or pulse slowed movements slurred speech small red or purple spots on the skin sore throat sores, ulcers, or white spots on the lips or tongue or inside the mouth stiffness of the arms and legs sudden loss of consciousness swelling of the breasts or breast soreness in both females and males swelling of the face, fingers, feet, ankles, or lower legs swollen or painful glands tightness in the chest trembling

Drug Side Effects Summary

and shaking of the fingers and hands troubled breathing at rest troubled breathing with exertion unexplained weight loss unpleasant breath odor unsteadiness or awkwardness unusual bleeding or bruising unusual tiredness or weakness vaginal bleeding vomiting of blood or material that looks like coffee grounds weakness in the arms, hands, legs, or feet weight gain yellow eyes or skin

Symptoms of overdose Confusion about identity, place, and time severe headache unusual drowsiness, dullness, or feeling of sluggishness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Mild headache Less common Continuing ringing or buzzing or other unexplained noise in the ears difficulty having a bowel movement (stool) discouragement feeling sad or empty general feeling of discomfort or illness hearing loss irritability loss of interest or pleasure sleepiness trouble with concentrating Rare Anxiety bloated or full feeling changes in patterns and rhythms of speech excess air or gas in the stomach or intestines feeling of constant movement of self or surroundings involuntary muscle movements lightheadedness passing gas sensation of spinning tiredness trouble sleeping trouble with speaking Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Indocin side effects

2157. This particular disease, Osteoarthritis, interacts with the drug indomethacin and has the follo

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. Stop using indomethacin and seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include skin rash, fever, swollen glands, muscle aches, severe weakness, unusual

Drug Side Effects Summary

bruising, or yellowing of your skin or eyes. Indomethacin may cause serious side effects. Stop using indomethacin and call your doctor at once if you have: a skin rash, no matter how mild

- changes in your vision
- heart problems--swelling, rapid weight gain, feeling short of breath
- signs of liver or pancreas problems--loss of appetite, upper stomach pain (that may spread to your back), nausea , vomiting, diarrhea , unusual tiredness, itching, flu-like symptoms, dark urine, jaundice (yellowing of the skin or eyes)
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Common side effects of indomethacin may include: indigestion , nausea
- headache
- or dizziness .

2158. This particular disease, Osteoarthritis, interacts with the drug Proprinal and has the following

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. Proprinal may cause serious side effects. Stop using Proprinal and call your doctor at once if you have: changes in your vision
- shortness of breath (even with mild exertion)
- swelling or rapid weight gain
- a skin rash, no matter how mild
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds

Drug Side Effects Summary

- liver problems--nausea, upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- low red blood cells (anemia)--pale skin, feeling light-headed or short of breath, rapid heart rate, trouble concentrating
- or kidney problems--little or no urinating, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath. Common side effects of Proprinal may include: nausea, vomiting, gas
- bleeding
- or dizziness , headache.

2159. This particular disease, Osteoarthritis, interacts with the drug A-G Profen and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Abdominal pain acid or sour stomach belching bloating cloudy urine decrease in amount of urine decrease in urine output or decrease in urine-concentrating ability diarrhea difficulty having a bowel movement (stool) excess air or gas in stomach or intestines full feeling heartburn indigestion itching skin pain or discomfort in chest, upper stomach, or throat pale skin passing gas nausea noisy, rattling breathing rash with flat lesions or small raised lesions on the skin shortness of breath swelling of face, fingers, hands, feet, lower legs, or ankles troubled breathing at rest troubled breathing with exertion unusual bleeding or bruising unusual tiredness or weakness vomiting weight gain Less common Abdominal cramps stomach soreness or discomfort Rare Agitation back, leg, or stomach pains bleeding gums blistering, peeling, loosening of skin blood in urine or stools bloody, black, or tarry stools blurred vision burning feeling in chest or stomach change in vision chest pain chills clay-colored stools coma confusion constipation cough or hoarseness dark urine decreased urine output depression difficulty breathing difficulty swallowing dilated neck veins dizziness dry mouth extreme fatigue fast, irregular,

Drug Side Effects Summary

pounding, or racing heartbeat or pulse fever with or without chills frequent urination general body swelling general feeling of tiredness or weakness hair loss, thinning of hair headache hives or welts hostility impaired vision increased blood pressure increased volume of pale, dilute urine irregular breathing irritability itching joint or muscle pain lab results that show problems with liver lethargy light-colored stools loss of appetite lower back or side pain muscle twitching nosebleeds painful or difficult urination pains in stomach, side, or abdomen, possibly radiating to the back pinpoint red spots on skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rash red skin lesions, often with a purple center red, irritated eyes redness of skin seizures severe abdominal pain, cramping, burning severe and continuing nausea sore throat sores, ulcers, or white spots in mouth or on lips stiff neck or back stomach upset stupor swollen or painful glands tenderness in stomach area thirst tightness in chest unpleasant breath odor upper right abdominal pain vomiting of blood vomiting of material that looks like coffee grounds wheezing yellow eyes and skin

Symptoms of overdose Bluish lips or skin difficulty sleeping disorientation dizziness, faintness, or lightheadedness when getting up from a lying or sitting position suddenly drowsiness to profound coma hallucination lightheadedness or fainting mood or other mental changes muscle tremors not breathing rapid, deep breathing restlessness slow or irregular heartbeat stomach cramps sudden fainting sweating

Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them:

More common Continuing ringing or buzzing or other unexplained noise in ears hearing loss nervousness

Rare Crying depersonalization discouragement dry eyes dysphoria euphoria feeling sad or empty lack of appetite loss of interest or pleasure mental depression paranoia quick to react or overreact rapidly changing moods runny nose sleepiness or unusual drowsiness sleeplessness sneezing stuffy nose trouble concentrating trouble

Drug Side Effects Summary

sleeping unable to sleep Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. A-G Profen side effects

2160. This particular disease, Osteoarthritis, interacts with the drug acetaminophen / codeine and h

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Acetaminophen and codeine may cause serious side effects. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops
- a light-headed feeling, like you might pass out
- confusion, severe drowsiness
- liver problems-- nausea , upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or high levels of serotonin in the body-- agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting , diarrhea . Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders. Common side effects include: drowsiness, dizziness, feeling tired
- nausea, vomiting, stomach pain

Drug Side Effects Summary

- constipation
- or headache .

2161. This particular disease, Osteoarthritis, interacts with the drug Actipufen and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Abdominal pain acid or sour stomach belching bloating cloudy urine decrease in amount of urine decrease in urine output or decrease in urine-concentrating ability diarrhea difficulty having a bowel movement (stool) excess air or gas in stomach or intestines full feeling heartburn indigestion itching skin pain or discomfort in chest, upper stomach, or throat pale skin passing gas nausea noisy, rattling breathing rash with flat lesions or small raised lesions on the skin shortness of breath swelling of face, fingers, hands, feet, lower legs, or ankles troubled breathing at rest troubled breathing with exertion unusual bleeding or bruising unusual tiredness or weakness vomiting weight gain Less common Abdominal cramps stomach soreness or discomfort Rare Agitation back, leg, or stomach pains bleeding gums blistering, peeling, loosening of skin blood in urine or stools bloody, black, or tarry stools blurred vision burning feeling in chest or stomach change in vision chest pain chills clay-colored stools coma confusion constipation cough or hoarseness dark urine decreased urine output depression difficulty breathing difficulty swallowing dilated neck veins dizziness dry mouth extreme fatigue fast, irregular, pounding, or racing heartbeat or pulse fever with or without chills frequent urination general body swelling general feeling of tiredness or weakness hair loss, thinning of hair headache hives or welts hostility impaired vision increased blood pressure increased volume of pale, dilute urine irregular breathing irritability itching joint or muscle pain lab results that show problems with liver lethargy light-colored stools loss of appetite lower back or side pain muscle twitching nosebleeds painful or difficult urination pains in stomach, side, or abdomen, possibly radiating to the back pinpoint red

Drug Side Effects Summary

spots on skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rash red skin lesions, often with a purple center red, irritated eyes redness of skin seizures severe abdominal pain, cramping, burning severe and continuing nausea sore throat sores, ulcers, or white spots in mouth or on lips stiff neck or back stomach upset stupor swollen or painful glands tenderness in stomach area thirst tightness in chest unpleasant breath odor upper right abdominal pain vomiting of blood vomiting of material that looks like coffee grounds wheezing yellow eyes and skin Symptoms of overdose Bluish lips or skin difficulty sleeping disorientation dizziness, faintness, or lightheadedness when getting up from a lying or sitting position suddenly drowsiness to profound coma hallucination lightheadedness or fainting mood or other mental changes muscle tremors not breathing rapid, deep breathing restlessness slow or irregular heartbeat stomach cramps sudden fainting sweating Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Continuing ringing or buzzing or other unexplained noise in ears hearing loss nervousness Rare Crying depersonalization discouragement dry eyes dysphoria euphoria feeling sad or empty lack of appetite loss of interest or pleasure mental depression paranoia quick to react or overreact rapidly changing moods runny nose sleepiness or unusual drowsiness sleeplessness sneezing stuffy nose trouble concentrating trouble sleeping unable to sleep Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Actiprofen side effects

2162. This particular disease, Osteoarthritis, interacts with the drug Addaprin and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of

Drug Side Effects Summary

these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Abdominal pain acid or sour stomach belching bloating cloudy urine decrease in amount of urine decrease in urine output or decrease in urine-concentrating ability diarrhea difficulty having a bowel movement (stool) excess air or gas in stomach or intestines full feeling heartburn indigestion itching skin pain or discomfort in chest, upper stomach, or throat pale skin passing gas nausea noisy, rattling breathing rash with flat lesions or small raised lesions on the skin shortness of breath swelling of face, fingers, hands, feet, lower legs, or ankles troubled breathing at rest troubled breathing with exertion unusual bleeding or bruising unusual tiredness or weakness vomiting weight gain Less common Abdominal cramps stomach soreness or discomfort Rare Agitation back, leg, or stomach pains bleeding gums blistering, peeling, loosening of skin blood in urine or stools bloody, black, or tarry stools blurred vision burning feeling in chest or stomach change in vision chest pain chills clay-colored stools coma confusion constipation cough or hoarseness dark urine decreased urine output depression difficulty breathing difficulty swallowing dilated neck veins dizziness dry mouth extreme fatigue fast, irregular, pounding, or racing heartbeat or pulse fever with or without chills frequent urination general body swelling general feeling of tiredness or weakness hair loss, thinning of hair headache hives or welts hostility impaired vision increased blood pressure increased volume of pale, dilute urine irregular breathing irritability itching joint or muscle pain lab results that show problems with liver lethargy light-colored stools loss of appetite lower back or side pain muscle twitching nosebleeds painful or difficult urination pains in stomach, side, or abdomen, possibly radiating to the back pinpoint red spots on skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rash red skin lesions, often with a purple center red, irritated eyes redness of skin seizures severe abdominal pain, cramping, burning severe and continuing nausea sore throat sores, ulcers, or white spots in mouth or on lips stiff neck or back stomach upset stupor swollen or painful glands tenderness in stomach area thirst tightness in chest unpleasant breath odor upper right abdominal pain vomiting of

Drug Side Effects Summary

blood vomiting of material that looks like coffee grounds wheezing yellow eyes and skin Symptoms of overdose Bluish lips or skin difficulty sleeping disorientation dizziness, faintness, or lightheadedness when getting up from a lying or sitting position suddenly drowsiness to profound coma hallucination lightheadedness or fainting mood or other mental changes muscle tremors not breathing rapid, deep breathing restlessness slow or irregular heartbeat stomach cramps sudden fainting sweating Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Continuing ringing or buzzing or other unexplained noise in ears hearing loss nervousness Rare Crying depersonalization discouragement dry eyes dysphoria euphoria feeling sad or empty lack of appetite loss of interest or pleasure mental depression paranoia quick to react or overreact rapidly changing moods runny nose sleepiness or unusual drowsiness sleeplessness sneezing stuffy nose trouble concentrating trouble sleeping unable to sleep Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Addaprin side effects

2163. This particular disease, Osteoarthritis, interacts with the drug Advil Liqui-Gels and has the fo

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching

Drug Side Effects Summary

- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of bleeding like throwing up or coughing up blood
- vomit that looks like coffee grounds
- blood in the urine
- black, red, or tarry stools
- bleeding from the gums
- abnormal vaginal bleeding
- bruises without a cause or that get bigger
- or bleeding you cannot stop. Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain. Signs of high potassium levels like a heartbeat that does not feel normal
- feeling confused
- feeling weak, lightheaded, or dizzy
- feeling like passing out
- numbness or tingling
- or shortness of breath. Signs of high blood pressure like very bad headache or dizziness , passing out, or change in eyesight. Shortness of breath, a big weight gain, or swelling in the arms or legs. Chest pain or pressure or a fast heartbeat . Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight. Feeling very tired or weak. Ringing in ears. Very bad back pain . Change in eyesight. A very bad skin reaction (

Drug Side Effects Summary

Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)

- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Advil Liqui-Gels side effects

2164. This particular disease, Osteoarthritis, interacts with the drug chondroitin / glucosamine / me

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Common side effects of chondroitin, glucosamine, and methylsulfonylmethane may include: gas, bloating , heartburn , stomach cramps, nausea
- headache , drowsiness
- diarrhea , constipation
- swelling in your ankles or feet
- itching, skin irritation
- or puffy eyelids.

2165. This particular disease, Osteoarthritis, interacts with the drug diclofenac / misoprostol and ha

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: the first sign of any skin rash, no matter how mild
- shortness of breath (even with mild exertion)
- swelling or rapid weight gain
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee

Drug Side Effects Summary

grounds

- liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
 - kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
 - or low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Common side effects of diclofenac and misoprostol may include:
- abnormal vaginal bleeding
- heartburn , indigestion stomach pain, gas
 - nausea , vomiting
 - diarrhea , constipation
 - or dizziness .

2166. This particular disease, Osteoarthritis, interacts with the drug Duexis and has the following side effects:

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. Stop using ibuprofen and call your doctor at once if you have: changes in your vision
- the first sign of any skin rash, no matter how mild
- fever, headache , neck stiffness, increased sensitivity to light, nausea, vomiting, confusion, drowsiness
- severe headache, blurred vision, pounding in your neck or ears
- little or no urination
- swelling, rapid weight gain
- liver problems - loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- low red blood cells (anemia) - pale skin, unusual tiredness, feeling light-headed or short of breath,

Drug Side Effects Summary

cold hands and feet

- or signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Some side effects may be more likely in older adults and in people who have severe kidney disease. Common Duexis side effects may include: nausea, stomach pain
- diarrhea , constipation
- or headache.

2167. This particular disease, Osteoarthritis, interacts with the drug Genpril and has the following s

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. Genpril may cause serious side effects. Stop using Genpril and call your doctor at once if you have: changes in your vision
- shortness of breath (even with mild exertion)
- swelling or rapid weight gain
- a skin rash, no matter how mild
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems--nausea, upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- low red blood cells (anemia)--pale skin, feeling light-headed or short of breath, rapid heart rate, trouble concentrating
- or kidney problems--little or no urinating, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath. Common side effects of Genpril may include: nausea, vomiting, gas
- bleeding
- or dizziness , headache.

Drug Side Effects Summary

2168. This particular disease, Osteoarthritis, interacts with the drug IBU-200 and has the following s

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Abdominal pain acid or sour stomach belching bloating cloudy urine decrease in amount of urine decrease in urine output or decrease in urine-concentrating ability diarrhea difficulty having a bowel movement (stool) excess air or gas in stomach or intestines full feeling heartburn indigestion itching skin pain or discomfort in chest, upper stomach, or throat pale skin passing gas nausea noisy, rattling breathing rash with flat lesions or small raised lesions on the skin shortness of breath swelling of face, fingers, hands, feet, lower legs, or ankles troubled breathing at rest troubled breathing with exertion unusual bleeding or bruising unusual tiredness or weakness vomiting weight gain Less common Abdominal cramps stomach soreness or discomfort Rare Agitation back, leg, or stomach pains bleeding gums blistering, peeling, loosening of skin blood in urine or stools bloody, black, or tarry stools blurred vision burning feeling in chest or stomach change in vision chest pain chills clay-colored stools coma confusion constipation cough or hoarseness dark urine decreased urine output depression difficulty breathing difficulty swallowing dilated neck veins dizziness dry mouth extreme fatigue fast, irregular, pounding, or racing heartbeat or pulse fever with or without chills frequent urination general body swelling general feeling of tiredness or weakness hair loss, thinning of hair headache hives or welts hostility impaired vision increased blood pressure increased volume of pale, dilute urine irregular breathing irritability itching joint or muscle pain lab results that show problems with liver lethargy light-colored stools loss of appetite lower back or side pain muscle twitching nosebleeds painful or difficult urination pains in stomach, side, or abdomen, possibly radiating to the back pinpoint red spots on skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rash red skin lesions, often with a purple center red, irritated eyes redness of skin seizures severe abdominal pain, cramping, burning severe and continuing nausea sore throat sores, ulcers, or white spots in

Drug Side Effects Summary

mouth or on lips stiff neck or back stomach upset stupor swollen or painful glands tenderness in stomach area thirst tightness in chest unpleasant breath odor upper right abdominal pain vomiting of blood vomiting of material that looks like coffee grounds wheezing yellow eyes and skin Symptoms of overdose Bluish lips or skin difficulty sleeping disorientation dizziness, faintness, or lightheadedness when getting up from a lying or sitting position suddenly drowsiness to profound coma hallucination lightheadedness or fainting mood or other mental changes muscle tremors not breathing rapid, deep breathing restlessness slow or irregular heartbeat stomach cramps sudden fainting sweating Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Continuing ringing or buzzing or other unexplained noise in ears hearing loss nervousness Rare Crying depersonalization discouragement dry eyes dysphoria euphoria feeling sad or empty lack of appetite loss of interest or pleasure mental depression paranoia quick to react or overreact rapidly changing moods runny nose sleepiness or unusual drowsiness sleeplessness sneezing stuffy nose trouble concentrating trouble sleeping unable to sleep Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. IBU-200 side effects

2169. This particular disease, Osteoarthritis, interacts with the drug Midol IB and has the following

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. Midol IB may cause serious side effects. Stop using Midol IB and call your doctor at once if you have: changes in your vision
- shortness of breath (even with mild exertion)

Drug Side Effects Summary

- swelling or rapid weight gain
- a skin rash, no matter how mild
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems--nausea, upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- low red blood cells (anemia)--pale skin, feeling light-headed or short of breath, rapid heart rate, trouble concentrating
- or kidney problems--little or no urinating, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath. Common side effects of Midol IB may include: nausea, vomiting, gas
- bleeding
- or dizziness , headache.

2170. This particular disease, Osteoarthritis, interacts with the drug Motrin IB and has the following

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. Stop using Motrin IB and call your doctor at once if you have: changes in your vision
- shortness of breath (even with mild exertion)
- swelling or rapid weight gain
- a skin rash, no matter how mild
- signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems - nausea, upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- low red blood cells (anemia) - pale skin, feeling light-headed or short of breath, rapid heart rate,

Drug Side Effects Summary

trouble concentrating

- or kidney problems - little or no urinating, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath. Common Motrin IB side effects may include: nausea, vomiting, gas
- bleeding
- or dizziness , headache.

2171. This particular disease, Osteoarthritis, interacts with the drug Naprelan and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Belching bruising chest tightness difficult or labored breathing feeling of indigestion headache itching skin large, flat, blue, or purplish patches in the skin pain in the chest below the breastbone skin eruptions stomach pain swelling Less common Bloating bloody or black, tarry stools blurred or loss of vision burning upper stomach pain cloudy urine constipation decrease in urine output or decrease in urine-concentrating ability disturbed color perception double vision fast, irregular, pounding, or racing heartbeat or pulse halos around lights indigestion loss of appetite nausea or vomiting night blindness overbright appearance of lights pale skin pinpoint red or purple spots on the skin severe and continuing nausea severe stomach burning, cramping, or pain skin rash swelling or inflammation of the mouth trouble breathing tunnel vision unusual bleeding or bruising unusual tiredness or weakness vomiting of material that looks like coffee grounds weight loss Rare Anxiety back or leg pains bleeding gums blindness blistering, peeling, or loosening of the skin blood in the urine or stools blue lips and fingernails canker sores change in the ability to see colors, especially blue or yellow chest pain or discomfort clay-colored stools cold sweats coma confusion cool, pale skin cough or hoarseness coughing that sometimes produces a pink frothy sputum cracks in the skin darkened urine

Drug Side Effects Summary

decreased vision depression diarrhea difficult, burning, or painful urination difficult, fast, or noisy breathing difficulty with swallowing dilated neck veins dizziness dry cough dry mouth early appearance of redness, or swelling of the skin excess air or gas in the stomach eye pain fainting fever with or without chills fluid-filled skin blisters flushed, dry skin frequent urination fruit-like breath odor greatly decreased frequency of urination or amount of urine hair loss high fever hives increased hunger increased sensitivity of the skin to sunlight increased sweating increased thirst increased urination increased volume of pale, dilute urine irregular breathing joint or muscle pain large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs late appearance of rash with or without weeping blisters that become crusted, especially in sun-exposed areas of skin, may extend to unexposed areas light-colored stools lightheadedness loss of heat from the body lower back or side pain nervousness nightmares no blood pressure no breathing no pulse nosebleeds numbness or tingling in the hands, feet, or lips pain in the ankles or knees pain or burning in the throat pain or discomfort in the arms, jaw, back, or neck painful, red lumps under the skin, mostly on the legs pains in the stomach, side, or abdomen, possibly radiating to the back pale or blue lips, fingernails, or skin pounding in the ears puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rapid, shallow breathing red, irritated eyes red skin lesions, often with a purple center red-green color blindness redness or other discoloration of the skin redness, swelling, or soreness of the tongue scaly skin seizures severe sunburn shakiness skin thinness slurred speech sneezing sore throat sores, ulcers, or white spots on the lips or tongue or inside the mouth sores, welting, or blisters spots on your skin resembling a blister or pimple stiff neck or back stomach cramps or tenderness stomach upset swelling in the legs and ankles swelling of the face, fingers, feet, or lower legs swollen, painful, or tender lymph glands in the neck, armpit, or groin tiny bumps on the inner lining of the eyelid unexplained weight loss unpleasant breath odor watery or bloody diarrhea weakness or heaviness of the legs weight gain yellow eyes or skin Symptoms of overdose Bleeding under the skin confusion about identity, place, and time muscle tremors

Drug Side Effects Summary

restlessness sleepiness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Continuing ringing or buzzing or other unexplained noise in the ears hearing loss Less common Acid or sour stomach change in hearing feeling of constant movement of self or surroundings passing gas sensation of spinning stomach soreness or discomfort Rare Appetite changes burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings burning, dry, or itching eyes difficulty with moving discharge, excessive tearing general feeling of discomfort or illness lack or loss of strength menstrual changes muscle aching, cramping, stiffness, or weakness not able to concentrate redness, pain, or swelling of the eye, eyelid, or inner lining of the eyelid seeing, hearing, or feeling things that are not there shakiness in the legs, arms, hands, or feet swollen joints trembling or shaking of the hands or feet trouble getting pregnant trouble performing routine tasks trouble sleeping unusual drowsiness, dullness, or feeling of sluggishness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Naprelan side effects

2172. This particular disease, Osteoarthritis, interacts with the drug Nuprin and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Abdominal pain acid or sour stomach belching bloating cloudy urine decrease in amount of urine decrease in urine output or decrease in urine-concentrating ability diarrhea difficulty having a bowel movement (stool) excess air

Drug Side Effects Summary

or gas in stomach or intestines full feeling heartburn indigestion itching skin pain or discomfort in chest, upper stomach, or throat pale skin passing gas nausea noisy, rattling breathing rash with flat lesions or small raised lesions on the skin shortness of breath swelling of face, fingers, hands, feet, lower legs, or ankles troubled breathing at rest troubled breathing with exertion unusual bleeding or bruising unusual tiredness or weakness vomiting weight gain Less common Abdominal cramps stomach soreness or discomfort Rare Agitation back, leg, or stomach pains bleeding gums blistering, peeling, loosening of skin blood in urine or stools bloody, black, or tarry stools blurred vision burning feeling in chest or stomach change in vision chest pain chills clay-colored stools coma confusion constipation cough or hoarseness dark urine decreased urine output depression difficulty breathing difficulty swallowing dilated neck veins dizziness dry mouth extreme fatigue fast, irregular, pounding, or racing heartbeat or pulse fever with or without chills frequent urination general body swelling general feeling of tiredness or weakness hair loss, thinning of hair headache hives or welts hostility impaired vision increased blood pressure increased volume of pale, dilute urine irregular breathing irritability itching joint or muscle pain lab results that show problems with liver lethargy light-colored stools loss of appetite lower back or side pain muscle twitching nosebleeds painful or difficult urination pains in stomach, side, or abdomen, possibly radiating to the back pinpoint red spots on skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rash red skin lesions, often with a purple center red, irritated eyes redness of skin seizures severe abdominal pain, cramping, burning severe and continuing nausea sore throat sores, ulcers, or white spots in mouth or on lips stiff neck or back stomach upset stupor swollen or painful glands tenderness in stomach area thirst tightness in chest unpleasant breath odor upper right abdominal pain vomiting of blood vomiting of material that looks like coffee grounds wheezing yellow eyes and skin Symptoms of overdose Bluish lips or skin difficulty sleeping disorientation dizziness, faintness, or lightheadedness when getting up from a lying or sitting position suddenly drowsiness to profound coma hallucination lightheadedness or fainting mood or other mental changes muscle tremors not

Drug Side Effects Summary

breathing rapid, deep breathing restlessness slow or irregular heartbeat stomach cramps sudden fainting sweating Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Continuing ringing or buzzing or other unexplained noise in ears hearing loss nervousness Rare Crying depersonalization discouragement dry eyes dysphoria euphoria feeling sad or empty lack of appetite loss of interest or pleasure mental depression paranoia quick to react or overreact rapidly changing moods runny nose sleepiness or unusual drowsiness sleeplessness sneezing stuffy nose trouble concentrating trouble sleeping unable to sleep Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Nuprin side effects

2173. This particular disease, Osteoarthritis, interacts with the drug Osteo Bi-Flex and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Although not all side effects are known, Osteo Bi-Flex is thought to be possibly safe when taken for up to 2 years. Stop using Osteo Bi-Flex and call your healthcare provider at once if you have: irregular heartbeats
- or swelling in your legs. Common side effects of Osteo Bi-Flex may include: nausea , diarrhea , constipation
- stomach pain, gas, bloating
- hair loss
- or puffy eyelids.

Drug Side Effects Summary

2174. This particular disease, Osteoarthritis, interacts with the drug piroxicam and has the following

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. Stop using piroxicam and seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Piroxicam may cause serious side effects. Stop using piroxicam and call your doctor at once if you have: severe headache , blurred vision, pounding in your neck or ears
- heart problems--swelling, rapid weight gain, feeling short of breath
- liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Common side effects of piroxicam may include: abnormal liver function tests
- urination problems
- upset stomach , heartburn , loss of appetite, stomach pain, nausea, vomiting
- gas, diarrhea , constipation
- dizziness , headache
- itching, rash
- or ringing in your ears.

2175. This particular disease, Osteoarthritis, interacts with the drug Aflaxen and has the following s

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of

Drug Side Effects Summary

these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Belching bruising chest tightness difficult or labored breathing feeling of indigestion headache itching skin large, flat, blue, or purplish patches in the skin pain in the chest below the breastbone skin eruptions stomach pain swelling Less common Bloating bloody or black, tarry stools blurred or loss of vision burning upper stomach pain cloudy urine constipation decrease in urine output or decrease in urine-concentrating ability disturbed color perception double vision fast, irregular, pounding, or racing heartbeat or pulse halos around lights indigestion loss of appetite nausea or vomiting night blindness overbright appearance of lights pale skin pinpoint red or purple spots on the skin severe and continuing nausea severe stomach burning, cramping, or pain skin rash swelling or inflammation of the mouth trouble breathing tunnel vision unusual bleeding or bruising unusual tiredness or weakness vomiting of material that looks like coffee grounds weight loss Rare Anxiety back or leg pains bleeding gums blindness blistering, peeling, or loosening of the skin blood in the urine or stools blue lips and fingernails canker sores change in the ability to see colors, especially blue or yellow chest pain or discomfort clay-colored stools cold sweats coma confusion cool, pale skin cough or hoarseness coughing that sometimes produces a pink frothy sputum cracks in the skin darkened urine decreased vision depression diarrhea difficult, burning, or painful urination difficult, fast, or noisy breathing difficulty with swallowing dilated neck veins dizziness dry cough dry mouth early appearance of redness, or swelling of the skin excess air or gas in the stomach eye pain fainting fever with or without chills fluid-filled skin blisters flushed, dry skin frequent urination fruit-like breath odor greatly decreased frequency of urination or amount of urine hair loss high fever hives increased hunger increased sensitivity of the skin to sunlight increased sweating increased thirst increased urination increased volume of pale, dilute urine irregular breathing joint or muscle pain large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs late appearance of rash with or without weeping blisters that become crusted, especially in sun-exposed

Drug Side Effects Summary

areas of skin, may extend to unexposed areas light-colored stools lightheadedness loss of heat from the body lower back or side pain nervousness nightmares no blood pressure no breathing no pulse nosebleeds numbness or tingling in the hands, feet, or lips pain in the ankles or knees pain or burning in the throat pain or discomfort in the arms, jaw, back, or neck painful, red lumps under the skin, mostly on the legs pains in the stomach, side, or abdomen, possibly radiating to the back pale or blue lips, fingernails, or skin pounding in the ears puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rapid, shallow breathing red, irritated eyes red skin lesions, often with a purple center red-green color blindness redness or other discoloration of the skin redness, swelling, or soreness of the tongue scaly skin seizures severe sunburn shakiness skin thinness slurred speech sneezing sore throat sores, ulcers, or white spots on the lips or tongue or inside the mouth sores, welting, or blisters spots on your skin resembling a blister or pimple stiff neck or back stomach cramps or tenderness stomach upset swelling in the legs and ankles swelling of the face, fingers, feet, or lower legs swollen, painful, or tender lymph glands in the neck, armpit, or groin tiny bumps on the inner lining of the eyelid unexplained weight loss unpleasant breath odor watery or bloody diarrhea weakness or heaviness of the legs weight gain yellow eyes or skin Symptoms of overdose Bleeding under the skin confusion about identity, place, and time muscle tremors restlessness sleepiness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Continuing ringing or buzzing or other unexplained noise in the ears hearing loss Less common Acid or sour stomach change in hearing feeling of constant movement of self or surroundings passing gas sensation of spinning stomach soreness or discomfort Rare Appetite changes burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings burning, dry, or itching eyes difficulty with moving

Drug Side Effects Summary

discharge, excessive tearing general feeling of discomfort or illness lack or loss of strength menstrual changes muscle aching, cramping, stiffness, or weakness not able to concentrate redness, pain, or swelling of the eye, eyelid, or inner lining of the eyelid seeing, hearing, or feeling things that are not there shakiness in the legs, arms, hands, or feet swollen joints trembling or shaking of the hands or feet trouble getting pregnant trouble performing routine tasks trouble sleeping unusual drowsiness, dullness, or feeling of sluggishness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Aflaxen side effects

2176. This particular disease, Osteoarthritis, interacts with the drug All Day Pain Relief and has the

- No side effects listed

2177. This particular disease, Osteoarthritis, interacts with the drug All Day Relief and has the follow

- No side effects listed

2178. This particular disease, Osteoarthritis, interacts with the drug Anaprox and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Belching bruising chest tightness difficult or labored breathing feeling of indigestion headache itching skin large, flat, blue, or purplish patches in the skin pain in the chest below the breastbone skin eruptions stomach pain swelling Less common Bloating bloody or black, tarry stools blurred or loss of vision burning upper stomach pain cloudy urine constipation decrease in urine output or decrease in urine-concentrating ability disturbed color perception double vision fast, irregular, pounding, or racing heartbeat or pulse

Drug Side Effects Summary

halos around lights indigestion loss of appetite nausea or vomiting night blindness overbright appearance of lights pale skin pinpoint red or purple spots on the skin severe and continuing nausea severe stomach burning, cramping, or pain skin rash swelling or inflammation of the mouth trouble breathing tunnel vision unusual bleeding or bruising unusual tiredness or weakness vomiting of material that looks like coffee grounds weight loss Rare Anxiety back or leg pains bleeding gums blindness blistering, peeling, or loosening of the skin blood in the urine or stools blue lips and fingernails canker sores change in the ability to see colors, especially blue or yellow chest pain or discomfort clay-colored stools cold sweats coma confusion cool, pale skin cough or hoarseness coughing that sometimes produces a pink frothy sputum cracks in the skin darkened urine decreased vision depression diarrhea difficult, burning, or painful urination difficult, fast, or noisy breathing difficulty with swallowing dilated neck veins dizziness dry cough dry mouth early appearance of redness, or swelling of the skin excess air or gas in the stomach eye pain fainting fever with or without chills fluid-filled skin blisters flushed, dry skin frequent urination fruit-like breath odor greatly decreased frequency of urination or amount of urine hair loss high fever hives increased hunger increased sensitivity of the skin to sunlight increased sweating increased thirst increased urination increased volume of pale, dilute urine irregular breathing joint or muscle pain large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs late appearance of rash with or without weeping blisters that become crusted, especially in sun-exposed areas of skin, may extend to unexposed areas light-colored stools lightheadedness loss of heat from the body lower back or side pain nervousness nightmares no blood pressure no breathing no pulse nosebleeds numbness or tingling in the hands, feet, or lips pain in the ankles or knees pain or burning in the throat pain or discomfort in the arms, jaw, back, or neck painful, red lumps under the skin, mostly on the legs pains in the stomach, side, or abdomen, possibly radiating to the back pale or blue lips, fingernails, or skin pounding in the ears puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rapid, shallow breathing red, irritated eyes red skin lesions, often with

Drug Side Effects Summary

a purple center red-green color blindness redness or other discoloration of the skin redness, swelling, or soreness of the tongue scaly skin seizures severe sunburn shakiness skin thinness slurred speech sneezing sore throat sores, ulcers, or white spots on the lips or tongue or inside the mouth sores, welting, or blisters spots on your skin resembling a blister or pimple stiff neck or back stomach cramps or tenderness stomach upset swelling in the legs and ankles swelling of the face, fingers, feet, or lower legs swollen, painful, or tender lymph glands in the neck, armpit, or groin tiny bumps on the inner lining of the eyelid unexplained weight loss unpleasant breath odor watery or bloody diarrhea weakness or heaviness of the legs weight gain yellow eyes or skin Symptoms of overdose Bleeding under the skin confusion about identity, place, and time muscle tremors restlessness sleepiness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Continuing ringing or buzzing or other unexplained noise in the ears hearing loss Less common Acid or sour stomach change in hearing feeling of constant movement of self or surroundings passing gas sensation of spinning stomach soreness or discomfort Rare Appetite changes burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings burning, dry, or itching eyes difficulty with moving discharge, excessive tearing general feeling of discomfort or illness lack or loss of strength menstrual changes muscle aching, cramping, stiffness, or weakness not able to concentrate redness, pain, or swelling of the eye, eyelid, or inner lining of the eyelid seeing, hearing, or feeling things that are not there shakiness in the legs, arms, hands, or feet swollen joints trembling or shaking of the hands or feet trouble getting pregnant trouble performing routine tasks trouble sleeping unusual drowsiness, dullness, or feeling of sluggishness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional.

Drug Side Effects Summary

Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Anaprox side effects

2179. This particular disease, Osteoarthritis, interacts with the drug Anaprox-DS and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Belching bruising chest tightness difficult or labored breathing feeling of indigestion headache itching skin large, flat, blue, or purplish patches in the skin pain in the chest below the breastbone skin eruptions stomach pain swelling Less common Bloating bloody or black, tarry stools blurred or loss of vision burning upper stomach pain cloudy urine constipation decrease in urine output or decrease in urine-concentrating ability disturbed color perception double vision fast, irregular, pounding, or racing heartbeat or pulse halos around lights indigestion loss of appetite nausea or vomiting night blindness overbright appearance of lights pale skin pinpoint red or purple spots on the skin severe and continuing nausea severe stomach burning, cramping, or pain skin rash swelling or inflammation of the mouth trouble breathing tunnel vision unusual bleeding or bruising unusual tiredness or weakness vomiting of material that looks like coffee grounds weight loss Rare Anxiety back or leg pains bleeding gums blindness blistering, peeling, or loosening of the skin blood in the urine or stools blue lips and fingernails canker sores change in the ability to see colors, especially blue or yellow chest pain or discomfort clay-colored stools cold sweats coma confusion cool, pale skin cough or hoarseness coughing that sometimes produces a pink frothy sputum cracks in the skin darkened urine decreased vision depression diarrhea difficult, burning, or painful urination difficult, fast, or noisy breathing difficulty with swallowing dilated neck veins dizziness dry cough dry mouth early appearance of redness, or swelling of the skin excess air or gas in the stomach eye pain fainting fever with or without chills fluid-filled skin blisters flushed, dry skin frequent urination fruit-like breath

Drug Side Effects Summary

odor greatly decreased frequency of urination or amount of urine hair loss high fever hives increased hunger increased sensitivity of the skin to sunlight increased sweating increased thirst increased urination increased volume of pale, dilute urine irregular breathing joint or muscle pain large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs late appearance of rash with or without weeping blisters that become crusted, especially in sun-exposed areas of skin, may extend to unexposed areas light-colored stools lightheadedness loss of heat from the body lower back or side pain nervousness nightmares no blood pressure no breathing no pulse nosebleeds numbness or tingling in the hands, feet, or lips pain in the ankles or knees pain or burning in the throat pain or discomfort in the arms, jaw, back, or neck painful, red lumps under the skin, mostly on the legs pains in the stomach, side, or abdomen, possibly radiating to the back pale or blue lips, fingernails, or skin pounding in the ears puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rapid, shallow breathing red, irritated eyes red skin lesions, often with a purple center red-green color blindness redness or other discoloration of the skin redness, swelling, or soreness of the tongue scaly skin seizures severe sunburn shakiness skin thinness slurred speech sneezing sore throat sores, ulcers, or white spots on the lips or tongue or inside the mouth sores, welting, or blisters spots on your skin resembling a blister or pimple stiff neck or back stomach cramps or tenderness stomach upset swelling in the legs and ankles swelling of the face, fingers, feet, or lower legs swollen, painful, or tender lymph glands in the neck, armpit, or groin tiny bumps on the inner lining of the eyelid unexplained weight loss unpleasant breath odor watery or bloody diarrhea weakness or heaviness of the legs weight gain yellow eyes or skin Symptoms of overdose Bleeding under the skin confusion about identity, place, and time muscle tremors restlessness sleepiness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are

Drug Side Effects Summary

bothersome or if you have any questions about them: More common Continuing ringing or buzzing or other unexplained noise in the ears hearing loss Less common Acid or sour stomach change in hearing feeling of constant movement of self or surroundings passing gas sensation of spinning stomach soreness or discomfort Rare Appetite changes burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings burning, dry, or itching eyes difficulty with moving discharge, excessive tearing general feeling of discomfort or illness lack or loss of strength menstrual changes muscle aching, cramping, stiffness, or weakness not able to concentrate redness, pain, or swelling of the eye, eyelid, or inner lining of the eyelid seeing, hearing, or feeling things that are not there shakiness in the legs, arms, hands, or feet swollen joints trembling or shaking of the hands or feet trouble getting pregnant trouble performing routine tasks trouble sleeping unusual drowsiness, dullness, or feeling of sluggishness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Anaprox-DS side effects

2180. This particular disease, Osteoarthritis, interacts with the drug Cosamin DS and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Cosamin DS and call your healthcare provider at once if you have: irregular heartbeats
- or swelling in your legs. Common side effects of Cosamin DS may include: nausea , diarrhea , constipation
- stomach pain, gas, bloating
- hair loss
- or puffy eyelids.

Drug Side Effects Summary

2181. This particular disease, Osteoarthritis, interacts with the drug Diclozor and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of bleeding like throwing up or coughing up blood
- vomit that looks like coffee grounds
- blood in the urine
- black, red, or tarry stools
- bleeding from the gums
- abnormal vaginal bleeding
- bruises without a cause or that get bigger
- or bleeding you cannot stop. Signs of high blood pressure like very bad headache or dizziness , passing out, or change in eyesight. Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain. Signs of high potassium levels like a heartbeat that does not feel normal
- feeling confused

Drug Side Effects Summary

- feeling weak, lightheaded, or dizzy
- feeling like passing out
- numbness or tingling
- or shortness of breath. Chest pain or pressure or a fast heartbeat . Shortness of breath, a big weight gain, or swelling in the arms or legs. Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight. Feeling very tired or weak. Flu-like signs. Very bad back pain . Very bad belly pain. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Liver problems have happened with drugs like this one. Sometimes, this has been deadly. Call your doctor right away if you have signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Diclozor side effects

2182. This particular disease, Osteoarthritis, interacts with the drug diflunisal and has the following

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, swelling in your legs, feeling short of breath. Diflunisal may cause serious side effects. Stop using diflunisal and call your doctor at once if you have: changes in your vision
- any skin rash, no matter how mild
- shortness of breath (even with mild exertion)
- swelling or rapid weight gain
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds

Drug Side Effects Summary

- liver problems-- nausea , upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired
- or low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed, cold hands and feet. Common side effects of diflunisal may include: nausea, vomiting, stomach pain, indigestion
- diarrhea, constipation , gas
- dizziness , drowsiness, headache , tired feeling
- rash
- or ringing in your ears.

2183. This particular disease, Osteoarthritis, interacts with the drug Dolobid and has the following s

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Stomach cramps or pain Less common Back or leg pains bleeding gums blistering, peeling, loosening of skin bloody or black, tarry stools bloody or cloudy urine blurred or loss of vision burning feeling in chest or stomach burning while urinating clay-colored stools constipation cough or hoarseness cracks in the skin dark-colored urine diarrhea difficult or painful urination difficulty breathing difficulty swallowing disturbed color perception dizziness double vision facial swelling fast heartbeat feeling of warmth fever with or without chills flu-like symptoms general body swelling greatly decreased frequency of urination or amount of urine halos around lights headache hives, itching, skin rash increased blood pressure increased thirst indigestion joint or muscle pain large, hive-like swelling on face, eyelids, lips, tongue, throat, hands, legs, feet, sex organs light-colored stools loss of appetite loss of heat from the body lower back or side pain muscle aches, pains, or weakness nausea or vomiting night blindness noisy breathing nosebleeds overbright appearance of lights pale skin pinpoint red spots

Drug Side Effects Summary

on skin puffiness or swelling of the eyelids or around the eyes, face, lips or tongue red, irritated eyes redness of the face, neck, arms and occasionally, upper chest red skin lesions, often with a purple center red, swollen skin scaly skin severe or continuing stomach pain sores, ulcers, or white spots in mouth or on lips sore throat stomach upset, tenderness swelling of the face, fingers, feet or lower legs swollen lymph nodes tightness in chest trouble breathing tunnel vision unpleasant breath odor unusual bleeding or bruising unusual tiredness or weakness upper right stomach pain vomiting of blood or material that looks like coffee grounds weight gain yellow eyes or skin Symptoms of overdose Change in consciousness confusion about identity, place, and time continuing ringing or buzzing or other unexplained noise in ears decreased awareness or responsiveness deep or fast breathing with dizziness fast, pounding, or irregular heartbeat or pulse hearing loss loss of consciousness numbness of feet, hands, and around mouth severe sleepiness sweating Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Acid or sour stomach belching heartburn indigestion stomach discomfort or upset Less common Bloated, full feeling burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings discouragement dry mucous membranes excess air or gas in stomach or intestines feeling of constant movement of self or surroundings feeling sad or empty increased sensitivity of skin to sunlight irritability lack or loss of strength lightheadedness loss of interest or pleasure mood or mental changes nervousness passing gas seeing, hearing, or feeling things that are not there sensation of spinning severe sunburn sleeplessness swelling or inflammation of the mouth tiredness trouble concentrating trouble sleeping unable to sleep weight loss Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You

Drug Side Effects Summary

may report side effects to the FDA at 1-800-FDA-1088. Dolobid side effects

2184. This particular disease, Osteoarthritis, interacts with the drug EC-Naprosyn and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Belching bruising chest tightness difficult or labored breathing feeling of indigestion headache itching skin large, flat, blue, or purplish patches in the skin pain in the chest below the breastbone skin eruptions stomach pain swelling Less common Bloating bloody or black, tarry stools blurred or loss of vision burning upper stomach pain cloudy urine constipation decrease in urine output or decrease in urine-concentrating ability disturbed color perception double vision fast, irregular, pounding, or racing heartbeat or pulse halos around lights indigestion loss of appetite nausea or vomiting night blindness overbright appearance of lights pale skin pinpoint red or purple spots on the skin severe and continuing nausea severe stomach burning, cramping, or pain skin rash swelling or inflammation of the mouth trouble breathing tunnel vision unusual bleeding or bruising unusual tiredness or weakness vomiting of material that looks like coffee grounds weight loss Rare Anxiety back or leg pains bleeding gums blindness blistering, peeling, or loosening of the skin blood in the urine or stools blue lips and fingernails canker sores change in the ability to see colors, especially blue or yellow chest pain or discomfort clay-colored stools cold sweats coma confusion cool, pale skin cough or hoarseness coughing that sometimes produces a pink frothy sputum cracks in the skin darkened urine decreased vision depression diarrhea difficult, burning, or painful urination difficult, fast, or noisy breathing difficulty with swallowing dilated neck veins dizziness dry cough dry mouth early appearance of redness, or swelling of the skin excess air or gas in the stomach eye pain fainting fever with or without chills fluid-filled skin blisters flushed, dry skin frequent urination fruit-like breath odor greatly decreased frequency of urination or amount of urine hair loss high fever hives increased

Drug Side Effects Summary

hunger increased sensitivity of the skin to sunlight increased sweating increased thirst increased urination increased volume of pale, dilute urine irregular breathing joint or muscle pain large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs late appearance of rash with or without weeping blisters that become crusted, especially in sun-exposed areas of skin, may extend to unexposed areas light-colored stools lightheadedness loss of heat from the body lower back or side pain nervousness nightmares no blood pressure no breathing no pulse nosebleeds numbness or tingling in the hands, feet, or lips pain in the ankles or knees pain or burning in the throat pain or discomfort in the arms, jaw, back, or neck painful, red lumps under the skin, mostly on the legs pains in the stomach, side, or abdomen, possibly radiating to the back pale or blue lips, fingernails, or skin pounding in the ears puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rapid, shallow breathing red, irritated eyes red skin lesions, often with a purple center red-green color blindness redness or other discoloration of the skin redness, swelling, or soreness of the tongue scaly skin seizures severe sunburn shakiness skin thinness slurred speech sneezing sore throat sores, ulcers, or white spots on the lips or tongue or inside the mouth sores, welting, or blisters spots on your skin resembling a blister or pimple stiff neck or back stomach cramps or tenderness stomach upset swelling in the legs and ankles swelling of the face, fingers, feet, or lower legs swollen, painful, or tender lymph glands in the neck, armpit, or groin tiny bumps on the inner lining of the eyelid unexplained weight loss unpleasant breath odor watery or bloody diarrhea weakness or heaviness of the legs weight gain yellow eyes or skin Symptoms of overdose Bleeding under the skin confusion about identity, place, and time muscle tremors restlessness sleepiness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Continuing ringing or buzzing

Drug Side Effects Summary

or other unexplained noise in the ears hearing loss Less common Acid or sour stomach change in hearing feeling of constant movement of self or surroundings passing gas sensation of spinning stomach soreness or discomfort Rare Appetite changes burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings burning, dry, or itching eyes difficulty with moving discharge, excessive tearing general feeling of discomfort or illness lack or loss of strength menstrual changes muscle aching, cramping, stiffness, or weakness not able to concentrate redness, pain, or swelling of the eye, eyelid, or inner lining of the eyelid seeing, hearing, or feeling things that are not there shakiness in the legs, arms, hands, or feet swollen joints trembling or shaking of the hands or feet trouble getting pregnant trouble performing routine tasks trouble sleeping unusual drowsiness, dullness, or feeling of sluggishness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. EC-Naprosyn side effects

2185. This particular disease, Osteoarthritis, interacts with the drug Feldene and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Bloating bloody or black, tarry stools burning upper stomach pain cloudy urine constipation decrease in urine output or decrease in urine-concentrating ability headache heartburn indigestion itching skin or rash loss of appetite nausea or vomiting pale skin severe stomach pain, cramping, or burning severe and continuing nausea swelling swelling of the face, fingers, feet, or lower legs trouble breathing unusual bleeding or bruising unusual tiredness or weakness vomiting of blood or material that looks like coffee grounds weight changes Less common Bleeding gums blood in the urine bloody nose blurred vision burning feeling in the chest or stomach burning, crawling, itching, numbness, prickling, "pins

Drug Side Effects Summary

and needles", or tingling feelings chest pain or tightness clay-colored stools confusion cough or hoarseness dark urine difficult or labored breathing difficult, burning, or painful urination difficulty with swallowing dilated neck veins dizziness fainting fever or chills flushing or redness of the skin frequent urge to urinate increased sensitivity of the skin to sunlight increased thirst increased volume of pale, dilute urine large, flat, blue, or purplish patches in the skin lightheadedness lower back or side pain nervousness noisy breathing numbness or tingling in the hands, feet, or lips pain or burning in the throat peeling of the skin pinpoint red or purple spots on the skin pounding in the ears rapid, shallow breathing redness or other discoloration of the skin redness, swelling, or soreness of the tongue severe sunburn slow, fast, pounding, or irregular heartbeat or pulse sore throat sores, ulcers, or white spots on the lips or tongue or inside the mouth stomach upset swelling or inflammation of the mouth swollen glands tenderness in the stomach area unpleasant breath odor unusually warm skin weakness or heaviness of the legs yellow eyes or skin Rare Anxiety back or leg pains burning, dry, or itching eyes cold sweats coma cracks in the skin diarrhea discharge or excessive tearing dizziness, faintness, or lightheadedness when getting up from a lying or sitting position dry mouth flushed, dry skin fruit-like breath odor general body swelling general feeling of discomfort or illness high fever increased hunger increased urination inflammation of the joints irregular, fast or slow, or shallow breathing joint pain large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs light-colored stools loss of heat from the body muscle aches and pains nightmares no blood pressure no breathing no pulse pain or discomfort in the arms, jaw, back, or neck pains in the stomach, side, or abdomen, possibly radiating to the back pale or blue lips, fingernails, or skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red skin lesions, often with a purple center red, irritated eyes redness, pain, or swelling of the eye, eyelid, or inner lining of the eyelid runny nose scaly skin seeing, hearing, or feeling things that are not there seizures severe headache shakiness shivering sleepiness slurred speech sneezing sores, welting, or blisters stiff neck or back stomach pain, continuing suddenly

Drug Side Effects Summary

sweating swollen, painful, or tender lymph glands in the neck, armpit, or groin trouble sleeping

Symptoms of overdose Agitation depression hives hostility irritability muscle twitching pain or discomfort in the chest, upper stomach, or throat rapid weight gain stupor swelling of the ankles or hands unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Acid or sour stomach belching continuing ringing or buzzing or other unexplained noise in the ears excess air or gas in the stomach or intestines hearing loss passing gas stomach discomfort or upset Less common Feeling of constant movement of self or surroundings hair loss or thinning of the hair lack or loss of strength sensation of spinning shakiness in the legs, arms, hands, or feet trembling or shaking of the hands or feet Rare Change in hearing changes in appetite inability to sit still mood alterations need to keep moving restlessness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Feldene side effects

2186. This particular disease, Osteoarthritis, interacts with the drug Flanax Pain Reliever and has the

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop taking Flanax and seek medical attention or call your doctor at once if you have any of these serious side effects: chest pain, weakness, shortness of breath, slurred speech, problems with vision or balance
- black, bloody, or tarry stools

Drug Side Effects Summary

- coughing up blood or vomit that looks like coffee grounds
- swelling or rapid weight gain
- urinating less than usual or not at all
- nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- bruising, severe tingling, numbness, pain, muscle weakness
- fever, headache, neck stiffness, chills, increased sensitivity to light, purple spots on the skin, and/or seizure (convulsions) or severe skin reaction -- fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain, followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Less serious Flanax side effects may include: upset stomach, mild heartburn or stomach pain, diarrhea, constipation
- bloating, gas
- dizziness, headache, nervousness
- skin itching or rash
- blurred vision
- or ringing in your ears.

2187. This particular disease, Osteoarthritis, interacts with the drug ketoprofen and has the following

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Ketoprofen may cause serious side effects. Stop using ketoprofen and call your doctor at once if you have: shortness of breath (even with mild exertion)
- swelling or rapid weight gain
- nausea, diarrhea , flu-like symptoms (fever, swollen glands, muscle aches, severe weakness)
- a skin rash, no matter how mild
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee

Drug Side Effects Summary

grounds

- liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Common side effects of ketoprofen may include: heartburn , stomach pain, gas, nausea, vomiting
- diarrhea, constipation
- or dizziness .

2188. This particular disease, Osteoarthritis, interacts with the drug Lofena and has the following side effects:

- No side effects listed

2189. This particular disease, Osteoarthritis, interacts with the drug oxaprozin and has the following side effects:

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Oxaprozin may cause serious side effects. Stop using oxaprozin and call your doctor at once if you have: a skin rash, no matter how mild
- heart problems--swelling, rapid weight gain, feeling short of breath
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems-- nausea , upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Common side effects of oxaprozin may include: nausea, upset stomach

Drug Side Effects Summary

- diarrhea , constipation
- or rash.

2190. This particular disease, Osteoarthritis, interacts with the drug Profinac and has the following

- No side effects listed

2191. This particular disease, Osteoarthritis, interacts with the drug salsalate and has the following

- swelling in your hands or feet, rapid weight gain
- severe headache, blurred vision, pounding in your neck or ears
- pale skin, unusual tiredness, cold hands and feet
- liver problems-- nausea , stomach pain (upper right side), tiredness, itching, jaundice (yellowing of the skin or eyes)
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- or signs of a stroke--sudden numbness or weakness (especially on one side of the body), severe headache, slurred speech, balance problems. Older adults may be more likely to have side effects from salsalate. Common side effects of salsalate may include: nausea
- hearing problems
- a spinning sensation
- rash
- or abnormal liver function tests .

2192. This particular disease, Osteoarthritis, interacts with the drug Schiff Move Free and has the following

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Although not all side effects are known, Schiff Move Free is thought to be possibly safe when taken for up to 2 years. Stop using Schiff Move Free and call your healthcare provider at once if you have: irregular heartbeats
- or swelling in your legs. Common side effects of Schiff Move Free may include: nausea , diarrhea , constipation
- stomach pain, gas, bloating
- hair loss
- or puffy eyelids.

2193. This particular disease, Osteoarthritis, interacts with the drug sulindac and has the following

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Sulindac may cause serious side effects. Stop using sulindac and call your doctor at once if you have: changes in your vision
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems-- nausea , upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- pancreatitis --severe pain in your upper stomach spreading to your back, nausea and vomiting
- kidney problems--little or no urinating, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- or low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Common side effects of sulindac may include: nausea, vomiting ,

Drug Side Effects Summary

stomach pain, indigestion , loss of appetite

- diarrhea , constipation , gas
- headache , dizziness , nervousness
- itching, rash
- or ringing in your ears.

2194. This particular disease, Osteoarthritis, interacts with the drug Xrylix and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of bleeding like throwing up or coughing up blood
- vomit that looks like coffee grounds
- blood in the urine
- black, red, or tarry stools
- bleeding from the gums
- abnormal vaginal bleeding

Drug Side Effects Summary

- bruises without a cause or that get bigger
- or bleeding you cannot stop. Signs of high blood pressure like very bad headache or dizziness , passing out, or change in eyesight. Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain. Signs of high potassium levels like a heartbeat that does not feel normal
- feeling confused
- feeling weak, lightheaded, or dizzy
- feeling like passing out
- numbness or tingling
- or shortness of breath. Chest pain or pressure or a fast heartbeat . Shortness of breath, a big weight gain, or swelling in the arms or legs. Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight. Feeling very tired or weak. Flu-like signs. Very bad back pain . Very bad belly pain. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Liver problems have happened with drugs like this one. Sometimes, this has been deadly. Call your doctor right away if you have signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Xrylix side effects

2195. This particular disease, Osteoarthritis, interacts with the drug betamethasone and has the fol

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Betamethasone may cause serious side effects. Call your doctor at once if you have: fast, slow, or irregular heartbeats
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- bumps on the skin, or raised blotches (purple, pink, brown, or red)
- chest pain, swollen glands
- a seizure
- swelling, rapid weight gain, shortness of breath
- any wound that will not heal
- pain and swelling or stiffness in your joints with fever or general ill feeling
- severe pain in your upper stomach spreading to your back
- severe depression , changes in personality, unusual thoughts or behavior
- or increased adrenal gland hormones --weight gain in your face and shoulders, slow wound healing, skin discoloration, thinning skin, increased body hair, tiredness, mood changes, menstrual changes, sexual changes. Betamethasone can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using betamethasone. Common side effects of betamethasone may include: headache , depression, mood swings
- sleep problems (insomnia)
- numbness , tingling, burning pain
- skin redness, itching, or discoloration
- thinning skin, stretch marks
- muscle weakness
- or nausea , bloating , stomach pain.

2196. This particular disease, Osteoarthritis, interacts with the drug bioflavonoids and has the follo

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Common side effects of bioflavonoids may include:
blurred vision
- fast or slow heart rate
- headache , pounding in your ears
- dizziness , nervousness
- skin redness or itching
- or swelling, pain, or a heavy feeling in your legs or feet. Less serious side effects may be more likely, and you may have none at all.

2197. This particular disease, Osteoarthritis, interacts with the drug capsaicin and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Wash the skin and get medical attention right away if you have severe burning, pain, swelling, or blistering of the skin where you applied capsaicin topical. Capsaicin topical may cause serious side effects. Stop using capsaicin topical and call your doctor at once if you have: severe pain or skin redness where the medicine was applied
- loss of feeling
- trouble breathing or swallowing (after accidental inhalation of capsaicin odor or dried residue). Common side effects may include pain or a burning sensation that can last for several hours or days (especially after your first use of capsaicin topical).

2198. This particular disease, Osteoarthritis, interacts with the drug Celestone Soluspan and has the

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Celestone Soluspan may cause serious side effects.

Call your doctor at once if you have: fast, slow, or irregular heartbeats

- blurred vision, tunnel vision, eye pain , or seeing halos around lights

- bumps on the skin, or raised blotches (purple, pink, brown, or red)

- chest pain, swollen glands

- a seizure

- swelling, rapid weight gain, shortness of breath

- any wound that will not heal

- pain and swelling or stiffness in your joints with fever or general ill feeling

- severe pain in your upper stomach spreading to your back

- severe depression , changes in personality, unusual thoughts or behavior

- or increased adrenal gland hormones --weight gain in your face and shoulders, slow wound healing, skin discoloration, thinning skin, increased body hair, tiredness, mood changes, menstrual changes, sexual changes. Celestone Soluspan can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using Celestone Soluspan. Common side effects of Celestone Soluspan may include: headache , depression, mood swings

- sleep problems (insomnia)

- numbness , tingling, burning pain

- skin redness, itching, or discoloration

- thinning skin, stretch marks

- muscle weakness

- or nausea , bloating , stomach pain.

2199. This particular disease, Osteoarthritis, interacts with the drug Clinoril and has the following s

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of

Drug Side Effects Summary

these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Acid or sour stomach belching constipation headache heartburn nausea or vomiting skin rash stomach pain Less common Excess air or gas in the stomach or intestines itching skin passing gas stomach cramps swelling weight loss Rare Back or leg pains bleeding gums blistering, peeling, or loosening of the skin bloating bloody or black, tarry stools bloody or cloudy urine blurred or loss of vision burning feeling in the chest or stomach burning while urinating chest pain chills clay-colored stools confusion cough or hoarseness cracks in the skin dark urine decreased urine output diarrhea difficult or labored breathing difficult or painful urination difficulty with swallowing dilated neck veins discoloration of urine disturbed color perception dizziness double vision extreme fatigue eye pain fainting feeling of warmth fever flu-like symptoms general body swelling general feeling of discomfort or illness groin pain halos around lights hives increased sensitivity of the skin to sunlight increased sweating increased thirst indigestion irregular breathing joint or muscle pain large, flat, blue, or purplish patches in the skin large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs light-colored stools loss of appetite loss of heat from the body lower back or side pain muscle aches, pains, or weakness nervousness night blindness noisy breathing nosebleeds numbness or tingling in the hands, feet, or lips overbright appearance of lights painful or difficult urination pains in the stomach, side, or abdomen, possibly radiating to the back pale skin persistent bleeding or oozing from puncture sites, mouth, or nose pinpoint red or purple spots on the skin pounding in the ears puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red skin lesions, often with a purple center red, irritated eyes redness of the face, neck, arms and occasionally, upper chest redness or other discoloration of the skin redness, swelling, or soreness of the tongue scaly skin seizures severe or continuing stomach pain severe sunburn shortness of breath slow or fast, irregular, pounding, or racing heartbeat or pulse sore throat sores, ulcers, or white spots in the mouth or on the lips stiff neck or back stomach upset swelling of the face, fingers,

Drug Side Effects Summary

feet, or lower legs swelling or inflammation of the mouth swollen or painful glands tenderness in the stomach area tightness in the chest troubled breathing tunnel vision unpleasant breath odor unusual bleeding or bruising unusual tiredness or weakness upper right abdominal pain vomiting of blood or material that looks like coffee grounds watery or bloody diarrhea weakness or heaviness of the legs weight gain wheezing yellow eyes or skin Symptoms of overdose Change in consciousness decreased awareness or responsiveness loss of consciousness severe sleepiness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Continuing ringing or buzzing or other unexplained noise in the ears hearing loss Rare Burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings burning, dry, or itching eyes decreased hearing difficulty with moving discharge, excessive tearing discouragement false sense of well-being feeling of constant movement of self or surroundings feeling sad or empty hair loss or thinning of the hair hallucinations irritability loss of interest or pleasure metallic or bitter taste mood swings muscle cramping or stiffness personality changes redness, pain, or swelling of the eye, eyelid, or inner lining of the eyelid restlessness sensation of spinning sleepiness or unusual drowsiness sleeplessness sore or dry mucous membranes swelling or redness in the joints trouble with concentrating trouble with sleeping unable to sleep vaginal bleeding Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Clinoril side effects

2200. This particular disease, Osteoarthritis, interacts with the drug cortisone and has the following

- hives

Drug Side Effects Summary

- difficulty breathing
- swelling of your face, lips, tongue, or throat. Cortisone may cause serious side effects. Call your doctor at once if you have: blurred vision, eye pain or redness, seeing halos around lights
- swelling, rapid weight gain, feeling short of breath
- unusual changes in mood or behavior
- skin lesions
- sudden unusual pain in a bone or joint
- severe headaches, ringing in your ears, pain behind your eyes
- stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- pancreatitis --severe pain in your upper stomach spreading to your back, nausea and vomiting
- or low blood potassium--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Cortisone can affect growth in children. Tell your doctor if your child is not growing at a normal rate. Common side effects of cortisone may include: high blood pressure
- muscle pain or weakness
- sleep problems (insomnia), mood changes
- thinning skin, bruising or discoloration
- increased sweating
- headache , dizziness , spinning sensation
- stomach pain, bloating
- or slow wound healing.

2201. This particular disease, Osteoarthritis, interacts with the drug Daypro and has the following s

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Daypro may cause serious side effects. Stop using

Drug Side Effects Summary

Daypro and call your doctor at once if you have: a skin rash, no matter how mild

- heart problems--swelling, rapid weight gain, feeling short of breath
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems-- nausea , upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Common side effects of Daypro may include: nausea, upset stomach
- diarrhea , constipation
- or rash.

2202. This particular disease, Osteoarthritis, interacts with the drug esomeprazole / naproxen and h

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Esomeprazole and naproxen may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- heart problems-- swelling, rapid weight gain, feeling short of breath
- kidney problems-- fever, rash, nausea , loss of appetite, joint pain, urinating less than usual, blood in your urine, weight gain
- liver problems--loss of appetite, stomach pain (upper right side), dark urine, jaundice (yellowing of the skin or eyes)
- low magnesium-- dizziness , fast or irregular heart rate, tremors (shaking) or jerking muscle movements, feeling jittery, muscle cramps, muscle spasms in your hands and feet, cough or choking feeling

Drug Side Effects Summary

- low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet
- or new or worsening symptoms of lupus--joint pain, and a skin rash on your cheeks or arms that worsens in sunlight. Taking esomeprazole and naproxen long-term may cause you to develop stomach growths called fundic gland polyps. Talk with your doctor about this risk. If you use esomeprazole and naproxen for longer than 3 years, you could develop a vitamin B-12 deficiency. Talk to your doctor about how to manage this condition if you develop it. Common side effects of esomeprazole and naproxen may include: stomach discomfort
- or diarrhea.

2203. This particular disease, Osteoarthritis, interacts with the drug Euflexxa and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Euflexxa may cause serious side effects. Call your doctor at once if you have: severe pain or swelling around the knee after the injection. Common side effects of Euflexxa may include: warmth, pain, redness, stiffness, bruising, or puffiness where the medicine was injected
- nausea , stomach pain
- trouble walking
- swelling in your hands or feet
- back pain , joint pain, muscle pain
- numbness or tingly feeling
- headache , dizziness
- or runny or stuffy nose, sneezing, sore throat .

Drug Side Effects Summary

2204. This particular disease, Osteoarthritis, interacts with the drug hyaluronan and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Hyaluronan may cause serious side effects. Call your doctor at once if you have: bleeding
- increased knee pain
- or signs of infection (warmth, swelling, redness) around your knee. Common side effects of hyaluronan may include: joint pain, stiffness, or swelling
- itching, numbness , or tingling
- headache , dizziness
- back pain
- swelling, pain, redness, or mild discomfort where the medicine was injected.

2205. This particular disease, Osteoarthritis, interacts with the drug Indocin SR and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Acid or sour stomach belching diarrhea heartburn indigestion nausea stomach discomfort, upset, or pain vomiting Rare Abdominal or stomach cramping, burning, or tenderness back or leg pains bleeding gums blistering, peeling, or loosening of the skin bloody or black, tarry stools blue lips and fingernails blurred vision breast enlargement and tenderness burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings burning upper abdominal or stomach pain canker sores change in consciousness change in hearing chest pain, discomfort, or burning clay colored stools cloudy or bloody urine confusion continuing diarrhea cough or hoarseness coughing that sometimes produces a pink frothy sputum cracks in the skin dark urine decreased appetite decreased vision or any

Drug Side Effects Summary

change in vision depression difficult or labored breathing difficulty with swallowing dilated neck veins dizziness, faintness, or lightheadedness when getting up from a lying or sitting position double vision dry mouth extreme fatigue false sense of well-being feeling of unreality feeling of warmth fever with or without chills flushed, dry skin fruit-like breath odor general body swelling greatly decreased frequency of urination or amount of urine hair loss headache heavier menstrual periods increased hunger increased sweating increased thirst increased urination irregular breathing irritation and swelling of the eye jerky movements of the head, face, mouth, and neck joint pain large, flat, blue or purplish patches in the skin large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs loss of appetite loss of balance control loss of bladder control loss of consciousness loss of hearing loss of heat from the body lower back or side pain mask-like face mood swings muscle aches, pains, or weakness muscle spasm or jerking of all extremities nervousness noisy, rattling breathing nosebleeds numbness or tingling in the hands, feet, or lips pain in the ankles or knees pain or discomfort in the upper stomach or throat pain with swallowing painful or difficult urination painful, red lumps under the skin, mostly on the legs pale skin persistent bleeding or oozing from puncture sites, mouth, or nose personality changes pinpoint red or purple spots on the skin pounding in the ears puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red skin lesions, often with a purple center red, irritated eyes red, swollen skin redness of the face, neck, arms and occasionally, upper chest scaly skin seeing double seeing, hearing, or feeling things that are not there seizures sense of detachment from self or body severe constipation severe mental changes severe or continuing stomach pain shuffling walk skin rash, hives or welts, itching slow, fast, irregular, pounding, or racing heartbeat or pulse slowed movements slurred speech small red or purple spots on the skin sore throat sores, ulcers, or white spots on the lips or tongue or inside the mouth stiffness of the arms and legs sudden loss of consciousness swelling of the breasts or breast soreness in both females and males swelling of the face, fingers, feet, ankles, or lower legs swollen or painful glands tightness in the chest trembling

Drug Side Effects Summary

and shaking of the fingers and hands troubled breathing at rest troubled breathing with exertion unexplained weight loss unpleasant breath odor unsteadiness or awkwardness unusual bleeding or bruising unusual tiredness or weakness vaginal bleeding vomiting of blood or material that looks like coffee grounds weakness in the arms, hands, legs, or feet weight gain yellow eyes or skin

Symptoms of overdose Confusion about identity, place, and time severe headache unusual drowsiness, dullness, or feeling of sluggishness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Mild headache Less common Continuing ringing or buzzing or other unexplained noise in the ears difficulty having a bowel movement (stool) discouragement feeling sad or empty general feeling of discomfort or illness hearing loss irritability loss of interest or pleasure sleepiness trouble with concentrating Rare Anxiety bloated or full feeling changes in patterns and rhythms of speech excess air or gas in the stomach or intestines feeling of constant movement of self or surroundings involuntary muscle movements lightheadedness passing gas sensation of spinning tiredness trouble sleeping trouble with speaking Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Indocin SR side effects

2206. This particular disease, Osteoarthritis, interacts with the drug Limbrel and has the following s

- Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your

Drug Side Effects Summary

health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Blurred vision dizziness dull ache or feeling of pressure or heaviness in legs fluid accumulation in the knee headache itching skin near damaged veins nervousness pounding in the ears red or scaling skin slow or fast heartbeat swollen feet and ankles Limbrel side effects

2207. This particular disease, Osteoarthritis, interacts with the drug Orthovisc and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: bleeding
- increased knee pain
- or signs of infection (warmth, swelling, redness) around your knee. Common Orthovisc side effects may include: joint pain, stiffness, or swelling
- itching, numbness, or tingling
- headache, dizziness
- back pain
- swelling, pain, redness, or mild discomfort where the medicine was injected.

2208. This particular disease, Osteoarthritis, interacts with the drug Osteo Bi-Flex Advanced and has

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Common side effects of Osteo Bi-Flex Advanced may include: gas, bloating , heartburn , stomach cramps, nausea
- headache , drowsiness
- diarrhea , constipation

Drug Side Effects Summary

- swelling in your ankles or feet
- itching, skin irritation
- or puffy eyelids.

2209. This particular disease, Osteoarthritis, interacts with the drug sodium hyaluronate and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Sodium hyaluronate may cause serious side effects. Call your doctor at once if you have: severe pain or swelling around the knee after the injection. Common side effects of sodium hyaluronate may include: warmth, pain, redness, stiffness, bruising, or puffiness where the medicine was injected
- nausea , stomach pain
- trouble walking
- swelling in your hands or feet
- back pain , joint pain, muscle pain
- numbness or tingly feeling
- headache , dizziness
- or runny or stuffy nose, sneezing, sore throat .

2210. This particular disease, Osteoarthritis, interacts with the drug Supartz and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Supartz may cause serious side effects. Call your doctor at once if you have: severe pain or swelling around the knee after the injection. Common side effects of Supartz may include: warmth, pain, redness, stiffness, bruising, or puffiness where the

Drug Side Effects Summary

medicine was injected

- nausea , stomach pain
- trouble walking
- swelling in your hands or feet
- back pain , joint pain, muscle pain
- numbness or tingly feeling
- headache , dizziness
- or runny or stuffy nose, sneezing, sore throat .

2211. This particular disease, Osteoarthritis, interacts with the drug Synvisc and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have severe pain or swelling around the knee after the injection. Common Synvisc side effects may include: warmth, redness, pain, stiffness, swelling, or puffiness where the medicine was injected
- muscle pain, trouble walking
- fever, chills, nausea
- prickly feeling in your skin
- headache, dizziness
- or itching or skin irritation around the knee.

2212. This particular disease, Osteoarthritis, interacts with the drug Synvisc-One and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have severe pain or

Drug Side Effects Summary

swelling around the knee after the injection. Less serious Synvisc-One side effects may include: warmth, pain, stiffness, swelling, or puffiness where the medicine was injected

- muscle pain, trouble walking
- fever, chills, nausea
- numbness or tingly feeling
- headache, dizziness
- tired feeling
- or itching or skin irritation around the knee.

2213. This particular disease, Osteoarthritis, interacts with the drug Zilretta and has the following s

- Rare serious allergic reactions Effects in the injected knee such as infection (with pain, swelling and restricted motion) or joint damage Increased chance of getting an infection, and a decreased ability to fight an infection Effects on hormone production. These effects can be reversible Elevated blood pressure, sodium and water retention, and potassium loss Intestinal perforation if you have certain gastrointestinal disorders Weakening of bones Changes in behavior or mood disturbances Increased pressure inside the eye Zilretta side effects

2214. This particular disease, Osteoarthritis, interacts with the drug Zostrix and has the following s

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Wash the skin and get medical attention right away if you have severe burning, pain, swelling, or blistering of the skin where you applied Zostrix. Zostrix may cause serious side effects. Stop using Zostrix and call your doctor at once if you have: severe pain or skin redness where the medicine was applied
- loss of feeling

Drug Side Effects Summary

- trouble breathing or swallowing (after accidental inhalation of capsaicin odor or dried residue).

Common side effects may include pain or a burning sensation that can last for several hours or days (especially after your first use of capsaicin topical).

2215. This particular disease, Osteoarthritis, interacts with the drug acetaminophen / aspirin and ha

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of bleeding like throwing up or coughing up blood
- vomit that looks like coffee grounds
- blood in the urine
- black, red, or tarry stools
- bleeding from the gums
- abnormal vaginal bleeding
- bruises without a cause or that get bigger

Drug Side Effects Summary

- or bleeding you cannot stop. Feeling confused. Feeling very tired or weak. Very bad belly pain. Ringing in ears. Hearing loss . Very bad headache or if headache is not better after the first dose. Pain for more than 10 days. Fever for more than 3 days. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Acetaminophen / aspirin side effects

2216. This particular disease, Osteoarthritis, interacts with the drug Arthricream and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Irritation where Arthricream (trolamine cream and lotion) is used. Redness. Arthricream side effects

2217. This particular disease, Osteoarthritis, interacts with the drug Asper-Flex and has the following

Drug Side Effects Summary

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Irritation where Asper-Flex (trolamine cream and lotion) is used. Redness. Asper-Flex side effects

2218. This particular disease, Osteoarthritis, interacts with the drug Axsain and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking

Drug Side Effects Summary

- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad burning or burning that does not go away. Axsain side effects

2219. This particular disease, Osteoarthritis, interacts with the drug Boroleum and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: redness or swelling of the treated area
- increased pain
- or severe burning or skin irritation such as a rash, itching, pain, or blistering. Less serious side effects may be more likely, and you may have none at all.

2220. This particular disease, Osteoarthritis, interacts with the drug Capsin and has the following s

- hives
 - difficulty breathing
 - swelling of your face, lips, tongue, or throat. Wash the skin and get medical attention right away if you have severe burning, pain, swelling, or blistering of the skin where you applied Capsin. Capsin may cause serious side effects. Stop using Capsin and call your doctor at once if you have: severe pain or skin redness where the medicine was applied
 - loss of feeling
 - trouble breathing or swallowing (after accidental inhalation of capsaicin odor or dried residue).
- Common side effects may include pain or a burning sensation that can last for several hours or days (especially after your first use of capsaicin topical).

Drug Side Effects Summary

2221. This particular disease, Osteoarthritis, interacts with the drug Capzasin and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Capzasin side effects

2222. This particular disease, Osteoarthritis, interacts with the drug Capzasin-HP and has the follow

- hives
 - difficulty breathing
 - swelling of your face, lips, tongue, or throat. Wash the skin and get medical attention right away if you have severe burning, pain, swelling, or blistering of the skin where you applied Capzasin-HP. Capzasin-HP may cause serious side effects. Stop using Capzasin-HP and call your doctor at once if you have: severe pain or skin redness where the medicine was applied
 - loss of feeling
 - trouble breathing or swallowing (after accidental inhalation of capsaicin odor or dried residue).
- Common side effects may include pain or a burning sensation that can last for several hours or days (especially after your first use of capsaicin topical).

Drug Side Effects Summary

2223. This particular disease, Osteoarthritis, interacts with the drug Capzasin-P and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Capzasin-P side effects

2224. This particular disease, Osteoarthritis, interacts with the drug Castiva Cooling and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: redness or swelling of the treated area
- increased pain
- or severe burning or skin irritation such as a rash, itching, pain, or blistering. Less serious side effects may be more likely, and you may have none at all.

2225. This particular disease, Osteoarthritis, interacts with the drug Castiva Warming and has the following side effects:

- hives

Drug Side Effects Summary

- difficulty breathing
 - swelling of your face, lips, tongue, or throat. Wash the skin and get medical attention right away if you have severe burning, pain, swelling, or blistering of the skin where you applied Castiva Warming. Castiva Warming may cause serious side effects. Stop using Castiva Warming and call your doctor at once if you have: severe pain or skin redness where the medicine was applied
 - loss of feeling
 - trouble breathing or swallowing (after accidental inhalation of capsaicin odor or dried residue).
- Common side effects may include pain or a burning sensation that can last for several hours or days (especially after your first use of capsaicin topical).

2226. This particular disease, Osteoarthritis, interacts with the drug Choline Magnesium Trisalicylate

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Choline magnesium trisalicylate may cause serious side effects. Stop using choline magnesium trisalicylate and call your doctor at once if you have: trouble breathing
- ringing in your ears, hearing loss
- behavior changes with nausea and vomiting in a child using choline magnesium trisalicylate
- worsening fever or pain
- or signs of stomach bleeding--feeling light-headed, ongoing stomach pain, bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Common side effects of choline magnesium trisalicylate may include: nausea , vomiting, stomach pain
- heartburn , indigestion
- or diarrhea, constipation .

Drug Side Effects Summary

2227. This particular disease, Osteoarthritis, interacts with the drug Cidaflex and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Although not all side effects are known, Cidaflex is thought to be possibly safe when taken for up to 2 years. Stop using Cidaflex and call your healthcare provider at once if you have: irregular heartbeats
- or swelling in your legs. Common side effects of Cidaflex may include: nausea , diarrhea , constipation
- stomach pain, gas, bloating
- hair loss
- or puffy eyelids.

2228. This particular disease, Osteoarthritis, interacts with the drug diclofenac / menthol and has the

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Also call your doctor at once if you have: a skin rash, no matter how mild
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- liver problems--nausea, diarrhea , stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Serious side effects may be more likely in older adults. Common side effects of diclofenac and menthol topical may include: skin dryness, redness, itching, or rash
- blistering or hardening of the treated skin

Drug Side Effects Summary

- numbness, tingling, burning pain
- acne
- flushing (sudden warmth, redness, or tingly feeling)
- stomach pain, constipation , diarrhea, gas, heartburn , nausea, vomiting
- or dizziness .

2229. This particular disease, Osteoarthritis, interacts with the drug DicloPR and has the following :

- No side effects listed

2230. This particular disease, Osteoarthritis, interacts with the drug DicloStream and has the following :

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Also call your doctor at once if you have: a skin rash, no matter how mild
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- liver problems--nausea, diarrhea , stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Serious side effects may be more likely in older adults. Common side effects of DicloStream may include: skin dryness, redness, itching, or rash
- blistering or hardening of the treated skin
- numbness, tingling, burning pain
- acne
- flushing (sudden warmth, redness, or tingly feeling)

Drug Side Effects Summary

- stomach pain, constipation , diarrhea, gas, heartburn , nausea, vomiting
- or dizziness .

2231. This particular disease, Osteoarthritis, interacts with the drug Durolane and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Durolane may cause serious side effects. Call your doctor at once if you have: severe pain or swelling around the knee after the injection. Common side effects of Durolane may include: warmth, pain, redness, stiffness, bruising, or puffiness where the medicine was injected
- nausea , stomach pain
- trouble walking
- swelling in your hands or feet
- back pain , joint pain, muscle pain
- numbness or tingly feeling
- headache , dizziness
- or runny or stuffy nose, sneezing, sore throat .

2232. This particular disease, Osteoarthritis, interacts with the drug Eucalyptamint and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: redness or swelling of the treated area
- increased pain
- or severe burning or skin irritation such as a rash, itching, pain, or blistering. Less serious side

Drug Side Effects Summary

effects may be more likely, and you may have none at all.

2233. This particular disease, Osteoarthritis, interacts with the drug Exocaine Plus and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: redness or swelling of the treated area
- increased pain
- or severe burning or skin irritation such as a rash, itching, pain, or blistering. Less serious side effects may be more likely, and you may have none at all.

2234. This particular disease, Osteoarthritis, interacts with the drug famotidine / ibuprofen and has the following side effects:

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. Ibuprofen may cause serious side effects. Stop using ibuprofen and call your doctor at once if you have: changes in your vision
- the first sign of any skin rash, no matter how mild
- fever, headache , neck stiffness, increased sensitivity to light, nausea, vomiting, confusion, drowsiness
- severe headache, blurred vision, pounding in your neck or ears
- little or no urination
- swelling, rapid weight gain
- liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet

Drug Side Effects Summary

- or signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Some side effects may be more likely in older adults and in people who have severe kidney disease. Common side effects of famotidine and ibuprofen may include: nausea, stomach pain
- diarrhea , constipation
- or headache.

2235. This particular disease, Osteoarthritis, interacts with the drug fenoprofen and has the following

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, swelling in your legs, feeling short of breath. Fenoprofen may cause serious side effects. Stop using fenoprofen and call your doctor at once if you have: changes in your vision
- any skin rash, no matter how mild
- shortness of breath (even with mild exertion)
- swelling or rapid weight gain
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems-- nausea , upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired
- or low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed, cold hands and feet. Common side effects of fenoprofen may include: nausea, vomiting, stomach pain, indigestion
- diarrhea , constipation
- headache , dizziness , drowsiness, tiredness
- feeling nervous
- itching, sweating

Drug Side Effects Summary

- or ringing in your ears.

2236. This particular disease, Osteoarthritis, interacts with the drug Fenorthol and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of bleeding like throwing up or coughing up blood
- vomit that looks like coffee grounds
- blood in the urine
- black, red, or tarry stools
- bleeding from the gums
- abnormal vaginal bleeding
- bruises without a cause or that get bigger
- or bleeding you cannot stop. Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain. Signs of high blood pressure like very bad headache or dizziness , passing out, or change in eyesight. Shortness of breath, a big weight

Drug Side Effects Summary

gain, or swelling in the arms or legs. Chest pain or pressure or a fast heartbeat . Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight. Feeling very tired or weak. Ringing in ears. Change in eyesight. Very upset stomach or throwing up. Very bad belly pain. Very bad back pain . A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)

- red or irritated eyes

- or sores in your mouth, throat, nose, or eyes. Fenortho side effects

2237. This particular disease, Osteoarthritis, interacts with the drug Fenovar and has the following

- No side effects listed

2238. This particular disease, Osteoarthritis, interacts with the drug flurbiprofen and has the following

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Flurbiprofen may cause serious side effects. Stop using flurbiprofen and call your doctor at once if you have: changes in your vision

- a skin rash, no matter how mild

- shortness of breath (even with mild exertion)

- swelling or rapid weight gain

- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds

- liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)

- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath

Drug Side Effects Summary

- or low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Common side effects of flurbiprofen may include: indigestion , stomach pain, nausea , vomiting
- confusion, tremors, feeling nervous
- diarrhea , constipation
- headache , dizziness , drowsiness
- increased sweating, itching or rash
- or ringing in your ears.

2239. This particular disease, Osteoarthritis, interacts with the drug Gelsyn-3 and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Gelsyn-3 may cause serious side effects. Call your doctor at once if you have: severe pain or swelling around the knee after the injection. Common side effects of Gelsyn-3 may include: warmth, pain, redness, stiffness, bruising, or puffiness where the medicine was injected
- nausea , stomach pain
- trouble walking
- swelling in your hands or feet
- back pain , joint pain, muscle pain
- numbness or tingly feeling
- headache , dizziness
- or runny or stuffy nose, sneezing, sore throat .

2240. This particular disease, Osteoarthritis, interacts with the drug Glucoten and has the following

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Although not all side effects are known, Glucoten is thought to be possibly safe when taken for up to 2 years. Stop using Glucoten and call your healthcare provider at once if you have: irregular heartbeats
- or swelling in your legs. Common side effects of Glucoten may include: nausea , diarrhea , constipation
- stomach pain, gas, bloating
- hair loss
- or puffy eyelids.

2241. This particular disease, Osteoarthritis, interacts with the drug Gordogesic and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: redness or swelling of the treated area
- increased pain
- or severe burning or skin irritation such as a rash, itching, pain, or blistering. Less serious side effects may be more likely, and you may have none at all.

2242. This particular disease, Osteoarthritis, interacts with the drug Heet Triple Action and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: redness or swelling of the treated area

Drug Side Effects Summary

- increased pain
- or severe burning or skin irritation such as a rash, itching, pain, or blistering. Less serious side effects may be more likely, and you may have none at all.

2243. This particular disease, Osteoarthritis, interacts with the drug Hyalgan and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Hyalgan may cause serious side effects. Call your doctor at once if you have: severe pain or swelling around the knee after the injection. Common side effects of Hyalgan may include: warmth, pain, redness, stiffness, bruising, or puffiness where the medicine was injected
- nausea , stomach pain
- trouble walking
- swelling in your hands or feet
- back pain , joint pain, muscle pain
- numbness or tingly feeling
- headache , dizziness
- or runny or stuffy nose, sneezing, sore throat .

2244. This particular disease, Osteoarthritis, interacts with the drug hylan g-f 20 and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have severe pain or swelling around the knee after the injection. Common side effects of hylan G-F 20 may include: warmth, redness, pain, stiffness, swelling, or puffiness where the medicine was injected

Drug Side Effects Summary

- muscle pain, trouble walking
- fever, chills, nausea
- prickly feeling in your skin
- headache , dizziness
- or itching or skin irritation around the knee.

2245. This particular disease, Osteoarthritis, interacts with the drug Hymovis and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Hymovis may cause serious side effects. Call your doctor at once if you have: bleeding
- increased knee pain
- or signs of infection (warmth, swelling, redness) around your knee. Common side effects of Hymovis may include: joint pain, stiffness, or swelling
- itching, numbness , or tingling
- headache , dizziness
- back pain
- swelling, pain, redness, or mild discomfort where the medicine was injected.

2246. This particular disease, Osteoarthritis, interacts with the drug meclofenamate and has the fol

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Meclofenamate may cause serious side effects. Stop using meclofenamate and call your doctor at once if you have: a skin rash, no matter how mild
- shortness of breath (even with mild exertion)
- swelling or rapid weight gain

Drug Side Effects Summary

- nausea, diarrhea , flu-like symptoms (fever, swollen glands, muscle aches, severe weakness)
 - signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
 - liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
 - kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
 - o low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Older adults may be more likely to have symptoms of stomach bleeding.
- Common side effects of meclizine may include: indigestion , stomach pain, nausea
- diarrhea, constipation
 - or dizziness .

2247. This particular disease, Osteoarthritis, interacts with the drug Mentha Arthritis Cream with Capsaicin

- hives
 - difficulty breathing
 - swelling of your face, lips, tongue, or throat. Wash the skin and get medical attention right away if you have severe burning, pain, swelling, or blistering of the skin where you applied Mentha Arthritis Cream with Capsaicin. Mentha Arthritis Cream with Capsaicin may cause serious side effects. Stop using Mentha Arthritis Cream with Capsaicin and call your doctor at once if you have: severe pain or skin redness where the medicine was applied
 - loss of feeling
 - trouble breathing or swallowing (after accidental inhalation of capsaicin odor or dried residue).
- Common side effects may include pain or a burning sensation that can last for several hours or days (especially after your first use of capsaicin topical).

Drug Side Effects Summary

2248. This particular disease, Osteoarthritis, interacts with the drug Mentholatum Deep Heating and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: redness or swelling of the treated area
- increased pain
- or severe burning or skin irritation such as a rash, itching, pain, or blistering. Less serious side effects may be more likely, and you may have none at all.

2249. This particular disease, Osteoarthritis, interacts with the drug methyl salicylate and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: redness or swelling of the treated area
- increased pain
- or severe burning or skin irritation such as a rash, itching, pain, or blistering. Less serious side effects may be more likely, and you may have none at all.

2250. This particular disease, Osteoarthritis, interacts with the drug Monovisc and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Monovisc may cause serious side effects. Call your doctor at once if you have: bleeding
- increased knee pain
- or signs of infection (warmth, swelling, redness) around your knee. Common side effects of

Drug Side Effects Summary

Monovisc may include: joint pain, stiffness, or swelling

- itching, numbness , or tingling
- headache , dizziness
- back pain
- swelling, pain, redness, or mild discomfort where the medicine was injected.

2251. This particular disease, Osteoarthritis, interacts with the drug Myoflex and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Irritation where Myoflex (trolamine cream and lotion) is used. Redness. Myoflex side effects

2252. This particular disease, Osteoarthritis, interacts with the drug Nalfon and has the following side effects:

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, swelling in your legs, feeling short of breath. Nalfon may cause serious side effects. Stop using Nalfon and call your doctor at once if you have: changes in your vision

Drug Side Effects Summary

- any skin rash, no matter how mild
- shortness of breath (even with mild exertion)
- swelling or rapid weight gain
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems-- nausea , upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired
- or low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed, cold hands and feet. Common side effects of Nalfon may include: nausea, vomiting, stomach pain, indigestion
- diarrhea , constipation
- headache , dizziness , drowsiness, tiredness
- feeling nervous
- itching, sweating
- or ringing in your ears.

2253. This particular disease, Osteoarthritis, interacts with the drug OptiFlex Complete and has the

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Although not all side effects are known, OptiFlex Complete is thought to be possibly safe when taken for up to 2 years. Stop using OptiFlex Complete and call your healthcare provider at once if you have: irregular heartbeats
- or swelling in your legs. Common side effects of OptiFlex Complete may include: nausea , diarrhea , constipation
- stomach pain, gas, bloating

Drug Side Effects Summary

- hair loss
- or puffy eyelids.

2254. This particular disease, Osteoarthritis, interacts with the drug Pryflex and has the following s

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Although not all side effects are known, Pryflex is thought to be possibly safe when taken for up to 2 years. Stop using Pryflex and call your healthcare provider at once if you have: irregular heartbeats
- or swelling in your legs. Common side effects of Pryflex may include: nausea , diarrhea , constipation
- stomach pain, gas, bloating
- hair loss
- or puffy eyelids.

2255. This particular disease, Osteoarthritis, interacts with the drug Relafen DS and has the followin

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat

Drug Side Effects Summary

- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of bleeding like throwing up or coughing up blood
- vomit that looks like coffee grounds
- blood in the urine
- black, red, or tarry stools
- bleeding from the gums
- abnormal vaginal bleeding
- bruises without a cause or that get bigger
- or bleeding you cannot stop. Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain. Signs of high potassium levels like a heartbeat that does not feel normal
- feeling confused
- feeling weak, lightheaded, or dizzy
- feeling like passing out
- numbness or tingling
- or shortness of breath. Signs of high blood pressure like very bad headache or dizziness , passing out, or change in eyesight. Shortness of breath, a big weight gain, or swelling in the arms or legs. Chest pain or pressure. Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight. Feeling very tired or weak. Ringing in ears. Flu-like signs. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)

Drug Side Effects Summary

- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Relafen DS side effects

2256. This particular disease, Osteoarthritis, interacts with the drug Relamine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Although not all side effects are known, Relamine is thought to be possibly safe when taken for up to 2 years. Stop using Relamine and call your healthcare provider at once if you have: irregular heartbeats
- or swelling in your legs. Common side effects of Relamine may include: nausea , diarrhea , constipation
- stomach pain, gas, bloating
- hair loss
- or puffy eyelids.

2257. This particular disease, Osteoarthritis, interacts with the drug Salonpas Pain Patch and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: redness or swelling of the treated area
- increased pain
- or severe burning or skin irritation such as a rash, itching, pain, or blistering. Less serious side effects may be more likely, and you may have none at all.

2258. This particular disease, Osteoarthritis, interacts with the drug Satogesic and has the following

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: redness or swelling of the treated area
- increased pain
- or severe burning or skin irritation such as a rash, itching, pain, or blistering. Less serious side effects may be more likely, and you may have none at all.

2259. This particular disease, Osteoarthritis, interacts with the drug Sloan's Liniment and has the following side effects:

- hives
 - difficulty breathing
 - swelling of your face, lips, tongue, or throat. Wash the skin and get medical attention right away if you have severe burning, pain, swelling, or blistering of the skin where you applied Sloan's Liniment. Stop using Sloan's Liniment and call your doctor at once if you have: severe pain or skin redness where the medicine was applied
 - loss of feeling
 - trouble breathing or swallowing (after accidental inhalation of capsaicin odor or dried residue).
- Common side effects may include pain or a burning sensation that can last for several hours or days (especially after your first use of capsaicin topical).

2260. This particular disease, Osteoarthritis, interacts with the drug Supartz FX and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: More common Difficulty with moving muscle pain or stiffness pain in the joints Less common Swelling or redness in the joints

Drug Side Effects Summary

Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Bleeding, blistering, burning, coldness, discoloration of the skin, feeling of pressure, hives, infection, inflammation, itching, lumps, numbness, pain, rash, redness, scarring, soreness, stinging, swelling, tenderness, tingling, ulceration, or warmth at the injection site Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Supartz FX side effects

2261. This particular disease, Osteoarthritis, interacts with the drug tolmetin and has the following

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Tolmetin may cause serious side effects. Stop using tolmetin and call your doctor at once if you have: changes in your vision
- shortness of breath (even with mild exertion)
- swelling or rapid weight gain
- any skin rash, no matter how mild
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems-- nausea , upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- kidney problems--little or no urinating, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath
- or low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of

Drug Side Effects Summary

breath, cold hands and feet. Common side effects of tolmetin may include: indigestion , stomach pain, nausea, vomiting

- diarrhea , gas
- headache , dizziness
- weakness
- or changes in weight.

2262. This particular disease, Osteoarthritis, interacts with the drug trolamine salicylate and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Irritation where trolamine cream and lotion is used. Redness. Trolamine salicylate topical side effects

2263. This particular disease, Osteoarthritis, interacts with the drug Zostrix Diabetic Foot Pain and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if

Drug Side Effects Summary

you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Zostrix Diabetic Foot Pain side effects

2264. This particular disease, Osteoarthritis, interacts with the drug Zostrix Sports and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Zostrix Sports side effects

Drug Side Effects Summary

2265. This particular disease, Osteoporosis, interacts with the drug alendronate and has the following side effects:

- hives
- wheezing, difficulty breathing
- swelling of your face, lips, tongue, or throat. Alendronate may cause serious side effects. Stop using alendronate and call your doctor at once if you have: chest pain, new or worsening heartburn
- difficulty or pain when swallowing
- pain or burning under the ribs or in the back
- severe heartburn, burning pain in your upper stomach, or coughing up blood
- new or unusual pain in your thigh or hip
- jaw pain, numbness, or swelling
- severe joint, bone, or muscle pain
- or low calcium levels--muscle spasms or contractions, numbness or tingly feeling (around your mouth, or in your fingers and toes). Common side effects of alendronate may include: heartburn, upset stomach
- stomach pain, nausea
- diarrhea , constipation
- or bone pain, muscle or joint pain.

2266. This particular disease, Osteoporosis, interacts with the drug Prolia and has the following side effects:

- hives, itching, rash
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: new or unusual pain in your thigh, hip, or groin
- severe pain in your joints, muscles, or bones
- skin problems such as dryness, peeling, redness, itching, blisters, bumps, oozing, or crusting

Drug Side Effects Summary

- or low levels of calcium in your blood (hypocalcemia) - numbness or tingly feeling around your mouth or in your fingers or toes, muscle tightness or contraction, overactive reflexes. Serious infections may occur during treatment with Prolia. Call your doctor right away if you have signs of infection such as : fever, chills, night sweats
- swelling, pain, tenderness, warmth, or redness anywhere on your body
- pain or burning when you urinate
- increased or urgent need to urinate
- severe stomach pain
- or cough, feeling short of breath. Common Prolia side effects may include: bladder infection (painful or difficult urination)
- back pain, muscle pain
- or pain in your arms or legs.

2267. This particular disease, Osteoporosis, interacts with the drug Fosamax and has the following

- hives
- wheezing, difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using Fosamax and call your doctor at once if you have: chest pain, new or worsening heartburn
- difficulty or pain when swallowing
- pain or burning under the ribs or in the back
- severe heartburn, burning pain in your upper stomach, or coughing up blood
- new or unusual pain in your thigh, hip or groin
- jaw pain, numbness, or swelling
- severe joint, bone, or muscle pain
- or signs of low calcium levels - muscle spasms or contractions, numbness or tingly feeling (around

Drug Side Effects Summary

your mouth, or in your fingers and toes). Common Fosamax side effects may include: heartburn, upset stomach

- stomach pain, nausea
- diarrhea, constipation
- or headaches
- or bone pain, muscle or joint pain.

2268. This particular disease, Osteoporosis, interacts with the drug calcium / vitamin d and has the

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have signs of too much calcium in your body, such as: nausea , vomiting , constipation
- increased thirst or urination
- muscle weakness, bone pain
- or confusion, lack of energy, or feeling tired. Common side effects of calcium and vitamin D combination may include: an irregular heartbeat
- weakness, drowsiness, headache
- dry mouth, or a metallic taste in your mouth
- or muscle or bone pain.

2269. This particular disease, Osteoporosis, interacts with the drug denosumab and has the followi

- hives , itching, rash
- difficult breathing, feeling light-headed
- swelling of your face, lips, tongue, or throat. Denosumab may cause serious side effects. Call your doctor at once if you have: new or unusual pain in your thigh, hip, or groin

Drug Side Effects Summary

- severe pain in your joints, muscles, or bones
- skin problems such as dryness, peeling, redness, itching, blisters, bumps, oozing, or crusting
- or low calcium level--muscle spasms or contractions, numbness or tingly feeling (around your mouth, or in your fingers and toes). Serious infections may occur during treatment with Prolia. Call your doctor right away if you have signs of infection such as : fever, chills, night sweats
- swelling, pain, tenderness, warmth, or redness anywhere on your body
- pain or burning when you urinate
- increased or urgent need to urinate
- severe stomach pain
- or cough, wheezing, feeling short of breath. Common side effects of denosumab may include:
bladder infection (painful or difficult urination)
- lung infection (cough, shortness of breath)
- headache
- back pain , muscle pain, joint pain
- increased blood pressure
- cold symptoms such as stuffy nose, sneezing, sore throat
- high cholesterol
- or pain in your arms or legs.

2270. This particular disease, Osteoporosis, interacts with the drug Boniva and has the following side effects:

- hives
- wheezing, difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: chest pain, new or worsening heartburn
- difficulty or pain when swallowing

Drug Side Effects Summary

- pain or burning under the ribs or in the back
- severe heartburn, burning pain in your upper stomach, or coughing up blood
- new or unusual pain in your thigh or hip
- jaw pain, numbness, or swelling
- severe joint, bone, or muscle pain
- or low calcium levels - muscle spasms or contractions, numbness or tingly feeling (around your mouth, or in your fingers and toes). Common Boniva side effects may include: heartburn, stomach pain, diarrhea
- back pain, bone pain, muscle or joint pain
- pain in your arms or legs
- headache
- or fever, chills, tiredness, flu-like symptoms.

2271. This particular disease, Osteoporosis, interacts with the drug Actonel and has the following s

- chest pain, new or worsening heartburn
- difficulty or pain when swallowing
- pain or burning under the ribs or in the back
- severe heartburn, burning pain in your upper stomach, or coughing up blood
- new or unusual pain in your thigh or hip
- jaw pain, numbness, or swelling
- severe joint, bone, or muscle pain
- or low calcium levels - muscle spasms or contractions, numbness or tingly feeling (around your mouth, or in your fingers and toes). Common Actonel side effects may include: heartburn, diarrhea, indigestion
- stomach pain

Drug Side Effects Summary

- back pain, joint pain, muscle pain
- or flu-like symptoms.

2272. This particular disease, Osteoporosis, interacts with the drug Evenity and has the following side effects:

- chest pain or pressure, shortness of breath, feeling light-headed, sudden numbness or weakness, problems with vision or speech, or loss of balance. Tell your doctor if you have: new or unusual pain in your thigh, hip, or groin
- jaw pain or numbness
- red or swollen gums, loose teeth, infected gums
- or low calcium level - muscle spasms or contractions, numbness or tingly feeling (around your mouth, or in your fingers and toes). Common Evenity side effects may include: headache
- or joint pain.

2273. This particular disease, Osteoporosis, interacts with the drug Forteo and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: bone pain
- new or unusual swelling or lumps under your skin
- a light-headed feeling, like you might pass out (may occur within 4 hours after injection)
- pounding heartbeats or fluttering in your chest after using an injection
- or high calcium levels - confusion, muscle weakness, nausea, vomiting, lack of energy, constipation, increased thirst or urination, weight loss. Common Forteo side effects may include: nausea
- joint pain
- or pain anywhere in your body.

Drug Side Effects Summary

2274. This particular disease, Osteoporosis, interacts with the drug Reclast and has the following side effects:

- hives
- wheezing, chest tightness, trouble breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: new or unusual pain in your thigh or hip
- jaw pain or numbness, red or swollen gums, loose teeth, or slow healing after dental work
- severe joint, bone, or muscle pain
- kidney problems - little or no urination, swelling in your feet or ankles, feeling tired
- low red blood cells (anemia) - pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet
- or low calcium levels - muscle spasms or contractions, numbness or tingly feeling (around your mouth, or in your fingers and toes). Serious side effects on the kidneys may be more likely in older adults. Common Reclast side effects may include: trouble breathing
- nausea , vomiting , diarrhea , constipation
- bone pain, muscle or joint pain
- fever or other flu symptoms
- tiredness
- eye pain or swelling
- pain in your arms or legs
- headache
- or anemia.

2275. This particular disease, Osteoporosis, interacts with the drug Evista and has the following side effects:

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: swelling, tenderness, or other changes in your breasts
- signs of a stroke - sudden numbness or weakness (especially on one side of the body), slurred speech, vision problems
- signs of a blood clot in the lung - chest pain, trouble breathing, coughing up blood
- or signs of a blood clot deep in the body - swelling, warmth, or redness in an arm or leg. Common Evista side effects may include: hot flashes
- leg cramps
- swelling in your hands, feet, or ankles
- joint pain
- flu symptoms
- or increased sweating.

2276. This particular disease, Osteoporosis, interacts with the drug Caltrate 600+D and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have signs of too much calcium in your body, such as: nausea , vomiting , constipation
- increased thirst or urination
- muscle weakness, bone pain
- or confusion, lack of energy, or feeling tired. Common side effects of Caltrate 600+D Soft Chews may include: an irregular heartbeat
- weakness, drowsiness, headache
- dry mouth, or a metallic taste in your mouth
- or muscle or bone pain.

Drug Side Effects Summary

2277. This particular disease, Osteoporosis, interacts with the drug Citracal + D and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have signs of too much calcium in your body, such as: nausea , vomiting , constipation
- increased thirst or urination
- muscle weakness, bone pain
- or confusion, lack of energy, or feeling tired. Common side effects of Citracal + D may include: an irregular heartbeat
- weakness, drowsiness, headache
- dry mouth, or a metallic taste in your mouth
- or muscle or bone pain.

2278. This particular disease, Osteoporosis, interacts with the drug risedronate and has the following side effects:

- chest pain, new or worsening heartburn
- difficulty or pain when swallowing
- pain or burning under the ribs or in the back
- severe heartburn, burning pain in your upper stomach, or coughing up blood
- new or unusual pain in your thigh or hip
- jaw pain, numbness, or swelling
- severe joint, bone, or muscle pain
- or low calcium levels--muscle spasms or contractions, numbness or tingly feeling (around your mouth, or in your fingers and toes). Common side effects of risedronate may include: heartburn, diarrhea , indigestion
- stomach pain

Drug Side Effects Summary

- back pain , joint pain, muscle pain
- or flu-like symptoms.

2279. This particular disease, Osteoporosis, interacts with the drug Caltrate and has the following s

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad constipation . Caltrate side effects

2280. This particular disease, Osteoporosis, interacts with the drug calcium citrate and has the follo

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Calcium citrate may cause serious side effects. Call your doctor at once if you have: little or no urination
- swelling, rapid weight gain
- or high levels of calcium in your blood-- nausea , vomiting , constipation , increased thirst or urination, muscle weakness, bone pain, confusion, lack of energy, or feeling tired. Common side

Drug Side Effects Summary

effects of calcium citrate may include: upset stomach

- or constipation.

2281. This particular disease, Osteoporosis, interacts with the drug ibandronate and has the following

- hives
- wheezing, difficult breathing
- swelling of your face, lips, tongue, or throat. Ibandronate may cause serious side effects. Stop using ibandronate and call your doctor at once if you have: chest pain, new or worsening heartburn
- difficulty or pain when swallowing
- pain or burning under the ribs or in the back
- severe heartburn, burning pain in your upper stomach, or coughing up blood
- new or unusual pain in your thigh or hip
- jaw pain, numbness, or swelling
- severe joint, bone, or muscle pain
- or low calcium levels--muscle spasms or contractions, numbness or tingly feeling (around your mouth, or in your fingers and toes). Common side effects of ibandronate may include: heartburn, stomach pain, diarrhea
- back pain , bone pain, muscle or joint pain
- pain in your arms or legs
- headache
- or fever, chills, tiredness, flu-like symptoms.

2282. This particular disease, Osteoporosis, interacts with the drug raloxifene and has the following

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Raloxifene may cause serious side effects. Stop using raloxifene and call your doctor at once if you have: swelling, tenderness, or other changes in your breasts
- signs of a stroke--sudden numbness or weakness (especially on one side of the body), slurred speech, vision problems
- signs of a blood clot in the lung--chest pain, trouble breathing, coughing up blood
- or signs of a blood clot deep in the body--swelling, warmth, or redness in an arm or leg. Common side effects of raloxifene may include: hot flashes
- leg cramps
- swelling in your hands, feet, or ankles
- joint pain
- flu symptoms
- or increased sweating.

2283. This particular disease, Osteoporosis, interacts with the drug Premarin and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: heart attack symptoms - chest pain or pressure, pain spreading to your jaw or shoulder, sweating
- signs of a stroke - sudden numbness or weakness (especially on one side of the body), sudden severe headache , slurred speech, problems with vision or balance
- signs of a blood clot - sudden vision loss, stabbing chest pain, coughing up blood , pain or warmth in one or both legs
- swelling, rapid weight gain
- jaundice (yellowing of the skin or eyes)

Drug Side Effects Summary

- memory problems, confusion, unusual behavior
- unusual vaginal bleeding, pelvic pain
- a lump in your breast
- or high levels of calcium in your blood - vomiting , constipation , increased thirst or urination, muscle weakness, bone pain, lack of energy. Common Premarin side effects may include: swelling
- hair loss
- numbness, tingling, burning pain
- back pain, leg cramps, pain
- bloating, gas, indigestion, nausea, vomiting, stomach pain
- dizziness, headache
- breast pain
- or vaginal itching or discharge, changes in your menstrual periods, breakthrough bleeding .

2284. This particular disease, Osteoporosis, interacts with the drug Atelvia and has the following side effects:

- chest pain, new or worsening heartburn
- difficulty or pain when swallowing
- pain or burning under the ribs or in the back
- severe heartburn, burning pain in your upper stomach, or coughing up blood
- new or unusual pain in your thigh or hip
- jaw pain, numbness, or swelling
- severe joint, bone, or muscle pain
- or low calcium levels - muscle spasms or contractions, numbness or tingly feeling (around your mouth, or in your fingers and toes). Common Atelvia side effects may include: heartburn, diarrhea, indigestion
- stomach pain

Drug Side Effects Summary

- back pain, joint pain, muscle pain
- or flu-like symptoms.

2285. This particular disease, Osteoporosis, interacts with the drug calcitonin and has the following

- You may have increased bone pain during the first few months of treatment with calcitonin injection. This is not a sign that the medication isn't working properly. hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Calcitonin injection may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- or low calcium levels--muscle spasms or contractions, numbness or tingly feeling (around your mouth, or in your fingers and toes). Common side effects of calcitonin injection may include: flushing (warmth, redness, or tingly feeling)
- nausea , vomiting
- or swelling where the injection was given.

2286. This particular disease, Osteoporosis, interacts with the drug Zometa and has the following s

- hives
- wheezing, chest tightness, trouble breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: new or unusual pain in your thigh or hip
- jaw pain or numbness, red or swollen gums, loose teeth, or slow healing after dental work
- severe joint, bone, or muscle pain
- kidney problems - little or no urination, swelling in your feet or ankles, feeling tired
- low red blood cells (anemia) - pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet

Drug Side Effects Summary

- or low calcium levels - muscle spasms or contractions, numbness or tingling feeling (around your mouth, or in your fingers and toes). Serious side effects on the kidneys may be more likely in older adults. Common Zometa side effects may include: trouble breathing
- nausea , vomiting , diarrhea , constipation
- bone pain, muscle or joint pain
- fever or other flu symptoms
- tiredness
- eye pain or swelling
- pain in your arms or legs
- headache
- or anemia.

2287. This particular disease, Osteoporosis, interacts with the drug teriparatide and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Teriparatide may cause serious side effects. Call your doctor at once if you have: bone pain
- new or unusual swelling or lumps under your skin
- a light-headed feeling, like you might pass out (may occur within 4 hours after injection)
- pounding heartbeats or fluttering in your chest after using an injection
- or high calcium levels--confusion, muscle weakness, nausea, vomiting, lack of energy, constipation , increased thirst or urination, weight loss . Common side effects of teriparatide may include: nausea
- joint pain
- or pain anywhere in your body.

Drug Side Effects Summary

2288. This particular disease, Osteoporosis, interacts with the drug zoledronic acid and has the following side effects:

- hives
- wheezing, chest tightness, trouble breathing
- swelling of your face, lips, tongue, or throat. Zoledronic acid may cause serious side effects. Call your doctor at once if you have: new or unusual pain in your thigh or hip
- jaw pain or numbness, red or swollen gums, loose teeth, or slow healing after dental work
- severe joint, bone, or muscle pain
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired
- low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet
- or low calcium levels--muscle spasms or contractions, numbness or tingly feeling (around your mouth, or in your fingers and toes). Serious side effects on the kidneys may be more likely in older adults. Common side effects of zoledronic acid may include: trouble breathing
- nausea , vomiting , diarrhea , constipation
- bone pain, muscle or joint pain
- fever or other flu symptoms
- tiredness
- eye pain or swelling
- pain in your arms or legs
- headache
- or anemia.

2289. This particular disease, Osteoporosis, interacts with the drug alendronate / cholecalciferol and has the following side effects:

- hives
- difficulty breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Alendronate and cholecalciferol may cause serious side effects. Stop using alendronate and cholecalciferol and call your doctor at once if you have: chest pain, new or worsening heartburn
- difficulty or pain when swallowing
- pain or burning under the ribs or in the back
- new or worsening heartburn
- severe joint, bone, or muscle pain
- new or unusual pain in your thigh or hip
- or jaw pain, numbness, or swelling. Common side effects of alendronate and cholecalciferol may include: mild heartburn, stomach upset
- diarrhea , gas, constipation
- mild joint or back pain
- or headache .

2290. This particular disease, Osteoporosis, interacts with the drug Fosamax Plus D and has the fo

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using Fosamax Plus D and call your doctor at once if you have: chest pain, new or worsening heartburn
- difficulty or pain when swallowing
- pain or burning under the ribs or in the back
- new or worsening heartburn
- severe joint, bone, or muscle pain
- new or unusual pain in your thigh or hip
- or jaw pain, numbness, or swelling. Common Fosamax Plus D side effects may include: mild

Drug Side Effects Summary

heartburn, stomach upset

- diarrhea, gas, constipation
- mild joint or back pain
- or headache.

2291. This particular disease, Osteoporosis, interacts with the drug Binosto and has the following side effects:

- hives
- wheezing, difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using Binosto and call your doctor at once if you have: chest pain, new or worsening heartburn
- difficulty or pain when swallowing
- pain or burning under the ribs or in the back
- severe heartburn, burning pain in your upper stomach, or coughing up blood
- new or unusual pain in your thigh or hip
- jaw pain, numbness, or swelling
- severe joint, bone, or muscle pain
- or low calcium levels - muscle spasms or contractions, numbness or tingly feeling (around your mouth, or in your fingers and toes). Common Binosto side effects may include: heartburn, upset stomach
- stomach pain, nausea
- diarrhea, constipation
- or bone pain, muscle or joint pain.

2292. This particular disease, Osteoporosis, interacts with the drug Miacalcin and has the following side effects:

- hives

Drug Side Effects Summary

- fast heartbeats
- feeling lightheaded
- chest pain, difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- severe nasal irritation
- or low levels of calcium in your blood--numbness or tingly feeling around your mouth, fast or slow heart rate, muscle tightness or contraction, overactive reflexes. Common Miacalcin side effects may include: bleeding from your nose
- runny or stuffy nose
- dryness, itching, tenderness, or general discomfort of your nose
- or crusting, scabs, or sores inside your nose.

2293. This particular disease, Osteoporosis, interacts with the drug Tymlos and has the following s

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: a light-headed feeling, like you might pass out (may occur within 4 hours after injection)
- bone pain
- unusual body aches
- a new or worsening lump or swelling under your skin
- pounding heartbeats or fluttering in your chest
- high calcium levels - confusion, tiredness, nausea , vomiting, loss of appetite, constipation , increased thirst or urination, weight loss
- or symptoms of a kidney stone - severe pain in your side or lower back, blood in your urine, painful

Drug Side Effects Summary

urination. Common Tymlos side effects may include: dizziness, spinning sensation

- headache
- fast heart rate
- nausea, upper stomach pain
- or feeling very tired.

2294. This particular disease, Osteoporosis, interacts with the drug Aclasta and has the following side effects:

- urinating less than usual or not at all
- muscle spasms, numb or tingly feeling (especially around your mouth)
- fever, chills, body aches, flu symptoms
- pale skin, easy bruising, unusual weakness
- or severe joint, bone, or muscle pain. Less serious Aclasta side effects may include: cough
- loss of appetite, nausea, vomiting
- diarrhea, constipation
- headache, dizziness, tired feeling
- low blood pressure, swelling in your legs or feet
- bone, joint or muscle pain
- or redness or swelling where the needle was placed.

2295. This particular disease, Osteoporosis, interacts with the drug Prempro and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: heart attack symptoms - chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating
- signs of a stroke - sudden numbness or weakness (especially on one side of the body), sudden

Drug Side Effects Summary

severe headache , slurred speech, problems with vision or balance

- signs of a blood clot - sudden vision loss, stabbing chest pain, feeling short of breath, coughing up

blood , pain or warmth in one or both legs

- swelling or tenderness in your stomach

- jaundice (yellowing of the skin or eyes)

- memory problems, confusion, unusual behavior

- unusual vaginal bleeding, pelvic pain

- a lump in your breast

- or high levels of calcium in your blood - nausea, vomiting , constipation , increased thirst or urination, muscle weakness, bone pain, lack of energy. Common Prempro side effects may include:

nausea, gas, stomach pain

- headache, back pain

- depression , weakness

- breast pain

- itching, thinning scalp hair

- or vaginal itching or discharge, changes in your menstrual periods, breakthrough bleeding .

2296. This particular disease, Osteoporosis, interacts with the drug abaloparatide and has the following side effects:

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Abaloparatide may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out (may occur within 4 hours after injection)

- bone pain

- unusual body aches

Drug Side Effects Summary

- a new or worsening lump or swelling under your skin
- pounding heartbeats or fluttering in your chest
- high calcium levels--confusion, tiredness, nausea , vomiting, loss of appetite, constipation , increased thirst or urination, weight loss
- or symptoms of a kidney stone--severe pain in your side or lower back, blood in your urine, painful urination. Common side effects of abaloparatide may include: dizziness, spinning sensation
- headache
- fast heart rate
- nausea, upper stomach pain
- or feeling very tired.

2297. This particular disease, Osteoporosis, interacts with the drug Calcitrate with D and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have signs of too much calcium in your body, such as: nausea , vomiting , constipation
- increased thirst or urination
- muscle weakness, bone pain
- or confusion, lack of energy, or feeling tired. Common side effects of Calcitrate with D may include: an irregular heartbeat
- weakness, drowsiness, headache
- dry mouth, or a metallic taste in your mouth
- or muscle or bone pain.

2298. This particular disease, Osteoporosis, interacts with the drug calcium gluconate and has the following side effects:

Drug Side Effects Summary

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of high calcium levels like weakness, confusion, feeling tired, headache , upset stomach and throwing up, constipation , or bone pain.

Calcium gluconate side effects

2299. This particular disease, Osteoporosis, interacts with the drug calcium lactate and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Calcium lactate may cause serious side effects. Stop using calcium lactate and call your doctor at once if you have: severe nausea, vomiting, or constipation
- or high blood calcium--confusion, tiredness, loss of appetite, increased thirst or urination, weight loss . Common side effects of calcium lactate may include: stomach pain, gas
- or constipation.

Drug Side Effects Summary

2300. This particular disease, Osteoporosis, interacts with the drug calcium phosphate, tribasic and

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Calcium phosphate may cause serious side effects. Stop using calcium phosphate and call your doctor at once if you have: severe nausea, vomiting, or constipation
- or high blood calcium--confusion, tiredness, loss of appetite, increased thirst or urination, weight loss
- or symptoms of a kidney stone--severe pain in your side or lower back, blood in your urine, painful urination. Less serious side effects may include: nausea, stomach bloating
- constipation
- or gas.

2301. This particular disease, Osteoporosis, interacts with the drug conjugated estrogens and has t

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Conjugated estrogens may cause serious side effects. Call your doctor at once if you have: heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, sweating
- signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache , slurred speech, problems with vision or balance
- signs of a blood clot--sudden vision loss, stabbing chest pain, coughing up blood , pain or warmth in one or both legs
- swelling, rapid weight gain
- jaundice (yellowing of the skin or eyes)

Drug Side Effects Summary

- memory problems, confusion, unusual behavior
- unusual vaginal bleeding, pelvic pain
- a lump in your breast
- or high levels of calcium in your blood-- vomiting , constipation , increased thirst or urination, muscle weakness, bone pain, lack of energy. Common side effects of conjugated estrogens may include: swelling
- hair loss
- numbness, tingling, burning pain
- back pain , leg cramps, pain
- bloating , gas, indigestion , nausea , vomiting, stomach pain
- dizziness , headache
- breast pain
- or vaginal itching or discharge, changes in your menstrual periods, breakthrough bleeding .

2302. This particular disease, Osteoporosis, interacts with the drug Os-Cal Calcium+D3 and has the

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking

Drug Side Effects Summary

- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad constipation . Os-Cal Calcium+D3 side effects

2303. This particular disease, Osteoporosis, interacts with the drug bazedoxifene / conjugated estrogens

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Bazedoxifene and conjugated estrogens may cause serious side effects. Call your doctor at once if you have: heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating
- signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache , slurred speech, problems with vision or balance
- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood , pain or warmth in one or both legs
- sudden vision loss
- swelling, rapid weight gain
- jaundice (yellowing of the skin or eyes)
- memory problems, confusion, unusual behavior
- unusual vaginal bleeding, pelvic pain
- a lump in your breast
- or low calcium level--muscle spasms or contractions, numbness or tingly feeling (around your mouth, or in your fingers and toes). Common side effects of bazedoxifene and conjugated estrogens may include: dizziness
- nausea, stomach pain or discomfort, diarrhea
- muscle spasm

Drug Side Effects Summary

- or throat pain, neck pain .

2304. This particular disease, Osteoporosis, interacts with the drug Cal-GLU and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of high calcium levels like weakness, confusion, feeling tired, headache , upset stomach and throwing up, constipation , or bone pain.

Cal-GLU side effects

2305. This particular disease, Osteoporosis, interacts with the drug Cal-Lac and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching

Drug Side Effects Summary

- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad constipation . Very upset stomach or throwing up. Feeling confused. Cal-Lac side effects

2306. This particular disease, Osteoporosis, interacts with the drug Calcitrate and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of high calcium levels like weakness, confusion, feeling tired, headache , upset stomach and throwing up, constipation , or bone pain. Mood changes. Calcitrate side effects

2307. This particular disease, Osteoporosis, interacts with the drug conjugated estrogens / medrox

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Conjugated estrogens and medroxyprogesterone may cause serious side effects. Call your doctor at once if you have: heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating
- signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache , slurred speech, problems with vision or balance
- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood , pain or warmth in one or both legs
- swelling or tenderness in your stomach
- jaundice (yellowing of the skin or eyes)
- memory problems, confusion, unusual behavior
- unusual vaginal bleeding, pelvic pain
- a lump in your breast
- or high levels of calcium in your blood--nausea, vomiting , constipation , increased thirst or urination, muscle weakness, bone pain, lack of energy. Common side effects of conjugated estrogens and medroxyprogesterone may include: nausea, gas, stomach pain
- headache, back pain
- depression , weakness
- breast pain
- itching, thinning scalp hair
- or vaginal itching or discharge, changes in your menstrual periods, breakthrough bleeding .

2308. This particular disease, Osteoporosis, interacts with the drug Duavee and has the following s

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: heart attack symptoms - chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating
- signs of a stroke - sudden numbness or weakness (especially on one side of the body), sudden severe headache , slurred speech, problems with vision or balance
- signs of a blood clot - sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood , pain or warmth in one or both legs
- sudden vision loss
- swelling, rapid weight gain
- jaundice (yellowing of the skin or eyes)
- memory problems, confusion, unusual behavior
- unusual vaginal bleeding, pelvic pain
- a lump in your breast
- or low calcium level - muscle spasms or contractions, numbness or tingly feeling (around your mouth, or in your fingers and toes). Common Duavee side effects may include: dizziness
- nausea, stomach pain or discomfort, diarrhea
- muscle spasm
- or throat pain, neck pain .

2309. This particular disease, Osteoporosis, interacts with the drug esterified estrogens and has the

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Esterified estrogens may cause serious side effects. Stop using esterified estrogens and call your doctor at once if you have: signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache , slurred

Drug Side Effects Summary

speech, problems with vision or balance

- signs of a blood clot in the lung--chest pain, sudden cough, wheezing, rapid breathing, coughing up blood
- signs of a blood clot in your leg--pain, swelling, warmth, or redness in one or both legs
- heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating
- liver problems--severe stomach pain, fever, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- high levels of calcium in your blood--nausea, vomiting , constipation , increased thirst or urination, muscle weakness, bone pain, confusion, lack of energy, or tired feeling
- a change in the pattern or severity of migraine headaches
- swelling in your hands, ankles, or feet
- a breast lump
- or severe skin reaction--fever, sore throat , swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of esterified estrogens may include: light vaginal bleeding or spotting
- breast pain or tenderness
- nausea, vomiting, bloating
- skin color changes, increased facial hair, thinning scalp hair
- headache, dizziness , mood changes, decreased sex drive
- vaginal itching or discharge, very light menstrual periods
- or problems with contact lenses.

2310. This particular disease, Osteoporosis, interacts with the drug estropipate and has the following

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Estropipate may cause serious side effects. Call your doctor at once if you have: heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating
- signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache , slurred speech, problems with vision or balance
- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up blood , pain or warmth in one or both legs
- swelling or tenderness in your stomach
- jaundice (yellowing of the skin or eyes)
- memory problems, confusion, unusual behavior
- unusual vaginal bleeding, pelvic pain
- a lump in your breast
- or high levels of calcium in your blood--nausea, vomiting , constipation , increased thirst or urination, muscle weakness, bone pain, lack of energy. Common side effects of estropipate may include: headache
- vaginal itching or discharge, changes in your menstrual periods, breakthrough bleeding
- bloating , stomach cramps, nausea, vomiting
- breast pain
- thinning scalp hair
- or fluid retention (swelling, rapid weight gain).

2311. This particular disease, Osteoporosis, interacts with the drug Menest and has the following s

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and

Drug Side Effects Summary

sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of gallbladder problems like pain in the upper right belly area, right shoulder area, or between the shoulder blades
- yellow skin or eyes
- fever with chills
- bloating
- or very upset stomach or throwing up. Signs of high blood pressure like very bad headache or dizziness , passing out, or change in eyesight. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of a pancreas problem (pancreatitis) like very bad stomach pain, very bad back pain , or very bad upset stomach or throwing up. Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight. Dizziness or passing out. Change in eyesight. Loss of eyesight. Bulging eyes. Change in how contact lenses feel in the eyes. A lump in the breast, breast pain or soreness, or nipple discharge. Breast pain. Vaginal itching or discharge. Vaginal bleeding that is not normal. Low mood (depression). Mood changes. Memory problems or loss. Swelling. Call your doctor right away if you have signs of a blood clot like chest

Drug Side Effects Summary

pain or pressure

- coughing up blood
- shortness of breath
- swelling, warmth, numbness , change of color, or pain in a leg or arm
- or trouble speaking or swallowing. Menest side effects

2312. This particular disease, Osteoporosis, interacts with the drug Ogen and has the following side effects

- No side effects listed

2313. This particular disease, Osteoporosis, interacts with the drug Os-Cal 500 Chewable and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad constipation . Os-Cal 500 Chewable side effects

2314. This particular disease, Osteoporosis, interacts with the drug Os-Cal Extra D3 and has the following side effects

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have signs of too much calcium in your body, such as: nausea , vomiting , constipation
- increased thirst or urination
- muscle weakness, bone pain
- or confusion, lack of energy, or feeling tired. Common side effects of Os-Cal Extra D3 may include: an irregular heartbeat
- weakness, drowsiness, headache
- dry mouth, or a metallic taste in your mouth
- or muscle or bone pain.

2315. This particular disease, Osteoporosis, interacts with the drug Oystercal-D and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have signs of too much calcium in your body, such as: nausea , vomiting , constipation
- increased thirst or urination
- muscle weakness, bone pain
- or confusion, lack of energy, or feeling tired. Common side effects of Oystercal-D may include: an irregular heartbeat
- weakness, drowsiness, headache
- dry mouth, or a metallic taste in your mouth
- or muscle or bone pain.

Drug Side Effects Summary

2316. This particular disease, Osteoporosis, interacts with the drug Posture and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Calcium phosphate may cause serious side effects. Stop using calcium phosphate and call your doctor at once if you have: severe nausea, vomiting, or constipation
- or high blood calcium--confusion, tiredness, loss of appetite, increased thirst or urination, weight loss
- or symptoms of a kidney stone--severe pain in your side or lower back, blood in your urine, painful urination. Less serious side effects may include: nausea, stomach bloating
- constipation
- or gas.

2317. This particular disease, Osteoporosis, interacts with the drug Premarin Intravenous and has the following side effects:

- No side effects listed

2318. This particular disease, Osteoporosis, interacts with the drug Premphase and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Premphase may cause serious side effects. Call your doctor at once if you have: heart attack symptoms--chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating
- signs of a stroke--sudden numbness or weakness (especially on one side of the body), sudden severe headache , slurred speech, problems with vision or balance
- signs of a blood clot--sudden vision loss, stabbing chest pain, feeling short of breath, coughing up

Drug Side Effects Summary

blood , pain or warmth in one or both legs

- swelling or tenderness in your stomach
- jaundice (yellowing of the skin or eyes)
- memory problems, confusion, unusual behavior
- unusual vaginal bleeding, pelvic pain
- a lump in your breast
- or high levels of calcium in your blood--nausea, vomiting , constipation , increased thirst or urination, muscle weakness, bone pain, lack of energy. Common side effects of Premphase may include: nausea, gas, stomach pain
- headache, back pain
- depression , weakness
- breast pain
- itching, thinning scalp hair
- or vaginal itching or discharge, changes in your menstrual periods, breakthrough bleeding .

2319. This particular disease, Osteoporosis, interacts with the drug romosozumab and has the following

- chest pain or pressure, shortness of breath, feeling light-headed, sudden numbness or weakness, problems with vision or speech, or loss of balance. Romosozumab may cause serious side effects. Call your doctor at once if you have: new or unusual pain in your thigh, hip, or groin
- jaw pain or numbness
- red or swollen gums, loose teeth, infected gums
- or low calcium level--muscle spasms or contractions, numbness or tingly feeling (around your mouth, or in your fingers and toes). Common side effects of romosozumab may include: headache
- or joint pain.

Drug Side Effects Summary

2320. This particular disease, Osteoporosis, interacts with the drug UPCal D and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have signs of too much calcium in your body, such as: nausea , vomiting , constipation
- increased thirst or urination
- muscle weakness, bone pain
- or confusion, lack of energy, or feeling tired. Common side effects of UpCal D may include: an irregular heartbeat
- weakness, drowsiness, headache
- dry mouth, or a metallic taste in your mouth
- or muscle or bone pain.

2321. This particular disease, Pain, interacts with the drug oxycodone and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine, including oxycodone, can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep
- a slow heart rate or weak pulse
- cold, clammy skin
- a light-headed feeling, like you might pass out
- confusion, unusual thoughts or behavior

Drug Side Effects Summary

- seizure (convulsions)
- low cortisol levels - nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness
- or high levels of serotonin in the body - agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, diarrhea. Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders. Long-term use of opioid medication, such as oxycodone, may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent. Common oxycodone side effects may include: drowsiness, dizziness, tiredness
- headache
- constipation , stomach pain, nausea, vomiting
- or itching, red eyes, or flushing.

2322. This particular disease, Pain, interacts with the drug acetaminophen and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal, even if you took acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Stop taking acetaminophen and call your doctor at once if you have signs of liver problems: stomach pain (upper right side)
- loss of appetite
- tiredness, itching
- dark urine, clay-colored stools
- or jaundice (yellowing of the skin or eyes). Less serious acetaminophen side effects may be more

Drug Side Effects Summary

likely, and you may have none at all.

2323. This particular disease, Pain, interacts with the drug hydrocodone and has the following side

- hives
 - difficult breathing
 - swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Stop using hydrocodone and call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep
 - a slow heart rate or weak pulse
 - pain or burning when you urinate
 - confusion, tremors, severe drowsiness
 - a light-headed feeling, like you might pass out
 - low cortisol levels - nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness
 - or high levels of serotonin in the body - agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, diarrhea. Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea .
- Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent. Common hydrocodone side effects may include: constipation , nausea, vomiting
- dizziness, drowsiness, feeling tired

Drug Side Effects Summary

- headache
- or cold symptoms such as stuffy nose, sneezing, sore throat .

2324. This particular disease, Pain, interacts with the drug Tylenol and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal, even if you took Tylenol in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Stop taking this medicine and call your doctor at once if you have signs of liver problems: stomach pain (upper right side)
- loss of appetite
- tiredness, itching
- dark urine, clay-colored stools
- or jaundice (yellowing of the skin or eyes). Less serious Tylenol side effects may be more likely, and you may have none at all.

2325. This particular disease, Pain, interacts with the drug methadone and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Call your doctor at once if you have: weak or shallow breathing, breathing that stops during sleep

Drug Side Effects Summary

- severe constipation
 - a light-headed feeling, like you might pass out
 - fast or pounding heartbeats, fluttering in your chest, shortness of breath
 - low cortisol levels-- nausea , vomiting , loss of appetite, dizziness, worsening tiredness or weakness
 - or serotonin syndrome - agitation , hallucinations, fever, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, diarrhea . Serious breathing problems may be more likely in older adults and people who are debilitated or have wasting syndrome or chronic breathing disorders.
- Common methadone side effects may include: dizziness, drowsiness
- nausea, vomiting
 - increased sweating
 - or pain, redness, or swelling where the medicine was injected.

2326. This particular disease, Pain, interacts with the drug acetaminophen / hydrocodone and has t

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep
- a light-headed feeling, like you might pass out

Drug Side Effects Summary

- liver problems - nausea, upper stomach pain, tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- low cortisol levels - nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness
- or high levels of serotonin in the body - agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, diarrhea. Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders. Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent. Common acetaminophen or hydrocodone side effects include: dizziness, drowsiness, feeling tired
- nausea, vomiting, stomach pain
- constipation
- or headache .

2327. This particular disease, Pain, interacts with the drug Dilaudid and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Like other narcotic medicines, hydromorphone can slow your breathing. Death may occur if breathing becomes too weak. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep
- confusion, feelings of extreme happiness or sadness
- severe weakness or drowsiness
- a light-headed feeling, like you might pass out
- infertility, missed menstrual periods
- impotence, sexual problems, loss of interest in sex

Drug Side Effects Summary

- or low cortisol levels - nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Hydromorphone is more likely to cause breathing problems in older adults and people who are severely ill, malnourished, or otherwise debilitated.

Common Dilaudid side effects may include: constipation, nausea, vomiting, stomach pain

- dizziness, drowsiness
- headache, tired feeling
- feelings of extreme happiness or sadness
- sweating, mild itching
- dry mouth
- or flushing (warmth, redness, or tingly feeling).

2328. This particular disease, Pain, interacts with the drug buprenorphine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Call your doctor at once if you have: opioid withdrawal symptoms - shivering, goose bumps, increased sweating, feeling hot or cold, runny nose , watery eyes , diarrhea , muscle pain
- noisy breathing, sighing, shallow breathing, breathing that stops during sleep
- slow heartbeat or weak pulse
- a light-headed feeling, like you might pass out
- chest pain, trouble breathing

Drug Side Effects Summary

- low cortisol levels - nausea , vomiting , loss of appetite, dizziness, worsening tiredness or weakness
- or liver problems - nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Common buprenorphine side effects may be more likely to occur, such as: constipation , nausea, vomiting
- headache
- increased sweating
- sleep problems (insomnia)
- or pain anywhere in your body.

2329. This particular disease, Pain, interacts with the drug OxyContin and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep
- a slow heart rate or weak pulse
- a light-headed feeling, like you might pass out
- confusion, unusual thoughts or behavior
- seizure (convulsions)
- low cortisol levels - nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness

Drug Side Effects Summary

- or high levels of serotonin in the body - agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, diarrhea. Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders. Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent. Common OxyContin side effects may include: drowsiness, headache , dizziness, tiredness
- or constipation , stomach pain, nausea, vomiting.

2330. This particular disease, Pain, interacts with the drug Buprenex and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep
- slow heartbeat or weak pulse
- blue lips or fingernails
- severe constipation
- confusion, feelings of extreme happiness
- little or no urination
- or low cortisol levels - nausea , vomiting , loss of appetite, dizziness, worsening tiredness or weakness. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss

Drug Side Effects Summary

of coordination, nausea, vomiting, or diarrhea . Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent. Common Buprenex side effects may include: drowsiness

- constipation
- dizziness, spinning sensation
- nausea, vomiting
- increased sweating
- headache
- or blurred vision, double vision.

2331. This particular disease, Pain, interacts with the drug Percocet and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen or Tylenol in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Oxycodone can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep
- slow heartbeat or weak pulse
- cold, clammy skin
- a light-headed feeling, like you might pass out

Drug Side Effects Summary

- weakness, tiredness, fever, unusual bruising or bleeding
- confusion, unusual thoughts or behavior
- seizure (convulsions)
- problems with urination
- signs of liver problems including nausea, upper stomach pain, tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness
- or high levels of serotonin in the body - agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, diarrhea. Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders. Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent. Common Percocet side effects include: dizziness, drowsiness, feeling tired
- headache
- constipation , nausea, vomiting, stomach pain
- blurred vision
- itching, red eyes, or flushing
- feelings of extreme happiness or sadness
- or dry mouth

2332. This particular disease, Pain, interacts with the drug Paracetamol and has the following side effects

- Serious side effects of paracetamol include: Allergic reactions, which may be severe and include: Skin rashes, itching or hives Swelling of the throat, tongue or face Shortness of breath or wheezing Skin rash or peeling, or mouth ulcers Breathing problems. This is more likely if you have experienced them before when taking other painkillers such as ibuprofen and aspirin Unexplained

Drug Side Effects Summary

bruising or bleeding or becoming unusually tired. Getting more infections than usual. Liver problems. Nausea, sudden weight loss, loss of appetite and yellowing of the eyes and skin can occur. If you get any of the rare side effects listed above, stop taking the medicine and contact your doctor immediately. Common side effect of paracetamol suppositories include: Redness or soreness in or around the rectum. Everyday, long-term use (several months or more) of paracetamol can cause liver or kidney damage. People taking this medication in the usual way for shorter periods of time have not had these problems. These are not all of the possible side effects of this medication. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Paracetamol side effects

2333. This particular disease, Pain, interacts with the drug ketorolac and has the following side effects

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. Ketorolac may cause serious side effects. Stop using ketorolac and call your doctor at once if you have: any skin rash, no matter how mild
- high blood pressure--severe headache , blurred vision, pounding in your neck or ears
- heart problems--swelling, rapid weight gain, feeling short of breath
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- liver problems-- nausea , upper stomach pain, itching, tiredness, flu-like symptoms, dark urine, jaundice (yellowing of the skin or eyes)
- low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet
- or signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Common side effects of ketorolac may include: heartburn , stomach pain, gas, nausea, vomiting
- diarrhea , constipation

Drug Side Effects Summary

- slow heartbeats
- decreased urination
- abnormal liver function tests
- increased blood pressure
- pain or irritation in your nose
- runny nose
- watery eyes
- throat irritation
- or rash.

2334. This particular disease, Pain, interacts with the drug Ultram and has the following side effects

- noisy breathing, sighing, shallow breathing, breathing that stops during sleep
- a slow heart rate or weak pulse
- a light-headed feeling, like you might pass out
- seizure (convulsions)
- or low cortisol levels - nausea , vomiting , loss of appetite, dizziness, worsening tiredness or weakness. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea . Serious breathing problems may be more likely in older adults and people who are debilitated or have wasting syndrome or chronic breathing disorders. Common Ultram side effects may include: constipation , nausea, vomiting, stomach pain
- dizziness, drowsiness, tiredness
- headache
- or itching.

Drug Side Effects Summary

2335. This particular disease, Pain, interacts with the drug acetaminophen / oxycodone and has the

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing
- a light-headed feeling, like you might pass out
- weakness, tiredness, fever, unusual bruising or bleeding
- confusion, unusual thoughts or behavior
- problems with urination
- liver problems - nausea, upper stomach pain, tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or low cortisol levels - nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea . Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent. Common acetaminophen and oxycodone side effects include: dizziness, drowsiness, feeling tired
- feelings of extreme happiness or sadness

Drug Side Effects Summary

- nausea, vomiting, stomach pain
- constipation
- or headache .

2336. This particular disease, Pain, interacts with the drug Demerol and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Stop using this medicine and call your doctor at once if you have: slow heartbeats, weak or shallow breathing, breathing that stops during sleep
- severe drowsiness, feeling like you might pass out
- confusion, mood changes
- severe constipation
- tremors, muscle movements you cannot control, or a seizure (convulsions)
- or low cortisol levels - nausea , vomiting , loss of appetite, dizziness, worsening tiredness or weakness. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea . Serious side effects may be more likely in older adults and those who are malnourished or debilitated. Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent. Common Demerol side effects may include: dizziness, drowsiness
- headache
- nausea, vomiting

Drug Side Effects Summary

- or sweating.

2337. This particular disease, Pain, interacts with the drug Nucynta and has the following side effects

- hives
- chest pain, fast heartbeats, difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops
- a light-headed feeling, like you might pass out
- agitation , feeling hot
- severe drowsiness or dizziness, confusion, problems with speech or balance
- a seizure
- serotonin syndrome - agitation, hallucinations, fever, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , diarrhea
- or low cortisol levels - nausea, vomiting , loss of appetite, dizziness, worsening tiredness or weakness. Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders. Common Nucynta side effects may include: constipation, nausea, vomiting, stomach pain
- headache , feeling tired
- or drowsiness, dizziness.

2338. This particular disease, Pain, interacts with the drug hydromorphone and has the following side effects

- hives

Drug Side Effects Summary

- difficulty breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Hydromorphone may cause serious side effects. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep
- a slow heart rate or weak pulse
- confusion, feelings of extreme happiness or sadness
- severe weakness or drowsiness
- a light-headed feeling, like you might pass out
- low cortisol levels-- nausea , vomiting , loss of appetite, dizziness, worsening tiredness or weakness. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea . Serious side effects may be more likely in older adults and those who are malnourished or debilitated. Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent. Common side effects of hydromorphone may include: drowsiness, tiredness
- dizziness
- headache
- or constipation , nausea, vomiting, stomach pain.

2339. This particular disease, Pain, interacts with the drug Roxycodone and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Oxycodone can slow or stop your breathing, and

Drug Side Effects Summary

death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep

- a slow heart rate or weak pulse
- a light-headed feeling, like you might pass out
- confusion, unusual thoughts or behavior
- seizure (convulsions)
- low cortisol levels - nausea , vomiting , loss of appetite, dizziness, worsening tiredness or weakness
- or high levels of serotonin in the body - agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, diarrhea . Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders. Common Roxycodone side effects may include: drowsiness, headache , dizziness, tiredness
- or constipation , stomach pain, nausea, vomiting.

2340. This particular disease, Pain, interacts with the drug Tylenol Arthritis Pain and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever

Drug Side Effects Summary

- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Tylenol Arthritis Pain side effects

2341. This particular disease, Pain, interacts with the drug morphine and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Call your doctor at once if you have: slow heart rate , sighing, weak or shallow breathing, breathing that stops
- chest pain, fast or pounding heartbeats
- extreme drowsiness, feeling like you might pass out
- serotonin syndrome - agitation , hallucinations, fever, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , diarrhea
- or low cortisol levels - nausea, vomiting , loss of appetite, dizziness, worsening tiredness or

Drug Side Effects Summary

weakness. Serious breathing problems may be more likely in older adults and people who are debilitated or have wasting syndrome or chronic breathing disorders. Common morphine side effects may include: drowsiness, dizziness, tiredness

- constipation , stomach pain, nausea, vomiting
- sweating
- or feelings of extreme happiness or sadness.

2342. This particular disease, Pain, interacts with the drug Fentanyl Transdermal System and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Feeling confused. Severe constipation or stomach pain. These may be signs of a severe bowel problem. Noisy breathing. Chest pain. Fast or slow heartbeat . Seizures. Low mood (depression). Trouble controlling body movements. Swelling in the arms or legs. A severe and sometimes deadly problem called serotonin syndrome may happen if you take fentanyl transdermal system with certain other drugs. Call your doctor right away if you have agitation

Drug Side Effects Summary

- change in balance
- confusion
- hallucinations
- fever
- fast or abnormal heartbeat
- flushing
- muscle twitching or stiffness
- seizures
- shivering or shaking
- sweating a lot
- severe diarrhea , upset stomach , or throwing up
- or severe headache . Taking an opioid drug like fentanyl transdermal system may lead to a rare but very bad adrenal gland problem. Call your doctor right away if you have very bad dizziness or passing out, very bad upset stomach or throwing up, or if you feel less hungry, very tired, or very weak. Fentanyl Transdermal System side effects

2343. This particular disease, Pain, interacts with the drug pregabalin and has the following side effects

- Pregabalin can cause a severe allergic reaction. Stop taking this medicine and get emergency medical help if you have: hives or blisters on your skin
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, depression, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself. Pregabalin may cause serious side effects. Call your doctor at once if you have: weak or shallow breathing

Drug Side Effects Summary

- blue-colored skin, lips, fingers, and toes
- confusion, extreme drowsiness or weakness
- vision problems
- skin sores (if you have diabetes)
- easy bruising, unusual bleeding
- swelling in your hands or feet, rapid weight gain (especially if you have diabetes or heart problems)
- or unexplained muscle pain, tenderness, or weakness (especially if you also have fever or don't feel well). Pregabalin can cause life-threatening breathing problems. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Breathing problems may be more likely in older adults or in people with COPD. If you have diabetes, tell your doctor right away if you have any new sores or other skin problems. Common side effects of pregabalin may include: dizziness , drowsiness
- swelling in your hands and feet
- trouble concentrating
- increased appetite
- weight gain
- dry mouth
- or blurred vision.

2344. This particular disease, Pain, interacts with the drug acetaminophen / tramadol and has the fo

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking acetaminophen and tramadol and call your doctor right away if you

Drug Side Effects Summary

have skin redness or a rash that spreads and causes blistering and peeling. This medicine can slow or stop your breathing, and death may occur. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep

- a slow heart rate or weak pulse
- a light-headed feeling, like you might pass out
- seizure (convulsions)
- chest pain
- liver problems--upper stomach pain, loss of appetite, dark urine, jaundice (yellowing of the skin or eyes)
- low cortisol levels-- nausea , vomiting , loss of appetite, dizziness, worsening tiredness or weakness
- or high levels of serotonin in the body-- agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, diarrhea . Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders. Common side effects of acetaminophen and tramadol may include: dizziness, drowsiness
- stomach pain, nausea, loss of appetite, diarrhea, constipation
- or sweating.

2345. This particular disease, Pain, interacts with the drug hydrocodone / ibuprofen and has the fol

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. Opioid medicine can slow or stop your

Drug Side Effects Summary

breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Hydrocodone and ibuprofen may cause serious side effects. Call your doctor at once if you have: weak or shallow breathing, sighing, slow heartbeat , breathing that stops during sleep

- swelling, rapid weight gain
- a skin rash, no matter how mild
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems--loss of appetite, diarrhea , upper stomach pain, tiredness, flu-like symptoms, dark urine, jaundice (yellowing of the skin or eyes)
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- low cortisol levels-- vomiting , dizziness, worsening tiredness or weakness
- or high levels of serotonin in the body-- agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, diarrhea. Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders. Common side effects of hydrocodone and ibuprofen may include: dizziness, drowsiness
- stomach pain, nausea, vomiting, heartburn , gas
- constipation , diarrhea
- or headache .

2346. This particular disease, Pain, interacts with the drug fentanyl and has the following side effects

- hives
- difficulty breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Like other narcotic medicines, fentanyl can slow your breathing. Death may occur if breathing becomes too weak. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Fentanyl may cause serious side effects. Call your doctor at once if you have : slow heart rate , sighing, shallow breathing, breathing that stops during sleep
- severe drowsiness, feeling like you might pass out
- confusion, extreme fear, unusual thoughts or behavior
- or low cortisol levels - nausea , vomiting , loss of appetite, dizziness, worsening tiredness or weakness. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea . Serious side effects may be more likely in older adults and those who are malnourished or debilitated. Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent. Common fentanyl side effects may include: headache , dizziness, drowsiness, pale skin, feeling weak or tired constipation , nausea, vomiting, stomach pain or sleep problems (insomnia) swelling in your hands or feet increased sweating, or cold feeling for patches itching, redness, or rash where a patch was worn.

2347. This particular disease, Pain, interacts with the drug oxymorphone and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Oxymorphone may cause serious side effects. Call your doctor at once if you have: weak or

Drug Side Effects Summary

shallow breathing, breathing that stops

- a light-headed feeling, like you might pass out
- seizure (convulsions)
- chest pain, wheezing, cough with yellow or green mucus
- severe vomiting
- high levels of serotonin in the body-- agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, diarrhea
- or low cortisol levels--nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness. Serious breathing problems may be more likely in older adults and people who are debilitated or have wasting syndrome or chronic breathing disorders. Common side effects of oxymorphone may include: dizziness, drowsiness
- headache , tiredness
- or stomach pain, nausea, vomiting, constipation .

2348. This particular disease, Pain, interacts with the drug tapentadol and has the following side effects

- hives
- chest pain, fast heartbeats, difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops
- a light-headed feeling, like you might pass out
- agitation , feeling hot
- severe drowsiness or dizziness, confusion, problems with speech or balance

Drug Side Effects Summary

- a seizure
- serotonin syndrome - agitation, hallucinations, fever, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , diarrhea
- or low cortisol levels - nausea, vomiting , loss of appetite, dizziness, worsening tiredness or weakness. Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders. Common tapentadol side effects may include: constipation, nausea, vomiting, stomach pain
- headache , feeling tired
- or drowsiness, dizziness.

2349. This particular disease, Pain, interacts with the drug Tylenol 8 Hour and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (

Drug Side Effects Summary

Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)

- red or irritated eyes

- or sores in your mouth, throat, nose, or eyes. Tylenol 8 Hour side effects

2350. This particular disease, Pain, interacts with the drug Altenol and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bloody or black, tarry stools bloody or cloudy urine fever with or without chills (not present before treatment and not caused by the condition being treated) pain in the lower back and/or side (severe and/or sharp) pinpoint red spots on the skin skin rash, hives, or itching sore throat (not present before treatment and not caused by the condition being treated) sores, ulcers, or white spots on the lips or in the mouth sudden decrease in the amount of urine unusual bleeding or bruising unusual tiredness or weakness yellow eyes or skin Symptoms of overdose Diarrhea increased sweating loss of appetite nausea or vomiting stomach cramps or pain swelling, pain, or tenderness in the upper abdomen or stomach area Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Altenol side effects

2351. This particular disease, Pain, interacts with the drug Arthritis Pain Relief and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bloody or black, tarry stools

Drug Side Effects Summary

bloody or cloudy urine fever with or without chills (not present before treatment and not caused by the condition being treated) pain in the lower back and/or side (severe and/or sharp) pinpoint red spots on the skin skin rash, hives, or itching sore throat (not present before treatment and not caused by the condition being treated) sores, ulcers, or white spots on the lips or in the mouth sudden decrease in the amount of urine unusual bleeding or bruising unusual tiredness or weakness yellow eyes or skin Symptoms of overdose Diarrhea increased sweating loss of appetite nausea or vomiting stomach cramps or pain swelling, pain, or tenderness in the upper abdomen or stomach area Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Arthritis Pain Relief side effects

2352. This particular disease, Pain, interacts with the drug Feverall and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, Feverall may cause a severe skin reaction that can be fatal. This could occur even if you have taken this medicine in the past and had no reaction. Stop using Feverall and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains this medicine. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: nausea, upper stomach pain, loss of appetite
- itching, dark urine, clay-colored stools
- or jaundice (yellowing of the skin or eyes). Less serious side effects may be more likely, and you may have none at all.

Drug Side Effects Summary

2353. This particular disease, Pain, interacts with the drug Flector Patch and has the following side effects:

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. Stop using Flector and call your doctor at once if you have: a skin rash, no matter how mild
- shortness of breath (even with mild exertion)
- swelling or rapid weight gain
- signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems - nausea , upper stomach pain, itching, tired feeling, flu-like symptoms, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or low red blood cells (anemia) - pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Common Flector side effects may include: heartburn , gas, stomach pain, nausea, vomiting
- diarrhea , constipation
- or mild itching, burning, redness, or other skin irritation where the patch was worn.

2354. This particular disease, Pain, interacts with the drug Mapap and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, Mapap may cause a severe skin reaction that can be fatal, even if you took this medicine in the past and had no reaction. Stop taking Mapap and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Stop taking Mapap and call your doctor at once if you have signs of liver

Drug Side Effects Summary

problems: stomach pain (upper right side)

- loss of appetite
- tiredness, itching
- dark urine, clay-colored stools
- or jaundice (yellowing of the skin or eyes). Less serious side effects may be more likely, and you may have none at all.

2355. This particular disease, Pain, interacts with the drug meperidine and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Meperidine may cause serious side effects. Stop using meperidine and call your doctor at once if you have: slow heartbeats, weak or shallow breathing, breathing that stops during sleep
- severe drowsiness, feeling like you might pass out
- confusion, mood changes
- severe constipation
- tremors, muscle movements you cannot control, or a seizure (convulsions)
- or low cortisol levels-- nausea , vomiting , loss of appetite, dizziness, worsening tiredness or weakness. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea . Serious side effects may be more likely in older adults and those who are malnourished or debilitated. Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on

Drug Side Effects Summary

fertility are permanent. Common side effects of meperidine may include: dizziness, drowsiness

- headache
- nausea, vomiting
- or sweating.

2356. This particular disease, Pain, interacts with the drug MS Contin and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Call your doctor at once if you have: slow heart rate , sighing, weak or shallow breathing, breathing that stops
- chest pain, fast or pounding heartbeats
- extreme drowsiness, feeling like you might pass out
- serotonin syndrome - agitation , hallucinations, fever, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , diarrhea
- or low cortisol levels - nausea, vomiting , loss of appetite, dizziness, worsening tiredness or weakness. Serious breathing problems may be more likely in older adults and people who are debilitated or have wasting syndrome or chronic breathing disorders. Common MS Contin side effects may include: drowsiness, dizziness, tiredness
- constipation , stomach pain, nausea, vomiting
- sweating
- or feelings of extreme happiness or sadness.

2357. This particular disease, Pain, interacts with the drug 7T Gummy ES Chewable Tablets and has

Drug Side Effects Summary

- No side effects listed

2358. This particular disease, Pain, interacts with the drug Abenol and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bloody or black, tarry stools bloody or cloudy urine fever with or without chills (not present before treatment and not caused by the condition being treated) pain in the lower back and/or side (severe and/or sharp) pinpoint red spots on the skin skin rash, hives, or itching sore throat (not present before treatment and not caused by the condition being treated) sores, ulcers, or white spots on the lips or in the mouth sudden decrease in the amount of urine unusual bleeding or bruising unusual tiredness or weakness yellow eyes or skin Symptoms of overdose Diarrhea increased sweating loss of appetite nausea or vomiting stomach cramps or pain swelling, pain, or tenderness in the upper abdomen or stomach area Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Abenol side effects

2359. This particular disease, Pain, interacts with the drug Acephen and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, Acephen may cause a severe skin reaction that can be fatal. This could occur even if you have taken this medicine in the past and had no reaction. Stop using Acephen and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. If you have this type of reaction, you should never again take any medicine that contains this medicine. This medicine may cause serious side effects.

Drug Side Effects Summary

Stop using this medicine and call your doctor at once if you have: nausea, upper stomach pain, loss of appetite

- itching, dark urine, clay-colored stools

- or jaundice (yellowing of the skin or eyes). Less serious side effects may be more likely, and you may have none at all.

2360. This particular disease, Pain, interacts with the drug Actamin and has the following side effects

- hives

- difficulty breathing

- swelling of your face, lips, tongue, or throat. In rare cases, Actamin may cause a severe skin reaction that can be fatal, even if you took this medicine in the past and had no reaction. Stop taking Actamin and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Stop taking Actamin and call your doctor at once if you have signs of liver problems: stomach pain (upper right side)

- loss of appetite

- tiredness, itching

- dark urine, clay-colored stools

- or jaundice (yellowing of the skin or eyes). Less serious side effects may be more likely, and you may have none at all.

2361. This particular disease, Pain, interacts with the drug Actimol Children's and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bloody or black, tarry stools bloody or cloudy urine fever with or without chills (not present before treatment and not caused by

Drug Side Effects Summary

the condition being treated) pain in the lower back and/or side (severe and/or sharp) pinpoint red spots on the skin skin rash, hives, or itching sore throat (not present before treatment and not caused by the condition being treated) sores, ulcers, or white spots on the lips or in the mouth sudden decrease in the amount of urine unusual bleeding or bruising unusual tiredness or weakness yellow eyes or skin Symptoms of overdose Diarrhea increased sweating loss of appetite nausea or vomiting stomach cramps or pain swelling, pain, or tenderness in the upper abdomen or stomach area Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Actimol Children's side effects

2362. This particular disease, Pain, interacts with the drug Actimol Infant and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bloody or black, tarry stools bloody or cloudy urine fever with or without chills (not present before treatment and not caused by the condition being treated) pain in the lower back and/or side (severe and/or sharp) pinpoint red spots on the skin skin rash, hives, or itching sore throat (not present before treatment and not caused by the condition being treated) sores, ulcers, or white spots on the lips or in the mouth sudden decrease in the amount of urine unusual bleeding or bruising unusual tiredness or weakness yellow eyes or skin Symptoms of overdose Diarrhea increased sweating loss of appetite nausea or vomiting stomach cramps or pain swelling, pain, or tenderness in the upper abdomen or stomach area Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Actimol Infant side effects

Drug Side Effects Summary

2363. This particular disease, Pain, interacts with the drug Advil Children's and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Abdominal pain acid or sour stomach belching bloating cloudy urine decrease in amount of urine decrease in urine output or decrease in urine-concentrating ability diarrhea difficulty having a bowel movement (stool) excess air or gas in stomach or intestines full feeling heartburn indigestion itching skin pain or discomfort in chest, upper stomach, or throat pale skin passing gas nausea noisy, rattling breathing rash with flat lesions or small raised lesions on the skin shortness of breath swelling of face, fingers, hands, feet, lower legs, or ankles troubled breathing at rest troubled breathing with exertion unusual bleeding or bruising unusual tiredness or weakness vomiting weight gain Less common Abdominal cramps stomach soreness or discomfort Rare Agitation back, leg, or stomach pains bleeding gums blistering, peeling, loosening of skin blood in urine or stools bloody, black, or tarry stools blurred vision burning feeling in chest or stomach change in vision chest pain chills clay-colored stools coma confusion constipation cough or hoarseness dark urine decreased urine output depression difficulty breathing difficulty swallowing dilated neck veins dizziness dry mouth extreme fatigue fast, irregular, pounding, or racing heartbeat or pulse fever with or without chills frequent urination general body swelling general feeling of tiredness or weakness hair loss, thinning of hair headache hives or welts hostility impaired vision increased blood pressure increased volume of pale, dilute urine irregular breathing irritability itching joint or muscle pain lab results that show problems with liver lethargy light-colored stools loss of appetite lower back or side pain muscle twitching nosebleeds painful or difficult urination pains in stomach, side, or abdomen, possibly radiating to the back pinpoint red spots on skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rash red skin lesions, often with a purple center red, irritated eyes redness of skin seizures severe abdominal pain, cramping, burning severe and continuing nausea sore throat sores, ulcers, or white spots in

Drug Side Effects Summary

mouth or on lips stiff neck or back stomach upset stupor swollen or painful glands tenderness in stomach area thirst tightness in chest unpleasant breath odor upper right abdominal pain vomiting of blood vomiting of material that looks like coffee grounds wheezing yellow eyes and skin Symptoms of overdose Bluish lips or skin difficulty sleeping disorientation dizziness, faintness, or lightheadedness when getting up from a lying or sitting position suddenly drowsiness to profound coma hallucination lightheadedness or fainting mood or other mental changes muscle tremors not breathing rapid, deep breathing restlessness slow or irregular heartbeat stomach cramps sudden fainting sweating Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Continuing ringing or buzzing or other unexplained noise in ears hearing loss nervousness Rare Crying depersonalization discouragement dry eyes dysphoria euphoria feeling sad or empty lack of appetite loss of interest or pleasure mental depression paranoia quick to react or overreact rapidly changing moods runny nose sleepiness or unusual drowsiness sleeplessness sneezing stuffy nose trouble concentrating trouble sleeping unable to sleep Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Advil Children's side effects

2364. This particular disease, Pain, interacts with the drug Advil Infant's Concentrated Drops and h

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Abdominal pain acid or

Drug Side Effects Summary

sour stomach belching bloating cloudy urine decrease in amount of urine decrease in urine output or decrease in urine-concentrating ability diarrhea difficulty having a bowel movement (stool) excess air or gas in stomach or intestines full feeling heartburn indigestion itching skin pain or discomfort in chest, upper stomach, or throat pale skin passing gas nausea noisy, rattling breathing rash with flat lesions or small raised lesions on the skin shortness of breath swelling of face, fingers, hands, feet, lower legs, or ankles troubled breathing at rest troubled breathing with exertion unusual bleeding or bruising unusual tiredness or weakness vomiting weight gain

Less common Abdominal cramps stomach soreness or discomfort Rare Agitation back, leg, or stomach pains bleeding gums blistering, peeling, loosening of skin blood in urine or stools bloody, black, or tarry stools blurred vision burning feeling in chest or stomach change in vision chest pain chills clay-colored stools coma confusion constipation cough or hoarseness dark urine decreased urine output depression difficulty breathing difficulty swallowing dilated neck veins dizziness dry mouth extreme fatigue fast, irregular, pounding, or racing heartbeat or pulse fever with or without chills frequent urination general body swelling general feeling of tiredness or weakness hair loss, thinning of hair headache hives or welts hostility impaired vision increased blood pressure increased volume of pale, dilute urine irregular breathing irritability itching joint or muscle pain lab results that show problems with liver lethargy light-colored stools loss of appetite lower back or side pain muscle twitching nosebleeds painful or difficult urination pains in stomach, side, or abdomen, possibly radiating to the back pinpoint red spots on skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rash red skin lesions, often with a purple center red, irritated eyes redness of skin seizures severe abdominal pain, cramping, burning severe and continuing nausea sore throat sores, ulcers, or white spots in mouth or on lips stiff neck or back stomach upset stupor swollen or painful glands tenderness in stomach area thirst tightness in chest unpleasant breath odor upper right abdominal pain vomiting of blood vomiting of material that looks like coffee grounds wheezing yellow eyes and skin

Symptoms of overdose Bluish lips or skin difficulty sleeping disorientation dizziness, faintness, or

Drug Side Effects Summary

lightheadedness when getting up from a lying or sitting position suddenly drowsiness to profound coma hallucination lightheadedness or fainting mood or other mental changes muscle tremors not breathing rapid, deep breathing restlessness slow or irregular heartbeat stomach cramps sudden fainting sweating Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Continuing ringing or buzzing or other unexplained noise in ears hearing loss nervousness Rare Crying depersonalization discouragement dry eyes dysphoria euphoria feeling sad or empty lack of appetite loss of interest or pleasure mental depression paranoia quick to react or overreact rapidly changing moods runny nose sleepiness or unusual drowsiness sleeplessness sneezing stuffy nose trouble concentrating trouble sleeping unable to sleep Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Advil Infant's Concentrated Drops side effects

2365. This particular disease, Pain, interacts with the drug Advil Migraine and has the following side effects:

- No side effects listed

2366. This particular disease, Pain, interacts with the drug Aminofen and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bloody or black, tarry stools bloody or cloudy urine fever with or without chills (not present before treatment and not caused by

Drug Side Effects Summary

the condition being treated) pain in the lower back and/or side (severe and/or sharp) pinpoint red spots on the skin skin rash, hives, or itching sore throat (not present before treatment and not caused by the condition being treated) sores, ulcers, or white spots on the lips or in the mouth sudden decrease in the amount of urine unusual bleeding or bruising unusual tiredness or weakness yellow eyes or skin Symptoms of overdose Diarrhea increased sweating loss of appetite nausea or vomiting stomach cramps or pain swelling, pain, or tenderness in the upper abdomen or stomach area Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Aminofen side effects

2367. This particular disease, Pain, interacts with the drug Anacin Aspirin Free and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, Anacin AF may cause a severe skin reaction that can be fatal, even if you took this medicine in the past and had no reaction. Stop taking Anacin AF and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Stop taking Anacin AF and call your doctor at once if you have signs of liver problems: stomach pain (upper right side)
- loss of appetite
- tiredness, itching
- dark urine, clay-colored stools
- or jaundice (yellowing of the skin or eyes). Less serious side effects may be more likely, and you may have none at all.

2368. This particular disease, Pain, interacts with the drug Apra and has the following side effects:

Drug Side Effects Summary

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bloody or black, tarry stools bloody or cloudy urine fever with or without chills (not present before treatment and not caused by the condition being treated) pain in the lower back and/or side (severe and/or sharp) pinpoint red spots on the skin skin rash, hives, or itching sore throat (not present before treatment and not caused by the condition being treated) sores, ulcers, or white spots on the lips or in the mouth sudden decrease in the amount of urine unusual bleeding or bruising unusual tiredness or weakness yellow eyes or skin Symptoms of overdose Diarrhea increased sweating loss of appetite nausea or vomiting stomach cramps or pain swelling, pain, or tenderness in the upper abdomen or stomach area Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Apra side effects

2369. This particular disease, Pain, interacts with the drug Atasol and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bloody or black, tarry stools bloody or cloudy urine fever with or without chills (not present before treatment and not caused by the condition being treated) pain in the lower back and/or side (severe and/or sharp) pinpoint red spots on the skin skin rash, hives, or itching sore throat (not present before treatment and not caused by the condition being treated) sores, ulcers, or white spots on the lips or in the mouth sudden decrease in the amount of urine unusual bleeding or bruising unusual tiredness or weakness yellow eyes or skin Symptoms of overdose Diarrhea increased sweating loss of appetite nausea or vomiting stomach cramps or pain swelling, pain, or tenderness in the upper abdomen or

Drug Side Effects Summary

stomach area Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Atasol side effects

2370. This particular disease, Pain, interacts with the drug Aurophen and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, Aurophen may cause a severe skin reaction that can be fatal, even if you took this medicine in the past and had no reaction. Stop taking Aurophen and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Stop taking Aurophen and call your doctor at once if you have signs of liver problems: stomach pain (upper right side)
- loss of appetite
- tiredness, itching
- dark urine, clay-colored stools
- or jaundice (yellowing of the skin or eyes). Less serious side effects may be more likely, and you may have none at all.

2371. This particular disease, Pain, interacts with the drug Caldolor and has the following side effects

- hives, difficult breathing, swelling in your face or throat) or a severe skin reaction : (fever, sore throat, burning eyes, skin pain, red or purple skin rash with blistering and peeling). Stop using Caldolor and seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Get emergency medical help if you have signs of a heart attack or stroke: chest pain spreading to your jaw or shoulder, sudden

Drug Side Effects Summary

numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath. Tell your caregivers at once if you have: changes in your vision

- swelling, rapid weight gain, feeling short of breath
- severe headache , pounding in your neck or ears
- little or no urination
- high potassium level - nausea , weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement
- liver problems - loss of appetite, stomach pain (upper right side), dark urine, jaundice (yellowing of the skin or eyes)
- low red blood cells (anemia) - pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet
- or signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. If your baby has been treated with Neoprofen , tell the doctor at once if the baby has: signs of infection such as fever, chills, fussiness
- unusual bleeding
- or bruising, swelling, warmth, redness, or irritation where the IV needle is placed. Common Caldolor side effects may include: nausea, vomiting , gas
- headache
- or dizziness .

2372. This particular disease, Pain, interacts with the drug Cetafen and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bloody or black, tarry stools
bloody or cloudy urine fever with or without chills (not present before treatment and not caused by

Drug Side Effects Summary

the condition being treated) pain in the lower back and/or side (severe and/or sharp) pinpoint red spots on the skin skin rash, hives, or itching sore throat (not present before treatment and not caused by the condition being treated) sores, ulcers, or white spots on the lips or in the mouth sudden decrease in the amount of urine unusual bleeding or bruising unusual tiredness or weakness yellow eyes or skin Symptoms of overdose Diarrhea increased sweating loss of appetite nausea or vomiting stomach cramps or pain swelling, pain, or tenderness in the upper abdomen or stomach area Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Cetafen side effects

2373. This particular disease, Pain, interacts with the drug Children's Motrin and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Abdominal pain acid or sour stomach belching bloating cloudy urine decrease in amount of urine decrease in urine output or decrease in urine-concentrating ability diarrhea difficulty having a bowel movement (stool) excess air or gas in stomach or intestines full feeling heartburn indigestion itching skin pain or discomfort in chest, upper stomach, or throat pale skin passing gas nausea noisy, rattling breathing rash with flat lesions or small raised lesions on the skin shortness of breath swelling of face, fingers, hands, feet, lower legs, or ankles troubled breathing at rest troubled breathing with exertion unusual bleeding or bruising unusual tiredness or weakness vomiting weight gain Less common Abdominal cramps stomach soreness or discomfort Rare Agitation back, leg, or stomach pains bleeding gums blistering, peeling, loosening of skin blood in urine or stools bloody, black, or tarry stools blurred vision burning feeling in chest or stomach change in vision chest pain chills clay-colored stools coma confusion constipation cough or hoarseness dark urine decreased urine output depression difficulty

Drug Side Effects Summary

breathing difficulty swallowing dilated neck veins dizziness dry mouth extreme fatigue fast, irregular, pounding, or racing heartbeat or pulse fever with or without chills frequent urination general body swelling general feeling of tiredness or weakness hair loss, thinning of hair headache hives or welts hostility impaired vision increased blood pressure increased volume of pale, dilute urine irregular breathing irritability itching joint or muscle pain lab results that show problems with liver lethargy light-colored stools loss of appetite lower back or side pain muscle twitching nosebleeds painful or difficult urination pains in stomach, side, or abdomen, possibly radiating to the back pinpoint red spots on skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rash red skin lesions, often with a purple center red, irritated eyes redness of skin seizures severe abdominal pain, cramping, burning severe and continuing nausea sore throat sores, ulcers, or white spots in mouth or on lips stiff neck or back stomach upset stupor swollen or painful glands tenderness in stomach area thirst tightness in chest unpleasant breath odor upper right abdominal pain vomiting of blood vomiting of material that looks like coffee grounds wheezing yellow eyes and skin

Symptoms of overdose Bluish lips or skin difficulty sleeping disorientation dizziness, faintness, or lightheadedness when getting up from a lying or sitting position suddenly drowsiness to profound coma hallucination lightheadedness or fainting mood or other mental changes muscle tremors not breathing rapid, deep breathing restlessness slow or irregular heartbeat stomach cramps sudden fainting sweating

Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them:

More common Continuing ringing or buzzing or other unexplained noise in ears hearing loss nervousness

Rare Crying depersonalization discouragement dry eyes dysphoria euphoria feeling sad or empty lack of appetite loss of interest or pleasure mental depression paranoia quick to react or overreact rapidly changing moods runny nose

Drug Side Effects Summary

sleepiness or unusual drowsiness sleeplessness sneezing stuffy nose trouble concentrating trouble sleeping unable to sleep Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Children's Motrin side effects

2374. This particular disease, Pain, interacts with the drug Children's Silapap and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
 - hives
 - itching
 - red, swollen, blistered, or peeling skin with or without fever
 - wheezing
 - tightness in the chest or throat
 - trouble breathing, swallowing, or talking
 - unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes

Drug Side Effects Summary

- or sores in your mouth, throat, nose, or eyes. Children's Silapap side effects

2375. This particular disease, Pain, interacts with the drug Children's Tylenol and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, Children's Tylenol may cause a severe skin reaction that can be fatal, even if you took this medicine in the past and had no reaction. Stop taking Children's Tylenol and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Stop taking Children's Tylenol and call your doctor at once if you have signs of liver problems: stomach pain (upper right side)
- loss of appetite
- tiredness, itching
- dark urine, clay-colored stools
- or jaundice (yellowing of the skin or eyes). Less serious side effects may be more likely, and you may have none at all.

2376. This particular disease, Pain, interacts with the drug ConZip and has the following side effect

- noisy breathing, sighing, shallow breathing, breathing that stops during sleep
- a slow heart rate or weak pulse
- a light-headed feeling, like you might pass out
- seizure (convulsions)
- or low cortisol levels-- nausea , vomiting , loss of appetite, dizziness, worsening tiredness or weakness. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea . Serious breathing problems may be more likely in

Drug Side Effects Summary

older adults and people who are debilitated or have wasting syndrome or chronic breathing disorders. Common side effects of ConZip may include: constipation , nausea, vomiting, stomach pain

- dizziness, drowsiness, tiredness
- headache
- or itching.

2377. This particular disease, Pain, interacts with the drug Dolono and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bloody or black, tarry stools bloody or cloudy urine fever with or without chills (not present before treatment and not caused by the condition being treated) pain in the lower back and/or side (severe and/or sharp) pinpoint red spots on the skin skin rash, hives, or itching sore throat (not present before treatment and not caused by the condition being treated) sores, ulcers, or white spots on the lips or in the mouth sudden decrease in the amount of urine unusual bleeding or bruising unusual tiredness or weakness yellow eyes or skin Symptoms of overdose Diarrhea increased sweating loss of appetite nausea or vomiting stomach cramps or pain swelling, pain, or tenderness in the upper abdomen or stomach area Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Dolono side effects

2378. This particular disease, Pain, interacts with the drug Duragesic and has the following side effects

- hives
- chest pain, difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Remove the Duragesic patch and call your doctor at once if you have: slow heart rate , sighing, weak or shallow breathing (up to several days after removing the skin patch)
- breathing that stops during sleep
- confusion, severe drowsiness, feeling like you might pass out
- chest pain, fast or pounding heartbeats
- or low cortisol levels - nausea , vomiting , loss of appetite, dizziness, worsening tiredness or weakness. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea . Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Common Duragesic side effects may include: headache , dizziness, drowsiness, tiredness
- nausea, vomiting, stomach pain, diarrhea, constipation
- itching, redness, or rash where a patch was worn
- sleep problems (insomnia)
- or increased sweating, or cold feeling.

2379. This particular disease, Pain, interacts with the drug Febrol Solution and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives

Drug Side Effects Summary

- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Febrol Solution side effects

2380. This particular disease, Pain, interacts with the drug Genapap and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bloody or black, tarry stools bloody or cloudy urine fever with or without chills (not present before treatment and not caused by the condition being treated) pain in the lower back and/or side (severe and/or sharp) pinpoint red spots on the skin skin rash, hives, or itching sore throat (not present before treatment and not caused by the condition being treated) sores, ulcers, or white spots on the lips or in the mouth sudden decrease in the amount of urine unusual bleeding or bruising unusual tiredness or weakness yellow eyes or skin Symptoms of overdose Diarrhea increased sweating loss of appetite

Drug Side Effects Summary

nausea or vomiting stomach cramps or pain swelling, pain, or tenderness in the upper abdomen or stomach area Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Genapap side effects

2381. This particular disease, Pain, interacts with the drug Genebs and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bloody or black, tarry stools bloody or cloudy urine fever with or without chills (not present before treatment and not caused by the condition being treated) pain in the lower back and/or side (severe and/or sharp) pinpoint red spots on the skin skin rash, hives, or itching sore throat (not present before treatment and not caused by the condition being treated) sores, ulcers, or white spots on the lips or in the mouth sudden decrease in the amount of urine unusual bleeding or bruising unusual tiredness or weakness yellow eyes or skin Symptoms of overdose Diarrhea increased sweating loss of appetite nausea or vomiting stomach cramps or pain swelling, pain, or tenderness in the upper abdomen or stomach area Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Genebs side effects

2382. This particular disease, Pain, interacts with the drug Infant's Tylenol and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

Drug Side Effects Summary

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Infant's Tylenol side effects

2383. This particular disease, Pain, interacts with the drug Jr. Tylenol and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing

Drug Side Effects Summary

- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Jr. Tylenol side effects

2384. This particular disease, Pain, interacts with the drug Kadian and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Call your doctor at once if you have: slow heart rate , sighing, weak or shallow breathing, breathing that stops
- chest pain, fast or pounding heartbeats
- extreme drowsiness, feeling like you might pass out
- serotonin syndrome - agitation , hallucinations, fever, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , diarrhea
- or low cortisol levels - nausea, vomiting , loss of appetite, dizziness, worsening tiredness or weakness. Serious breathing problems may be more likely in older adults and people who are

Drug Side Effects Summary

debilitated or have wasting syndrome or chronic breathing disorders. Common Kadian side effects may include: drowsiness, dizziness, tiredness

- constipation , stomach pain, nausea, vomiting
- sweating
- or feelings of extreme happiness or sadness.

2385. This particular disease, Pain, interacts with the drug M-Pap and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, M-Pap may cause a severe skin reaction that can be fatal, even if you took this medicine in the past and had no reaction. Stop taking M-Pap and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Stop taking M-Pap and call your doctor at once if you have signs of liver problems: stomach pain (upper right side)
- loss of appetite
- tiredness, itching
- dark urine, clay-colored stools
- or jaundice (yellowing of the skin or eyes). Less serious side effects may be more likely, and you may have none at all.

2386. This particular disease, Pain, interacts with the drug Mapap Arthritis Pain and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bloody or black, tarry stools bloody or cloudy urine fever with or without chills (not present before treatment and not caused by

Drug Side Effects Summary

the condition being treated) pain in the lower back and/or side (severe and/or sharp) pinpoint red spots on the skin skin rash, hives, or itching sore throat (not present before treatment and not caused by the condition being treated) sores, ulcers, or white spots on the lips or in the mouth sudden decrease in the amount of urine unusual bleeding or bruising unusual tiredness or weakness yellow eyes or skin Symptoms of overdose Diarrhea increased sweating loss of appetite nausea or vomiting stomach cramps or pain swelling, pain, or tenderness in the upper abdomen or stomach area Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Mapap Arthritis Pain side effects

2387. This particular disease, Pain, interacts with the drug Motrin Childrens and has the following s

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Abdominal pain acid or sour stomach belching bloating cloudy urine decrease in amount of urine decrease in urine output or decrease in urine-concentrating ability diarrhea difficulty having a bowel movement (stool) excess air or gas in stomach or intestines full feeling heartburn indigestion itching skin pain or discomfort in chest, upper stomach, or throat pale skin passing gas nausea noisy, rattling breathing rash with flat lesions or small raised lesions on the skin shortness of breath swelling of face, fingers, hands, feet, lower legs, or ankles troubled breathing at rest troubled breathing with exertion unusual bleeding or bruising unusual tiredness or weakness vomiting weight gain Less common Abdominal cramps stomach soreness or discomfort Rare Agitation back, leg, or stomach pains bleeding gums blistering, peeling, loosening of skin blood in urine or stools bloody, black, or tarry stools blurred vision burning feeling in chest or stomach change in vision chest pain chills clay-colored stools coma

Drug Side Effects Summary

confusion constipation cough or hoarseness dark urine decreased urine output depression difficulty breathing difficulty swallowing dilated neck veins dizziness dry mouth extreme fatigue fast, irregular, pounding, or racing heartbeat or pulse fever with or without chills frequent urination general body swelling general feeling of tiredness or weakness hair loss, thinning of hair headache hives or welts hostility impaired vision increased blood pressure increased volume of pale, dilute urine irregular breathing irritability itching joint or muscle pain lab results that show problems with liver lethargy light-colored stools loss of appetite lower back or side pain muscle twitching nosebleeds painful or difficult urination pains in stomach, side, or abdomen, possibly radiating to the back pinpoint red spots on skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rash red skin lesions, often with a purple center red, irritated eyes redness of skin seizures severe abdominal pain, cramping, burning severe and continuing nausea sore throat sores, ulcers, or white spots in mouth or on lips stiff neck or back stomach upset stupor swollen or painful glands tenderness in stomach area thirst tightness in chest unpleasant breath odor upper right abdominal pain vomiting of blood vomiting of material that looks like coffee grounds wheezing yellow eyes and skin

Symptoms of overdose Bluish lips or skin difficulty sleeping disorientation dizziness, faintness, or lightheadedness when getting up from a lying or sitting position suddenly drowsiness to profound coma hallucination lightheadedness or fainting mood or other mental changes muscle tremors not breathing rapid, deep breathing restlessness slow or irregular heartbeat stomach cramps sudden fainting sweating

Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them:

More common Continuing ringing or buzzing or other unexplained noise in ears hearing loss nervousness

Rare Crying depersonalization discouragement dry eyes dysphoria euphoria feeling sad or empty lack of appetite loss of interest or

Drug Side Effects Summary

pleasure mental depression paranoia quick to react or overreact rapidly changing moods runny nose sleepiness or unusual drowsiness sleeplessness sneezing stuffy nose trouble concentrating trouble sleeping unable to sleep Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Motrin Childrens side effects

2388. This particular disease, Pain, interacts with the drug Motrin Junior Strength and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of bleeding like throwing up or coughing up blood
- vomit that looks like coffee grounds
- blood in the urine
- black, red, or tarry stools
- bleeding from the gums

Drug Side Effects Summary

- abnormal vaginal bleeding
- bruises without a cause or that get bigger
- or bleeding you cannot stop. Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain. Signs of high potassium levels like a heartbeat that does not feel normal
- feeling confused
- feeling weak, lightheaded, or dizzy
- feeling like passing out
- numbness or tingling
- or shortness of breath. Signs of high blood pressure like very bad headache or dizziness , passing out, or change in eyesight. Shortness of breath, a big weight gain, or swelling in the arms or legs. Chest pain or pressure or a fast heartbeat . Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight. Feeling very tired or weak. Ringing in ears. Very bad back pain . Change in eyesight. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Motrin Junior Strength side effects

2389. This particular disease, Pain, interacts with the drug Nortemp Children's and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bloody or black, tarry stools bloody or cloudy urine fever with or without chills (not present before treatment and not caused by

Drug Side Effects Summary

the condition being treated) pain in the lower back and/or side (severe and/or sharp) pinpoint red spots on the skin skin rash, hives, or itching sore throat (not present before treatment and not caused by the condition being treated) sores, ulcers, or white spots on the lips or in the mouth sudden decrease in the amount of urine unusual bleeding or bruising unusual tiredness or weakness yellow eyes or skin Symptoms of overdose Diarrhea increased sweating loss of appetite nausea or vomiting stomach cramps or pain swelling, pain, or tenderness in the upper abdomen or stomach area Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Nortemp Children's side effects

2390. This particular disease, Pain, interacts with the drug Nucynta ER and has the following side effects

- hives
- chest pain, fast heartbeats, difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Nucynta ER may cause serious side effects. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops
- a light-headed feeling, like you might pass out
- agitation , feeling hot
- severe drowsiness or dizziness, confusion, problems with speech or balance
- a seizure
- serotonin syndrome --agitation, hallucinations, fever, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , diarrhea
- or low cortisol levels-- nausea, vomiting , loss of appetite, dizziness, worsening tiredness or

Drug Side Effects Summary

weakness. Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders. Common side effects of Nucynta ER may include: constipation, nausea, vomiting, stomach pain

- headache , feeling tired
- or drowsiness, dizziness.

2391. This particular disease, Pain, interacts with the drug Ofirmev and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal, even if you took acetaminophen in the past and had no reaction. Tell your caregivers or call your doctor if you have skin redness or a rash that spreads and causes blistering and peeling. Stop using Ofirmev and call your doctor at once if you have: fever, cough, chest pain, trouble breathing
- or liver problems - nausea, upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common Ofirmev side effects may include: cough, breathing problems
- nausea, vomiting
- constipation
- agitation
- itching
- headache
- or sleep problems (insomnia).

2392. This particular disease, Pain, interacts with the drug Pharbetol and has the following side effects

Drug Side Effects Summary

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, Pharbetol may cause a severe skin reaction that can be fatal, even if you took this medicine in the past and had no reaction. Stop taking Pharbetol and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Stop taking Pharbetol and call your doctor at once if you have signs of liver problems: stomach pain (upper right side)
- loss of appetite
- tiredness, itching
- dark urine, clay-colored stools
- or jaundice (yellowing of the skin or eyes). Less serious side effects may be more likely, and you may have none at all.

2393. This particular disease, Pain, interacts with the drug Qdolo and has the following side effects

- The possible side effects of Qdolo include: constipation, nausea, sleepiness, vomiting, tiredness, headache, dizziness, abdominal pain. Call your healthcare provider if you have any of these symptoms and they are severe. trouble breathing, shortness of breath, fast heartbeat, chest pain, swelling of your face, tongue, or throat, extreme drowsiness, light-headedness when changing positions, feeling faint, agitation, high body temperature, trouble walking, stiff muscles, or mental changes such as confusion. These are not all the possible side effects of Qdolo. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. For more information go to dailymed.nlm.nih.gov. Qdolo side effects

2394. This particular disease, Pain, interacts with the drug Silapap Childrens and has the following

- hives

Drug Side Effects Summary

- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, Silapap Childrens may cause a severe skin reaction that can be fatal, even if you took this medicine in the past and had no reaction. Stop taking Silapap Childrens and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Stop taking Silapap Childrens and call your doctor at once if you have signs of liver problems: stomach pain (upper right side)
- loss of appetite
- tiredness, itching
- dark urine, clay-colored stools
- or jaundice (yellowing of the skin or eyes). Less serious side effects may be more likely, and you may have none at all.

2395. This particular disease, Pain, interacts with the drug Tactinal and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, Tactinal may cause a severe skin reaction that can be fatal, even if you took this medicine in the past and had no reaction. Stop taking Tactinal and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Stop taking Tactinal and call your doctor at once if you have signs of liver problems: stomach pain (upper right side)
- loss of appetite
- tiredness, itching
- dark urine, clay-colored stools
- or jaundice (yellowing of the skin or eyes). Less serious side effects may be more likely, and you may have none at all.

Drug Side Effects Summary

2396. This particular disease, Pain, interacts with the drug Tempra Quicklets and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, Tempra Quicklets may cause a severe skin reaction that can be fatal, even if you took this medicine in the past and had no reaction. Stop taking Tempra Quicklets and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Stop taking Tempra Quicklets and call your doctor at once if you have signs of liver problems: stomach pain (upper right side)
- loss of appetite
- tiredness, itching
- dark urine, clay-colored stools
- or jaundice (yellowing of the skin or eyes). Less serious side effects may be more likely, and you may have none at all.

2397. This particular disease, Pain, interacts with the drug Tycolene and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, Tycolene may cause a severe skin reaction that can be fatal, even if you took this medicine in the past and had no reaction. Stop taking Tycolene and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Stop taking Tycolene and call your doctor at once if you have signs of liver problems: stomach pain (upper right side)
- loss of appetite
- tiredness, itching
- dark urine, clay-colored stools

Drug Side Effects Summary

- or jaundice (yellowing of the skin or eyes). Less serious side effects may be more likely, and you may have none at all.

2398. This particular disease, Pain, interacts with the drug Tylophen and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

- itching

- red, swollen, blistered, or peeling skin with or without fever

- wheezing

- tightness in the chest or throat

- trouble breathing, swallowing, or talking

- unusual hoarseness

- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Not able to pass urine or change in how much urine is passed. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)

- red or irritated eyes

- or sores in your mouth, throat, nose, or eyes. Tylophen side effects

2399. This particular disease, Pain, interacts with the drug Ultracet and has the following side effects:

Drug Side Effects Summary

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. This medicine can slow or stop your breathing, and death may occur. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Stop using this medicine and call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep
- a slow heart rate or weak pulse
- a light-headed feeling, like you might pass out
- seizure (convulsions)
- chest pain
- liver problems - upper stomach pain, loss of appetite, dark urine, jaundice (yellowing of the skin or eyes)
- low cortisol levels - nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness
- or high levels of serotonin in the body - agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, diarrhea. Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders. Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent. Common Ultracet side effects may include: dizziness, drowsiness
- stomach pain, nausea, loss of appetite
- diarrhea, constipation

Drug Side Effects Summary

- or sweating.

2400. This particular disease, Pain, interacts with the drug Vitapap and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. In rare cases, Vitapap may cause a severe skin reaction that can be fatal, even if you took this medicine in the past and had no reaction. Stop taking Vitapap and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Stop taking Vitapap and call your doctor at once if you have signs of liver problems: stomach pain (upper right side)
- loss of appetite
- tiredness, itching
- dark urine, clay-colored stools
- or jaundice (yellowing of the skin or eyes). Less serious side effects may be more likely, and you may have none at all.

2401. This particular disease, Pain, interacts with the drug Actiq and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Like other narcotic medicines, fentanyl can slow your breathing. Death may occur if breathing becomes too weak. Remove the Actiq unit from your mouth and call your doctor at once if you have: slow heart rate, sighing, shallow breathing, breathing that stops during sleep
- severe drowsiness, feeling like you might pass out
- dizziness or an upset stomach before the medicine has completely dissolved

Drug Side Effects Summary

- confusion, extreme fear, unusual thoughts or behavior
- or low cortisol levels - nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever , sweating , shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting , or diarrhea . Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent. Common Actiq side effects may include: dizziness, mild drowsiness, depressed mood
- sleep problems (insomnia)
- headache , weakness, anxiety
- nausea, vomiting, constipation
- or mild rash .

2402. This particular disease, Pain, interacts with the drug Anjeso and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: Less common Abdominal or stomach pain, cramping, burning, or tenderness bloody or black, tarry stools constipation dark urine decreased appetite diarrhea fever headache heartburn indigestion itching or skin rash loss of appetite nausea and vomiting swelling of the feet or lower legs trouble breathing unusual tiredness or weakness vomiting of material that looks like coffee grounds, severe and continuing yellow eyes or skin Incidence not known Blistering, peeling, or loosening of the skin bloody or cloudy urine cough or hoarseness decrease in the frequency of urination decrease in urine volume diarrhea difficulty in passing urine (dribbling) difficulty swallowing dizziness fast heartbeat fever with or

Drug Side Effects Summary

without chills general feeling of tiredness or weakness greatly decreased frequency of urination or amount of urine hives lower back or side pain pains in the chest, groin, or legs, especially calves of the legs painful or difficult urination puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red, irritated eyes severe headaches of sudden onset sore throat sores, ulcers, or white spots on the lips or in the mouth sudden loss of coordination sudden onset of shortness of breath for no apparent reason sudden onset of slurred speech sudden vision changes swelling of the feet or lower legs tightness in the chest unusual bleeding or bruising Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Bleeding, blistering, burning, coldness, discoloration of the skin, feeling of pressure, hives, infection, inflammation, itching, lumps, numbness, pain, rash, redness, scarring, soreness, stinging, swelling, tenderness, tingling, ulceration, or warmth at the injection site Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Anjeso side effects

2403. This particular disease, Pain, interacts with the drug Dazidox and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Chills cold sweats confusion difficult or labored breathing dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position fever tightness in the chest twitching Rare Bloating or swelling of the face, arms, hands, lower legs, or feet blood in the urine burning while urinating

Drug Side Effects Summary

burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain cough decrease in the frequency of urination decreased urine output difficult or painful urination difficulty in passing urine (dribbling) difficulty with swallowing dizziness dry mouth fainting fast, irregular, pounding, or racing heartbeat or pulse feeling of warmth or heat flushing or redness of the skin, especially on the face and neck frequent urination headache hives, itching, or skin rash increase in heart rate increased thirst increased volume of pale, dilute urine lightheadedness muscle pain or cramps nausea puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rapid breathing rapid weight gain seizures severe constipation severe vomiting stomach pain sunken eyes sweating swollen, painful, or tender lymph glands in the neck, armpit, or groin thirst trembling or shaking of the hands or feet unusual tiredness or weakness unusual weight gain or loss vomiting wrinkled skin Incidence not known Blurred vision choking clay-colored stools cold, clammy skin dark urine darkening of the skin diarrhea fast, weak pulse gagging irregular, fast, slow, or shallow breathing loss of appetite loss of consciousness pale or blue lips, fingernails, or skin unpleasant breath odor very slow heartbeat yellow eyes or skin Symptoms of overdose Change in consciousness chest pain or discomfort cold, clammy skin constricted, pinpoint, or small pupils (black part of the eye) coughing that sometimes produces a pink frothy sputum decreased awareness or responsiveness increased sweating irregular, fast, or slow, or shallow breathing loss of consciousness no muscle tone or movement pale or blue lips, fingernails, or skin sleepiness or unusual drowsiness slow or irregular heartbeat swelling in the legs and ankles, Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Difficulty having a bowel movement drowsiness lack or loss of strength relaxed and calm feeling Less common Abnormal dreams anxiety belching burning feeling

Drug Side Effects Summary

in the chest or stomach false or unusual sense of well-being heartburn hiccups indigestion stomach discomfort or upset tenderness in the stomach area trouble sleeping weight loss Rare Absent, missed, or irregular menstrual periods bad, unusual or unpleasant (after) taste bloated or full feeling body aches or pain change in taste change in walking and balance changes in vision clumsiness or unsteadiness continuous ringing or buzzing or other unexplained noise in the ears crying decreased interest in sexual intercourse deep or fast breathing with dizziness delusions of persecution, mistrust, suspiciousness, or combativeness dental caries or tooth decay depression difficulty with speaking dry skin dryness or soreness of the throat excess air or gas in the stomach or bowels excessive muscle tone feeling of constant movement of self or surroundings feeling of unreality general feeling of discomfort or illness headache, severe and throbbing hearing loss hoarseness inability to have or keep an erection increase in body movements increased appetite increased cough irritability loss in sexual ability, desire, drive, or performance loss of heat from the body loss of memory loss of strength or energy muscle stiffness, tension, tightness, pain, or weakness neck pain numbness of the feet, hands, and around the mouth passing of gas problems with memory quick to react or overreact emotionally rapidly changing moods red, swollen skin restlessness runny nose scaly skin sensation of spinning sense of detachment from self or body swelling or inflammation of the mouth voice changes Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Dazidox side effects

2404. This particular disease, Pain, interacts with the drug ETH-Oxydose and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Chills cold sweats confusion difficult or labored breathing dizziness, faintness, or lightheadedness when getting up

Drug Side Effects Summary

suddenly from a lying or sitting position fever tightness in the chest twitching Rare Bloating or swelling of the face, arms, hands, lower legs, or feet blood in the urine burning while urinating burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain cough decrease in the frequency of urination decreased urine output difficult or painful urination difficulty in passing urine (dribbling) difficulty with swallowing dizziness dry mouth fainting fast, irregular, pounding, or racing heartbeat or pulse feeling of warmth or heat flushing or redness of the skin, especially on the face and neck frequent urination headache hives, itching, or skin rash increase in heart rate increased thirst increased volume of pale, dilute urine lightheadedness muscle pain or cramps nausea puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rapid breathing rapid weight gain seizures severe constipation severe vomiting stomach pain sunken eyes sweating swollen, painful, or tender lymph glands in the neck, armpit, or groin thirst trembling or shaking of the hands or feet unusual tiredness or weakness unusual weight gain or loss vomiting wrinkled skin Incidence not known Blurred vision choking clay-colored stools cold, clammy skin dark urine darkening of the skin diarrhea fast, weak pulse gagging irregular, fast, slow, or shallow breathing loss of appetite loss of consciousness pale or blue lips, fingernails, or skin unpleasant breath odor very slow heartbeat yellow eyes or skin Symptoms of overdose Change in consciousness chest pain or discomfort cold, clammy skin constricted, pinpoint, or small pupils (black part of the eye) coughing that sometimes produces a pink frothy sputum decreased awareness or responsiveness increased sweating irregular, fast, or slow, or shallow breathing loss of consciousness no muscle tone or movement pale or blue lips, fingernails, or skin sleepiness or unusual drowsiness slow or irregular heartbeat swelling in the legs and ankles, Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any

Drug Side Effects Summary

questions about them: More common Difficulty having a bowel movement drowsiness lack or loss of strength relaxed and calm feeling Less common Abnormal dreams anxiety belching burning feeling in the chest or stomach false or unusual sense of well-being heartburn hiccups indigestion stomach discomfort or upset tenderness in the stomach area trouble sleeping weight loss Rare Absent, missed, or irregular menstrual periods bad, unusual or unpleasant (after) taste bloated or full feeling body aches or pain change in taste change in walking and balance changes in vision clumsiness or unsteadiness continuous ringing or buzzing or other unexplained noise in the ears crying decreased interest in sexual intercourse deep or fast breathing with dizziness delusions of persecution, mistrust, suspiciousness, or combativeness dental caries or tooth decay depression difficulty with speaking dry skin dryness or soreness of the throat excess air or gas in the stomach or bowels excessive muscle tone feeling of constant movement of self or surroundings feeling of unreality general feeling of discomfort or illness headache, severe and throbbing hearing loss hoarseness inability to have or keep an erection increase in body movements increased appetite increased cough irritability loss in sexual ability, desire, drive, or performance loss of heat from the body loss of memory loss of strength or energy muscle stiffness, tension, tightness, pain, or weakness neck pain numbness of the feet, hands, and around the mouth passing of gas problems with memory quick to react or overreact emotionally rapidly changing moods red, swollen skin restlessness runny nose scaly skin sensation of spinning sense of detachment from self or body swelling or inflammation of the mouth voice changes Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. ETH-Oxydose side effects

2405. This particular disease, Pain, interacts with the drug Hysingla ER and has the following side effects

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Stop using Hysingla ER and call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep

- a slow heart rate or weak pulse

- pain or burning when you urinate

- confusion, tremors, severe drowsiness

- a light-headed feeling, like you might pass out

- low cortisol levels --nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness

- or high levels of serotonin in the body - agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, diarrhea. Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea .

Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent. Common Hysingla ER side effects may include: constipation , nausea, vomiting

- dizziness, drowsiness, feeling tired

- headache

- or cold symptoms such as stuffy nose, sneezing, sore throat .

2406. This particular disease, Pain, interacts with the drug Oxaydo and has the following side effects

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Oxaydo may cause serious side effects. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep
- a slow heart rate or weak pulse
- a light-headed feeling, like you might pass out
- confusion, unusual thoughts or behavior
- seizure (convulsions)
- low cortisol levels-- nausea , vomiting , loss of appetite, dizziness, worsening tiredness or weakness
- or high levels of serotonin in the body-- agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, diarrhea . Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders. Common side effects of Oxaydo may include: drowsiness, headache , dizziness, tiredness
- or constipation , stomach pain, nausea, vomiting.

2407. This particular disease, Pain, interacts with the drug Oxydose and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Chills cold sweats confusion difficult or labored breathing dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position fever tightness in the chest twitching Rare Bloating or

Drug Side Effects Summary

swelling of the face, arms, hands, lower legs, or feet blood in the urine burning while urinating burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain cough decrease in the frequency of urination decreased urine output difficult or painful urination difficulty in passing urine (dribbling) difficulty with swallowing dizziness dry mouth fainting fast, irregular, pounding, or racing heartbeat or pulse feeling of warmth or heat flushing or redness of the skin, especially on the face and neck frequent urination headache hives, itching, or skin rash increase in heart rate increased thirst increased volume of pale, dilute urine lightheadedness muscle pain or cramps nausea puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rapid breathing rapid weight gain seizures severe constipation severe vomiting stomach pain sunken eyes sweating swollen, painful, or tender lymph glands in the neck, armpit, or groin thirst trembling or shaking of the hands or feet unusual tiredness or weakness unusual weight gain or loss vomiting wrinkled skin Incidence not known Blurred vision choking clay-colored stools cold, clammy skin dark urine darkening of the skin diarrhea fast, weak pulse gagging irregular, fast, slow, or shallow breathing loss of appetite loss of consciousness pale or blue lips, fingernails, or skin unpleasant breath odor very slow heartbeat yellow eyes or skin Symptoms of overdose Change in consciousness chest pain or discomfort cold, clammy skin constricted, pinpoint, or small pupils (black part of the eye) coughing that sometimes produces a pink frothy sputum decreased awareness or responsiveness increased sweating irregular, fast, or slow, or shallow breathing loss of consciousness no muscle tone or movement pale or blue lips, fingernails, or skin sleepiness or unusual drowsiness slow or irregular heartbeat swelling in the legs and ankles, Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Difficulty having a bowel movement drowsiness lack or loss of

Drug Side Effects Summary

strength relaxed and calm feeling Less common Abnormal dreams anxiety belching burning feeling in the chest or stomach false or unusual sense of well-being heartburn hiccups indigestion stomach discomfort or upset tenderness in the stomach area trouble sleeping weight loss Rare Absent, missed, or irregular menstrual periods bad, unusual or unpleasant (after) taste bloated or full feeling body aches or pain change in taste change in walking and balance changes in vision clumsiness or unsteadiness continuous ringing or buzzing or other unexplained noise in the ears crying decreased interest in sexual intercourse deep or fast breathing with dizziness delusions of persecution, mistrust, suspiciousness, or combativeness dental caries or tooth decay depression difficulty with speaking dry skin dryness or soreness of the throat excess air or gas in the stomach or bowels excessive muscle tone feeling of constant movement of self or surroundings feeling of unreality general feeling of discomfort or illness headache, severe and throbbing hearing loss hoarseness inability to have or keep an erection increase in body movements increased appetite increased cough irritability loss in sexual ability, desire, drive, or performance loss of heat from the body loss of memory loss of strength or energy muscle stiffness, tension, tightness, pain, or weakness neck pain numbness of the feet, hands, and around the mouth passing of gas problems with memory quick to react or overreact emotionally rapidly changing moods red, swollen skin restlessness runny nose scaly skin sensation of spinning sense of detachment from self or body swelling or inflammation of the mouth voice changes Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Oxydose side effects

2408. This particular disease, Pain, interacts with the drug Oxyfast and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and

Drug Side Effects Summary

death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Oxyfast may cause serious side effects. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep

- a slow heart rate or weak pulse
- a light-headed feeling, like you might pass out
- confusion, unusual thoughts or behavior
- seizure (convulsions)
- low cortisol levels-- nausea , vomiting , loss of appetite, dizziness, worsening tiredness or weakness
- or high levels of serotonin in the body-- agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, diarrhea . Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders. Common side effects of Oxyfast may include: drowsiness, headache , dizziness, tiredness
- or constipation , stomach pain, nausea, vomiting.

2409. This particular disease, Pain, interacts with the drug RoxyBond and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Chills cold sweats confusion difficult or labored breathing dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position fever tightness in the chest twitching Rare Bloating or swelling of the face, arms, hands, lower legs, or feet blood in the urine burning while urinating burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain

Drug Side Effects Summary

cough decrease in the frequency of urination decreased urine output difficult or painful urination difficulty in passing urine (dribbling) difficulty with swallowing dizziness dry mouth fainting fast, irregular, pounding, or racing heartbeat or pulse feeling of warmth or heat flushing or redness of the skin, especially on the face and neck frequent urination headache hives, itching, or skin rash increase in heart rate increased thirst increased volume of pale, dilute urine lightheadedness muscle pain or cramps nausea puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rapid breathing rapid weight gain seizures severe constipation severe vomiting stomach pain sunken eyes sweating swollen, painful, or tender lymph glands in the neck, armpit, or groin thirst trembling or shaking of the hands or feet unusual tiredness or weakness unusual weight gain or loss vomiting wrinkled skin Incidence not known Blurred vision choking clay-colored stools cold, clammy skin dark urine darkening of the skin diarrhea fast, weak pulse gagging irregular, fast, slow, or shallow breathing loss of appetite loss of consciousness pale or blue lips, fingernails, or skin unpleasant breath odor very slow heartbeat yellow eyes or skin Symptoms of overdose Change in consciousness chest pain or discomfort cold, clammy skin constricted, pinpoint, or small pupils (black part of the eye) coughing that sometimes produces a pink frothy sputum decreased awareness or responsiveness increased sweating irregular, fast, or slow, or shallow breathing loss of consciousness no muscle tone or movement pale or blue lips, fingernails, or skin sleepiness or unusual drowsiness slow or irregular heartbeat swelling in the legs and ankles, Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Difficulty having a bowel movement drowsiness lack or loss of strength relaxed and calm feeling Less common Abnormal dreams anxiety belching burning feeling in the chest or stomach false or unusual sense of well-being heartburn hiccups indigestion stomach

Drug Side Effects Summary

discomfort or upset tenderness in the stomach area trouble sleeping weight loss Rare Absent, missed, or irregular menstrual periods bad, unusual or unpleasant (after) taste bloated or full feeling body aches or pain change in taste change in walking and balance changes in vision clumsiness or unsteadiness continuous ringing or buzzing or other unexplained noise in the ears crying decreased interest in sexual intercourse deep or fast breathing with dizziness delusions of persecution, mistrust, suspiciousness, or combativeness dental caries or tooth decay depression difficulty with speaking dry skin dryness or soreness of the throat excess air or gas in the stomach or bowels excessive muscle tone feeling of constant movement of self or surroundings feeling of unreality general feeling of discomfort or illness headache, severe and throbbing hearing loss hoarseness inability to have or keep an erection increase in body movements increased appetite increased cough irritability loss in sexual ability, desire, drive, or performance loss of heat from the body loss of memory loss of strength or energy muscle stiffness, tension, tightness, pain, or weakness neck pain numbness of the feet, hands, and around the mouth passing of gas problems with memory quick to react or overreact emotionally rapidly changing moods red, swollen skin restlessness runny nose scaly skin sensation of spinning sense of detachment from self or body swelling or inflammation of the mouth voice changes Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. RoxyBond side effects

2410. This particular disease, Pain, interacts with the drug Xtampza ER and has the following side effects

- The possible side effects of are: constipation, nausea, sleepiness, vomiting, tiredness, headache, dizziness, abdominal pain. Call your healthcare provider if you have any of these symptoms and they are severe. trouble breathing, shortness of breath, fast heartbeat, chest pain, swelling of your face, tongue, or throat, extreme drowsiness, lightheadedness when changing positions, feeling faint, agitation, high body temperature, trouble walking, stiff muscles, or mental changes such as

Drug Side Effects Summary

confusion. These are not all the possible side effects. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Xtampza ER side effects

2411. This particular disease, Pain, interacts with the drug Zipsor and has the following side effects

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Stop using Zipsor and call your doctor at once if you have: the first sign of any skin rash, no matter how mild
- flu-like symptoms
- heart problems - swelling, rapid weight gain, feeling short of breath
- kidney problems - little or no urinating, painful or difficult urination, swelling in your arms or legs, feeling tired or short of breath
- liver problems - nausea , diarrhea , stomach pain (upper right side), tiredness, itching, dark urine, jaundice (yellowing of the skin or eyes)
- or signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Common Zipsor side effects may include: indigestion , gas, nausea, vomiting , stomach pain
- diarrhea, constipation
- headache, dizziness , drowsiness
- abnormal lab tests
- itching, sweating
- stuffy nose
- increased blood pressure
- or swelling or pain in your arms or legs.

2412. This particular disease, Pain, interacts with the drug Aloe Vera Burn Relief Spray with Lidocaine

Drug Side Effects Summary

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

- itching

- red, swollen, blistered, or peeling skin with or without fever

- wheezing

- tightness in the chest or throat

- trouble breathing, swallowing, or talking

- unusual hoarseness

- or swelling of the mouth, face, lips, tongue, or throat. Signs of infection like fever, chills, very bad sore throat, ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal. Signs of too much acid in the blood (acidosis) like confusion

- fast breathing

- fast heartbeat

- a heartbeat that does not feel normal

- very bad stomach pain, upset stomach, or throwing up

- feeling very sleepy

- shortness of breath

- or feeling very tired or weak. Signs of methemoglobinemia like a blue or gray color of the lips, nails, or skin

- a heartbeat that does not feel normal

- seizures

Drug Side Effects Summary

- very bad dizziness or passing out
- very bad headache
- feeling very sleepy
- feeling tired or weak
- or shortness of breath. This effect is rare but may be deadly if it happens. Trouble breathing, slow breathing, or shallow breathing. Very bad numbness and tingling. Feeling lightheaded, sleepy, confused, or having blurred eyesight. Seizures. Change in eyesight. Feeling nervous and excitable. Dizziness or passing out. Ringing in ears. Upset stomach or throwing up. Feeling hot or cold. Shakiness. Twitching. Slow heartbeat . Chest pain. Aloe Vera Burn Relief Spray with Lidocaine side effects

2413. This particular disease, Pain, interacts with the drug AneCream and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. AneCream may cause serious side effects. Call your doctor at once if you have: severe headache or vomiting
- severe burning, stinging, or irritation where the medicine was applied
- swelling or redness
- sudden dizziness or drowsiness after medicine is applied
- confusion, problems with speech or vision, ringing in your ears
- or unusual sensations of temperature. Common side effects include: mild irritation where the medication is applied
- or numbness in places where the medicine is accidentally applied.

2414. This particular disease, Pain, interacts with the drug Anestacon and has the following side effects

Drug Side Effects Summary

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Blistering, crusting, irritation, itching, or reddening of the skin cough cracked, dry, or scaly skin fast heartbeat fever hives or welts, itching, skin rash hoarseness irritation joint pain, stiffness, or swelling large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or genitals noisy breathing swelling of the eyelids, face, lips, hands, or feet tightness in the chest trouble breathing or swallowing Incidence not known Blistering, bruising, burning, discoloration, itching, redness, or swelling at the application site bluish-colored lips, fingernails, or palms blurred vision chest pain or discomfort cold, clammy, or pale skin confusion dark urine dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position drowsiness or dizziness false or unusual sense of well-being fear or nervousness headache irregular, fast or slow, or shallow breathing loss of consciousness no blood pressure or pulse pale or blue lips, fingernails, or skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rapid heart rate seizures shakiness in the legs, arms, hands, or feet slow or irregular heartbeat sore throat stopping of the heart trembling or shaking of the hands or feet twitching unusual bleeding or bruising unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Bad, unusual, or unpleasant (after) taste change in taste cold or numbness confusion about identity, place, and time continuing ringing or buzzing or other unexplained noise in the ears double vision hearing loss heat sensation lack or loss of strength metallic taste sleepiness or unusual drowsiness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your

Drug Side Effects Summary

healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Anestacon side effects

2415. This particular disease, Pain, interacts with the drug Astero and has the following side effects

- No side effects listed

2416. This particular disease, Pain, interacts with the drug Astramorph PF and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: More common Difficult or troubled breathing irregular, fast or slow, or shallow breathing pale or blue lips, fingernails, or skin shortness of breath very slow breathing Incidence not known Blurred vision convulsions decrease in frequency of urination decrease in the amount of urine difficulty in passing urine (dribbling) dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position painful urination sweating unusual tiredness or weakness Symptoms of overdose Bluish lips or skin dizziness fainting irregular heartbeat lightheadedness low blood pressure or pulse slow heartbeat unconsciousness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Absent, missed, or irregular menstrual periods anxiety confusion decreased interest in sexual intercourse delusions depersonalization difficulty having a bowel movement (stool) false or unusual sense of well-being hallucinations headache inability to have or keep an erection itching skin loss in sexual ability, desire, drive, or performance menstrual changes nausea and vomiting stopping of menstrual

Drug Side Effects Summary

bleeding Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Astramorph PF side effects

2417. This particular disease, Pain, interacts with the drug Bactine and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Bactine may cause serious side effects. Call your doctor at once if you have: severe headache or vomiting
- severe burning, stinging, or irritation where the medicine was applied
- swelling or redness
- sudden dizziness or drowsiness after medicine is applied
- confusion, problems with speech or vision, ringing in your ears
- or unusual sensations of temperature. Common side effects include: mild irritation where the medication is applied
- or numbness in places where the medicine is accidentally applied.

2418. This particular disease, Pain, interacts with the drug Belbuca and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Like other narcotic medicines, buprenorphine can slow your breathing. Death may occur if breathing becomes too weak. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep

Drug Side Effects Summary

- slow heartbeat or weak pulse
- a light-headed feeling, like you might pass out
- chest pain, fast heart rate , trouble breathing
- severe constipation
- opioid withdrawal symptoms - shivering, goose bumps, increased sweating , feeling hot or cold, runny nose , watery eyes, diarrhea , muscle pain
- low cortisol levels - nausea , vomiting , loss of appetite, dizziness, worsening tiredness or weakness
- or liver problems - nausea, upper stomach pain, itching , loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent. Common Belbuca side effects may include: constipation, nausea, vomiting
- headache , dizziness, drowsiness
- increased sweating
- sleep problems (insomnia)
- or pain anywhere in your body.

2419. This particular disease, Pain, interacts with the drug Biofreeze and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Biofreeze may cause serious side effects. Stop using

Drug Side Effects Summary

Biofreeze and call your doctor at once if you have: severe burning, stinging, redness, or irritation after using this medicine

- or pain, swelling, or blistering where the medicine was applied. Less serious side effects may be more likely, and you may have none at all.

2420. This particular disease, Pain, interacts with the drug butorphanol and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Butorphanol may cause serious side effects. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops
- a slow heart rate or weak pulse
- fast or pounding heartbeats
- problems with urination
- confusion, feeling like you are floating
- a light-headed feeling, like you might pass out
- or high levels of serotonin in the body-- agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting , diarrhea
- Serious breathing problems may be more likely in older adults and those who are debilitated or have wasting syndrome or chronic breathing disorders. Common side effects include: nausea, vomiting
- drowsiness, dizziness
- dry mouth

Drug Side Effects Summary

- or warmth or redness under the skin.

2421. This particular disease, Pain, interacts with the drug Butrans and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Stop using Butrans and call your doctor at once if you have: weak or shallow breathing, deep sighs, snoring that is new or unusual
- breathing that stops during sleep
- chest pain, fast heart rate , seizure (convulsions)
- a light-headed feeling, like you might pass out
- blisters, swelling, or severe irritation where the patch was worn
- adrenal gland problems - nausea , vomiting , loss of appetite, dizziness, feeling weak or tired
- or liver problems - upper stomach pain, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Seek medical attention right away if you have symptoms of serotonin syndrome , such as: agitation , hallucinations, fever, sweating , shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea . Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent. Common Butrans side effects may include: constipation , nausea, vomiting
- headache , dizziness, drowsiness, tiredness
- or redness, itching, or rash where the patch was worn.

Drug Side Effects Summary

2422. This particular disease, Pain, interacts with the drug CidalEaze and has the following side effects

- No side effects listed

2423. This particular disease, Pain, interacts with the drug DermacinRx Lidotral? and has the following side effects

- No side effects listed

2424. This particular disease, Pain, interacts with the drug DermacinRx Lido V Pak and has the following side effects

- No side effects listed

2425. This particular disease, Pain, interacts with the drug diphenhydramine and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
 - hives
 - itching
 - red, swollen, blistered, or peeling skin with or without fever
 - wheezing
 - tightness in the chest or throat
 - trouble breathing, swallowing, or talking
 - unusual hoarseness
 - or swelling of the mouth, face, lips, tongue, or throat. Very bad skin irritation. Diphenhydramine topical side effects

2426. This particular disease, Pain, interacts with the drug Duraclon and has the following side effects

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Duraclon may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- slow heartbeats
- weak or shallow breathing
- signs of infection around the infusion catheter, such as warmth, redness, or swelling
- fever
- or headache, neck stiffness, vomiting , increased sensitivity to light. Common side effects of Duraclon may include: dizziness, drowsiness
- depressed mood
- nausea , dry mouth
- or confusion.

2427. This particular disease, Pain, interacts with the drug Duramorph and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: More common Difficult or troubled breathing irregular, fast or slow, or shallow breathing pale or blue lips, fingernails, or skin shortness of breath very slow breathing Incidence not known Blurred vision convulsions decrease in frequency of urination decrease in the amount of urine difficulty in passing urine (dribbling) dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position painful urination sweating unusual tiredness or weakness Symptoms of overdose Bluish lips or skin dizziness fainting irregular heartbeat lightheadedness low blood pressure or pulse slow heartbeat unconsciousness Some side effects may occur that usually do not need medical attention. These

Drug Side Effects Summary

side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Absent, missed, or irregular menstrual periods anxiety confusion decreased interest in sexual intercourse delusions depersonalization difficulty having a bowel movement (stool) false or unusual sense of well-being hallucinations headache inability to have or keep an erection itching skin loss in sexual ability, desire, drive, or performance menstrual changes nausea and vomiting stopping of menstrual bleeding Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Duramorph side effects

2428. This particular disease, Pain, interacts with the drug Duramorph PF and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Duramorph PF may cause serious side effects. Call your doctor at once if you have: slow heart rate , sighing, shallow breathing, breathing that stops
- extreme drowsiness, feeling like you might pass out
- flushing (sudden warmth, redness, or tingly feeling)
- a seizure
- high levels of serotonin in the body-- agitation , hallucinations, fever, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , diarrhea

Drug Side Effects Summary

- or low cortisol levels-- nausea, vomiting , loss of appetite, dizziness, worsening tiredness or weakness. Serious breathing problems may be more likely in older adults and people who are debilitated or have wasting syndrome or chronic breathing disorders. Common side effects of Duramorph PF may include: breathing problems

- drowsiness, dizziness
- constipation , nausea, vomiting
- sweating
- or numbness , tingling, or cold feeling in your hands and feet.

2429. This particular disease, Pain, interacts with the drug Eha Lotion and has the following side effects:

- No side effects listed

2430. This particular disease, Pain, interacts with the drug Fentora and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Like other narcotic medicines, fentanyl can slow your breathing. Death may occur if breathing becomes too weak. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Call your doctor at once if you have: slow heart rate, sighing, shallow breathing, breathing that stops during sleep
- severe drowsiness, feeling like you might pass out
- confusion, extreme fear, unusual thoughts or behavior
- or low cortisol levels - nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss

Drug Side Effects Summary

of coordination, nausea, vomiting, or diarrhea. Serious side effects may be more likely in older adults and those who are malnourished or debilitated. Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent. Common Fentora side effects may include: headache, dizziness, drowsiness, pale skin, feeling weak or tired

- constipation, nausea, vomiting
- or swelling in your hands or feet.

2431. This particular disease, Pain, interacts with the drug Infumorph and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Infumorph may cause serious side effects. Call your doctor at once if you have: slow heart rate , sighing, shallow breathing, breathing that stops
- extreme drowsiness, feeling like you might pass out
- flushing (sudden warmth, redness, or tingly feeling)
- a seizure
- high levels of serotonin in the body-- agitation , hallucinations, fever, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , diarrhea
- or low cortisol levels-- nausea, vomiting , loss of appetite, dizziness, worsening tiredness or weakness. Serious breathing problems may be more likely in older adults and people who are debilitated or have wasting syndrome or chronic breathing disorders. Common side effects of Infumorph may include: breathing problems

Drug Side Effects Summary

- drowsiness, dizziness
- constipation , nausea, vomiting
- sweating
- or numbness , tingling, or cold feeling in your hands and feet.

2432. This particular disease, Pain, interacts with the drug Itch Relief and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad skin irritation. Itch Relief side effects

2433. This particular disease, Pain, interacts with the drug levorphanol and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake

Drug Side Effects Summary

up. Levorphanol may cause serious side effects. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops

- a slow heart rate or weak pulse
- a light-headed feeling, like you might pass out
- mood changes, unusual thoughts or behavior
- severe stomach pain, severe constipation
- high levels of serotonin in the body-- agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting , diarrhea
- or low cortisol levels-- nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness. Serious breathing problems may be more likely in older adults and those who are debilitated or have wasting syndrome or chronic breathing disorders. Common side effects of levorphanol may include: nausea, vomiting, stomach pain
- constipation
- dizziness, drowsiness
- tiredness
- or headache .

2434. This particular disease, Pain, interacts with the drug LidaMantle and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. LidaMantle may cause serious side effects. Call your doctor at once if you have: severe headache or vomiting
- severe burning, stinging, or irritation where the medicine was applied
- swelling or redness
- sudden dizziness or drowsiness after medicine is applied

Drug Side Effects Summary

- confusion, problems with speech or vision, ringing in your ears
- or unusual sensations of temperature. Common side effects include: mild irritation where the medication is applied
- or numbness in places where the medicine is accidentally applied.

2435. This particular disease, Pain, interacts with the drug Lidopac and has the following side effects

- No side effects listed

2436. This particular disease, Pain, interacts with the drug LidoRx and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. LidoRx may cause serious side effects. Call your doctor at once if you have: severe headache or vomiting
- severe burning, stinging, or irritation where the medicine was applied
- swelling or redness
- sudden dizziness or drowsiness after medicine is applied
- confusion, problems with speech or vision, ringing in your ears
- or unusual sensations of temperature. Common side effects include: mild irritation where the medication is applied
- or numbness in places where the medicine is accidentally applied.

2437. This particular disease, Pain, interacts with the drug Lidotrans 5 Pak and has the following side effects

- No side effects listed

2438. This particular disease, Pain, interacts with the drug Lidozion and has the following side effects

Drug Side Effects Summary

- No side effects listed

2439. This particular disease, Pain, interacts with the drug LMX 4 and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Blistering, crusting, irritation, itching, or reddening of the skin cough cracked, dry, or scaly skin fast heartbeat fever hives or welts, itching, skin rash hoarseness irritation joint pain, stiffness, or swelling large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or genitals noisy breathing swelling of the eyelids, face, lips, hands, or feet tightness in the chest trouble breathing or swallowing Incidence not known Blistering, bruising, burning, discoloration, itching, redness, or swelling at the application site bluish-colored lips, fingernails, or palms blurred vision chest pain or discomfort cold, clammy, or pale skin confusion dark urine dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position drowsiness or dizziness false or unusual sense of well-being fear or nervousness headache irregular, fast or slow, or shallow breathing loss of consciousness no blood pressure or pulse pale or blue lips, fingernails, or skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rapid heart rate seizures shakiness in the legs, arms, hands, or feet slow or irregular heartbeat sore throat stopping of the heart trembling or shaking of the hands or feet twitching unusual bleeding or bruising unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Bad, unusual, or unpleasant (after) taste change in taste cold or numbness confusion about identity, place, and time

Drug Side Effects Summary

continuing ringing or buzzing or other unexplained noise in the ears double vision hearing loss heat sensation lack or loss of strength metallic taste sleepiness or unusual drowsiness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. LMX 4 side effects

2440. This particular disease, Pain, interacts with the drug LMX 5 and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Blistering, crusting, irritation, itching, or reddening of the skin cough cracked, dry, or scaly skin fast heartbeat fever hives or welts, itching, skin rash hoarseness irritation joint pain, stiffness, or swelling large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or genitals noisy breathing swelling of the eyelids, face, lips, hands, or feet tightness in the chest trouble breathing or swallowing Incidence not known Blistering, bruising, burning, discoloration, itching, redness, or swelling at the application site bluish-colored lips, fingernails, or palms blurred vision chest pain or discomfort cold, clammy, or pale skin confusion dark urine dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position drowsiness or dizziness false or unusual sense of well-being fear or nervousness headache irregular, fast or slow, or shallow breathing loss of consciousness no blood pressure or pulse pale or blue lips, fingernails, or skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rapid heart rate seizures shakiness in the legs, arms, hands, or feet slow or irregular heartbeat sore throat stopping of the heart trembling or shaking of the hands or feet twitching unusual bleeding or bruising unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care

Drug Side Effects Summary

professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Bad, unusual, or unpleasant (after) taste change in taste cold or numbness confusion about identity, place, and time continuing ringing or buzzing or other unexplained noise in the ears double vision hearing loss heat sensation lack or loss of strength metallic taste sleepiness or unusual drowsiness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. LMX 5 side effects

2441. This particular disease, Pain, interacts with the drug Lydexa and has the following side effects

- No side effects listed

2442. This particular disease, Pain, interacts with the drug Medi-Quik Spray and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Medi-Quik Spray may cause serious side effects. Call your doctor at once if you have: severe headache or vomiting
- severe burning, stinging, or irritation where the medicine was applied
- swelling or redness
- sudden dizziness or drowsiness after medicine is applied
- confusion, problems with speech or vision, ringing in your ears
- or unusual sensations of temperature. Common side effects include: mild irritation where the medication is applied
- or numbness in places where the medicine is accidentally applied.

Drug Side Effects Summary

2443. This particular disease, Pain, interacts with the drug mefenamic acid and has the following side effects:

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. Stop using mefenamic acid and seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Mefenamic acid may cause serious side effects. Stop using mefenamic acid and call your doctor at once if you have: shortness of breath (even with mild exertion)
- swelling or rapid weight gain
- nausea , diarrhea , flu-like symptoms (fever, swollen glands, muscle aches, severe weakness)
- a skin rash, no matter how mild
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- o low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Common side effects of mefenamic acid may include: heartburn , stomach pain, gas, nausea, vomiting
- diarrhea, constipation
- or dizziness .

2444. This particular disease, Pain, interacts with the drug Methadone Diskets and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your

Drug Side Effects Summary

doctor immediately if any of the following side effects occur: Incidence not known Black, tarry stools bleeding gums blood in the urine or stools blurred vision bulging soft spot on the head of an infant change in the ability to see colors, especially blue or yellow changes in skin color chest discomfort or pain confusion cough coughing that sometimes produces a pink frothy sputum darkening of the skin decreased urine output diarrhea difficult, fast, noisy breathing difficulty with swallowing dilated neck veins dizziness dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position dry mouth extreme fatigue fainting fast, slow, or irregular heartbeat headache hives, itching, or skin rash increased sweating increased thirst irregular heartbeat irregular, fast or slow, or shallow breathing loss of appetite mental depression muscle pain or cramps nausea or vomiting numbness or tingling in the hands, feet, or lips pain pale or blue lips, fingernails, or skin pinpoint red spots on the skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue seizures sweating swelling of the face, fingers, feet, or lower legs tenderness trouble breathing trouble sleeping trouble urinating unusual bleeding or bruising unusual tiredness or weakness weight gain Symptoms of overdose Change in consciousness chest pain or discomfort cold, clammy skin constricted, pinpoint, or small pupils (black part of the eye) coughing that sometimes produces a pink frothy sputum decreased awareness or responsiveness increased sweating irregular, fast, or slow, or shallow breathing loss of consciousness no muscle tone or movement pale or blue lips, fingernails, or skin sleepiness or unusual drowsiness slow or irregular heartbeat swelling in the legs and ankles, Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Absent, missed, or irregular menstrual periods anxiety blurred or loss of vision confusion about identity, place, and time constipation decreased interest in sexual intercourse disturbed color

Drug Side Effects Summary

perception double vision false or unusual sense of well-being halos around lights inability to have or keep an erection irritability lack or loss of strength loss in sexual ability, desire, drive, or performance night blindness overbright appearance of lights redness, swelling, or soreness of the tongue restlessness stopping of menstrual bleeding tunnel vision weight changes welts Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Methadone Diskets side effects

2445. This particular disease, Pain, interacts with the drug Methadose and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Black, tarry stools bleeding gums blood in the urine or stools blurred vision bulging soft spot on the head of an infant change in the ability to see colors, especially blue or yellow changes in skin color chest discomfort or pain confusion cough coughing that sometimes produces a pink frothy sputum darkening of the skin decreased urine output diarrhea difficult, fast, noisy breathing difficulty with swallowing dilated neck veins dizziness dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position dry mouth extreme fatigue fainting fast, slow, or irregular heartbeat headache hives, itching, or skin rash increased sweating increased thirst irregular heartbeat irregular, fast or slow, or shallow breathing loss of appetite mental depression muscle pain or cramps nausea or vomiting numbness or tingling in the hands, feet, or lips pain pale or blue lips, fingernails, or skin pinpoint red spots on the skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue seizures sweating swelling of the face, fingers, feet, or lower legs tenderness trouble breathing trouble sleeping trouble urinating unusual bleeding or bruising unusual tiredness or weakness weight gain Symptoms of overdose Change in consciousness chest pain or discomfort

Drug Side Effects Summary

cold, clammy skin constricted, pinpoint, or small pupils (black part of the eye) coughing that sometimes produces a pink frothy sputum decreased awareness or responsiveness increased sweating irregular, fast, or slow, or shallow breathing loss of consciousness no muscle tone or movement pale or blue lips, fingernails, or skin sleepiness or unusual drowsiness slow or irregular heartbeat swelling in the legs and ankles, Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Absent, missed, or irregular menstrual periods anxiety blurred or loss of vision confusion about identity, place, and time constipation decreased interest in sexual intercourse disturbed color perception double vision false or unusual sense of well-being halos around lights inability to have or keep an erection irritability lack or loss of strength loss in sexual ability, desire, drive, or performance night blindness overbright appearance of lights redness, swelling, or soreness of the tongue restlessness stopping of menstrual bleeding tunnel vision weight changes welts Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Methadose side effects

2446. This particular disease, Pain, interacts with the drug Mitigo and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake

Drug Side Effects Summary

up. Mitigo may cause serious side effects. Call your doctor at once if you have: slow heart rate , sighing, shallow breathing, breathing that stops

- extreme drowsiness, feeling like you might pass out
- flushing (sudden warmth, redness, or tingly feeling)
- a seizure
- high levels of serotonin in the body-- agitation , hallucinations, fever, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , diarrhea
- or low cortisol levels-- nausea, vomiting , loss of appetite, dizziness, worsening tiredness or weakness. Serious breathing problems may be more likely in older adults and people who are debilitated or have wasting syndrome or chronic breathing disorders. Common side effects of Mitigo may include: breathing problems
- drowsiness, dizziness
- constipation , nausea, vomiting
- sweating
- or numbness , tingling, or cold feeling in your hands and feet.

2447. This particular disease, Pain, interacts with the drug Moxicaine and has the following side effects

- No side effects listed

2448. This particular disease, Pain, interacts with the drug nalbuphine and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Tell your caregiver right

Drug Side Effects Summary

away if you have: shallow breathing, breathing that stops during sleep

- a light-headed feeling, like you might pass out
- severe drowsiness
- severe constipation
- low cortisol levels-- nausea , vomiting , loss of appetite, dizziness, worsening tiredness or weakness
- or high levels of serotonin in the body-- agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, diarrhea . Serious breathing problems may be more likely in older adults and those who are debilitated or have wasting syndrome or chronic breathing disorders. Common side effects of nalbuphine may include:
drowsiness
- dizziness, spinning sensation
- dry mouth
- headache
- sweating
- cold, clammy skin
- or nausea, vomiting.

2449. This particular disease, Pain, interacts with the drug RadiaGuard and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. RadiaGuard may cause serious side effects. Call your doctor at once if you have: severe headache or vomiting
- severe burning, stinging, or irritation where the medicine was applied
- swelling or redness

Drug Side Effects Summary

- sudden dizziness or drowsiness after medicine is applied
- confusion, problems with speech or vision, ringing in your ears
- or unusual sensations of temperature. Common side effects include: mild irritation where the medication is applied
- or numbness in places where the medicine is accidentally applied.

2450. This particular disease, Pain, interacts with the drug RectiCare and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. RectiCare may cause serious side effects. Call your doctor at once if you have: severe headache or vomiting
- severe burning, stinging, or irritation where the medicine was applied
- swelling or redness
- sudden dizziness or drowsiness after medicine is applied
- confusion, problems with speech or vision, ringing in your ears
- or unusual sensations of temperature. Common side effects include: mild irritation where the medication is applied
- or numbness in places where the medicine is accidentally applied.

2451. This particular disease, Pain, interacts with the drug Regenecare HA Spray and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Regenecare HA Spray may cause serious side effects. Call your doctor at once if you have: severe headache or vomiting
- severe burning, stinging, or irritation where the medicine was applied

Drug Side Effects Summary

- swelling or redness
- sudden dizziness or drowsiness after medicine is applied
- confusion, problems with speech or vision, ringing in your ears
- or unusual sensations of temperature. Common side effects include: mild irritation where the medication is applied
- or numbness in places where the medicine is accidentally applied.

2452. This particular disease, Pain, interacts with the drug Sprix and has the following side effects:

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling , feeling short of breath. Stop using Sprix and call your doctor at once if you have: any skin rash, no matter how mild
- high blood pressure - severe headache , blurred vision, pounding in your neck or ears
- heart problems - swelling, rapid weight gain, feeling short of breath
- kidney problems - little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- liver problems - nausea , upper stomach pain, itching, tiredness, flu-like symptoms, dark urine, jaundice (yellowing of the skin or eyes)
- low red blood cells (anemia) - pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet
- or signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Common Sprix side effects may include: heartburn , stomach pain, gas, nausea, vomiting
- diarrhea , constipation
- slow heartbeats
- decreased urination

Drug Side Effects Summary

- abnormal liver function tests
- increased blood pressure
- pain or irritation in your nose
- runny nose
- watery eyes
- throat irritation
- or rash.

2453. This particular disease, Pain, interacts with the drug Topicaïne and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bluish color of the fingernails, lips, skin, palms, or nail beds Incidence not known Blistering, burning, crusting, dryness, or flaking of the skin cracking, itching, redness, or stinging of the skin dark urine difficulty with breathing difficulty with walking dizziness or lightheadedness fainting fever headache inability to feel hands and feet irritability irritation of the nose itching, scaling, severe redness, soreness, or swelling of the skin pale skin rapid heart rate red, sore eyes shortness of breath sore throat unusual bleeding or bruising unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness unusual tiredness or weakness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Benzocaine topical side effects

2454. This particular disease, Pain, interacts with the drug Tranzarel and has the following side effects

- No side effects listed

Drug Side Effects Summary

2455. This particular disease, Pain, interacts with the drug Xylocaine Topical and has the following

- No side effects listed

2456. This particular disease, Pain, interacts with the drug Zionodil and has the following side effects

- No side effects listed

2457. This particular disease, Pain, interacts with the drug acetaminophen / benzhydrocodone and

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking acetaminophen and benzhydrocodone and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Acetaminophen and benzhydrocodone may cause serious side effects. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops
- a light-headed feeling, like you might pass out
- liver problems--nausea, upper stomach pain, tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- low cortisol levels-- nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness
- or high levels of serotonin in the body-- agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, diarrhea . Serious breathing problems may be more likely in older adults and in those who are debilitated or have

Drug Side Effects Summary

wasting syndrome or chronic breathing disorders. Common side effects include: dizziness, drowsiness, feeling tired

- nausea, vomiting, stomach pain
- constipation
- or headache .

2458. This particular disease, Pain, interacts with the drug acetaminophen / caffeine / dihydrocodeine

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking acetaminophen, caffeine, and dihydrocodeine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Acetaminophen, caffeine, and dihydrocodeine may cause serious side effects. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep
- chest pain, fast or pounding heartbeats, feeling light-headed, fainting
- confusion, extreme drowsiness, unusual thoughts or behavior
- painful or difficult urination
- severe constipation
- liver problems-- nausea , upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or low cortisol levels-- nausea, vomiting , loss of appetite, dizziness, worsening tiredness or

Drug Side Effects Summary

weakness. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea . Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent. Common side effects include: dizziness, drowsiness, tiredness, headache

- nausea, vomiting, stomach pain, constipation
- sweating, itching
- or skin rash .

2459. This particular disease, Pain, interacts with the drug acetaminophen / ibuprofen and has the following side effects:

- swelling, rapid weight gain, feeling short of breath
- little or no urination
- loss of appetite, new or worsening stomach pain
- bloody or tarry stools
- coughing up blood or vomit that looks like coffee grounds
- or dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Less serious side effects may be more likely, and you may have none at all.

2460. This particular disease, Pain, interacts with the drug acetaminophen / phenyltoloxamine / salicylates and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

Drug Side Effects Summary

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of bleeding like throwing up or coughing up blood
- vomit that looks like coffee grounds
- blood in the urine
- black, red, or tarry stools
- bleeding from the gums
- abnormal vaginal bleeding
- bruises without a cause or that get bigger
- or bleeding you cannot stop. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. A fast heartbeat . A heartbeat that does not feel normal. Very bad dizziness or passing out. Ringing in ears. Not able to pass urine or change in how much urine is passed. Change in balance. Not able to focus. Seizures . Blurred eyesight. Shakiness. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Acetaminophen / phenyltoloxamine / salicylamide

Drug Side Effects Summary

side effects

2461. This particular disease, Pain, interacts with the drug ActivICE and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. ActivICE may cause serious side effects. Stop using ActivICE and call your doctor at once if you have: severe burning, stinging, redness, or irritation after using this medicine
- or pain, swelling, or blistering where the medicine was applied. Less serious side effects may be more likely, and you may have none at all.

2462. This particular disease, Pain, interacts with the drug alfentanil and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: More common Blurred vision chest pain or discomfort confusion difficult or troubled breathing dizziness dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position fainting fast, slow, pounding, or irregular heartbeat or pulse headache irregular, fast or slow, or shallow breathing lightheadedness, dizziness, or fainting nervousness not breathing pale or blue lips, fingernails, or skin pounding in the ears sweating trouble breathing unusual tiredness or weakness Rare Cough noisy breathing rigid muscles of the neck, arms, and legs tightness in the chest Incidence not known Agitation cough darkening of the skin diarrhea difficulty with swallowing fainting fever hives, itching, or skin rash loss of appetite nausea overactive reflexes poor coordination puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue restlessness shivering sweating talking or acting with excitement you cannot control trembling or shaking twitching vomiting Some side effects may

Drug Side Effects Summary

occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Drowsiness sleepiness Rare False or unusual sense of well-being Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Alfentanil side effects

2463. This particular disease, Pain, interacts with the drug Alka-Seltzer Wake-Up Call! and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Alka-Seltzer Morning Relief may cause serious side effects. Stop using Alka-Seltzer Morning Relief and call your doctor at once if you have: ringing in your ears, confusion, hallucinations, rapid breathing, seizure (convulsions)
- severe nausea , vomiting , or stomach pain
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- fever lasting longer than 3 days
- or swelling, or pain lasting longer than 10 days. Common side effects of Alka-Seltzer Morning Relief may include: upset stomach , heartburn
- drowsiness
- or mild headache .

2464. This particular disease, Pain, interacts with the drug Anacin and has the following side effects:

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Anacin and call your doctor at once if you have: ringing in your ears, confusion, hallucinations, rapid breathing, seizure (convulsions)
- severe nausea, vomiting, or stomach pain
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- fever lasting longer than 3 days
- or swelling, or pain lasting longer than 10 days. Common Anacin side effects may include: upset stomach, heartburn
- drowsiness
- or mild headache.

2465. This particular disease, Pain, interacts with the drug Anbesol and has the following side effects

- Anbesol Gel used in the mouth may cause a condition in which the oxygen in your body tissues can become dangerously low. This is a potentially fatal condition called methemoglobinemia (met-HEEM-oh glo-bin-EE-mee-a). This condition may occur after only one use of benzocaine or after several uses. Signs and symptoms may occur within minutes or up to 2 hours after using Anbesol Gel in the mouth or throat. GET EMERGENCY MEDICAL HELP IF YOU HAVE: a headache, tiredness, confusion
- fast heartbeats
- feeling light-headed or short of breath
- and pale, blue, or gray appearance of your skin, lips, or fingernails. hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Anbesol Gel and call your doctor at once if you have: severe burning, stinging, or sensitivity where the medicine is applied
- swelling, warmth, or redness

Drug Side Effects Summary

- or oozing, blistering, or any signs of infection. Common Anbesol Gel side effects may include: mild stinging, burning, or itching where the medicine is applied
- skin tenderness or redness
- or dry white flakes where the medicine was applied.

2466. This particular disease, Pain, interacts with the drug Apadaz and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Apadaz and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep
- a light-headed feeling, like you might pass out
- liver problems - nausea, upper stomach pain, tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- low cortisol levels - nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness
- or high levels of serotonin in the body - agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, diarrhea. Serious side effects may be more likely in older adults and those who are malnourished or debilitated. Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent. Common Apadaz side effects include:

Drug Side Effects Summary

dizziness, drowsiness, feeling tired

- nausea, vomiting, stomach pain
- constipation
- or headache .

2467. This particular disease, Pain, interacts with the drug Arctic Relief and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Arctic Relief may cause serious side effects. Stop using Arctic Relief and call your doctor at once if you have: severe burning, stinging, or irritation after using this medicine
- or breathing problems after using this medicine on the chest or throat. Less serious side effects may be more likely, and you may have none at all.

2468. This particular disease, Pain, interacts with the drug Aspercreme Max No Mess Roll-On and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking

Drug Side Effects Summary

- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad skin irritation. Aspercreme Max No Mess Roll-On side effects

2469. This particular disease, Pain, interacts with the drug Aspercreme Pain Relieving Heat Gel and

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Aspercreme Heat Pain Relieving may cause serious side effects. Stop using Aspercreme Heat Pain Relieving and call your doctor at once if you have: severe burning, stinging, redness, or irritation after using this medicine
- or pain, swelling, or blistering where the medicine was applied. Less serious side effects may be more likely, and you may have none at all.

2470. This particular disease, Pain, interacts with the drug aspirin / caffeine / dihydrocodeine and h

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Agitation anxiety bleeding gums bloating bloody or black, tarry stools bloody or cloudy urine blue lips and fingernails blurred vision chest pain or discomfort chills cold sweats confusion constipation continuing ringing or buzzing or other unexplained noise in the ears cool, pale skin coughing that sometimes produces a pink frothy sputum dark urine darkening of the skin depression diarrhea difficult, fast, noisy breathing dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position drowsiness dry mouth fast, pounding, or irregular heartbeat or pulse fast, slow, or irregular heartbeat fever flushed, dry skin fruit-like breath odor general tiredness and weakness greatly decreased frequency of urination or amount of urine headache headache, sudden and severe hearing loss

Drug Side Effects Summary

heartburn increased hunger increased sweating increased thirst increased urination indigestion large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or genitals light-colored stools loss of appetite loss of consciousness muscle cramps, spasms, pain, weakness, or stiffness muscle tremors nausea nervousness nightmares numbness and tingling of the face, lips, hands, and feet overactive reflexes pains in the stomach, side, or abdomen, possibly radiating to the back pinpoint red spots on the skin poor coordination rapid, deep breathing restlessness seizures severe stomach pain shakiness shivering skin rash slurred speech stomach pain, cramping, or burning sunken eyes swelling in the legs, feet, or ankles trembling or shaking trouble with breathing twitching unexplained weight loss unusual bleeding or bruising unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness upper right abdominal or stomach pain vomiting vomiting of blood or material that looks like coffee grounds vomiting, severe or continuous weakness or heaviness of the legs wrinkled skin yellow eyes or skin Symptoms of overdose Chest pain or discomfort cold and clammy skin coughing that sometimes produces a pink frothy sputum difficult, fast, noisy breathing extreme sleepiness increased sweating irregular, fast or slow, or shallow breathing lightheadedness, dizziness or fainting pale or blue lips, fingernails, or skin sleepiness or unusual drowsiness swelling in the legs and ankles Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Belching clumsiness low body temperature sleepiness weak or feeble pulse weight gain Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Aspirin / caffeine / dihydrocodeine side effects

Drug Side Effects Summary

2471. This particular disease, Pain, interacts with the drug aspirin / caffeine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Aspirin and caffeine may cause serious side effects. Stop using aspirin and caffeine and call your doctor at once if you have: ringing in your ears, confusion, hallucinations, rapid breathing, seizure (convulsions)
- severe nausea , vomiting , or stomach pain
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- fever lasting longer than 3 days
- or swelling, or pain lasting longer than 10 days. Common side effects of aspirin and caffeine may include: upset stomach , heartburn
- drowsiness
- or mild headache .

2472. This particular disease, Pain, interacts with the drug aspirin / carisoprodol / codeine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Aspirin may cause stomach or intestinal bleeding, which can be fatal. This can occur without warning while you are taking this medicine. Aspirin, carisoprodol, and codeine may cause serious side effects. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing
- a slow heart rate or weak pulse
- a light-headed feeling, like you might pass out

Drug Side Effects Summary

- confusion, unusual thoughts or behavior
- easy bruising or bleeding (nosebleeds , bleeding gums)
- severe constipation
- symptoms of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- or low cortisol levels-- nausea , vomiting, loss of appetite, dizziness, worsening tiredness or weakness. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea . Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Common side effects of aspirin, carisoprodol, and codeine may include: dizziness, drowsiness, tiredness
- nausea, vomiting, stomach pain, constipation
- or headache .

2473. This particular disease, Pain, interacts with the drug aspirin / oxycodone and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Agitation bleeding gums bloating blood in the urine or stools bloody, black, or tarry stools blue lips, fingernails, or skin blurred vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings change in consciousness or confusion chest pain, tightness, or discomfort chills clay-colored stools confusion cough coughing or vomiting blood dark urine darkening of the skin decrease in urine volume or frequency decreased appetite depression difficult, fast, noisy breathing difficulty in passing urine (dribbling) difficulty having a bowel movement (stool) difficulty with swallowing dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position

Drug Side Effects Summary

drowsiness dry mouth fainting fast, slow, irregular, pounding, or racing heartbeat or pulse feeling of hostility or irritability feeling of warmth feeling that something terrible will happen fever headache, sudden, severe heartburn hives, itching, or skin rash increased menstrual flow or vaginal bleeding increased sweating indigestion irregular, fast, slow, or shallow breathing large, flat, blue or purplish patches on the skin large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or genitals loss of consciousness muscle cramps, pain, stiffness, weakness, or tremors nausea nosebleeds numbness or tingling in the hands, feet, or lips overactive reflexes painful or difficult urination pains in the stomach, side, or abdomen, possibly radiating to the back pale skin pinpoint red or purple spots on the skin poor coordination prolonged bleeding from cuts puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red or black, tarry stools or dark urine restlessness seizures shivering sleepiness sunken eyes sweating swelling of the face, ankles, hands, feet, or lower legs stomach pain, cramping, or tenderness talking or acting with excitement you cannot control thirst trembling or shaking twitching unusual bleeding or bruising unusual tiredness or weakness vomiting vomiting of material that looks like coffee grounds, severe and continuing weak or feeble pulse weakness or heaviness of the legs weight gain wrinkled skin yellow eyes or skin

Symptoms of overdose Blue lips and fingernails blurred vision change in consciousness chest pain cold, clammy skin confusion continuing ringing or buzzing or other unexplained noise in the ears coughing that sometimes produces a pink frothy sputum decreased awareness or responsiveness diarrhea dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position drowsiness enlarged pupils extremely high fever or body temperature fast, weak heartbeat hearing loss increase in heart rate increased sweating loss of consciousness pale skin restlessness severe sleepiness swelling in legs and ankles unusual tiredness or weakness

Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your

Drug Side Effects Summary

health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Relaxed and calm feeling sleepiness Incidence not known Belching bloated, full feeling blurred or loss of vision change in color perception cold sweats constricted, pinpoint, or small pupils (black part of the eye) cool, pale skin double vision excess air or gas in the stomach false or unusual sense of well-being flushed, dry skin fruit-like breath odor halos around lights increased hunger or thirst increased urination lack or loss of strength night blindness nightmares overbright appearance of lights red eyes redness of the skin seeing, hearing, or feeling things that are not there shakiness slurred speech trouble sleeping tunnel vision unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness weight loss Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Aspirin / oxycodone side effects

2474. This particular disease, Pain, interacts with the drug Back & Body Extra Strength and has the

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness

Drug Side Effects Summary

- or swelling of the mouth, face, lips, tongue, or throat. Signs of bleeding like throwing up or coughing up blood
- vomit that looks like coffee grounds
- blood in the urine
- black, red, or tarry stools
- bleeding from the gums
- abnormal vaginal bleeding
- bruises without a cause or that get bigger
- or bleeding you cannot stop. Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of high potassium levels like a heartbeat that does not feel normal
- feeling confused
- feeling weak, lightheaded, or dizzy
- feeling like passing out
- numbness or tingling
- or shortness of breath. Signs of too much acid in the blood (acidosis) like confusion
- fast breathing
- fast heartbeat
- a heartbeat that does not feel normal
- very bad stomach pain, upset stomach, or throwing up
- feeling very sleepy
- shortness of breath
- or feeling very tired or weak. Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight. Very bad dizziness or passing out.

Drug Side Effects Summary

Very bad headache . Ringing in the ears, hearing loss , or any other changes in hearing. Feeling agitated. Seizures . A fast heartbeat. Back & Body Extra Strength side effects

2475. This particular disease, Pain, interacts with the drug Bayer Back & Body and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Bayer Back & Body and call your doctor at once if you have: ringing in your ears, confusion, hallucinations, rapid breathing, seizure (convulsions)
- severe nausea , vomiting , or stomach pain
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- fever lasting longer than 3 days
- or swelling, or pain lasting longer than 10 days. Common side effects of Bayer Back & Body may include: upset stomach , heartburn
- drowsiness
- or mild headache .

2476. This particular disease, Pain, interacts with the drug Bayer Extra Strength Back & Body and h

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever

Drug Side Effects Summary

- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of bleeding like throwing up or coughing up blood
- vomit that looks like coffee grounds
- blood in the urine
- black, red, or tarry stools
- bleeding from the gums
- abnormal vaginal bleeding
- bruises without a cause or that get bigger
- or bleeding you cannot stop. Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of high potassium levels like a heartbeat that does not feel normal
- feeling confused
- feeling weak, lightheaded, or dizzy
- feeling like passing out
- numbness or tingling
- or shortness of breath. Signs of too much acid in the blood (acidosis) like confusion
- fast breathing
- fast heartbeat
- a heartbeat that does not feel normal
- very bad stomach pain, upset stomach, or throwing up

Drug Side Effects Summary

- feeling very sleepy
- shortness of breath
- or feeling very tired or weak. Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight. Very bad dizziness or passing out. Very bad headache . Ringing in the ears, hearing loss , or any other changes in hearing. Feeling agitated. Seizures . A fast heartbeat. Bayer Extra Strength Back & Body side effects

2477. This particular disease, Pain, interacts with the drug BC Arthritis and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. BC Arthritis may cause serious side effects. Stop using BC Arthritis and call your doctor at once if you have: ringing in your ears, confusion, hallucinations, rapid breathing, seizure (convulsions)
- severe nausea , vomiting , or stomach pain
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- fever lasting longer than 3 days
- or swelling, or pain lasting longer than 10 days. Common side effects of BC Arthritis may include: upset stomach , heartburn
- drowsiness
- or mild headache .

2478. This particular disease, Pain, interacts with the drug BC Fast Pain Relief and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs

Drug Side Effects Summary

of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of bleeding like throwing up or coughing up blood
- vomit that looks like coffee grounds
- blood in the urine
- black, red, or tarry stools
- bleeding from the gums
- abnormal vaginal bleeding
- bruises without a cause or that get bigger
- or bleeding you cannot stop. Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of high potassium levels like a heartbeat that does not feel normal
- feeling confused
- feeling weak, lightheaded, or dizzy
- feeling like passing out
- numbness or tingling
- or shortness of breath. Signs of too much acid in the blood (acidosis) like confusion

Drug Side Effects Summary

- fast breathing
- fast heartbeat
- a heartbeat that does not feel normal
- very bad stomach pain, upset stomach, or throwing up
- feeling very sleepy
- shortness of breath
- or feeling very tired or weak. Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight. Very bad dizziness or passing out. Very bad headache . Ringing in the ears, hearing loss , or any other changes in hearing. Feeling agitated. Seizures . A fast heartbeat. BC Fast Pain Relief side effects

2479. This particular disease, Pain, interacts with the drug Bengay Cold Therapy Gel and has the fo

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad skin irritation. Bengay Cold Therapy Gel side effects

Drug Side Effects Summary

2480. This particular disease, Pain, interacts with the drug Bengay Ultra Strength Pain Relieving Pa

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. BENGAY Pain Relieving may cause serious side effects. Stop using BENGAY Pain Relieving and call your doctor at once if you have: severe burning, stinging, redness, or irritation after using this medicine
- or pain, swelling, or blistering where the medicine was applied. Less serious side effects may be more likely, and you may have none at all.

2481. This particular disease, Pain, interacts with the drug BENGAY Vanishing Scent and has the fo

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad skin irritation. BENGAY Vanishing Scent side effects

2482. This particular disease, Pain, interacts with the drug Bengay Zero Degrees and has the follow

Drug Side Effects Summary

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad skin irritation. Bengay Zero Degrees side effects

2483. This particular disease, Pain, interacts with the drug Benzo-Jel and has the following side effects

- No side effects listed

2484. This particular disease, Pain, interacts with the drug benzocaine and has the following side effects

- Benzocaine topical used in the mouth may cause a condition in which the oxygen in your body tissues can become dangerously low. This is a potentially fatal condition called methemoglobinemia (met-HEEM-oh glo-bin-EE-mee-a). This condition may occur after only one use of benzocaine or after several uses. Signs and symptoms may occur within minutes or up to 2 hours after using benzocaine topical in the mouth or throat. GET EMERGENCY MEDICAL HELP IF YOU HAVE: a headache, tiredness, confusion
- fast heartbeats

Drug Side Effects Summary

- feeling light-headed or short of breath
- and pale, blue, or gray appearance of your skin, lips, or fingernails. hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Benzocaine topical may cause serious side effects. Stop using benzocaine topical and call your doctor at once if you have: severe burning, stinging, or sensitivity where the medicine is applied
- swelling, warmth, or redness
- or oozing, blistering, or any signs of infection. Common side effects of benzocaine topical may include: mild stinging, burning, or itching where the medicine is applied
- skin tenderness or redness
- or dry white flakes where the medicine was applied.

2485. This particular disease, Pain, interacts with the drug Bio-Therm Pain Relieving Lotion and has

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Irritation where Bio-Therm Pain Relieving

Drug Side Effects Summary

Lotion (methyl salicylate, menthol, and capsaicin lotion) is used. Very bad burning or burning that does not go away. Bio-Therm Pain Relieving Lotion side effects

2486. This particular disease, Pain, interacts with the drug Blue-Emu Maximum Strength Arthritis, M

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad skin irritation. Blue-Emu Maximum Strength Arthritis, Muscle & Joint Pain Relief side effects

2487. This particular disease, Pain, interacts with the drug Blue Gel and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Blue Gel may cause serious side effects. Stop using Blue Gel and call your doctor at once if you have: severe burning, stinging, redness, or irritation after using this medicine
- or pain, swelling, or blistering where the medicine was applied. Less serious side effects may be

Drug Side Effects Summary

more likely, and you may have none at all.

2488. This particular disease, Pain, interacts with the drug bupivacaine liposome and has the following side effects:

- hives , red rash, itching
- sneezing, difficulty breathing
- severe dizziness , vomiting
- swelling of your face, lips, tongue, or throat. You will be watched closely after receiving bupivacaine liposome, to make sure you do not have a reaction to the medicine. Tell your caregivers at once if you have any of these signs of a serious side effect: ringing in your ears
- drowsiness, feeling restless or anxious
- feeling like you might pass out
- speech or vision problems, a metallic taste in your mouth
- numbness or tingling around your mouth
- fast or slow heart rate , feeling short of breath, feeling unusually hot or cold
- tremors, twitching, mood changes
- ongoing numbness, weakness, or loss of movement where the medicine was injected
- or joint pain or stiffness, or weakness in any part of your body for months after your surgery. You may still feel numb or be unable to move the numbed area for up to 5 days after you are treated with bupivacaine liposome. Common side effects include: nausea , vomiting
- constipation
- or fever.

2489. This particular disease, Pain, interacts with the drug Capsagel Extra Strength and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if

Drug Side Effects Summary

you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Capsagel Extra Strength side effects

2490. This particular disease, Pain, interacts with the drug capsaicin / diclofenac and has the following

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: a skin rash, no matter how mild
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Common side effects of capsaicin and diclofenac may include: skin rash, itching, redness, blistering, tingling, or other irritation where the medicine was applied
- or dryness or hardening of treated skin.

Drug Side Effects Summary

2491. This particular disease, Pain, interacts with the drug capsaicin / lidocaine / menthol / methyl

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Wash the skin and get medical attention right away if you have severe burning, pain, swelling, or blistering of the skin where you applied this medicine. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: pale skin, blue-colored lips
- headache , confusion
- or rapid heartbeats. Common side effects may include a mild burning sensation that can last for several hours or days, especially after your first use of this medicine.

2492. This particular disease, Pain, interacts with the drug capsaicin / menthol / methyl salicylate and

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Irritation where methyl salicylate, menthol, and capsaicin cream and ointment is used. Very bad burning or burning that does not go away.

Drug Side Effects Summary

Capsaicin / menthol / methyl salicylate topical side effects

2493. This particular disease, Pain, interacts with the drug Capsinac and has the following side effects

- No side effects listed

2494. This particular disease, Pain, interacts with the drug celecoxib / tramadol and has the following side effects

- agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting , or diarrhea . Stop using celecoxib and tramadol and seek medical treatment right away if you have a serious drug reaction that can affect many parts of your body. Symptoms may include skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Celecoxib and tramadol can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Celecoxib and tramadol may cause serious side effects. Call your doctor at once if you have: severe headache , blurred vision, pounding in your neck or ears, or severe constipation

- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath

- decreased adrenal gland hormones --nausea, vomiting, stomach pain, loss of appetite, feeling tired or light-headed, muscle or joint pain, skin discoloration, craving salty foods

- signs of a heart attack or stroke--chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, leg swelling, feeling short of breath

- or low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Celecoxib and tramadol may cause serious side effects. Stop using celecoxib and tramadol and call your doctor at once if you have: low blood sodium--headache, confusion, problems with thinking or memory, weakness, feeling unsteady

Drug Side Effects Summary

- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems--nausea, stomach pain (upper right side), itching, tiredness, dark urine, jaundice (yellowing of the skin or eyes)
- or heart problems--swelling, rapid weight gain, feeling short of breath. Common side effects may include nausea, vomiting, dizziness, headache, and drowsiness.

2495. This particular disease, Pain, interacts with the drug Cepacol Fizzlers and has the following s

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of methemoglobinemia like a blue or gray color of the lips, nails, or skin
- a heartbeat that does not feel normal
- seizures
- very bad dizziness or passing out
- very bad headache

Drug Side Effects Summary

- feeling very sleepy
- feeling tired or weak
- or shortness of breath. This effect is rare but may be deadly if it happens. Very bad mouth pain. When treating a very sore throat , if signs last more than 2 days, or happen with fever, headache, rash, upset stomach , or throwing up. Cepacol Fizzlers side effects

2496. This particular disease, Pain, interacts with the drug Chiggerex and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bluish color of the fingernails, lips, skin, palms, or nail beds Incidence not known Blistering, burning, crusting, dryness, or flaking of the skin cracking, itching, redness, or stinging of the skin dark urine difficulty with breathing difficulty with walking dizziness or lightheadedness fainting fever headache inability to feel hands and feet irritability irritation of the nose itching, scaling, severe redness, soreness, or swelling of the skin pale skin rapid heart rate red, sore eyes shortness of breath sore throat unusual bleeding or bruising unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness unusual tiredness or weakness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Chiggerex side effects

2497. This particular disease, Pain, interacts with the drug Cool 'n Heat Patch and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

Drug Side Effects Summary

- hives
 - itching
 - red, swollen, blistered, or peeling skin with or without fever
 - wheezing
 - tightness in the chest or throat
 - trouble breathing, swallowing, or talking
 - unusual hoarseness
 - or swelling of the mouth, face, lips, tongue, or throat. Very bad skin irritation. Cool 'n Heat Patch
- side effects

2498. This particular disease, Pain, interacts with the drug Cope and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Cope may cause serious side effects. Stop using Cope and call your doctor at once if you have: ringing in your ears, confusion, hallucinations, rapid breathing, seizure (convulsions)
- severe nausea , vomiting , or stomach pain
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- fever lasting longer than 3 days
- or swelling, or pain lasting longer than 10 days. Common side effects of Cope may include: upset stomach , heartburn
- drowsiness
- or mild headache .

2499. This particular disease, Pain, interacts with the drug Dendracin Neurodendraxcin and has the

Drug Side Effects Summary

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Irritation where Dendracin Neurodendraxcin (methyl salicylate, menthol, and capsaicin lotion) is used. Very bad burning or burning that does not go away. Dendracin Neurodendraxcin side effects

2500. This particular disease, Pain, interacts with the drug Denti-Care Denti-Freeze and has the following side effects:

- No side effects listed

2501. This particular disease, Pain, interacts with the drug DermacinRx Lexitral and has the following side effects:

- No side effects listed

2502. This particular disease, Pain, interacts with the drug DermacinRx Lexitral PharmaPak and has the following side effects:

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: a skin rash, no matter how mild

Drug Side Effects Summary

- shortness of breath (even with mild exertion), swelling, rapid weight gain
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Common side effects of DermacinRx Lexitral PharmaPak may include: skin rash, itching, redness, blistering, tingling, or other irritation where the medicine was applied
- or dryness or hardening of treated skin.

2503. This particular disease, Pain, interacts with the drug Dermoplast and has the following side effects

- Dermoplast is not for use in the mouth. Other brands of benzocaine can be used in the mouth may cause a condition in which the oxygen in your body tissues can become dangerously low. This is a potentially fatal condition called methemoglobinemia (met-HEEM-oh glo-bin-EE-mee-a). This condition may occur after only one use of benzocaine or after several uses. Signs and symptoms may occur within minutes or up to 2 hours after using benzocaine in the mouth or throat. GET EMERGENCY MEDICAL HELP IF YOU HAVE: a headache, tiredness, confusion
- fast heartbeats
- feeling light-headed or short of breath
- and pale, blue, or gray appearance of your skin, lips, or fingernails. hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Dermoplast and call your doctor at once if you have: severe burning, stinging, or sensitivity where the medicine is applied
- swelling, warmth, or redness

Drug Side Effects Summary

- or oozing, blistering, or any signs of infection. Common Dermoplast side effects may include: mild stinging, burning, or itching where the medicine is applied
- skin tenderness or redness
- or dry white flakes where the medicine was applied.

2504. This particular disease, Pain, interacts with the drug Diclofex DC and has the following side effects

- No side effects listed

2505. This particular disease, Pain, interacts with the drug Dicloheal-60 and has the following side effects

- No side effects listed

2506. This particular disease, Pain, interacts with the drug Diclosaicin and has the following side effects

- No side effects listed

2507. This particular disease, Pain, interacts with the drug Diclotral Pak and has the following side effects

- chest pain spreading to your jaw or shoulder, sudden numbness or weakness on one side of the body, slurred speech, feeling short of breath. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: a skin rash, no matter how mild
- shortness of breath (even with mild exertion), swelling, rapid weight gain
- signs of stomach bleeding--bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of

Drug Side Effects Summary

breath, cold hands and feet. Common side effects of Diclofenac Pak may include: skin rash, itching, redness, blistering, tingling, or other irritation where the medicine was applied

- or dryness or hardening of treated skin.

2508. This particular disease, Pain, interacts with the drug Doan's Pills and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

- itching

- red, swollen, blistered, or peeling skin with or without fever

- wheezing

- tightness in the chest or throat

- trouble breathing, swallowing, or talking

- unusual hoarseness

- or swelling of the mouth, face, lips, tongue, or throat. Signs of bleeding like throwing up or coughing up blood

- vomit that looks like coffee grounds

- blood in the urine

- black, red, or tarry stools

- bleeding from the gums

- abnormal vaginal bleeding

- bruises without a cause or that get bigger

- or bleeding you cannot stop. Very bad belly pain. Feeling confused. Ringing in ears. Hearing loss .

Drug Side Effects Summary

Doan's Pills side effects

2509. This particular disease, Pain, interacts with the drug Doans Pills Extra Strength and has the following side effects:

- sneezing, runny or stuffy nose
- wheezing or trouble breathing
- hives
- swelling of your face, lips, tongue, or throat. Doans Pills Extra Strength may cause serious side effects. Stop using Doans Pills Extra Strength and call your doctor at once if you have: new symptoms such as redness or swelling
- ringing in your ears, hearing loss
- a light-headed feeling, like you might pass out
- new or worsening stomach pain
- or bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Less serious side effects may be more likely, and you may have none at all.

2510. This particular disease, Pain, interacts with the drug Dsuvia and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: Less common Blurred vision confusion dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position sweating unusual tiredness or weakness Rare Bloody urine blue lips, fingernails, or skin burning feeling in the chest or stomach burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain or discomfort decreased frequency or amount of urine decreased urine output difficult or troubled breathing difficulty in passing urine (dribbling) extremely shallow or slow breathing fainting fast heartbeat increased blood pressure increased thirst

Drug Side Effects Summary

indigestion irregular, fast or slow, or shallow breathing lightheadedness, dizziness or fainting loss of appetite lower back or side pain nausea painful urination seeing, hearing, or feeling things that are not there severe constipation severe vomiting slow or irregular heartbeat stomach pain or upset swelling of the face, fingers, or lower legs tenderness in the stomach area troubled breathing unusual drowsiness, dullness, or feeling of sluggishness vomiting weakness weight gain Incidence not known Agitation cough darkening of the skin diarrhea difficulty swallowing fever hives, itching, skin rash mental depression overactive reflexes poor coordination puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue restlessness shivering talking or acting with excitement you cannot control tightness in the chest trembling or shaking twitching Symptoms of overdose Blue lips and fingernails blurred vision chest pain choking cold and clammy skin constricted pupils coughing that sometimes produces a pink frothy sputum decreased awareness or responsiveness difficult or troubled breathing irregular, fast or slow, or shallow breathing loss of consciousness pale or blue lips, fingernails, or skin severe sleepiness slow heartbeat swelling in the legs and ankles Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Headache Rare Anxiety belching bloated confusion about identity, place, and time difficulty having a bowel movement (stool) drowsiness dry mouth excess air or gas in the stomach or bowels false or unusual sense of well-being feeling of warmth forgetfulness full feeling heartburn indigestion muscle spasm passing gas redness of the face, neck, arms and occasionally, upper chest sleepiness or unusual drowsiness trouble sleeping Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Dsuvia side effects

Drug Side Effects Summary

2511. This particular disease, Pain, interacts with the drug Duraxin and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

- itching

- red, swollen, blistered, or peeling skin with or without fever

- wheezing

- tightness in the chest or throat

- trouble breathing, swallowing, or talking

- unusual hoarseness

- or swelling of the mouth, face, lips, tongue, or throat. Signs of bleeding like throwing up or coughing up blood

- vomit that looks like coffee grounds

- blood in the urine

- black, red, or tarry stools

- bleeding from the gums

- abnormal vaginal bleeding

- bruises without a cause or that get bigger

- or bleeding you cannot stop. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. A fast heartbeat . A heartbeat that does not feel normal. Very bad dizziness or passing out. Ringing in ears. Not able to pass urine or change in how much urine is passed. Change in balance. Not able to focus. Seizures . Blurred eyesight. Shakiness. A very bad skin reaction (Stevens-Johnson syndrome /

Drug Side Effects Summary

toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)

- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Duraxin side effects

2512. This particular disease, Pain, interacts with the drug Dvorah and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Dvorah and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Dvorah may cause serious side effects. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep
- chest pain, fast or pounding heartbeats, feeling light-headed, fainting
- confusion, extreme drowsiness, unusual thoughts or behavior
- painful or difficult urination
- severe constipation
- liver problems-- nausea , upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or low cortisol levels-- nausea, vomiting , loss of appetite, dizziness, worsening tiredness or weakness. Seek medical attention right away if you have symptoms of serotonin syndrome, such as:

Drug Side Effects Summary

agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea . Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent. Common side effects include: dizziness, drowsiness, tiredness, headache

- nausea, vomiting, stomach pain, constipation
- sweating, itching
- or skin rash .

2513. This particular disease, Pain, interacts with the drug dyclonine and has the following side effects

- No side effects listed

2514. This particular disease, Pain, interacts with the drug Ed-Flex and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness

Drug Side Effects Summary

- or swelling of the mouth, face, lips, tongue, or throat. Signs of bleeding like throwing up or coughing up blood
- vomit that looks like coffee grounds
- blood in the urine
- black, red, or tarry stools
- bleeding from the gums
- abnormal vaginal bleeding
- bruises without a cause or that get bigger
- or bleeding you cannot stop. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. A fast heartbeat . A heartbeat that does not feel normal. Very bad dizziness or passing out. Ringing in ears. Not able to pass urine or change in how much urine is passed. Change in balance. Not able to focus. Seizures . Blurred eyesight. Shakiness. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Ed-Flex side effects

2515. This particular disease, Pain, interacts with the drug Endodan and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Agitation bleeding gums bloating blood in the urine or stools bloody, black, or tarry stools blue lips, fingernails, or skin blurred vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings

Drug Side Effects Summary

change in consciousness or confusion chest pain, tightness, or discomfort chills clay-colored stools confusion cough coughing or vomiting blood dark urine darkening of the skin decrease in urine volume or frequency decreased appetite depression difficult, fast, noisy breathing difficulty in passing urine (dribbling) difficulty having a bowel movement (stool) difficulty with swallowing dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position drowsiness dry mouth fainting fast, slow, irregular, pounding, or racing heartbeat or pulse feeling of hostility or irritability feeling of warmth feeling that something terrible will happen fever headache, sudden, severe heartburn hives, itching, or skin rash increased menstrual flow or vaginal bleeding increased sweating indigestion irregular, fast, slow, or shallow breathing large, flat, blue or purplish patches on the skin large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or genitals loss of consciousness muscle cramps, pain, stiffness, weakness, or tremors nausea nosebleeds numbness or tingling in the hands, feet, or lips overactive reflexes painful or difficult urination pains in the stomach, side, or abdomen, possibly radiating to the back pale skin pinpoint red or purple spots on the skin poor coordination prolonged bleeding from cuts puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red or black, tarry stools or dark urine restlessness seizures shivering sleepiness sunken eyes sweating swelling of the face, ankles, hands, feet, or lower legs stomach pain, cramping, or tenderness talking or acting with excitement you cannot control thirst trembling or shaking twitching unusual bleeding or bruising unusual tiredness or weakness vomiting vomiting of material that looks like coffee grounds, severe and continuing weak or feeble pulse weakness or heaviness of the legs weight gain wrinkled skin yellow eyes or skin

Symptoms of overdose Blue lips and fingernails blurred vision change in consciousness chest pain cold, clammy skin confusion continuing ringing or buzzing or other unexplained noise in the ears coughing that sometimes produces a pink frothy sputum decreased awareness or responsiveness diarrhea dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position drowsiness enlarged pupils extremely high fever or body temperature fast,

Drug Side Effects Summary

weak heartbeat hearing loss increase in heart rate increased sweating loss of consciousness pale skin restlessness severe sleepiness swelling in legs and ankles unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Relaxed and calm feeling sleepiness Incidence not known Belching bloated, full feeling blurred or loss of vision change in color perception cold sweats constricted, pinpoint, or small pupils (black part of the eye) cool, pale skin double vision excess air or gas in the stomach false or unusual sense of well-being flushed, dry skin fruit-like breath odor halos around lights increased hunger or thirst increased urination lack or loss of strength night blindness nightmares overbright appearance of lights red eyes redness of the skin seeing, hearing, or feeling things that are not there shakiness slurred speech trouble sleeping tunnel vision unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness weight loss Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Endodan side effects

2516. This particular disease, Pain, interacts with the drug Exoten-C Pain Relief and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching

Drug Side Effects Summary

- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Irritation where Exoten-C Pain Relief (methyl salicylate, menthol, and capsaicin lotion) is used. Very bad burning or burning that does not go away. Exoten-C Pain Relief side effects

2517. This particular disease, Pain, interacts with the drug Exparel and has the following side effects

- hives , red rash, itching
- sneezing, difficulty breathing
- severe dizziness , vomiting
- swelling of your face, lips, tongue, or throat. You will be watched closely after receiving Exparel, to make sure you do not have a reaction to the medicine. Tell your caregivers at once if you have any of these signs of a serious side effect: ringing in your ears
- drowsiness, feeling restless or anxious
- feeling like you might pass out
- speech or vision problems, a metallic taste in your mouth
- numbness or tingling around your mouth
- fast or slow heart rate , feeling short of breath, feeling unusually hot or cold
- tremors, twitching, mood changes
- ongoing numbness, weakness, or loss of movement where the medicine was injected
- or joint pain or stiffness, or weakness in any part of your body for months after your surgery. You may still feel numb or be unable to move the numbed area for up to 5 days after you are treated with

Drug Side Effects Summary

bupivacaine. Common Exparel side effects include: nausea , vomiting

- constipation
- or fever .

2518. This particular disease, Pain, interacts with the drug Flexall and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Flexall may cause serious side effects. Stop using Flexall and call your doctor at once if you have: severe burning, stinging, redness, or irritation after using this medicine
- or pain, swelling, or blistering where the medicine was applied. Less serious side effects may be more likely, and you may have none at all.

2519. This particular disease, Pain, interacts with the drug Gebauer's Spray and Stretch and has the

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness

Drug Side Effects Summary

- or swelling of the mouth, face, lips, tongue, or throat. A burning, numbness , or tingling feeling that is not normal. Irritation where Gebauer's Spray and Stretch (pentafluoropropane and tetrafluoroethane) is used. Gebauer's Spray and Stretch side effects

2520. This particular disease, Pain, interacts with the drug Gold Bond Medicated Maximum Strength

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Gold Bond Medicated Maximum Strength Foot may cause serious side effects. Stop using Gold Bond Medicated Maximum Strength Foot and call your doctor at once if you have: severe burning, stinging, redness, or irritation after using this medicine
- or pain, swelling, or blistering where the medicine was applied. Less serious side effects may be more likely, and you may have none at all.

2521. This particular disease, Pain, interacts with the drug Icy Hot Advanced Relief Pain Relief Patc

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness

Drug Side Effects Summary

- or swelling of the mouth, face, lips, tongue, or throat. Very bad skin irritation. Icy Hot Advanced Relief Pain Relief Patch side effects

2522. This particular disease, Pain, interacts with the drug Icy Hot Extra Strength and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Icy Hot Extra Strength may cause serious side effects. Stop using Icy Hot Extra Strength and call your doctor at once if you have: severe burning, stinging, redness, or irritation after using this medicine
- or pain, swelling, or blistering where the medicine was applied. Less serious side effects may be more likely, and you may have none at all.

2523. This particular disease, Pain, interacts with the drug Icy Hot Medicated Spray and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad skin irritation. Icy Hot Medicated

Drug Side Effects Summary

Spray side effects

2524. This particular disease, Pain, interacts with the drug Icy Hot Vanishing Scent Pain Relieving Ointment and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad skin irritation. Icy Hot Vanishing Scent Pain Relieving Gel side effects

2525. This particular disease, Pain, interacts with the drug Inavix and has the following side effects:

- No side effects listed

2526. This particular disease, Pain, interacts with the drug Kank-a and has the following side effects:

- KANK-A used in the mouth may cause a condition in which the oxygen in your body tissues can become dangerously low. This is a potentially fatal condition called methemoglobinemia (met-HEEM-oh glo-bin-EE-mee-a). This condition may occur after only one use of benzocaine or after several uses. Signs and symptoms may occur within minutes or up to 2 hours after using

Drug Side Effects Summary

KANK-A in the mouth or throat. GET EMERGENCY MEDICAL HELP IF YOU HAVE: a headache, tiredness, confusion

- fast heartbeats
- feeling light-headed or short of breath
- and pale, blue, or gray appearance of your skin, lips, or fingernails. hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: severe burning, stinging, or sensitivity where the medicine is applied
- swelling, warmth, or redness
- or oozing, blistering, or any signs of infection. Common KANK-A side effects may include: mild stinging, burning, or itching where the medicine is applied
- skin tenderness or redness
- or dry white flakes where the medicine was applied.

2527. This particular disease, Pain, interacts with the drug Lanacane and has the following side effects

- Lanacane should not be used in the mouth. Other products containing benzocaine are used in the mouth and may cause a condition in which the oxygen in your body tissues can become dangerously low. This is a potentially fatal condition called methemoglobinemia (met-HEEM-oh glo-bin-EE-mee-a). This condition may occur after only one use of benzocaine or after several uses. Signs and symptoms may occur within minutes or up to 2 hours after using benzocaine in the mouth or throat. GET EMERGENCY MEDICAL HELP IF YOU HAVE: a headache, tiredness, confusion
- fast heartbeats
- feeling light-headed or short of breath
- and pale, blue, or gray appearance of your skin, lips, or fingernails. hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Stop using Lanacane and call your doctor at once if you have: severe burning, stinging, or sensitivity where the medicine is applied
- swelling, warmth, or redness
- or oozing, blistering, or any signs of infection. Common Lanacane side effects may include: mild stinging, burning, or itching where the medicine is applied
- skin tenderness or redness
- or dry white flakes where the medicine was applied.

2528. This particular disease, Pain, interacts with the drug Laryngesic and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of methemoglobinemia like a blue or gray color of the lips, nails, or skin
- a heartbeat that does not feel normal
- seizures
- very bad dizziness or passing out

Drug Side Effects Summary

- very bad headache
 - feeling very sleepy
 - feeling tired or weak
 - or shortness of breath. This effect is rare but may be deadly if it happens. Very bad mouth pain.
- When treating a very sore throat , if signs last more than 2 days, or happen with fever, headache, rash, upset stomach , or throwing up. Laryngesic side effects

2529. This particular disease, Pain, interacts with the drug LidoPro and has the following side effects

- hives
 - difficulty breathing
 - swelling of your face, lips, tongue, or throat. Wash the skin and get medical attention right away if you have severe burning, pain, swelling, or blistering of the skin where you applied this medicine.
- This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: pale skin, blue-colored lips
- headache , confusion
 - or rapid heartbeats. Common side effects may include a mild burning sensation that can last for several hours or days, especially after your first use of this medicine.

2530. This particular disease, Pain, interacts with the drug magnesium salicylate and has the following

- sneezing, runny or stuffy nose
 - wheezing or trouble breathing
 - hives
 - swelling of your face, lips, tongue, or throat. magnesium salicylate may cause serious side effects.
- Stop using magnesium salicylate and call your doctor at once if you have: new symptoms such as redness or swelling

Drug Side Effects Summary

- ringing in your ears, hearing loss
- a light-headed feeling, like you might pass out
- new or worsening stomach pain
- or bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Less serious side effects may be more likely, and you may have none at all.

2531. This particular disease, Pain, interacts with the drug Max-Freeze and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Max-Freeze may cause serious side effects. Stop using Max-Freeze and call your doctor at once if you have: severe burning, stinging, redness, or irritation after using this medicine
- or pain, swelling, or blistering where the medicine was applied. Less serious side effects may be more likely, and you may have none at all.

2532. This particular disease, Pain, interacts with the drug Medi-Derm Rx and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat

Drug Side Effects Summary

- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Irritation where Medi-Derm Rx (methyl salicylate, menthol, and capsaicin cream and ointment) is used. Very bad burning or burning that does not go away. Medi-Derm Rx side effects

2533. This particular disease, Pain, interacts with the drug Medicone and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bluish color of the fingernails, lips, skin, palms, or nail beds Incidence not known Blistering, burning, crusting, dryness, or flaking of the skin cracking, itching, redness, or stinging of the skin dark urine difficulty with breathing difficulty with walking dizziness or lightheadedness fainting fever headache inability to feel hands and feet irritability irritation of the nose itching, scaling, severe redness, soreness, or swelling of the skin pale skin rapid heart rate red, sore eyes shortness of breath sore throat unusual bleeding or bruising unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness unusual tiredness or weakness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Medicone side effects

2534. This particular disease, Pain, interacts with the drug Medrox and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

Drug Side Effects Summary

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Irritation where Medrox (methyl salicylate, menthol, and capsaicin cream and ointment) is used. Very bad burning or burning that does not go away. Medrox side effects

2535. This particular disease, Pain, interacts with the drug Medrox-Rx and has the following side effects

- No side effects listed

2536. This particular disease, Pain, interacts with the drug menthol and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. menthol topical may cause serious side effects. Stop using menthol topical and call your doctor at once if you have: severe burning, stinging, redness, or irritation after using this medicine
- or pain, swelling, or blistering where the medicine was applied. Less serious side effects may be more likely, and you may have none at all.

2537. This particular disease, Pain, interacts with the drug Meperitab and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of

Drug Side Effects Summary

these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Agitation blurred vision chest pain or discomfort cold, clammy skin confusion cough darkening of the skin decrease in the frequency of urination decrease in urine volume diarrhea difficult or trouble breathing difficulty in passing urine (dribbling) difficulty with swallowing dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position fast, slow, pounding, or irregular heartbeat feeling of warmth fever hives, itching, or skin rash irregular, fast, slow, or shallow breathing lightheadedness, dizziness, or fainting loss of appetite mental depression muscle twitching or jerking nausea overactive reflexes painful urination pale or blue lips, fingernails, or skin poor coordination puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue redness of the face, neck, arms, and occasionally, upper chest restlessness rhythmic movement of the muscles seizures shakiness in the legs, arms, hands, or feet shivering sweating talking or acting with excitement you cannot control tightness in the chest trembling or shaking of the hands or feet twitching unusual tiredness or weakness upper abdominal or stomach pain vomiting Symptoms of overdose Bluish color of the fingernails, lips, skin, palms, or nail beds change in consciousness decreased awareness or responsiveness loss of consciousness severe sleepiness sleepiness or unusual drowsiness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Drowsiness relaxed and calm feeling Incidence not known Blurred or loss of vision confusion about identity, place, and time deep or fast breathing with dizziness difficulty having a bowel movement disturbed color perception double vision dry mouth false or unusual sense of well-being halos around lights headache irritability nervousness night blindness numbness of the feet, hands, and around the mouth overbright

Drug Side Effects Summary

appearance of lights redness of the skin seeing, hearing, or feeling things that are not there trouble sleeping tunnel vision Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Meperitab side effects

2538. This particular disease, Pain, interacts with the drug Mineral Ice and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Mineral Ice may cause serious side effects. Stop using Mineral Ice and call your doctor at once if you have: severe burning, stinging, redness, or irritation after using this medicine
- or pain, swelling, or blistering where the medicine was applied. Less serious side effects may be more likely, and you may have none at all.

2539. This particular disease, Pain, interacts with the drug MST and has the following side effects:

- sneezing, runny or stuffy nose
- wheezing or trouble breathing
- hives
- swelling of your face, lips, tongue, or throat. MST may cause serious side effects. Stop using MST and call your doctor at once if you have: new symptoms such as redness or swelling
- ringing in your ears, hearing loss
- a light-headed feeling, like you might pass out
- new or worsening stomach pain
- or bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Less serious side effects may be more likely, and you may have none at all.

Drug Side Effects Summary

2540. This particular disease, Pain, interacts with the drug Nalocet and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Nalocet and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Nalocet may cause serious side effects. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops
- a light-headed feeling, like you might pass out
- weakness, tiredness, fever, unusual bruising or bleeding
- confusion, unusual thoughts or behavior
- problems with urination
- liver problems--nausea, upper stomach pain, tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- low cortisol levels-- nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness
- or high levels of serotonin in the body-- agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, diarrhea . Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders. Common side effects include: dizziness, drowsiness, feeling tired
- feelings of extreme happiness or sadness
- nausea, vomiting, stomach pain

Drug Side Effects Summary

- constipation
- or headache .

2541. This particular disease, Pain, interacts with the drug naloxone / pentazocine and has the following side effects:

- noisy breathing, sighing, shallow breathing, breathing that stops
 - a slow heart rate or weak pulse
 - a light-headed feeling, like you might pass out
 - severe constipation
 - confusion, hallucinations, unusual thoughts or behavior
 - a seizure
 - low cortisol levels-- nausea , vomiting , loss of appetite, dizziness, worsening tiredness or weakness
 - or serotonin syndrome -- agitation , hallucinations, fever, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, diarrhea . Serious breathing problems may be more likely in older adults and those who are debilitated or have wasting syndrome or chronic breathing disorders.
- Common side effects of naloxone and pentazocine may include: dizziness, drowsiness
- constipation
 - stomach pain, nausea, vomiting
 - or feeling tired.

2542. This particular disease, Pain, interacts with the drug New Terocin and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

Drug Side Effects Summary

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Irritation where New Terocin (methyl salicylate, menthol, and capsaicin lotion) is used. Very bad burning or burning that does not go away. New Terocin side effects

2543. This particular disease, Pain, interacts with the drug NuDiclo SoluPak and has the following side effects:

- No side effects listed

2544. This particular disease, Pain, interacts with the drug Nuprin Backache and has the following side effects:

- sneezing, runny or stuffy nose
- wheezing or trouble breathing
- hives
- swelling of your face, lips, tongue, or throat. Nuprin Backache may cause serious side effects. Stop using Nuprin Backache and call your doctor at once if you have: new symptoms such as redness or swelling
- ringing in your ears, hearing loss
- a light-headed feeling, like you might pass out
- new or worsening stomach pain
- or bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds. Less serious

Drug Side Effects Summary

side effects may be more likely, and you may have none at all.

2545. This particular disease, Pain, interacts with the drug oliceridine and has the following side effects:

- hives
 - difficult breathing
 - swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Oliceridine may cause serious side effects. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep
 - a slow heart rate or weak pulse
 - a light-headed feeling, like you might pass out
 - severe constipation
 - low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet
 - low cortisol levels-- nausea , vomiting , loss of appetite, dizziness, worsening tiredness or weakness
 - or low oxygen in your blood--confusion, feeling restless or short of breath, blue-colored lips or skin.
- Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea . Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders.
- Common side effects of oliceridine may include: nausea, vomiting, constipation
- headache , dizziness
 - itching

Drug Side Effects Summary

- or low oxygen in your blood.

2546. This particular disease, Pain, interacts with the drug Olinvyk and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Olinvyk may cause serious side effects. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep
- a slow heart rate or weak pulse
- a light-headed feeling, like you might pass out
- severe constipation
- low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet
- low cortisol levels-- nausea , vomiting , loss of appetite, dizziness, worsening tiredness or weakness
- or low oxygen in your blood--confusion, feeling restless or short of breath, blue-colored lips or skin. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea . Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders. Common side effects of Olinvyk may include: nausea, vomiting, constipation
- headache , dizziness
- itching

Drug Side Effects Summary

- or low oxygen in your blood.

2547. This particular disease, Pain, interacts with the drug P-A-C and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of bleeding like throwing up or coughing up blood
- vomit that looks like coffee grounds
- blood in the urine
- black, red, or tarry stools
- bleeding from the gums
- abnormal vaginal bleeding
- bruises without a cause or that get bigger
- or bleeding you cannot stop. Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow

Drug Side Effects Summary

skin or eyes. Signs of high potassium levels like a heartbeat that does not feel normal

- feeling confused
- feeling weak, lightheaded, or dizzy
- feeling like passing out
- numbness or tingling
- or shortness of breath. Signs of too much acid in the blood (acidosis) like confusion
- fast breathing
- fast heartbeat
- a heartbeat that does not feel normal
- very bad stomach pain, upset stomach, or throwing up
- feeling very sleepy
- shortness of breath
- or feeling very tired or weak. Weakness on 1 side of the body, trouble speaking or thinking, change in balance, drooping on one side of the face, or blurred eyesight. Very bad dizziness or passing out. Very bad headache . Ringing in the ears, hearing loss , or any other changes in hearing. Feeling agitated. Seizures . A fast heartbeat. P-A-C side effects

2548. This particular disease, Pain, interacts with the drug P-A-C Analgesic and has the following s

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. P-A-C Analgesic may cause serious side effects. Stop using P-A-C Analgesic and call your doctor at once if you have: ringing in your ears, confusion, hallucinations, rapid breathing, seizure (convulsions)
- severe nausea , vomiting , or stomach pain
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds

Drug Side Effects Summary

- fever lasting longer than 3 days
- or swelling, or pain lasting longer than 10 days. Common side effects of P-A-C Analgesic may include: upset stomach , heartburn
- drowsiness
- or mild headache .

2549. This particular disease, Pain, interacts with the drug pentafluoropropane / tetrafluoroethane a

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
 - hives
 - itching
 - red, swollen, blistered, or peeling skin with or without fever
 - wheezing
 - tightness in the chest or throat
 - trouble breathing, swallowing, or talking
 - unusual hoarseness
 - or swelling of the mouth, face, lips, tongue, or throat. A burning, numbness , or tingling feeling that is not normal. Irritation where pentafluoropropane and tetrafluoroethane is used.
- Pentafluoropropane / tetrafluoroethane topical side effects

2550. This particular disease, Pain, interacts with the drug Percodan and has the following side effe

- hives
- difficulty breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Oxycodone can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Call your doctor at once if you have: fast or slow heartbeats, weak or shallow breathing, breathing that stops during sleep
- severe stomach pain or constipation, vomiting
- confusion, unusual thoughts or behavior, feeling like you might pass out
- decreased hearing or ringing in the ears
- a seizure (convulsions)
- signs of stomach bleeding - bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- or low cortisol levels - nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Percodan is more likely to cause breathing problems in older adults and people who are severely ill, malnourished, or otherwise debilitated. Common Percodan side effects may include: headache, dizziness, drowsiness
- constipation, heartburn, upset stomach, bloating, gas, diarrhea
- or dry mouth.

2551. This particular disease, Pain, interacts with the drug Pramox and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

Drug Side Effects Summary

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Irritation where Pramox (pramoxine gel) is used. Swelling, warmth, or redness of the skin. Pramox side effects

2552. This particular disease, Pain, interacts with the drug pramoxine and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: any new redness or swelling where the medicine was applied
- or severe pain, burning, or stinging where the medicine was applied. Less serious side effects are more likely, and you may have none at all.

2553. This particular disease, Pain, interacts with the drug Prax and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: For all uses of Prax (pramoxine lotion): Signs of an allergic reaction, like rash
- hives

Drug Side Effects Summary

- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Irritation where Prax (pramoxine lotion) is used. Swelling, warmth, or redness of the skin. Rectal: Bleeding from rectum or rectal pain. Prax side effects

2554. This particular disease, Pain, interacts with the drug Precaine B and has the following side effects:

- No side effects listed

2555. This particular disease, Pain, interacts with the drug Prialt and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Ziconotide can affect your central nervous system and you may feel less alert. A person caring for you should seek emergency medical attention if you are hard to wake up. Call your doctor at once if you have: new or worsening muscle pain, soreness, or weakness, and/or dark urine
- a light-headed feeling (like you might pass out)
- strange sensations in your mouth
- skin sores, itching, blisters, breakdown of the outer layer of skin
- confusion (especially in older adults)
- problems with memory or thought

Drug Side Effects Summary

- unusual changes in mood or behavior - anger, aggression, paranoia, hallucinations, racing thoughts, risk-taking behavior
- symptoms of depression - feelings of low self-worth, loss of interest in things you once enjoyed, new sleep problems, thoughts about hurting yourself
- or symptoms of meningitis - fever, headache, neck stiffness, increased sensitivity to light, nausea, vomiting, confusion, or drowsiness. Common Prial side effects may include: dizziness
- confusion
- nausea
- or unusual or involuntary eye movements.

2556. This particular disease, Pain, interacts with the drug Primlev and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Primlev and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Primlev may cause serious side effects. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops
- a light-headed feeling, like you might pass out
- weakness, tiredness, fever, unusual bruising or bleeding
- confusion, unusual thoughts or behavior
- problems with urination

Drug Side Effects Summary

- liver problems--nausea, upper stomach pain, tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- low cortisol levels-- nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness
- or high levels of serotonin in the body-- agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, diarrhea . Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders. Common side effects include: dizziness, drowsiness, feeling tired
- feelings of extreme happiness or sadness
- nausea, vomiting, stomach pain
- constipation
- or headache .

2557. This particular disease, Pain, interacts with the drug Prolate and has the following side effects:

- No side effects listed

2558. This particular disease, Pain, interacts with the drug Qutenza and has the following side effects:

- hives
 - difficulty breathing
 - swelling of your face, lips, tongue, or throat. Wash the skin and get medical attention right away if you have severe burning, pain, swelling, or blistering of the skin where you applied this medication. Stop using Qutenza and call your doctor at once if you have: skin redness where the medicine was applied
 - or trouble breathing or swallowing (after accidental inhalation of capsaicin odor or dried residue).
- Less serious side effects may include a mild burning sensation that can last for several hours or

Drug Side Effects Summary

days, especially after your first use of Qutenza.

2559. This particular disease, Pain, interacts with the drug Rematex and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Irritation where Rematex (methyl salicylate, menthol, and capsaicin cream and ointment) is used. Very bad burning or burning that does not go away. Rematex side effects

2560. This particular disease, Pain, interacts with the drug Salonpas-Hot and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching

Drug Side Effects Summary

- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Salonpas-Hot side effects

2561. This particular disease, Pain, interacts with the drug Sarna Sensitive and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: For all uses of Sarna Sensitive (pramoxine lotion): Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Irritation where Sarna Sensitive (pramoxine lotion) is used. Swelling, warmth, or redness of the skin. Rectal: Bleeding from rectum or rectal pain.

Sarna Sensitive side effects

2562. This particular disease, Pain, interacts with the drug Seglantis and has the following side effects:

- The possible side effects of Seglantis: constipation, nausea, sleepiness, vomiting, tiredness,

Drug Side Effects Summary

headache, dizziness, abdominal pain. Call your healthcare provider if you have any of these symptoms and they are severe. NSAIDs can cause serious side effects, including: new or worse high blood pressure, heart failure, liver problems including liver failure, kidney problems including kidney failure, low red blood cells (anemia), life-threatening skin reactions, lifethreatening allergic reactions. Other side effects of NSAIDs include: stomach pain, constipation, diarrhea, gas, heartburn, nausea, vomiting, and dizziness. trouble breathing, shortness of breath, fast heartbeat, chest pain, swelling of your face, tongue, or throat, extreme drowsiness, light-headedness when changing positions, feeling faint, agitation, high body temperature, trouble walking, stiff muscles, or mental changes such as confusion. Stop taking Seglentis and call your healthcare provider right away if you get any of the following symptoms: nausea, more tired or weaker than usual, diarrhea, itching, your skin or eyes look yellow, indigestion or stomach pain, flulike symptoms, vomit blood, there is blood in your bowel movement or it is black and sticky like tar, unusual weight gain, skin rash or blisters with fever, swelling of the arms, legs, hands and feet. Seglentis may cause fertility problems in males and females, which may affect the ability to have children. Talk to your healthcare provider if you have concerns about fertility. These are not all the possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. For more information go to dailymed.nlm.nih.gov. Other information about NSAIDs: Aspirin is an NSAID but it does not increase the chance of a heart attack. Aspirin can cause bleeding in the brain, stomach, and intestines. Aspirin can also cause ulcers in the stomach and intestines. Some NSAIDs are sold in lower doses without a prescription (over the counter). Talk to your healthcare provider before using over-the-counter NSAIDs for more than 10 days. Seglentis side effects

2563. This particular disease, Pain, interacts with the drug Solarcaine First Aid Medicated Spray and

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of

Drug Side Effects Summary

these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Bluish color of the fingernails, lips, skin, palms, or nail beds Incidence not known Blistering, burning, crusting, dryness, or flaking of the skin cracking, itching, redness, or stinging of the skin dark urine difficulty with breathing difficulty with walking dizziness or lightheadedness fainting fever headache inability to feel hands and feet irritability irritation of the nose itching, scaling, severe redness, soreness, or swelling of the skin pale skin rapid heart rate red, sore eyes shortness of breath sore throat unusual bleeding or bruising unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness unusual tiredness or weakness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Solarcaine First Aid Medicated Spray side effects

2564. This particular disease, Pain, interacts with the drug Soltice Quick-Rub and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness

Drug Side Effects Summary

- or swelling of the mouth, face, lips, tongue, or throat. Soltice Quick-Rub side effects

2565. This particular disease, Pain, interacts with the drug Sombra Cool Therapy and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

- itching

- red, swollen, blistered, or peeling skin with or without fever

- wheezing

- tightness in the chest or throat

- trouble breathing, swallowing, or talking

- unusual hoarseness

- or swelling of the mouth, face, lips, tongue, or throat. Very bad skin irritation. Sombra Cool Therapy side effects

2566. This particular disease, Pain, interacts with the drug Sombra Warm Therapy and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

- itching

- red, swollen, blistered, or peeling skin with or without fever

Drug Side Effects Summary

- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Sombra Warm Therapy side effects

2567. This particular disease, Pain, interacts with the drug Stopain and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stopain may cause serious side effects. Stop using Stopain and call your doctor at once if you have: severe burning, stinging, redness, or irritation after using this medicine
- or pain, swelling, or blistering where the medicine was applied. Less serious side effects may be more likely, and you may have none at all.

2568. This particular disease, Pain, interacts with the drug sufentanil and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: Incidence not known Agitation blurred vision confusion cough darkening of the skin diarrhea difficult or troubled breathing difficulty with swallowing dizziness dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position fainting fast heartbeat fever hives, itching, or skin rash irregular, fast or slow, or shallow breathing loss of appetite mental depression muscle stiffness nausea overactive reflexes pale or blue lips, fingernails, or skin poor coordination puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue restlessness shivering sweating talking or acting with excitement you

Drug Side Effects Summary

cannot control tightness in the chest trembling or shaking twitching unusual tiredness or weakness vomiting Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Sufentanil side effects

2569. This particular disease, Pain, interacts with the drug Sure Result DSS Premium Pak and has the following side effects:

- No side effects listed

2570. This particular disease, Pain, interacts with the drug Theragen HP and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
 - hives
 - itching
 - red, swollen, blistered, or peeling skin with or without fever
 - wheezing
 - tightness in the chest or throat
 - trouble breathing, swallowing, or talking
 - unusual hoarseness
 - or swelling of the mouth, face, lips, tongue, or throat. Theragen HP side effects

2571. This particular disease, Pain, interacts with the drug Therapeutic Ice and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if

Drug Side Effects Summary

you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Very bad skin irritation. Therapeutic Ice side effects

2572. This particular disease, Pain, interacts with the drug Tiger Balm and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tiger Balm may cause serious side effects. Stop using Tiger Balm and call your doctor at once if you have: severe burning, stinging, or irritation after using this medicine
- or breathing problems after using this medicine on the chest or throat. Less serious side effects may be more likely, and you may have none at all.

2573. This particular disease, Pain, interacts with the drug Tiger Balm Pain Relieving Patch and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs

Drug Side Effects Summary

of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Tiger Balm Pain Relieving Patch side effects

2574. This particular disease, Pain, interacts with the drug Tivorbex and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Acid or sour stomach belching diarrhea heartburn indigestion nausea stomach discomfort, upset, or pain vomiting Rare Abdominal or stomach cramping, burning, or tenderness back or leg pains bleeding gums blistering, peeling, or loosening of the skin bloody or black, tarry stools blue lips and fingernails blurred vision breast enlargement and tenderness burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings burning upper abdominal or stomach pain canker sores change in consciousness change in hearing chest pain, discomfort, or burning clay colored stools cloudy or bloody urine confusion continuing diarrhea cough or hoarseness coughing that sometimes produces a pink frothy sputum cracks in the skin dark urine decreased appetite decreased vision or any change in vision depression difficult or labored breathing difficulty with swallowing dilated neck veins dizziness, faintness, or lightheadedness when getting up from a lying or sitting position double vision dry mouth extreme fatigue false sense of well-being feeling of unreality feeling of warmth fever with

Drug Side Effects Summary

or without chills flushed, dry skin fruit-like breath odor general body swelling greatly decreased frequency of urination or amount of urine hair loss headache heavier menstrual periods increased hunger increased sweating increased thirst increased urination irregular breathing irritation and swelling of the eye jerky movements of the head, face, mouth, and neck joint pain large, flat, blue or purplish patches in the skin large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs loss of appetite loss of balance control loss of bladder control loss of consciousness loss of hearing loss of heat from the body lower back or side pain mask-like face mood swings muscle aches, pains, or weakness muscle spasm or jerking of all extremities nervousness noisy, rattling breathing nosebleeds numbness or tingling in the hands, feet, or lips pain in the ankles or knees pain or discomfort in the upper stomach or throat pain with swallowing painful or difficult urination painful, red lumps under the skin, mostly on the legs pale skin persistent bleeding or oozing from puncture sites, mouth, or nose personality changes pinpoint red or purple spots on the skin pounding in the ears puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red skin lesions, often with a purple center red, irritated eyes red, swollen skin redness of the face, neck, arms and occasionally, upper chest scaly skin seeing double seeing, hearing, or feeling things that are not there seizures sense of detachment from self or body severe constipation severe mental changes severe or continuing stomach pain shuffling walk skin rash, hives or welts, itching slow, fast, irregular, pounding, or racing heartbeat or pulse slowed movements slurred speech small red or purple spots on the skin sore throat sores, ulcers, or white spots on the lips or tongue or inside the mouth stiffness of the arms and legs sudden loss of consciousness swelling of the breasts or breast soreness in both females and males swelling of the face, fingers, feet, ankles, or lower legs swollen or painful glands tightness in the chest trembling and shaking of the fingers and hands troubled breathing at rest troubled breathing with exertion unexplained weight loss unpleasant breath odor unsteadiness or awkwardness unusual bleeding or bruising unusual tiredness or weakness vaginal bleeding vomiting of blood or material that looks like

Drug Side Effects Summary

coffee grounds weakness in the arms, hands, legs, or feet weight gain yellow eyes or skin
Symptoms of overdose Confusion about identity, place, and time severe headache unusual
drowsiness, dullness, or feeling of sluggishness Some side effects may occur that usually do not
need medical attention. These side effects may go away during treatment as your body adjusts to
the medicine. Also, your health care professional may be able to tell you about ways to prevent or
reduce some of these side effects. Check with your health care professional if any of the following
side effects continue or are bothersome or if you have any questions about them: More common
Mild headache Less common Continuing ringing or buzzing or other unexplained noise in the ears
difficulty having a bowel movement (stool) discouragement feeling sad or empty general feeling of
discomfort or illness hearing loss irritability loss of interest or pleasure sleepiness trouble with
concentrating Rare Anxiety bloated or full feeling changes in patterns and rhythms of speech excess
air or gas in the stomach or intestines feeling of constant movement of self or surroundings
involuntary muscle movements lightheadedness passing gas sensation of spinning tiredness trouble
sleeping trouble with speaking Other side effects not listed may also occur in some patients. If you
notice any other effects, check with your healthcare professional. Call your doctor for medical advice
about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Tivorbex side effects

2575. This particular disease, Pain, interacts with the drug Topical Anesthetic Dental Gel and has the following side effects:

- No side effects listed

2576. This particular disease, Pain, interacts with the drug Transderm-iQ and has the following side effects:

- hives

- difficulty breathing

- swelling of your face, lips, tongue, or throat. Wash the skin and get medical attention right away if you have severe burning, pain, swelling, or blistering of the skin where you applied this medicine.

Drug Side Effects Summary

This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: pale skin, blue-colored lips

- headache , confusion
- or rapid heartbeats. Common side effects may include a mild burning sensation that can last for several hours or days, especially after your first use of this medicine.

2577. This particular disease, Pain, interacts with the drug Trezix and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Trezix and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Trezix may cause serious side effects. Call your doctor at once if you have: noisy breathing, sighing, shallow breathing, breathing that stops during sleep
- chest pain, fast or pounding heartbeats, feeling light-headed, fainting
- confusion, extreme drowsiness, unusual thoughts or behavior
- painful or difficult urination
- severe constipation
- liver problems-- nausea , upper stomach pain, itching, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or low cortisol levels-- nausea, vomiting , loss of appetite, dizziness, worsening tiredness or weakness. Seek medical attention right away if you have symptoms of serotonin syndrome, such as:

Drug Side Effects Summary

agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea . Serious side effects may be more likely in older adults and those who are overweight , malnourished, or debilitated. Long-term use of opioid medication may affect fertility (ability to have children) in men or women. It is not known whether opioid effects on fertility are permanent. Common side effects include: dizziness, drowsiness, tiredness, headache

- nausea, vomiting, stomach pain, constipation
- sweating, itching
- or skin rash .

2578. This particular disease, Pain, interacts with the drug Tru-micin and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Irritation where Tru-micin (trolamine cream and lotion) is used. Redness. Tru-micin side effects

Drug Side Effects Summary

2579. This particular disease, Pain, interacts with the drug Ultracin T and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Irritation where Ultracin T (trolamine cream and lotion) is used. Redness. Ultracin T side effects

2580. This particular disease, Pain, interacts with the drug Xodol and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Opioid medicine can slow or stop your breathing, and death may occur. A person caring for you should give naloxone and/or seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. In rare cases, acetaminophen may cause a severe skin reaction that can be fatal. This could occur even if you have taken acetaminophen in the past and had no reaction. Stop taking Xodol and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling. Xodol may cause serious side effects. Call your doctor at once if you have: noisy breathing,

Drug Side Effects Summary

sighing, shallow breathing, breathing that stops

- a light-headed feeling, like you might pass out

- liver problems--nausea, upper stomach pain, tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)

- low cortisol levels-- nausea, vomiting, loss of appetite, dizziness, worsening tiredness or weakness

- o high levels of serotonin in the body-- agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, diarrhea . Serious breathing problems may be more likely in older adults and in those who are debilitated or have wasting syndrome or chronic breathing disorders. Common side effects include: dizziness, drowsiness, feeling tired

- nausea, vomiting, stomach pain

- constipation

- or headache .

2581. This particular disease, Pain, interacts with the drug Xolox and has the following side effects.

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Black, tarry stools chills dark urine dizziness fever headache itching, skin rash light-colored stools loss of appetite nausea stomach pain unpleasant breath odor unusual tiredness or weakness vomiting of blood yellow eyes or skin Rare Cough fever with or without chills general feeling of tiredness or weakness hoarseness lower back or side pain painful or difficult urination sore throat sores, ulcers, or white spots on the lips or in the mouth unusual bleeding or bruising Incidence not known Back, leg, or stomach pains bleeding gums bloating blood in the urine or stools blue lips and fingernails blurred vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain, discomfort,

Drug Side Effects Summary

or tightness cloudy urine clumsiness confusion constipation coughing that sometimes produces a pink frothy sputum darkening of the skin decreased awareness or responsiveness decreased frequency or amount of urine diarrhea difficult, fast, noisy breathing difficulty in passing urine (dribbling) difficulty with swallowing dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position drowsiness dry mouth extremely shallow or slow breathing fainting fast or deep breathing fast, slow, irregular, pounding, or racing heartbeat feeling of warmth general body swelling hives or welts increased sweating increased thirst indigestion large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or genitals lightheadedness mental depression muscle aches, tremors, or weakness nervousness nosebleeds pains in the stomach, side, or abdomen, possibly radiating to the back pale skin pinpoint red spots on the skin pounding in the ears puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rapid, deep or shallow breathing redness of the face, neck, arms, and occasionally, upper chest restlessness seizures severe constipation severe sleepiness severe vomiting skin blisters sleepiness stomach cramps sunken eyes sweating swelling of the face, fingers, lower legs, or ankles thirst tiredness trouble breathing vomiting weakness or heaviness of the legs weight gain

Symptoms of overdose Bluish lips or skin change in consciousness cold, clammy skin extreme sleepiness general feeling of discomfort or illness loss of consciousness low blood pressure or pulse slow breathing unconsciousness

Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them:

More common

Relaxed and calm feeling

Incidence not known

Agitation bad, unusual or unpleasant (after) taste belching change in taste cold sweats constricted, pinpoint, or small pupils (black part of the eye) continuing ringing or buzzing or other unexplained noise in the ears cool, pale skin depression difficulty with moving

Drug Side Effects Summary

disturbed color perception double vision excess air or gas in the stomach or bowels false or unusual sense of well-being flushed, dry skin fruit-like breath odor full feeling general feeling of discomfort or illness halos around lights hearing loss heartburn increased hunger increased urination indigestion joint pain lack or loss of strength muscle cramps, spasms, pain, or stiffness night blindness nightmares overbright appearance of lights seeing, hearing, or feeling things that are not there seizures shakiness slurred speech swollen joints trouble sleeping tunnel vision unexplained weight loss unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Xolox side effects

2582. This particular disease, Pain, interacts with the drug Ziclopro and has the following side effects

- No side effects listed

2583. This particular disease, Pain, interacts with the drug ziconotide and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Ziconotide can affect your central nervous system and you may feel less alert. A person caring for you should seek emergency medical attention if you are hard to wake up. Call your doctor at once if you have: new or worsening muscle pain, soreness, or weakness, and/or dark urine
- a light-headed feeling (like you might pass out)
- strange sensations in your mouth
- skin sores, itching, blisters, breakdown of the outer layer of skin
- confusion (especially in older adults)

Drug Side Effects Summary

- problems with memory or thought
- unusual changes in mood or behavior--anger, aggression, paranoia, hallucinations, racing thoughts, risk-taking behavior
- symptoms of depression --feelings of low self-worth, loss of interest in things you once enjoyed, new sleep problems, thoughts about hurting yourself
- or symptoms of meningitis --fever, headache, neck stiffness, increased sensitivity to light, nausea, vomiting, confusion, or drowsiness. Common ziconotide side effects may include: dizziness
- confusion
- nausea
- or unusual or involuntary eye movements.

2584. This particular disease, Pain, interacts with the drug Zyfrel and has the following side effects.

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or

Drug Side Effects Summary

eyes. Very bad dizziness or passing out. Clammy skin. Sweating a lot. Feeling confused. Trouble walking. Severe constipation or stomach pain. These may be signs of a severe bowel problem. Not able to pass urine or change in how much urine is passed. Hearing loss . Change in hearing. Chest pain or pressure. Fast or slow heartbeat . Fever, chills, or sore throat

- any unexplained bruising or bleeding

- or feeling very tired or weak. Mood changes. Trouble passing urine. Change in eyesight. Seizures.

A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)

- red or irritated eyes

- or sores in your mouth, throat, nose, or eyes. A severe and sometimes deadly problem called serotonin syndrome may happen if you take Zylrel (hydrocodone and acetaminophen liquid) with certain other drugs. Call your doctor right away if you have agitation

- change in balance

- confusion

- hallucinations

- fever

- fast or abnormal heartbeat

- flushing

- muscle twitching or stiffness

- seizures

- shivering or shaking

- sweating a lot

- severe diarrhea , upset stomach, or throwing up

- or severe headache . Zylrel side effects

Drug Side Effects Summary

2585. This particular disease, Pneumonia, interacts with the drug clindamycin and has the following

- Antibiotic medicines can cause diarrhea, which may be a sign of a new infection. If you have diarrhea that is watery or bloody, stop using clindamycin and call your doctor. Do not use anti-diarrhea medicine unless your doctor tells you to. (hives, difficult breathing, swelling in your face or throat) or a severe skin reaction (fever, sore throat, burning in your eyes, skin pain, red or purple skin rash that spreads and causes blistering and peeling). Seek medical treatment if you have symptoms of a serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash, fever, swollen glands, flu-like symptoms, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. This reaction may occur several weeks after you began using clindamycin. Clindamycin can cause diarrhea, which may be severe or lead to serious, life-threatening intestinal problems. If you have diarrhea that is watery or bloody, stop using clindamycin and call your doctor. Call your doctor at once if you have: any change in bowel habits
- severe stomach pain, diarrhea that is watery or bloody
- little or no urination
- or a metallic taste in your mouth (after clindamycin injection). Common clindamycin side effects may include: nausea, vomiting, stomach pain
- mild skin rash
- or vaginal itching or discharge
-

2586. This particular disease, Pneumonia, interacts with the drug cefotaxime and has the following

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- burning, irritation, or skin changes where the injection was given
- dark urine, jaundice (yellowing of the skin or eyes)

Drug Side Effects Summary

- a seizure
- fever, chills, tiredness
- or easy bruising, unusual bleeding, pale skin, cold hands and feet. Common side effects of cefotaxime may include: pain, bruising, swelling, or other irritation where the injection was given
- diarrhea
- fever
- or rash, itching.

2587. This particular disease, Pneumonia, interacts with the drug metronidazole and has the following

- new or worsening symptoms of infection
- painful or difficult urination
- confusion
- a light-headed feeling (like you might pass out)
- vaginal itching or discharge
- or blisters or ulcers in your mouth, red or swollen gums, trouble swallowing. Stop taking the medicine and call your doctor right away if you have neurologic side effects (more likely to occur while taking metronidazole long term): numbness, tingling, or burning pain in your hands or feet
- vision problems, pain behind your eyes, seeing flashes of light
- muscle weakness, problems with speech or coordination
- trouble speaking or understanding what is said to you
- a seizure
- or fever, neck stiffness, and increased sensitivity to light. Metronidazole can cause life-threatening liver problems in people with Cockayne syndrome. If you have this condition, stop taking this medicine and contact your doctor if you have signs of liver failure --nausea, stomach pain (upper right side), dark urine, clay-colored stools, or jaundice (yellowing of the skin or eyes). Side effects

Drug Side Effects Summary

may be more likely in older adults. Common metronidazole side effects may include: depression , trouble sleeping, feeling irritable

- headache , dizziness , weakness
- nausea, vomiting, loss of appetite, stomach pain
- diarrhea , constipation
- unpleasant metallic taste
- rash, itching
- vaginal itching or discharge, pain during sex
- mouth sores
- or swollen, red, or "hairy" tongue.

2588. This particular disease, Pneumonia, interacts with the drug vancomycin and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Certain side effects may occur if your body absorbs vancomycin through the intestinal walls. Call your doctor at once if you have: increased diarrhea that is watery or bloody
- hearing loss , ringing in your ears
- kidney problems - swelling, rapid weight gain, pain in your side or lower back, little or no urination
- or low potassium level - leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Side effects on the kidneys may be more likely in older adults taking this medicine. Common vancomycin side effects may include: low potassium
- nausea
- or stomach pain.

Drug Side Effects Summary

2589. This particular disease, Pneumonia, interacts with the drug cefepime and has the following side effects:

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- confusion, hallucinations
- trouble speaking, reading, or understanding the words of other people
- seizure (blackout or convulsions)
- pale skin, easy bruising, unusual bleeding
- or dark colored urine, jaundice (yellowing of the skin or eyes). Common side effects of cefepime may include: nausea , vomiting , diarrhea
- fever
- headache
- itching, rash
- or abnormal blood tests.

2590. This particular disease, Pneumonia, interacts with the drug ceftazidime and has the following side effects:

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- pale or yellowed skin, dark colored urine, fever, or weakness
- confusion, hallucinations, severe weakness
- involuntary muscle movement
- seizure (black-out or convulsions)
- or a cold feeling, discoloration, or skin changes in your fingers. Common side effects of ceftazidime injection may include: allergic reaction
- numbness , tingling, burning pain
- headache , dizziness

Drug Side Effects Summary

- nausea , vomiting , diarrhea, stomach pain
- or signs of a fungal infection--sores or white patches in your mouth or throat, vaginal itching or discharge.

2591. This particular disease, Pneumonia, interacts with the drug Flagyl and has the following side

- new or worsening symptoms of infection
- painful or difficult urination
- confusion
- a light-headed feeling (like you might pass out)
- vaginal itching or discharge
- or blisters or ulcers in your mouth, red or swollen gums, trouble swallowing. Stop taking the medicine and call your doctor right away if you have neurologic side effects (more likely to occur while taking metronidazole long term): numbness, tingling, or burning pain in your hands or feet
- vision problems, pain behind your eyes, seeing flashes of light
- muscle weakness, problems with speech or coordination
- trouble speaking or understanding what is said to you
- a seizure
- or fever, neck stiffness, and increased sensitivity to light. Metronidazole can cause life-threatening liver problems in people with Cockayne syndrome. If you have this condition, stop taking Flagyl and contact your doctor if you have signs of liver failure --nausea, stomach pain (upper right side), dark urine, clay-colored stools, or jaundice (yellowing of the skin or eyes). Side effects may be more likely in older adults. Common Flagyl side effects may include: depression , trouble sleeping, feeling irritable
- headache , dizziness , weakness
- nausea, vomiting, loss of appetite, stomach pain

Drug Side Effects Summary

- diarrhea , constipation
- unpleasant metallic taste
- rash, itching
- vaginal itching or discharge, pain during sex
- mouth sores
- or swollen, red, or "hairy" tongue.

2592. This particular disease, Pneumonia, interacts with the drug amikacin and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Amikacin may cause serious side effects. Call your doctor at once if you have: hearing loss , or a roaring sound in your ears
- severe or ongoing dizziness
- kidney problems--little or no urinating
- painful or difficult urination
- swelling in your feet or ankles
- feeling tired or short of breath
- weak or shallow breathing
- numbness or tingly feeling
- muscle twitching or seizure (convulsions)
- or severe stomach pain, diarrhea that is watery or bloody.

2593. This particular disease, Pneumonia, interacts with the drug Ancef and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your

Drug Side Effects Summary

doctor or nurse immediately if any of the following side effects occur: Rare Bluish color changes in skin color pain swelling of the foot or leg tenderness Incidence not known Abdominal or stomach cramps or tenderness back, leg, or stomach pains black, tarry stools bleeding gums blistering, peeling, or loosening of the skin bloating blood in the urine or stools bloody or cloudy urine chest pain chills clay-colored stools cloudy urine cough coughing up blood dark urine decrease in urine output or decrease in urine-concentrating ability decreased frequency or amount of urine diarrhea diarrhea, watery and severe, which may also be bloody difficult or painful urination difficulty with breathing or swallowing dizziness excessive muscle tone fast heartbeat feeling of discomfort fever general body swelling general tiredness and weakness headache hives increased blood pressure increased menstrual flow or vaginal bleeding increased thirst inflammation of the joints itching itching of the vagina or genital area joint or muscle pain light-colored stools loss of appetite lower back or side pain muscle aches or stiffness muscle tension or tightness nausea or vomiting nosebleeds pain during sexual intercourse pain, warmth, or burning in the fingers, toes, and legs pale skin paralysis pinpoint red spots on the skin problems with vision or hearing prolonged bleeding from cuts puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rash red irritated eyes red or black, tarry stools red or dark brown urine red skin lesions, often with a purple center red, irritated eyes restlessness seizures skin rash sore throat sores, ulcers, or white spots on the lips or in the mouth stomach cramps sudden decrease in the amount of urine swelling of the face, fingers, or lower legs swollen lymph glands swollen or painful glands thick, white vaginal discharge with no odor or with a mild odor tightness in the chest trouble sitting still troubled breathing unpleasant breath odor unusual bleeding or bruising unusual tiredness or weakness unusual weight loss upper right abdominal pain vomiting vomiting of blood weight gain wheezing yellowing of the eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of

Drug Side Effects Summary

these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Bleeding, blistering, burning, coldness, discoloration of the skin, feeling of pressure, hives, infection, inflammation, itching, lumps, numbness, pain, rash, redness, scarring, soreness, stinging, swelling, tenderness, tingling, ulceration, or warmth at the injection site hives or welts redness of the skin sore mouth or tongue weight loss white patches in the mouth, tongue, or throat Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Ancef side effects

2594. This particular disease, Pneumonia, interacts with the drug Zyvox and has the following side

- vision problems, changes in color vision
- severe stomach pain, diarrhea that is watery or bloody
- a seizure
- sweating, feeling anxious or shaky (may be signs of low blood sugar)
- high levels of serotonin in the body - agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting , diarrhea
- lactic acidosis - unusual muscle pain , trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness , feeling cold, or feeling very weak or tired
- or low blood cell counts - fever, chills, tiredness, weakness, confusion, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation, hallucinations, fever, sweating, shivering, fast heart rate, muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea. Common Zyvox side effects may include: nausea, vomiting, diarrhea

Drug Side Effects Summary

- skin rash
- anemia (low red blood cells)
- or headache , dizziness.

2595. This particular disease, Pneumonia, interacts with the drug Cleocin and has the following side effects:

- skin rash, fever, swollen glands, flu-like symptoms, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Call your doctor at once if you have: any change in bowel habits
- severe stomach pain, diarrhea that is watery or bloody
- little or no urination
- or a metallic taste in your mouth (after Cleocin injection). Common Cleocin side effects may include: nausea , vomiting , stomach pain
- mild skin rash
- or vaginal itching or discharge
-

2596. This particular disease, Pneumonia, interacts with the drug Flagyl IV and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: More common Agitation back pain blindness blurred vision burning, crawling, itching, numbness, pain, prickling, "pins and needles", or tingling feelings change in taste clay-colored stools confusion dark-colored urine decreased vision dizziness drowsiness eye pain fever headache irritability itching of the vagina or outside genitals light-colored stools loss of appetite loss of taste mood or mental changes nausea pain during sexual intercourse seeing or hearing things that are not there seizures stiff neck or back

Drug Side Effects Summary

stomach pain thick, white curd-like vaginal discharge without odor or with mild odor uncontrolled eye movements unpleasant breath odor unsteadiness or awkwardness unusual tiredness or weakness vomiting vomiting of blood weakness in the arms, hands, legs, or feet yellow eyes or skin Incidence not known Black, tarry stools blistering, peeling, or loosening of the skin changes in skin color changes in speech patterns chest pain chills cough diarrhea fainting fast, pounding, or irregular heartbeat or pulse feeling of warmth itching joint or muscle pain lower back or side pain pain, tenderness, or swelling of the foot or leg painful or difficult urination pale skin red skin lesions, often with a purple center red, irritated eyes redness of the face, neck, arms, and occasionally, upper chest shakiness and unsteady walk slurred speech sore throat sores, ulcers, or white spots in the mouth or on the lips trouble with speaking unsteadiness, trembling, or other problems with muscle control or coordination unusual bleeding or bruising Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Dry mouth hives or welts, skin rash redness of the skin stomach discomfort stuffy nose unpleasant metallic taste Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Flagyl IV side effects

2597. This particular disease, Pneumonia, interacts with the drug linezolid and has the following side effects:

- vision problems, changes in color vision
- severe stomach pain, diarrhea that is watery or bloody
- a seizure
- sweating, feeling anxious or shaky (may be signs of low blood sugar)

Drug Side Effects Summary

- lactic acidosis --unusual muscle pain, trouble breathing, stomach pain, vomiting, irregular heart rate, dizziness , feeling cold, or feeling very weak or tired
- or low blood cell counts--fever, chills, tiredness, weakness, confusion, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea , vomiting, or diarrhea. Common side effects of linezolid may include: nausea, vomiting, diarrhea
- rash
- anemia (low red blood cells)
- or headache , dizziness.

2598. This particular disease, Pneumonia, interacts with the drug piperacillin / tazobactam and has

- skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Piperacillin and tazobactam may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- confusion
- muscle twitching or stiffness, trouble walking
- a seizure
- low white blood cell counts--fever, mouth sores, skin sores, sore throat, cough, trouble breathing
- or low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of piperacillin and tazobactam may include: diarrhea, constipation
- nausea

Drug Side Effects Summary

- headache
- or sleep problems (insomnia).

2599. This particular disease, Pneumonia, interacts with the drug Zosyn and has the following side

- skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. To make sure Zosyn is safe for you, tell your doctor if you have ever had: severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- confusion
- muscle twitching or stiffness, trouble walking
- a seizure
- low white blood cell counts: may experience fever, mouth sores, skin sores, sore throat, cough, trouble breathing
- or low potassium: may experience leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common Zosyn side effects may include: diarrhea, constipation
- nausea
- headache
- or sleep problems (insomnia).

2600. This particular disease, Pneumonia, interacts with the drug cefazolin and has the following side

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- white patches or sores inside your mouth or on your lips
- fever, swollen glands, itching, joint pain, or not feeling well
- a seizure

Drug Side Effects Summary

- or liver problems--upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of cefazolin may include: nausea , vomiting , diarrhea
- headache
- pain where the medicine was injected
- rash
- or an allergic reaction.

2601. This particular disease, Pneumonia, interacts with the drug Cleocin HCl and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Cracks in the skin loss of heat from the body red, swollen skin scaly skin Incidence not known Abdominal or stomach cramps, pain, or tenderness black, tarry stools bleeding gums blistering, peeling, or loosening of the skin bloating blood in the urine or stools blurred vision chest pain chills clay-colored stools cloudy urine confusion cough or hoarseness dark urine decrease in the amount of urine diarrhea diarrhea, watery and severe, which may also be bloody difficulty with swallowing dizziness dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position dry mouth fast heartbeat fever with or without chills general feeling of tiredness or weakness headache heartburn heart stops hives or welts, itching, or skin rash increased thirst itching of the vagina or genital area joint or muscle pain loss of appetite lower back or side pain nausea or vomiting no breathing no pulse or blood pressure pain during sexual intercourse pain in the lower back or side painful or difficult urination pinpoint red spots on the skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rash with flat lesions or small raised lesions on the skin red skin lesions, often with a purple center red, irritated eyes redness of the skin sore throat sores, ulcers, or white spots in the

Drug Side Effects Summary

mouth or on the lips swollen glands thick, white vaginal discharge with no odor or with a mild odor thirst tightness in the chest trouble breathing unconscious unpleasant breath odor unusual bleeding or bruising unusual tiredness or weakness unusual weight loss vomiting of blood yellow eyes or skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Cleocin HCl side effects

2602. This particular disease, Pneumonia, interacts with the drug Cleocin Pediatric and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Cracks in the skin loss of heat from the body red, swollen skin scaly skin Incidence not known Abdominal or stomach cramps, pain, or tenderness black, tarry stools bleeding gums blistering, peeling, or loosening of the skin bloating blood in the urine or stools blurred vision chest pain chills clay-colored stools cloudy urine confusion cough or hoarseness dark urine decrease in the amount of urine diarrhea diarrhea, watery and severe, which may also be bloody difficulty with swallowing dizziness dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position dry mouth fast heartbeat fever with or without chills general feeling of tiredness or weakness headache heartburn heart stops hives or welts, itching, or skin rash increased thirst itching of the vagina or genital area joint or muscle pain loss of appetite lower back or side pain nausea or vomiting no breathing no pulse or blood pressure pain during sexual intercourse pain in the lower back or side painful or difficult urination pinpoint red spots on the skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rash with flat lesions or small raised lesions on the skin red skin lesions, often with a purple center red, irritated eyes redness of the skin sore throat sores, ulcers, or white spots in the mouth or on the lips swollen glands thick, white vaginal discharge with no odor or with a mild odor

Drug Side Effects Summary

thirst tightness in the chest trouble breathing unconscious unpleasant breath odor unusual bleeding or bruising unusual tiredness or weakness unusual weight loss vomiting of blood yellow eyes or skin Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Cleocin Pediatric side effects

2603. This particular disease, Pneumonia, interacts with the drug Cleocin Phosphate and has the fo

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: Rare Cracks in the skin loss of heat from the body red, swollen skin scaly skin Incidence not known Agitation black, tarry stools bleeding gums blistering, peeling, or loosening of the skin bloating blood in the urine or stools blurred vision chest pain or tightness chills clay-colored stools cloudy urine coma confusion cough or hoarseness dark urine decrease in the amount of urine depression diarrhea diarrhea, watery and severe, which may also be bloody difficulty with swallowing dizziness dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position dry mouth fast heartbeat fever with or without chills general feeling of tiredness or weakness headache heart stops hives or welts, itching, or skin rash hostility increased thirst irritability itching of the vagina or genital area joint or muscle pain lethargy loss of appetite lower back or side pain muscle twitching nausea or vomiting no breathing no pulse or blood pressure pain during sexual intercourse pain in the lower back or side pain, tenderness, or swelling of the foot or leg painful or difficult urination pinpoint red spots on the skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rapid weight gain rash with flat lesions or small raised lesions on the skin red skin lesions, often with a purple center red, irritated eyes redness of the skin seizures sore throat sores, ulcers, or white spots in the mouth or on the lips stomach cramps, pain, or tenderness swelling of the face, ankles, or hands

Drug Side Effects Summary

swollen glands thick, white vaginal discharge with no odor or with a mild odor thirst unconscious unpleasant breath odor unusual bleeding or bruising unusual tiredness or weakness unusual weight loss vomiting of blood yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Bitter taste changes in skin color pain, lump, or irritation at the injection site Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Cleocin Phosphate side effects

2604. This particular disease, Pneumonia, interacts with the drug Flagyl 375 and has the following side effects:

- No side effects listed

2605. This particular disease, Pneumonia, interacts with the drug tobramycin and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tobramycin may cause serious side effects. Call your doctor at once if you have: new or worsening breathing problems, such as wheezing, cough, chest tightness, or trouble breathing
- hearing problems, ringing in your ears
- hoarse voice
- severe dizziness , spinning sensation, balance problems
- weak or shallow breathing

Drug Side Effects Summary

- muscle weakness
- or kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath. Common side effects of tobramycin may include: cough, sore throat , hoarse voice
- feeling short of breath
- noisy breathing
- worsening of lung problems or cystic fibrosis
- coughing up mucus or blood
- altered sense of taste
- fever
- headache
- or rash.

2606. This particular disease, Pneumonia, interacts with the drug aztreonam and has the following :

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Aztreonam may cause serious side effects. Call your doctor at once if you have: bronchospasm (wheezing, cough, chest tightness, trouble breathing)
- or any new or worsening breathing problems. Common side effects of aztreonam may include: wheezing, chest discomfort
- stomach pain, vomiting
- fever (more common in children)
- or cough, sore throat , stuffy nose.

2607. This particular disease, Pneumonia, interacts with the drug ertapenem and has the following

- hives

Drug Side Effects Summary

- difficulty breathing
- swelling of your face, lips, tongue, or throat. Ertapenem may cause serious side effects. Call your doctor at once if you have: a seizure (convulsions)
- or severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose). Common side effects include: nausea , vomiting
- diarrhea
- headache
- or pain or redness where the injection was given.

2608. This particular disease, Pneumonia, interacts with the drug Garamycin and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Garamycin may cause serious side effects. Call your doctor at once if you have: hearing loss , or a roaring sound in your ears
- severe or ongoing dizziness
- weak or shallow breathing
- numbness or tingly feeling
- twitching, muscle tightness or contraction
- seizure (convulsions)
- severe stomach pain, diarrhea that is watery or bloody
- fever, blisters or ulcers in your mouth, red or swollen gums, trouble swallowing
- kidney problems--little or no urinating
- painful or difficult urination
- swelling in your feet or ankles
- feeling tired or short of breath

Drug Side Effects Summary

- signs of an electrolyte imbalance--confusion, weakness, bone pain, increased urination
- or increased pressure inside the skull--severe headaches, ringing in your ears, dizziness, nausea , vision problems, pain behind your eyes. Side effects may be more likely in older adults. Common side effects of Garamycin may include: vision problems
- nausea, vomiting , loss of appetite, weight loss
- a light-headed feeling, like you might pass out
- itching or rash
- pain where the medicine was injected
- headache , mood changes
- or joint pain.

2609. This particular disease, Pneumonia, interacts with the drug gentamicin and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Gentamicin may cause serious side effects. Call your doctor at once if you have: hearing loss , or a roaring sound in your ears
- severe or ongoing dizziness
- weak or shallow breathing
- numbness or tingly feeling
- twitching, muscle tightness or contraction
- seizure (convulsions)
- severe stomach pain, diarrhea that is watery or bloody
- fever, blisters or ulcers in your mouth, red or swollen gums, trouble swallowing
- kidney problems--little or no urinating
- painful or difficult urination

Drug Side Effects Summary

- swelling in your feet or ankles
- feeling tired or short of breath
- signs of an electrolyte imbalance--confusion, weakness, bone pain, increased urination
- or increased pressure inside the skull--severe headaches, ringing in your ears, dizziness, nausea , vision problems, pain behind your eyes. Side effects may be more likely in older adults. Common side effects of gentamicin may include: vision problems
- nausea, vomiting , loss of appetite, weight loss
- a light-headed feeling, like you might pass out
- itching or rash
- pain where the medicine was injected
- headache , mood changes
- or joint pain.

2610. This particular disease, Pneumonia, interacts with the drug Invanz and has the following side

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Invanz may cause serious side effects. Call your doctor at once if you have: a seizure (convulsions)
- or severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose). Common side effects include: nausea , vomiting
- diarrhea
- headache
- or pain or redness where the injection was given.

2611. This particular disease, Pneumonia, interacts with the drug Maxipime and has the following s

Drug Side Effects Summary

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- confusion, hallucinations
- trouble speaking, reading, or understanding the words of other people
- seizure (blackout or convulsions)
- pale skin, easy bruising, unusual bleeding
- or dark colored urine, jaundice (yellowing of the skin or eyes). Common side effects of Maxipime may include: nausea , vomiting , diarrhea
- fever
- headache
- itching, rash
- or abnormal blood tests.

2612. This particular disease, Pneumonia, interacts with the drug Penicillin VK and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness

Drug Side Effects Summary

- or swelling of the mouth, face, lips, tongue, or throat. Diarrhea is common with antibiotics . Rarely, a severe form called C diff-associated diarrhea (CDAD) may happen. Sometimes, this has led to a deadly bowel problem (colitis). CDAD may happen during or a few months after taking antibiotics. Call your doctor right away if you have stomach pain, cramps, or very loose, watery, or bloody stools. Check with your doctor before treating diarrhea. Penicillin VK side effects

2613. This particular disease, Pneumonia, interacts with the drug penicillin v potassium and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have any of these serious side effects: diarrhea that is watery or bloody
- fever, chills, body aches, flu symptoms
- easy bruising or bleeding, unusual weakness
- urinating less than usual or not at all
- severe skin rash, itching, or peeling
- agitation, confusion, unusual thoughts or behavior
- or seizure (black-out or convulsions). Less serious penicillin V side effects are more likely to occur, such as: nausea, vomiting, stomach pain
- vaginal itching or discharge
- headache
- swollen, black, or "hairy" tongue
- or thrush (white patches or inside your mouth or throat).

2614. This particular disease, Pneumonia, interacts with the drug Teflaro and has the following side effects:

- hives

Drug Side Effects Summary

- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- little or no urination
- a seizure
- drowsiness, tiredness, confusion, thinking problems
- low potassium - leg cramps, constipation , irregular heartbeats, fluttering in your chest, muscle weakness or limp feeling
- or low blood cell counts - sudden weakness or ill feeling, fever, chills, cold or flu symptoms, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath. Common Teflaro side effects may include: nausea
- diarrhea
- or rash.

2615. This particular disease, Pneumonia, interacts with the drug tigecycline and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Tigecycline may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- increased pressure inside the brain--severe headache , ringing in your ears, dizziness , nausea , vision problems, pain behind your eyes
- pancreas problems--severe pain in your upper stomach spreading to your back, nausea and vomiting , fast heart rate
- or liver problems--nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of tigecycline may

Drug Side Effects Summary

include: nausea, vomiting , stomach pain

- diarrhea
- headache
- or abnormal liver function tests .

2616. This particular disease, Pneumonia, interacts with the drug Tygacil and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- increased pressure inside the brain - severe headache, ringing in your ears, dizziness, nausea, vision problems, pain behind your eyes
- pancreas problems - severe pain in your upper stomach spreading to your back, nausea and vomiting, fast heart rate
- or liver problems - nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common Tygacil side effects may include: nausea, vomiting, stomach pain
- diarrhea
- headache
- or abnormal liver function tests.

2617. This particular disease, Pneumonia, interacts with the drug Vancocin and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs

Drug Side Effects Summary

of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of kidney problems like unable to pass urine, change in how much urine is passed, blood in the urine, or a big weight gain. Signs of low potassium levels like muscle pain or weakness, muscle cramps, or a heartbeat that does not feel normal. Swelling in the arms or legs. Flushing. Pain when passing urine. This medicine may cause hearing loss that may not go away. The chance may be higher if you use Vancocin (vancomycin capsules) for a long time. Do not use Vancocin (vancomycin capsules) longer than you have been told by your doctor. Tell your doctor right away if you have a change in hearing or hearing loss.

Vancocin side effects

2618. This particular disease, Pneumonia, interacts with the drug Vancocin HCl and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: Rare Black, tarry stools bleeding gums blood in the urine or stools continuing ringing or buzzing or other unexplained noise in the ears cough or hoarseness dizziness or lightheadedness feeling of constant movement of self or surroundings feeling of fullness in the ears fever with or without chills general feeling of tiredness or weakness hearing loss loss of balance lower back or side pain painful or difficult urination pale

Drug Side Effects Summary

skin pinpoint red spots on the skin ringing or buzzing in the ears sensation of spinning sore throat sores, ulcers, or white spots on the lips or in the mouth trouble in hearing unusual bleeding or bruising unusual tiredness or weakness Incidence not known Abdominal or stomach cramps or tenderness back pain blistering, peeling, loosening of the skin bloating bluish color blurred vision change in frequency of urination or amount of urine changes in skin color chest pain chest tightness confusion cracks in the skin diarrhea, watery and severe, which may also be bloody difficulty with swallowing dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position drowsiness fast heartbeat flushing headache hives, itching, or skin rash increased thirst joint or muscle pain loss of appetite loss of heat from the body nausea or vomiting pain, tenderness, or swelling of the foot or leg puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red skin lesions, often with a purple center red, irritated eyes red, swollen skin redness, soreness, or itching skin scaly skin sores, welting, or blisters sweating swelling of the feet or lower legs swollen glands unusual weight loss Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Vancocin HCl side effects

2619. This particular disease, Pneumonia, interacts with the drug Vancocin HCl Pulvules and has the

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Bladder pain bloating or swelling of the face, arms, hands, lower legs, or feet bloody or cloudy urine decreased urine difficult, burning, or painful urination dry mouth fever frequent urge to urinate increased thirst irregular heartbeat loss of appetite lower back or side pain mood changes muscle pain or cramps nausea numbness or tingling in the hands, feet, or lips seizures unusual tiredness or weakness unusual

Drug Side Effects Summary

weight gain or loss vomiting Less common Change in the frequency of urination or amount of urine drowsiness trouble breathing weakness Rare Hives, skin rash redness or other discoloration of the skin scaling or welting of the skin Incidence not known Black, tarry stools bleeding gums blistering, peeling, or loosening of the skin blurred vision chest pain or tightness chills confusion continuing ringing or buzzing or other unexplained noise in the ears cough difficulty with swallowing dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position fast heartbeat feeling of constant movement of self or surroundings feeling of fullness in the ears hearing loss itching joint or muscle pain loss of balance lower back or side pain pale skin pinpoint red spots on the skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue red skin lesions, often with a purple center red, irritated eyes ringing or buzzing in the ears sensation of spinning sore throat sores, ulcers, or white spots in the mouth or on the lips sweating swollen glands unusual bleeding or bruising Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Back pain bitter or unpleasant taste excess air or gas in the stomach or bowels headache mouth irritation passing gas stomach pain Incidence not known Depression difficulty having a bowel movement trouble sleeping Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Vancocin HCl Pulvules side effects

2620. This particular disease, Pneumonia, interacts with the drug Amikin and has the following side

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Amikin may cause serious side effects. Call your doctor at once if you have: hearing loss , or a roaring sound in your ears
- severe or ongoing dizziness
- kidney problems--little or no urinating
- painful or difficult urination
- swelling in your feet or ankles
- feeling tired or short of breath
- weak or shallow breathing
- numbness or tingly feeling
- muscle twitching or seizure (convulsions)
- or severe stomach pain, diarrhea that is watery or bloody.

2621. This particular disease, Pneumonia, interacts with the drug Amikin Pediatric and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: Incidence not known Agitation black, tarry stools bloody or cloudy urine bluish lips or skin blurred vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings chest pain chills coma confusion cough decrease in the amount of urine decreased urine output depression difficulty with breathing difficulty with moving dizziness dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position drowsiness dry mouth feeling of fullness in the ears fever headache hearing loss irritability lethargy loss of balance loss or change in hearing muscle pain or stiffness muscle twitching nausea not breathing pain in the joints pain in the lower back or side painful or difficult urination pale skin rapid weight gain ringing or buzzing in the ears seizures shakiness in the legs, arms, hands, or feet shortness of breath sore throat sores, ulcers, or white spots on the lips or in the

Drug Side Effects Summary

mouth stupor sweating swelling of the face, ankles, or hands swollen glands thirst trembling or shaking of the hands or feet trouble with hearing troubled breathing with exertion unusual bleeding or bruising unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Skin rash vomiting Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Amikin Pediatric side effects

2622. This particular disease, Pneumonia, interacts with the drug ampicillin / sulbactam and has the

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- thrush (white patches inside your mouth or throat)
- heart problems--fast heartbeats, chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating
- liver problems--nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or signs of a new infection--fever, headache , tiredness, ear pain or drainage, eye pain or redness, cough, skin rash, sores or pimples with pus, joint pain, ongoing neck or back pain , diarrhea, pain or burning when you urinate. Common side effects of ampicillin and sulbactam may include: diarrhea
- rash
- swollen, black, or "hairy" tongue

Drug Side Effects Summary

- vaginal itching or discharge
- pain where the medicine was injected.

2623. This particular disease, Pneumonia, interacts with the drug avibactam / ceftazidime and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Avibactam and cefTAZidime may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- a seizure (convulsions)
- rigid muscles, overactive reflexes
- confusion, hallucinations
- or a change in your mental state. Common side effects of avibactam and cefTAZidime may include: nausea , vomiting
- diarrhea
- or bruising where the IV needle is placed.

2624. This particular disease, Pneumonia, interacts with the drug Avycaz and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Avycaz may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- a seizure (convulsions)
- rigid muscles, overactive reflexes
- confusion, hallucinations
- or a change in your mental state. Common side effects of Avycaz may include: nausea , vomiting

Drug Side Effects Summary

- diarrhea
- or bruising where the IV needle is placed.

2625. This particular disease, Pneumonia, interacts with the drug Azactam and has the following side effects:

- severe stomach pain, diarrhea that is watery or bloody
- wheezing, chest pain
- easy bruising or bleeding
- a seizure
- or liver problems--loss of appetite, stomach pain (upper right side), dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Azactam may include: nausea , vomiting , diarrhea
- rash
- vaginal itching or discharge
- or pain, bruising, swelling, or irritation where the medicine was injected.

2626. This particular disease, Pneumonia, interacts with the drug BACiiM and has the following side effects:

- hives
- difficult breathing
- swelling of the face, lips, tongue, or throat. While receiving Baci-IM, your baby will be watched for the following side effects: diarrhea that is watery or bloody
- urinating more than usual or more often
- little or no urinating (fewer wet diapers)
- blood in the urine
- lower back pain
- or painful urination. Common side effects include: nausea , vomiting

Drug Side Effects Summary

- mild skin rash
- or pain, burning, or swelling where the medicine was injected.

2627. This particular disease, Pneumonia, interacts with the drug bacitracin and has the following side effects:

- hives
- difficult breathing
- swelling of the face, lips, tongue, or throat. While receiving bacitracin injection, your baby will be watched for the following side effects: diarrhea that is watery or bloody
- urinating more than usual or more often
- little or no urinating (fewer wet diapers)
- blood in the urine
- lower back pain
- or painful urination. Common side effects include: nausea , vomiting
- mild skin rash
- or pain, burning, or swelling where the medicine was injected.

2628. This particular disease, Pneumonia, interacts with the drug Baxdela and has the following side effects:

- hives , or the first sign of a skin rash
- fast heartbeats, difficult breathing
- swelling of your face, lips, tongue, or throat. Delafloxacin can cause serious side effects, including tendon problems, side effects on your nerves (which may cause permanent nerve damage), serious mood or behavior changes (after just one dose), or low blood sugar (which can lead to coma). Stop using this medicine and call your doctor at once if you have: low blood sugar - headache, hunger, sweating, irritability, dizziness , nausea , fast heart rate , or feeling anxious or shaky
- nerve symptoms in your hands, arms, legs, or feet - numbness, weakness, tingling, burning pain

Drug Side Effects Summary

- serious mood or behavior changes - nervousness, confusion, agitation, paranoia, hallucinations, memory problems, trouble concentrating, thoughts of suicide
- or signs of tendon rupture - sudden pain, swelling, bruising, tenderness, stiffness, movement problems, or a snapping or popping sound in any of your joints (rest the joint until you receive medical care or instructions). In rare cases, delafloxacin may cause damage to your aorta, the main blood artery of the body. This could lead to dangerous bleeding or death. Get emergency medical help if you have severe and constant pain in your chest, stomach, or back. Also, stop using this medicine and call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- a seizure
- muscle weakness or trouble breathing
- a light-headed feeling, like you might pass out
- the first sign of any skin rash, no matter how mild
- or increased pressure inside the skull - severe headaches, ringing in your ears, dizziness, nausea, vision problems, pain behind your eyes. Common Baxdela side effects may include: nausea, vomiting, diarrhea
- headache
- or abnormal liver function tests.

2629. This particular disease, Pneumonia, interacts with the drug Bicillin C-R and has the following

- slow heart rate , weak pulse, slow breathing (breathing may stop)
- pain, numbness , tingling, burning, or feeling cold
- pale or mottled skin, blue-colored lips, fingers, or toes
- severe pain or swelling in your lower leg
- weakness in your arms or legs

Drug Side Effects Summary

- or blistering, peeling, discoloration, or painful skin changes where the medicine was injected. Also call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- confusion, agitation , hallucinations (seeing or hearing things that are not real), extreme fear
- a seizure
- pain, swelling, warmth, redness, bruising, bleeding, or a lump where the medicine was injected
- pale or yellowed skin, bruising or bleeding, dark colored urine
- urination problems
- or signs of a new infection--fever, chills, mouth sores, vaginal itching or discharge. Seek medical attention right away if you develop a serious condition called methemoglobinemia . Symptoms include headache , fast heartbeats, shortness of breath, feeling tired or light-headed, or skin that looks pale, gray, or blue-colored. Common side effects of Bicillin C-R may include: nausea , vomiting , diarrhea
- itching, sweating, allergic reaction
- flushing (sudden warmth, redness, or tingly feeling)
- feeling anxious or nervous
- headache, dizziness , drowsiness
- or pain, swelling, or bruising where an injection was given.

2630. This particular disease, Pneumonia, interacts with the drug Bicillin C-R 900/300 and has the following side effects:

- slow heart rate , weak pulse, slow breathing (breathing may stop)
- pain, numbness , tingling, burning, or feeling cold
- pale or mottled skin, blue-colored lips, fingers, or toes
- severe pain or swelling in your lower leg
- weakness in your arms or legs

Drug Side Effects Summary

- or blistering, peeling, discoloration, or painful skin changes where the medicine was injected. Also call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- confusion, agitation , hallucinations (seeing or hearing things that are not real), extreme fear
- a seizure
- pain, swelling, warmth, redness, bruising, bleeding, or a lump where the medicine was injected
- pale or yellowed skin, bruising or bleeding, dark colored urine
- urination problems
- or signs of a new infection--fever, chills, mouth sores, vaginal itching or discharge. Seek medical attention right away if you develop a serious condition called methemoglobinemia . Symptoms include headache , fast heartbeats, shortness of breath, feeling tired or light-headed, or skin that looks pale, gray, or blue-colored. Common side effects of Bicillin C-R 900/300 may include: nausea , vomiting , diarrhea
- itching, sweating, allergic reaction
- flushing (sudden warmth, redness, or tingly feeling)
- feeling anxious or nervous
- headache, dizziness , drowsiness
- or pain, swelling, or bruising where an injection was given.

2631. This particular disease, Pneumonia, interacts with the drug Cefotan and has the following side effects:

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- fever, chills, body aches, flu symptoms
- pale skin, easy bruising, unusual bleeding
- a seizure

Drug Side Effects Summary

- fever, weakness, confusion
- dark colored urine, jaundice (yellowing of the skin or eyes)
- or kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath. Common side effects of Cefotan may include: nausea , vomiting
- diarrhea
- or vaginal itching or discharge.

2632. This particular disease, Pneumonia, interacts with the drug cefotetan and has the following side effects:

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- fever, chills, body aches, flu symptoms
- pale skin, easy bruising, unusual bleeding
- a seizure
- fever, weakness, confusion
- dark colored urine, jaundice (yellowing of the skin or eyes)
- or kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath. Common side effects of cefotetan may include: nausea , vomiting
- diarrhea
- or vaginal itching or discharge.

2633. This particular disease, Pneumonia, interacts with the drug cefoxitin and has the following side effects:

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- a light-headed feeling, like you might pass out
- little or no urination

Drug Side Effects Summary

- jaundice (yellowing of the skin or eyes)
- a seizure
- fever, chills, tiredness
- or easy bruising, unusual bleeding, pale skin, cold hands and feet. Common side effects of cefoxitin may include: pain, bruising, swelling, or other irritation where the injection was given
- diarrhea
- fever
- or rash, itching.

2634. This particular disease, Pneumonia, interacts with the drug ceftaroline and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Ceftaroline may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- little or no urination
- a seizure
- drowsiness, tiredness, confusion, thinking problems
- low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, muscle weakness or limp feeling
- or low blood cell counts-- sudden weakness or ill feeling, fever, chills, cold or flu symptoms, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath. Common side effects of ceftaroline may include: nausea
- diarrhea
- or rash.

Drug Side Effects Summary

2635. This particular disease, Pneumonia, interacts with the drug ceftolozane / tazobactam and has

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Ceftolozane and tazobactam may cause serious side effects. Call your doctor at once if you have: severe stomach pain
- diarrhea that is watery or bloody (even if it occurs months after your last dose)
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or symptoms of bleeding in the brain--sudden numbness or weakness, problems with vision or speech. Side effects may be more likely in older adults. Common side effects of ceftolozane and tazobactam may include: nausea , diarrhea
- headache
- fever
- kidney problems
- or abnormal liver function tests .

2636. This particular disease, Pneumonia, interacts with the drug Claforan and has the following side

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- burning, irritation, or skin changes where the injection was given
- dark urine, jaundice (yellowing of the skin or eyes)
- a seizure
- fever, chills, tiredness
- or easy bruising, unusual bleeding, pale skin, cold hands and feet. Common side effects of Claforan may include: pain, bruising, swelling, or other irritation where the injection was given
- diarrhea

Drug Side Effects Summary

- fever
- or rash, itching.

2637. This particular disease, Pneumonia, interacts with the drug delafloxacin and has the following

- hives , or the first sign of a skin rash
- fast heartbeats, difficult breathing
- swelling of your face, lips, tongue, or throat. Delafloxacin can cause serious side effects, including tendon problems, side effects on your nerves (which may cause permanent nerve damage), serious mood or behavior changes (after just one dose), or low blood sugar (which can lead to coma). Stop using this medicine and call your doctor at once if you have: low blood sugar--headache, hunger, sweating, irritability, dizziness , nausea , fast heart rate , or feeling anxious or shaky
- nerve symptoms in your hands, arms, legs, or feet--numbness, weakness, tingling, burning pain
- serious mood or behavior changes-- nervousness , confusion, agitation, paranoia, hallucinations, memory problems, trouble concentrating, thoughts of suicide
- or signs of tendon rupture--sudden pain, swelling, bruising, tenderness, stiffness, movement problems, or a snapping or popping sound in any of your joints (rest the joint until you receive medical care or instructions). In rare cases, delafloxacin may cause damage to your aorta, the main blood artery of the body. This could lead to dangerous bleeding or death. Get emergency medical help if you have severe and constant pain in your chest, stomach, or back. Also, stop using delafloxacin and call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- a seizure
- muscle weakness or trouble breathing
- a light-headed feeling, like you might pass out
- the first sign of any skin rash, no matter how mild

Drug Side Effects Summary

- or increased pressure inside the skull--severe headaches, ringing in your ears, dizziness, nausea, vision problems, pain behind your eyes. Common side effects of delafloxacin may include: nausea, vomiting , diarrhea
- headache
- or abnormal liver function tests .

2638. This particular disease, Pneumonia, interacts with the drug Fortaz and has the following side

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- pale or yellowed skin, dark colored urine, fever, or weakness
- confusion, hallucinations, severe weakness
- involuntary muscle movement
- seizure (black-out or convulsions)
- or a cold feeling, discoloration, or skin changes in your fingers. Common side effects of Fortaz may include: allergic reaction
- numbness , tingling, burning pain
- headache , dizziness
- nausea , vomiting , diarrhea, stomach pain
- or signs of a fungal infection--sores or white patches in your mouth or throat, vaginal itching or discharge.

2639. This particular disease, Pneumonia, interacts with the drug lefamulin and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Lefamulin may cause serious side effects. Call your

Drug Side Effects Summary

doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)

- or low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of lefamulin may include: nausea , vomiting , diarrhea
- sleep problems (insomnia)
- low potassium
- headache
- abnormal liver function tests
- or pain, bruising, swelling, or irritation where the medicine was injected.

2640. This particular disease, Pneumonia, interacts with the drug nafcillin and has the following side effects:

- hives , itching
- fever, stomach pain, muscle or joint pain
- difficulty breathing
- swelling of your face, lips, tongue, or throat. A delayed allergic reaction to nafcillin can occur as few as 2 days to as many as 4 weeks after you use the medication. Nafcillin may cause serious side effects. Call your doctor at once if you have: burning, pain, or swelling around the IV needle when the medicine is injected
- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- little or no urination
- painful urination, blood or pus in your urine
- pain in your side or lower back
- pain, swelling, bruising, or skin changes where the medicine was injected

Drug Side Effects Summary

- skin rash , itching, joint pain, or not feeling well
- fever, chills, swollen glands, sore throat , cough, trouble breathing
- or blisters or ulcers in your mouth, red or swollen gums, trouble swallowing. Common side effects of nafcillin may include: nausea , vomiting , diarrhea
- skin rash
- black or "hairy" tongue
- or tenderness or irritation around the IV needle.

2641. This particular disease, Pneumonia, interacts with the drug Nuzyra and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Nuzyra may cause serious side effects. Call your doctor at once if you have: little or no urination
- or increased pressure inside the skull--severe headaches, ringing in your ears, dizziness , nausea , vision problems, pain behind your eyes. Common side effects of Nuzyra may include: pain, irritation, redness, swelling, or a hard lump where the medicine was injected
- nausea, vomiting , diarrhea, constipation
- headache
- trouble sleeping
- increased blood pressure
- or abnormal liver function tests .

2642. This particular disease, Pneumonia, interacts with the drug omadacycline and has the following side effects:

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Omadacycline may cause serious side effects. Call your doctor at once if you have: little or no urination
- or increased pressure inside the skull--severe headaches, ringing in your ears, dizziness , nausea , vision problems, pain behind your eyes. Common side effects of omadacycline may include: pain, irritation, redness, swelling, or a hard lump where the medicine was injected
- nausea, vomiting , diarrhea, constipation
- headache
- trouble sleeping
- increased blood pressure
- or abnormal liver function tests .

2643. This particular disease, Pneumonia, interacts with the drug oxacillin and has the following side effects:

- hives , itching
- wheezing, difficult breathing
- feeling light-headed
- swelling of your face, lips, tongue, or throat. An allergic reaction to oxacillin can occur right after you use the medicine, or you may have a delayed reaction. An immediate allergic reaction can occur within 48 hours after you use oxacillin, and may cause fever with an itchy skin rash. A delayed reaction may occur past 48 hours and up to 4 weeks after you use oxacillin. Symptoms of a delayed reaction may include fever with swollen glands, rash or itching, joint pain, stomach pain, and a general ill feeling. Call your doctor or seek medical attention if you think you may be having an allergic reaction. Also call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody
- nausea , vomiting
- bruising or swelling around the IV needle

Drug Side Effects Summary

- kidney problems--little or no urination, painful or difficult urination, red or pink urine
- liver problems--nausea, upper stomach pain, itching, tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or low white blood cell counts--fever, chills, sore throat , swollen gums, mouth sores, pain when swallowing, skin sores, cold or flu symptoms, cough, trouble breathing. Common side effects of oxacillin may include: nausea, upset stomach , vomiting, mild diarrhea
- vaginal itching or discharge
- swollen, black, or "hairy" tongue
- or thrush (white patches or inside your mouth or throat).

2644. This particular disease, Pneumonia, interacts with the drug penicillin g benzathine / procaine

- slow heart rate , weak pulse, slow breathing (breathing may stop)
- pain, numbness , tingling, burning, or feeling cold
- pale or mottled skin, blue-colored lips, fingers, or toes
- severe pain or swelling in your lower leg
- weakness in your arms or legs
- or blistering, peeling, discoloration, or painful skin changes where the medicine was injected. Also call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- confusion, agitation , hallucinations (seeing or hearing things that are not real), extreme fear
- a seizure
- pain, swelling, warmth, redness, bruising, bleeding, or a lump where the medicine was injected
- pale or yellowed skin, bruising or bleeding, dark colored urine
- urination problems
- or signs of a new infection--fever, chills, mouth sores, vaginal itching or discharge. Seek medical

Drug Side Effects Summary

attention right away if you develop a serious condition called methemoglobinemia . Symptoms include headache , fast heartbeats, shortness of breath, feeling tired or light-headed, or skin that looks pale, gray, or blue-colored. Common side effects of penicillin G benzathine and penicillin G procaine may include: nausea , vomiting , diarrhea

- itching, sweating, allergic reaction
- flushing (sudden warmth, redness, or tingly feeling)
- feeling anxious or nervous
- headache, dizziness , drowsiness
- or pain, swelling, or bruising where an injection was given.

2645. This particular disease, Pneumonia, interacts with the drug penicillin g potassium and has the

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Penicillin G potassium may cause serious side effects.

Call your doctor at once if you have: the first sign of any skin rash , no matter how mild

- red or scaly skin
- fever, chills, swollen glands, muscle or joint pain, fast heartbeats, general ill feeling
- a light-headed feeling, like you might pass out
- severe stomach pain, diarrhea that is watery or bloody
- little or no urinating
- bruising, severe tingling, numbness , pain, muscle weakness
- seizure (convulsions)
- or unusual changes in mood or behavior. Common side effects of penicillin G potassium may include: mild diarrhea
- headache

Drug Side Effects Summary

- black or hairy tongue
- or pain, swelling, bruising, or irritation around the IV needle.

2646. This particular disease, Pneumonia, interacts with the drug penicillin g sodium and has the following side effects:

- hives , itching
- feeling like you might pass out
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Penicillin G sodium may cause serious side effects.

Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody

- fever, chills, swollen glands, skin sores, muscle pain, feeling short of breath, warmth or redness under your skin, severe dizziness
- white patches or sores inside your mouth or on your lips
- easy bruising, skin rash , unusual bleeding, pale or yellowed skin, severe tingling, numbness , muscle weakness
- red or pink urine, dark colored urine
- swelling in your hands or feet
- muscle twitching , seizures (convulsions)
- or signs of an electrolyte imbalance--dry mouth, increased thirst, mood changes, confusion, stomach pain, vomiting , muscle pain or weakness, lack of energy, irregular heartbeats, dark urine.

Common side effects of penicillin G sodium may include: mild diarrhea

- nausea , vomiting
- black or hairy tongue
- or pain, swelling, bruising, or irritation around the IV needle.

2647. This particular disease, Pneumonia, interacts with the drug Pfizerpen and has the following side effects:

Drug Side Effects Summary

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Pfizerpen may cause serious side effects. Call your doctor at once if you have: the first sign of any skin rash , no matter how mild
- red or scaly skin
- fever, chills, swollen glands, muscle or joint pain, fast heartbeats, general ill feeling
- a light-headed feeling, like you might pass out
- severe stomach pain, diarrhea that is watery or bloody
- little or no urinating
- bruising, severe tingling, numbness , pain, muscle weakness
- seizure (convulsions)
- or unusual changes in mood or behavior. Common side effects of Pfizerpen may include: mild diarrhea
- headache
- black or hairy tongue
- or pain, swelling, bruising, or irritation around the IV needle.

2648. This particular disease, Pneumonia, interacts with the drug piperacillin and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever

Drug Side Effects Summary

- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Any unexplained bruising or bleeding. Fever or chills. Sore throat . Seizures . Extra muscle action. Diarrhea is common with antibiotics . Rarely, a severe form called C diff-associated diarrhea (CDAD) may happen. Sometimes, this has led to a deadly bowel problem (colitis). CDAD may happen during or a few months after taking antibiotics. Call your doctor right away if you have stomach pain, cramps, or very loose, watery, or bloody stools. Check with your doctor before treating diarrhea. Piperacillin side effects

2649. This particular disease, Pneumonia, interacts with the drug procaine penicillin and has the fo

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have a serious side effect such as: diarrhea that is watery or bloody
- peeling skin , severe pain, or changes in skin color where the medicine was injected
- dizziness , joint or muscle pain
- fast or pounding heartbeats
- numbness , tingling, pain, swelling, or redness in your arms or legs
- confusion, agitation, depression , unusual thoughts or behavior
- chest pain, problems with vision or speech
- feeling like you might pass out
- fever, chills, dizziness, muscle pain, rapid breathing or heart rate
- uncontrolled muscle movements, problems with balance or walking

Drug Side Effects Summary

- pale skin, easy bruising or bleeding, unusual weakness
- sore throat , flu symptoms
- urinating less than usual or not at all
- rash or itching with swollen glands, joint pain, or general ill feeling
- or slow heart rate , weak pulse, fainting, slow breathing. Less serious side effects may include:
pain where the injection was given
- vaginal itching or discharge
- white patches in your mouth or throat
- nausea , vomiting
- blurred vision, ringing in your ears
- headache , dizziness
- or mild skin rash ,

2650. This particular disease, Pneumonia, interacts with the drug Tazicef and has the following side

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- pale or yellowed skin, dark colored urine, fever, or weakness
- confusion, hallucinations, severe weakness
- involuntary muscle movement
- seizure (black-out or convulsions)
- or a cold feeling, discoloration, or skin changes in your fingers. Common side effects of Tazicef may include: allergic reaction
- numbness , tingling, burning pain
- headache , dizziness
- nausea , vomiting , diarrhea, stomach pain

Drug Side Effects Summary

- or signs of a fungal infection--sores or white patches in your mouth or throat, vaginal itching or discharge.

2651. This particular disease, Pneumonia, interacts with the drug telavancin and has the following side effects:

- Some side effects may occur during the injection. Tell your caregiver if you feel itchy or tingly, or have a red rash during the injection. hives
- rapid heartbeats
- difficult breathing, fainting
- swelling of your face, lips, tongue, or throat. Telavancin may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- or signs of kidney problems--little or no urinating, swelling in your ankles or feet, weight gain, or urine that looks foamy. Common side effects of telavancin may include: foamy urine
- nausea , vomiting
- diarrhea
- or changes in your sense of taste.

2652. This particular disease, Pneumonia, interacts with the drug Tobo and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tobo may cause serious side effects. Call your doctor at once if you have: new or worsening breathing problems, such as wheezing, cough, chest tightness, or trouble breathing

Drug Side Effects Summary

- hearing problems, ringing in your ears
- hoarse voice
- severe dizziness , spinning sensation, balance problems
- weak or shallow breathing
- muscle weakness
- or kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath. Common side effects of Tobin may include: cough, sore throat , hoarse voice
- feeling short of breath
- noisy breathing
- worsening of lung problems or cystic fibrosis
- coughing up mucus or blood
- altered sense of taste
- fever
- headache
- or rash.

2653. This particular disease, Pneumonia, interacts with the drug Unasyn and has the following side effects:

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- thrush (white patches inside your mouth or throat)
- heart problems--fast heartbeats, chest pain or pressure, pain spreading to your jaw or shoulder, nausea , sweating
- liver problems--nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or signs of a new infection--fever, headache , tiredness, ear pain or drainage, eye pain or redness,

Drug Side Effects Summary

cough, skin rash, sores or pimples with pus, joint pain, ongoing neck or back pain , diarrhea, pain or burning when you urinate. Common side effects of Unasyn may include: diarrhea

- rash
- swollen, black, or "hairy" tongue
- vaginal itching or discharge
- pain where the medicine was injected.

2654. This particular disease, Pneumonia, interacts with the drug Vibativ and has the following side effects:

- Some side effects may occur during the injection. Tell your caregiver if you feel itchy or tingly, or have a red rash during the injection. hives
- rapid heartbeats
- difficult breathing, fainting
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- or signs of kidney problems - little or no urinating, swelling in your ankles or feet, weight gain, or urine that looks foamy. Common Vibativ side effects may include: foamy urine
- nausea , vomiting
- diarrhea
- or changes in your sense of taste.

2655. This particular disease, Pneumonia, interacts with the drug Xenleta and has the following side effects:

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Xenleta may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- or low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of Xenleta may include: nausea , vomiting , diarrhea
- sleep problems (insomnia)
- low potassium
- headache
- abnormal liver function tests
- or pain, bruising, swelling, or irritation where the medicine was injected.

2656. This particular disease, Pneumonia, interacts with the drug Zerbaxa and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Zerbaxa may cause serious side effects. Call your doctor at once if you have: severe stomach pain
- diarrhea that is watery or bloody (even if it occurs months after your last dose)
- kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath
- or symptoms of bleeding in the brain--sudden numbness or weakness, problems with vision or speech. Side effects may be more likely in older adults. Common side effects of Zerbaxa may include: nausea , diarrhea
- headache
- fever
- kidney problems

Drug Side Effects Summary

- or abnormal liver function tests .

2657. This particular disease, Psoriasis, interacts with the drug Humira and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor right away if you have any symptoms of lymphoma : fever, swollen glands, night sweats , general feeling of illness
- joint and muscle pain , skin rash , easy bruising or bleeding
- pale skin, feeling light-headed or short of breath, cold hands and feet
- pain in your upper stomach that may spread to your shoulder
- or loss of appetite, feeling full after eating only a small amount, weight loss . Also call your doctor at once if you have: new or worsening psoriasis (raised, silvery flaking of the skin)
- a sore or bump on your skin that does not heal
- symptoms of sepsis - confusion, chills, severe drowsiness, fast heartbeats, rapid breathing, feeling very ill
- liver problems - body aches, tiredness, stomach pain, right-sided upper stomach pain, vomiting, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- lupus -like syndrome - joint pain or swelling, chest pain, shortness of breath, patchy skin color that worsens in sunlight
- nerve problems - numbness, tingling, dizziness , vision problems, weakness in your arms or legs
- or signs of tuberculosis - fever with ongoing cough, weight loss (fat or muscle). Older adults may be more likely to develop infections or cancer while using adalimumab. Common side effects of Humira may include: headache
- cold symptoms such as stuffy nose, sinus pain, sneezing, sore throat
- rash

Drug Side Effects Summary

- or redness, bruising, itching, or swelling where the injection was given.

2658. This particular disease, Psoriasis, interacts with the drug Stelara and has the following side effects:

- hives
- chest pain, difficult breathing
- feeling light-headed
- swelling of your face, lips, tongue, or throat. Serious infections may occur during treatment with Stelara. Call your doctor right away if you have signs of infection such as: fever, chills, muscle pain, shortness of breath, weight loss, diarrhea or stomach pain, burning when you urinate, feeling very tired, skin warmth or redness, painful skin sores, or coughing up blood. Also call your doctor at once if you have: a mole that has changed in size or color
- swelling, pain, warmth, or redness anywhere on your body
- stomach pain that is sudden and severe or comes on slowly, changes in bowel habits (diarrhea or constipation)
- new or worsening cough, sudden chest pain, feeling short of breath
- pain or burning when you urinate
- or severe headache , confusion, change in mental status, vision problems, and/or seizure (convulsions). Common Stelara side effects may include: fever, flu-like symptoms
- infection
- itching
- stomach pain, nausea, vomiting , diarrhea
- vaginal itching or discharge
- pain or burning when you urinate
- cough with mucus, shortness of breath, chest discomfort
- headache, tiredness

Drug Side Effects Summary

- or redness at the injection site.

2659. This particular disease, Psoriasis, interacts with the drug ustekinumab and has the following

- What are the most common side effects of ustekinumab ? cough with mucus, shortness of breath, chest discomfort Infection fever, flu-like symptoms headache, tiredness Itching redness at the injection site vaginal itching or discharge pain or burning when you urinate stomach pain, nausea, vomiting, diarrhea. What are the serious (bad) side effects of ustekinumab ? hives
- chest pain, difficult breathing
- feeling light-headed
- swelling of your face, lips, tongue, or throat. Serious infections may occur during treatment with ustekinumab . Call your doctor right away if you have signs of infection such as: fever, chills, muscle pain, shortness of breath, weight loss, diarrhea or stomach pain, burning when you urinate, feeling very tired, skin warmth or redness, painful skin sores, or coughing up blood. Also call your doctor at once if you have: a mole that has changed in size or color swelling, pain, warmth, or redness anywhere on your body stomach pain that is sudden and severe or comes on slowly, changes in bowel habits (diarrhea or constipation) new or worsening cough, sudden chest pain, feeling short of breath pain or burning when you urinate severe headache, confusion, change in mental status, vision problems, and/or seizure (convulsions).

2660. This particular disease, Psoriasis, interacts with the drug adalimumab and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor right away if you have any symptoms of lymphoma : fever, swollen glands, night sweats , general feeling of illness
- joint and muscle pain, skin rash , easy bruising or bleeding

Drug Side Effects Summary

- pale skin, feeling light-headed or short of breath, cold hands and feet
- pain in your upper stomach that may spread to your shoulder
- or loss of appetite, feeling full after eating only a small amount, weight loss . Also call your doctor at once if you have: new or worsening psoriasis (raised, silvery flaking of the skin)
- a sore or bump on your skin that does not heal
- symptoms of sepsis --confusion, chills, severe drowsiness, fast heartbeats, rapid breathing, feeling very ill
- liver problems--body aches, tiredness, stomach pain, right-sided upper stomach pain, vomiting , loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- lupus -like syndrome--joint pain or swelling, chest pain, shortness of breath, patchy skin color that worsens in sunlight
- nerve problems--numbness, tingling, dizziness , vision problems, weakness in your arms or legs
- or signs of tuberculosis--fever with ongoing cough, weight loss (fat or muscle). Older adults may be more likely to develop infections or cancer while using adalimumab. Common side effects of adalimumab may include: headache
- cold symptoms such as stuffy nose, sinus pain, sneezing, sore throat
- rash
- or redness, bruising, itching, or swelling where the injection was given.

2661. This particular disease, Psoriasis, interacts with the drug Cosentyx and has the following side

- hives
- chest tightness, difficult breathing
- feeling like you might pass out
- swelling of your face, lips, tongue, or throat. Call your doctor right away if you have signs of infection such as: redness, warmth, or painful sores on your skin

Drug Side Effects Summary

- cough, shortness of breath, cough with red or pink mucus
- increased urination, burning when you urinate
- sores or white patches in your mouth or throat (yeast infection or "thrush")
- new or worsening diarrhea or stomach pain
- or fever, chills, sweating, muscle pain, weight loss. Further doses may be delayed until your infection clears up. Common Cosentyx side effects may include: diarrhea
- or cold symptoms such as stuffy nose, sneezing, sore throat .

2662. This particular disease, Psoriasis, interacts with the drug Taltz and has the following side effects:

- rash
- chest tightness, difficult breathing
- feeling like you might pass out
- swelling of your face, eyelids, lips, mouth, tongue, or throat. Call your doctor at once if you have: fever, chills, sweating, muscle pain, weight loss
- diarrhea (may be bloody), stomach cramps
- painful skin sores
- cough, shortness of breath, cough with red or pink mucus
- sores or white patches in your mouth or throat (yeast infection or "thrush")
- increased urination, pain or burning when you urinate
- eye swelling, redness, crusting or drainage (may be signs of infection)
- or signs of a fungal infection (rash or red patches, itching, burning, cracked skin, hair loss).

Common Taltz side effects may include: pain or redness where the medicine was injected

- nausea
- ear infection
- or cold symptoms such as stuffy nose, sneezing, sore throat.

Drug Side Effects Summary

2663. This particular disease, Psoriasis, interacts with the drug Skyrizi and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor right away if you have signs of infection such as: fever, chills, sweating, body aches
- shortness of breath, cough, bloody mucus
- mouth sores, red or swollen gums
- stomach pain, diarrhea
- increased urination, burning when you urinate
- pale skin, easy bruising, unusual bleeding
- a fungal skin infection - skin sores different from psoriasis, rash or redness, blisters, itching, burning, cracking or peeling, changes in skin color
- or signs of tuberculosis: fever, cough, night sweats , loss of appetite, weight loss, and feeling very tired. Further doses may be delayed until your infection clears up. Skyrizi may cause serious side effects including liver problems in adults with Crohn's disease. Your healthcare provider will do blood tests to check your liver before, during, and up to 12 weeks of treatment. Your healthcare provider may stop treatment if you develop liver problems. Tell your healthcare provider right away if you notice any of the following symptoms: unexplained rash
- nausea
- vomiting
- stomach (abdominal) pain
- tiredness (fatigue)
- loss of appetite
- yellowing of the skin and eyes (jaundice)
- dark urine. Common Skyrizi side effects in adults treated for Crohn's disease may include: cold

Drug Side Effects Summary

symptoms such as stuffy nose, sneezing, sore throat

- pain, redness, itching, bruising, swelling, warmth, bleeding, burning, or other skin irritation where the medicine was injected

- fever

- headache

- stomach (abdominal) pain

- back pain

- joint pain

- low red blood cells (anemia). Common Skyrizi side effects in adults treated for plaque psoriasis and psoriatic arthritis may include: cold symptoms such as stuffy nose, sneezing, sore throat

- headache

- feeling tired

- fungal skin infections

- injection site reactions.

2664. This particular disease, Psoriasis, interacts with the drug Tremfya and has the following side

- hives, rash, itching

- chest tightness, difficult breathing

- feeling light-headed

- swelling of your face, lips, tongue, or throat. Guselkumab can weaken (suppress) your immune system, and you may get an infection more easily. Call your doctor at once if you have signs of infection, such as: fever, chills, body aches, night sweats

- weight loss, feeling very tired

- cough (may contain blood or mucus), shortness of breath

- pain or burning when you urinate

Drug Side Effects Summary

- severe diarrhea or stomach cramps
- or skin redness, tingling, blisters, oozing, or sores that look different from psoriasis. Common Tremfya side effects may include: headache, joint pain
- diarrhea, stomach pain
- cold symptoms such as stuffy nose, sneezing, sore throat
- cough, feeling short of breath
- skin infections
- or pain, itching, swelling, redness, or bruising where the medicine was injected.

2665. This particular disease, Psoriasis, interacts with the drug Ilumya and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. You may get infections more easily, even serious or fatal infections. Call your doctor right away if you have signs of infection such as: fever, chills, sweating
- skin sores
- muscle pain
- increased urination, pain or burning when you urinate
- stomach pain, diarrhea, weight loss
- or cough, shortness of breath, coughing up pink or red mucus. Common Ilumya side effects may include: pain, itching, rash, redness, swelling, bruising, or bleeding where the medicine was injected
- diarrhea
- or cold symptoms such as stuffy nose, sneezing, sore throat .

2666. This particular disease, Psoriasis, interacts with the drug Avsola and has the following side effects

Drug Side Effects Summary

- Avsola can cause serious side effects, including: See "Important information" above. Serious Infections Some patients, especially those 65 years and older have had serious infections while receiving infliximab products, such as Avsola. These serious infections include TB and infections caused by viruses, fungi, or bacteria that have spread throughout the body or cause infections in certain areas (such as skin). Some patients die from these infections. If you get an infection while receiving treatment with Avsola your doctor will treat your infection and may need to stop your Avsola treatment. Tell your doctor right away if you have any of the following signs of an infection while receiving or after receiving Avsola: a fever feel very tired have a cough have flu-like symptoms warm, red, or painful skin Your doctor will examine you for TB and perform a test to see if you have TB. If your doctor feels that you are at risk for TB, you may be treated with medicine for TB before you begin treatment with Avsola and during treatment with Avsola. Even if your TB test is negative, your doctor should carefully monitor you for TB infections while you are receiving Avsola. Patients who had a negative TB skin test before receiving infliximab products have developed active TB. If you are a chronic carrier of the hepatitis B virus, the virus can become active while you are being treated with Avsola. In some cases, patients have died as a result of hepatitis B virus being reactivated. Your doctor should do a blood test for hepatitis B virus before you start treatment with Avsola and occasionally while you are being treated. Tell your doctor if you have any of the following symptoms: feel unwell poor appetite tiredness (fatigue) fever, skin rash, or joint pain Heart Failure If you have a heart problem called congestive heart failure, your doctor should check you closely while you are receiving Avsola. Your congestive heart failure may get worse while you are receiving Avsola. Be sure to tell your doctor of any new or worse symptoms including: shortness of breath swelling of ankles or feet sudden weight gain Treatment with Avsola may need to be stopped if you get new or worse congestive heart failure. Other Heart Problems Some patients have experienced a heart attack (some of which led to death), low blood flow to the heart, or abnormal heart rhythm within 24 hours of beginning their infusion of infliximab products. Symptoms may include chest

Drug Side Effects Summary

discomfort or pain, arm pain, stomach pain, shortness of breath, anxiety, lightheadedness, dizziness, fainting, sweating, nausea, vomiting, fluttering or pounding in your chest, and/or a fast or a slow heartbeat. Tell your doctor right away if you have any of these symptoms. Liver Injury Some patients receiving infliximab products have developed serious liver problems. Tell your doctor if you have: jaundice (skin and eyes turning yellow) dark brown-colored urine pain on the right side of your stomach area (right-sided abdominal pain) fever extreme tiredness (severe fatigue) Blood Problems In some patients receiving infliximab products, the body may not make enough of the blood cells that help fight infections or help stop bleeding. Tell your doctor if you: have a fever that does not go away bruise or bleed very easily look very pale Nervous System Disorders Some patients receiving infliximab products have developed problems with their nervous system. Tell your doctor if you have: changes in your vision weakness in your arms or legs numbness or tingling in any part of your body seizures Some patients have experienced a stroke within approximately 24 hours of their infusion with infliximab products. Tell your doctor right away if you have symptoms of a stroke which may include: numbness or weakness of the face, arm or leg, especially on one side of the body

- sudden confusion, trouble speaking or understanding

- sudden trouble seeing in one or both eyes, sudden trouble walking, dizziness, loss of balance or coordination or a sudden, severe headache. Allergic Reactions Some patients have had allergic reactions to infliximab products. Some of these reactions were severe. These reactions can happen while you are getting your Avsola treatment or shortly afterward. Your doctor may need to stop or pause your treatment with Avsola and may give you medicines to treat the allergic reaction. Signs of an allergic reaction can include: hives (red, raised, itchy patches of skin) difficulty breathing chest pain high or low blood pressure fever chills Some patients treated with infliximab products have had delayed allergic reactions. The delayed reactions occurred 3 to 12 days after receiving treatment with infliximab products. Tell your doctor right away if you have any of these signs of delayed allergic reaction to Avsola: fever rash headache sore throat muscle or joint pain swelling of the face and

Drug Side Effects Summary

hands difficulty swallowing Lupus-like Syndrome Some patients have developed symptoms that are like the symptoms of Lupus. If you develop any of the following symptoms, your doctor may decide to stop your treatment with Avsola. chest discomfort or pain that does not go away shortness of breath joint pain rash on the cheeks or arms that gets worse in the sun Psoriasis Some people receiving infliximab products had new psoriasis or worsening of psoriasis they already had. Tell your doctor if you develop red scaly patches or raised bumps on the skin that are filled with pus. Your doctor may decide to stop your treatment with Avsola. The most common side effects of infliximab products include: respiratory infections, such as sinus infections and sore throat headache coughing stomach pain Infusion reactions can happen up to 2 hours after your infusion of Avsola. Symptoms of infusion reactions may include: fever chills chest pain low blood pressure or high blood pressure shortness of breath rash itching Children with Crohn's disease showed some differences in side effects of treatment compared with adults with Crohn's disease. The side effects that happened more in children were: anemia (low red blood cells), leukopenia (low white blood cells), flushing (redness or blushing), viral infections, neutropenia (low neutrophils, the white blood cells that fight infection), bone fracture, bacterial infection and allergic reactions of the breathing tract. Among patients who received infliximab for ulcerative colitis in clinical studies, more children had infections as compared with adults. Tell your doctor about any side effect that bothers you or does not go away. These are not all of the side effects with Avsola. Ask your doctor or pharmacist for more information. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Avsola side effects

2667. This particular disease, Psoriasis, interacts with the drug Duobrii and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: irritation of

Drug Side Effects Summary

treated skin

- a sunburn
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin - weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Common Duobrii side effects may include: burning, stinging, redness, itching, pain, or swelling of treated skin
- redness or crusting around your hair follicles
- rash
- peeling skin
- or thinning skin.

2668. This particular disease, Psoriasis, interacts with the drug guselkumab and has the following side effects:

- Guselkumab may cause serious side effects including: See "Important information" above. The most common side effects of guselkumab include: upper respiratory infections joint pain (arthralgia) fungal skin infections headache diarrhea herpes simplex infections injection site reactions stomach flu (gastroenteritis) bronchitis These are not all the possible side effects of guselkumab. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Guselkumab side effects

2669. This particular disease, Psoriasis, interacts with the drug halobetasol / tazarotene and has the following side effects:

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Halobetasol and tazarotene topical may cause serious side effects. Call your doctor at once if you have: irritation of treated skin
- a sunburn
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Common side effects of halobetasol and tazarotene topical may include: burning, stinging, redness, itching, pain, or swelling of treated skin
- redness or crusting around your hair follicles
- rash
- peeling skin
- or thinning skin.

2670. This particular disease, Psoriasis, interacts with the drug Inflectra and has the following side

- Serious side effects of Inflectra include: See "Important information" above. Serious Infections Some patients, especially those 65 years and older have had serious infections while receiving infliximab products, such as Inflectra. These serious infections include TB and infections caused by viruses, fungi, or bacteria that have spread throughout the body. Some patients die from these infections. If you get an infection while receiving treatment with Inflectra your doctor will treat your infection and may need to stop your Inflectra treatment. Tell your doctor right away if you have any of the following signs of an infection while receiving or after receiving Inflectra: a fever feel very tired have a cough have flu-like symptoms warm, red, or painful skin Your doctor will examine you for TB and perform a test to see if you have TB. If your doctor feels that you are at risk for TB, you may be

Drug Side Effects Summary

treated with medicine for TB before you begin treatment with Inflectra and during treatment with Inflectra. Even if your TB test is negative, your doctor should carefully monitor you for TB infections while you are receiving Inflectra. Patients who had a negative TB skin test before receiving infliximab products have developed active TB. If you are a chronic carrier of the hepatitis B virus, the virus can become active while you are being treated with Inflectra. In some cases, patients have died as a result of hepatitis B virus being reactivated. Your doctor should do a blood test for hepatitis B virus before you start treatment with Inflectra and occasionally while you are being treated. Tell your doctor if you have any of the following symptoms: feel unwell tiredness (fatigue) poor appetite fever, skin rash, or joint pain

Heart Failure If you have a heart problem called congestive heart failure, your doctor should check you closely while you are receiving Inflectra. Your congestive heart failure may get worse while you are receiving Inflectra. Be sure to tell your doctor of any new or worse symptoms including: shortness of breath swelling of ankles or feet sudden weight gain Treatment with Inflectra may need to be stopped if you get new or worse congestive heart failure.

Other Heart Problems Some patients have experienced a heart attack (some of which led to death), low blood flow to the heart, or abnormal heart rhythm within 24 hours of beginning their infusion of an infliximab product. Symptoms may include chest discomfort or pain, arm pain, stomach pain, shortness of breath, anxiety, lightheadedness, dizziness, fainting, sweating, nausea, vomiting, fluttering or pounding in your chest, and/or a fast or a slow heartbeat. Tell your doctor right away if you have any of these symptoms.

Liver Injury Some patients receiving infliximab products have developed serious liver problems. Tell your doctor if you have: jaundice (skin and eyes turning yellow) dark brown-colored urine pain on the right side of your stomach area (right-sided abdominal pain) fever extreme tiredness (severe fatigue)

Blood Problems In some patients receiving infliximab products, the body may not make enough of the blood cells that help fight infections or help stop bleeding. Tell your doctor if you: have a fever that does not go away bruise or bleed very easily look very pale

Nervous System Disorders Some patients receiving infliximab products have developed

Drug Side Effects Summary

problems with their nervous system. Tell your doctor if you have: changes in your vision numbness or tingling in any part of your body weakness in your arms or legs seizures Some patients have experienced a stroke within approximately 24 hours of their infusion with infliximab products. Tell your doctor right away if you have symptoms of a stroke which may include: numbness or weakness of the face, arm or leg, especially on one side of the body

- sudden confusion, trouble speaking or understanding

- sudden trouble seeing in one or both eyes, sudden trouble walking, dizziness, loss of balance or coordination or a sudden, severe headache.

Allergic Reactions Some patients have had allergic reactions to infliximab products. Some of these reactions were severe. These reactions can happen while you are getting your Inflectra treatment or shortly afterward. Your doctor may need to stop or pause your treatment with Inflectra and may give you medicines to treat the allergic reaction. Signs of an allergic reaction can include: hives (red, raised, itchy patches of skin) difficulty breathing chest pain high or low blood pressure fever chills Some patients treated with infliximab products have had delayed allergic reactions. The delayed reactions occurred 3 to 12 days after receiving treatment with an infliximab product. Tell your doctor right away if you have any of these signs of delayed allergic reaction to Inflectra: fever rash headache sore throat muscle or joint pain swelling of the face and hands difficulty swallowing

Lupus-like Syndrome Some patients have developed symptoms that are like the symptoms of Lupus. If you develop any of the following symptoms, your doctor may decide to stop your treatment with Inflectra. chest discomfort or pain that does not go away shortness of breath joint pain rash on the cheeks or arms that gets worse in the sun

Psoriasis Some people receiving infliximab products had new psoriasis or worsening of psoriasis they already had. Tell your doctor if you develop red scaly patches or raised bumps on the skin that are filled with pus. Your doctor may decide to stop your treatment with Inflectra. The most common side effects of infliximab products include: respiratory infections, such as sinus infections and sore throat headache coughing stomach pain

Infusion reactions can happen up to 2 hours after your infusion of Inflectra.

Drug Side Effects Summary

Symptoms of infusion reactions may include: fever chills chest pain low blood pressure or high blood pressure shortness of breath rash itching Children with Crohn's disease showed some differences in side effects of treatment compared with adults with Crohn's disease. The side effects that happened more in children were: anemia (low red blood cells), leukopenia (low white blood cells), flushing (redness or blushing), viral infections, neutropenia (low neutrophils, the white blood cells that fight infection), bone fracture, bacterial infection and allergic reactions of the breathing tract. Among patients who took infliximab for ulcerative colitis in clinical studies, more children had infections as compared with adults. Tell your doctor about any side effect that bothers you or does not go away. These are not all of the side effects with Inflectra. Ask your doctor or pharmacist for more information. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA1088. Inflectra side effects

2671. This particular disease, Psoriasis, interacts with the drug ixekizumab and has the following side effects:

- Ixekizumab may cause serious side effects, including: See "Important information" above Serious allergic reactions. If you have a severe allergic reaction, do not give another injection of ixekizumab. feel faint swelling of your face, eyelids, lips, mouth, tongue, or throat trouble breathing or throat tightness chest tightness skin rash Crohn's disease or ulcerative colitis (inflammatory bowel disease) can happen during treatment with ixekizumab, including worsening symptoms. Tell your healthcare provider if you have new or worsening symptoms of inflammatory bowel disease during treatment with ixekizumab, including: stomach (abdomen) pain diarrhea with or without blood weight loss The most common side effects of ixekizumab in adults and children include: injection site reactions upper respiratory infections nausea fungal infections These are not all of the possible side effects of ixekizumab. Tell your healthcare provider about any side effect that bothers you or that does not go away. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Ixekizumab side effects

Drug Side Effects Summary

2672. This particular disease, Psoriasis, interacts with the drug Renflexis and has the following side effects:

- Serious side effects of Renflexis include: See "Important information" above. Serious Infections Some patients, especially those 65 years and older have had serious infections while receiving infliximab products, such as Renflexis. These serious infections include TB and infections caused by viruses, fungi, or bacteria that have spread throughout the body. Some patients die from these infections. If you get an infection while receiving treatment with Renflexis your doctor will treat your infection and may need to stop your Renflexis treatment. Tell your doctor right away if you have any of the following signs of an infection while receiving or after receiving Renflexis: a fever feel very tired have a cough have flu-like symptoms warm, red, or painful skin Your doctor will examine you for TB and perform a test to see if you have TB. If your doctor feels that you are at risk for TB, you may be treated with medicine for TB before you begin treatment with Renflexis and during treatment with Renflexis. Even if your TB test is negative, your doctor should carefully monitor you for TB infections while you are receiving Renflexis. Patients who had a negative TB skin test before receiving infliximab products have developed active TB. If you are a chronic carrier of the hepatitis B virus, the virus can become active while you are being treated with Renflexis. In some cases, patients have died as a result of hepatitis B virus being reactivated. Your doctor should do a blood test for hepatitis B virus before you start treatment with Renflexis and occasionally while you are being treated. Tell your doctor if you have any of the following symptoms: feel unwell tiredness (fatigue) poor appetite fever, skin rash, or joint pain Heart Failure If you have a heart problem called congestive heart failure, your doctor should check you closely while you are receiving Renflexis. Your congestive heart failure may get worse while you are receiving Renflexis. Be sure to tell your doctor of any new or worse symptoms including: shortness of breath swelling of ankles or feet sudden weight gain Treatment with Renflexis may need to be stopped if you get new or worse congestive heart failure. Other Heart Problems Some patients have experienced a heart attack (some of which led to death), low blood flow to the heart, or abnormal heart rhythm within 24 hours

Drug Side Effects Summary

of beginning their infusion of an infliximab product. Symptoms may include chest discomfort or pain, arm pain, stomach pain, shortness of breath, anxiety, lightheadedness, dizziness, fainting, sweating, nausea, vomiting, fluttering or pounding in your chest, and/or a fast or a slow heartbeat. Tell your doctor right away if you have any of these symptoms.

Liver Injury Some patients receiving infliximab products have developed serious liver problems. Tell your doctor if you have: jaundice (skin and eyes turning yellow) dark brown-colored urine pain on the right side of your stomach area (right-sided abdominal pain) fever extreme tiredness (severe fatigue)

Blood Problems In some patients receiving infliximab products, the body may not make enough of the blood cells that help fight infections or help stop bleeding. Tell your doctor if you: have a fever that does not go away bruise or bleed very easily look very pale

Nervous System Disorders Some patients receiving infliximab products have developed problems with their nervous system. Tell your doctor if you have: changes in your vision numbness or tingling in any part of your body weakness in your arms or legs seizures

Some patients have experienced a stroke within approximately 24 hours of their infusion with infliximab products. Tell your doctor right away if you have symptoms of a stroke which may include: numbness or weakness of the face, arm or leg, especially on one side of the body

- sudden confusion, trouble speaking or understanding

- sudden trouble seeing in one or both eyes, sudden trouble walking, dizziness, loss of balance or coordination or a sudden, severe headache.

Allergic Reactions Some patients have had allergic reactions to infliximab products. Some of these reactions were severe. These reactions can happen while you are getting your Renflexis treatment or shortly afterward. Your doctor may need to stop or pause your treatment with Renflexis and may give you medicines to treat the allergic reaction. Signs of an allergic reaction can include: hives (red, raised, itchy patches of skin) difficulty breathing chest pain high or low blood pressure fever chills

Some patients treated with infliximab products have had delayed allergic reactions. The delayed reactions occurred 3 to 12 days after receiving treatment with an infliximab product. Tell your doctor right away if you have any of these signs of delayed

Drug Side Effects Summary

allergic reaction to Renflexis: fever rash headache sore throat muscle or joint pain swelling of the face and hands difficulty swallowing Lupus-like Syndrome Some patients have developed symptoms that are like the symptoms of Lupus. If you develop any of the following symptoms, your doctor may decide to stop your treatment with Renflexis. chest discomfort or pain that does not go away shortness of breath joint pain rash on the cheeks or arms that gets worse in the sun Psoriasis Some people receiving infliximab products had new psoriasis or worsening of psoriasis they already had. Tell your doctor if you develop red scaly patches or raised bumps on the skin that are filled with pus. Your doctor may decide to stop your treatment with Renflexis. The most common side effects of infliximab products include: respiratory infections, such as sinus infections and sore throat headache coughing stomach pain Infusion reactions can happen up to 2 hours after your infusion of Renflexis. Symptoms of infusion reactions may include: fever chills chest pain low blood pressure or high blood pressure shortness of breath rash itching Children with Crohn's disease showed some differences in side effects of treatment compared with adults with Crohn's disease. The side effects that happened more in children were: anemia (low red blood cells), leukopenia (low white blood cells), flushing (redness or blushing), viral infections, neutropenia (low neutrophils, the white blood cells that fight infection), bone fracture, bacterial infection and allergic reactions of the breathing tract. Among patients who took infliximab for ulcerative colitis in clinical studies, more children had infections as compared with adults. Tell your doctor about any side effect that bothers you or does not go away. These are not all of the side effects with Renflexis. Ask your doctor or pharmacist for more information. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA1088. Renflexis side effects

2673. This particular disease, Psoriasis, interacts with the drug risankizumab and has the following

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Call your doctor right away if you have signs of infection such as: fever, chills, sweating, body aches
- shortness of breath, cough, bloody mucus
- mouth sores, red or swollen gums
- stomach pain, diarrhea
- increased urination, burning when you urinate
- pale skin, easy bruising, unusual bleeding
- a fungal skin infection --skin sores different from psoriasis, rash or redness, blisters, itching, burning, cracking or peeling, changes in skin color
- or signs of tuberculosis: fever, cough, night sweats , loss of appetite, weight loss, and feeling very tired. Further doses may be delayed until your infection clears up. Common side effects of risankizumab may include: headache
- tiredness
- cold symptoms such as stuffy nose, sneezing, sore throat
- fungal skin infections
- or pain, redness, itching, bruising, swelling, warmth, bleeding, burning, or other skin irritation where the medicine was injected.

2674. This particular disease, Psoriasis, interacts with the drug secukinumab and has the following

- Secukinumab may cause serious side effects including: See "important information" above on this page. Inflammatory bowel disease. New cases of inflammatory bowel disease or "flare-ups" can happen with secukinumab, and can sometimes be serious. If you have inflammatory bowel disease (ulcerative colitis or Crohn's disease), tell your healthcare provider if you have worsening disease symptoms during treatment with secukinumab or develop new symptoms of stomach pain or diarrhea. Serious allergic reactions. swelling of your face, eyelids, lips, mouth, tongue, or throat feel

Drug Side Effects Summary

faint chest tightness trouble breathing or throat tightness hives (red, itchy bumps) skin rash If you have a severe allergic reaction, you should not have another injection of secukinumab. The most common side effects of secukinumab include: upper respiratory infections cold symptoms diarrhea These are not all of the possible side effects of secukinumab. Call your healthcare provider for medical advice and about side effects. You can report side effects to FDA at 1-800-FDA-1088.

Secukinumab side effects

2675. This particular disease, Psoriasis, interacts with the drug tildrakizumab and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. You may get infections more easily, even serious or fatal infections. Call your doctor right away if you have signs of infection such as: fever, chills, sweating
- skin sores
- muscle pain
- increased urination, pain or burning when you urinate
- stomach pain, diarrhea, weight loss
- or cough, shortness of breath, coughing up pink or red mucus. Common side effects of tildrakizumab may include: pain, itching, rash, redness, swelling, bruising, or bleeding where the medicine was injected
- diarrhea
- or cold symptoms such as stuffy nose, sneezing, sore throat .

2676. This particular disease, Psoriasis, interacts with the drug calcipotriene and has the following

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Calcipotriene topical may cause serious side effects. Stop using calcipotriene topical and call your doctor at once if you have: severe burning, stinging, skin rash , or other irritation after applying the medicine
- worsening of your skin condition
- or high calcium levels--confusion, tiredness, nausea , vomiting , loss of appetite, constipation , increased thirst or urination, weight loss . Common side effects of calcipotriene topical may include: mild skin irritation
- skin rash
- or itching.

2677. This particular disease, Psoriasis, interacts with the drug Dovonex and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Dovonex and call your doctor at once if you have: severe burning, stinging, skin rash, or other irritation after applying the medicine
- worsening of your skin condition
- or high calcium levels - confusion, tiredness, nausea, vomiting, loss of appetite, constipation, increased thirst or urination, weight loss. Common Dovonex side effects may include: mild skin irritation
- skin rash
- or itching.

2678. This particular disease, Psoriasis, interacts with the drug acitretin and has the following side effects:

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Acitretin may cause serious side effects. Stop using acitretin and call your doctor at once if you have: mood changes--depression, aggression, unusual thoughts or behavior, thoughts of hurting yourself
- heart attack or stroke symptoms--chest pain, dizziness, nausea, feeling short of breath, sudden numbness or weakness (especially on one side of the body), sudden severe headache , problems with speech or balance, swelling or warmth in one or both legs
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor, headache, blurred vision
- increased pressure inside the skull--severe headaches, ringing in your ears, dizziness, nausea, vision problems, pain behind your eyes
- liver problems--nausea, vomiting, loss of appetite, dark urine, or jaundice (yellowing of your skin or eyes)
- problems with your bones or muscles--loss of feeling in your hands or feet, trouble moving, pain in your back, joints, muscles, or bones
- serious skin problems--itching, redness, pain, swelling or peeling of your skin
- or signs of a blood vessel problem--sudden swelling, rapid weight gain, fever, muscle pain, feeling light-headed. Common side effects of acitretin may include: chapped lips, dry mouth
- itchy or scaly skin
- weak nails, fragile skin
- peeling skin on your hands and feet
- hair loss
- dry eyes, discomfort while wearing contact lenses
- dry or runny nose , nosebleeds
- or joint pain, tight muscles.

Drug Side Effects Summary

2679. This particular disease, Psoriasis, interacts with the drug betamethasone and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using betamethasone and call your doctor at once if you have: severe skin irritation where the medicine was applied
- or signs of skin infection (swelling, redness, warmth, oozing). Your skin can absorb topical steroid medicine, which may cause steroid side effects throughout the body. Stop using betamethasone and call your doctor if you have: blurred vision, tunnel vision, eye pain , or seeing halos around lights
- slow wound healing, thinning skin, increased body hair
- increased thirst or urination, dry mouth, fruity breath odor
- weight gain, puffiness in your face
- or muscle weakness, tired feeling, depression , anxiety , feeling irritable. Steroids can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common betamethasone side effects may include: itching, redness, burning, stinging, or blistering of treated skin
- skin bruising or shiny appearance
- or folliculitis (redness or crusting around your hair follicles).

2680. This particular disease, Psoriasis, interacts with the drug Soriatane and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Soriatane may cause serious side effects. Stop using Soriatane and call your doctor at once if you have: mood changes--depression, aggression, unusual thoughts or behavior, thoughts of hurting yourself

Drug Side Effects Summary

- heart attack or stroke symptoms--chest pain, dizziness, nausea, feeling short of breath, sudden numbness or weakness (especially on one side of the body), sudden severe headache , problems with speech or balance, swelling or warmth in one or both legs
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor, headache, blurred vision
- increased pressure inside the skull--severe headaches, ringing in your ears, dizziness, nausea, vision problems, pain behind your eyes
- liver problems--nausea, vomiting, loss of appetite, dark urine, or jaundice (yellowing of your skin or eyes)
- problems with your bones or muscles--loss of feeling in your hands or feet, trouble moving, pain in your back, joints, muscles, or bones
- serious skin problems--itching, redness, pain, swelling or peeling of your skin
- or signs of a blood vessel problem--sudden swelling, rapid weight gain, fever, muscle pain, feeling light-headed. Common side effects of Soriatane may include: chapped lips, dry mouth
- itchy or scaly skin
- weak nails, fragile skin
- peeling skin on your hands and feet
- hair loss
- dry eyes, discomfort while wearing contact lenses
- dry or runny nose , nosebleeds
- or joint pain, tight muscles.

2681. This particular disease, Psoriasis, interacts with the drug Taclonex and has the following side

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high calcium levels - confusion, tiredness, nausea , vomiting , loss of appetite, constipation , increased thirst or urination, weight loss
- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin - weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea, diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Common Taclonex side effects may include: skin color changes
- worsening of psoriasis
- skin irritation, rash, itching, redness, or hives
- redness or crusting around your hair follicles
- or headache .

2682. This particular disease, Psoriasis, interacts with the drug betamethasone / calcipotriene and I

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Betamethasone and calcipotriene topical may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high calcium levels--confusion, tiredness, nausea , vomiting , loss of appetite, constipation , increased thirst or urination, weight loss

Drug Side Effects Summary

- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea, diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Common side effects of betamethasone and calcipotriene topical may include: skin color changes
- worsening of psoriasis
- skin irritation, rash, itching, redness, or hives
- redness or crusting around your hair follicles
- or headache .

2683. This particular disease, Psoriasis, interacts with the drug Neoral and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Abdominal or stomach pain or tenderness back pain black, tarry stools blurred vision chest pain chills clay colored stools cloudy urine cough dark urine decrease in urine output or decrease in urine-concentrating ability decreased appetite dizziness drowsiness fever headache headache, severe and throbbing itching loss of appetite muscle spasms (tetany) or twitching nausea and vomiting nervousness painful or difficult urination pounding in the ears shakiness in the legs, arms, hands, or feet shortness of breath skin rash slow or fast heartbeat sore throat sores, ulcers, or white spots on the lips or in the mouth swelling of the feet or lower legs swollen glands trembling or shaking of the hands or feet unusual bleeding or bruising unusual tiredness or weakness yellow eyes or skin Less common Bleeding gums blood in the urine blood in the vomit burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings convulsions difficulty swallowing hives pale skin pinpoint red

Drug Side Effects Summary

spots on the skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue severe or continuing stomach pain tightness in the chest troubled breathing with exertion Rare Bloating chest discomfort constipation darkened urine hoarseness indigestion lower back or side pain night sweats pain or discomfort in the arms, jaw, back, or neck pains in the stomach, side, or abdomen, possibly radiating to the back sweating vomiting of blood or material that looks like coffee grounds Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Abdominal or stomach discomfort bleeding, tender, or enlarged gums blemishes on the skin increased hair growth, especially on the face pain or tenderness around the eyes and cheekbones pimples stuffy or runny nose Less common Brittle fingernails burning feeling in the chest or stomach burning, dry, or itching eyes continuing ringing or buzzing or other unexplained noise in the ears cramps discharge or excessive tearing feeling of warmth hearing loss redness of the face, neck, arms, and occasionally, upper chest redness, pain, swelling of the eye, eyelid, or inner lining of the eyelid swelling of the breasts or breast soreness in both females and males weight loss Rare Blurred or loss of vision discouragement disturbed color perception double vision fear feeling sad or empty halos around lights irritability joint pain loss of interest or pleasure night blindness overbright appearance of lights tiredness trouble concentrating trouble sleeping tunnel vision unusual drowsiness, dullness, tiredness, weakness, or feeling of sluggishness weakness weight loss Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Neoral side effects

Drug Side Effects Summary

2684. This particular disease, Psoriasis, interacts with the drug Otrexup and has the following side

- sudden chest pain, wheezing, dry cough, cough with mucus, feeling short of breath
- swollen lymph glands, night sweats , weight loss
- blisters or ulcers in your mouth, red or swollen gums, trouble swallowing
- vomiting , diarrhea, blood in your urine or stools
- skin changes such as redness, warmth, swelling, or oozing
- signs of metabolic acidosis --confusion, vomiting, lack of energy, irregular heartbeats
- low blood cell counts--fever, chills, bruising or bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath
- kidney problems--little or no urination, swelling in your feet or ankles
- liver problems--swelling around your midsection, right-sided upper stomach pain, nausea , loss of appetite, dark urine, jaundice (yellowing of the skin or eyes)
- nerve problems--confusion, weakness, drowsiness, coordination problems, feeling irritable, headache , neck stiffness, vision problems, loss of movement in any part of your body, seizure
- or signs of tumor cell breakdown--tiredness, muscle cramps, nausea, vomiting, diarrhea, fast or slow heart rate , tingling in your hands and feet or around your mouth. Common side effects of Otrexup may include: low blood cell counts, fever, chills, tiredness, not feeling well
- headache, dizziness , cough, chest tightness, trouble breathing
- mouth sores, nausea, vomiting, stomach pain, indigestion , diarrhea
- abnormal liver function tests
- runny or stuffy nose, sore throat
- or rash, hair loss , burning skin lesions, being more sensitive to light.

2685. This particular disease, Psoriasis, interacts with the drug Trexall and has the following side e

- sudden chest pain, wheezing, dry cough, cough with mucus, chest pain, feeling short of breath

Drug Side Effects Summary

- fever, chills, swollen lymph glands, night sweats , weight loss
- blisters or ulcers in your mouth, red or swollen gums, trouble swallowing
- vomiting, diarrhea, blood in your urine or stools
- skin changes such as redness, warmth, swelling, or oozing
- low blood cell counts--fever, chills, tiredness, mouth sores, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath
- kidney problems--little or no urination, swelling in your feet or ankles
- liver problems--swelling around your midsection, right-sided upper stomach pain, nausea , loss of appetite, dark urine, jaundice (yellowing of the skin or eyes)
- nerve problems--confusion, weakness, drowsiness, coordination problems, feeling irritable, headache , neck stiffness, vision problems, loss of movement in any part of your body, seizure
- or signs of tumor cell breakdown--tiredness, weakness, muscle cramps, nausea, vomiting, diarrhea, fast or slow heart rate , tingling in your hands and feet or around your mouth. Common side effects of Trexall may include: fever, chills, tiredness, not feeling well
- low blood cell counts
- mouth sores
- nausea, stomach pain
- abnormal liver function tests
- hair loss
- burning skin lesions
- or being more sensitive to light.

2686. This particular disease, Psoriasis, interacts with the drug Enstilar and has the following side

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Enstilar may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high calcium levels--confusion, tiredness, nausea , vomiting , loss of appetite, constipation , increased thirst or urination, weight loss
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea, diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Common side effects of Enstilar may include: skin color changes
- worsening of psoriasis
- skin irritation, rash, itching, redness, or hives
- redness or crusting around your hair follicles
- or headache .

2687. This particular disease, Psoriasis, interacts with the drug Rasuvo and has the following side effects

- sudden chest pain, wheezing, dry cough, cough with mucus, feeling short of breath
- swollen lymph glands, night sweats , weight loss
- blisters or ulcers in your mouth, red or swollen gums, trouble swallowing
- vomiting , diarrhea, blood in your urine or stools
- skin changes such as redness, warmth, swelling, or oozing
- signs of metabolic acidosis --confusion, vomiting, lack of energy, irregular heartbeats
- low blood cell counts--fever, chills, bruising or bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath

Drug Side Effects Summary

- kidney problems--little or no urination, swelling in your feet or ankles
- liver problems--swelling around your midsection, right-sided upper stomach pain, nausea , loss of appetite, dark urine, jaundice (yellowing of the skin or eyes)
- nerve problems--confusion, weakness, drowsiness, coordination problems, feeling irritable, headache , neck stiffness, vision problems, loss of movement in any part of your body, seizure
- or signs of tumor cell breakdown--tiredness, muscle cramps, nausea, vomiting, diarrhea, fast or slow heart rate , tingling in your hands and feet or around your mouth. Common side effects of Rasuvo may include: low blood cell counts, fever, chills, tiredness, not feeling well
- headache, dizziness , cough, chest tightness, trouble breathing
- mouth sores, nausea, vomiting, stomach pain, indigestion , diarrhea
- abnormal liver function tests
- runny or stuffy nose, sore throat
- or rash, hair loss , burning skin lesions, being more sensitive to light.

2688. This particular disease, Psoriasis, interacts with the drug tacrolimus and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tacrolimus topical may cause serious side effects. Call your doctor at once if you have: severe pain, burning, itching, or stinging
- new or worsened skin symptoms (warts , unusual rash or skin lesions, blistering or oozing)
- a mole that has changed in size or color
- or symptoms of a viral skin infection--cold sores around your mouth, skin sores or blisters, scabs, itching, tingling. Common side effects of tacrolimus topical may include: burning, stinging, itching, or tingling of treated skin (especially during the first few days of treatment)
- viral skin infection

Drug Side Effects Summary

- your skin being more sensitive to hot or cold temperatures
- acne , redness or crusting around your hair follicles
- upset stomach
- headache , muscle pain
- runny or stuffy nose
- or fever, chills, body aches, sore throat , cough, swollen glands.

2689. This particular disease, Psoriasis, interacts with the drug calcitriol and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: blistering, or severe redness, itching, or other irritation of treated areas
- or worsened symptoms or no improvement in psoriasis. Using calcitriol topical can affect your body's ability to metabolize calcium. Tell your doctor right away if you have symptoms of high levels of calcium in your blood (hypercalcemia), including: nausea , vomiting , loss of appetite, constipation
- increased thirst or urination
- lack of energy, feeling tired
- muscle weakness
- confusion
- or bone pain. Common side effects of calcitriol may include: high levels of calcium in your blood or urine
- itching, skin irritation
- or abnormal lab tests.

Drug Side Effects Summary

2690. This particular disease, Psoriasis, interacts with the drug coal tar and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. coal tar topical may cause serious side effects. Stop using coal tar topical and call your doctor at once if you have: severe stinging, burning, swelling, or other irritation of the treated skin. Common side effects may include mild skin irritation or skin rash .

2691. This particular disease, Psoriasis, interacts with the drug prednisolone and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Prednisolone may cause serious side effects. Call your doctor at once if you have: shortness of breath (even with mild exertion), swelling, rapid weight gain
- bruising, thinning skin, or any wound that will not heal
- severe depression, changes in personality, unusual thoughts or behavior
- new or unusual pain in an arm or leg or in your back
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- severe pain in your upper stomach spreading to your back, nausea and vomiting
- a seizure (convulsions)
- or low potassium --leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling. Steroids can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using prednisolone. Common side effects of prednisolone may include: fluid retention (swelling in your hands or ankles)
- dizziness , spinning sensation
- changes in your menstrual periods

Drug Side Effects Summary

- headache
- muscle pain or weakness
- or stomach discomfort, bloating .

2692. This particular disease, Psoriasis, interacts with the drug RediTrex and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: More common Black, tarry stools blood in the urine or stools bloody vomit diarrhea joint pain reddening of the skin sores in the mouth or lips stomach pain swelling of the feet or lower legs Less common Back pain bleeding, blistering, burning, coldness, discoloration of the skin, feeling of pressure, hives, infection, inflammation, itching, lumps, numbness, pain, rash, redness, scarring, soreness, stinging, swelling, tenderness, tingling, or warmth at the injection site blurred vision confusion convulsions (seizures) cough or hoarseness dark urine dizziness drowsiness fever or chills headache lower back or side pain painful or difficult urination pinpoint red spots on the skin shortness of breath unusual bleeding or bruising unusual tiredness or weakness yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Hair loss, temporary loss of appetite nausea or vomiting Less common Acne boils on skin pale skin skin rash or itching Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. RediTrex side effects

Drug Side Effects Summary

2693. This particular disease, Psoriasis, interacts with the drug Taclonex Scalp and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Burning, itching, and pain in hairy areas burning sensation of the skin flushing or redness of the skin itching skin pus at the root of the hair red, scaling, or crusted skin scaly rash skin irritation thinning of the skin with easy bruising unusually warm skin Rare Body aches or pain chills cough darkening of the skin diarrhea difficulty with breathing ear congestion fever general feeling of discomfort or illness headache joint pain lightening of normal skin color lightening of the treated areas of dark skin loss of appetite loss of voice muscle aches and pains nausea raised, dark red, or wart-like spots on the skin red, sore eyes shivering sneezing sore throat spots on your skin that look like a blister or pimple stuffy or runny nose sweating trouble sleeping unusual tiredness or weakness vomiting worsening of psoriasis Incidence not known Blindness blurred vision burning and itching of the skin with pinhead-sized red blisters decreased vision eye pain tearing Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Bruising large, flat, blue, or purplish patches in the skin pain Rare Blemishes on the skin pimples Incidence not known Dryness of the skin reddish purple lines on the arms, face, legs, trunk, or groin redness and scaling around the mouth Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Taclonex Scalp side effects

Drug Side Effects Summary

2694. This particular disease, Psoriasis, interacts with the drug Vectical and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have: blistering, or severe redness, itching, or other irritation of treated areas
- or worsened symptoms or no improvement in psoriasis. Using Vectical can affect your body's ability to metabolize calcium. Tell your doctor right away if you have symptoms of high levels of calcium in your blood (hypercalcemia), including: nausea , vomiting , loss of appetite, constipation
- increased thirst or urination
- lack of energy, feeling tired
- muscle weakness
- confusion
- or bone pain. Common Vectical side effects may include: calcium in your urine
- itching
- or abnormal lab tests.

2695. This particular disease, Psoriasis, interacts with the drug Gengraf and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Abdominal or stomach pain or tenderness back pain black, tarry stools blurred vision chest pain chills clay colored stools cloudy urine cough dark urine decrease in urine output or decrease in urine-concentrating ability decreased appetite dizziness drowsiness fever headache headache, severe and throbbing itching loss of appetite muscle spasms (tetany) or twitching nausea and vomiting nervousness painful or difficult urination pounding in the ears shakiness in the legs, arms, hands, or feet shortness of

Drug Side Effects Summary

breath skin rash slow or fast heartbeat sore throat sores, ulcers, or white spots on the lips or in the mouth swelling of the feet or lower legs swollen glands trembling or shaking of the hands or feet unusual bleeding or bruising unusual tiredness or weakness yellow eyes or skin Less common Bleeding gums blood in the urine blood in the vomit burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings convulsions difficulty swallowing hives pale skin pinpoint red spots on the skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue severe or continuing stomach pain tightness in the chest troubled breathing with exertion Rare Bloating chest discomfort constipation darkened urine hoarseness indigestion lower back or side pain night sweats pain or discomfort in the arms, jaw, back, or neck pains in the stomach, side, or abdomen, possibly radiating to the back sweating vomiting of blood or material that looks like coffee grounds Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Abdominal or stomach discomfort bleeding, tender, or enlarged gums blemishes on the skin increased hair growth, especially on the face pain or tenderness around the eyes and cheekbones pimples stuffy or runny nose Less common Brittle fingernails burning feeling in the chest or stomach burning, dry, or itching eyes continuing ringing or buzzing or other unexplained noise in the ears cramps discharge or excessive tearing feeling of warmth hearing loss redness of the face, neck, arms, and occasionally, upper chest redness, pain, swelling of the eye, eyelid, or inner lining of the eyelid swelling of the breasts or breast soreness in both females and males weight loss Rare Blurred or loss of vision discouragement disturbed color perception double vision fear feeling sad or empty halos around lights irritability joint pain loss of interest or pleasure night blindness overbright appearance of lights tiredness trouble concentrating trouble sleeping tunnel vision unusual drowsiness, dullness, tiredness, weakness, or feeling of

Drug Side Effects Summary

sluggishness weakness weight loss Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Gengraf side effects

2696. This particular disease, Psoriasis, interacts with the drug Oxсорalen-Ultra and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Oxсорalen-Ultra may cause serious side effects. Call your doctor at once if you have: severe skin redness within 24 hours after UVA treatment
- swelling, severe itching, or severe skin discomfort
- skin redness and swelling with blisters
- worsening of your psoriasis
- a new skin lesion, or a mole that has changed in size or color
- or blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights. Mild skin redness may occur and can last for 1 or 2 days. This is a normal side effect after UVA treatment and may not be a sign of severe sunburn. Ask your doctor if you have concerns about any redness or swelling. Common side effects of Oxсорalen-Ultra may include: itching, redness
- nausea
- feeling nervous
- or sleep problems (insomnia).

2697. This particular disease, Psoriasis, interacts with the drug Psoriasin and has the following side effects:

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Psoriasin may cause serious side effects. Stop using Psoriasin and call your doctor at once if you have: severe stinging, burning, swelling, or other irritation of the treated skin. Common side effects may include mild skin irritation or skin rash .

2698. This particular disease, Psoriasis, interacts with the drug Sernivo and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Betamethasone may cause serious side effects. Stop using betamethasone and call your doctor at once if you have: severe skin irritation where the medicine was applied
- or signs of skin infection (swelling, redness, warmth, oozing). Your skin can absorb topical steroid medicine, which may cause steroid side effects throughout the body. Stop using betamethasone and call your doctor if you have: blurred vision, tunnel vision, eye pain , or seeing halos around lights
- slow wound healing, thinning skin, increased body hair
- increased thirst or urination, dry mouth, fruity breath odor
- weight gain, puffiness in your face
- or muscle weakness, tired feeling, depression , anxiety , feeling irritable. Steroids can affect growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common side effects of Sernivo may include: itching, redness, burning, stinging, or blistering of treated skin
- skin bruising or shiny appearance
- or folliculitis (redness or crusting around your hair follicles).

2699. This particular disease, Psoriasis, interacts with the drug Abrilada and has the following side effects:

Drug Side Effects Summary

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Bladder pain bloody or cloudy urine body aches or pain chest pain or tightness chills cough difficult, burning, or painful urination ear congestion or pain fever frequent urge to urinate head congestion headache hoarseness or other voice changes lower back or side pain pain or tenderness around the eyes and cheekbones sneezing sore throat stuffy or runny nose trouble breathing Less common Black, tarry stools blindness blue or pale skin blurred vision bone fractures confusion difficulty with moving dizziness fast, slow, or irregular heartbeat or pulse increased thirst increased urination itching, skin rash indigestion light-colored stools loss of appetite loss of strength or energy muscle pain or stiffness nervousness pain in the arms or legs pounding in the ears seizures slow or fast heartbeat sores, ulcers, or white spots on the lips or in the mouth trouble thinking unpleasant breath odor unusual bleeding or bruising vomiting vomiting of blood yellow eyes or skin Incidence not known Blistering, peeling, or loosening of the skin decreased urine output diarrhea dilated neck veins irregular breathing irregular heartbeat itching numbness or tingling in the fingers, face, or feet red, irritated eyes red skin lesions, often with a purple center sores, ulcers, or white spots in the mouth or on the lips swelling of the face, fingers, feet, or lower legs unusual tiredness or weakness weight gain Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Back pain bleeding, blistering, burning, coldness, discoloration of the skin, feeling of pressure, hives, infection, inflammation, itching, lumps, numbness, pain, rash, redness, scarring, soreness, stinging, swelling, tenderness, tingling, ulceration, or warmth at the injection site Incidence not known Hair loss or thinning of the

Drug Side Effects Summary

hair Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Abrilada side effects

2700. This particular disease, Psoriasis, interacts with the drug Amjevita and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Bladder pain body aches or pain bloody or cloudy urine chest tightness chills cough difficult, burning, or painful urination ear congestion fever frequent urge to urinate headache hoarseness or other voice changes lower back or side pain pain or tenderness around the eyes and cheekbones sneezing sore throat stuffy or runny nose sweating trouble breathing unusual tiredness or weakness Less common Black, tarry stools blindness blue or pale skin bone fractures confusion dark urine decreased vision difficulty with moving dizziness eye pain fainting fast, slow, or irregular heartbeat or pulse hoarseness increased thirst increased urination itching, skin rash indigestion light-colored stools loss of appetite loss of strength or energy muscle cramps, pain, spasms, or stiffness nausea nervousness pain in the arms or legs pounding in the ears seizures slow or fast heartbeat sores, ulcers, or white spots on the lips or in the mouth stomach pain trouble thinking unpleasant breath odor unusual bleeding or bruising vomiting vomiting of blood yellow eyes or skin Incidence not known Blistering, peeling, or loosening of the skin decreased urine output dilated neck veins diarrhea irregular breathing irregular heartbeat joint pain numbness or tingling in the fingers, face, or feet red, irritated eyes red skin lesions, often with a purple center swelling of the face, fingers, feet, or lower legs weight gain Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care

Drug Side Effects Summary

professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Back pain bleeding, blistering, burning, coldness, discoloration of the skin, feeling of pressure, hives, infection, inflammation, itching, lumps, numbness, pain, rash, redness, scarring, soreness, stinging, swelling, tenderness, tingling, ulceration, or warmth at the injection site Incidence not known Hair loss or thinning of the hair Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Amjevita side effects

2701. This particular disease, Psoriasis, interacts with the drug Analpram-HC and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of a weak adrenal gland like a very bad upset stomach or throwing up, very bad dizziness or passing out, muscle weakness, feeling very tired, mood changes, not hungry, or weight loss . Signs of Cushing's disease like weight gain in the upper back or belly, moon face, very bad headache , or slow healing. Signs of high blood sugar like

Drug Side Effects Summary

confusion, feeling sleepy, more thirst, more hungry, passing urine more often, flushing, fast breathing, or breath that smells like fruit. Irritation where Analpram HC (pramoxine and hydrocortisone rectal cream) is used. Very bad bleeding from rectum or rectal pain. Analpram-HC side effects

2702. This particular disease, Psoriasis, interacts with the drug Analpram E and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of a weak adrenal gland like a very bad upset stomach or throwing up, very bad dizziness or passing out, muscle weakness, feeling very tired, mood changes, not hungry, or weight loss . Signs of Cushing's disease like weight gain in the upper back or belly, moon face, very bad headache , or slow healing. Signs of high blood sugar like confusion, feeling sleepy, more thirst, more hungry, passing urine more often, flushing, fast breathing, or breath that smells like fruit. Irritation where Analpram E (pramoxine and hydrocortisone rectal cream) is used. Very bad bleeding from rectum or rectal pain. Analpram E side effects

Drug Side Effects Summary

2703. This particular disease, Psoriasis, interacts with the drug Anthraforte and has the following side effects

- Anthralin has been shown to cause tumors in animals. However, there have been no reports of anthralin causing tumors in humans. Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: More common Redness or other skin irritation of treated or uninvolved skin not present before use of this medicine Rare Skin rash Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Anthraforte side effects

2704. This particular disease, Psoriasis, interacts with the drug anthralin and has the following side effects

- Anthralin has been shown to cause tumors in animals. However, there have been no reports of anthralin causing tumors in humans. Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: More common Redness or other skin irritation of treated or uninvolved skin not present before use of anthralin Rare Skin rash Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Anthralin topical side effects

2705. This particular disease, Psoriasis, interacts with the drug Anthrascalp and has the following side effects

- Anthralin has been shown to cause tumors in animals. However, there have been no reports of anthralin causing tumors in humans. Along with its needed effects, a medicine may cause some

Drug Side Effects Summary

unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: More common Redness or other skin irritation of treated or uninvolved skin not present before use of this medicine Rare Skin rash Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Anthrascalp side effects

2706. This particular disease, Psoriasis, interacts with the drug Balnetar and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Balnetar may cause serious side effects. Stop using Balnetar and call your doctor at once if you have: severe stinging, burning, swelling, or other irritation of the treated skin. Common side effects may include mild skin irritation or skin rash .

2707. This particular disease, Psoriasis, interacts with the drug Betatar Gel and has the following side

- In animal studies, coal tar has been shown to increase the chance of skin cancer. Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: Rare Skin irritation not present before use of this medicine skin rash Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Stinging (mild)--especially for

Drug Side Effects Summary

gel and solution dosage forms Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Betatar Gel side effects

2708. This particular disease, Psoriasis, interacts with the drug brodalumab and has the following s

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Brodalumab comes with a Medication Guide and a Patient Wallet Card listing symptoms of a serious side effect. Read this information and learn what symptoms to watch for. Keep the Wallet Card with you at all times. Brodalumab may cause serious side effects. Call your doctor at once if you have: fever, chills, sweating, tiredness
- sore throat , cough, trouble breathing or swallowing
- skin sores with warmth, redness, or swelling
- stomach cramps, loss of appetite, constipation
- painful diarrhea, bloody stools, weight loss
- increased urination, pain or burning when you urinate
- unusual changes in mood or behavior
- suicidal thoughts
- or new or worsening anxiety or depression. Common side effects of brodalumab may include:
fever, chills, flu-like symptoms
- itching, skin redness
- muscle or joint pain
- nausea , diarrhea
- headache , tiredness

Drug Side Effects Summary

- mouth pain
- or pain, redness, bruising, itching, or bleeding where the medicine was injected.

2709. This particular disease, Psoriasis, interacts with the drug Bryhali and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Bryhali may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea , diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Common side effects of Bryhali may include: burning, stinging, itching, or dryness of treated skin
- pain where the foam was applied
- redness or crusting around your hair follicles
- stretch marks
- spider veins
- headache
- or cold symptoms such as stuffy nose, sneezing, sore throat .

2710. This particular disease, Psoriasis, interacts with the drug Calcitrene and has the following side effects:

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Calcitrene may cause serious side effects. Stop using Calcitrene and call your doctor at once if you have: severe burning, stinging, skin rash , or other irritation after applying the medicine
- worsening of your skin condition
- or high calcium levels--confusion, tiredness, nausea , vomiting , loss of appetite, constipation , increased thirst or urination, weight loss . Common side effects of Calcitrene may include: mild skin irritation
- skin rash
- or itching.

2711. This particular disease, Psoriasis, interacts with the drug coal tar/salicylic acid/sulfur and has the following side effects:

- In animal studies, coal tar (contained in this combination medicine) has been shown to increase the chance of skin cancer. Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: Skin irritation not present before use of salicylic acid, sulfur, and coal tar Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Coal tar/salicylic acid/sulfur topical side effects

2712. This particular disease, Psoriasis, interacts with the drug coal tar/salicylic acid and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs

Drug Side Effects Summary

of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness

- or swelling of the mouth, face, lips, tongue, or throat. Very bad skin irritation. What are some other side effects of Coal Tar and Salicylic Acid? All drugs may cause side effects. However, many people have no side effects or only have minor side effects. Call your doctor or get medical help if any of these side effects or any other side effects bother you or do not go away: Irritation where coal tar and salicylic acid is used. These are not all of the side effects that may occur. If you have questions about side effects, call your doctor. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-332-1088. You may also report side effects at <https://www.fda.gov/medwatch>. If OVERDOSE is suspected: If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened. How do I store and/or throw out Coal Tar and Salicylic Acid? Store at room temperature. Store in a dry place. Do not store in a bathroom. Keep all drugs in a safe place. Keep all drugs out of the reach of children and pets. Throw away unused or expired drugs. Do not flush down a toilet or pour down a drain unless you are told to do so. Check with your pharmacist if you have questions about the best way to throw out drugs. There may be drug take-back programs in your area. Consumer Information Use and Disclaimer If your symptoms or health problems do not get better or if they become worse, call your doctor. Do not share your drugs with others and do not take anyone else's drugs. Some drugs may have another patient

Drug Side Effects Summary

information leaflet. Check with your pharmacist. If you have any questions about coal tar and salicylic acid, please talk with your doctor, nurse, pharmacist, or other health care provider. If you think there has been an overdose, call your poison control center or get medical care right away. Be ready to tell or show what was taken, how much, and when it happened. More about coal tar/salicylic acid topical Check interactions Drug class: miscellaneous topical agents Related treatment guides Psoriasis Seborrheic Dermatitis Further information Always consult your healthcare provider to ensure the information displayed on this page applies to your personal circumstances. Medical Disclaimer Drug Status Availability Over the counter OTC CSA Schedule* Not a controlled drug N/A Loading... User Reviews & Ratings Review this drug Related Drugs Cosentyx , Humira , Stelara , Taltz , prednisone , methotrexate , dexamethasone , hydrocortisone topical , ketoconazole topical , Decadron Drugs.com Mobile Apps The easiest way to lookup drug information, identify pills, check interactions and set up your own personal medication records. Available for Android and iOS devices. Explore Apps Support Help Center Sitemap Contact Us About About Drugs.com Advertising Policy Attribution & Citations Terms & Privacy Editorial Policy Privacy Policy Terms of Use Follow Drugs.com on Facebook Follow Drugs.com on Twitter Follow Drugs.com on Pinterest Follow Drugs.com on YouTube Subscribe to Drugs.com newsletters Subscribe to Drugs.com newsletters for the latest medication news, new drug approvals, alerts and updates. Drugs.com provides accurate and independent information on more than 24,000 prescription drugs, over-the-counter medicines and natural products. This material is provided for educational purposes only and is not intended for medical advice, diagnosis or treatment. Data sources include IBM Watson Micromedex (updated 2 Dec 2022), Cerner Multum? (updated 7 Dec 2022), ASHP (updated 11 Nov 2022) and others. Ad Choices We comply with the HONcode standard for trustworthy health information. Copyright © 2000-2022 Drugs.com. All rights reserved.

SITECONFIG

=

```
{"agent":{"isMobile":false},"user":{"userid":0,"modal":{"login":"/api/user/modal/login/"},"consent":{"skip
```

Drug Side Effects Summary

```
":true,"api":{"url":"/api/user/"}}, "page":{"section":"cdi", "subsection":"","doc":"content","delayedCss":["/
bundle/css/component/component-share.min.9d6804aa.css"],"delayedScripts":[],"image":{"lazyLoad
":{"selector":"[data-lazy-load]","screens":1,"delay":2000}}, "mailto":{"delay":5000,"location":"/support/
email-mailto.html"}}, "form":{"recaptcha":{"sitekey":"6Ld-iSYTAAAAA00VTEwVI71HtvrFoSIPSibGr4
g","callback":"DDC_RECAPTCHA_CALLBACK"}}, "dwLog":{"a":714,"b":"null","c":"54,860","d":"132,1
34","e":"136","f":1213,"h":0,"i":"default","k":"54,860","l":"","m":"","n":"","o":"","p":0,"q":"drugs/cdi"},"dru
gs":[{"ddc_id":714,"brand_name_id":0,"mtm_id":"d04352","generic_name":"coal tar/salicylic acid
topical","name":"coal tar/salicylic acid
topical"}], "options":{"recordSelection":false,"classifyLists":{"wrapper":"#content","countHtmlItems":fal
se,"numItems":{"short":15,"medium":30,"long":60},"charLimit":{"paragraph":30,"content":350}}}, "subs
cribe":null,"modal":null,"dictionary":{},"share":{"sticky":{"target":".ddc-social-share"},"types":[{"title":"F
acebook","label":"Share on Facebook","cssClass":"ddc-share-icon
ddc-share-icon-facebook","labelClass":"ddc-share-icon-label","icon":"<svg class='ddc-icon
ddc-icon-facebook' width='20' height='20' viewBox='0 0 24 24' aria-hidden='true' focusable='false'
xmlns='http://www.w3.org/2000/svg'><path d='M14 6.54c-2.06 0-3.4 1.25-3.4
3.52v2H8.3v2.61h2.3V21A9.04 9.04 0 013 12.05C3 7.05 7.03 3 12 3s9 4.05 9 9.05c0 4.53-3.3
8.27-7.6 8.95v-6.33h2.1l4-2.62h-2.5v-1.7c0-.7.35-1.4 1.47-1.4h1.14V6.7s-1.03-.17-2.02-.17z'
/></svg>"},"shareUrl":"https://www.facebook.com/sharer/sharer.php?u=SHAREURL","width":600,"hei
ght":500}},{"title":"Twitter","label":"Share on Twitter","cssClass":"ddc-share-icon
ddc-share-icon-twitter","labelClass":"ddc-share-icon-label","icon":"<svg class='ddc-icon
ddc-icon-twitter' width='20' height='20' viewBox='0 0 24 24' aria-hidden='true' focusable='false'
xmlns='http://www.w3.org/2000/svg'><path d='M8.66 19.63a10.44 10.44 0 0010.5-10.5v-.5A7.7 7.7
0 0021 6.74c-.67.3-1.39.5-2.12.58a3.73 3.73 0 001.64-2.04c-.72.43-1.52.74-2.34.9a3.7 3.7 0 00-6.3
3.36 10.46 10.46 0 01-7.61-3.85 3.69 3.69 0 001.14 4.9 3.66 3.66 0 01-1.63-.45v.04a3.7 3.7 0 002.9
3.64 3.7 3.7 0 01-1.63.06 3.69 3.69 0 003.42 2.53 7.4 7.4 0 01-4.6 1.64 8 8 0 01-.87-.07 10.45
```

Drug Side Effects Summary

10.45 0 005.66 1.63'

/></svg>","shareUrl":"https://twitter.com/intent/tweet?text=SHARETITLE&url=SHAREURL","width":600,"height":420},{ "title":"Email","label":"Email to a friend","cssClass":"ddc-share-icon-ddc-share-icon-email","labelClass":"ddc-share-icon-label","icon":"<svg class='ddc-icon-ddc-icon-email' width='20' height='20' viewBox='0 0 24 24' aria-hidden='true' focusable='false' xmlns='http://www.w3.org/2000/svg'><path d='M20 4.25c.92 0 1.67.7 1.74 1.6l.01.15v12c0 .92-.7 1.67-1.6 1.74l-.15.01H4c-.92 0-1.67-.7-1.74-1.6L2.25 18V6c0-.92-.7-1.6-1.74L4 4.25h16zm-17 1.77a.75.75 0 00-1.06-.1L12 11.52l-6.77-5.6-.1-.06a.75.75 0 00-.86 1.22l7.25 6.1.07c.27.15.61.13.86-.07l7.25-6L19.8 7a.75.75 0 00.02-.98z' /></svg>","mailUrl":"mailto:?body=https%3A%2F%2Fwww.drugs.com%2Fcdi%2Fcoal-tar-and-salicylic-acid.html","width":600,"height":420}},{"mednotes":{"apiUrl":"/mednotes/api/modal","saveButton":"data-modal-mednotes-save"}},{"ads":{"section":"drugs/cdi","ugc":false,"minWidth728":1120,"minWidth970":1540,"display":{"confiant":false,"mobile":false,"timeout":900,"logUrl":"/api/log/ads/","targeting":{"ugc":"0","rc":"0","c1":["4223","6067"],"d":"714","b":"0","c2":["54","860"],"dc":["136"],"mi":"1","rx":"2","c2o":["54","860"]},"units":[{"code":"display-ad-1","section":"/7146/drugs/cdi","lazy":0,"sizes":[[728,90]],"sizeMappings":[[[0,0],[1],[[320,100],[[320,50],[300,50]]],[[768,300],[[728,90]]]],"targeting":{"uid":"d-leaderboard-1","mobile":0,"prebid":1,"size":"728x90","tier":"static","loading":"eager","position":1},"layout":{"prebid":1},"code":"display-ad-2","section":"/7146/drugs/cdi","lazy":0.75,"sizes":[[300,250],[300,600],[1,1]],"sizeMappings":[[[0,0],[1],[[320,100],[[300,250],[1,1]]],[[700,300],[[300,600],[300,250],[1,1]]]],"targeting":{"uid":"d-300x600-static-1","mobile":0,"prebid":1,"size":"300x600","tier":"static","loading":"lazy","position":2},"layout":{"prebid":1},"code":"display-ad-3","section":"/7146/drugs/cdi","lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[1],[[320,100],[[300,250],[1,1]]]],"targeting":{"uid":"d-300x250-static-1","mobile":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","position":3},"layout":{"wrappingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSidebarFloat":true},"prebid":1},"code":"display-ad-4","section":"/7146/drugs/cdi","lazy":0.75,"sizes":[[300,250],[1,1]]

Drug Side Effects Summary

, "sizeMappings": [[[0,0],[]], [[320,100],[[300,250],[1,1]]]], "targeting": {"uid": "d-300x250-static-2", "mobile": 0, "prebid": 1, "size": "300x250", "tier": "static", "loading": "lazy", "position": 4, "layout": {"wrappingClass": "sideBoxStackedAdWrap", "requiresContentHeight": true, "requiresSidebarFloat": true}, "prebid": 1}, {"code": "display-ad-5", "section": "/7146/drugs/cdi", "lazy": 0.75, "sizes": [[300,250],[1,1]], "sizeMappings": [[[0,0],[]], [[320,100],[[300,250],[1,1]]]], "targeting": {"uid": "d-300x250-static-3", "mobile": 0, "prebid": 1, "size": "300x250", "tier": "static", "loading": "lazy", "position": 5, "layout": {"wrappingClass": "sideBoxStackedAdWrap", "requiresContentHeight": true, "requiresSidebarFloat": true}, "prebid": 1}, {"code": "display-ad-6", "section": "/7146/drugs/cdi", "lazy": 0.75, "sizes": [[300,250],[1,1]], "sizeMappings": [[[0,0],[]], [[320,100],[[300,250],[1,1]]]], "targeting": {"uid": "d-300x250-static-4", "mobile": 0, "prebid": 1, "size": "300x250", "tier": "static", "loading": "lazy", "position": 6, "layout": {"wrappingClass": "sideBoxStackedAdWrap", "requiresContentHeight": true, "requiresSidebarFloat": true}, "prebid": 1}, {"code": "display-ad-7", "section": "/7146/drugs/cdi", "lazy": 0.75, "sizes": [[300,250],[1,1]], "sizeMappings": [[[0,0],[]], [[320,100],[[300,250],[1,1]]]], "targeting": {"uid": "d-300x250-static-5", "mobile": 0, "prebid": 1, "size": "300x250", "tier": "static", "loading": "lazy", "position": 7, "layout": {"wrappingClass": "sideBoxStackedAdWrap", "requiresContentHeight": true, "requiresSidebarFloat": true}, "prebid": 1}, {"code": "display-ad-8", "section": "/7146/drugs/cdi", "lazy": 0.75, "sizes": [[300,250],[1,1]], "sizeMappings": [[[0,0],[]], [[320,100],[[300,250],[1,1]]]], "targeting": {"uid": "d-300x250-static-6", "mobile": 0, "prebid": 1, "size": "300x250", "tier": "static", "loading": "lazy", "position": 8, "layout": {"wrappingClass": "sideBoxStackedAdWrap", "requiresContentHeight": true, "requiresSidebarFloat": true}, "prebid": 1}, {"code": "display-ad-9", "section": "/7146/drugs/cdi", "lazy": 0.75, "sizes": [[300,250],[1,1]], "sizeMappings": [[[0,0],[]], [[320,100],[[300,250],[1,1]]]], "targeting": {"uid": "d-300x250-static-7", "mobile": 0, "prebid": 1, "size": "300x250", "tier": "static", "loading": "lazy", "position": 9, "layout": {"wrappingClass": "sideBoxStackedAdWrap", "requiresContentHeight": true, "requiresSidebarFloat": true}, "prebid": 1}, {"code": "display-ad-10", "section": "/7146/drugs/cdi", "lazy": 0.75, "sizes": [[300,250],[1,1]], "sizeMappings": [[[0,0],[]], [[320,100],[[300,250],[1,1]]]], "targeting": {"uid": "d-300x250-static-8", "mobile": 0, "prebid": 1, "size": "300x250", "tier": "static", "loading": "lazy", "position": 10, "layout": {"wrappingClass": "sideBoxStackedAdWrap", "requiresContentHeight": true, "requiresSidebarFloat": true}, "prebid": 1}

Drug Side Effects Summary

ContentHeight":true,"requiresSidebarFloat":true},"prebid":1},{ "code":"display-ad-11","section":"/7146/drugs/cdi","lazy":0.75,"sizes":[[300,250],[1,1]],"sizeMappings":[[[0,0],[1,1]],[[320,100],[[300,250],[1,1]]]],"targeting":{"uid":"d-300x250-static-9","mobile":0,"prebid":1,"size":"300x250","tier":"static","loading":"lazy","position":11},"layout":{"wrappingClass":"sideBoxStackedAdWrap","requiresContentHeight":true,"requiresSidebarFloat":true},"prebid":1}},"injection":{"lazy":0.75,"maximum":{"ads":10,"tags":1000},"container":"#content","start":[".nav-tabs","h2","h1"],"stop":[".footer-feature","#sidebar"],"last":{"buffer":500},"insert":["section","h2","h3","h4","h5","p","ol","ul","table",".ddc-legend",".ddc-media",".nav-tabs",".nav-tabs-group",".responsive-table-wrap"],"blacklist":["form","ol","ul","dl","table",".blackboxWarning",".ddc-accordion",".ddc-box",".ddc-card",".ddc-card-group",".ddc-form-inline",".ddc-expand-read-more",".ddc-grid",".ddc-inset-text",".ddc-legend",".ddc-media",".ddc-notification",".ddc-paging",".ddc-reference-list",".ddc-toc",".ddc-widget-calculator",".injection-ignore-inside",".nav-tabs",".nav-tabs-group",".paging-list-wrap",".responsive-table-wrap",".referenceList",".snippet",".ddc-official-answer"],"list":{"position":1,"type":"Prebid","width":728,"height":90,"buffer":-1,"viewportPercent":85,"minBuffer":800,"lazy":0.75,"data":{"section":"/7146/drugs/cdi","lazy":0,"sizes":[[728,90]],"sizeMappings":[],"targeting":{"uid":"d-728x90-auto-1","mobile":0,"prebid":1,"size":"728x90","tier":"auto"},"prebid":1},"conditional":{"minWidth":1120,"fallback":{"position":1,"type":"Prebid","width":300,"height":250,"buffer":-1,"viewportPercent":85,"minBuffer":800,"lazy":0.75,"data":{"section":"/7146/drugs/cdi","lazy":0,"sizes":[[300,250]],"sizeMappings":[],"targeting":{"uid":"d-mrec-1","mobile":0,"prebid":1,"size":"300x250","tier":"auto"},"prebid":1}}}},"position":2,"type":"Prebid","width":728,"height":90,"buffer":1200,"lazy":0.75,"data":{"section":"/7146/drugs/cdi","lazy":0,"sizes":[[728,90]],"sizeMappings":[],"targeting":{"uid":"d-728x90-auto-2","mobile":0,"prebid":1,"size":"728x90","tier":"auto"},"prebid":1},"conditional":{"minWidth":1120,"fallback":{"position":2,"type":"Prebid","width":300,"height":250,"buffer":1200,"lazy":0.75,"data":{"section":"/7146/drugs/cdi","lazy":0,"sizes":[[300,250]],"sizeMappings":[],"targeting":{"uid":"d-300x250-auto-1","mobile":0,"prebid":1,"size":"300x250","tier":"auto"},"prebid":1}}}},"repeat":{"position":3,"type":"Prebid","width":728,"height":90,"buffer":1800,"lazy":0.75,"data":{"section":"/7146/drugs/cdi","lazy":0,"sizes":[[72

Drug Side Effects Summary

```
8,90]], "sizeMappings": [], "targeting": {"uid": "d-728x90-auto-3", "mobile": 0, "prebid": 1, "size": "728x90", "tier": "auto"}, "prebid": 1}, "conditional": {"minWidth": 1120, "fallback": {"position": 3, "type": "Prebid", "width": 300, "height": 250, "buffer": 1800, "lazy": 0.75, "data": {"section": "/7146/drugs/cdi", "lazy": 0, "sizes": [[300, 250]]}, "sizeMappings": [], "targeting": {"uid": "d-300x250-auto-2", "mobile": 0, "prebid": 1, "size": "300x250", "tier": "auto"}, "prebid": 1}}}, {"nativeApp": {"version": 0, "appSrc": "", "memoryWarning": {"log": 0, "ads": 1}}, "fixable": {"items": [{"targetSelector": ".topbanner-wrap", "delay": 2500, "leaderboard": true}, {"containerSelector": "#contentWrap", "targetSelector": "#sideBoxStackedAd-1", "delay": 2500, "maximizeStickyHeight": true}, {"containerSelector": "#contentWrap", "targetSelector": "#sideBoxStackedAd-2", "delay": 3000, "maximizeStickyHeight": true}, {"containerSelector": "#contentWrap", "targetSelector": "#sideBoxStackedAd-3", "delay": 3500, "maximizeStickyHeight": true}, {"containerSelector": "#contentWrap", "targetSelector": "#sideBoxStackedAd-4", "delay": 4000, "maximizeStickyHeight": true}, {"containerSelector": "#contentWrap", "targetSelector": "#sideBoxStackedAd-5", "delay": 4500, "maximizeStickyHeight": true}, {"containerSelector": "#contentWrap", "targetSelector": "#sideBoxStackedAd-6", "delay": 5000, "maximizeStickyHeight": true}, {"containerSelector": "#contentWrap", "targetSelector": "#sideBoxStackedAd-7", "delay": 5500, "maximizeStickyHeight": true}, {"containerSelector": "#contentWrap", "targetSelector": "#sideBoxStackedAd-8", "delay": 6000, "maximizeStickyHeight": true}, {"containerSelector": "#contentWrap", "targetSelector": "#sideBoxStackedAd-9", "delay": 6500, "maximizeStickyHeight": true}, {"containerSelector": "#contentWrap", "targetSelector": "#sideBoxStackedAd-10", "delay": 7000, "maximizeStickyHeight": true}]}], "toc": {}, "interaction": {"isDeferredSave": false, "url": "/interaction/list/", "param": {"drugList": "drug_list", "listId": "interaction_list_id"}, "identityLink": {"api": {"url": "/api/user/"}, "cookie": {"link": "pbjs_identitylink", "data": "pbjs_identitydata"}}
```

```
- window.dataLayer = window.dataLayer || []
```

```
- window.dataLayer.push({'gtm.start': new Date().getTime(), event: 'gtm.js'})
```

```
- window.dataLayer.push({"event": "pageview", "drug_name": "coal tar/salicylic acid topical", "drug_id": "714-0", "ddc_id": "714", "condition_ids": ""
```

Drug Side Effects Summary

```
- 54
- 860
- ", "zone_ids":
- 4223
- 6067
-
", "user_signed_in": "No", "user_ilink": "No", "page_index_status": "index/follow", "page_orphan": "No", "page_traffic_op": "No", "content_type": "Drug", "content_section": "Medfacts
Consumer", "content_sub_section": "Medfacts
Consumer", "content_audience": "Consumer", "content_language": "English", "content_size": "Medium
Page", "content_ugc": "No", "app_source": "Web", "ads_pro": "No"))
- var _comscore = _comscore || []
- _comscore.push({c1: '2', c2: '9355535'})
- (function(){/* Copyright The Closure Library Authors. SPDX-License-Identifier: Apache-2.0 */ 'use
strict'
- var g=function(a){var b=0
-                                     return                                     function(){return
b<a.length?{done:!1,value:a[b++]}:{done:!0}}},l=this||self,m=/^[w+/_-]+[=]{0,2}$/ ,p=null,q=function(){
,r=function(a){var b=typeof a
- if("object"==b)if(a){if(a instanceof Array)return"array"
- if(a instanceof Object)return b
- var c=Object.prototype.toString.call(a)
- if("[object Window]"==c)return"object"
- if("[object Array]"==c||"number"==typeof a.length&&"undefined"!=typeof
a.splice&&"undefined"!=typeof
```

Drug Side Effects Summary

a.propertyIsEnumerable&&!a.propertyIsEnumerable("splice"))return"array"

- if("[object Function]"==c||"undefined"!=typeof a.call&&"undefined"!=typeof

a.propertyIsEnumerable&&!a.propertyIsEnumerable("call"))return"function"}else return"null"

- else if("function"==b&&"undefined"==typeof a.call)return"object"

- return b},u=function(a,b){function c(){c.prototype=b.prototype

- a.prototype=new c

- a.prototype.constructor=a}

- var v=function(a,b){Object.defineProperty(l,a,{configurable:!1,get:function(){return b},set:q}})

- var y=function(a,b){this.b=a===w&&b||""

- this.a=x,x={},w={}

- var aa=function(a,b){a.src=b instanceof

y&&b.constructor===y&&b.a===x?b.b:"type_error:TrustedResourceUrl"

- if(null===p)b:{b=l.document

-

if((b=b.querySelector&&b.querySelector("script[nonce]"))&&(b=b.nonce||b.getAttribute("nonce"))&&

m.test(b)){p=b

- break b}p=""b=p

- b&&a.setAttribute("nonce",b)}

- var z=function(){return

Math.floor(2147483648*Math.random()).toString(36)+Math.abs(Math.floor(2147483648*Math.random())^+new Date).toString(36)}

- var A=function(a,b){b=String(b)

- "application/xhtml+xml"===a.contentType&&(b=b.toLowerCase())

- return a.createElement(b)},B=function(a){this.a=a||l.document||document}

- B.prototype.appendChild=function(a,b){a.appendChild(b)}

Drug Side Effects Summary

```
- var C=function(a,b,c,d,e,f){try{var k=a.a,h=A(a.a,"SCRIPT")

- h.async=!0

- aa(h,b)

- k.head.appendChild(h)

- h.addEventListener("load",function(){e()

- d&&k.head.removeChild(h)})}

-

h.addEventListener("error",function(){0<c?C(a,b,c-1,d,e,f):(d&&k.head.removeChild(h),f())}})catch(n)

{f()}}

-

var

ba=l.atob("aHR0cHM6Ly93d3cuZ3N0YXRpYy5jb20vaW1hZ2VzL2ljb25zL21hdGVyaWFsL3N5c3Rl

bS8xeC93YXJuaW5nX2FtYmVyXzI0ZHAucG5n"),ca=l.atob("WW91IGFyZSBzZWVpbmcgdGhpcyBt

ZXNzYWdlIGJlY2F1c2UgYWQgb3lgc2NyaXB0IGJsb2NraW5nIHNVZnR3YXJlIGlzlGludGVyZmVya

W5nIHdpdGggdGhpcyBwYWdlLg=="),da=l.atob("RGlzYWJsZSBhbnkgYWQgb3lgc2NyaXB0IGJsb2

NraW5nIHNVZnR3YXJlCB0aGVuIHJlbG9hZCB0aGlzIHBoZ2Uu"),ea=function(a,b,c){this.b=a

- this.f=new B(this.b)

- this.a=null

- this.c=[]

- this.g=!1

- this.i=b

- this.h=c},F=function(a){if(a.b.body&&!a.g){var b= function(){D(a)

- l.setTimeout(function(){return E(a,3)},50)}

- C(a.f,a.i,2,!0,function(){l[a.h]||b()},b)

- a.g=!0}},D=function(a){for(var b=G(1,5),c=0

- c<b
```

Drug Side Effects Summary

- c++){var d=H(a)
- a.b.body.appendChild(d)
- a.c.push(d)}b=H(a)
- b.style.bottom="0"
- b.style.left="0"
- b.style.position="fixed"
- b.style.width=G(100,110).toString()+"%"
- b.style.zIndex=G(2147483544,2147483644).toString()
- b.style["background-color"]=l(249,259,242,252,219,229)
- b.style["box-shadow"]="0 0 12px #888"
- b.style.color=l(0,10,0,10,0,10)
- b.style.display= "flex"
- b.style["justify-content"]="center"
- b.style["font-family"]="Roboto, Arial"
- c=H(a)
- c.style.width=G(80,85).toString()+"%"
- c.style.maxWidth=G(750,775).toString()+"px"
- c.style.margin="24px"
- c.style.display="flex"
- c.style["align-items"]="flex-start"
- c.style["justify-content"]="center"
- d=A(a.f.a,"IMG")
- d.className=z()
- d.src=ba
- d.style.height="24px"

Drug Side Effects Summary

- d.style.width="24px"
- d.style["padding-right"]="16px"
- var e=H(a),f=H(a)
- f.style["font-weight"]="bold"
- f.textContent=ca
- var k=H(a)
- k.textContent=da
- J(a, e,f)
- J(a,e,k)
- J(a,c,d)
- J(a,c,e)
- J(a,b,c)
- a.a=b
- a.b.body.appendChild(a.a)
- b=G(1,5)
- for(c=0
- c<b
- c++)d=H(a),a.b.body.appendChild(d),a.c.push(d)},J=function(a,b,c){for(var d=G(1,5),e=0
- e<d
- e++){var f=H(a)
- b.appendChild(f)}b.appendChild(c)
- c=G(1,5)
- for(d=0
- d<c
- d++)e=H(a),b.appendChild(e)},G=function(a,b){return

Drug Side Effects Summary

```
Math.floor(a+Math.random()*(b-a))),l=function(a,b,c,d,e,f){return"rgb("+G(Math.max(a,0),Math.min(b,255)).toString()+","+G(Math.max(c,0),Math.min(d,255)).toString()+","+G(Math.max(e,0),Math.min(f,255)).toString()+")"},H=function(a){a=A(a.f.a,"DIV")
```

```
- a.className=z()
```

```
-
```

```
return
```

```
a},E=function(a,b){0>=b||null!=a.a&&0!=a.a.offsetHeight&&0!=a.a.offsetWidth||(fa(a),D(a),l.setTimeout(function(){return E(a,b-1)},50))),fa=function(a){var b=a.c
```

```
- var c="undefined"!=typeof Symbol&&Symbol.iterator&&b[Symbol.iterator]
```

```
- b=c?c.call(b):{next:g(b)}
```

```
- for(c=b.next()
```

```
- !c.done
```

```
- c=b.next()(c=c.value)&&c.parentNode&&c.parentNode.removeChild(c)
```

```
- a.c=[]
```

```
- (b=a.a)&&b.parentNode&&b.parentNode.removeChild(b)
```

```
- a.a=null}
```

```
- var ia=function(a,b,c,d,e){var f=ha(c),k=function(n){n.appendChild(f)
```

```
-
```

```
l.setTimeout(function(){f?(0!==f.offsetHeight&&0!==f.offsetWidth?b():a(),f.parentNode&&f.parentNode.removeChild(f)):a()},d)},h=function(n){document.body?k(document.body):0<n?!.setTimeout(function(){h(n-1)},e):b()}
```

```
- h(3)},ha=function(a){var b=document.createElement("div")
```

```
- b.className=a
```

```
- b.style.width="1px"
```

```
- b.style.height="1px"
```

```
- b.style.position="absolute"
```

Drug Side Effects Summary

```
- b.style.left="-10000px"
- b.style.top="-10000px"
- b.style.zIndex="-10000"
- return b}
- var K={},L=null
- var M=function() {},N="function"==typeof Uint8Array,O=function(a,b){a.b=null
- b||(b=[])
- a.j=void 0
- a.f=-1
- a.a=b
- a:{if(b=a.a.length){--b
- var c=a.a[b]
- if(!(null===c||"object"!=typeof c||Array.isArray(c)||N&&c instanceof Uint8Array)){a.g=b-a.f
- a.c=c
- break a}}a.g=Number.MAX_VALUE}a.i={},P=[],Q=function(a,b){if(b<a.g){b+=a.f
- var c=a.a[b]
- return c===P?a.a[b]=[]:c}if(a.c)return c=a.c[b],c===P?a.c[b]=[]:c,R=function(a,b,c){a.b||(a.b={})
- if(!a.b[c]){var d=Q(a,c)
- d&&(a.b[c]=new b(d))}return a.b[c]}
- M.prototype.h=N?function(){var a=Uint8Array.prototype.toJSON
- Uint8Array.prototype.toJSON=function(){var b
- void 0===b&&(b=0)
- if(!L){L={}
-
for(var
c="ABCDEFGHJKLMNOPQRSTUVWXYZabcdefghijklmnopqrstuvwxyz0123456789".split(""),d=["+/"
```


Drug Side Effects Summary

=", "+/", "-_=", "-_.", "-_"], e=0

- 5>e

- e++){var f=c.concat(d[e].split(""))

- K[e]=f

- for(var k=0

- k<f.length

- k++){var h=f[k]

- void 0===L[h]&&(L[h]=k)}}b=K[b]

- c=[]

- for(d=0

- d<this.length

- d+=3){var n=this[d],t=(e=d+1<this.length)?this[d+1]:0

- h=(f=d+2<this.length)?this[d+2]:0

- k=n>>2

- n=(n&3)<<4|t>>4

- t=(t&15)<<2|h>>6

- h&=63

- f||(h=64,e||(t=64))

- c.push(b[k],b[n],b[t]||"",b[h]||"")}return c.join("")}

-

try{return

JSON.stringify(this.a&&this.a,S)}finally{Uint8Array.prototype.toJSON=a}}:function(){return

JSON.stringify(this.a&&this.a,S)}

- var S=function(a,b){return"number"!==typeof b||!isNaN(b)&&Infinity!==b&&-Infinity!==b?b:String(b)}

- M.prototype.toString=function(){return this.a.toString()}

- var T=function(a){O(this,a)}

Drug Side Effects Summary

- u(T,M)
- var U=function(a){O(this,a)}
- u(U,M)
- var ja=function(a,b){this.c=new B(a)
- var c=R(b,T,5)
- c=new y(w,Q(c,4)||"")
- this.b=new ea(a,c,Q(b,4))
- this.a=b},ka=function(a,b,c,d){b=new T(b?JSON.parse(b):null)
- b=new y(w,Q(b,4)||"")
- C(a.c,b,3,!1,c,function()){ja(function(){F(a.b
- d(!1)},function(){d(!0)},Q(a.a,2),Q(a.a,3),Q(a.a,1))}}}
- var la=function(a,b){V(a,"internal_api_load_with_sb",function(c,d,e){ka(b,c,d,e)})
- V(a,"internal_api_sb",function(){F(b.b)}),V=function(a,b,c){a=l.btoa(a+b)
- v(a,c)},W=function(a,b,c){for(var d=[],e=2
- e<arguments.length
- ++e)d[e-2]=arguments[e]
- e=l.btoa(a+b)
- e=l[e]
- if("function"==r(e))e.apply(null,d)
- else throw Error("API not exported.")
- }
- var X=function(a){O(this,a)}
- u(X,M)
- var Y=function(a){this.h=window
- this.a=a

Drug Side Effects Summary

- this.b=Q(this.a,1)
- this.f=R(this.a,T,2)
- this.g=R(this.a,U,3)
- this.c=!1}
- Y.prototype.start=function(){ma()
- var a=new ja(this.h.document,this.g)
- la(this.b,a)
- na(this)}
- var ma=function(){var a=function(){if(!l.frames.googlefcPresent)if(document.body){var b=document.createElement("iframe")
- b.style.display="none"
- b.style.width="0px"
- b.style.height="0px"
- b.style.border="none"
- b.style.zIndex="-1000"
- b.style.left="-1000px"
- b.style.top="-1000px"
- b.name="googlefcPresent"
- document.body.appendChild(b)}else l.setTimeout(a,5)}
- a()},na=function(a){var b=Date.now()
- W(a.b,"internal_api_load_with_sb",a.f.h(),function(){var c
- var d=a.b,e=l[l.btoa(d+"loader_js")]
- if(e){e=l.atob(e)
- e=parseInt(e,10)
- d=l.btoa(d+"loader_js").split(".")

Drug Side Effects Summary

- var f=1

```
- d[0]in f||"undefined"==typeof f.execScript||f.execScript("var "+d[0])
```

- for(

```
- d.length&&(c=d.shift())
```

```
- )d.length?f[c]&&f[c]!==Object.prototype[c]?f[c]:f[c]={}:f[c]=null
```

```
- c=Math.abs(b-e)
```

```
- c=1728E5>c?0:c}else c=-1
```

—

```
0!=c&&(W(a.b,"internal_api_sb"),Z(a,Q(a.a,6))),function(c){Z(a,c?Q(a.a,4):Q(a.a,5))}},Z=function(a,
b){a.c||(a.c=!0,a=new XMLHttpRequest,a.open("GET",b,!0),a.send())}
```

```
- (function(a,b){l[a]=function(c){for(var d=[],e=0
```

- e<arguments.length

```
- ++e)d[e-0]=arguments[e]
```

- $I[a]=q$

```
- b.apply(null,d)}}("__d3lUW8vwsKIB__",function(a){"function"==typeof
window.atob&&(a=window.atob(a),a=new X(a?JSON.parse(a):null),(new Y(a)).start()))}
```

- }).call(this)

—

window.__d3IUW8vwsKIB__("WlyYzZIMmQxOTc1M2U5ZGEzlixbbnVsbCxudWxsLG51bGwslmh0
 dHBzOi8vZnVuZGluZ2Nob2ljZXNtZXNzYWdlcy5nb29nbGUuY29tL2YvQUdTS1d4VmZTeHR0VzVf
 LXo0NIYxeGhodEk4M053bE41MjJtZkdM0V1plV2xnQ2w1cW50NkRsM1luZ3QtSmQ
 1bXZsc2s4QXJnUWJMRnBtbW1ZY1x1MDAzZCJdCixbMjAsImRpd1ncHQyYVQlDEwMCwiTW1N
 MlpUSmtNVGszTIRObE9XUmhNd1x1MDAzZFx1MDAzZCIsW251bGwsbnVsbCxudWxsLCJodHRw
 czovL3d3dy5nc3RhdGljLmNvbS8wZW1uL2YvcC8yYzZIMmQxOTc1M2U5ZGEzLmpzP3VzcXBcdT
 AwM2RDQkkiXQpdCiwiaHR0cHM6Ly9mdW5kaW5nY2hvaWNlc21lc3NhZ2VzLmdvb2dsZS5jb20vb

Drug Side Effects Summary

C9BR1NLV3hVQIJCUV9HOGIYM0NDQUpKT09RM3ILNG5LYzNrUkNyeW13Tk1JVkhZUGo1WUsw
WndVRVI0aHI2N3FBRnQ1cl9IODJRWWU1Zkp0S184M0o2dVpqP2FiXHUwMDNkMSIsImh0dHBzO
i8vZnVuZGluZ2Nob2ljZXNtZXNzYWdlcy5nb29nbGUuY29tL2wvQUdTS1d4WE9yRjJUOWFjVE5kQz
B5UXE5UVczNjZDd3h4X1dMZVpLYk4wcVAyM1c2ZXd1UHM4ajdCNTIsRIM0RVZERUVhUWRGN
UF5SW1vM2RVQzcyUWg5VD9hYlx1MDAzZDJcdTAwMjZzYmZcdTAwM2QxliwiaHR0cHM6Ly9md
W5kaW5nY2hvaWNlc21lc3NhZ2VzLmdvb2dsZS5jb20vbC9BR1NLV3hXR1lwVEdYWUxWU3FKQlo
xQTJkUjk5S2t2MHg1NjBJelpzLW5ZZHlITjRSUDNfNIZnTzMxRG5xU0VjTzNsSIIyaFpuckpQeFNwe
UNfSGluMVVxP3NiZlx1MDAzZDliXQo=")

-

2713. This particular disease, Psoriasis, interacts with the drug Cutar and has the following side effects:

- In animal studies, coal tar has been shown to increase the chance of skin cancer. Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: Rare Skin irritation not present before use of this medicine skin rash Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Stinging (mild)--especially for gel and solution dosage forms Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Cutar side effects

2714. This particular disease, Psoriasis, interacts with the drug Cyltezo and has the following side effects:

Drug Side Effects Summary

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Bladder pain bloody or cloudy urine body aches or pain chills cough difficult, burning, or painful urination difficulty in breathing ear congestion fever frequent urge to urinate headache loss of voice lower back or side pain runny or stuffy nose sneezing sore throat unusual tiredness or weakness Less common Abnormal healing anxiety black, tarry stools bloating or swelling of the face, arms, hands, lower legs, or feet blue or pale skin blurred vision chest pain, possibly moving to the left arm, neck, or shoulder dark urine decreased vision diarrhea dizziness fainting fast, pounding, or irregular heartbeat or pulse hoarseness indigestion itching, skin rash light-colored stools loss of appetite loss of strength or energy muscle pain or weakness nausea pain in the throat, arms, legs, or pelvis pain or discomfort in the arms, jaw, back, or neck painful or difficult urination rapid weight gain severe nausea or vomiting sores, ulcers, or white spots on the lips or in the mouth stomach pain sweating tingling of the hands or feet unpleasant breath odor unusual bleeding or bruising unusual weight gain or loss vomiting vomiting of blood yellow eyes or skin Incidence not known Blindness blistering, peeling, or loosening of the skin blue-yellow color blindness blurred vision constipation darkened urine eye pain indigestion joint or muscle pain pain, redness, or swelling in the arm or leg pains in the stomach, side, or abdomen, possibly radiating to the back red, irritated eyes red skin lesions, often with a purple center trouble breathing Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Back pain bleeding, blistering, burning, coldness, discoloration of skin, feeling of pressure, hives, infection, inflammation, itching, lumps, numbness, pain, rash, redness, scarring, soreness, stinging, swelling,

Drug Side Effects Summary

tenderness, tingling, ulceration, or warmth at the injection site Incidence not known Loss or thinning of the hair Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Cyltezo side effects

2715. This particular disease, Psoriasis, interacts with the drug DHS Tar Shampoo and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. DHS Tar Shampoo may cause serious side effects. Stop using DHS Tar Shampoo and call your doctor at once if you have: severe stinging, burning, swelling, or other irritation of the treated skin. Common side effects may include mild skin irritation or skin rash .

2716. This particular disease, Psoriasis, interacts with the drug Doak Tar and has the following side effects:

- In animal studies, coal tar has been shown to increase the chance of skin cancer. Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: Rare Skin irritation not present before use of this medicine skin rash Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Stinging (mild)--especially for gel and solution dosage forms Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice

Drug Side Effects Summary

about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Doak Tar side effects

2717. This particular disease, Psoriasis, interacts with the drug Dritho-Scalp and has the following side effects:

- Anthralin has been shown to cause tumors in animals. However, there have been no reports of anthralin causing tumors in humans. Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: More common Redness or other skin irritation of treated or uninvolved skin not present before use of this medicine Rare Skin rash Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Dritho-Scalp side effects

2718. This particular disease, Psoriasis, interacts with the drug Drithocrema and has the following side effects:

- Anthralin has been shown to cause tumors in animals. However, there have been no reports of anthralin causing tumors in humans. Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: More common Redness or other skin irritation of treated or uninvolved skin not present before use of this medicine Rare Skin rash Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Drithocrema side effects

Drug Side Effects Summary

2719. This particular disease, Psoriasis, interacts with the drug Elta Tar and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Elta Tar may cause serious side effects. Stop using Elta Tar and call your doctor at once if you have: severe stinging, burning, swelling, or other irritation of the treated skin. Common side effects may include mild skin irritation or skin rash .

2720. This particular disease, Psoriasis, interacts with the drug Epifoam and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of a weak adrenal gland like a very bad upset stomach or throwing up, very bad dizziness or passing out, muscle weakness, feeling very tired, mood changes, not hungry, or weight loss . Signs of Cushing's disease like weight gain in the upper back or belly, moon face, very bad headache , or slow healing. Signs of high blood sugar like confusion, feeling sleepy, more thirst, more hungry, passing urine more often, flushing, fast breathing, or breath that smells like fruit. Irritation where Epifoam (pramoxine and hydrocortisone

Drug Side Effects Summary

foam) is used. Skin changes (pimples , stretch marks, slow healing, hair growth). Thinning of the skin. Epifoam side effects

2721. This particular disease, Psoriasis, interacts with the drug Estar and has the following side effects

- In animal studies, coal tar has been shown to increase the chance of skin cancer. Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: Rare Skin irritation not present before use of this medicine skin rash Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Stinging (mild)--especially for gel and solution dosage forms Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Estar side effects

2722. This particular disease, Psoriasis, interacts with the drug Fototar and has the following side effects

- In animal studies, coal tar has been shown to increase the chance of skin cancer. Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: Rare Skin irritation not present before use of this medicine skin rash Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side

Drug Side Effects Summary

effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Stinging (mild)--especially for gel and solution dosage forms Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Fototar side effects

2723. This particular disease, Psoriasis, interacts with the drug Hadlima and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Bladder pain bloody or cloudy urine blurred vision chest pain or tightness chills cough difficult, burning, or painful urination dizziness ear congestion or pain fever frequent urge to urinate head congestion headache hoarseness or other voice changes lower back or side pain nervousness pain or tenderness around the eyes and cheekbones pounding in the ears slow or fast heartbeat sneezing sore throat stuffy or runny nose trouble breathing Less common Abnormal healing agitation anxiety arm, back, jaw pain bleeding from the gums or nose blindness bloating or swelling of the face, arms, hands, lower legs, or feet bloody or black, tarry stools bloody or cloudy urine blue or pale skin bone pain or fractures burning, crawling, itching, numbness, prickling, "pins and needles" , or tingling feelings chest pain, discomfort, tightness, or heaviness coma confusion cough or hoarseness dark urine decreased vision depression diarrhea difficulty in moving difficulty in swallowing drowsiness dry mouth dry skin eye pain fainting feeling of illness frequent urge to urinate gaseous stomach pain hair loss hallucinations heartburn increased thirst increased urination indigestion irritability itching, skin rash loss of appetite loss of strength or energy light-colored stools lower back or side pain mood or mental changes muscle pain, stiffness, cramps, tightness, rigidity, or spasms nausea no blood pressure or pulse pain, swelling, or redness in the joints pain in the arms or legs pain in the groin or

Drug Side Effects Summary

genitals pain or burning in the throat painful or difficult urination pale skin pelvic pain rapid breathing rapid weight gain or loss recurrent fever redness or swelling of the lower leg ringing in the ears unconsciousness seizures shakiness in the legs, arms, hands, or feet sharp back pain just below ribs sore throat sores, ulcers, or white spots on the lips or tongue or inside the mouth stiff neck stomach pain stopping of the heart sunken eyes sweating tingling of the hands or feet trembling or shaking of the hands or feet trouble thinking unpleasant breath odor unusual bleeding or bruising unusual weight gain or loss visual disturbances vomiting vomiting of blood or material that looks like coffee grounds wrinkled skin yellow eyes or skin Incidence not known Blindness blistering, peeling, or loosening of the skin decreased urine output decreased vision dilated neck veins irregular breathing irregular heartbeat red, irritated eyes red skin lesions, often with a purple center swelling of the face, fingers, feet, or lower legs unusual tiredness or weakness weight gain Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Bleeding, blistering, burning, coldness, discoloration of the skin, feeling of pressure, hives, infection, inflammation, itching, lumps, numbness, pain, rash, redness, scarring, soreness, stinging, swelling, tenderness, tingling, ulceration, or warmth at the injection site Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Hadlima side effects

2724. This particular disease, Psoriasis, interacts with the drug Hulio and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your

Drug Side Effects Summary

doctor immediately if any of the following side effects occur: More common Bladder pain bloody or cloudy urine chest pain or tightness chills cough difficult, burning, or painful urination ear congestion or pain fever frequent urge to urinate headache hoarseness or other voice changes lower back or side pain pain or tenderness around the eyes and cheekbones sneezing sore throat stuffy or runny nose trouble breathing Less common Black, tarry stools blindness blurred vision bone fractures confusion constipation dark urine decreased vision difficulty with moving dizziness dry skin eye pain hair loss increased thirst increased urination itching, skin rash joint pain light-colored stools loss of appetite muscle cramps, pain, spasms, or stiffness nausea nervousness pain in the arms or legs pounding in the ears seizures slow or fast heartbeat sores, ulcers, or white spots in the mouth or on the lips stomach pain trouble thinking unpleasant breath odor unusual bleeding or bruising unusual tiredness or weakness vomiting vomiting of blood yellow eyes or skin Incidence not known Blistering, peeling, or loosening of the skin decreased urine output diarrhea dilated neck veins irregular breathing irregular heartbeat joint pain numbness or tingling in the fingers, face, or feet. red, irritated eyes red skin lesions, often with a purple center swelling of the face, fingers, feet, or lower legs weight gain Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Bleeding, blistering, burning, coldness, discoloration of the skin, feeling of pressure, hives, infection, inflammation, itching, lumps, numbness, pain, rash, redness, scarring, soreness, stinging, swelling, tenderness, tingling, ulceration, or warmth at the injection site Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Hulo side effects

Drug Side Effects Summary

2725. This particular disease, Psoriasis, interacts with the drug hydrocortisone / pramoxine and has

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop using this medication and call your doctor at once if you show signs of absorbing too much of the medicine through your skin , such as: blurred vision, eye pain , or seeing halos around lights
- mood changes
- sleep problems (insomnia)
- high blood sugar (increased thirst or urination, hunger, dry mouth, fruity breath odor, weight loss)
- weight gain, puffiness in your face
- or muscle weakness, feeling tired. Stop using the medicine and call your doctor if you have rectal bleeding or severe irritation in any treated skin . Common side effects of hydrocortisone and pramoxine may include: mild redness or swelling of treated skin
- thinning of treated skin
- or numbness on areas where the medicine is accidentally applied.

2726. This particular disease, Psoriasis, interacts with the drug hydroxyurea and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Hydroxyurea may cause serious side effects. Call your doctor at once if you have: numbness , tingling, or burning pain in your hands or feet
- skin numbness or purple discoloration
- skin ulcers or open sores
- sudden chest pain, wheezing, dry cough, feeling short of breath
- low blood cell counts--fever, chills, tiredness, mouth sores, easy bruising, unusual bleeding, pale

Drug Side Effects Summary

skin, cold hands and feet, feeling light-headed

- or signs of liver or pancreas problems--loss of appetite, upper stomach pain (that may spread to your back), nausea or vomiting , fast heart rate , dark urine, jaundice (yellowing of the skin or eyes).

Common side effects of hydroxyurea may include: loss of appetite, nausea

- constipation , diarrhea

- low blood cell counts

- bleeding

- or mouth sores.

2727. This particular disease, Psoriasis, interacts with the drug Hyrimoz and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Bladder pain bloody or cloudy urine body aches or pain chest pain or tightness chills cough difficult, burning, or painful urination ear congestion fever frequent urge to urinate headache hoarseness or other voice changes lower back or side pain pain or tenderness around the eyes and cheekbones sneezing sore throat stuffy or runny nose trouble breathing unusual tiredness or weakness Less common Black, tarry stools blindness blue or pale skin bone fractures confusion dark urine decreased vision difficulty with moving dizziness dry skin eye pain fainting fast, pounding, or irregular heartbeat or pulse hoarseness increased thirst increased urination itching, skin rash indigestion light-colored stools loss of appetite loss of strength or energy muscle cramps, pain, spasms, or stiffness nausea nervousness pain in the arms or legs pounding in the ears seizures slow or fast heartbeat sores, ulcers, or white spots on the lips or in the mouth stomach pain trouble thinking unpleasant breath odor unusual bleeding or bruising vomiting vomiting of blood yellow eyes or skin Incidence not known Blistering, peeling, or loosening of the skin decreased urine output diarrhea dilated neck

Drug Side Effects Summary

veins irregular breathing irregular heartbeat joint pain numbness or tingling in the fingers, face, or feet. red, irritated eyes red skin lesions, often with a purple center swelling of the face, fingers, feet, or lower legs weight gain Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Back pain bleeding, blistering, burning, coldness, discoloration of the skin, feeling of pressure, hives, infection, inflammation, itching, lumps, numbness, pain, rash, redness, scarring, soreness, stinging, swelling, tenderness, tingling, ulceration, or warmth at the injection site Incidence not known Hair loss, thinning of hair Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Hyrimoz side effects

2728. This particular disease, Psoriasis, interacts with the drug Impoyz and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Impoyz may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain in your face and shoulders, slow wound healing, skin discoloration, thinning skin, increased body hair, tiredness, mood changes, menstrual changes, sexual changes. Common side effects of Impoyz may include:

Drug Side Effects Summary

burning, itching, swelling, or irritation of treated skin

- dry or cracking skin
- redness or crusting around your hair follicles
- spider veins
- stretch marks, thinning skin
- rash or hives
- acne
- or temporary hair loss .

2729. This particular disease, Psoriasis, interacts with the drug Lonil T and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Lonil T may cause serious side effects. Stop using Lonil T and call your doctor at once if you have: severe stinging, burning, swelling, or other irritation of the treated skin. Common side effects may include mild skin irritation or skin rash .

2730. This particular disease, Psoriasis, interacts with the drug Ixifi and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor or nurse immediately if any of the following side effects occur: More common Black, tarry stools bladder pain bloody or cloudy urine blurred vision body aches or pain chest pain chills cough cough producing mucus difficult, burning, or painful urination difficulty breathing dizziness dryness or soreness of the throat ear congestion fever frequent urge to urinate headache hoarseness itching, rash loss of voice lower back or side pain nervousness pain or swelling in the arms or legs pain or tenderness around the eyes and cheekbones painful or difficult urination pale skin pounding in the

Drug Side Effects Summary

ears slow or fast heartbeat sneezing sores, ulcers, or white spots on the lips or in the mouth stuffy or runny nose swollen glands tender, swollen glands in the neck tightness in the chest trouble swallowing troubled breathing with exertion unusual bleeding or bruising unusual tiredness or weakness voice changes Rare Bleeding gums blood in the stool blue lips and fingernails changes in skin color or tenderness of the foot or leg chest discomfort confusion coughing that sometimes produces a pink frothy sputum dark urine decreased urination dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position dry mouth fast or noisy breathing feeling of discomfort general feeling of illness general tiredness and weakness high fever hives increased sweating irregular heartbeat light-colored stools muscle aches nausea pinpoint red spots on the skin puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue rapid, shallow breathing swelling of the joints sunken eyes thirst upper right abdominal or stomach pain vomiting weight loss wrinkled skin yellow eyes and skin Incidence not known Back pain, sudden and severe blistering, peeling, loosening of the skin bloody nose burning, tingling, numbness or pain in the hands, arms, feet, or legs change in mental status clay-colored stools diarrhea difficulty speaking dilated neck veins double vision heavier menstrual periods inability to move the arms, legs, or facial muscles inability to speak loss of appetite muscle weakness, sudden and progressing red, irritated eyes red skin lesions, often with a purple center seizures sensation of pins and needles slow or irregular breathing slow speech stabbing pain temporary vision loss vomiting of blood weight gain Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Belching diarrhea difficulty in moving feeling of warmth heartburn indigestion muscle stiffness redness of the face, neck, arms, and occasionally, upper chest Less common Constipation, severe Other side effects not listed may

Drug Side Effects Summary

also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Ixifi side effects

2731. This particular disease, Psoriasis, interacts with the drug Kalosar and has the following side effects:

- No side effects listed

2732. This particular disease, Psoriasis, interacts with the drug Medotar and has the following side effects:

- In animal studies, coal tar has been shown to increase the chance of skin cancer. Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: Rare Skin irritation not present before use of this medicine skin rash Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Stinging (mild)--especially for gel and solution dosage forms Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Medotar side effects

2733. This particular disease, Psoriasis, interacts with the drug methoxsalen and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Methoxsalen may cause serious side effects. Call your

Drug Side Effects Summary

doctor at once if you have: severe skin redness within 24 hours after UVA treatment

- swelling, severe itching, or severe skin discomfort
- skin redness and swelling with blisters
- worsening of your psoriasis
- a new skin lesion, or a mole that has changed in size or color
- or blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights. Mild skin redness may occur and can last for 1 or 2 days. This is a normal side effect after UVA treatment and may not be a sign of severe sunburn. Ask your doctor if you have concerns about any redness or swelling. Common side effects of methoxsalen may include: itching, redness
- nausea
- feeling nervous
- or sleep problems (insomnia).

2734. This particular disease, Psoriasis, interacts with the drug MG217 Medicated Tar and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. MG217 Medicated Tar may cause serious side effects. Stop using MG217 Medicated Tar and call your doctor at once if you have: severe stinging, burning, swelling, or other irritation of the treated skin. Common side effects may include mild skin irritation or skin rash .

2735. This particular disease, Psoriasis, interacts with the drug Neutrogena T/Derm and has the following side effects:

- In animal studies, coal tar has been shown to increase the chance of skin cancer. Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as

Drug Side Effects Summary

possible if any of the following side effects occur: Rare Skin irritation not present before use of this medicine skin rash Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Stinging (mild)--especially for gel and solution dosage forms Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Neutrogena T/Derm side effects

2736. This particular disease, Psoriasis, interacts with the drug Neutrogena T/Gel and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Neutrogena T/Gel and call your doctor at once if you have: severe stinging, burning, swelling, or other irritation of the treated skin. Common side effects may include mild skin irritation or skin rash .

2737. This particular disease, Psoriasis, interacts with the drug Novacort and has the following side effects:

- No side effects listed

2738. This particular disease, Psoriasis, interacts with the drug Oxipor VHC and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Oxipor VHC may cause serious side effects. Stop

Drug Side Effects Summary

using Oxipor VHC and call your doctor at once if you have: severe stinging, burning, swelling, or other irritation of the treated skin. Common side effects may include mild skin irritation or skin rash .

2739. This particular disease, Psoriasis, interacts with the drug Pramoxone and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives

- itching

- red, swollen, blistered, or peeling skin with or without fever

- wheezing

- tightness in the chest or throat

- trouble breathing, swallowing, or talking

- unusual hoarseness

- or swelling of the mouth, face, lips, tongue, or throat. Signs of a weak adrenal gland like a very bad upset stomach or throwing up, very bad dizziness or passing out, muscle weakness, feeling very tired, mood changes, not hungry, or weight loss . Signs of Cushing's disease like weight gain in the upper back or belly, moon face, very bad headache , or slow healing. Signs of high blood sugar like confusion, feeling sleepy, more thirst, more hungry, passing urine more often, flushing, fast breathing, or breath that smells like fruit. Irritation where Pramoxone (pramoxine and hydrocortisone cream, gel, and ointment) is used. Skin changes (pimples , stretch marks, slow healing, hair growth). Thinning of the skin. Pramoxone side effects

2740. This particular disease, Psoriasis, interacts with the drug Proctofoam HC and has the following side effects:

Drug Side Effects Summary

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of a weak adrenal gland like a very bad upset stomach or throwing up, very bad dizziness or passing out, muscle weakness, feeling very tired, mood changes, not hungry, or weight loss . Signs of Cushing's disease like weight gain in the upper back or belly, moon face, very bad headache , or slow healing. Signs of high blood sugar like confusion, feeling sleepy, more thirst, more hungry, passing urine more often, flushing, fast breathing, or breath that smells like fruit. Irritation where Proctofoam HC (pramoxine and hydrocortisone rectal foam) is used. Very bad bleeding from rectum or rectal pain. Proctofoam HC side effects

2741. This particular disease, Psoriasis, interacts with the drug Scytera and has the following side effects

- No side effects listed

2742. This particular disease, Psoriasis, interacts with the drug Siliq and has the following side effects

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Siliq comes with a Medication Guide and a Patient Wallet Card listing symptoms of a serious side effect. Read this information and learn what symptoms to watch for. Keep the Wallet Card with you at all times. Call your doctor at once if you have: fever, chills, sweating, tiredness
- sore throat , cough, trouble breathing or swallowing
- skin sores with warmth, redness, or swelling
- stomach cramps, loss of appetite, constipation
- painful diarrhea, bloody stools, weight loss
- increased urination, pain or burning when you urinate
- unusual changes in mood or behavior
- suicidal thoughts
- or new or worsening anxiety or depression. Common Siliq side effects may include: fever, chills, flu-like symptoms
- itching, skin redness
- muscle or joint pain
- nausea , diarrhea
- headache , tiredness
- mouth pain
- or pain, redness, bruising, itching, or bleeding where the medicine was injected.

2743. This particular disease, Psoriasis, interacts with the drug Sorilux and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Sorilux and call your doctor at once if you

Drug Side Effects Summary

have a serious side effect such as: severe burning, stinging, or irritation after using this medication

- worsening of your skin condition

- or high calcium levels - confusion, tiredness, nausea, vomiting, loss of appetite, constipation, increased thirst or urination, weight loss. Less serious Sorilux side effects may include: mild skin redness

- dry or peeling skin

- or mild rash or itching.

2744. This particular disease, Psoriasis, interacts with the drug Tarsum and has the following side effects:

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Tarsum may cause serious side effects. Stop using Tarsum and call your doctor at once if you have: severe stinging, burning, swelling, or other irritation of the treated skin. Common side effects may include mild skin irritation or skin rash .

2745. This particular disease, Psoriasis, interacts with the drug Theraplex T and has the following side effects:

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Theraplex T may cause serious side effects. Stop using Theraplex T and call your doctor at once if you have: severe stinging, burning, swelling, or other irritation of the treated skin. Common side effects may include mild skin irritation or skin rash .

2746. This particular disease, Psoriasis, interacts with the drug Wyzora and has the following side effects:

- hives

- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Wyzora may cause serious side effects. Call your doctor at once if you have: worsening of your skin condition
- redness, warmth, swelling, oozing, or severe irritation of any treated skin
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- high calcium levels--confusion, tiredness, nausea , vomiting , loss of appetite, constipation , increased thirst or urination, weight loss
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor
- or possible signs of absorbing this medicine through your skin--weight gain (especially in your face or your upper back and torso), slow wound healing, thinning or discolored skin, increased body hair, muscle weakness, nausea, diarrhea , tiredness, mood changes, menstrual changes, sexual changes. Common side effects of Wyzora may include: skin color changes
- worsening of psoriasis
- skin irritation, rash, itching, redness, or hives
- redness or crusting around your hair follicles
- or headache .

2747. This particular disease, Psoriasis, interacts with the drug Yusimry and has the following side

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Blurred vision body aches or pain chills cough dizziness ear congestion fever headache loss of voice pounding in the ears slow or fast heartbeat sneezing stuffy or runny nose trouble breathing unusual tiredness or weakness Less common Abnormal healing agitation anxiety arm, back, jaw pain bleeding from the gums or nose blindness bloating or swelling of the face, arms, hands, lower legs, or feet bloody or black, tarry stools bloody or cloudy urine blue or pale skin bone pain or fractures burning, crawling,

Drug Side Effects Summary

itching, numbness, prickling, "pins and needles" , or tingling feelings chest pain, discomfort, tightness, or heaviness coma confusion cough or hoarseness dark urine decreased vision depression diarrhea difficulty in moving difficulty in swallowing drowsiness dry mouth dry skin eye pain fainting feeling of illness frequent urge to urinate gaseous stomach pain hair loss hallucinations heartburn increased thirst increased urination indigestion irritability itching loss of appetite loss of strength or energy light-colored stools lower back or side pain mood or mental changes muscle pain, stiffness, cramps, tightness, rigidity, or spasms nausea no blood pressure or pulse pain, swelling, or redness in the joints pain in the arms or legs pain in the groin or genitals pain or burning in the throat painful or difficult urination pale skin pelvic pain rapid breathing rapid weight gain or loss rash recurrent fever redness or swelling of the lower leg ringing in the ears unconsciousness seizures shakiness in the legs, arms, hands, or feet sharp back pain just below ribs sore throat sores, ulcers, or white spots on the lips or tongue or inside the mouth stiff neck stomach pain stopping of the heart sunken eyes sweating tingling of the hands or feet trembling or shaking of the hands or feet trouble thinking unpleasant breath odor unusual bleeding or bruising unusual weight gain or loss visual disturbances vomiting vomiting of blood or material that looks like coffee grounds wrinkled skin yellow eyes or skin Incidence not known Blistering, peeling, or loosening of skin blisters on the skin blue-yellow color blindness inability to move arms and legs red irritated eyes red skin lesions, often with a purple center sudden numbness and weakness in the arms and legs swelling in the arm or leg Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Bleeding, blistering, burning, coldness, discoloration of skin, feeling of pressure, hives, infection, inflammation, itching, lumps, numbness, pain, rash, redness, scarring, soreness, stinging, swelling, tenderness, tingling, ulceration, or warmth at the

Drug Side Effects Summary

injection site pain or tenderness around eyes and cheekbones Incidence not known Loss or thinning of the hair Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Yusimry side effects

2748. This particular disease, Psoriasis, interacts with the drug Zithranol and has the following side effects

- Anthralin has been shown to cause tumors in animals. However, there have been no reports of anthralin causing tumors in humans. Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: More common Redness or other skin irritation of treated or uninvolved skin not present before use of this medicine Rare Skin rash Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Zithranol side effects

2749. This particular disease, Psoriasis, interacts with the drug Zithranol-RR and has the following side effects

- Anthralin has been shown to cause tumors in animals. However, there have been no reports of anthralin causing tumors in humans. Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: More common Redness or other skin irritation of treated or uninvolved skin not present before use of this medicine Rare Skin rash Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Zithranol-RR

Drug Side Effects Summary

side effects

2750. This particular disease, Rheumatoid Arthritis, interacts with the drug Enbrel and has the follo

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Serious and sometimes fatal infections may occur. Call your doctor right away if you have signs of infection such as: fever, chills, cough, sweating, body aches, skin sores or redness, diarrhea, weight loss, burning when you urinate, coughing up blood, or feeling very tired or short of breath. Also call your doctor at once if you have: fever, chills, flu symptoms
- pale skin, easy bruising or bleeding
- pain, redness, or swelling where etanercept was injected (for longer than 5 days after injection)
- signs of lymphoma - fever, night sweats, weight loss, stomach pain or swelling, swollen glands (in your neck, armpits, or groin)
- signs of tuberculosis - cough, night sweats, loss of appetite, weight loss, feeling very tired
- new or worsening psoriasis - skin redness or scaly patches, raised bumps filled with pus
- nerve problems - dizziness, numbness or tingling, problems with vision, or weak feeling in your arms or legs
- signs of heart failure - shortness of breath, swelling in your lower legs
- lupus-like syndrome - joint pain or swelling, chest discomfort, feeling short of breath, skin rash on your cheeks or arms (worsens in sunlight)
- or liver problems - right-sided upper stomach pain, vomiting, tiredness, loss of appetite, yellowing of your skin or eyes. Common Enbrel side effects may include: pain, swelling, itching, or redness where the medicine was injected
- headache

Drug Side Effects Summary

- or cold symptoms such as stuffy nose, sneezing, sore throat.

2751. This particular disease, Rheumatoid Arthritis, interacts with the drug hydroxychloroquine and

- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out). Hydroxychloroquine may cause serious side effects. Call your doctor at once if you have: a seizure

- yellowing of your eyes

- ringing in your ears, trouble hearing

- unusual mood changes

- severe muscle weakness, loss of coordination, underactive reflexes

- any sudden changes in mood or behavior, or thoughts about suicide

- low blood cell counts - fever, chills, tiredness, sore throat, mouth sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath

- low blood sugar - headache , hunger, sweating, irritability, dizziness, fast heart rate , and feeling anxious or shaky

- or a serious drug reaction that can affect many parts of your body - skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Taking hydroxychloroquine long-term or at high doses may cause irreversible damage to the retina of your eye. Stop taking hydroxychloroquine and tell your doctor if you have: blurred vision, trouble focusing, trouble reading

- distorted vision, blind spots

- changes in your color vision

- hazy or cloudy vision

- seeing light flashes or streaks, seeing halos around lights

- or increased sensitivity to light. Common side effects of hydroxychloroquine may include:

Drug Side Effects Summary

headache

- dizziness
- nausea or vomiting
- stomach pain
- loss of appetite
- weight loss
- feeling nervous or irritable
- skin rash or itching
- or hair loss .

2752. This particular disease, Rheumatoid Arthritis, interacts with the drug Plaquenil and has the fo

- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness

(like you might pass out). Call your doctor at once if you have: a seizure

- yellowing of your eyes
- ringing in your ears, trouble hearing
- unusual mood changes
- severe muscle weakness, loss of coordination, underactive reflexes
- any sudden changes in mood or behavior, or thoughts about suicide
- low blood cell counts - fever, chills, tiredness, sore throat, mouth sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath
- low blood sugar - headache , hunger, sweating, irritability, dizziness, fast heart rate , and feeling anxious or shaky
- or a serious drug reaction that can affect many parts of your body - skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Taking Plaquenil long-term or at high doses may cause irreversible damage to the retina of your eye. Stop

Drug Side Effects Summary

taking Plaquenil and tell your doctor if you have: blurred vision, trouble focusing, trouble reading

- distorted vision, blind spots
- changes in your color vision
- hazy or cloudy vision
- seeing light flashes or streaks, seeing halos around lights
- or increased sensitivity to light. Common Plaquenil side effects may include: headache, dizziness
- nausea , vomiting, stomach pain
- loss of appetite, weight loss
- feeling nervous or irritable
- skin rash or itching
- or hair loss .

2753. This particular disease, Rheumatoid Arthritis, interacts with the drug Cimzia and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Serious and sometimes fatal infections may occur during treatment with Cimzia. Tell your doctor if you have signs of infection, such as: fever, chills, cough, sweating, muscle pain , open sores or skin wounds, unusual tiredness, feeling short of breath, painful urination, diarrhea, or weight loss . Call your doctor right away if you have any of these symptoms of lymphoma : chest pain, cough, feeling short of breath
- swelling in your neck, underarm, or groin (this swelling may come and go)
- fever, night sweats, itching, weight loss, feeling tired
- feeling full after eating only a small amount
- or pain in your upper stomach that may spread to your back or shoulder. Stop using Cimzia and call your doctor at once if you have: shortness of breath (even with mild exertion), swelling, rapid

Drug Side Effects Summary

weight gain

- pale skin, easy bruising or bleeding
- a new growth on your skin (may be red or purple), or any change in the size or color of a mole, freckle, or bump on your skin
- nerve problems - vision problems, dizziness , numbness or tingly feeling, muscle weakness in your arms or legs
- liver problems - loss of appetite, right-sided stomach pain, tiredness, jaundice (yellowing of the skin or eyes)
- or new or worsening symptoms of lupus - joint pain, and a skin rash on your cheeks or arms that worsens in sunlight. Common Cimzia side effects may include: pain or burning when you urinate
- rash
- or cold symptoms such as stuffy nose, sneezing, sore throat .

2754. This particular disease, Rheumatoid Arthritis, interacts with the drug Simponi and has the fol

- hives , itching
- nausea
- chest pain, difficulty breathing
- swelling of your face, lips, tongue, or throat. You may get infections more easily, even serious or fatal infections. Call your doctor right away if you have signs of infection such as: fever, chills, night sweats, muscle aches, feeling very tired
- cough, bloody mucus, shortness of breath
- weight loss
- skin sores with pain, warmth, or redness
- diarrhea, stomach pain
- or increased urination, or burning when you urinate. Also call your doctor at once if you have: skin

Drug Side Effects Summary

growths or changes in skin appearance

- swelling in your lower legs
- vision changes
- numbness or tingly feeling, weakness in your arms or legs
- pale skin, easy bruising or bleeding
- liver problems - right-sided upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- new or worsening symptoms of lupus - muscle or joint pain, and a skin rash on your cheeks or arms that worsens in sunlight
- or signs of psoriasis - red or scaly patches of skin, flaking, pus. Common Simponi side effects may include: infections, cold or flu symptoms
- abnormal liver function tests
- high blood pressure
- rash
- or pain, itching, redness, or swelling where the medicine was injected.

2755. This particular disease, Rheumatoid Arthritis, interacts with the drug Rinvoq and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some people taking Rinvoq have developed heart attacks, strokes, or serious blood clots. Stop taking Rinvoq and seek emergency medical attention if you have: sudden shortness of breath
- chest pain or pressure that may spread to your jaw, shoulder, arms, or back
- nausea, vomiting , cold sweat
- a light-headed feeling, like you might pass out

Drug Side Effects Summary

- weakness on one side of your body
- slurred speech, drooping on one side of your mouth
- or pain, swelling, or redness in an arm or a leg. Call your doctor at once if you have: fever, chills, night sweats, constant tiredness
- wheezing, trouble breathing, severe or worsening cough
- increased urination, pain or burning when you urinate
- unexplained weight loss
- lumps in your neck, armpits, or groin
- signs of shingles - flu-like symptoms, tingly or painful blistering rash on one side of your body
- a hole (perforation) in your digestive tract - fever, severe stomach pain, diarrhea or changes in bowel habits
- or signs of hepatitis - loss of appetite, vomiting, stomach pain (upper right side), dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common Rinvoq side effects may include: fever
- cough
- nausea
- or cold symptoms such as stuffy nose, sneezing, sore throat .

2756. This particular disease, Rheumatoid Arthritis, interacts with the drug rituximab and has the fo

- Rituximab can cause serious side effects, including: See "Important information" above. Tumor Lysis Syndrome (TLS). TLS is caused by the fast breakdown of cancer cells. TLS can cause you or your child to have: kidney failure and the need for dialysis treatment abnormal heart rhythm TLS can happen within 12 to 24 hours after an infusion of rituximab. Your healthcare provider may do blood tests to check you or your child for TLS. Your healthcare provider may give you or your child medicine to help prevent TLS. Tell your healthcare provider right away if you or your child have any

Drug Side Effects Summary

of the following signs or symptoms or TLS: nausea vomiting diarrhea lack of energy Serious infections. Serious infections can happen during and after treatment with rituximab, and can lead to death. Rituximab can increase your or your child's risk of getting infections and can lower the ability of your or your child's immune system to fight infections. Types of serious infections that can happen with rituximab include bacterial, fungal, and viral infections. After receiving rituximab, some people have developed low levels of certain antibodies in their blood for a long period of time (longer than 11 months). Some of these people with low antibody levels developed infections. People with serious infections should not receive rituximab. Tell your healthcare provider right away if you or your child have any symptoms of infection: fever cold symptoms, such as runny nose or sore throat that do not go away flu symptoms, such as cough, tiredness, and body aches earache or headache pain during urination cold sores in the mouth or throat cuts, scrapes or incisions that are red, warm, swollen or painful Heart problems. Rituximab may cause chest pain, irregular heartbeats, and heart attack. Your healthcare provider may monitor your or your child's heart during and after treatment with rituximab if you or your child have symptoms or heart problems or have a history of heart problems. Tell your healthcare provider right away if you or your child have chest pain or irregular heartbeats during treatment with rituximab. Kidney problems, especially if you or your child are receiving rituximab for NHL. Rituximab can cause severe kidney problems that lead to death. Your healthcare provider should do blood tests to check how well your or your child's kidneys are working. Stomach and Serious bowel problems that can sometimes lead to death. Bowel problems, including blockage or tears in the bowel can happen if you or your child receive rituximab with chemotherapy medicines. Tell your healthcare provider right away if you or your child have any severe stomach-area (abdomen) pain or repeated vomiting during treatment with rituximab. Your healthcare provider will stop treatment with rituximab if you have severe, serious or life-threatening side effects. The most common side effects of rituximab include: infusion-related reactions. (see "Important information") infections (may include fever, chills) body aches tiredness nausea In adult

Drug Side Effects Summary

patients with GPA or MPA the most common side effects of rituximab also include: low white and red blood cells swelling diarrhea muscle spasms In children with B-cell NHL or B-AL who receive rituximab with chemotherapy, the most common side effects include: decreased white blood cells with fever mouth sores inflammation of the upper intestine serious infection throughout the body and organs (sepsis) changes in liver function blood tests low level of potassium in the blood Other side effects with rituximab include: aching joints during or within hours of receiving an infusion more frequent upper respiratory tract infection These are not all of the possible side effects with rituximab. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Interactions Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Especially tell your doctor if you take or have taken: a Tumor Necrosis Factor (TNF) inhibitor medicine a Disease Modifying Anti-Rheumatic Drug (DMARD) If you are not sure if your medicine is one listed above, ask your healthcare provider. Rituximab drug interactions

2757. This particular disease, Rheumatoid Arthritis, interacts with the drug certolizumab and has the

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Serious and sometimes fatal infections may occur during treatment with certolizumab. Tell your doctor if you have signs of infection, such as: fever, chills, cough, sweating, muscle pain, open sores or skin wounds, unusual tiredness, feeling short of breath, painful urination, diarrhea, or weight loss. Call your doctor right away if you have any of these symptoms of lymphoma : chest pain, cough, feeling short of breath
- swelling in your neck, underarm, or groin (this swelling may come and go)
- fever, night sweats, itching, weight loss, feeling tired
- feeling full after eating only a small amount

Drug Side Effects Summary

- or pain in your upper stomach that may spread to your back or shoulder. Certolizumab may cause serious side effects. Stop using certolizumab and call your doctor at once if you have: shortness of breath (even with mild exertion), swelling, rapid weight gain
- pale skin, easy bruising or bleeding
- a new growth on your skin (may be red or purple), or any change in the size or color of a mole, freckle, or bump on your skin
- nerve problems--vision problems, dizziness , numbness or tingly feeling, muscle weakness in your arms or legs
- liver problems--loss of appetite, right-sided stomach pain, tiredness, jaundice (yellowing of the skin or eyes)
- or new or worsening symptoms of lupus --joint pain, and a skin rash on your cheeks or arms that worsens in sunlight. Common side effects of certolizumab may include: pain or burning when you urinate
- rash
- or cold symptoms such as stuffy nose, sneezing, sore throat .

2758. This particular disease, Rheumatoid Arthritis, interacts with the drug golimumab and has the

- hives , itching
- nausea
- chest pain, difficulty breathing
- swelling of your face, lips, tongue, or throat. You may get infections more easily, even serious or fatal infections. Call your doctor right away if you have signs of infection such as: fever, chills, night sweats , muscle aches, feeling very tired
- cough, bloody mucus, shortness of breath
- weight loss

Drug Side Effects Summary

- skin sores with pain, warmth, or redness
- diarrhea, stomach pain
- or increased urination, or burning when you urinate. Also call your doctor at once if you have: skin growths or changes in skin appearance
- swelling in your lower legs
- vision changes
- numbness or tingly feeling, weakness in your arms or legs
- pale skin, easy bruising or bleeding
- liver problems--right-sided upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- new or worsening symptoms of lupus --muscle or joint pain, and a skin rash on your cheeks or arms that worsens in sunlight
- or signs of psoriasis--red or scaly patches of skin, flaking, pus. Common side effects of golimumab may include: infections, cold or flu symptoms
- abnormal liver function tests
- high blood pressure
- rash
- or pain, itching, redness, or swelling where the medicine was injected.

2759. This particular disease, Rheumatoid Arthritis, interacts with the drug upadacitinib and has the

- Upadacitinib may cause serious side effects, including: See "Important information" above on this page for serious side effect information. The most common side effects for rheumatoid arthritis patients being treated with upadacitinib include: upper respiratory tract infections (common cold, sinus infections) shingles (herpes zoster) herpes simplex virus infections, including cold sores Bronchitis nausea cough fever acne The most common side effects for atopic dermatitis patients

Drug Side Effects Summary

include: upper respiratory tract infections (common cold, sinus infections) acne herpes simplex virus infections, including cold sores headache increased blood levels of creatine phosphokinase cough allergic reactions inflammation of hair follicles nausea stomach-area (abdominal) pain fever increased weight shingles (herpes zoster) flu tiredness low white blood cell count (neutropenia) muscle pain flu-like illness The most common side effects for ulcerative colitis patients include: upper respiratory tract infections (common cold, sinus infections) Acne herpes simplex virus infections, including cold sores inflammation of the hair follicles rash flu shingles (herpes zoster) increased blood cholesterol levels increased blood levels of creatine phosphokinase increased liver enzyme levels low number of certain types of white blood cells (neutropenia, lymphopenia) Separation or tear to the lining of the back part of the eye (retinal detachment) has happened in people with atopic dermatitis treated with upadacitinib. Call your healthcare provider right away if you have any sudden changes in your vision during treatment with upadacitinib. These are not all the possible side effects of upadacitinib. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Upadacitinib side effects

2760. This particular disease, Rheumatoid Arthritis, interacts with the drug leflunomide and has the

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Leflunomide may cause serious side effects. Call your doctor at once if you have: signs of infection--sudden weakness or ill feeling, fever, chills, sore throat , mouth sores, red or swollen gums, trouble swallowing
- sudden chest pain or discomfort, wheezing, dry cough, feeling short of breath
- easy bruising, unusual bleeding (nose, mouth, vagina, or rectum), purple or red pinpoint spots under your skin
- numbness , tingling, or burning pain in your hands or feet

Drug Side Effects Summary

- liver problems-- nausea , upper stomach pain, itching, tiredness, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or severe skin reaction--fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain, followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of leflunomide may include: nausea, diarrhea, stomach pain
- headache
- abnormal liver function tests
- thinning hair
- back pain
- weakness
- rash
- or high blood pressure .

2761. This particular disease, Rheumatoid Arthritis, interacts with the drug Arava and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: signs of infection
- sudden weakness or ill feeling, fever, chills, sore throat, mouth sores, red or swollen gums, trouble swallowing
- sudden chest pain or discomfort, wheezing, dry cough, feeling short of breath
- easy bruising, unusual bleeding (nose, mouth, vagina, or rectum), purple or red pinpoint spots under your skin
- numbness, tingling, or burning pain in your hands or feet
- liver problems - nausea, upper stomach pain, itching, tiredness, loss of appetite, dark urine,

Drug Side Effects Summary

clay-colored stools, jaundice (yellowing of the skin or eyes)

- or severe skin reaction - fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain, followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common Arava side effects may include: nausea, diarrhea, stomach pain
- headache
- abnormal liver function tests
- thinning hair
- back pain
- weakness
- rash
- or high blood pressure.

2762. This particular disease, Rheumatoid Arthritis, interacts with the drug Orencia and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Some side effects may occur during the injection. Tell your caregiver right away if you feel dizzy, light-headed, itchy, or have a severe headache or trouble breathing within 1 hour after receiving the injection. You may get infections more easily, even serious or fatal infections. Call your doctor right away if you have signs of infection such as: fever, chills, night sweats , flu symptoms, weight loss
- feeling very tired
- dry cough, sore throat
- or warmth, pain, or redness of your skin. Call your doctor at once if you have any of these other serious side effects: trouble breathing

Drug Side Effects Summary

- stabbing chest pain, wheezing, cough with yellow or green mucus
- pain or burning when you urinate
- or signs of skin infection such as itching, swelling, warmth, redness, or oozing. Common Orencia side effects may include: fever
- nausea , diarrhea , stomach pain
- headache
- or cold symptoms such as stuffy nose, sneezing, sore throat, cough.

2763. This particular disease, Rheumatoid Arthritis, interacts with the drug Rituxan and has the following side effects:

- hives, difficult breathing, swelling in your face or throat) or a severe skin reaction (fever, sore throat, burning eyes, skin pain, red or purple skin rash with blistering and peeling). Some side effects may occur during the injection (or within 24 hours afterward). Tell your caregiver right away if you feel itchy, dizzy, weak, light-headed, short of breath, or if you have a rash or hives, chest pain, wheezing, sudden cough, or pounding heartbeats or fluttering in your chest. Rituximab may cause a serious brain infection that can lead to disability or death. Call your doctor right away if you have any of the following symptoms (which may start gradually and get worse quickly): confusion, memory problems, or other changes in your mental state
- weakness on one side of your body
- vision changes
- or problems with speech or walking. Rituxan can lead to other serious side effects including infections, heart attacks or other heart problems, kidney failure, and intestinal blockage and/or damage. Talk to your doctor about your risk. Call your doctor at once if you have any of these other side effects, even if they occur several months after you receive Rituxan, or after your treatment ends. painful skin or mouth sores, or a severe skin rash with blistering, peeling, or pus
- redness, warmth, or swelling of the skin

Drug Side Effects Summary

- severe stomach pain, vomiting , constipation , bloody or tarry stools
- irregular heartbeats, chest pain or pressure, pain spreading to your jaw or shoulder
- tiredness or jaundice (yellowing of the skin or eyes)
- signs of infection - fever, chills, cold or flu symptoms, cough, sore throat, mouth sores, headache, earache, pain or burning when you urinate
- or signs of tumor cell breakdown - confusion, weakness, muscle cramps, nausea, vomiting, fast or slow heart rate, decreased urination, tingling in your hands and feet or around your mouth. Common Rituxan side effects may include: low white and red blood cells (fever, chills, body aches, pale skin, unusual tiredness, infections)
- nausea, diarrhea
- swelling in your hands or feet
- headache, weakness
- painful urination
- muscle spasms
- depressed mood
- or cold symptoms such as stuffy nose, sneezing, sore throat.

2764. This particular disease, Rheumatoid Arthritis, interacts with the drug Xeljanz and has the following

- Xeljanz, Xeljanz XR and Xeljanz oral solution may cause serious side effects, including: See "Important information" above. Hepatitis B or C activation infection in people who carry the virus in their blood. If you are a carrier of the hepatitis B or C virus (viruses that affect the liver), the virus may become active while you use these medications. Your healthcare provider may do blood tests before you start treatment and while you are taking the medication. Tell your healthcare provider if you have any of the following symptoms of a possible hepatitis B or C infection: feel very tired little or no appetite clay-colored bowel movements chills muscle aches skin rash skin or eyes look yellow

Drug Side Effects Summary

vomiting fevers stomach discomfort dark urine Common side effects of Xeljanz and Xeljanz XR in people with rheumatoid arthritis, psoriatic arthritis, and ankylosing spondylitis include: upper respiratory tract infections (common cold, sinus infections) headache diarrhea nasal congestion, sore throat, and runny nose (nasopharyngitis) high blood pressure (hypertension) Common side effects of Xeljanz and Xeljanz XR in people with ulcerative colitis include: nasal congestion, sore throat, and runny nose (nasopharyngitis) increased cholesterol levels headache upper respiratory tract infections (common cold, sinus infections) increased muscle enzyme levels rash diarrhea shingles (herpes zoster) Common side effects of Xeljanz and Xeljanz oral solution in people with polyarticular course juvenile arthritis include: upper respiratory tract infections (common cold, sinus infections) nasal congestion, sore throat, and runny nose (nasopharyngitis) headache fever nausea vomiting Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects. For more information, ask your healthcare provider or pharmacist. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Pfizer at 1-800-438-1985.

Xeljanz side effects

2765. This particular disease, Rheumatoid Arthritis, interacts with the drug Xeljanz XR and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing

Drug Side Effects Summary

- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Signs of liver problems like dark urine, feeling tired, not hungry, upset stomach or stomach pain, light-colored stools, throwing up, or yellow skin or eyes. Signs of infection like fever, chills, very bad sore throat , ear or sinus pain, cough, more sputum or change in color of sputum, pain with passing urine, mouth sores, or wound that will not heal. Warm, red, or painful skin or sores on the body. Shingles . Change in color or size of a mole. A skin lump or growth. Feeling very tired or weak. Change in bowel habits. Muscle pain or weakness. Slow heartbeat . A heartbeat that does not feel normal. Call your doctor right away if you have a swollen gland, night sweats , shortness of breath, or weight loss without trying. Tears in the stomach or bowel wall have happened in certain people taking Xeljanz XR (tofacitinib extended-release tablets). Call your doctor right away if you have swelling or pain in your stomach that is very bad, gets worse, or does not go away. Call your doctor right away if you throw up blood or have throw up that looks like coffee grounds
- upset stomach or throwing up that does not go away
- or black, tarry, or bloody stools. Some people have had lung problems with Xeljanz XR (tofacitinib extended-release tablets). Sometimes, this has been deadly. Call your doctor right away if you have signs of lung problems like shortness of breath or other trouble breathing, cough that is new or worse, or fever. Xeljanz XR side effects

2766. This particular disease, Rheumatoid Arthritis, interacts with the drug Simponi Aria and has th

- Serious side effects of Simponi Aria include: See "Important information" above. Serious Infections Some patients have an increased chance of getting serious infections while receiving this medication. These serious infections include TB and infections caused by viruses, fungi, or bacteria

Drug Side Effects Summary

that have spread throughout the body. Some patients die from these infections. If you get an infection while receiving treatment with this medication your doctor will treat your infection and may need to stop your treatment. Tell your doctor right away if you have any of the following signs of an infection while receiving or after receiving this medication: a fever feel very tired have a cough have flu-like symptoms warm, red, or painful skin Your doctor will examine you for TB and perform a test to see if you have TB. If your doctor feels that you are at risk for TB, you may be treated with medicine for TB before you begin treatment with Simponi Aria and during treatment with it. Even if your TB test is negative your doctor should carefully monitor you for TB infections while you are receiving this medication. People who had a negative TB skin test before receiving this medication have developed active TB. Tell your doctor if you have any of the following symptoms while receiving or after receiving Simponi Aria: cough that does not go away low grade fever weight loss loss of body fat and muscle (wasting) Hepatitis B infection in people who carry the virus in their blood. If you are a carrier of the hepatitis B virus (a virus that affects the liver), the virus can become active while you use this medication. Your doctor should do blood tests before you start treatment with and while you are receiving it. Tell your doctor if you have any of the following symptoms of a possible hepatitis B infection: feel very tired dark urine skin or eyes look yellow little or no appetite vomiting muscle aches clay-colored bowel movements fevers chills stomach discomfort skin rash Heart failure, including new heart failure or worsening of heart failure that you already have can happen in people who use TNF-blocker medicines, including this medication. If you develop new or worsening heart failure with Simponi Aria, you may need to be treated in a hospital, and it may result in death. If you have heart failure before starting this medication, your condition should be watched closely during treatment. Call your doctor right away if you get new or worsening symptoms of heart failure during treatment with this medication (such as shortness of breath or swelling of your lower legs or feet, or sudden weight gain). Nervous System Problems. Rarely, people receiving TNF-blocker medicines, including this one, have nervous system problems such as multiple

Drug Side Effects Summary

sclerosis or Guillain-Barré syndrome. Tell your doctor right away if you get any of these symptoms: vision changes weakness in your arms or legs numbness or tingling in any part of your body

Immune System Problems. Rarely, people receiving TNF-blocker medicines have developed symptoms that are like the symptoms of lupus. Tell your doctor if you have any of these symptoms: a rash on your cheeks or other parts of the body sensitivity to the sun new joint or muscle pains becoming very tired chest pain or shortness of breath swelling of the feet, ankles, or legs

Liver Problems. Liver problems can happen in people who receive TNF-blocker medicines, including this medication. These problems can lead to liver failure and death. Call your doctor right away if you have any of these symptoms: feel very tired skin or eyes look yellow poor appetite or vomiting pain on the right side of your stomach (abdomen)

Blood Problems. Low blood counts have been seen with Simponi Aria. Your body may not make enough blood cells that help fight infections or help stop bleeding. Symptoms include fever, bruising or bleeding easily, or looking pale. Your doctor will check your blood counts before and during treatment.

Allergic Reactions. Allergic reactions can happen in people who receive TNF-blocker medicines, including this medication. Some reactions may be serious and can be life-threatening. Some of these reactions can happen after receiving your first dose. Call your doctor right away if you have any of these symptoms of an allergic reaction: hives swollen face breathing trouble chest pain

The most common side effects of Simponi Aria include: upper respiratory infection (runny nose, sore throat, and hoarseness or laryngitis) abnormal liver tests decreased blood cells that fight infection viral infections, such as flu and cold sores in the mouth bronchitis high blood pressure rash

These are not all of the possible side effects of. Tell your doctor about any side effect that bothers you or does not go away. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Simponi Aria side effects

2767. This particular disease, Rheumatoid Arthritis, interacts with the drug abatacept and has the f

Drug Side Effects Summary

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Some side effects may occur during the injection. Tell your caregiver right away if you feel dizzy, light-headed, itchy, or have a severe headache or trouble breathing within 1 hour after receiving the injection. You may get infections more easily, even serious or fatal infections. Call your doctor right away if you have signs of infection such as: fever, chills, night sweats , flu symptoms, weight loss
- feeling very tired
- dry cough, sore throat
- or warmth, pain, or redness of your skin. Call your doctor at once if you have any of these other serious side effects: trouble breathing
- stabbing chest pain, wheezing, cough with yellow or green mucus
- pain or burning when you urinate
- or signs of skin infection such as itching, swelling, warmth, redness, or oozing. Common side effects of abatacept may include: fever
- nausea , diarrhea , stomach pain
- headache
- or cold symptoms such as stuffy nose, sneezing, sore throat, cough.

2768. This particular disease, Rheumatoid Arthritis, interacts with the drug auranofin and has the fo

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Auranofin may cause serious side effects. Call your doctor at once if you have: pale skin, easy bruising, unusual bleeding (nosebleeds , bleeding gums), purple or red spots under your skin

Drug Side Effects Summary

- unusual tiredness, feeling short of breath
 - severe skin rash or itching
 - sudden chest pain, wheezing, dry cough, feeling short of breath
 - severe or ongoing diarrhea, nausea , vomiting , or loss of appetite
 - black, bloody, or tarry stools
 - or kidney problems--puffy eyes, swelling in your ankles or feet, blood in your urine, urine that looks foamy. Auranofin may cause more side effects on the stomach than gold injections cause. If you were switched from gold injections to auranofin capsules, tell your doctor if you have concerns about these side effects. You may need to use auranofin for a few months before your symptoms improve.
- Common side effects of auranofin may include: diarrhea, loose stools
- rash, itching
 - or blisters or ulcers in your mouth, red or swollen gums, trouble swallowing.

2769. This particular disease, Rheumatoid Arthritis, interacts with the drug Kevzara and has the following side effects:

- hives
- chest pain, difficult breathing, feeling like you might pass out
- swelling of your face, lips, tongue, or throat. Serious and sometimes fatal infections may occur during treatment with Kevzara. Stop using this medicine and call your doctor right away if you have signs of infection such as: fever, chills, sweating, body aches
- cough with bloody mucus
- feeling short of breath
- diarrhea, stomach pain, weight loss
- sores on your skin
- pain or burning when you urinate
- or feeling very tired. Also call your doctor at once if you have signs of perforation (a hole or tear) in

Drug Side Effects Summary

your stomach or intestines : fever

- ongoing stomach pain
- or a change in bowel habits. Common Kevzara side effects may include: runny or stuffy nose, sinus pain, sore throat
- abnormal liver function tests
- painful urination
- or skin redness where an injection was given.

2770. This particular disease, Rheumatoid Arthritis, interacts with the drug Ridaura and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Ridaura may cause serious side effects. Call your doctor at once if you have: pale skin, easy bruising, unusual bleeding (nosebleeds , bleeding gums), purple or red spots under your skin
- unusual tiredness, feeling short of breath
- severe skin rash or itching
- sudden chest pain, wheezing, dry cough, feeling short of breath
- severe or ongoing diarrhea, nausea , vomiting , or loss of appetite
- black, bloody, or tarry stools
- or kidney problems--puffy eyes, swelling in your ankles or feet, blood in your urine, urine that looks foamy. Ridaura may cause more side effects on the stomach than gold injections cause. If you were switched from gold injections to this medicine capsules, tell your doctor if you have concerns about these side effects. You may need to use this medicine for a few months before your symptoms improve. Common side effects of Ridaura may include: diarrhea, loose stools
- rash, itching

Drug Side Effects Summary

- or blisters or ulcers in your mouth, red or swollen gums, trouble swallowing.

2771. This particular disease, Rheumatoid Arthritis, interacts with the drug tofacitinib and has the f

- Tofacitinib may cause serious side effects, including: See "Important information" above. Hepatitis B or C activation infection in people who carry the virus in their blood. If you are a carrier of the hepatitis B or C virus (viruses that affect the liver), the virus may become active while you use tofacitinib. Your healthcare provider may do blood tests before you start treatment with tofacitinib and while you are taking tofacitinib. Tell your healthcare provider if you have any of the following symptoms of a possible hepatitis B or C infection: feel very tired little or no appetite clay-colored bowel movements chills muscle aches skin rash skin or eyes look yellow vomiting fevers stomach discomfort dark urine Common side effects of tofacitinib tablets and tofacitinib XR in people with rheumatoid arthritis, psoriatic arthritis, and ankylosing spondylitis include: upper respiratory tract infections (common cold, sinus infections) headache diarrhea nasal congestion, sore throat, and runny nose (nasopharyngitis) high blood pressure (hypertension) Common side effects of tofacitinib tablets and tofacitinib XR in people with ulcerative colitis include: nasal congestion, sore throat, and runny nose (nasopharyngitis) increased cholesterol levels headache upper respiratory tract infections (common cold, sinus infections) increased muscle enzyme levels rash diarrhea shingles (herpes zoster) Common side effects of tofacitinib tablets and tofacitinib oral solution in people with polyarticular course juvenile arthritis include: upper respiratory tract infections (common cold, sinus infections) nasal congestion, sore throat, and runny nose (nasopharyngitis) headache fever nausea vomiting Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of tofacitinib. For more information, ask your healthcare provider or pharmacist. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Pfizer at 1-800-438-1985. Tofacitinib side effects

Drug Side Effects Summary

2772. This particular disease, Rheumatoid Arthritis, interacts with the drug alemtuzumab and has the

- hives

- difficult breathing

- swelling of your face, lips, tongue, or throat. Get emergency medical help if you have symptoms of thrombotic thrombocytopenic purpura: purple spots on the skin or in the mouth, jaundice (yellowing of the skin or eyes), tiredness, fever, feeling short of breath, fast heart rate , speech or vision changes, confusion, seizure, dark or blood in your urine, stomach pain, nausea , vomiting , or diarrhea . Alemtuzumab may cause a brain infection that can lead to disability or death. Tell your doctor if you have problems with speech, thought, vision, or muscle movement. These symptoms can get worse quickly. Some side effects may occur during or shortly after the injection. Tell your caregiver if you feel weak, feverish, chilled, dizzy, nauseated, light-headed, or have a rash, wheezing, chest pain, trouble breathing, swelling in your mouth or throat, or fast, slow, or irregular heartbeats. Alemtuzumab can cause your immune system to attack cells and organs in your body. This can lead to serious medical problems that may occur months to years after you receive Campath or Lemtrada. Call your doctor at once if you have: unusual bleeding or bruising, nosebleeds , bleeding gums, heavy menstrual periods, blood in your urine or stools, coughing up blood , problems with speech, weakness on one side, neck pain , severe headache , drooping of your face

- a mole that has changed in size or color, cough, wheezing, chest pain, feeling short of breath, coughing up blood

- an overactive immune system--fever, swollen glands, rash, feeling unsteady or less alert, trouble waking, seizure

- liver problems--loss of appetite, stomach pain (upper right side), dark urine, jaundice (yellowing of the skin or eyes)

- kidney problems--swelling in your lower legs, weight gain, loss of appetite, sudden pain in your

Drug Side Effects Summary

stomach and back, urine that looks pink/brown or foamy

- signs of infection--fever, chills, sore throat , cough, mouth sores, skin sores or blisters, tingling, burning pain, pale or yellowed skin, pain or burning when you urinate, dark urine, feeling light-headed, cold hands and feet
- signs of tuberculosis--cough, night sweats , loss of appetite, weight loss , and feeling very tired
- signs of a stroke or tear in an artery--sudden severe headache, weakness on one side of your body, drooping in your face, slurred speech
- gallbladder problems--fever, nausea, vomiting, and stomach pain
- symptoms of Autoimmune encephalitis --personality or mood changes, hallucinations, agitation , confusion, short term memory loss , movement disorders
- symptoms of Adult onset Still's disease--high fever, pain, stiffness with or without swelling in multiple joints, rash
- thyroid problems--sweating, feeling cold, fast heartbeats, feeling nervous or tired, eye swelling, weight gain or loss, constipation
- or symptoms of thyroid cancer --a lump or swelling in your neck or throat, trouble swallowing, hoarse voice, or a new cough (not caused by a cold). Common side effects of alemtuzumab may include: reactions to the injection, rash, itching, tingling, hives
- stomach pain, nausea, vomiting, diarrhea
- infections (fever, chills, runny or stuffy nose, mouth or throat pain, painful urination)
- chest pain or tightness, coughing up blood
- dizziness , tiredness, trouble sleeping
- headache, joint pain, back pain, pain in your arms or legs
- or thyroid problems, flushing (sudden warmth, redness, or tingly feeling).

2773. This particular disease, Rheumatoid Arthritis, interacts with the drug camphor / menthol / me

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using this medicine and call your doctor at once if you have severe burning, stinging, irritation, pain, swelling, rash, or blistering where the medicine was applied. Less serious side effects may be more likely, and you may have none at all.

2774. This particular disease, Rheumatoid Arthritis, interacts with the drug Cuprimine and has the f

- hives , rash
- swollen glands
- difficult breathing
- swelling of your face, lips, tongue, or throat. Cuprimine may cause serious side effects. Call your doctor at once if you have: new or worsening joint pain
- muscle weakness in your arms and legs
- muscle weakness in your face, drooping eyelids, double vision, trouble chewing or swallowing
- new or worsening cough, fever, trouble breathing
- blisters or ulcers in your mouth, red or swollen gums, trouble swallowing
- skin rash , peeling, or watery blisters
- pain or burning when you urinate, foamy or bloody urine, lower back pain
- swelling in your hands, legs, and feet
- or low blood cell counts--fever, chills, tiredness, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath. Common side effects of Cuprimine may include: decreased sense of taste
- skin changes such as wrinkling or pimples
- stomach pain, nausea , vomiting , diarrhea , loss of appetite
- numbness or tingly feeling

Drug Side Effects Summary

- ringing in your ears
- or a wound that will not heal.

2775. This particular disease, Rheumatoid Arthritis, interacts with the drug D-Penamine and has the

- hives , rash
- swollen glands
- difficult breathing
- swelling of your face, lips, tongue, or throat. D-Penamine may cause serious side effects. Call your doctor at once if you have: new or worsening joint pain
- muscle weakness in your arms and legs
- muscle weakness in your face, drooping eyelids, double vision, trouble chewing or swallowing
- new or worsening cough, fever, trouble breathing
- blisters or ulcers in your mouth, red or swollen gums, trouble swallowing
- skin rash , peeling, or watery blisters
- pain or burning when you urinate, foamy or bloody urine, lower back pain
- swelling in your hands, legs, and feet
- or low blood cell counts--fever, chills, tiredness, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath. Common side effects of D-Penamine may include: decreased sense of taste
- skin changes such as wrinkling or pimples
- stomach pain, nausea , vomiting , diarrhea , loss of appetite
- numbness or tingly feeling
- ringing in your ears
- or a wound that will not heal.

Drug Side Effects Summary

2776. This particular disease, Rheumatoid Arthritis, interacts with the drug Depen and has the following side effects:

- hives , rash
- swollen glands
- difficult breathing
- swelling of your face, lips, tongue, or throat. Depen may cause serious side effects. Call your doctor at once if you have: new or worsening joint pain
- muscle weakness in your arms and legs
- muscle weakness in your face, drooping eyelids, double vision, trouble chewing or swallowing
- new or worsening cough, fever, trouble breathing
- blisters or ulcers in your mouth, red or swollen gums, trouble swallowing
- skin rash , peeling, or watery blisters
- pain or burning when you urinate, foamy or bloody urine, lower back pain
- swelling in your hands, legs, and feet
- or low blood cell counts--fever, chills, tiredness, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath. Common side effects of Depen may include: decreased sense of taste
- skin changes such as wrinkling or pimples
- stomach pain, nausea , vomiting , diarrhea , loss of appetite
- numbness or tingly feeling
- ringing in your ears
- or a wound that will not heal.

2777. This particular disease, Rheumatoid Arthritis, interacts with the drug Erelzi and has the following side effects:

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Serious and sometimes fatal infections may occur.

Call your doctor right away if you have: fever, chills, flu symptoms

- pale skin, easy bruising or bleeding

- pain, redness, or swelling where Erelzi Prefilled Syringe was injected (for longer than 5 days after injection)

- signs of lymphoma--fever, night sweats, weight loss, stomach pain or swelling, swollen glands (in your neck, armpits, or groin)

- signs of tuberculosis--cough, night sweats, loss of appetite, weight loss, feeling very tired

- new or worsening psoriasis --skin redness or scaly patches, raised bumps filled with pus

- nerve problems-- dizziness , numbness or tingling, problems with vision, or weak feeling in your arms or legs

- signs of heart failure--shortness of breath, swelling in your lower legs

- lupus -like syndrome--joint pain or swelling, chest discomfort, feeling short of breath, skin rash on your cheeks or arms (worsens in sunlight)

- o liver problems--right-sided upper stomach pain, vomiting , tiredness, loss of appetite, yellowing of your skin or eyes. Common side effects of Erelzi Prefilled Syringe may include: pain, swelling, itching, or redness where the medicine was injected

- or cold symptoms such as stuffy nose, sneezing, sore throat .

2778. This particular disease, Rheumatoid Arthritis, interacts with the drug Eticovo and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Body aches or pain chills cough difficulty in breathing ear congestion fever headache loss of voice sneezing sore throat stuffy or runny nose unusual tiredness or weakness Less common Accumulation of pus chest pain

Drug Side Effects Summary

confusion cough producing mucus diarrhea dizziness fainting fast heartbeat frequent or painful urination hives or welts, itching, skin rash hoarseness increase in bone pain irritation itching, pain, redness, swelling, tenderness, or warmth on the skin joint pain, stiffness, or swelling lightheadedness loss of appetite nausea rapid, shallow breathing redness of the skin stomach pain swelling of the eyelids, face, lips, hands, or feet swollen, red, or tender area of infection tightness in the chest trouble with swallowing Rare Rectal bleeding severe diarrhea severe stomach pain Incidence not known Back pain, sudden and severe being forgetful black, tarry stools bleeding gums blistering, peeling, or loosening of the skin blood in the urine or stools blue-yellow color blindness blurred vision or other change in vision burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings change in size, shape, or color of existing mole dark urine decreased urine output difficulty controlling your bladder or bowels difficulty walking dilated neck veins extreme tiredness or weakness eye pain, redness, or tenderness feeling sad or depressed general feeling of discomfort, illness, or weakness high fever increased tearing irregular breathing irregular heartbeat large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or genitals light-colored stools mole that leaks fluid or bleeds muscle cramps or pain muscle weakness, sudden and progressing new mole painful blisters on the trunk of the body pale skin pinpoint red spots on the skin red skin lesions, often with a purple center red, irritated eyes seizures sensitivity of the eye or skin to light severe eye pain slurred speech sores, ulcers, or white spots on the lips or in the mouth swollen glands swollen, painful, or tender lymph glands in the neck, armpit, or groin troubled breathing with exertion unexplained bleeding or bruising upper right abdominal pain vomiting weight gain yellow eyes and skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Bleeding, blistering, burning,

Drug Side Effects Summary

coldness, discoloration of the skin, feeling of pressure, hives, infection, inflammation, itching, lumps, numbness, pain, rash, redness, scarring, soreness, stinging, swelling, tenderness, tingling, ulceration, or warmth at the injection site Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Eticovo side effects

2779. This particular disease, Rheumatoid Arthritis, interacts with the drug interferon gamma-1b and

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Interferon gamma-1b may cause serious side effects.

Call your doctor at once if you have: confusion, hallucinations

- a seizure (convulsions)
- low blood cell counts--fever, chills, flu-like symptoms, swollen gums, mouth sores, skin sores, easy bruising, unusual bleeding
- or kidney problems--little or no urination, painful or difficult urination, swelling in your feet or ankles, feeling tired or short of breath. Your doses may be delayed or reduced if you have certain side effects. Common side effects of interferon gamma-1b may include: fever, chills
- diarrhea
- headache
- feeling tired
- rash
- or redness or tenderness where the medicine was injected.

2780. This particular disease, Rheumatoid Arthritis, interacts with the drug penicillamine and has th

Drug Side Effects Summary

- hives , rash
- swollen glands
- difficult breathing
- swelling of your face, lips, tongue, or throat. Penicillamine may cause serious side effects. Call your doctor at once if you have: new or worsening joint pain
- muscle weakness in your arms and legs
- muscle weakness in your face, drooping eyelids, double vision, trouble chewing or swallowing
- new or worsening cough, fever, trouble breathing
- blisters or ulcers in your mouth, red or swollen gums, trouble swallowing
- skin rash , peeling, or watery blisters
- pain or burning when you urinate, foamy or bloody urine, lower back pain
- swelling in your hands, legs, and feet
- or low blood cell counts--fever, chills, tiredness, skin sores, easy bruising, unusual bleeding, pale skin, cold hands and feet, feeling light-headed or short of breath. Common side effects of penicillamine may include: decreased sense of taste
- skin changes such as wrinkling or pimples
- stomach pain, nausea , vomiting , diarrhea , loss of appetite
- numbness or tingly feeling
- ringing in your ears
- or a wound that will not heal.

2781. This particular disease, Rheumatoid Arthritis, interacts with the drug Riabni and has the follo

- confusion, memory problems, or other changes in your mental state
- weakness on one side of your body
- vision changes

Drug Side Effects Summary

- or problems with speech or walking. Call your doctor at once if you have any of these other side effects, even if they occur several months after you receive Riabni, or after your treatment ends.

painful skin or mouth sores, or a severe skin rash with blistering, peeling, or pus

- redness, warmth, or swelling of the skin

- severe stomach pain, vomiting , constipation , bloody or tarry stools

- irregular heartbeats, chest pain or pressure, pain spreading to your jaw or shoulder

- tiredness or jaundice (yellowing of the skin or eyes)

- signs of infection--fever, chills, cold or flu symptoms, cough, sore throat, mouth sores, headache , earache, pain or burning when you urinate

- or signs of tumor cell breakdown--confusion, weakness, muscle cramps, nausea , vomiting, fast or slow heart rate , decreased urination, tingling in your hands and feet or around your mouth.

Common side effects of Riabni may include: low white and red blood cells (fever, chills, body aches, pale skin, unusual tiredness, infections)

- nausea, diarrhea

- swelling in your hands or feet

- headache, weakness

- painful urination

- muscle spasms

- depressed mood

- or cold symptoms such as stuffy nose, sneezing, sore throat.

2782. This particular disease, Rheumatoid Arthritis, interacts with the drug sarilumab and has the following side effects:

- hives

- chest pain, difficult breathing, feeling like you might pass out

- swelling of your face, lips, tongue, or throat. Serious and sometimes fatal infections may occur

Drug Side Effects Summary

during treatment with sarilumab. Stop using sarilumab and call your doctor right away if you have signs of infection such as: fever, chills, sweating, body aches

- cough with bloody mucus
- feeling short of breath
- diarrhea, stomach pain, weight loss
- sores on your skin
- pain or burning when you urinate
- or feeling very tired. Also call your doctor at once if you have signs of perforation (a hole or tear) in your stomach or intestines : fever
- ongoing stomach pain
- or a change in bowel habits. Common side effects of sarilumab may include: runny or stuffy nose, sinus pain, sore throat
- abnormal liver function tests
- painful urination
- or skin redness where an injection was given.

2783. This particular disease, Schizophrenia, interacts with the drug clozapine and has the following

- fever, flu-like symptoms, extreme weakness
- mouth sores, skin sores
- new or worsening cough, trouble breathing
- pain or burning when you urinate
- or vaginal itching or discharge. Further doses may be delayed until your infection clears up. High doses or long-term use of clozapine can cause a serious movement disorder that may not be reversible. The longer you use clozapine, the more likely you are to develop this disorder, especially if you are a woman or an older adult. Clozapine can increase your risk of seizure, especially at high

Drug Side Effects Summary

doses. Avoid any activity that could be dangerous if you have a seizure or lose consciousness. Call your doctor at once if you have: uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)

- a seizure (blackout-out or convulsions)
- severe constipation
- dry or hard bowel movements, or painful gas
- nausea , vomiting , stomach pain or bloating
- heart problems - chest pain, fast or pounding heartbeats, fluttering in your chest, slow heartbeats, shortness of breath, and sudden dizziness (like you might pass out)
- liver problems - loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out
- or signs of a blood clot in the lung - chest pain, sudden cough, wheezing, rapid breathing, coughing up blood . Untreated constipation may lead to serious bowel complications, hospitalization, or death. Tell your doctor right away if you are not having bowel movements at least 3 times per week. Common clozapine side effects may include: weight gain
- dizziness, tremor
- fast heart rate
- headache , drowsiness
- nausea, constipation
- dry mouth, or increased salivation
- vision problems
- or fever, increased sweating.

Drug Side Effects Summary

2784. This particular disease, Schizophrenia, interacts with the drug Fanapt and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. High doses or long-term use of Fanapt can cause a serious movement disorder that may not be reversible. The longer you use iloperidone, the more likely you are to develop this disorder, especially if you are an older adult. To make sure this medicine is safe for you, tell your doctor if you have: fast or pounding heartbeats, fluttering in your chest
- a light-headed feeling, like you might pass out
- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- tremors, slow muscle movement, muscles pain or stiffness
- confusion, agitation , thoughts of hurting yourself
- loss of bladder control
- penis erection that is painful or lasts 4 hours or longer
- high blood sugar - increased thirst, increased urination, dry mouth, fruity breath odor
- low white blood cell counts - fever, chills, sore throat , mouth sores, skin sores, cough, trouble breathing
- or severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out
- Older adults may be more likely to have side effects from this medicine. Common Fanapt side effects may include: weight gain
- dizziness, drowsiness, tired feeling
- dry mouth, stuffy nose
- or fast heart rate .

Drug Side Effects Summary

2785. This particular disease, Schizophrenia, interacts with the drug Aristada and has the following

- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- feeling uncomfortably warm or hot
- trouble swallowing
- a seizure
- severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out
- low blood cell counts - sudden weakness or ill feeling, fever, chills, sore throat, painful mouth sores, swollen gums, skin sores, cold or flu symptoms, cough, trouble breathing
- or high blood sugar - increased thirst, increased urination, hunger, dry mouth, fruity breath odor. You may have increased sexual urges, unusual urges to gamble, or other intense urges while using this medicine. Talk with your doctor if this occurs. Common Aristada side effects may include: feeling restless
- pain where the injection was given
- weight gain
- or drowsiness.

2786. This particular disease, Schizophrenia, interacts with the drug Clozaril and has the following

- fever, flu-like symptoms, extreme weakness
- mouth sores, skin sores
- new or worsening cough, trouble breathing
- pain or burning when you urinate
- or vaginal itching or discharge. Further doses may be delayed until your infection clears up. High doses or long-term use of Clozaril can cause a serious movement disorder that may not be

Drug Side Effects Summary

reversible. The longer you use Clozaril, the more likely you are to develop this disorder, especially if you are a woman or an older adult. Clozapine can increase your risk of seizure, especially at high doses. Avoid any activity that could be dangerous if you have a seizure or lose consciousness. Call your doctor at once if you have: uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)

- a seizure (blackout-out or convulsions)
- severe constipation
- dry or hard bowel movements, or painful gas
- nausea , vomiting , stomach pain or bloating
- heart problems - chest pain, fast or pounding heartbeats, fluttering in your chest, slow heartbeats, shortness of breath, and sudden dizziness (like you might pass out)
- liver problems - loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out
- or signs of a blood clot in the lung - chest pain, sudden cough, wheezing, rapid breathing, coughing up blood . Untreated constipation may lead to serious bowel complications, hospitalization, or death. Tell your doctor right away if you are not having bowel movements at least 3 times per week. Common Clozaril side effects may include: weight gain
- dizziness, tremor
- fast heart rate
- headache , drowsiness
- nausea, constipation
- dry mouth, or increased salivation
- vision problems

Drug Side Effects Summary

- or fever, increased sweating.

2787. This particular disease, Schizophrenia, interacts with the drug Invega and has the following s

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Stop taking Invega and call your doctor at once if you have any of these signs of a serious movement disorder: tremors or shaking in your arms or legs
- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- or any new or unusual muscle movements you cannot control. Call your doctor at once if you have: fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- breast swelling (in women or men), nipple discharge
- changes in menstrual periods
- impotence , penis erection that is painful or lasts 4 hours or longer
- weight gain
- low white blood cell counts - fever, chills, mouth sores, skin sores, sore throat , cough, trouble breathing
- high blood sugar - increased thirst, increased urination, hunger, fruity breath odor
- or severe nervous system reaction - very stiff (rigid) muscles, high fever, fast or pounding heartbeats, fainting. Common Invega side effects may include: drowsiness
- anxiety
- muscle stiffness, tremors or shaking
- uncontrolled muscle movements, trouble with walking, balance, or speech
- weight gain

Drug Side Effects Summary

- upset stomach , constipation
- fast heart rate
- or stuffy nose, sore throat.

2788. This particular disease, Schizophrenia, interacts with the drug chlorpromazine and has the fo

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Long-term use of chlorpromazine can cause a serious movement disorder that may not be reversible. The longer you use chlorpromazine, the more likely you are to develop this disorder, especially if you are a an older adult. Chlorpromazine may cause serious side effects. Call your doctor at once if you have: uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- stiffness in your neck, tightness in your throat, trouble breathing or swallowing
- a light-headed feeling, like you might pass out
- confusion, agitation , feeling jittery, trouble sleeping
- weakness
- breast swelling or discharge
- a seizure
- jaundice (yellowing of the skin or eyes)
- low white blood cell counts--fever, chills, mouth sores, skin sores, sore throat , cough, trouble breathing, feeling light-headed
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Some side effects may be more likely in older adults. Common side effects of chlorpromazine may include: drowsiness
- dry mouth or stuffy nose

Drug Side Effects Summary

- blurred vision
- constipation
- or impotence , trouble having an orgasm.

2789. This particular disease, Schizophrenia, interacts with the drug Invega Sustenna and has the f

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tell your doctor right away if you have any of these signs of a serious movement disorder: tremors or shaking in your arms or legs
- uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- or any new or unusual muscle movements you cannot control. Call your doctor at once if you have: fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)
- trouble swallowing
- a seizure (convulsions)
- breast swelling (in women or men), nipple discharge
- changes in menstrual periods
- impotence , or penis erection that is painful or lasts 4 hours or longer
- weight gain
- fever, chills, sore throat , mouth sores, feeling light-headed
- high blood sugar - increased thirst, increased urination, hunger, fruity breath odor
- or severe nervous system reaction - very stiff (rigid) muscles, high fever, fast or pounding heartbeats, fainting. Common Invega Sustenna side effects may include: dizziness, drowsiness
- cold symptoms such as stuffy nose, sneezing, sore throat

Drug Side Effects Summary

- feeling restless or anxious
- muscle stiffness, tremors or shaking
- uncontrolled muscle movements, trouble with walking, balance, or speech
- abnormal movements of your eyes
- weight gain
- upset stomach , constipation
- fast heart rate
- or pain or tenderness where the medicine was injected.

2790. This particular disease, Schizophrenia, interacts with the drug loxapine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. High doses or long-term use of loxapine can cause a serious movement disorder that may not be reversible. The longer you use loxapine, the more likely you are to develop this disorder, especially if you are a woman or an older adult. Loxapine may cause serious side effects. Call your doctor at once if you have: uncontrolled muscle movements in your arms or legs, or your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- fast heart rate
- a light-headed feeling, like you might pass out
- confusion, slurred speech
- agitation , trouble sleeping
- seizure (convulsions)
- little or no urinating
- severe constipation

Drug Side Effects Summary

- low white blood cell counts--fever, chills, mouth sores, skin sores, sore throat , cough, trouble breathing, feeling light-headed
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Common side effects of loxapine may include: dizziness, problems with balance or walking
- swelling in your face
- itching or rash
- tremors, muscle twitching or stiffness
- numbness , weakness
- blurred vision
- feeling restless or agitated
- nausea , vomiting , constipation
- dry mouth, stuffy nose
- or insomnia .

2791. This particular disease, Schizophrenia, interacts with the drug Zyprexa Relprevv and has the

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash , fever, swollen glands, flu-like symptoms, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Zyprexa Relprevv can cause serious symptoms if the medicine gets into your bloodstream too fast. During the first 3 hours after your injection you will be watched for the following signs: severe dizziness, drowsiness, or weakness
- confusion, anxiety , feeling angry or hostile

Drug Side Effects Summary

- feeling nervous or shaky
- trouble walking or talking
- seizure (convulsions)
- or a light-headed feeling, like you might pass out. Call your doctor at once if you have: twitching or uncontrollable movements of your eyes, lips, tongue, face, arms, or legs
- trouble speaking or swallowing
- signs of dehydration - if you feel very thirsty or hot, are unable to urinate, and have heavy sweating or hot and dry skin
- high blood sugar - increased thirst, increased urination, hunger, dry mouth, fruity breath odor
- or severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors. Common Zyprexa side effects may include: headache , back pain
- weight gain (more likely in teenagers), increased appetite
- drowsiness
- dry mouth, nausea , vomiting , diarrhea
- sinus pain, runny or stuffy nose, cough
- or pain where the injection was given.

2792. This particular disease, Schizophrenia, interacts with the drug Invega Trinza and has the following

- Invega Trinza may cause serious side effects, including: See "Important information" Stroke in elderly people (cerebrovascular problems) that can lead to death Neuroleptic Malignant Syndrome (NMS). NMS is a rare but very serious problem that can happen in people who receive Invega Trinza. NMS can cause death and must be treated in a hospital. Call your healthcare provider right away if you become severely ill and have any of these symptoms: high fever severe muscle stiffness confusion loss of consciousness changes in your breathing, heartbeat and blood pressure problems with your heartbeat. These heart problems can cause death. Call your healthcare provider right

Drug Side Effects Summary

away if you have any of these symptoms: passing out or feeling like you will pass out dizziness feeling as if your heart is pounding or missing beats uncontrolled movements of your tongue, face, mouth, or jaw (tardive dyskinesia) metabolic changes. Metabolic changes may include high blood sugar (hyperglycemia), diabetes mellitus and changes in the fat levels in your blood (dyslipidemia), and weight gain. low blood pressure and fainting changes in your blood cell counts high level of prolactin in your blood (hyperprolactinemia). Invega Trinza may cause a rise in the blood levels of a hormone called prolactin (hyperprolactinemia) that may cause side effects including missed menstrual periods, leakage of milk from the breasts, development of breasts in men, or problems with erection. problems thinking clearly and moving your body seizures difficulty swallowing that can cause food or liquid to get into your lungs prolonged or painful erection lasting more than 4 hours. Call your healthcare provider or go to your nearest emergency room right away if you have an erection that lasts more than 4 hours. problems with control of your body temperature especially when you exercise a lot or spend time doing things that make you warm. It is important for you to drink water to avoid dehydration. The most common side effects of Invega Trinza include: injection site reactions, weight gain, headache, upper respiratory tract infections, feeling restlessness or difficulty sitting still, slow movements, tremors, stiffness and shuffling walk. Tell your healthcare provider if you have any side effect that bothers you or does not go away. These are not all the possible side effects of Invega Trinza. For more information, ask your healthcare provider or pharmacist. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Invega Trinza side effects

2793. This particular disease, Schizophrenia, interacts with the drug thioridazine and has the follow

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. High doses or long-term use of thioridazine can cause

Drug Side Effects Summary

a serious movement disorder that may not be reversible. The longer you use thioridazine, the more likely you are to develop this disorder, especially if you are a woman or an older adult. Thioridazine may cause serious side effects. Call your doctor at once if you have: uncontrolled muscle movements in your arms, legs, or face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)

- fast or pounding heartbeats, fluttering in your chest, shortness of breath, and sudden dizziness (like you might pass out)

- agitation , confusion

- seizure (convulsions)

- decreased night vision, tunnel vision, watery eyes , increased sensitivity to light

- little or no urinating

- low white blood cell counts--fever, chills, mouth sores, skin sores, sore throat , cough, trouble breathing, feeling light-headed

- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Common side effects of thioridazine may include: drowsiness

- dry mouth, blurred vision

- nausea , vomiting , constipation , diarrhea

- breast swelling or discharge

- changes in your menstrual periods

- or swelling in your hands or feet.

2794. This particular disease, Schizophrenia, interacts with the drug Aristada Initio and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your

Drug Side Effects Summary

doctor or nurse immediately if any of the following side effects occur: More common Difficulty with speaking drooling loss of balance control muscle trembling, jerking, or stiffness restlessness shuffling walk stiffness of the limbs twisting movements of the body uncontrolled movements, especially of the face, neck, and back Less common Blurred vision dizziness fast, pounding, or irregular heartbeat or pulse headache inability to move the eyes increased blinking or spasms of the eyelid nervousness pounding in the ears sticking out the tongue trouble breathing or swallowing unusual facial expressions Rare Fast heartbeat high fever increased sweating lip smacking or puckering loss of bladder control muscle spasm or jerking of the arms or legs puffing of the cheeks rapid or worm-like movements of the tongue seizures severe muscle stiffness sudden loss of consciousness tiredness uncontrolled chewing movements uncontrolled movements of the arms and legs unusually pale skin Incidence not known Hives or welts, itching, or skin rash itching, puffiness, or swelling of the eyelids or around the eyes, face, lips, or tongue large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or genitals redness of the skin tightness in the chest unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Belching deep or fast breathing with dizziness difficulty having a bowel movement dry mouth fear fever headache heartburn inability to sit still indigestion irritability lightheadedness need to keep moving numbness of the feet, hands, and around the mouth runny nose shaking sore throat stomach discomfort, upset, or pain trouble sleeping weight gain Less common Accidental injury bloating or swelling of the face, arms, hands, lower legs, or feet body aches or pain congestion coughing difficulty with moving hoarseness increased appetite joint pain muscle ache, cramps, pain, or stiffness sneezing stuffy nose swollen joints tender, swollen glands in the neck tingling of the hands or feet tremor unusual

Drug Side Effects Summary

weight gain or loss voice changes Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Aristada Initio side effects

2795. This particular disease, Schizophrenia, interacts with the drug FazaClo and has the following

- fever, flu-like symptoms, extreme weakness
- mouth sores, skin sores
- new or worsening cough, trouble breathing
- pain or burning when you urinate
- or vaginal itching or discharge. Further doses may be delayed until your infection clears up. High doses or long-term use of FazaClo can cause a serious movement disorder that may not be reversible. The longer you use clozapine, the more likely you are to develop this disorder, especially if you are a woman or an older adult. Clozapine can increase your risk of seizure, especially at high doses. Avoid any activity that could be dangerous if you have a seizure or lose consciousness. Call your doctor at once if you have: uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- a seizure (blackout-out or convulsions)
- severe constipation
- dry or hard bowel movements, or painful gas
- nausea , vomiting , stomach pain or bloating
- heart problems - chest pain, fast or pounding heartbeats, fluttering in your chest, slow heartbeats, shortness of breath, and sudden dizziness (like you might pass out)
- liver problems - loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)

Drug Side Effects Summary

- severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out
- or signs of a blood clot in the lung - chest pain, sudden cough, wheezing, rapid breathing, coughing up blood . Untreated constipation may lead to serious bowel complications, hospitalization, or death. Tell your doctor right away if you are not having bowel movements at least 3 times per week. Common FazaClo side effects may include: weight gain
- dizziness, tremor
- fast heart rate
- headache , drowsiness
- nausea, constipation
- dry mouth, or increased salivation
- vision problems
- or fever, increased sweating.

2796. This particular disease, Schizophrenia, interacts with the drug Perseris and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Perseris may cause serious side effects. Call your doctor at once if you have: uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- breast swelling or tenderness (in men or women), nipple discharge, impotence , lack of interest in sex, missed menstrual periods
- severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out
- high blood sugar--increased thirst, increased urination, dry mouth, fruity breath odor

Drug Side Effects Summary

- low blood cell counts--fever, mouth sores, skin sores, sore throat , cough, easy bruising, unusual bleeding, trouble breathing, feeling light-headed
- or penis erection that is painful or lasts 4 hours or longer. Serious side effects may be more likely in older adults. Common side effects of Perseris may include: headache
- dizziness, drowsiness, tired feeling
- tremors, twitching or uncontrollable muscle movements
- depressed mood, agitation , anxiety , restless feeling
- muscle or joint pain
- dry mouth, upset stomach , constipation
- weight gain
- or pain in your arms or legs.

2797. This particular disease, Schizophrenia, interacts with the drug thiothixene and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. High doses or long-term use of thiothixene can cause a serious movement disorder that may not be reversible. The longer you use thiothixene, the more likely you are to develop this disorder, especially if you are a woman or an older adult. Thiothixene may cause serious side effects. Call your doctor at once if you have: uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- stiffness in your neck, tightness in your throat, trouble breathing or swallowing
- seizure (convulsions)
- a light-headed feeling, like you might pass out
- severe constipation

Drug Side Effects Summary

- low white blood cell counts--fever, chills, mouth sores, skin sores, sore throat , cough, trouble breathing
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Common side effects of thiothixene may include: dry mouth
- blurred vision
- nausea , vomiting , diarrhea , constipation
- fast heartbeats, feeling restless
- breast swelling or discharge
- changes in weight or appetite
- or swelling in your hands or feet.

2798. This particular disease, Schizophrenia, interacts with the drug amisulpride and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Amisulpride may cause serious side effects. Call your doctor at once if you have: sudden dizziness (like you might pass out)
- fast or pounding heartbeats, fluttering in your chest
- shortness of breath
- or low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of amisulpride may include: low potassium
- feeling light-headed
- stomach bloating
- or pain where the medicine was injected.

Drug Side Effects Summary

2799. This particular disease, Schizophrenia, interacts with the drug Versacloz and has the following

- fever, flu-like symptoms, extreme weakness
- mouth sores, skin sores
- new or worsening cough, trouble breathing
- pain or burning when you urinate
- or vaginal itching or discharge. Further doses may be delayed until your infection clears up. High doses or long-term use of clozapine can cause a serious movement disorder that may not be reversible. The longer you use Versacloz, the more likely you are to develop this disorder, especially if you are a woman or an older adult. Versacloz can increase your risk of seizure, especially at high doses. Avoid any activity that could be dangerous if you have a seizure or lose consciousness. Versacloz may cause serious side effects. Call your doctor at once if you have: uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- a seizure (blackout-out or convulsions)
- severe constipation
- dry or hard bowel movements, or painful gas
- nausea , vomiting , stomach pain or bloating
- heart problems--chest pain, fast or pounding heartbeats, fluttering in your chest, slow heartbeats, shortness of breath, and sudden dizziness (like you might pass out)
- liver problems--loss of appetite, stomach pain (upper right side), tiredness, itching, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out
- or signs of a blood clot in the lung--chest pain, sudden cough, wheezing, rapid breathing, coughing up blood . Untreated constipation may lead to serious bowel complications, hospitalization, or death.

Drug Side Effects Summary

Tell your doctor right away if you are not having bowel movements at least 3 times per week.

Common side effects of Versacloz may include: weight gain

- dizziness, tremor
- fast heart rate
- headache , drowsiness
- nausea, constipation
- dry mouth, or increased salivation
- vision problems
- or fever, increased sweating.

2800. This particular disease, Schizophrenia, interacts with the drug Adasuve and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Tell your caregivers right away if you have: a light-headed feeling, like you might pass out
- blurred vision, tunnel vision, eye pain , or seeing halos around lights
- bronchospasm (wheezing, cough, chest tightness, trouble breathing)
- little or no urination
- a seizure (convulsions)
- or severe nervous system reaction - very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Common Adasuve side effects may include: drowsiness
- sore throat
- or an unusual or unpleasant taste in the mouth.

Drug Side Effects Summary

2801. This particular disease, Schizophrenia, interacts with the drug Secuado and has the following

- hives
- fast heartbeats, feeling light-headed
- wheezing, difficult breathing
- swelling of your face, lips, tongue, or throat. High doses or long-term use of asenapine can cause a serious movement disorder that may not be reversible. The longer you use Secuado, the more likely you are to develop this disorder, especially if you are a woman or an older adult. Secuado may cause serious side effects. Call your doctor at once if you have: uncontrolled muscle movements in your face (chewing, lip smacking, frowning, tongue movement, blinking or eye movement)
- a light-headed feeling, like you might pass out
- breast pain or swelling, nipple discharge
- low white blood cell counts--fever, chills, mouth sores, skin sores, sore throat , cough, trouble breathing
- or severe nervous system reaction--very stiff (rigid) muscles, high fever, sweating, confusion, fast or uneven heartbeats, tremors, feeling like you might pass out. Common side effects of Secuado may include: muscle stiffness, jerky muscle movements
- weight gain
- or skin pain, redness, itching, swelling, or other irritation where the patch was worn.

2802. This particular disease, Schizophrenia, interacts with the drug Invega Hafyera and has the fol

- Invega Hafyera may cause serious side effects, including: See Important information. Cerebrovascular problems (including stroke) in elderly people with dementia-related psychosis that can lead to death. Neuroleptic Malignant Syndrome (NMS), a serious condition that can lead to death. Call your healthcare provider or go to your nearest hospital emergency room right away if you have some or all of the following signs and symptoms of NMS: high fever confusion changes in your

Drug Side Effects Summary

breathing, pulse, heart rate, and blood pressure stiff muscles sweating Problems with your heartbeat. These heart problems can cause death. Call your healthcare provider right away if you have any of these symptoms: passing out or feeling like you will pass out dizziness feeling as if your heart is pounding or missing beats Uncontrolled body movements (tardive dyskinesia). Invega Hafyera may cause movements that you cannot control in your face, tongue, or other body parts. Tardive dyskinesia may not go away, even if you stop treatment. Tardive dyskinesia may also start after you stop treatment. Problems with your metabolism such as high blood sugar (hyperglycemia) and diabetes. Increases in blood sugar can happen in some people who receive Invega Hafyera. Extremely high blood sugar can lead to coma or death. Your healthcare provider should check your blood sugar before you start and regularly during treatment. Call your healthcare provider if you have any of these symptoms of high blood sugar during treatment: feel very thirsty need to urinate more than usual feel very hungry feel weak or tired feel sick to your stomach feel confused, or your breath smells fruity increased fat levels (cholesterol and triglycerides) in your blood. Your healthcare provider should check the fat levels in your blood before you start and regularly during treatment. weight gain. You and your healthcare provider should check your weight before you start and often during treatment. Decreased blood pressure (orthostatic hypotension) and fainting. You may feel lightheaded or faint when you rise too quickly from a sitting or lying position, especially early in treatment or when the dose is changed. Falls. Invega Hafyera may make you sleepy or dizzy, may cause a decrease in your blood pressure when changing position (orthostatic hypotension), and can slow your thinking and motor skills which may lead to falls that can cause fractures or other injuries. Low white blood cell count. Your healthcare provider may do blood tests during the first few months of treatment. Increased prolactin levels in your blood (hyperprolactinemia). Invega Hafyera may cause a rise in the blood levels of a hormone called prolactin (hyperprolactinemia) that may cause side effects including missed menstrual periods, a reversible reduction in fertility in females who are able to become pregnant, leakage of milk from the breasts, development of breasts in men, or

Drug Side Effects Summary

problems with erection. Invega Hafyera can make you sleepy or dizzy, and can slow your thinking and motor skills. Do not drive, operate heavy machinery, or do other dangerous activities until you know how Invega Hafyera affects you. Seizures (convulsions). Difficulty swallowing that can cause food or liquid to get into your lungs. Prolonged or painful erection lasting more than 4 hours (priapism). Call your healthcare provider or go to your nearest emergency room right away if you have an erection that lasts more than 4 hours. Problems controlling your body temperature so that you feel too warm. See, What should I avoid while receiving Invega Hafyera? The most common side effects include: upper respiratory tract infections weight gain feeling restless or difficulty sitting still tremors shuffling walk injection site reactions headache slow movements stiffness These are not all the possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800 FDA-1088. Invega Hafyera side effects

2803. This particular disease, Seizures, interacts with the drug Keppra and has the following side effects

- mood or behavior changes, depression, anxiety , panic attacks , trouble sleeping, or if you feel agitated, hostile, irritable, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: unusual changes in mood or behavior (unusual risk-taking behavior, being irritable or talkative)
- confusion, hallucinations, extreme drowsiness, feeling very weak or tired
- loss of balance or coordination, problems with walking or movement
- a skin rash, no matter how mild
- easy bruising, unusual bleeding
- or fever, chills, weakness, or other signs of infection. Common Keppra side effects may include: dizziness, drowsiness, tiredness, weakness
- feeling aggressive or irritable
- loss of appetite

Drug Side Effects Summary

- stuffy nose
- or infection.

2804. This particular disease, Seizures, interacts with the drug Diastat and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Less common Anxiety blurred vision changes in patterns and rhythms of speech confusion cough crying delusions dementia depersonalization difficulty breathing difficulty in speaking dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position dry mouth dysphoria false or unusual sense of well-being feeling of warmth or heat flushing or redness of the skin, especially on the face and neck headache irregular heartbeat irritability lack of coordination mental depression mood or mental changes nervousness noisy breathing paranoia quick to react or overreact emotionally rapidly changing moods restlessness seizures shakiness and unsteady walk slurred speech sweating tightness in chest trouble breathing trouble in speaking trouble sleeping unsteadiness, trembling, or other problems with muscle control or coordination unusual tiredness or weakness Rare Bladder pain bloody or cloudy urine difficult, burning, or painful urination fever or chills frequent urge to urinate increase in body movements lower back or side pain painful or difficult urination pale skin swollen, painful, or tender lymph glands in the neck, armpit, or groin unusual bleeding or bruising Symptoms of overdose Change or loss of consciousness sleepiness or unusual drowsiness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Diarrhea feeling of constant movement of self or surroundings

Drug Side Effects Summary

hiccups lack or loss of strength rash runny or stuffy nose sensation of spinning sneezing Rare Bigger, dilated, or enlarged pupils (black part of eye) increased sensitivity of the eyes to light itching skin loss of appetite vomiting weight loss Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

Diastat side effects

2805. This particular disease, Seizures, interacts with the drug Dilantin and has the following side effects:

- skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: slow or uneven heartbeats, chest pain, fluttering in your chest, and dizziness (like you might pass out)
- any skin rash, no matter how mild
- fever, chills, sore throat, swollen glands
- red or swollen gums, mouth sores
- easy bruising, unusual bleeding, purple or red spots under your skin
- or liver problems - loss of appetite, upper stomach pain, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common Dilantin side effects may include: drowsiness, confusion
- slurred speech
- abnormal eye movement
- or problems with balance, coordination, or muscle movement.

2806. This particular disease, Seizures, interacts with the drug Trokendi XR and has the following side effects:

Drug Side Effects Summary

- mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: a skin rash, no matter how mild
- vision problems, blurred vision, eye pain or redness, sudden vision loss (can be permanent if not treated quickly)
- confusion, problems with thinking or memory, trouble concentrating, problems with speech
- dehydration symptoms--decreased sweating, high fever, hot and dry skin
- signs of a kidney stone--severe pain in your side or lower back, painful or difficult urination
- signs of too much acid in your blood--irregular heartbeats, feeling tired, loss of appetite, trouble thinking, feeling short of breath
- or signs of too much ammonia in your blood-- vomiting , unexplained weakness, feeling like you might pass out. Common Trokendi XR side effects may include include: dizziness , drowsiness, tiredness, slow reactions
- problems with speech or memory, abnormal vision
- numbness or tingling in your arms and legs, decreased sensation (especially in the skin)
- changes in your sense of taste, feeling nervous
- nausea , diarrhea, stomach pain, loss of appetite
- fever, weight loss
- or cold symptoms such as stuffy nose, sneezing, sore throat.

2807. This particular disease, Seizures, interacts with the drug Valtoco and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Diazepam can slow or stop your breathing, especially

Drug Side Effects Summary

if you have recently used an opioid medication or alcohol. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Call your doctor at once if the person receiving this medicine has: extreme weakness or drowsiness

- unusual changes in mood or behavior
- new or worsening symptoms of depression or anxiety
- thoughts of suicide or hurting yourself
- sleep problems
- or new or worsening seizures. Drowsiness or dizziness may last longer in older adults. Use caution to avoid falling or accidental injury. Common Valtoco side effects may include: drowsiness
- headache
- or discomfort in your nose. After you stop using Valtoco, get medical help right away if you have symptoms such as: unusual muscle movements, being more active or talkative, sudden and severe changes in mood or behavior, confusion, hallucinations, seizures, suicidal thoughts or actions. Some withdrawal symptoms may last up to 12 months or longer after stopping this medicine suddenly. Tell your doctor if you have ongoing anxiety, depression, problems with memory or thinking, trouble sleeping, ringing in your ears, a burning or prickly feeling, or a crawling sensation under your skin.

2808. This particular disease, Seizures, interacts with the drug Vimpat and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash , fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, depression,

Drug Side Effects Summary

anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: a light-headed feeling, like you might pass out

- severe dizziness
- problems with balance or muscle movement
- chest pain, shortness of breath
- fast or pounding heartbeats
- very slow heartbeats
- or dark urine. Common Vimpat side effects may include: headache , dizziness
- drowsiness
- double vision
- or nausea .

2809. This particular disease, Seizures, interacts with the drug midazolam and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Midazolam can slow or stop your breathing, especially if you have recently used an opioid medication, alcohol, or other drugs that can slow your breathing. Your caregivers will watch you for symptoms such as weak or shallow breathing. Tell your medical caregivers right away if you have: cough, wheezing, trouble breathing
- slow heart rate
- a light-headed feeling, like you might pass out
- tremors
- or confusion, agitation , hallucinations, unusual thoughts or behavior. Drowsiness or dizziness may last longer in older adults. Use caution to avoid falling or accidental injury. Common side effects of

Drug Side Effects Summary

midazolam may include: amnesia or forgetfulness after your procedure

- drowsiness, dizziness
- nausea , vomiting
- or blurred vision.

2810. This particular disease, Seizures, interacts with the drug Nayzilam and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Nayzilam can slow or stop your breathing, especially if you have recently used an opioid medication or alcohol. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Tell your doctor right away if you have new or sudden changes in mood or behavior, including new or worse depression or anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, more active or talkative, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: weak or shallow breathing
- confusion, paranoia, thoughts about suicide or hurting yourself
- a seizure
- or blurred vision, eye pain or redness, seeing halos around lights. Common Nayzilam side effects may include: drowsiness
- headache
- runny nose , discomfort in your nose
- or throat irritation. After you stop using Nayzilam, get medical help right away if you have symptoms such as: unusual muscle movements, being more active or talkative, sudden and severe changes in mood or behavior, confusion, hallucinations, seizures, suicidal thoughts or actions. Some

Drug Side Effects Summary

withdrawal symptoms may last up to 12 months or longer after stopping this medicine suddenly. Tell your doctor if you have ongoing anxiety, depression, problems with memory or thinking, trouble sleeping, ringing in your ears, a burning or prickly feeling, or a crawling sensation under your skin.

2811. This particular disease, Seizures, interacts with the drug Sabril and has the following side effects

- Tell the doctor right away if the child shows signs of vision changes, such as bumping into things or being easily startled or surprised. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: increased or worsening seizures
- any change in your vision, no matter how mild
- unusual or involuntary eye movements
- severe drowsiness, fussiness, or feeding problems (in a baby taking vigabatrin)
- numbness , tingling, or burning pain in your hands or feet
- weight gain with or without swelling
- signs of an ear infection - fever, ear pain or full feeling, trouble hearing, drainage from the ear, fussiness in a child
- or low red blood cells (anemia) - pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Some babies treated with Sabril have had abnormal brain changes seen on magnetic resonance imaging (MRI). It is not known if these changes are caused by vigabatrin or if they are harmful. Tell any doctor who treats your baby that the child is taking this medicine. Common Sabril side effects may include: blurred vision or other eye problems
- drowsiness, dizziness, feeling tired
- problems with walking or coordination

Drug Side Effects Summary

- tremors or shaking
- aggressive behavior
- confusion, problems with thinking or memory
- weight gain
- joint pain
- cold symptoms such as stuffy nose, sneezing, sore throat
- or (in babies) fussiness, ear infection, cough, breathing problems.

2812. This particular disease, Seizures, interacts with the drug vigabatrin and has the following side effects:

- Vigabatrin can cause serious side effects, including: See "Important information" above. Sleepiness and tiredness. See "What should I avoid while taking vigabatrin?" above. Vigabatrin may cause your baby to be sleepy. Sleepy babies may have a harder time suckling and feeding, or may be irritable. Weight gain that happens without swelling. The following serious side effects happen in adults. It is not known if these side effects also happen in babies who take vigabatrin. low red blood cell counts (anemia). nerve problems. Symptoms of a nerve problem can include numbness and tingling in your toes or feet. It is not known if nerve problems will go away after you stop taking this medication. swelling. If you or your child has complex partial seizures, vigabatrin may make certain types of seizures worse. Tell your healthcare provider right away if your (or your child's) seizures get worse. The most common side effect of vigabatrin in adults include: blurred vision, sleepiness, dizziness, problems walking or feeling uncoordinated, shaking (tremor), and tiredness. The most common side effect of vigabatrin in children 3 to 16 years of age is weight gain. Also expect side effects like those seen in adults. If you are giving vigabatrin to your baby for infantile spasms: Vigabatrin may make certain types of seizures worse. You should tell your baby's healthcare provider right away if your baby's seizures get worse. Tell your baby's healthcare provider if you see any changes in your baby's behavior. The most common side effects of vigabatrin in babies include:

Drug Side Effects Summary

Sleepiness - this medication may cause your baby to be sleepy. Sleepy babies may have a harder time suckling and feeding or may be irritable. Swelling in the bronchial tubes (bronchitis) Ear infection Irritability Tell your healthcare provider if you or your child have any side effect that bothers you or that does not go away. These are not all the possible side effects of this medication. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Vigabatrin side effects

2813. This particular disease, Seizures, interacts with the drug Trileptal and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash, fever, swollen glands, flu-like symptoms, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Oxcarbazepine can reduce the sodium in your body to dangerously low levels, which can cause a life-threatening electrolyte imbalance. Call your doctor right away if you have nausea, lack of energy, confusion, feeling tired or irritable, severe weakness, muscle pain, or increased seizures. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, depression , anxiety , or if you feel agitated, hostile, restless, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself. Common Trileptal side effects may include: dizziness , drowsiness, tiredness
- balance or coordination problems
- nausea, vomiting
- tremors or shaking
- double vision
- or rash.

Drug Side Effects Summary

2814. This particular disease, Seizures, interacts with the drug zonisamide and has the following side effects:

- any form of skin rash , hives
- fever, swollen glands, feeling weak or tired, severe muscle pain, unusual bruising or bleeding
- yellowing of your skin or eyes
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: sudden mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), depressed, or have thoughts about suicide or hurting yourself. Zonisamide may cause serious side effects. Call your doctor at once if you have: eye pain or redness or any changes in your vision
- decreased sweating, feeling very hot
- signs of metabolic acidosis--confusion, vomiting , lack of energy, irregular heartbeats
- symptoms of a blood cell disorder--fever, chills, body aches, flu symptoms, sores in your mouth and throat
- symptoms of a kidney stone--severe pain in your stomach or lower back, blood in your urine
- increased or worsening seizures
- the first sign of any skin rash, no matter how mild
- or severe skin reaction--fever, sore throat , swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of zonisamide may include: drowsiness, dizziness
- problems with memory or concentration
- feeling agitated or irritable
- loss of coordination, trouble walking

Drug Side Effects Summary

- or loss of appetite.

2815. This particular disease, Seizures, interacts with the drug Zonegran and has the following side effects:

- any form of skin rash , hives
- fever, swollen glands, feeling weak or tired, severe muscle pain, unusual bruising or bleeding
- yellowing of your skin or eyes
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: sudden mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: eye pain or redness or any changes in your vision
- decreased sweating, feeling very hot
- signs of metabolic acidosis - confusion, vomiting , lack of energy, irregular heartbeats
- symptoms of a blood cell disorder - fever, chills, body aches, flu symptoms, sores in your mouth and throat
- symptoms of a kidney stone - severe pain in your stomach or lower back, blood in your urine
- increased or worsening seizures
- the first sign of any skin rash, no matter how mild
- or severe skin reaction - fever, sore throat , swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common Zonegran side effects may include: drowsiness, dizziness
- problems with memory or concentration
- feeling agitated or irritable

Drug Side Effects Summary

- loss of coordination, trouble walking
- or loss of appetite.

2816. This particular disease, Seizures, interacts with the drug Keppra XR and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Aggressive or angry anxiety change in personality chills cough crying deep or fast breathing with dizziness delusions of persecution, mistrust, suspiciousness, or combativeness diarrhea dry mouth false or unusual sense of well-being feeling of unreality fever general feeling of discomfort or illness headache hoarseness irregular heartbeat irritability joint pain loss of appetite lower back or side pain mental depression muscle aches and pains nausea numbness of the feet, hands, and around the mouth painful or difficult urination quick to react or overreact emotionally rapidly changing moods restlessness sense of detachment from self or body shaking shivering sleepiness or unusual drowsiness sore throat stuffy or runny nose sweating trouble sleeping unusual tiredness or weakness vomiting Less common Bloody nose burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings clumsiness or unsteadiness discouragement dizziness or lightheadedness double vision earache feeling of constant movement of self or surroundings feeling sad or empty increase in body movements loss of bladder control loss of memory mood or mental changes outburst of anger pain or tenderness around the eyes and cheekbones problems with memory redness or swelling in the ear seizures sensation of spinning shakiness and unsteady walk shakiness in the legs, arms, hands, or feet tightness of the chest trembling or shaking of the hands or feet trouble concentrating unsteadiness, trembling, or other problems with muscle control or coordination Incidence not known Attempts at killing oneself being forgetful bleeding gums blistering, peeling, or loosening of the skin bloating blood in the urine or stools bloody, black, or tarry stools blurred vision changes in vision

Drug Side Effects Summary

chest pain coma confusion constipation dark urine difficulty with moving fast heartbeat fever with or without chills general feeling of tiredness or weakness high fever increased thirst indigestion itching large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs light-colored stools muscle pains or stiffness pains in the stomach, side, or abdomen, possibly radiating to the back pale skin pinpoint red spots on the skin red skin lesions, often with a purple center red, irritated eyes skin rash, encrusted, scaly, and oozing sores, ulcers, or white spots on the lips or in the mouth stomach pain, continuing swelling of the face, ankles, or hands swollen glands swollen joints thoughts or attempts at killing oneself trouble with balance twitching, twisting, or uncontrolled repetitive movements of the tongue, lips, face, arms, or legs uncontrolled jerking or twisting movements of the hands, arms, or legs uncontrolled movements of the lips, tongue, or cheeks unexplained bleeding or bruising unusual bleeding or bruising upper right abdominal or stomach pain weight loss yellow eyes or skin Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: More common Loss of strength or energy muscle pain or weakness pain tender, swollen glands in the neck trouble swallowing unusual weak feeling voice changes Less common Body aches or pain burning, dry, or itching eyes change in the color of the skin congestion cough increased sneezing Incidence not known Hair loss or thinning of the hair Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Keppra XR side effects

2817. This particular disease, Seizures, interacts with the drug Topamax and has the following side

Drug Side Effects Summary

- mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: a skin rash, no matter how mild
- vision problems, blurred vision, eye pain or redness, sudden vision loss (can be permanent if not treated quickly)
- confusion, problems with thinking or memory, trouble concentrating, problems with speech
- dehydration symptoms--decreased sweating, high fever, hot and dry skin
- signs of a kidney stone--severe pain in your side or lower back, painful or difficult urination
- signs of too much acid in your blood--irregular heartbeats, feeling tired, loss of appetite, trouble thinking, feeling short of breath
- or signs of too much ammonia in your blood-- vomiting , unexplained weakness, feeling like you might pass out. Common Topamax side effects may include include: dizziness , drowsiness, tiredness, slow reactions
- problems with speech or memory, abnormal vision
- numbness or tingling in your arms and legs, decreased sensation (especially in the skin)
- changes in your sense of taste, feeling nervous
- nausea , diarrhea, stomach pain, loss of appetite
- fever, weight loss
- or cold symptoms such as stuffy nose, sneezing, sore throat.

2818. This particular disease, Seizures, interacts with the drug primidone and has the following side effects:

- hives , mouth sores, skin rash with blistering and peeling
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your

Drug Side Effects Summary

doctor, such as: mood or behavior changes, depression, anxiety , or if you feel agitated, hostile, restless, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself. Primidone may cause serious side effects. Call your doctor at once if you have: unusual thoughts or behavior

- loss of balance or coordination
- fever, chills, sore throat
- easy bruising or bleeding
- or pale skin, unusual tiredness. Common side effects of primidone may include: dizziness , drowsiness, spinning sensation
- problems with balance or muscle movement
- nausea , vomiting , loss of appetite
- feeling tired or irritable
- blurred vision
- rash
- or impotence , sexual problems.

2819. This particular disease, Seizures, interacts with the drug Mysoline and has the following side

- hives , mouth sores, skin rash with blistering and peeling
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, depression, anxiety , or if you feel agitated, hostile, restless, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself. Mysoline may cause serious side effects. Call your doctor at once if you have: unusual thoughts or behavior
- loss of balance or coordination

Drug Side Effects Summary

- fever, chills, sore throat
- easy bruising or bleeding
- or pale skin, unusual tiredness. Common side effects of Mysoline may include: dizziness , drowsiness, spinning sensation
- problems with balance or muscle movement
- nausea , vomiting , loss of appetite
- feeling tired or irritable
- blurred vision
- rash
- or impotence , sexual problems.

2820. This particular disease, Seizures, interacts with the drug Aptiom and has the following side effects

- skin rash, fever, swollen glands, flu-like symptoms, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. This reaction may occur several weeks after you began using eslicarbazepine. Eslicarbazepine can reduce the sodium in your body to dangerously low levels, which can cause a life-threatening electrolyte imbalance. Call your doctor right away if you have nausea, lack of energy, confusion, feeling tired or irritable, severe weakness, muscle pain, or increased seizures. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, depression, anxiety, or if you feel agitated, hostile, restless, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself. Also call your doctor at once if you have: severe dizziness or drowsiness, vision changes, thinking problems, trouble with walking or coordination
- sudden weakness or ill feeling, fever, chills, sore throat, mouth sores
- or liver problems - nausea, loss of appetite, upper stomach pain, dark urine, jaundice (yellowing of the skin or eyes). Common Aptiom side effects may include: dizziness, drowsiness, headache

Drug Side Effects Summary

- feeling tired
- nausea, vomiting
- coordination problems, feeling shaky
- or double vision.

2821. This particular disease, Seizures, interacts with the drug Diastat AcuDial and has the following side effects:

- Suddenly stopping or reducing the dose of Diastat AcuDial very quickly may precipitate acute withdrawal reactions, which can be life-threatening. In some cases, patients have developed withdrawal symptoms lasting weeks to more than 12 months, including but not limited to: anxiety, difficulty thinking, mental changes, depression, insomnia, abnormal skin sensations, muscle weakness, tremors, twitching, ringing in your ears, burning or prickling feeling in your hands, arms, or feet. The most frequent side effect reported for Diastat AcuDial in clinical studies was somnolence (sleepiness or drowsiness). Other side effects included dizziness, headache, pain, abdominal pain, nervousness, vasodilation (increase in diameter of blood vessel), diarrhea, ataxia/incoordination (lack of coordination), euphoria (feeling of great happiness or well-being), asthma, rhinitis (irritation of the nose similar to an allergy or a cold), and rash. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088. You may also contact Bausch Health Customer Service at 1-800-321-4576. Diastat AcuDial side effects

2822. This particular disease, Seizures, interacts with the drug Zarontin and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety, panic attacks, trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically),

Drug Side Effects Summary

depressed, or have thoughts about suicide or hurting yourself. Zarontin may cause serious side effects. Call your doctor at once if you have: fever, chills, flu symptoms, sore throat , feeling very weak

- confusion, hallucinations, unusual thoughts or behavior, extreme fear
- worsening seizures
- lupus-like syndrome--joint pain or swelling with fever, swollen glands, muscle aches, chest pain, vomiting, unusual thoughts or behavior, and patchy skin color
- signs of inflammation in your body--swollen glands, flu symptoms, easy bruising or bleeding, severe tingling or numbness , muscle weakness, upper stomach pain, jaundice (yellowing of the skin or eyes), chest pain, new or worsening cough, trouble breathing
- or severe skin reaction--fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of Zarontin may include: upset stomach , nausea, stomach pain, loss of appetite
- diarrhea , weight loss
- hiccups
- swelling in your tongue or gums
- headache , dizziness , trouble concentrating
- or feeling tired.

2823. This particular disease, Seizures, interacts with the drug lacosamide and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash , fever, swollen

Drug Side Effects Summary

glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, depression, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself. Lacosamide may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out

- severe dizziness
- problems with balance or muscle movement
- chest pain, shortness of breath
- fast or pounding heartbeats
- very slow heartbeats
- or dark urine. Common side effects of lacosamide may include: headache , dizziness
- drowsiness
- double vision
- or nausea .

2824. This particular disease, Seizures, interacts with the drug Phenytoin Sodium and has the following side effects:

- No side effects listed

2825. This particular disease, Seizures, interacts with the drug Gabitril and has the following side effects:

- hives
- fever
- swollen glands
- painful sores in or around your eyes or mouth
- difficulty breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. New seizures have occurred in some people who are not epileptic who take medicine for partial seizures. Call your doctor at once if you have new or worsening seizures. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Gabitril may cause serious side effects. Call your doctor at once if you have: new or worsened seizures
- confusion, extreme weakness
- a light-headed feeling, like you might pass out
- vision problems
- or severe skin reaction--fever, sore throat , swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of Gabitril may include: tiredness, lack of energy
- dizziness , drowsiness
- nausea , stomach pain
- feeling nervous or irritable
- tremors
- or trouble concentrating.

2826. This particular disease, Seizures, interacts with the drug Oxtellar XR and has the following side effects:

- See " Important information" above. Oxtellar XR may cause other serious side effects including seizures that can happen more often or become worse, especially in children. The most common side effects of Oxtellar XR include: dizziness sleepiness headache balance problems tremors vomiting double vision weakness or lack of energy (asthenia) These are not all the possible side

Drug Side Effects Summary

effects of Oxtellar XR. For more information, ask your healthcare provider or pharmacist. Tell your healthcare provider if you have any side effect that bothers you or does not go away. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Oxtellar XR side effects

2827. This particular disease, Seizures, interacts with the drug Diastat Pediatric and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Diazepam can slow or stop your breathing, especially if you have recently used an opioid medication or alcohol. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, blue colored lips, or if you are hard to wake up. Call your doctor at once if the person receiving this medicine has: new or worsening seizures
- unusual changes in mood or behavior
- thoughts of suicide or hurting yourself
- confusion, hallucinations
- sleep problems
- or anxiety , excitement, anger, or feeling restless. Drowsiness or dizziness may last longer in older adults. Use caution to avoid falling or accidental injury. Common side effects of Diastat Pediatric may include: drowsiness, dizziness, headache , feeling nervous
- problems with coordination or muscle movement
- flushing (sudden warmth, redness, or tingly feeling)
- stomach pain, diarrhea
- or rash. After you stop using diazepam, get medical help right away if you have symptoms such as: unusual muscle movements, being more active or talkative, sudden and severe changes in mood or

Drug Side Effects Summary

behavior, confusion, hallucinations, seizures, suicidal thoughts or actions. Some withdrawal symptoms may last up to 12 months or longer after stopping this medicine suddenly. Tell your doctor if you have ongoing anxiety, depression, problems with memory or thinking, trouble sleeping, ringing in your ears, a burning or prickly feeling, or a crawling sensation under your skin.

2828. This particular disease, Seizures, interacts with the drug Qudexy XR and has the following side effects:

- mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), depressed, or have thoughts about suicide or hurting yourself. Qudexy XR Sprinkle may cause serious side effects. Call your doctor at once if you have: a skin rash, no matter how mild
- vision problems, blurred vision, eye pain or redness, sudden vision loss (can be permanent if not treated quickly)
- confusion, problems with thinking or memory, trouble concentrating, problems with speech
- dehydration symptoms--decreased sweating, high fever, hot and dry skin
- signs of a kidney stone--severe pain in your side or lower back, painful or difficult urination
- signs of too much acid in your blood--irregular heartbeats, feeling tired, loss of appetite, trouble thinking, feeling short of breath
- or signs of too much ammonia in your blood-- vomiting , unexplained weakness, feeling like you might pass out. Common side effects of Qudexy XR Sprinkle may include: dizziness , drowsiness, tiredness, slow reactions
- problems with speech or memory, abnormal vision
- numbness or tingling in your arms and legs, decreased sensation (especially in the skin)
- changes in your sense of taste, feeling nervous
- nausea , diarrhea, stomach pain, loss of appetite
- fever, weight loss

Drug Side Effects Summary

- or cold symptoms such as stuffy nose, sneezing, sore throat.

2829. This particular disease, Seizures, interacts with the drug Vitamin B6 and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Vitamin B6 may cause serious side effects. Call your doctor at once if you have: decreased sensation to touch, temperature, and vibration
- loss of balance or coordination
- numbness in your feet or around your mouth
- clumsiness in your hands
- or feeling tired. Common side effects of Vitamin B6 may include: nausea
- headache
- drowsiness
- or mild numbness or tingling.

2830. This particular disease, Seizures, interacts with the drug Briviact and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: severe dizziness or drowsiness
- a light-headed feeling, like you might pass out
- loss of balance or coordination

Drug Side Effects Summary

- unusual thoughts or behavior
- or hallucinations (seeing or hearing things that are not real). Common Briviact side effects may include: dizziness, drowsiness
- nausea , vomiting
- or feeling tired.

2831. This particular disease, Seizures, interacts with the drug Elepsia XR and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Aggressive or angry anxiety change in personality chills cough crying deep or fast breathing with dizziness delusions of persecution, mistrust, suspiciousness, or combativeness diarrhea dry mouth false or unusual sense of well-being feeling of unreality fever general feeling of discomfort or illness headache hoarseness irregular heartbeat irritability joint pain loss of appetite lower back or side pain mental depression muscle aches and pains nausea numbness of the feet, hands, and around the mouth painful or difficult urination quick to react or overreact emotionally rapidly changing moods restlessness sense of detachment from self or body shaking shivering sleepiness or unusual drowsiness sore throat stuffy or runny nose sweating trouble sleeping unusual tiredness or weakness vomiting Less common Bloody nose burning, crawling, itching, numbness, prickling, "pins and needles", or tingling feelings clumsiness or unsteadiness discouragement dizziness or lightheadedness double vision earache feeling of constant movement of self or surroundings feeling sad or empty increase in body movements loss of bladder control loss of memory mood or mental changes outburst of anger pain or tenderness around the eyes and cheekbones problems with memory redness or swelling in the ear seizures sensation of spinning shakiness and unsteady walk shakiness in the legs, arms, hands, or feet tightness of the chest trembling or shaking of the hands or feet trouble concentrating

Drug Side Effects Summary

unsteadiness, trembling, or other problems with muscle control or coordination Incidence not known
Attempts at killing oneself being forgetful bleeding gums blistering, peeling, or loosening of the skin
bloating blood in the urine or stools bloody, black, or tarry stools blurred vision changes in vision
chest pain coma confusion constipation dark urine difficulty with moving fast heartbeat fever with or
without chills general feeling of tiredness or weakness high fever increased thirst indigestion itching
large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs
light-colored stools muscle pains or stiffness pains in the stomach, side, or abdomen, possibly
radiating to the back pale skin pinpoint red spots on the skin red skin lesions, often with a purple
center red, irritated eyes skin rash, encrusted, scaly, and oozing sores, ulcers, or white spots on the
lips or in the mouth stomach pain, continuing swelling of the face, ankles, or hands swollen glands
swollen joints thoughts or attempts at killing oneself trouble with balance twitching, twisting, or
uncontrolled repetitive movements of the tongue, lips, face, arms, or legs uncontrolled jerking or
twisting movements of the hands, arms, or legs uncontrolled movements of the lips, tongue, or
cheeks unexplained bleeding or bruising unusual bleeding or bruising upper right abdominal or
stomach pain weight loss yellow eyes or skin Some side effects may occur that usually do not need
medical attention. These side effects may go away during treatment as your body adjusts to the
medicine. Also, your health care professional may be able to tell you about ways to prevent or
reduce some of these side effects. Check with your health care professional if any of the following
side effects continue or are bothersome or if you have any questions about them: More common
Loss of strength or energy muscle pain or weakness pain tender, swollen glands in the neck trouble
swallowing unusual weak feeling voice changes Less common Body aches or pain burning, dry, or
itching eyes change in the color of the skin congestion cough increased sneezing Incidence not
known Hair loss or thinning of the hair Other side effects not listed may also occur in some patients.
If you notice any other effects, check with your healthcare professional. Call your doctor for medical
advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Elepsia XR

Drug Side Effects Summary

side effects

2832. This particular disease, Seizures, interacts with the drug Eprontia and has the following side

- Eprontia may cause serious side effects including: See Important information High blood ammonia levels. High ammonia in the blood can affect your mental activities, slow your alertness, make you feel tired, or cause vomiting. This has happened when Eprontia is taken with a medicine called valproic acid (Depakote). Call your healthcare provider right away if you develop unexplained tiredness, vomiting, slowing of your thinking or reaction time, or changes in your mental activities. Effects on thinking and alertness. Eprontia may affect how you think and cause confusion, problems with concentration, attention, memory, or speech. Eprontia may cause depression or mood problems, tiredness, and sleepiness. Dizziness or loss of muscle coordination. Serious skin reactions. Eprontia may cause a severe rash with blisters and peeling skin, especially around the mouth, nose, eyes, and genitals (Stevens-Johnson syndrome). Eprontia may also cause a rash with blisters and peeling skin over much of the body that may cause death (toxic epidermal necrolysis). Call your healthcare provider right away if you develop a skin rash or blisters. Kidney stones. Drink plenty of fluids when taking Eprontia to decrease your chances of getting kidney stones. Low body temperature. Taking Eprontia when you are also taking valproic acid can cause a drop in body temperature to less than 95°F, or can cause tiredness, confusion, or coma. Call your healthcare provider right away if you have any of the symptoms above. The most common side effects include: tingling of the arms and legs nervousness slow reactions (paresthesia) nausea diarrhea nervousness speech problems dizziness slow reactions pain in the abdomen decreased feeling or sensitivity, especially in the skin not feeling hungry a change in the way foods taste weight loss upper respiratory tract infection tiredness sleepiness/drowsiness difficulty with memory fever abnormal vision Tell your healthcare provider about any side effect that bothers you or that does not go away. These are not all the possible side effects of Eprontia. Call your doctor for medical advice

Drug Side Effects Summary

about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Azurity Pharmaceuticals, Inc. at 1-855-379-0383. Eprontia side effects

2833. This particular disease, Seizures, interacts with the drug ethosuximide and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), depressed, or have thoughts about suicide or hurting yourself. Ethosuximide may cause serious side effects. Call your doctor at once if you have: fever, chills, flu symptoms, sore throat , feeling very weak
- confusion, hallucinations, unusual thoughts or behavior, extreme fear
- worsening seizures
- lupus-like syndrome--joint pain or swelling with fever, swollen glands, muscle aches, chest pain, vomiting, unusual thoughts or behavior, and patchy skin color
- signs of inflammation in your body--swollen glands, flu symptoms, easy bruising or bleeding, severe tingling or numbness , muscle weakness, upper stomach pain, jaundice (yellowing of the skin or eyes), chest pain, new or worsening cough, trouble breathing
- or severe skin reaction--fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of ethosuximide may include: upset stomach , nausea, stomach pain, loss of appetite
- diarrhea , weight loss
- hiccups

Drug Side Effects Summary

- swelling in your tongue or gums
- headache , dizziness , trouble concentrating
- or feeling tired.

2834. This particular disease, Seizures, interacts with the drug Roweepra and has the following side effects:

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. If seizures are worse or not the same after starting Roweepra (levetiracetam tablets). Hallucinations (seeing or hearing things that are not there). Very bad dizziness or passing out. Change in balance. Trouble walking. Patients who take Roweepra (levetiracetam tablets) may be at a greater risk of having thoughts or actions of suicide. The risk may be greater in people who have had these thoughts or actions in the past. Call the doctor right away if signs like low mood (depression), nervousness , restlessness, grouching, panic attacks , or changes in mood or actions are new or worse. Call the doctor right away if any thoughts or actions of suicide occur. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away,

Drug Side Effects Summary

and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)

- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Low blood cell counts have happened with Roweepra (levetiracetam tablets). If blood cell counts get very low, this can lead to bleeding problems, infections, or anemia. Call your doctor right away if you have signs of infection like fever, chills, or sore throat
- any unexplained bruising or bleeding
- or if you feel very tired or weak. Roweepra side effects

2835. This particular disease, Seizures, interacts with the drug Spritam and has the following side effects

- See Important information. Spritam can cause serious side effects. Call your healthcare provider right away if you have any of these symptoms: mood and behavior changes such as aggression, agitation, anger, anxiety, apathy, mood swings, depression, hostility, and irritability. A few people may get psychotic symptoms such as hallucinations (seeing or hearing things that are really not there), delusions (false or strange thoughts or beliefs), and unusual behavior. extreme sleepiness, tiredness, and weakness. allergic reactions such as swelling of the face, lips, eyes, tongue, and throat, trouble swallowing or breathing, and hives. a skin rash. Serious skin rashes can happen after you start taking Spritam. There is no way to tell if a mild rash will become a serious reaction. Call your healthcare provider right away if you get a rash while taking Spritam. problems with muscle coordination (problems walking and moving). The most common side effects seen in adults include: sleepiness infection weakness dizziness The most common side effects seen in children include the side effects above and: tiredness decreased appetite irritability acting aggressive nasal congestion Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects. For more information, ask your healthcare provider or

Drug Side Effects Summary

pharmacist. Call your doctor for medical advice about side effects. You may report side effects to Aprelia Pharmaceuticals, LLC at 1-844-882-7732 or FDA at 1-800-FDA-1088. Spritam side effects

2836. This particular disease, Seizures, interacts with the drug Fycompa and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash , fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , fear, panic attacks , trouble sleeping, or if you feel irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself or someone else. Call your doctor at once if you have: severe dizziness, spinning sensation, feeling like you might pass out
- trouble walking, loss of balance or coordination
- feeling very weak or tired
- an accidental fall
- or memory problems, confusion, hallucinations. Accidental falls may occur more often in elderly patients who take Fycompa. Use caution to avoid falling or accidental injury while taking this medicine. Common Fycompa side effects may include: headache , dizziness, drowsiness
- feeling anxious, tired, or irritable
- nausea , vomiting , stomach pain
- bruising
- weight gain
- or loss of coordination.

Drug Side Effects Summary

2837. This particular disease, Seizures, interacts with the drug Phenytek and has the following side

- skin rash, fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Phenytek may cause serious side effects. Call your doctor at once if you have: slow or uneven heartbeats, chest pain, fluttering in your chest, and dizziness (like you might pass out)
- any skin rash, no matter how mild
- fever, chills, sore throat, swollen glands
- red or swollen gums, mouth sores
- easy bruising, unusual bleeding, purple or red spots under your skin
- or liver problems--loss of appetite, upper stomach pain, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common side effects of Phenytek may include: drowsiness, confusion
- slurred speech
- abnormal eye movement
- or problems with balance, coordination, or muscle movement.

2838. This particular disease, Seizures, interacts with the drug Roweepra XR and has the following

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives

Drug Side Effects Summary

- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. If seizures are worse or not the same after starting Roweepra XR (levetiracetam extended-release tablets). Hallucinations (seeing or hearing things that are not there). Very bad dizziness or passing out. Change in balance. Trouble walking. Patients who take Roweepra XR (levetiracetam extended-release tablets) may be at a greater risk of having thoughts or actions of suicide. The risk may be greater in people who have had these thoughts or actions in the past. Call the doctor right away if signs like low mood (depression), nervousness , restlessness, grouchiness, panic attacks , or changes in mood or actions are new or worse. Call the doctor right away if any thoughts or actions of suicide occur. A very bad skin reaction (Stevens-Johnson syndrome / toxic epidermal necrolysis) may happen. It can cause very bad health problems that may not go away, and sometimes death. Get medical help right away if you have signs like red, swollen, blistered, or peeling skin (with or without fever)
- red or irritated eyes
- or sores in your mouth, throat, nose, or eyes. Low blood cell counts have happened with Roweepra XR (levetiracetam extended-release tablets). If blood cell counts get very low, this can lead to bleeding problems, infections, or anemia. Call your doctor right away if you have signs of infection like fever, chills, or sore throat
- any unexplained bruising or bleeding
- or if you feel very tired or weak. Roweepra XR side effects

Drug Side Effects Summary

2839. This particular disease, Seizures, interacts with the drug Xcopri and has the following side effects:

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash , fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Xcopri can cause serious or life-threatening allergic reactions that can affect your liver, blood cells, or other parts of the body. Call your doctor or get emergency medical help if you have symptoms such as: problems with the electrical system of the heart - fast or pounding heartbeats, fluttering in your chest, and sudden dizziness (like you might pass out)
- feeling very weak or tired
- severe muscle pain
- fever, swollen glands, sore throat
- unusual bruising or bleeding
- painful sores in your mouth or around your eyes
- swelling in your face, mouth, or throat
- trouble breathing or swallowing
- hives or a rash
- yellowing of your skin or eyes
- any infection or illness that does not get better
- or nervous system problems-dizziness, trouble walking, loss of coordination, vision problems, drowsiness, tiredness, problems with thinking or memory. Common Xcopri side effects may include:

Drug Side Effects Summary

feeling tired

- dizziness, drowsiness
- double vision
- or headache .

2840. This particular disease, Seizures, interacts with the drug progesterone and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: unusual vaginal bleeding
- pain or burning when you urinate
- a breast lump
- sudden vision problems, severe headache or pain behind your eyes
- symptoms of depression (sleep problems, weakness, mood changes)
- severe dizziness or drowsiness, spinning sensation, confusion, shortness of breath
- heart attack symptoms - chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating
- liver problems - nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- signs of a stroke - sudden numbness or weakness (especially on one side of the body), sudden severe headache, slurred speech, problems with speech or balance
- signs of a blood clot in the lung - chest pain, sudden cough, wheezing, rapid breathing, coughing up blood
- or signs of a blood clot in your leg - pain, swelling, warmth, or redness in one or both legs.

Common progesterone side effects may include: drowsiness, dizziness

Drug Side Effects Summary

- breast pain
- mood changes
- headache
- constipation, diarrhea, heartburn
- bloating, swelling in your hands or feet
- joint pain
- hot flashes
- or vaginal discharge.

2841. This particular disease, Seizures, interacts with the drug pyridoxine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Pyridoxine may cause serious side effects. Call your doctor at once if you have: decreased sensation to touch, temperature, and vibration
- loss of balance or coordination
- numbness in your feet or around your mouth
- clumsiness in your hands
- or feeling tired. Common side effects of pyridoxine may include: nausea
- headache
- drowsiness
- or mild numbness or tingling.

2842. This particular disease, Seizures, interacts with the drug Aminoxin and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your

Drug Side Effects Summary

doctor as soon as possible if any of the following side effects occur: With large doses Clumsiness numbness of hands or feet Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Aminoxin side effects

2843. This particular disease, Seizures, interacts with the drug brivaracetam and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), depressed, or have thoughts about suicide or hurting yourself. Brivaracetam may cause serious side effects. Call your doctor at once if you have: severe dizziness or drowsiness
- a light-headed feeling, like you might pass out
- loss of balance or coordination
- unusual thoughts or behavior
- or hallucinations (seeing or hearing things that are not real). Common side effects of brivaracetam may include: dizziness, drowsiness
- nausea , vomiting
- or feeling tired.

2844. This particular disease, Seizures, interacts with the drug Celontin and has the following side effects:

- hives
- difficult breathing

Drug Side Effects Summary

- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), depressed, or have thoughts about suicide or hurting yourself. Serious and sometimes fatal infections may occur during treatment with Celontin. Call your doctor right away if you have signs of infection such as: fever
- swollen glands, sore throat
- easy bruising, unusual bleeding (nose, mouth)
- or feeling very weak or tired. Also call your doctor at once if you have: confusion, hallucinations, unusual thoughts or behavior, extreme fear
- worsening seizures
- lupus-like syndrome--joint pain or swelling with fever, swollen glands, muscle aches, chest pain, vomiting, unusual thoughts or behavior, and patchy skin color
- or severe skin reaction--fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of Celontin may include: nausea, vomiting, stomach pain, loss of appetite
- diarrhea , constipation
- weight loss
- headache , dizziness , drowsiness
- blurred vision
- or loss of balance or coordination.

2845. This particular disease, Seizures, interacts with the drug cenobamate and has the following s

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash , fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Cenobamate can cause serious or life-threatening allergic reactions that can affect your liver, blood cells, or other parts of the body. Call your doctor or get emergency medical help if you have symptoms such as: fast or pounding heartbeats, fluttering in your chest, and sudden dizziness (like you might pass out)
- feeling very weak or tired
- severe muscle pain
- fever, swollen glands, sore throat
- unusual bruising or bleeding
- painful sores in your mouth or around your eyes
- swelling in your face, mouth, or throat
- trouble breathing or swallowing
- hives or a rash
- yellowing of your skin or eyes
- any infection or illness that does not get better
- or nervous system problems--dizziness, trouble walking, loss of coordination, vision problems, drowsiness, tiredness, problems with thinking or memory. Common side effects of cenobamate may include: feeling tired
- dizziness, drowsiness

Drug Side Effects Summary

- double vision
- or headache .

2846. This particular disease, Seizures, interacts with the drug eslicarbazepine and has the following

- skin rash, fever, swollen glands, flu-like symptoms, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. This reaction may occur several weeks after you began using eslicarbazepine. Eslicarbazepine can reduce the sodium in your body to dangerously low levels, which can cause a life-threatening electrolyte imbalance. Call your doctor right away if you have nausea, lack of energy, confusion, feeling tired or irritable, severe weakness, muscle pain, or increased seizures. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, depression , anxiety , or if you feel agitated, hostile, restless, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself. Also call your doctor at once if you have: severe dizziness or drowsiness, vision changes, thinking problems, trouble with walking or coordination
- sudden weakness or ill feeling, fever, chills, sore throat, mouth sores
- or liver problems--nausea, loss of appetite, upper stomach pain, dark urine, jaundice (yellowing of the skin or eyes). Common side effects of eslicarbazepine may include: dizziness, drowsiness, headache
- feeling tired
- nausea, vomiting
- coordination problems, feeling shaky
- or double vision.

2847. This particular disease, Seizures, interacts with the drug methsuximide and has the following

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), depressed, or have thoughts about suicide or hurting yourself. Serious and sometimes fatal infections may occur during treatment with methsuximide. Call your doctor right away if you have signs of infection such as: fever
- swollen glands, sore throat
- easy bruising, unusual bleeding (nose, mouth)
- or feeling very weak or tired. Also call your doctor at once if you have: confusion, hallucinations, unusual thoughts or behavior, extreme fear
- worsening seizures
- lupus-like syndrome--joint pain or swelling with fever, swollen glands, muscle aches, chest pain, vomiting, unusual thoughts or behavior, and patchy skin color
- or severe skin reaction--fever, sore throat, swelling in your face or tongue, burning in your eyes, skin pain followed by a red or purple skin rash that spreads (especially in the face or upper body) and causes blistering and peeling. Common side effects of methsuximide may include: nausea, vomiting, stomach pain, loss of appetite
- diarrhea , constipation
- weight loss
- headache , dizziness , drowsiness
- blurred vision
- or loss of balance or coordination.

2848. This particular disease, Seizures, interacts with the drug perampanel and has the following side effects:

Drug Side Effects Summary

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Seek medical treatment if you have a serious drug reaction that can affect many parts of your body. Symptoms may include: skin rash , fever, swollen glands, muscle aches, severe weakness, unusual bruising, or yellowing of your skin or eyes. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , fear, panic attacks , trouble sleeping, or if you feel irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), or have thoughts about suicide or hurting yourself or someone else. Perampanel may cause serious side effects. Call your doctor at once if you have: severe dizziness, spinning sensation, feeling like you might pass out
- trouble walking, loss of balance or coordination
- feeling very weak or tired
- an accidental fall
- or memory problems, confusion, hallucinations. Accidental falls may occur more often in elderly patients who take perampanel. Use caution to avoid falling or accidental injury while taking this medicine. Common side effects of perampanel may include: headache , dizziness, drowsiness
- feeling anxious, tired, or irritable
- nausea , vomiting , stomach pain
- bruising
- weight gain
- or loss of coordination.

2849. This particular disease, Seizures, interacts with the drug Zonisade and has the following side

- Zonisade can cause serious side effects including: See Important information high blood ammonia levels. High ammonia in the blood can affect your mental status, slow your alertness, make you feel

Drug Side Effects Summary

tired, or cause vomiting. Call your healthcare provider right away if you develop unexplained tiredness, vomiting, slow alertness or changes in your mental status. kidney stones. Drink plenty of fluids while you take Zonisade to decrease your chances of getting kidney stones. Call your healthcare provider right away if you get back pain, stomach pain, or blood in your urine. decrease in kidney function. Zonisade may cause a decrease in kidney function. Your healthcare provider should do a blood test to measure your kidney function before and during treatment. The most common side effects include: drowsiness dizziness agitation or irritability loss of appetite trouble with walking and coordination difficulty with memory or concentration These are not all of the possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Azurity Pharmaceuticals, Inc. at 1-855-379-0383. Zonisade side effects

2850. This particular disease, Seizures, interacts with the drug ganaxolone and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have new or sudden changes in mood or behavior, including new or worse depression or anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, more active or talkative, or have thoughts about suicide or hurting yourself. Common side effects of ganaxolone may include: drowsiness
- fever
- excessive saliva or drooling
- or seasonal allergy.

2851. This particular disease, Seizures, interacts with the drug Vigadrone and has the following side effects:

Drug Side Effects Summary

- Tell the doctor right away if the child shows signs of vision changes, such as bumping into things or being easily startled or surprised. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), depressed, or have thoughts about suicide or hurting yourself. Vigadrone may cause serious side effects. Call your doctor at once if you have: increased or worsening seizures

- any change in your vision, no matter how mild

- unusual or involuntary eye movements

- severe drowsiness, fussiness, or feeding problems (in a baby taking Vigadrone)

- numbness , tingling, or burning pain in your hands or feet

- weight gain with or without swelling

- signs of an ear infection--fever, ear pain or full feeling, trouble hearing, drainage from the ear, fussiness in a child

- or low red blood cells (anemia)--pale skin, unusual tiredness, feeling light-headed or short of breath, cold hands and feet. Some babies treated with Vigadrone have had abnormal brain changes seen on magnetic resonance imaging (MRI). It is not known if these changes are caused by this medicine or if they are harmful. Tell any doctor who treats your baby that the child is taking this medicine. Common side effects of Vigadrone may include: blurred vision or other eye problems

- drowsiness, dizziness, feeling tired

- problems with walking or coordination

- tremors or shaking

- aggressive behavior

- confusion, problems with thinking or memory

- weight gain

- joint pain

Drug Side Effects Summary

- cold symptoms such as stuffy nose, sneezing, sore throat
- or (in babies) fussiness, ear infection, cough, breathing problems.

2852. This particular disease, Seizures, interacts with the drug Ztalmy and has the following side effects

- Ztalmy can cause serious side effects, including: See Important information . The most common side effects include: sleepiness fever excessive saliva or drooling seasonal allergy These are not all of the possible side effects. For more information ask your healthcare provider or pharmacist. Tell your healthcare provider about any side effect that bothers you or your child, or that does not go away. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also contact Marinus Pharmaceuticals at 844-627-4687. Ztalmy side effects

2853. This particular disease, Stroke, interacts with the drug Plavix and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Plavix increases your risk of bleeding, which can be severe or life-threatening. Call your doctor or seek emergency medical attention if you have bleeding that will not stop, if you have blood in your urine, black or bloody stools, or if you cough up blood or vomit that looks like coffee grounds. Also call your doctor at once if you have: nosebleeds, pale skin, easy bruising, purple spots under your skin or in your mouth
- jaundice (yellowing of your skin or eyes)
- fast heartbeats, shortness of breath
- headache , fever, weakness, feeling tired
- little or no urination
- a seizure

Drug Side Effects Summary

- low blood sugar - headache, hunger, sweating, irritability, dizziness, fast heart rate, and feeling anxious or shaky
- or signs of a blood clot - sudden numbness or weakness, confusion, problems with vision or speech. Common Plavix side effects may include: bleeding.

2854. This particular disease, Stroke, interacts with the drug clopidogrel and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Clopidogrel increases your risk of bleeding, which can be severe or life-threatening. Call your doctor or seek emergency medical attention if you have bleeding that will not stop, if you have blood in your urine, black or bloody stools, or if you cough up blood or vomit that looks like coffee grounds. Also call your doctor at once if you have: nosebleeds , pale skin, easy bruising, purple spots under your skin or in your mouth
- jaundice (yellowing of your skin or eyes)
- fast heartbeats, shortness of breath
- headache , fever, weakness, feeling tired
- little or no urination
- a seizure
- low blood sugar --headache, hunger, sweating, irritability, dizziness , fast heart rate , and feeling anxious or shaky
- or signs of a blood clot--sudden numbness or weakness, confusion, problems with vision or speech. Common side effects of clopidogrel may include: bleeding.

2855. This particular disease, Stroke, interacts with the drug Activase and has the following side effects:

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. Activase increases your risk of bleeding, which can be severe or fatal. Call your doctor or seek emergency medical attention if you have bleeding that will not stop. Bleeding may occur from a surgical incision, or from the skin where a needle was inserted during a blood test or while receiving injectable medication. You may also have bleeding on the inside of your body, such as in your stomach or intestines, kidneys or bladder, brain, or within the muscles. Call your doctor or get emergency medical help if you have signs of bleeding, such as: sudden headache , feeling very weak or dizzy
- bleeding gums, nosebleeds
- easy bruising
- bleeding from a wound, incision, catheter, or needle injection
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- red or pink urine
- heavy menstrual periods or abnormal vaginal bleeding
- or sudden numbness or weakness (especially on one side of the body), slurred speech, problems with vision or balance. Also call your doctor at once if you have: chest pain or heavy feeling, pain spreading to the jaw or shoulder, nausea , sweating, general ill feeling
- swelling, rapid weight gain, little or no urination
- severe stomach pain, nausea, and vomiting
- darkening or purple discoloration of your fingers or toes
- very slow heartbeats, shortness of breath, feeling light-headed
- sudden severe back pain , muscle weakness, numbness or loss of feeling in your arms or legs
- increased blood pressure--severe headache, blurred vision, pounding in your neck or ears, anxiety , nosebleed
- or pancreatitis --severe pain in your upper stomach spreading to your back, nausea and vomiting .

Drug Side Effects Summary

Bleeding is the most common side effect of alteplase .

2856. This particular disease, Stroke, interacts with the drug alteplase and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Alteplase increases your risk of bleeding, which can be severe or fatal. Call your doctor or seek emergency medical attention if you have bleeding that will not stop. Bleeding may occur from a surgical incision, or from the skin where a needle was inserted during a blood test or while receiving injectable medication. You may also have bleeding on the inside of your body, such as in your stomach or intestines, kidneys or bladder, brain, or within the muscles. Call your doctor or get emergency medical help if you have signs of bleeding, such as: sudden headache , feeling very weak or dizzy
- bleeding gums, nosebleeds
- easy bruising
- bleeding from a wound, incision, catheter, or needle injection
- bloody or tarry stools, coughing up blood or vomit that looks like coffee grounds
- red or pink urine
- heavy menstrual periods or abnormal vaginal bleeding
- or sudden numbness or weakness (especially on one side of the body), slurred speech, problems with vision or balance. Also call your doctor at once if you have: chest pain or heavy feeling, pain spreading to the jaw or shoulder, nausea , sweating, general ill feeling
- swelling, rapid weight gain, little or no urination
- severe stomach pain, nausea, and vomiting
- darkening or purple discoloration of your fingers or toes
- very slow heartbeats, shortness of breath, feeling light-headed

Drug Side Effects Summary

- sudden severe back pain , muscle weakness, numbness or loss of feeling in your arms or legs
- increased blood pressure--severe headache, blurred vision, pounding in your neck or ears, anxiety , nosebleed
- or pancreatitis --severe pain in your upper stomach spreading to your back, nausea and vomiting .

Bleeding is the most common side effect of alteplase.

2857. This particular disease, Stroke, interacts with the drug nimodipine and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Nimodipine may cause serious side effects. Call your doctor at once if you have: a light-headed feeling, like you might pass out
- fast or slow heart rate
- or swelling in your ankles or feet. Common side effects of nimodipine may include: low blood pressure (feeling light-headed)
- nausea , upset stomach
- slow heartbeats
- or muscle pain.

2858. This particular disease, Swine Flu, interacts with the drug Tamiflu and has the following side

- sudden confusion
- tremors or shaking
- unusual behavior
- or hallucinations (hearing or seeing things that are not there). Common Tamiflu side effects may include: nausea, vomiting
- headache

Drug Side Effects Summary

- or pain.

2859. This particular disease, Swine Flu, interacts with the drug oseltamivir and has the following side effects:

- sudden confusion
- tremors or shaking
- unusual behavior
- or hallucinations (hearing or seeing things that are not there). Common oseltamivir side effects may include: nausea, vomiting
- headache
- or pain.

2860. This particular disease, Swine Flu, interacts with the drug Relenza and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some people using Relenza (especially children) have had sudden unusual changes in mood or behavior. It is not certain that zanamivir is the exact cause of these symptoms. Even without using Relenza, anyone with influenza can have neurologic or behavioral symptoms. Call your doctor right away if the person using this medicine has: sudden confusion
- trouble speaking
- tremors or shaking
- a seizure
- or hallucinations (hearing or seeing things that are not there). Stop using this medicine and call your doctor at once if you have wheezing or severe breathing problems, or if you feel like you might pass out. Common Relenza side effects may include: headache , dizziness

Drug Side Effects Summary

- nausea , vomiting , diarrhea
- cough, wheezing, trouble breathing
- ear pain
- or cold symptoms such as stuffy nose, sneezing, sore throat.

2861. This particular disease, Swine Flu, interacts with the drug zanamivir and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some people using zanamivir (especially children) have had sudden unusual changes in mood or behavior. It is not certain that zanamivir is the exact cause of these symptoms. Even without using zanamivir, anyone with influenza can have neurologic or behavioral symptoms. Call your doctor right away if the person using this medicine has: sudden confusion
- trouble speaking
- tremors or shaking
- a seizure
- or hallucinations (hearing or seeing things that are not there). Stop using zanamivir and call your doctor at once if you have wheezing or severe breathing problems, or if you feel like you might pass out. Common side effects of zanamivir may include: headache , dizziness
- nausea , vomiting , diarrhea
- cough, wheezing, trouble breathing
- ear pain
- or cold symptoms such as stuffy nose, sneezing, sore throat.

2862. This particular disease, Swine Flu, interacts with the drug peramivir and has the following side effects:

Drug Side Effects Summary

- new or worsening flu symptoms
- or confusion, hallucinations, unusual thoughts or behavior. Common side effects of peramivir may include: diarrhea
- constipation
- or sleep problems (insomnia).

2863. This particular disease, UTI, interacts with the drug nitrofurantoin and has the following side

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: diarrhea that is watery or bloody
- sudden chest pain or discomfort, wheezing, dry cough or hack
- new or worsening cough, trouble breathing
- fever, chills, body aches, tiredness, unexplained weight loss
- numbness, tingling, or pain in your hands or feet
- liver problems--nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or lupus-like syndrome--joint pain or swelling with fever, swollen glands, muscle aches, chest pain, vomiting, unusual thoughts or behavior, and patchy skin color. Serious side effects may be more likely in older adults and those who are ill or debilitated. Common nitrofurantoin side effects may include: headache, dizziness
- gas, upset stomach
- mild diarrhea
- or vaginal itching or discharge.

Drug Side Effects Summary

2864. This particular disease, UTI, interacts with the drug Macrobid and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: diarrhea that is watery or bloody
- sudden chest pain or discomfort, wheezing, dry cough or hack
- new or worsening cough, trouble breathing
- fever, chills, body aches, tiredness, unexplained weight loss
- numbness, tingling, or pain in your hands or feet
- liver problems--nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes)
- or lupus-like syndrome--joint pain or swelling with fever, swollen glands, muscle aches, chest pain, vomiting, unusual thoughts or behavior, and patchy skin color. Serious side effects may be more likely in older adults and those who are ill or debilitated. Common Macrobid side effects may include: headache, dizziness
- gas, upset stomach
- mild diarrhea
- or vaginal itching or discharge.

2865. This particular disease, UTI, interacts with the drug Macrodantin and has the following side effects:

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- vision problems
- fever, chills, cough, chest pain, trouble breathing
- numbness , tingling, or burning pain in your hands or feet

Drug Side Effects Summary

- severe pain behind your eyes
 - pale skin, weakness
 - joint pain or swelling with fever, swollen glands, and muscle aches
 - pain, redness, or swelling in your lower jaw
 - increased pressure inside the skull--severe headaches, ringing in your ears, dizziness , nausea , vision problems, pain behind your eyes
 - or signs of liver or pancreas problems--upper stomach pain (that may spread to your back), nausea or vomiting, dark urine, yellowing of the skin or eyes. Side effects may be more likely in older adults.
- Common side effects of Macrochantin may include: headache , dizziness, drowsiness, weakness
- gas, indigestion , loss of appetite
 - nausea, vomiting
 - muscle or joint pain
 - rash, itching
 - or temporary hair loss .

2866. This particular disease, UTI, interacts with the drug methenamine and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Methenamine may cause serious side effects. Call your doctor at once if you have: increased urination
- painful urination
- blood in your urine
- or swelling, shortness of breath. Common side effects of methenamine may include: painful urination
- nausea , vomiting

Drug Side Effects Summary

- or rash, itching.

2867. This particular disease, UTI, interacts with the drug trimethoprim and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Trimethoprim may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- pale, gray, or bluish skin
- fever, weakness
- sore or swollen tongue
- easy bruising, purple or red spots under your skin
- a skin rash , no matter how mild
- or high potassium level--nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common side effects of trimethoprim may include: vomiting, diarrhea, stomach pain
- rash, itching
- or swelling in your tongue.

2868. This particular disease, UTI, interacts with the drug Uribel and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe dizziness , blurred vision, fast heart rate
- painful or difficult urination

Drug Side Effects Summary

- or shortness of breath. Older adults may be more likely to have side effects from this medication. Methylene blue will most likely cause your urine or stools to appear blue or green in color. This is a normal side effect of the medication and will not cause any harm. Common side effects of Uribel may include: dizziness, drowsiness
- dry mouth
- or nausea , vomiting .

2869. This particular disease, UTI, interacts with the drug Hiprex and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Hiprex may cause serious side effects. Call your doctor at once if you have: increased urination
- painful urination
- blood in your urine
- or swelling, shortness of breath. Common side effects of Hiprex may include: painful urination
- nausea , vomiting
- or rash, itching.

2870. This particular disease, UTI, interacts with the drug Furadantin and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: More common Changes in facial skin color chest pain chills cough fever general feeling of discomfort or illness hives hoarseness itching joint or muscle pain shortness of breath skin rash sudden trouble in swallowing or breathing swelling of the face, mouth, hands, or feet troubled breathing Less common Black, tarry stools blood in the

Drug Side Effects Summary

urine or stools burning, numbness, tingling, or painful sensations dizziness drowsiness headache
pinpoint red spots on the skin sore throat unsteadiness or awkwardness unusual bleeding or
bruising unusual tiredness or weakness weakness in the arms, hands, legs, or feet Rare Abdominal
or stomach pain blindness blistering, peeling, or loosening of the skin and mucous membranes
blue-yellow color blindness bluish color of the fingernails, lips, skin, palms, or nail beds blurred
vision or loss of vision, with or without eye pain bulging soft spot on the head of an infant change in
the ability to see colors, especially blue or yellow confusion cracks in the skin darkening of the urine
decreased vision diarrhea diarrhea, watery and severe, which may also be bloody eye pain general
tiredness and weakness light-colored stools loss of appetite loss of heat from the body mental
depression mood or mental changes nausea or vomiting pale skin pale stools red skin lesions, often
with a purple center red, irritated eyes red, swollen skin red, thickened, or scaly skin skin rash sores,
ulcers, or white spots on the lips or in the mouth swollen or painful glands tenderness of salivary
glands unpleasant breath odor upper right abdominal pain visual changes vomiting of blood
wheezing or tightness in the chest yellow eyes or skin Some side effects may occur that usually do
not need medical attention. These side effects may go away during treatment as your body adjusts
to the medicine. Also, your health care professional may be able to tell you about ways to prevent or
reduce some of these side effects. Check with your health care professional if any of the following
side effects continue or are bothersome or if you have any questions about them: More common
Diarrhea gas Incidence not known Dizziness or lightheadedness feeling of constant movement of
self or surroundings lack or loss of strength loss of hair, temporary sensation of spinning
uncontrolled eye movements Other side effects not listed may also occur in some patients. If you
notice any other effects, check with your healthcare professional. Call your doctor for medical advice
about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Furadantin side
effects

Drug Side Effects Summary

2871. This particular disease, UTI, interacts with the drug cefadroxil and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Cefadroxil may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- fever, chills, body aches, flu symptoms
- pale skin, easy bruising, unusual bleeding
- a seizure
- fever, weakness, confusion
- dark colored urine, jaundice (yellowing of the skin or eyes)
- or kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath. Common side effects of cefadroxil may include: diarrhea
- stomach pain
- or vaginal itching or discharge.

2872. This particular disease, UTI, interacts with the drug Duricef and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Black, tarry stools bleeding gums blistering, peeling, or loosening of the skin blood in the urine or stools chills clay-colored stools cough or hoarseness dark-colored urine diarrhea difficulty with moving feeling of discomfort fever fever with or without chills flushing or redness of the skin general feeling of tiredness or weakness headache inflammation of the joints itching joint or muscle pain light-colored stools lower back or side pain muscle aches muscle pain or stiffness painful or difficult urination pinpoint red spots on the

Drug Side Effects Summary

skin rash red skin lesions, often with a purple center red, irritated eyes sore throat sores, ulcers, or white spots on the lips or in the mouth stomach pain, continuing swollen lymph glands unusual bleeding or bruising unusually warm skin vomiting yellow eyes or skin Incidence not known Abdominal or stomach cramps or tenderness back, leg, or stomach pains bloating chest pain coughing up blood diarrhea, watery and severe, which may also be bloody difficulty with breathing or swallowing dizziness fast heartbeat general body swelling high fever hives increased menstrual flow or vaginal bleeding increased thirst itching large, hive-like swelling on the face, eyelids, lips, tongue, throat, hands, legs, feet, or sex organs loss of appetite nausea nosebleeds pain pale skin paralysis prolonged bleeding from cuts puffiness or swelling of the eyelids or around the eyes, face, lips, or tongue seizures shortness of breath sudden decrease in the amount of urine swollen or painful glands tightness in the chest unpleasant breath odor unusual tiredness or weakness unusual weight loss vomiting of blood wheezing Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Rare Acid or sour stomach belching heartburn indigestion stomach discomfort, upset, or pain Incidence not known Hives or welts itching of the vagina or genital area itching or pain of the genital area pain during sexual intercourse redness of the skin thick, white vaginal discharge with no odor or with a mild odor Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Duricef side effects

2873. This particular disease, UTI, interacts with the drug Hyophen and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and

Drug Side Effects Summary

sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. Shortness of breath. Trouble passing urine. A fast heartbeat . Blurred eyesight. Dizziness or passing out. A severe and sometimes deadly problem called serotonin syndrome may happen if you take Hyophen (methenamine, phenyl salicylate, methylene blue, benzoic acid, hyoscyamine) with certain other drugs. Call your doctor right away if you have agitation
- change in balance
- confusion
- hallucinations
- fever
- fast or abnormal heartbeat
- flushing
- muscle twitching or stiffness
- seizures
- shivering or shaking
- sweating a lot

Drug Side Effects Summary

- severe diarrhea , upset stomach , or throwing up
- or severe headache . Hyophen side effects

2874. This particular disease, UTI, interacts with the drug Urelle and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe dizziness , blurred vision, fast heart rate
- painful or difficult urination
- or shortness of breath. Older adults may be more likely to have side effects from this medication. Methylene blue will most likely cause your urine or stools to appear blue or green in color. This is a normal side effect of the medication and will not cause any harm. Common side effects of urelle may include: dizziness, drowsiness
- dry mouth
- or nausea , vomiting .

2875. This particular disease, UTI, interacts with the drug Uro-MP and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever

Drug Side Effects Summary

- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. A fast heartbeat . Blurred eyesight. Dizziness
- . Shortness of breath. Trouble passing urine. Uro-MP side effects

2876. This particular disease, UTI, interacts with the drug Urogesic Blue and has the following side

- Methylene blue will most likely cause your urine or stools to appear blue or green in color. This is a normal side effect of the medication and will not cause any harm. hives
 - difficult breathing
 - swelling of your face, lips, tongue, or throat. Urogesic-Blue may cause serious side effects. Stop using Urogesic-Blue and call your doctor at once if you have: severe dizziness , blurred vision, fast heart rate
 - agitation , confusion, feeling restless or excited
 - painful or difficult urination
 - or shortness of breath. Older adults may be more likely to have side effects from this medication.
- Common side effects of Urogesic-Blue may include: mild dizziness

- dry mouth
- or nausea , vomiting .

2877. This particular disease, UTI, interacts with the drug Ustell and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop

Drug Side Effects Summary

using this medicine and call your doctor at once if you have: severe dizziness , blurred vision, fast heart rate

- painful or difficult urination
 - or shortness of breath. Older adults may be more likely to have side effects from this medication.
- Methylene blue will most likely cause your urine or stools to appear blue or green in color. This is a normal side effect of the medication and will not cause any harm. Common side effects of Ustell may include: dizziness, drowsiness
- dry mouth
 - or nausea , vomiting .

2878. This particular disease, UTI, interacts with the drug Uta and has the following side effects:

- Methylene blue will most likely cause your urine or stools to appear blue or green in color. This is a normal side effect of the medication and will not cause any harm. hives
 - difficult breathing
 - swelling of your face, lips, tongue, or throat. UTA may cause serious side effects. Stop using UTA and call your doctor at once if you have: severe dizziness , blurred vision, fast heart rate
 - agitation , confusion, feeling restless or excited
 - painful or difficult urination
 - or shortness of breath. Older adults may be more likely to have side effects from this medication.
- Common side effects of UTA may include: mild dizziness
- dry mouth
 - or nausea , vomiting .

2879. This particular disease, UTI, interacts with the drug Utira-C and has the following side effects

- hives

Drug Side Effects Summary

- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe dizziness , blurred vision, fast heart rate
- painful or difficult urination
- or shortness of breath. Older adults may be more likely to have side effects from this medication. Methylene blue will most likely cause your urine or stools to appear blue or green in color. This is a normal side effect of the medication and will not cause any harm. Common side effects of Utira-C may include: dizziness, drowsiness
- dry mouth
- or nausea , vomiting .

2880. This particular disease, UTI, interacts with the drug acetohydroxamic acid and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Acetohydroxamic acid may cause serious side effects. Call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- signs of a blood clot in your leg--pain, swelling, warmth, or redness in one or both legs
- or signs of a red blood cell disorder--pale or yellowed skin, dark colored urine, fever, confusion or weakness. Common side effects of acetohydroxamic acid may include: headache during the first 2 days of treatment
- skin rash, warmth, tingling or redness (especially if you drink alcohol while taking acetohydroxamic acid)
- upset stomach , nausea , loss of appetite
- depressed mood

Drug Side Effects Summary

- anxiety , tremors, nervousness
- or hair loss .

2881. This particular disease, UTI, interacts with the drug Azuphen MB and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe dizziness , blurred vision, fast heart rate
- painful or difficult urination
- or shortness of breath. Older adults may be more likely to have side effects from this medication. Methylene blue will most likely cause your urine or stools to appear blue or green in color. This is a normal side effect of the medication and will not cause any harm. Common side effects of Azuphen MB may include: dizziness, drowsiness
- dry mouth
- or nausea , vomiting .

2882. This particular disease, UTI, interacts with the drug benzoic acid / hyoscyamine / methenamine:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: Less common or rare Blurred vision eye pain skin rash or hives Symptoms of overdose Blood in urine and/or stools diarrhea dizziness drowsiness (severe) fast heartbeat flushing or redness of face headache (severe or continuing) lower back pain pain or burning while urinating ringing or buzzing in the ears shortness of breath or troubled breathing sweating unusual tiredness or weakness Some side effects may

Drug Side Effects Summary

occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Difficult urination (more common with large doses taken over a prolonged period of time) dryness of mouth, nose, or throat nausea or vomiting stomach upset or pain (more common with large doses taken over a prolonged period of time) Atropine, hyoscyamine, methenamine, methylene blue, phenyl salicylate, and benzoic acid may cause your urine and/or stools to turn blue or blue-green. This is to be expected while you are taking atropine, hyoscyamine, methenamine, methylene blue, phenyl salicylate, and benzoic acid. Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Benzoic acid / hyoscyamine / methenamine / methylene blue / phenyl salicylate side effects

2883. This particular disease, UTI, interacts with the drug cefiderocol and has the following side effects

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- tremor, rigid muscles, or a seizure
- white patches or sores inside your mouth or on your lips
- or low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of cefiderocol may include: nausea , vomiting , diarrhea, constipation
- mouth sores
- cough

Drug Side Effects Summary

- abnormal liver function tests
- headache
- rash
- or pain, redness, itching, bruising, or swelling around the IV needle.

2884. This particular disease, UTI, interacts with the drug cilastatin / imipenem / relebactam and has

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Imipenem, cilastatin, and relebactam may cause serious side effects. Call your doctor at once if you have: tremors
- a seizure
- or severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose). Common side effects of imipenem, cilastatin, and relebactam may include: increased blood pressure
- nausea , diarrhea, vomiting
- abnormal liver function tests
- fever
- headache
- or pain, bruising, swelling, or irritation where the medicine was injected.

2885. This particular disease, UTI, interacts with the drug Fetroja and has the following side effects

- severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- tremor, rigid muscles, or a seizure
- white patches or sores inside your mouth or on your lips

Drug Side Effects Summary

- or low potassium level--leg cramps, constipation , irregular heartbeats, fluttering in your chest, increased thirst or urination, numbness or tingling, muscle weakness or limp feeling. Common side effects of Fetroja may include: nausea , vomiting , diarrhea, constipation
- mouth sores
- cough
- abnormal liver function tests
- headache
- rash
- or pain, redness, itching, bruising, or swelling around the IV needle.

2886. This particular disease, UTI, interacts with the drug Hyolev MB and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe dizziness , blurred vision, fast heart rate
- painful or difficult urination
- or shortness of breath. Older adults may be more likely to have side effects from this medication. Methylene blue will most likely cause your urine or stools to appear blue or green in color. This is a normal side effect of the medication and will not cause any harm. Common side effects of Hyolev MB may include: dizziness, drowsiness
- dry mouth
- or nausea , vomiting .

2887. This particular disease, UTI, interacts with the drug hyoscyamine / methenamine / methylene

Drug Side Effects Summary

- hives
 - difficult breathing
 - swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe dizziness , blurred vision, fast heart rate
 - painful or difficult urination
 - or shortness of breath. Older adults may be more likely to have side effects from this medication.
- Methylene blue will most likely cause your urine or stools to appear blue or green in color. This is a normal side effect of the medication and will not cause any harm. Common side effects of hyoscyamine, methenamine, methylene blue, phenyl salicylate, sodium biphosphate may include: dizziness, drowsiness
- dry mouth
 - or nausea , vomiting .

2888. This particular disease, UTI, interacts with the drug hyoscyamine / methenamine / methylene

- Methylene blue will most likely cause your urine or stools to appear blue or green in color. This is a normal side effect of the medication and will not cause any harm. hives
 - difficult breathing
 - swelling of your face, lips, tongue, or throat. Stop using hyoscyamine, methenamine, methylene blue, and sodium biphosphate and call your doctor at once if you have: severe dizziness , blurred vision, fast heart rate
 - agitation , confusion, feeling restless or excited
 - painful or difficult urination
 - or shortness of breath. Older adults may be more likely to have side effects from this medication.
- Common side effects of hyoscyamine, methenamine, methylene blue, and sodium biphosphate may

Drug Side Effects Summary

include: mild dizziness

- dry mouth
- or nausea , vomiting .

2889. This particular disease, UTI, interacts with the drug Lithostat and has the following side effects

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Lithostat may cause serious side effects. Call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- signs of a blood clot in your leg--pain, swelling, warmth, or redness in one or both legs
- or signs of a red blood cell disorder--pale or yellowed skin, dark colored urine, fever, confusion or weakness. Common side effects of Lithostat may include: headache during the first 2 days of treatment
- skin rash, warmth, tingling or redness (especially if you drink alcohol while taking Lithostat)
- upset stomach , nausea , loss of appetite
- depressed mood
- anxiety , tremors, nervousness
- or hair loss .

2890. This particular disease, UTI, interacts with the drug meropenem / vaborbactam and has the following side effects

- severe stomach pain, diarrhea that is watery or bloody
- a seizure (convulsions)
- sores or white patches in your mouth or throat (yeast infection or "thrush")
- severe tingling or numbness
- or unusual bleeding or bruising. Common side effects of meropenem and vaborbactam may

Drug Side Effects Summary

include: headache

- diarrhea
- or swelling or irritation where the medicine was injected.

2891. This particular disease, UTI, interacts with the drug Phosphasal and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe dizziness, blurred vision, fast heart rate
- painful or difficult urination
- or shortness of breath. Older adults may be more likely to have side effects from this medication. Methylene blue will most likely cause your urine or stools to appear blue or green in color. This is a normal side effect of the medication and will not cause any harm. Common side effects of Phosphasal may include: dizziness, drowsiness
- dry mouth
- or nausea, vomiting.

2892. This particular disease, UTI, interacts with the drug plazomicin and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Plazomicin can harm your kidneys, and may also cause nerve damage or hearing loss. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- dizziness, spinning sensation

Drug Side Effects Summary

- a light-headed feeling, like you might pass out
- hearing loss, a buzzing or roaring sound in your ears (during or after treatment with plazomicin)
- muscle weakness
- or kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath. Serious side effects may be more likely in older adults. Common side effects of plazomicin may include: nausea , vomiting , diarrhea
- headache
- or feeling light-headed.

2893. This particular disease, UTI, interacts with the drug Primsol and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Primsol may cause serious side effects. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- pale, gray, or bluish skin
- fever, weakness
- sore or swollen tongue
- easy bruising, purple or red spots under your skin
- a skin rash , no matter how mild
- or high potassium level--nausea, weakness, tingly feeling, chest pain, irregular heartbeats, loss of movement. Common side effects of Primsol may include: vomiting, diarrhea, stomach pain
- rash, itching
- or swelling in your tongue.

Drug Side Effects Summary

2894. This particular disease, UTI, interacts with the drug Recarbrio and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Recarbrio may cause serious side effects. Call your doctor at once if you have: tremors
- a seizure
- or severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose). Common side effects of Recarbrio may include: increased blood pressure
- nausea , diarrhea, vomiting
- abnormal liver function tests
- fever
- headache
- or pain, bruising, swelling, or irritation where the medicine was injected.

2895. This particular disease, UTI, interacts with the drug Uretron DS and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blurred vision dizziness rapid pulse Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Blue-colored urine Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report

Drug Side Effects Summary

side effects to the FDA at 1-800-FDA-1088. Uretron DS side effects

2896. This particular disease, UTI, interacts with the drug Urimar-T and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe dizziness , blurred vision, fast heart rate
- painful or difficult urination
- or shortness of breath. Older adults may be more likely to have side effects from this medication. Methylene blue will most likely cause your urine or stools to appear blue or green in color. This is a normal side effect of the medication and will not cause any harm. Common side effects of Urimar-T may include: dizziness, drowsiness
- dry mouth
- or nausea , vomiting .

2897. This particular disease, UTI, interacts with the drug Urin D/S and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. This medicine may cause serious side effects. Stop using this medicine and call your doctor at once if you have: severe dizziness , blurred vision, fast heart rate
- painful or difficult urination
- or shortness of breath. Older adults may be more likely to have side effects from this medication. Methylene blue will most likely cause your urine or stools to appear blue or green in color. This is a

Drug Side Effects Summary

normal side effect of the medication and will not cause any harm. Common side effects of Urin D/S may include: dizziness, drowsiness

- dry mouth
- or nausea , vomiting .

2898. This particular disease, UTI, interacts with the drug Urised and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor as soon as possible if any of the following side effects occur: Less common or rare Blurred vision eye pain skin rash or hives Symptoms of overdose Blood in urine and/or stools diarrhea dizziness drowsiness (severe) fast heartbeat flushing or redness of face headache (severe or continuing) lower back pain pain or burning while urinating ringing or buzzing in the ears shortness of breath or troubled breathing sweating unusual tiredness or weakness Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Less common Difficult urination (more common with large doses taken over a prolonged period of time) dryness of mouth, nose, or throat nausea or vomiting stomach upset or pain (more common with large doses taken over a prolonged period of time) This medicine may cause your urine and/or stools to turn blue or blue-green. This is to be expected while you are taking this medicine. Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Urised side effects

Drug Side Effects Summary

2899. This particular disease, UTI, interacts with the drug Uro-458 and has the following side effects

- No side effects listed

2900. This particular disease, UTI, interacts with the drug Uro-SP and has the following side effects

- No side effects listed

2901. This particular disease, UTI, interacts with the drug UroAv-81 and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs of an allergic reaction, like rash
 - hives
 - itching
 - red, swollen, blistered, or peeling skin with or without fever
 - wheezing
 - tightness in the chest or throat
 - trouble breathing, swallowing, or talking
 - unusual hoarseness
 - or swelling of the mouth, face, lips, tongue, or throat. A fast heartbeat . Blurred eyesight. Dizziness . Shortness of breath. Trouble passing urine. UroAv-81 side effects

2902. This particular disease, UTI, interacts with the drug UroAv-B and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs

Drug Side Effects Summary

of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. A fast heartbeat . Blurred eyesight. Dizziness . Shortness of breath. Trouble passing urine. UroAv-B side effects

2903. This particular disease, UTI, interacts with the drug UTICAP and has the following side effects

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blurred vision dizziness rapid pulse Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Blue-colored urine Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. UTICAP side effects

2904. This particular disease, UTI, interacts with the drug Utrona-C and has the following side effects

Drug Side Effects Summary

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Incidence not known Blurred vision dizziness rapid pulse Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Blue-colored urine Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Utrona-C side effects

2905. This particular disease, UTI, interacts with the drug Vabomere and has the following side effects

- severe stomach pain, diarrhea that is watery or bloody
- a seizure (convulsions)
- sores or white patches in your mouth or throat (yeast infection or "thrush")
- severe tingling or numbness
- or unusual bleeding or bruising. Common side effects of Vabomere may include: headache
- diarrhea
- or swelling or irritation where the medicine was injected.

2906. This particular disease, UTI, interacts with the drug Vilamit MB and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs

Drug Side Effects Summary

of an allergic reaction, like rash

- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. A fast heartbeat . Blurred eyesight. Dizziness
- . Shortness of breath. Trouble passing urine. Vilamit MB side effects

2907. This particular disease, UTI, interacts with the drug Vilevev MB and has the following side effects

- WARNING/CAUTION: Even though it may be rare, some people may have very bad and sometimes deadly side effects when taking a drug. Tell your doctor or get medical help right away if you have any of the following signs or symptoms that may be related to a very bad side effect: Signs
- of an allergic reaction, like rash
- hives
- itching
- red, swollen, blistered, or peeling skin with or without fever
- wheezing
- tightness in the chest or throat
- trouble breathing, swallowing, or talking
- unusual hoarseness
- or swelling of the mouth, face, lips, tongue, or throat. A fast heartbeat . Blurred eyesight. Dizziness
- . Shortness of breath. Trouble passing urine. Vilevev MB side effects

Drug Side Effects Summary

2908. This particular disease, UTI, interacts with the drug Zemdri and has the following side effects

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Zemdri can harm your kidneys, and may also cause nerve damage or hearing loss. Call your doctor at once if you have: severe stomach pain, diarrhea that is watery or bloody (even if it occurs months after your last dose)
- dizziness , spinning sensation
- a light-headed feeling, like you might pass out
- hearing loss, a buzzing or roaring sound in your ears (during or after treatment with Zemdri)
- muscle weakness
- or kidney problems--little or no urination, swelling in your feet or ankles, feeling tired or short of breath. Serious side effects may be more likely in older adults. Common side effects of Zemdri may include: nausea , vomiting , diarrhea
- headache
- or feeling light-headed.

2909. This particular disease, Weight Loss, interacts with the drug phentermine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Phentermine may cause serious side effects. Call your doctor at once if you have: feeling short of breath, even with mild exertion
- chest pain, feeling like you might pass out
- swelling in your ankles or feet
- pounding heartbeats or fluttering in your chest
- tremors, feeling restless, trouble sleeping

Drug Side Effects Summary

- unusual changes in mood or behavior
- or increased blood pressure - severe headache , blurred vision, pounding in your neck or ears, anxiety , nosebleed. Common side effects of phentermine may include: itching
- dizziness headache
- dry mouth
- unpleasant taste
- diarrhea
- constipation
- stomach pain
- or increased or decreased interest in sex.

2910. This particular disease, Weight Loss, interacts with the drug Adipex-P and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: feeling short of breath, even with mild exertion
- chest pain, feeling like you might pass out
- swelling in your ankles or feet
- pounding heartbeats or fluttering in your chest
- unusual changes in mood or behavior
- or severe headache , blurred vision, pounding in your neck or ears, anxiety , nosebleed. Common Adipex-P side effects may include: restlessness, tremors
- trouble sleeping
- dizziness , headache
- dry mouth, unpleasant taste

Drug Side Effects Summary

- diarrhea, constipation , stomach pain
- itching
- or increased or decreased interest in sex.

2911. This particular disease, Weight Loss, interacts with the drug Contrave and has the following side effects:

- fever, swollen glands, mouth sores, muscle or joint pain
- hives, rash or itching
- chest pain, difficult breathing
- swelling of your face, lips, tongue, or throat. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, severe drowsiness, or if you are hard to wake up. Stop taking this medicine and call your doctor right away if you have: severe headache, blurred vision, pounding in your neck or ears, fast heartbeats
- a seizure (convulsions)
- blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights
- changes in mood or behavior - anxiety, depression, panic attacks, trouble sleeping, agitation, thoughts about suicide or hurting yourself
- a manic episode - racing thoughts, increased energy, unusual risk-taking behavior, extreme happiness, being irritable or talkative
- liver problems - upper stomach pain, tiredness, dark urine, jaundice (yellowing of the skin or eyes)
- or severe skin reaction - fever, sore throat, burning eyes, skin pain, red or purple skin rash with blistering and peeling. Older adults may be more likely to have certain side effects. Common Contrave side effects may include: nausea, vomiting, diarrhea, constipation
- headache, dizziness
- dry mouth
- or sleep problems (insomnia).

Drug Side Effects Summary

2912. This particular disease, Weight Loss, interacts with the drug Alli and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using alli and call your doctor at once if you have:
severe stomach pain
- severe pain in your lower back
- blood in your urine, painful or difficult urination
- kidney problems - little or no urinating
- swelling in your feet or ankles
- feeling tired or short of breath
- or liver problems - nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common alli side effects caused by its fat-blocking action. These are signs that the medicine is working properly. These side effects are usually temporary and may lessen as you continue using alli: oily or fatty stools
- oily spotting in your undergarments
- orange or brown colored oil in your stool
- gas and oily discharge
- loose stools, or an urgent need to go to the bathroom, inability to control bowel movements
- an increased number of bowel movements
- or stomach pain, nausea, rectal pain.

2913. This particular disease, Weight Loss, interacts with the drug orlistat and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using orlistat and call your doctor at once if you

Drug Side Effects Summary

have: severe stomach pain

- severe pain in your lower back
- blood in your urine, painful or difficult urination
- signs of kidney problems including little or no urinating
- swelling in your feet or ankles
- feeling tired or short of breath
- or signs of liver problems including nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common orlistat side effects are caused by its fat-blocking action. These are signs that the medicine is working properly. These side effects are usually temporary and may lessen as you continue using this medicine: oily or fatty stools

- oily spotting in your undergarments
- orange or brown colored oil in your stool
- gas and oily discharge
- loose stools, or an urgent need to go to the bathroom, inability to control bowel movements
- an increased number of bowel movements
- or stomach pain, nausea, rectal pain.

2914. This particular disease, Weight Loss, interacts with the drug Xenical and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Stop using Xenical and call your doctor at once if you have: severe stomach pain
- severe pain in your lower back
- blood in your urine, painful or difficult urination

Drug Side Effects Summary

- kidney problems - little or no urinating
- swelling in your feet or ankles
- feeling tired or short of breath
- or liver problems - nausea, upper stomach pain, itching, tired feeling, loss of appetite, dark urine, clay-colored stools, jaundice (yellowing of the skin or eyes). Common Xenical side effects are caused by orlistat's fat-blocking action. These are signs that the medicine is working properly. These side effects are usually temporary and may lessen as you continue using Xenical: oily or fatty stools
- oily spotting in your undergarments
- orange or brown colored oil in your stool
- gas and oily discharge
- loose stools, or an urgent need to go to the bathroom, inability to control bowel movements
- an increased number of bowel movements
- or stomach pain, nausea, rectal pain.

2915. This particular disease, Weight Loss, interacts with the drug phendimetrazine and has the fol

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: pounding heartbeats or fluttering in your chest
- tremors, severe agitation, feeling restless, trouble sleeping
- unusual changes in mood or behavior
- little or no urination
- or increased blood pressure - severe headache , blurred vision, pounding in your neck or ears, anxiety , nosebleed. Common phendimetrazine side effects may include: flushing (warmth, redness, or tingly feeling)

Drug Side Effects Summary

- increased sweating or urination
- dizziness , headache
- blurred vision
- dry mouth, nausea
- diarrhea, constipation , stomach pain
- or increased or decreased interest in sex.

2916. This particular disease, Weight Loss, interacts with the drug diethylpropion and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Diethylpropion may cause serious side effects. Call your doctor at once if you have: chest pain, feeling short of breath (especially with exertion)
- swelling in your ankles or feet
- anxiety , feeling nervous or jittery
- muscle twitches
- feelings of extreme happiness or sadness
- fast or pounding heartbeats, fluttering in your chest
- or a light-headed feeling, like you might pass out. Common side effects of diethylpropion may include: nausea , vomiting , diarrhea , upset stomach , constipation
- headache , blurred vision
- sleep problems (insomnia)
- dizziness , drowsiness, tired feeling
- depressed mood
- dry mouth, unpleasant taste in your mouth
- decreased sex drive

Drug Side Effects Summary

- or redness, bruising, or rash.

2917. This particular disease, Weight Loss, interacts with the drug Qsymia and has the following side effects:

- hives
- difficult breathing
- swelling in your face or throat), or a severe skin reaction (fever, sore throat, burning eyes, skin pain, red or purple skin rash with blistering and peeling). Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), depressed, or have thoughts about suicide or hurting yourself. Call your doctor at once if you have: unusual changes in mood or behavior
- confusion, trouble concentrating, problems with speech or memory
- fast or pounding heartbeats while you are resting
- a seizure
- signs of a kidney stone - severe pain in your side or lower back, painful or difficult urination
- or signs of too much acid in your blood - loss of appetite, tiredness, thinking problems, irregular heartbeats. Common Qsymia side effects may include: joint pain, ankle sprain
- flu-like symptoms
- dizziness
- constipation
- numbness or tingly feeling
- sleep problems (insomnia)
- dry mouth, changes in your sense of taste
- or mood changes, feelings of low self-worth, loss of interest in things you once enjoyed, thoughts about hurting yourself.

Drug Side Effects Summary

2918. This particular disease, Weight Loss, interacts with the drug Saxenda and has the following side effects:

- hives
- fast heartbeats
- dizziness
- trouble breathing or swallowing
- swelling of your face, lips, tongue, or throat. Call your doctor at once if you have: racing or pounding heartbeats
- sudden changes in mood or behavior, suicidal thoughts
- severe ongoing nausea, vomiting, or diarrhea
- signs of a thyroid tumor - swelling or a lump in your neck, trouble swallowing, a hoarse voice, feeling short of breath
- gallbladder problems - fever, upper stomach pain, clay-colored stools, jaundice (yellowing of your skin or eyes)
- symptoms of pancreatitis - severe pain in your upper stomach spreading to your back, nausea with or without vomiting, fast heart rate
- severely low blood sugar - extreme weakness, confusion, tremors, sweating, fast heart rate, trouble speaking, nausea, vomiting, rapid breathing, fainting, and seizure (convulsions)
- or kidney problems - little or no urination
- painful or difficult urination
- swelling in your feet or ankles
- feeling tired or short of breath. Common Saxenda side effects may include: nausea (especially when you start using Saxenda), vomiting, stomach pain
- increased heart rate
- diarrhea, constipation
- headache, dizziness

Drug Side Effects Summary

- or feeling tired.

2919. This particular disease, Weight Loss, interacts with the drug Pregnyl and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some women using Pregnyl develop ovarian hyperstimulation syndrome (OHSS), a potentially life-threatening condition. Call your doctor right away if you have symptoms of OHSS: severe stomach pain or pelvic pain
- rapid weight gain, swelling around your waist, feeling short of breath
- severe nausea and vomiting , diarrhea
- or little or no urination. Pregnyl may cause serious side effects. Call your doctor at once if you have: fluid build-up around the lungs or stomach--rapid weight gain, stomach pain and bloating, pain when you breathe, feeling short of breath while lying down, cough with foamy mucus, rapid heartbeats
- signs of a blood clot--sudden numbness or weakness on one side of the body, chest pain, problems with vision or speech, pain or swelling in one leg
- or early puberty in boys --enlarged testicles and penis, facial hair, a deepened voice. Common side effects of Pregnyl may include: headache , depression
- feeling restless or irritable
- swelling
- breast tenderness or swelling
- or pain where the medicine was injected.

2920. This particular disease, Weight Loss, interacts with the drug Wegovy and has the following side effects:

- Wegovy may cause serious side effects, including: See Important information . inflammation of

Drug Side Effects Summary

your pancreas (pancreatitis). Stop using Wegovy and call your healthcare provider right away if you have severe pain in your stomach area (abdomen) that will not go away, with or without vomiting. You may feel the pain from your abdomen to your back. gallbladder problems. Wegovy may cause gallbladder problems including gallstones. Some gallbladder problems need surgery. Call your healthcare provider if you have any of the following symptoms: pain in your upper stomach (abdomen) fever yellowing of skin or eyes (jaundice) clay-colored stools increased risk of low blood sugar (hypoglycemia) in patients with type 2 diabetes, especially those who also take medicines to treat type 2 diabetes mellitus such as sulfonylureas or insulin. Low blood sugar in patients with type 2 diabetes who receive Wegovy can be both a serious and common side effect. Talk to your healthcare provider about how to recognize and treat low blood sugar. You should check your blood sugar before you start, and during treatment. Signs and symptoms of low blood sugar may include: dizziness or light-headedness blurred vision anxiety irritability or mood changes sweating slurred speech hunger confusion or drowsiness shakiness weakness headache fast heartbeat feeling jittery kidney problems (kidney failure). In people who have kidney problems, diarrhea, nausea, and vomiting may cause a loss of fluids (dehydration) which may cause kidney problems to get worse. It is important for you to drink fluids to help reduce your chance of dehydration. serious allergic reactions. Stop using Wegovy and get medical help right away, if you have any symptoms of a serious allergic reaction including: swelling of your face, lips, tongue or throat problems breathing or swallowing severe rash or itching fainting or feeling dizzy very rapid heartbeat change in vision in people with type 2 diabetes. Tell your healthcare provider if you have changes in vision during treatment. increased heart rate. Wegovy can increase your heart rate while you are at rest. Your healthcare provider should check your heart rate while you take Wegovy. Tell your healthcare provider if you feel your heart racing or pounding in your chest and it lasts for several minutes. depression or thoughts of suicide. You should pay attention to any mental changes, especially sudden changes in your mood, behaviors, thoughts, or feelings. Call your healthcare provider right

Drug Side Effects Summary

away if you have any mental changes that are new, worse, or worry you. The most common side effects may include: nausea diarrhea vomiting constipation stomach (abdomen) pain headache tiredness (fatigue) upset stomach dizziness feeling bloated belching gas stomach flu heartburn Talk to your healthcare provider about any side effect that bothers you or does not go away. These are not all the possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. Wegovy side effects

2921. This particular disease, Weight Loss, interacts with the drug bupropion / naltrexone and has the following side effects:

- fever, swollen glands, mouth sores, muscle or joint pain
- hives , rash or itching
- chest pain, difficult breathing
- swelling of your face, lips, tongue, or throat. A person caring for you should seek emergency medical attention if you have slow breathing with long pauses, severe drowsiness, or if you are hard to wake up. Stop taking bupropion and naltrexone and call your doctor right away if you have: severe headache , blurred vision, pounding in your neck or ears, fast heartbeats
- a seizure (convulsions)
- blurred vision, tunnel vision, eye pain or swelling, or seeing halos around lights
- changes in mood or behavior-- anxiety , depression, panic attacks , trouble sleeping, agitation , thoughts about suicide or hurting yourself
- a manic episode--racing thoughts, increased energy, unusual risk-taking behavior, extreme happiness, being irritable or talkative
- liver problems--upper stomach pain, tiredness, dark urine, jaundice (yellowing of the skin or eyes)
- or severe skin reaction--fever, sore throat , burning eyes, skin pain, red or purple skin rash with blistering and peeling. Older adults may be more likely to have certain side effects. Common side effects of bupropion and naltrexone may include: nausea , vomiting , diarrhea , constipation

Drug Side Effects Summary

- headache, dizziness
- dry mouth
- or sleep problems (insomnia).

2922. This particular disease, Weight Loss, interacts with the drug phentermine / topiramate and ha

- mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), depressed, or have thoughts about suicide or hurting yourself. Phentermine and topiramate may cause serious side effects. Call your doctor at once if you have: unusual changes in mood or behavior
- confusion, trouble concentrating, problems with speech or memory
- fast or pounding heartbeats while you are resting
- a seizure
- signs of a kidney stone--severe pain in your side or lower back, painful or difficult urination
- or signs of too much acid in your blood--loss of appetite, tiredness, thinking problems, irregular heartbeats. Common side effects of phentermine and topiramate may include: joint pain, ankle sprain
- flu-like symptoms
- dizziness
- constipation
- numbness or tingly feeling
- sleep problems (insomnia)
- dry mouth, changes in your sense of taste
- or mood changes, feelings of low self-worth, loss of interest in things you once enjoyed, thoughts about hurting yourself.

Drug Side Effects Summary

2923. This particular disease, Weight Loss, interacts with the drug Bontril PDM and has the following

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Seeing, hearing, or feeling things that are not there severe mental changes Incidence not known Anxiety burning while urinating chest pain or discomfort decreased ability to exercise difficult or painful urination dizziness dry mouth fainting fast, irregular, pounding, or racing heartbeat or pulse feeling of warmth headache hyperventilation increased need to urinate irritability nausea numbness or tingling in the arms or legs pain or discomfort in the arms, jaw, back, or neck passing urine more often redness of the face, neck, arms, and occasionally, upper chest restlessness shakiness in the legs, arms, hands, or feet shortness of breath sweating swelling of the feet or lower legs trembling or shaking of the hands or feet trouble with breathing trouble with sleeping trouble with thinking, speaking, or walking vomiting weakness Symptoms of overdose Abdominal or stomach cramps blurred vision change in consciousness convulsions diarrhea discouragement dizziness, faintness, or lightheadedness when getting up suddenly from a lying or sitting position fast, slow, or irregular heartbeat feeling sad or empty lack of appetite lightheadedness loss of consciousness loss of interest or pleasure overactive reflexes panic physical attempt to injure pounding in the ears rapid breathing sweating tiredness trouble with concentrating unusual tiredness or weakness violent actions Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Decreased interest in sexual intercourse difficulty having a bowel movement (stool) inability to have or keep an erection increased in sexual ability, desire, drive, or performance increased interest in sexual intercourse loss in sexual ability, desire,

Drug Side Effects Summary

drive, or performance sleeplessness stomach pain unable to sleep Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Bontril PDM side effects

2924. This particular disease, Weight Loss, interacts with the drug chorionic gonadotropin (hcg) and

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some women using human chorionic gonadotropin (HCG) develop ovarian hyperstimulation syndrome (OHSS), a potentially life-threatening condition. Call your doctor right away if you have symptoms of OHSS: severe stomach pain or pelvic pain
- rapid weight gain, swelling around your waist, feeling short of breath
- severe nausea and vomiting , diarrhea
- or little or no urination. Human chorionic gonadotropin (HCG) may cause serious side effects. Call your doctor at once if you have: fluid build-up around the lungs or stomach--rapid weight gain, stomach pain and bloating, pain when you breathe, feeling short of breath while lying down, cough with foamy mucus, rapid heartbeats
- signs of a blood clot--sudden numbness or weakness on one side of the body, chest pain, problems with vision or speech, pain or swelling in one leg
- or early puberty in boys --enlarged testicles and penis, facial hair, a deepened voice. Common side effects of human chorionic gonadotropin (HCG) may include: headache , depression
- feeling restless or irritable
- swelling
- breast tenderness or swelling
- or pain where the medicine was injected.

Drug Side Effects Summary

2925. This particular disease, Weight Loss, interacts with the drug Lomaira and has the following side effects:

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Lomaira may cause serious side effects. Call your doctor at once if you have: feeling short of breath, even with mild exertion
- chest pain, feeling like you might pass out
- swelling in your ankles or feet
- pounding heartbeats or fluttering in your chest
- tremors, feeling restless, trouble sleeping
- unusual changes in mood or behavior
- or increased blood pressure--severe headache , blurred vision, pounding in your neck or ears, anxiety , nosebleed. Common side effects of Lomaira may include: itching
- dizziness , headache
- dry mouth, unpleasant taste
- diarrhea, constipation , stomach pain
- or increased or decreased interest in sex.

2926. This particular disease, Weight Loss, interacts with the drug Atti-Plex P and has the following side effects:

- Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention. Check with your doctor immediately if any of the following side effects occur: Rare Seeing, hearing, or feeling things that are not there severe mental changes Incidence not known Chest pain decreased ability to exercise dizziness fainting fast, irregular, pounding, or racing heartbeat or pulse headache numbness or tingling in the arms or legs swelling of the feet or lower legs trembling or shaking of the legs, arms, hands, or feet trouble breathing trouble with thinking, speaking, or walking weakness

Drug Side Effects Summary

Some side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. Also, your health care professional may be able to tell you about ways to prevent or reduce some of these side effects. Check with your health care professional if any of the following side effects continue or are bothersome or if you have any questions about them: Incidence not known Decreased interest in sexual intercourse difficulty having a bowel movement (stool) dry mouth false or unusual sense of well-being hives or welts, itching, or skin rash inability to have or keep an erection increased in sexual ability, desire, drive, or performance increased interest in sexual intercourse loss in sexual ability, desire, drive, or performance redness of the skin unpleasant taste Other side effects not listed may also occur in some patients. If you notice any other effects, check with your healthcare professional. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. Atti-Plex P side effects

2927. This particular disease, Weight Loss, interacts with the drug benzphetamine and has the following

- hives
- difficulty breathing
- swelling of your face, lips, tongue, or throat. Benzphetamine may cause serious side effects. Stop using benzphetamine and call your doctor at once if you have: shortness of breath (even with mild exertion), swelling, rapid weight gain
- chest pain, feeling like you might pass out
- pounding heartbeats or fluttering in your chest
- confusion or irritability, unusual thoughts or behavior
- or dangerously high blood pressure (severe headache , blurred vision, buzzing in your ears, anxiety , confusion, chest pain, shortness of breath, uneven heartbeats, seizure). Common side effects of benzphetamine may include: feeling restless or hyperactive

Drug Side Effects Summary

- headache, dizziness , tremors
- sleep problems (insomnia)
- increased sweating
- dry mouth or an unpleasant taste in your mouth
- nausea, diarrhea upset stomach
- or skin rash .

2928. This particular disease, Weight Loss, interacts with the drug fenfluramine and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Report any new or worsening symptoms to your doctor, such as: mood or behavior changes, anxiety , panic attacks , trouble sleeping, or if you feel impulsive, irritable, agitated, hostile, aggressive, restless, hyperactive (mentally or physically), more depressed, or have thoughts about suicide or hurting yourself. Fenfluramine can cause serious side effects on your heart and lungs. Call your doctor right away if you have: chest pain, pounding heartbeats or fluttering in your chest
- shortness or breath
- blue colored skin or lips
- swelling in your lower legs
- or unusual tiredness or weakness, feeling like you might pass out. Also call your doctor at once if you have: loss of appetite and weight loss
- worsening seizures
- blurred vision, tunnel vision, eye pain or redness, or seeing halos around lights
- nausea or vomiting
- or increased blood pressure--severe headache , blurred vision, pounding in your neck or ears,

Drug Side Effects Summary

anxiety, nosebleed. Seek medical attention right away if you have symptoms of serotonin syndrome, such as: agitation , hallucinations, fever, sweating, shivering, fast heart rate , muscle stiffness, twitching, loss of coordination, nausea, vomiting, or diarrhea . Fenfluramine can affect weight or growth in children. Tell your doctor if your child is not growing at a normal rate while using this medicine. Common side effects of fenfluramine may include: loss of appetite, vomiting, diarrhea, constipation

- seizures that do not stop
- feeling weak or tired
- fever, infections
- abnormal heart function tests
- problems with balance, walking, or muscle movement
- drooling
- or cold symptoms such as stuffy nose, sneezing, sore throat .

2929. This particular disease, Weight Loss, interacts with the drug Fendique ER and has the following s

- No side effects listed

2930. This particular disease, Weight Loss, interacts with the drug Imcivree and has the following s

- Imcivree may cause serious side effects, including: Male and female sexual function problems. Imcivree can cause an erection that happens without any sexual activity in males (spontaneous penile erection) and unwanted sexual reactions (changes in sexual arousal that happen without any sexual activity) in females. If you have an erection lasting longer than 4 hours, get emergency medical help right away. Depression and suicidal thoughts or actions. You or a caregiver should call your healthcare provider right away if you have any new or worsening symptoms of depression. Increased skin pigmentation and darkening of skin lesions (moles or nevi) you already have. These

Drug Side Effects Summary

changes happen because of how Imcivree works in the body and will go away when you stop using Imcivree. You should have a full body skin exam before starting and during treatment to check for skin changes. Benzyl alcohol toxicity. Benzyl alcohol is a preservative in Imcivree. Benzyl alcohol can cause serious side effects, including death, in premature and low-birth weight infants, who have received medicines that contain benzyl alcohol. Imcivree should not be used in premature and low-birth weight infants. The most common side effects of include: injection site reactions darkening of the skin nausea vomiting headache erection that happens without any sexual activity in males diarrhea stomach pain. These are not all the possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or Rhythm Pharmaceuticals at 1-833-789-6337. Imcivree side effects

2931. This particular disease, Weight Loss, interacts with the drug setmelanotide and has the following

- hives
- difficult breathing
- swelling of your face, lips, tongue, or throat. Some people have thoughts about suicide while using setmelanotide. Tell your doctor right away if you have any sudden changes in mood or behavior, or thoughts about suicide. Setmelanotide may cause serious side effects. Call your doctor at once if you have: an erection lasting longer than 4 hours. Common side effects of setmelanotide may include: bruising, swelling, warmth, redness, oozing, or bleeding where an injection was given
- skin darkening
- headache
- erection without any sexual activity in males
- or nausea , vomiting , diarrhea , or stomach pain.