

Paper IV Repair Note: Normalizations, Fixed Parts, and the Relative Period Torsor

Christopher D. Long Antoine-Auguste Le Blanc*

September 2026

galizur@gmail.com

Abstract

This is a standalone set of replacement arguments for the pre-repair version of Paper IV, *Relative Exponential Fixed Parts and Functional Periods: An Exponential Nori–Ayoub Theorem*. It fixes the exponential graph sign and the passage from perverse to lisse tensor conventions; replaces two overly general categorical arguments by affine-line adjunction; makes the unipotent Fourier direct-image argument use actual exact sequences; and separates algebraic and analytic fiber functors in the Galois argument. The last section constructs the period comparison first over $\mathbb{C}(S)$, proves equivariance, and then descends injectivity, differential simplicity, and the exact field of constants. In particular, no assertion that the actual comparison matrix is already Picard–Vessiot over the smaller period-constant field is used to prove that assertion. External coefficient-system, Fourier, and 1-motivic inputs are identified separately from the replacement proofs.

Frozen baseline and status. The repository path, frozen commit, and Paper IV source-blob identifier are, respectively,

```
long-mathematics/exponential-integral-theorem  
0bb151018dc73c9e48b847fff31fb82fd2cb9072  
4cc45c6dd428e09cf5db940632106fc394d01393.
```

The reference `baseline/paper-iv-pre-repair-2026-09-28` points to that commit. The machine-readable baseline record contains the source and PDF SHA-256 hashes. This note does not modify the baseline manuscript or its 26-page PDF. Its proofs are submitted for a fresh adversarial audit; they are not an independently checked certification or an integrated revision.

Reading contract. The coefficient-system and arithmetic theorems cited below are external inputs, not conclusions of the desired period theorem. Their compatibility interfaces are spelled out so that an auditor can challenge an input or its application independently. The final torsor result also has an abstract form: once the fixed-part and rank-one lifting hypotheses are supplied, its proof uses no further claim about the construction of exponential motives. The accompanying repair ledger distinguishes these layers.

*Le Blanc cryptographic author identifier: `b604812530b3535b23d6dbb0f5abe32d1ed1a9de35fb4f7d04d62cff08ea19c8`

Contents

1	Normalization conventions	2
1.1	The exponential sign	2
1.2	The lisse tensor normalization	3
2	The coefficient system and the affine-line quotient	4
3	Fixed parts and extension from dense opens	5
4	Fourier fibers and ordinary geometric constituents	7
5	The arithmetic kernel and rank-one lifts	9
6	Compatible fibers and Galois exactness	11
6.1	Changing to Betti coordinates	12
7	Normalized solutions and extension of constants	12
8	The torsor and the exact constant-field argument	14
8.1	The actual constant-period base	14
8.2	An explicit equivariant comparison over $\mathbb{C}(S)$	15
8.3	Descent of differential simplicity and constants	16

1 Normalization conventions

Put $k = \overline{\mathbb{Q}} \subset \mathbb{C}$. Let S/k be smooth, connected, and quasi-projective, of dimension d , and write $a : S \rightarrow \mathrm{Spec} k$ and $K = k(S)$. Unless otherwise stated, $\lambda \in k^\times$. Geometric ground field, coefficient field, de Rham field, and analytic constant field are not silently identified: algebraic connections are defined over k , the motivic categories are k -linear, and analytic comparison is taken after extension to $\mathbb{C}$.

All six operations in this note use sheaf-compatible perverse normalizations. Derived functors are derived even when no R or L is written. A vector bundle with connection M on a smooth d -fold corresponds to a perverse-normalized object, while its unshifted de Rham complex has terms in degrees $0, \dots, d$.

1.1 The exponential sign

We fix

$$\mathcal{E}^f = (\mathcal{O}_S, d - df), \quad \mathcal{L}_\lambda = (\mathcal{O}_{\mathbb{A}_t^1}, d - \lambda dt). \quad (1.1)$$

In left differential-operator notation, the latter has $\partial_t \cdot p = p' - \lambda p$. A horizontal section of this connection is $e^{\lambda t}$; an integral pairing with the twisted de Rham complex uses the dual exponential. This distinction prevents a solution-function sign from being substituted for a connection sign. We use the explicit operator convention, as in Virk's realization construction [20, §6.7].

Let $\mathbf{Q}_S$ be the exponential quotient functor, and let $i_h : S \rightarrow S \times \mathbb{A}_t^1$ be the graph of $h \in \mathcal{O}(S)$.

Lemma 1.1 (Graph realization). *The normalized graph object*

$$\mathbf{G}_S(h) := \mathbf{Q}_S(i_{h,*} \mathbf{1}_S[d])$$

realizes the connection $\mathcal{E}^{\lambda h}$ at scale λ . In particular, the graph potential realizing $\mathcal{E}^f$ is $h = f/\lambda$.

Proof. In the realization, tensor the graph object with the pullback of $\mathcal{L}_\lambda$ and push forward along $S \times \mathbb{A}^1 \rightarrow S$. The projection restricted to the graph is the identity, and the graph is proper over S . Proper base change and the projection formula therefore reduce the calculation to $i_h^* \mathcal{L}_\lambda$. Its connection is $d - \lambda dh$. The normalization $[d]$ puts this connection in the perverse heart and introduces no change in its connection form. $\square$

For example, on $\mathbb{A}_s^1$ at scale one, the potential s gives $d - ds$, not $d + ds$. The two connections are not isomorphic over $k(s)$: a rational gauge between them would have logarithmic derivative 2 or -2 , whereas a rational logarithmic derivative tends to zero at infinity.

1.2 The lisse tensor normalization

Write $\star$ for the derived additive convolution over S , including the ordinary derived tensor product in the base direction. Its derived unit is $\mathbf{1}_S^{\text{exp}} := \mathbf{Q}_S(i_{0,*} \mathbf{1}_S)$. For perverse-lisse objects put

$$N \otimes_{\text{lisse}} M = (N \star M)[-d], \quad \mathbf{1}_{S,\text{lisse}} = \mathbf{1}_S^{\text{exp}}[d]. \quad (1.2)$$

The associativity, symmetry, and unit constraints are transported from the unshifted lisse normalization by $N \mapsto N[-d]$ and its inverse. This transport is part of the definition; the symmetry is not an unadjusted restriction of the derived Koszul symmetry.

Proposition 1.2 (Normalized tensor and fiber). *On the lisse subcategory the operations in (1.2) realize ordinary tensor products of connections. For an algebraic point $s \in S(k)$, the normalized absolute restriction is $i_s^* N[-d]$. Its absolute rapid-decay Betti fiber, in cohomological variance, is*

$$\omega_{B,s}(N) = \Psi_\infty(\text{rat}^{\text{exp}}(i_s^* N[-d])). \quad (1.3)$$

In any strictly full rigid subquotient-closed lisse tensor subcategory this is an exact tensor fiber functor. In homological notation one must dualize consistently, rather than identify homology with cohomology.

Proof. For two connections, the sheaf-compatible derived tensor of their perverse-normalized objects is their usual tensor connection shifted by d . Shifting back by $[-d]$ therefore gives the usual tensor connection in the heart. Realization compatibility with derived convolution gives the first assertion, and reflection of the heart by exact faithful realization gives lisseness of the result. The same calculation for the unit gives $\mathbf{1}_{S,\text{lisse}}$.

A constant perverse-lisse object is $a^* C[d]$, and $i_s^* a^* C[d][-d] = C$. For a general lisse object, normalized restriction is the vector-space fiber of its connection, hence is in degree zero. Exact faithful realization reflects that concentration, and the usual fiber functor on connections is exact. Tensor compatibility follows by applying $i_s^*[-d]$ to (1.2). Finally the absolute nearby-fiber functor is exact, faithful, and tensor, with the rapid-decay comparison; these are the absolute convolution inputs of Katz and Fresán–Jossen, in the form recalled by [17, Corollary 2.17 and Theorem 3.6]. $\square$

Remark 1.3 (Fixed-scale duality). If $\epsilon(s, t) = (s, -t)$, then $\rho_\lambda \epsilon^* = \rho_{-\lambda}$. Virk’s actual duality formula is

$$\mathbb{D} \rho_\lambda = \rho_{-\lambda} \mathbb{D},$$

not same-scale commutation [20, Theorem 6.13]. On the rigid lisse subcategory the usual expression for normalized tensor dual is $\epsilon^* \mathbb{D} N(-d)$: the Tate twist removes the smooth-base orientation twist.

Evaluation and coevaluation are understood in the transported normalization. In particular the tensor inverse of $G_S(h)$ is $G_S(-h)$; this follows directly from graph addition and does not rely on a bare Verdier-duality assertion. We do not infer that every object of the entire perverse heart is rigid.

2 The coefficient system and the affine-line quotient

External input 2.1 (Ordinary coefficient-system interface). Use the rational perverse-Nori system with exact scalar extension to k . Its constructible derived categories are identified with the bounded derived categories of the perverse hearts. They have adjunction, the six operations, projection formula, smooth purity, and conservative derived Betti realization; Betti realization is exact and faithful on the heart. There are compatible Hodge and algebraic de Rham realizations, the latter defined over k on the geometric objects in use. After extension to $\mathbb{C}$ it is the Hodge underlying connection realization. The operations and their adjunction maps commute with realizations. The sources are Ivorra–Morel, Terenzi, and Tubach [9, 18, 19].

This input includes the *algebraic descent* of de Rham realization; a functor only to complex analytic connections would not suffice for the arithmetic step in Section 5. The note does not reconstruct the universal Nori diagram category. This interface, together with its scalar extension, is a specific external-dependency audit item.

Fix X and $\pi : X \times \mathbb{A}^1 \rightarrow X$. Let $\mathcal{A}_X = \text{MN}(X)$ and $\mathcal{C}_X = \pi^*[1]\mathcal{A}_X \subset \mathcal{A}_{X \times \mathbb{A}^1}$.

Lemma 2.2 (Derived affine-line full faithfulness). *The adjunction unit $\text{id} \rightarrow R\pi_*\pi^*$ is an isomorphism. In particular π^* is fully faithful on the derived categories, in all degrees.*

Proof. Under Betti realization the unit is the usual pullback–pushforward unit for the product with the contractible complex affine line, and is an isomorphism. The cone has zero derived Betti realization; conservativity makes the cone zero. Adjunction now gives

$$\text{Hom}(\pi^*A, \pi^*B[n]) = \text{Hom}(A, B[n])$$

for every n . This is an assertion about the actual adjunction unit, not merely an unspecified equivalence of objects. $\square$

Lemma 2.3 (Reflection of pulled-back heart objects). *If $A \in \mathcal{A}_{X \times \mathbb{A}^1}$ and $\text{rat}(A) \simeq \pi^*L[1]$ for a perverse sheaf L on X , then $A \simeq \pi^*B[1]$, where $B = {}^pH^{-1}(R\pi_*A)$.*

Proof. The Betti realization of $R\pi_*A$ is $L[1]$. Its other perverse cohomology groups vanish, and exact faithfulness reflects their vanishing. Thus $R\pi_*A \simeq B[1]$. The counit $\pi^*B[1] \rightarrow A$ realizes as an isomorphism. Its cone is zero by conservativity. $\square$

Proposition 2.4 (Serre closure and the derived essential image). *The subcategory $\mathcal{C}_X$ is Serre. The essential image of derived π^* consists precisely of the bounded complexes whose perverse cohomology belongs to $\mathcal{C}_X$. It is thick and stable under perverse truncations.*

Proof. Smooth pullback with connected fibers has subquotient-closed image on perverse sheaves, by BBD, Corollaire 4.2.6.2. Exact Betti realization and Lemma 2.3 give subquotient closure in the Nori heart.

For extension closure, an extension of $\pi^*B[1]$ by $\pi^*A[1]$ is represented by a connecting morphism $\pi^*B[1] \rightarrow \pi^*A[2]$. By Lemma 2.2 this is the pullback of a morphism $B \rightarrow A[1]$. Its extension

triangle is a pullback triangle, and $\pi^*[1]$ is t -exact; hence the middle heart object belongs to $\mathcal{C}_X$. This proves the Serre property using the special affine-line adjunction, not a claim about arbitrary connected smooth fibers.

The pullback of a bounded complex has cohomology in $\mathcal{C}_X$, up to the fixed shift. Conversely, induct on the number of nonzero perverse cohomology objects of a complex K . A single-cohomology object is in the image. In a truncation triangle, the two shorter objects are in the image by induction. Derived full faithfulness lifts the connecting morphism, so its cone is again a pullback. This proves the essential-image description. It also gives truncation stability. A summand is in the image because its cohomology is a summand of the pulled-back cohomology, hence belongs to the Serre subcategory. Thus the image is thick. $\square$

Remark 2.5 (Two invalid general principles are not used). The pullback image for $\mathbb{G}_m \rightarrow \mathrm{Spec} k$ is not extension-closed: the connection $d - \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} dx/x$ is a nonconstant extension of two trivial connections. Likewise a Serre inclusion need not induce a fully faithful bounded derived functor. The proof above uses derived affine-line full faithfulness before truncation induction. It agrees with the division of responsibilities in [20, Proposition 7.2 and §7.3].

Theorem 2.6 (The quotient heart and faithful realization). *The Verdier quotient*

$$D^b(\mathcal{A}_{X \times \mathbb{A}^1}) / \pi^* D^b(\mathcal{A}_X)$$

has the induced bounded t -structure with heart $\mathrm{EM}_N(X) = \mathcal{A}_{X \times \mathbb{A}^1} / \mathcal{C}_X$. Exponentiation supplies the derived six operations and derived convolution. Hodge realization descends to an exact faithful functor on the quotient heart. Composing with nonzero exponential realization gives an exact faithful ρ_λ , which reflects the bounded t -structure.

Proof. By Proposition 2.4, the localization subcategory is stable under truncations. The localization theorem for t -structures identifies the heart with the indicated Serre quotient. The derived operations are those of exponentiation [7]; this statement does not claim that all six operations are exact on every heart.

The Hodge functor maps the constant Serre subcategory into the corresponding Hodge subcategory. It thus induces an exact quotient functor. If a quotient object has zero Hodge image, any heart representative has Hodge realization pulled back from X , hence Betti realization pulled back from X . Lemma 2.3 puts the representative in $\mathcal{C}_X$. The induced exact functor therefore detects zero objects, and is faithful: apply detection to the image of any morphism it kills. This argument does not require an arbitrary quotient morphism to lift to a morphism between particular chosen representatives.

Virk's Theorems 7.5–7.6 give exactness and faithfulness of nonzero realization on the exponential Hodge heart. Composition gives the same for ρ_λ . Finally $\rho_\lambda({}^p H^i K) = H^i(\rho_\lambda K)$, so exact faithfulness detects all cohomology vanishings and reflects the bounded t -structure. $\square$

3 Fixed parts and extension from dense opens

All tensor operations in this section are normalized as in (1.2). Call $N \in \mathrm{EM}_N(S)$ lisse when $\rho_\lambda(N)$ is an algebraic vector bundle with integrable connection. Exactness, faithfulness, and the usual closure properties of algebraic connections make this a Serre subcategory. Work in its dualizable rigid envelope when using Tannakian language.

Lemma 3.1 (Lowest direct image and independent horizontal sections). *For a connection M on a smooth connected d -fold,*

$$H^j(a_+M) = 0 \ (j < -d), \quad H^{-d}(a_+M) = H_{\text{dR}}^0(S, M) = \Gamma(S, M)^\nabla.$$

Evaluation $\mathcal{O}_S \otimes_k \Gamma(S, M)^\nabla \rightarrow M$ is injective and its image is the maximal trivial algebraic subconnection.

Proof. Algebraic de Rham hypercohomology has no negative ordinary degree; the perverse shift is $[d]$. Its lowest group is the kernel of the degree-zero connection operator on global sections. This gives the displayed formula also for a nonaffine base.

Horizontal sections independent over k are independent over $K = k(S)$. Indeed, in a shortest nontrivial rational relation normalize one coefficient to one and differentiate. Minimality forces every remaining coefficient to be a common differential constant, hence to lie in k . This contradicts k -independence. Locally analytically, a horizontal section vanishing at one point vanishes identically; equivalently a fundamental solution matrix makes independent horizontal sections a constant full-rank matrix. Evaluation is therefore a bundle injection. Every trivial subconnection has a basis of global algebraic horizontal sections, proving maximality. $\square$

Theorem 3.2 (Algebraic exponential fixed part). *Let N be lisse and set*

$$C_N = {}^pH^{-d}(a_*N), \quad N^{\text{fix}} = \text{im}(a^*C_N[d] \rightarrow N).$$

*Then $\rho_\lambda(N^{\text{fix}})$ is the maximal trivial algebraic subconnection of $\rho_\lambda(N)$. An object has trivial connection realization if and only if it is constant, namely of the form $a^*C[d]$. The object N^{fix} is the maximal constant subobject of N .*

Proof. Operation compatibility and Lemma 3.1 make the perverse cohomology of a_*N vanish below $-d$ after realization. Reflection gives that vanishing in the motivic category. The lowest truncation map and adjunction therefore define the displayed morphism. It realizes as horizontal evaluation. Exact realization takes its image to the maximal trivial subconnection.

If Q realizes a trivial connection, its own evaluation map $a^*C_Q[d] \rightarrow Q$ realizes an isomorphism, and hence is an isomorphism. Conversely a constant object has constant connection realization. Apply this to N^{fix} . A morphism $a^*C[d] \rightarrow N$ corresponds by adjunction to $C[d] \rightarrow a_*N$. The lowest-cohomology truncation and the t -structure factor this map through $C_N[d]$. Thus every constant subobject is contained in N^{fix} . $\square$

Corollary 3.3 (Constants and their fiber functors). *Constant lisse objects are closed under subquotients, finite direct sums, normalized tensor products, and tensor duals. The functor $a^*[d]$ is fully faithful on the absolute heart. On the constant subcategory, the generic de Rham fiber is scalar extension from a k -valued absolute fiber functor, and its actual comparison matrix is independent of s .*

Proof. Subquotients of a trivial connection are trivial, so use Theorem 3.2. The other closure properties follow in the same way or from normalized pullback. For full faithfulness, adjunction computes $\text{Hom}(a^*C[d], a^*D[d])$ as $\text{Hom}(C, a_*a^*D)$. By the projection formula, $a_*a^*D = D \otimes a_*\mathbf{1}_S$. Connectedness and the degree-zero direct image give $H^0(a_*\mathbf{1}_S) = \mathbf{1}$ and no negative cohomology. Maps from C see just that degree-zero term, by truncation orthogonality. This gives $\text{Hom}(C, D)$. Pullback and comparison for the absolute object now give the fiber-functor statement. $\square$

Remark 3.4 (Algebraic, not merely analytic, invariants). On $\mathbb{A}_s^1$, $(\mathcal{O}, d - ds)$ has the analytic horizontal section e^s but no nonzero algebraic horizontal section: $p' = p$ has no nonzero polynomial solution. The fixed-part theorem takes algebraic horizontals. Finite Betti monodromy alone does not make an algebraic connection constant.

Let $j : U \hookrightarrow S$ be a dense open immersion. Intermediate extension in the exponential heart is the image of ${}^pH^0 j_! \rightarrow {}^pH^0 j_*$; its formation commutes with exact realization.

Lemma 3.5 (Lisse intermediate extension). *A lisse exponential object N is canonically $j_{!*} j^* N$. Restriction between the lisse subcategories is fully faithful. If $Q_U \in \mathbf{EM}_N(U)$ realizes $L|_U$ for a connection L lisse on S , then $Q = j_{!*} Q_U$ is lisse and realizes L .*

Proof. A lisse connection on S has no nonzero holonomic subobject or quotient supported in $S \setminus U$, and is the minimal extension of its restriction. Exact realization reflects the absence of such boundary objects. The natural adjunction morphisms therefore identify N with $j_{!*} j^* N$.

A morphism on U induces a morphism of intermediate extensions. The canonical identifications just proved extend it to S ; a morphism with zero restriction has boundary-supported image, hence is zero. This proves full faithfulness. Finally $\rho_\lambda(j_{!*} Q_U) = j_*(L|_U) = L$. Reflection places Q in the lisse heart. $\square$

Proposition 3.6 (Extending invertible lifts). *In the last assertion of Lemma 3.5, suppose Q_U is invertible. Then Q is invertible on S .*

Proof. Extend Q_U^{-1} by intermediate extension to Q' . Its realization is L^{-1} , so Q' is lisse. The products $Q \otimes_{\text{lis}} Q'$ and $Q' \otimes_{\text{lis}} Q$ are lisse and restrict to the unit on U . Applying Lemma 3.5 to these products and to the unit identifies both products with $1_{S, \text{lis}}$. These isomorphisms extend the inverse isomorphisms on U and satisfy their coherence identities by faithful restriction. $\square$

Remark 3.7 (What this does and does not say about opens). Every generic differential subquotient of a fixed lisse connection has a unique lisse model on S : take the saturated differential subobject in the holonomic category, then its quotient. Coherent modules with integrable connection on a smooth characteristic-zero variety are locally free, so no interior singularity is introduced. Morphisms extend by the same minimal-extension property. The preceding proposition allows an individually constructed motivic rank-one lift to be returned to that fixed base. It does not assert that the intersection of infinitely many dense opens is open, or that finite tensor generation means finitely many tensor constructions. The corresponding lisse-extension and full-faithfulness arguments apply to finite diagrams in the generic motivic category.

4 Fourier fibers and ordinary geometric constituents

For this section an ordinary geometric object means one in the ordinary Nori/Hodge realization framework, before exponentiation. On smooth dense opens we use the abelian subquotient-closed category of ordinary geometric local systems; Jacobsen–Terenzi, Corollary 5.15, identifies the two standard Nori and Ayoub descriptions [10]. No irregular Riemann–Hilbert correspondence is being used to algebraize a connection.

External input 4.1 (The Fourier comparison used here). For regular geometric input on $S \times \mathbb{A}_t^1$, use the normalized partial Fourier–Laplace transform. The finite-graph microlocalization result gives a connection on $S \times \mathbb{G}_m$ after shrinking S . For a pure polarizable minimal extension $\mathcal{T}$ on $S \times \mathbb{P}_t^1$, Sabbah’s Proposition 5.8, Corollary 5.20, and sequences (5.21)–(5.22) compare the Fourier

cycles at zero with ordinary cycles at $t = \infty$ [15]. Mixed input is reduced using its strict weight filtration and exact transform/cycle functors.

Lemma 4.2 (Finite graph reduction and nonzero-scale lissity). *For regular geometric perverse input K_* on $S \times \mathbb{A}_t^1$, after shrinking S there is a finite etale cover on which all dominating singular multisections split into pairwise disjoint graphs $t = u_i(s)$. The localized partial transform is a connection on $S \times \mathbb{G}_m$.*

Proof. Take a finite Whitney stratification adapted to the regular input. Delete the closures of the images of strata that do not dominate S . Each remaining proper stratum has dimension d : it dominates a d -dimensional base and lies properly in a $(d + 1)$ -fold. Its closure is generically finite over S . Normalize these finitely many multisections and shrink so that they are finite etale. A common finite Galois refinement splits them into sections. Delete the images of intersections of distinct sections. The characteristic variety is then contained in the zero section and the conormals of the resulting graphs.

Apply the parameterized localized Laplace/microlocalization input with $z = \lambda^{-1}$. In Sabbah's Appendix B the localized module is locally free over the parameter ring $\mathcal{O}[z, z^{-1}]$, with integrable connection; Proposition B.1 identifies its residual systems. The reference is [16, Appendix B, preprint p. 12, immediately before Proposition B.1], not p. 40 of that preprint. The underpinning general parameter statement is [6, Propositions 1.18 and 1.20]. This gives nonzero-scale lissity locally on the cover, which descends in the finite etale topology. $\square$

Lemma 4.3 (Fixed fiber and nearby local system). *After this shrinking, a fixed nonzero Fourier fiber and the nearby-cycle local system at $\lambda = 0$ have isomorphic underlying local systems on S , after forgetting the commuting nearby monodromy.*

Proof. The local system on $S^{\text{an}} \times \mathbb{C}^\times$ is a representation of $\pi_1(S) \times \mathbb{Z}$. Restriction to a fiber forgets the second operator. Nearby cycles of that local system have the same representation of $\pi_1(S)$ and retain the second operator as nearby monodromy. Choose a path from the fixed scale to a punctured neighborhood of zero. Changing its winding changes the isomorphism by a power of the commuting operator, not the underlying isomorphism class or its constituents. $\square$

Proposition 4.4 (Unipotent comparison through two triangles). *Let $t' = t^{-1}$ and let $q : S \times \mathbb{P}_t^1 \rightarrow S$ denote the proper projection after taking cycles in the Fourier coordinate. Set*

$$V = \psi_{\lambda, -1}(\mathcal{FT}), \quad W = \ker N_\lambda.$$

Every simple constituent of every perverse cohomology of q_+V occurs among the perverse-cohomology constituents of the proper direct images of

$$i_{\infty, +} \ker N_{t'}, \quad \mathcal{T}, \quad i_{\infty, +} \psi_{t', -1} \mathcal{T}.$$

All cycles here are perverse-normalized.

Proof. Before q_+ , Sabbah's (5.22) and the first sequence of (5.21) give

$$0 \longrightarrow i_{\infty, +} \ker N_{t'} \longrightarrow W \longrightarrow \mathcal{T} \longrightarrow 0, \quad (4.1)$$

$$0 \longrightarrow W \longrightarrow V \longrightarrow i_{\infty, +} \psi_{t', -1} \mathcal{T} \longrightarrow 0. \quad (4.2)$$

Those sequences are stated before the subsequent dimension-one specialization in Sabbah's argument. Apply derived proper direct image to obtain two distinguished triangles. For any triangle $A \rightarrow$

$B \rightarrow C \rightarrow$, the segment ${}^p H^j A \rightarrow {}^p H^j B \rightarrow {}^p H^j C$ expresses ${}^p H^j B$ as an extension of a quotient of ${}^p H^j A$ by a subobject of ${}^p H^j C$. Apply this observation first to (4.1), then to (4.2), for each j . Finite length gives exactly the asserted constituent containment. No exactness of q_+ on the heart, and no interchange of a constituent filtration with a derived functor, is asserted. $\square$

Theorem 4.5 (Finite Betti monodromy of rank-one factors). *A rank-one differential subquotient, defined over k , of a fixed nonzero Fourier fiber of regular geometric input has finite Betti monodromy on a suitable dense smooth open.*

Proof. For pure input, the non-unipotent Fourier comparisons identify the relevant constituents with ordinary nearby-cycle constituents at infinity. The unipotent case follows from Proposition 4.4 and proper compatibility of nearby cycles. Minimal extension, ordinary cycles, proper direct image, and perverse cohomology remain ordinary geometric operations. For mixed input, use the strict finite weight filtration on the Hodge realization, exactness of normalized Fourier transformation and nearby cycles, and semisimplicity of each pure graded Hodge object. Its simple constituent local systems belong to the ordinary geometric-origin category; the abelian subquotient closure and its compatibility with the Nori description are the comparison input of [10, Corollary 5.15]. Apply the pure argument to the graded constituents.

By Lemma 4.3, a rank-one Betti constituent of the fixed fiber is one of these ordinary geometric local systems. The local system of a rank-one differential subquotient is such a constituent, because analytic horizontal realization is exact on connections. Lam–Litt’s published Lemma 3.5 says that an ordinary geometric rank-one local system is torsion [12]. Its image is therefore finite. This gives finite Betti monodromy, not regular singularity or finite tensor order of the original algebraic connection. $\square$

5 The arithmetic kernel and rank-one lifts

External input 5.1 (Degree-one arithmetic realization). Use the cohomological Picard 1-motive

$$\mathrm{Pic}^+(U) = [\mathrm{Div}_D^0(\overline{U}) \rightarrow \mathrm{Pic}^0(\overline{U})]$$

for a smooth compactification with normal-crossings boundary. Its realization is $H^1(U)(1)$. A rational boundary divisor giving a morphism from the weight-zero lattice object realizes, when made principal, as its logarithmic de Rham class [3]. The de Rham–Betti realization of 1-motives over $\overline{\mathbb{Q}}$ is fully faithful with image closed under subquotients [8, Theorem 9.14]. The comparison with degree-one Nori motives is [2].

Theorem 5.2 (Arithmetic exponential kernel). *Let U/k be smooth and connected and L a rank-one algebraic integrable connection over k with finite Betti monodromy. On a dense affine open U° there are $f \in \mathcal{O}(U^\circ)$ and a finite-order algebraic rank-one connection A such that*

$$L|_{U^\circ} \simeq \mathcal{E}^f \otimes A.$$

Proof. The finite Betti character defines a finite étale cover and its character idempotent over k ; algebraically closed extension does not change finite étale covers in characteristic zero. Let A_0 be its finite Artin connection and replace L by $L_0 = L \otimes A_0^{-1}$. On a dense affine open trivialize its underlying line bundle and write $\nabla = d + \omega$. Integrability gives $d\omega = 0$, and trivial monodromy gives

$$\int_\gamma \omega \in 2\pi i\mathbb{Z} \quad (\gamma \in H_1(U^{\mathrm{an}}, \mathbb{Z})).$$

Define the integral Betti class η by division of these periods by $2\pi i$. Comparison sends $[\omega]$ to $2\pi i\eta$. If $\eta = 0$, comparison implies $[\omega] = 0$, and affine algebraic de Rham theory supplies $F \in \mathcal{O}(U)$ with $\omega = dF$. Then $L_0 = \mathcal{E}^{-F}$.

Otherwise the *specified* pair $([\omega], \eta)$ gives a morphism $\mathbb{Q}(-1) \rightarrow H_{\text{dRB}}^1(U)$, with our convention that this Tate period is $2\pi i$. Twist to obtain a specified morphism $\mathbb{Q}(0) \rightarrow H_{\text{dRB}}^1(U)(1)$. By External input 5.1, it lifts to the specified morphism $\mathbb{Q}(0) \rightarrow \text{Pic}^+(U)_{\mathbb{Q}}$. This is a rational boundary divisor Δ whose Picard class is zero in $\text{Pic}^0(\bar{U}) \otimes \mathbb{Q}$.

Choose $N > 0$ clearing its denominators and killing its torsion Picard class. Then $N\Delta = \text{div}(u)$ for a function with no zeros or poles on U , so $u \in \mathcal{O}(U)^\times$. Compatibility with the specified morphism gives

$$N[\omega] = [d \log u] \quad \text{in } H_{\text{dR}}^1(U/k).$$

Since U is affine, closed one-forms modulo exact global forms compute this cohomology. Thus $N\omega - d \log u = dF$ for some regular F , and

$$L_0 \simeq \mathcal{E}^{-F/N} \otimes (\mathcal{O}_U, d + N^{-1}d \log u).$$

The N th power of the second factor has horizontal algebraic generator u^{-1} and is trivial. Tensor back by A_0 . This proves the result without a relative functional-period theorem. $\square$

Proposition 5.3 (Realizing the algebraic factors). *At scale $\lambda \in k^\times$, $\mathcal{E}^f \otimes A$ in Theorem 5.2 has an invertible exponential motivic lift on U° . If the connection extends lisse to a given smooth base, so does an invertible lift.*

Proof. Use the graph potential f/λ in Lemma 1.1. Its inverse is the graph with the negative potential. A finite-order connection is a rank-one summand of a finite etale pushforward; over k its character idempotent defines a rank-one Artin motive. Insert it on the zero graph and take its normalized tensor product with the exponential graph object. Both factors are invertible and realize the desired factors. The extension assertion is Proposition 3.6. $\square$

Theorem 5.4 (Rank-one saturation on a fixed lisse model). *Let $\mathcal{T}$ be a finitely generated rigid subquotient-closed lisse subcategory of the generic exponential Nori envelope, modeled on S . Suppose its objects have regular geometric pre-Fourier representatives, as supplied by the exponential quotient and the ordinary geometric coefficient system. Let $\mathcal{D}$ be the differential tensor category generated over K by their realizations and differential subquotients. Every invertible object of $\mathcal{D}$ has an ambient invertible exponential motivic lift. The lift can be modeled on S whenever the differential object is modeled lisse there.*

Proof. An invertible object $L \in \mathcal{D}$ is a differential subquotient of a finite tensor construction in realized generators. Choose its motivic counterpart B and a common open model. The quotient description is essentially surjective on the heart, so choose an ordinary geometric representative K_B for B . At nonzero scale the representative realizes as its partial Fourier fiber. Ordinary tensor constructions, convolution, and the six operations stay within regular geometric input before exponentiation; no claim of regularity of the exponential output is made.

Apply Theorem 4.5 to L after shrinking. It has finite Betti monodromy there. It is algebraic over k , so Theorem 5.2 applies. Proposition 5.3 gives the ambient invertible lift on a dense affine open, and Proposition 3.6 returns it to S if needed. This is an object-by-object argument on one fixed lisse model, not an intersection of infinitely many opens. Rank-one objects with transcendental connection coefficients are not fed into the arithmetic theorem; extension of differential constants is handled later in Section 7. $\square$

Remark 5.5 (The optional split Fourier-geometric route). If a separate construction supplies an actual isomorphism $q^*L \simeq \mathcal{E}^g \otimes A$ with finite A on a finite cover and an effective motivic descent datum, its lift is obtained from the potential g/λ and the Artin idempotent, followed by descent and Proposition 3.6. Such a statement requires the actual global rank-one comparison and effective descent hypotheses. A formal Stokes expression alone is not substituted for them. This optional route is not used in Theorem 5.4; the full nonzero-fiber argument above is what supplies saturation for the generic envelope.

6 Compatible fibers and Galois exactness

Choose $s_0 \in S(k)$ where a tensor generator is regular and fix the algebraic de Rham evaluation fiber

$$\omega_v : \mathcal{D} \longrightarrow \mathbf{Vec}_k, \quad M \longmapsto M_{s_0}.$$

All objects of $\mathcal{D}$ have their unique lisse model on S , as explained after Proposition 3.6. The same functor composed with realization is a fiber functor on $\mathcal{T}$. Set

$$G_v = \mathrm{Aut}^\otimes(\omega_v \rho_\lambda|_{\mathcal{T}}), \quad H_v = \mathrm{Aut}^\otimes(\omega_v|_{\mathcal{D}}).$$

The differential category is generated under subquotients by the image of $\mathcal{T}$, so the induced morphism $H_v \rightarrow G_v$ is a closed immersion, by [5, Proposition 2.21(b)]. This is not a consequence of exact faithfulness alone; the generation statement is also used.

Proposition 6.1 (Motivic semi-invariant subspaces). *For $B \in \mathcal{T}$ and a character $\chi : H_v \rightarrow \mathbb{G}_m$, the χ -semi-invariant subspace of $\omega_v \rho_\lambda(B)$ is the fiber of a subobject of B in $\mathcal{T}$.*

Proof. The character defines an invertible differential object L_χ . By Theorem 5.4, choose an ambient invertible lift $\mathcal{L}_\chi$, on a common lisse base if necessary. Take

$$C_\chi = (B \otimes_{\mathrm{lis}} \mathcal{L}_\chi^{-1})^{\mathrm{fix}}.$$

By Theorem 3.2 its connection realization is the maximal trivial subconnection of $\rho_\lambda(B) \otimes L_\chi^{-1}$. Tensor its inclusion by $\mathcal{L}_\chi$ and take the image $W_\chi \subset B$. Its fiber is exactly

$$L_\chi \otimes (\rho_\lambda(B) \otimes L_\chi^{-1})^{H_v},$$

namely the desired semi-invariant subspace. Although the lift was ambient, W_χ belongs to $\mathcal{T}$ because that category is closed under subobjects of its objects. $\square$

Lemma 6.2 (Normality from semi-invariants). *Let $H \subset G$ be algebraic groups over an algebraically closed field of characteristic zero. If V^χ is G -stable for every G -representation V and every character χ of H , then H is normal in G .*

Proof. By the stabilizer-of-a-line theorem for linear algebraic groups, choose a G -representation V and a line ℓ with stabilizer H . The line has an H -character χ and lies in $W = V^\chi$. By hypothesis W is G -stable and H acts on it by scalars. For $g \in G$ and $h \in H$, the element $g^{-1}hg$ therefore preserves ℓ , so it lies in H . Apply the same argument to g^{-1} to get equality. This proves normality on geometric points, hence on the characteristic-zero algebraic groups. This is the semi-invariant form of André's criterion used in [1, 13]. $\square$

Theorem 6.3 (Exactness in algebraic evaluation coordinates). *Let $\mathcal{T}_0$ be the full constant subcategory of $\mathcal{T}$, and let $G_{0,v}$ be its group under the evaluation fiber. Then*

$$1 \longrightarrow H_v \longrightarrow G_v \longrightarrow G_{0,v} \longrightarrow 1$$

is exact.

Proof. Proposition 6.1 and Lemma 6.2 make H_v normal. By Corollary 3.3, $\mathcal{T}_0$ is a full tensor subcategory closed under subquotients. Its inclusion gives a faithfully flat quotient of G_v [5, Proposition 2.21(a)]. A $\mathcal{T}$ -object is trivial as an H_v -representation precisely when its connection is trivial. By Theorem 3.2 this is equivalent to being constant. Thus the representations of G_v/H_v are precisely $\mathcal{T}_0$, proving exactness. No period-map injectivity has been used. $\square$

6.1 Changing to Betti coordinates

Let $\omega_B = \omega_{B,s_0}$ and let ω_{dR} be the generic K -valued de Rham fiber. Define over k

$$G_B = \text{Aut}^\otimes(\omega_B|\mathcal{T}), \quad G_{0,B} = \text{Aut}^\otimes(\omega_B|\mathcal{T}_0), \quad H_B = \ker(G_B \rightarrow G_{0,B}). \quad (6.1)$$

Let $c : \omega_B \otimes_k \mathbb{C} \xrightarrow{\sim} \omega_v \rho_\lambda \otimes_k \mathbb{C}$ be the actual comparison at s_0 . It is an isomorphism of fiber functors, not an assertion that its entries are algebraic.

Lemma 6.4 (The group change is conjugation). *The comparison c identifies $(G_B)_\mathbb{C}$ with $(G_v)_\mathbb{C}$, and*

$$(H_B)_\mathbb{C} = c^{-1}(H_v)_\mathbb{C}c.$$

Proof. An isomorphism of fiber functors conjugates their tensor automorphism functors. Its restriction to $\mathcal{T}_0$ conjugates the quotient groups as well. Kernels are preserved by this commutative diagram. Use Theorem 6.3 to identify the evaluation-coordinate kernel with H_v . This proves the formula before any analytic torsor comparison. $\square$

7 Normalized solutions and extension of constants

The next argument establishes the differential base-change statement needed in the last section. It prevents arbitrary complex differential subobjects from being passed unexamined to an arithmetic theorem over k .

Shrink around s_0 to choose algebraic etale coordinates $x_1, \dots, x_d$ and a frame of a tensor-generator connection. The corresponding derivations ∂_i extend uniquely from $k(x_1, \dots, x_d)$ to K and commute. They span the tangent derivations over K . Write the system as

$$\partial_i Y = A_i Y, \quad A_i \in M_n(K), \quad Y(s_0) = I. \quad (7.1)$$

Here $\nabla = d - \sum_i A_i dx_i$; integrability is $\partial_i A_j - \partial_j A_i - [A_i, A_j] = 0$.

Lemma 7.1 (Algebraic Taylor normalization). *The solution Y of (7.1) exists on a sufficiently small analytic polydisc B and has Taylor coefficients in k .*

Proof. In etale coordinates, the completed local ring at s_0 is $k[[x_1, \dots, x_d]]$, with the coordinates translated to zero. The A_i have coefficients there. Coefficient comparison in (7.1), starting from I , determines each Taylor coefficient using only coefficients already determined and division by positive integers. Integrability is precisely the consistency of the different recursions. Analytic existence for a nonsingular flat system gives a convergent solution; uniqueness identifies its Taylor series with this formal solution. Thus all its coefficients lie in k . $\square$

External input 7.2 (Picard–Vessiot facts for integrable systems). For finitely many commuting derivations and an algebraically closed common constant field, a field generated by a fundamental matrix with no new common constants is Picard–Vessiot; its fundamental-matrix algebra is differentially simple. Its spectrum is the differential comparison torsor, and its group is the tensor automorphism group of the horizontal fiber functor. Use classical Picard–Vessiot theory and its integrable-system form [14, 11, 4]. This is ordinary Picard–Vessiot theory with all base derivations, not parameterized Picard–Vessiot theory with some derivatives omitted.

Put

$$R_k = K[Y, \det(Y)^{-1}] \subset \mathcal{M}(B), \quad E = \mathbb{C}(S).$$

The fraction field of R_k has common constants k . Indeed, a horizontal meromorphic germ is complex constant; after writing the germ as a quotient of two formal series over k , a nonzero coefficient of the denominator shows that this constant belongs to k . Thus External input 7.2 applies to R_k , and its solution fiber is the evaluation fiber ω_v , since $Y(s_0) = I$. Its group is H_v .

Lemma 7.3 (Finite coefficient descent). *For $f_1, \dots, f_r \in k[[x_1, \dots, x_d]]$, the space of complex constant relations among the f_j is the complex scalar extension of the space of k -relations.*

Proof. Expand $f_j = \sum_{\alpha} a_{\alpha j} x^{\alpha}$. A relation is the kernel of the matrix $(a_{\alpha j})$ with r columns. Its possibly infinite family of rows spans a finite-dimensional subspace of k^r . Choose finitely many rows that are a basis of this row span. They define the same kernel, and kernels of finite matrices commute with field extension. This proves the claim without a field automorphism or a continuity assumption on one. $\square$

Proposition 7.4 (No extra relations after complex constant extension). *Multiplication of meromorphic functions gives an isomorphism*

$$R_k \otimes_K E \xrightarrow{\sim} E[Y, \det(Y)^{-1}] =: R_E. \quad (7.2)$$

The algebra R_E is Picard–Vessiot over E with group $(H_v)_{\mathbb{C}}$.

Proof. Surjectivity is immediate. To prove injectivity, take a finite expression in the kernel. Choose an affine neighborhood of s_0 defined over k . Every coefficient in E is a quotient of finite complex linear combinations of regular k -functions. Clear a common nonzero denominator of this type. The alleged relation becomes a finite expression

$$\sum_{j=1}^r c_j r_j = 0, \quad c_j \in \mathbb{C}, \quad r_j \in R_k.$$

Multiply once more by a nonzero element of K , if necessary, to remove the finitely many poles at s_0 of the r_j . Their expansions are now in $k[[x_1, \dots, x_d]]$ by Lemma 7.1. By Lemma 7.3, the vector (c_j) is a complex linear combination of k -relation vectors. Those relations already vanish in R_k . Therefore the tensor expression was zero. Reversing the invertible denominator multiplications proves injectivity.

The fraction field of R_E is a subfield of $\mathcal{M}(B)$ containing $\mathbb{C}$, so its common constants are exactly $\mathbb{C}$. It is generated by a fundamental matrix, hence is Picard–Vessiot by External input 7.2. For the group identification, the Picard–Vessiot torsor identity for R_k is

$$R_k \otimes_K R_k \simeq R_k \otimes_k k[H_v].$$

Tensor it with E and use (7.2). This is the torsor identity for R_E with group $(H_v)_{\mathbb{C}}$. The action sends Y to Yh . Equivalently, the matrix relation ideal and its tensor symmetries have just been shown to commute with this constant extension. $\square$

Remark 7.5 (Why this base-change argument is needed). An exact faithful functor on the original k -category does not by itself identify every later complex differential subobject. Here that step is proved using a normalized solution with algebraic Taylor coefficients. Actual period matrices need not have algebraic Taylor coefficients; they are introduced only after Proposition 7.4.

8 The torsor and the exact constant-field argument

We now end with the complete comparison and descent argument. The mathematics of this section is also an abstract implication: it applies to any finite-type neutral Tannakian system with the fiber functors and the exactness established in Section 6, and with the algebraic connection realization used in Section 7.

8.1 The actual constant-period base

Let $\omega_{\mathrm{dR},0} : \mathcal{T}_0 \rightarrow \mathrm{Vec}_k$ be the constant de Rham fiber of Corollary 3.3. Define the comparison torsors

$$\begin{aligned} X &= \mathrm{Isom}^{\otimes}(\omega_{\mathrm{B}} \otimes_k K, \omega_{\mathrm{dR}}), \\ X_0 &= \mathrm{Isom}^{\otimes}(\omega_{\mathrm{B}}|_{\mathcal{T}_0}, \omega_{\mathrm{dR},0}). \end{aligned}$$

Thus X is over K , whereas X_0 is over k . In a fiber product the base change $X_{0,K}$ is always understood explicitly. These are affine torsors by [5, Theorem 3.2].

Let $p_0 \in X_0(\mathbb{C})$ be the actual constant comparison point and put

$$\begin{aligned} Z_0 &= \overline{\{p_0\}}^{\mathrm{Zar}}, & A_0 &= k[Z_0] = k[X_0]/\ker(\mathrm{ev}_{p_0}), \\ F_0 &= \mathrm{Frac}(A_0), & R_0 &= K \otimes_k A_0, & K_0 &= \mathrm{Frac}(R_0) = F_0(S). \end{aligned} \tag{8.1}$$

Both A_0 and F_0 are identified with their actual embedded values in $\mathbb{C}$. This imposes their relations without asserting that the unknown constant-period relations are absent. All derivations kill F_0 .

Lemma 8.1 (Domains, embeddings, and base constants). *There are injective maps*

$$R_0 \hookrightarrow K_0 \hookrightarrow E = \mathbb{C}(S),$$

and the common constant field of K_0 is exactly F_0 .

Proof. Evaluation makes A_0 a domain. The smooth connected variety S is geometrically integral over the algebraically closed field k . After any field extension its function-field construction is therefore a domain. In particular $K \otimes_k \mathbb{C}$ is a domain whose fraction field is E . Flatness of K over k embeds R_0 in $K \otimes_k \mathbb{C}$, giving the claimed injections and identifying K_0 with $F_0(S)$.

The derivations ∂_i form a basis of the tangent derivations of K_0/F_0 . If they all kill $u \in K_0$, then $du = 0$ in Ω_{K_0/F_0}^1 . In characteristic zero this implies that u is algebraic over F_0 . Geometric integrality of S_{F_0} implies that F_0 is relatively algebraically closed in $F_0(S)$, so $u \in F_0$. The reverse inclusion is part of the definition of the derivations. $\square$

Set

$$X_{\mathrm{rel}} = X \times_{X_{0,K}} (\mathrm{Spec} K \times_k Z_0), \quad B_0 = \mathcal{O}(X_{\mathrm{rel}}). \tag{8.2}$$

Proposition 8.2 (The correct group and its torsor). *The map $X_{\mathrm{rel}} \rightarrow \mathrm{Spec} R_0$ is a torsor under H_{B,R_0} . Its coordinate algebra B_0 is faithfully flat and torsion-free over R_0 .*

Proof. The normal subgroup H_B of (6.1) is defined over k . The map $X \rightarrow X_{0,K}$ is an $H_{B,K}$ -torsor: after a faithfully flat trivialization of X , it is the quotient map $G_B \rightarrow G_{0,B}$, whose kernel is H_B . Descend this assertion and base change along $\mathrm{Spec} K \times_k Z_0 \rightarrow X_{0,K}$. An affine torsor under an affine algebraic group is faithfully flat. Since R_0 is a domain, flatness implies torsion-freeness. $\square$

There is a natural differential structure on B_0 . For a tensor-generator connection in the frame of (7.1), its universal comparison matrix U satisfies

$$\partial_i U = A_i U. \quad (8.3)$$

Naturality and tensor relations are stable under these operators because the de Rham morphisms and tensor structure are horizontal. Constant comparison coefficients are horizontal; hence restriction along Z_0 also respects the derivations. Thus actual analytic comparison defines a differential map $B_0 \rightarrow \mathcal{M}(B)$ without yet claiming it injective.

8.2 An explicit equivariant comparison over $\mathbb{C}(S)$

Use the tensor generator, algebraic frame, and normalized matrix Y of Section 7. Let P be its actual comparison matrix from Betti to de Rham coordinates, and let $c = P(s_0)$ be the matrix of the comparison in Lemma 6.4. Uniqueness of solutions gives

$$P = Yc. \quad (8.4)$$

The coefficients of c belong to $\mathbb{C}$ and need not belong to F_0 . This fact is allowed; those coefficients are not adjoined to the smaller solution field in the descent argument.

Lemma 8.3 (The analytic algebra and equivariance). *Evaluation gives a surjective E -algebra map*

$$B_0 \otimes_{R_0} E \longrightarrow E[P, \det(P)^{-1}] = R_E. \quad (8.5)$$

The induced closed immersion of schemes is equivariant for $H_{B,E}$, under the identification $(H_B)_{\mathbb{C}} = c^{-1}(H_v)_{\mathbb{C}}c$.

Proof. The actual comparison is natural and tensor-compatible. Its restriction to constant objects is p_0 , so it factors through X_{rel} . Because $\mathcal{T}$ has a tensor generator, its comparison algebra is generated by the entries of the generator's universal matrix and its inverse determinant, with its tensor relations. Evaluation therefore has precisely the image on the right of (8.5), and is surjective. By (8.4), $E[P, \det(P)^{-1}] = E[Y, \det(Y)^{-1}]$.

The group of the latter Picard–Vessiot algebra acts by $Y \mapsto Yh$, $h \in (H_v)_{\mathbb{C}}$. Consequently its action on P is

$$P = Yc \longmapsto Yhc = P(c^{-1}hc).$$

By Lemma 6.4, $c^{-1}hc$ ranges over $(H_B)_{\mathbb{C}}$. On the motivic comparison torsor, precomposition by a Betti tensor automorphism is exactly right multiplication $U \mapsto Ug$. The two formulas coincide. They are identities for the universal matrix coordinate functions, so they establish equivariance as group-scheme actions, not only on a selected set of analytic points. $\square$

Lemma 8.4 (Equivariant torsor maps). *An equivariant morphism between two nonempty torsors under the same affine group over a field is an isomorphism.*

Proof. Make a faithfully flat extension that trivializes both torsors and choose origins. An equivariant map between the two right regular group torsors is left translation by the image of the identity, hence is invertible. The inverse descends, since invertibility of a morphism is faithfully flat local on the base. $\square$

Theorem 8.5 (Injective relative period comparison). *The actual map $B_0 \rightarrow \mathcal{M}(B)$ is injective. Moreover*

$$B_0 \otimes_{R_0} \mathbb{C}(S) \simeq R_E \quad (8.6)$$

as differential algebras and equivariant torsor coordinate algebras.

Proof. By Proposition 7.4, the source scheme of the closed immersion in Lemma 8.3 is a Picard–Vessiot torsor under $H_{B,E}$, in the Betti coordinates just computed. By Proposition 8.2, its target is a torsor under the same group. Lemma 8.4 makes that immersion an isomorphism. This proves (8.6).

By Proposition 8.2 and Lemma 8.1, the actual period map is the composite of injections

$$B_0 \hookrightarrow B_0 \otimes_{R_0} K_0 \hookrightarrow B_0 \otimes_{R_0} E \xrightarrow{\sim} R_E \hookrightarrow \mathcal{M}(B).$$

The first injection is localization of a torsion-free module over a domain. The second is scalar extension along the field embedding $K_0 \hookrightarrow E$, hence is injective. The last is the concrete analytic Picard–Vessiot embedding. This proves injectivity without assuming anything about the constant field of the smaller period-generated solution field. $\square$

8.3 Descent of differential simplicity and constants

Put

$$R = B_0 \otimes_{R_0} K_0, \quad L = \text{Frac}(R).$$

These are domains by Theorem 8.5. In particular $R = K_0[P, \det(P)^{-1}]$ in the actual analytic embedding, not an algebra obtained by adjoining the separate entries of c .

Lemma 8.6 (Intersection in a faithfully embedded tensor product). *Let $F \subset L_1, L_2$ be fields. In $L_1 \otimes_F L_2$, equality $x \otimes 1 = 1 \otimes y$ with $x \in L_1$ and $y \in L_2$ implies $x = y \in F$. The same conclusion holds if the equality is checked in any field into which this tensor product injects.*

Proof. If $x \notin F$, the vectors $1, x$ are linearly independent over F . Choose an F -linear functional $\ell : L_1 \rightarrow F$ with $\ell(1) = 0$ and $\ell(x) = 1$, extending from their span to a basis. Applying $\ell \otimes \text{id}$ to the equality gives $1 = 0$, a contradiction. Thus $x \in F$, and the equality then implies $y = x$ by injectivity of the field into its scalar extension. The last assertion reduces to this one by the assumed injection. $\square$

Theorem 8.7 (Picard–Vessiot descent over the actual constant field). *The algebra R is differentially simple, its fraction field satisfies*

$$L^{\partial_1 = \dots = \partial_d = 0} = F_0, \quad (8.7)$$

and R is the Picard–Vessiot algebra for the realized tensor generator over $K_0 = F_0(S)$, with group $(H_B)_{F_0}$.

Proof. By Theorem 8.5, $R \otimes_{K_0} E \simeq R_E$ is differentially simple. If $I \subset R$ is a nonzero differential ideal, faithful flatness of the field extension makes $I \otimes E$ nonzero. Simplicity makes it all of R_E . Applying faithful flatness to $(R/I) \otimes E = 0$ gives $I = R$. Thus R is differentially simple.

To prove (8.7), localize the isomorphism $R \otimes_{K_0} E \simeq R_E$ at every nonzero element of R . Since all those elements map to nonzero elements of the domain R_E , this gives an injection

$$L \otimes_{K_0} E \hookrightarrow \text{Frac}(R_E) \hookrightarrow \mathcal{M}(B). \quad (8.8)$$

Let $b \in L$ be killed by every ∂_i . Its analytic image is a constant $\alpha \in \mathbb{C} \subset E$: on a connected polydisc a meromorphic function with all partial derivatives zero is constant. In the last field of (8.8) the two elements $b \otimes 1$ and $1 \otimes \alpha$ have equal image. The injection gives their equality in the tensor product. Lemma 8.6 implies $b \in K_0$. Its derivatives are zero there, so Lemma 8.1 gives $b \in F_0$. Conversely, F_0 is killed by all the derivations.

Finally R is generated over K_0 by the actual fundamental matrix P and its inverse determinant, it is differentially simple, and its fraction field has exactly the constants F_0 . These are the Picard–Vessiot properties over this possibly non-algebraically-closed constant field. The already established torsor coaction is defined over F_0 and identifies its group with $(H_B)_{F_0}$; equivalently this identification follows after the faithfully flat constant extension to $\mathbb{C}$ from Lemma 8.3 and descends. This completes the constant-field argument rather than presupposing it. $\square$

For $d = 0$, $S = \text{Spec } k$, the relative group is trivial and the argument reduces to the embedding $A_0 \hookrightarrow \mathbb{C}$; set $K_0 = L = F_0$. For $d > 0$, the proofs above use every base derivation. Injectivity on B implies injectivity on any common analytic continuation domain, so the same result holds on the universal-cover formulation of the period map. Every finite packet in the generic envelope lies in a category of the form used here. Thus these replacement arguments give the intended generic functional conclusion, subject to the explicitly identified external inputs and their realization interfaces. They give neither a new algebraic-fiber specialization theorem nor an elementary presentation of all motivic relations. A fresh audit must check those interfaces and the complete sequence of implications before this note is integrated into Paper IV.

Research disclosure

The authors used AI-assisted tools for proof development, source checking, and document preparation. The authors are responsible for the note. The frozen baseline and the repair ledger record the distinction between a proposed replacement proof, a build check, and an independent mathematical audit.

References

- [1] Y. André, *Mumford–Tate groups of mixed Hodge structures and the theorem of the fixed part*, *Compositio Math.* **82** (1992), 1–24.
- [2] J. Ayoub and L. Barbieri-Viale, *Nori 1-motives*, *Math. Ann.* **361** (2015), 367–402. [arXiv:1206.5923](https://arxiv.org/abs/1206.5923).
- [3] L. Barbieri-Viale and V. Srinivas, *Albanese and Picard 1-motives*, *Mém. Soc. Math. Fr. (N.S.)* **87** (2001), vi+104 pp.
- [4] T. Crespo, Z. Hajto, and R. Mohseni, *Real Liouvillian extensions of partial differential fields*, *SIGMA* **17** (2021), 095, especially §2.1 and its comparison with Kolchin’s integrable-system theory. [doi:10.3842/SIGMA.2021.095](https://doi.org/10.3842/SIGMA.2021.095).

- [5] P. Deligne and J. S. Milne, *Tannakian categories*, in *Hodge Cycles, Motives, and Shimura Varieties*, Lecture Notes in Math. **900**, Springer, 1982, 101–228. Author-hosted corrected text: <https://www.jmilne.org/math/xnotes/tc.pdf>.
- [6] A. Douai and C. Sabbah, *Gauss–Manin systems, Brieskorn lattices and Frobenius structures (I)*, Ann. Inst. Fourier **53** (2003), 1055–1116, Propositions 1.18 and 1.20.
- [7] M. Gallauer and S. Pepin Lehalleur, *Exponentiation of coefficient systems and exponential motives*, [arXiv:2211.17247](https://arxiv.org/abs/2211.17247).
- [8] A. Huber and G. Wüstholz, *Transcendence and Linear Relations of 1-Periods*, Cambridge Tracts in Mathematics **227**, Cambridge University Press, 2022. [arXiv:1805.10104](https://arxiv.org/abs/1805.10104), Theorem 9.14.
- [9] F. Ivorra and S. Morel, *The four operations on perverse motives*, J. Eur. Math. Soc. **26** (2024), 4191–4272.
- [10] E. Jacobsen and L. Terenzi, *A comparison of categories of Nori motivic sheaves*, [arXiv:2509.21476](https://arxiv.org/abs/2509.21476), Corollary 5.15.
- [11] E. R. Kolchin, *Picard–Vessiot theory of partial differential fields*, Proc. Amer. Math. Soc. **3** (1952), 596–603. doi:10.1090/S0002-9939-1952-0049883-8.
- [12] Y.-H. J. Lam and D. Litt, *Geometric local systems on the projective line minus four points*, Compos. Math. **161** (2025), 536–554, Lemma 3.5. doi:10.1112/S0010437X24007620.
- [13] A. Mostaedi, *On the invariant cycle theorem for families of Nori motives*, [arXiv:2206.05582](https://arxiv.org/abs/2206.05582), §4.
- [14] M. van der Put and M. F. Singer, *Galois Theory of Linear Differential Equations*, Grundlehren Math. Wiss. **328**, Springer, 2003, Chapter 1. doi:10.1007/978-3-642-55750-7.
- [15] C. Sabbah, *Monodromy at infinity and Fourier transform II*, Publ. Res. Inst. Math. Sci. **42** (2006), 803–835. Proposition 5.8, Corollary 5.20, and (5.21)–(5.22). Author-hosted text: https://perso.pages.math.cnrs.fr/users/claude.sabbah/articles/sabbah_Fourier-twistor.pdf.
- [16] C. Sabbah, *A short proof of a theorem of Cotti, Dubrovin and Guzzetti*, Port. Math. **79** (2022), 29–43. [arXiv:2103.16878v3](https://arxiv.org/abs/2103.16878v3), Appendix B, preprint pp. 12–13 (the assertion immediately before Proposition B.1 and that proposition).
- [17] B. Snodgrass, *Periods of E-operators*, [arXiv:2608.06005](https://arxiv.org/abs/2608.06005) (2026), Corollary 2.17 and Theorem 3.6.
- [18] L. Terenzi, *Tensor structure on perverse Nori motives*, Ann. K-Theory **11** (2026), 47–170. [arXiv:2401.13547](https://arxiv.org/abs/2401.13547).
- [19] S. Tubach, *On the Nori and Hodge realisations of Voevodsky motives*, Compos. Math. **161** (2025), 2155–2201. [arXiv:2309.11999](https://arxiv.org/abs/2309.11999).
- [20] R. Virk, *Realizations of exponential sheaves and Fourier transform*, [arXiv:2605.14904v4](https://arxiv.org/abs/2605.14904v4) (2026), §§2.6–2.7, 6.7, Theorems 6.13, 7.5–7.6, and Proposition 7.2.