Optimal Allocations for Gamma Completion and MDS Recovery

Christopher D. Long

October 11, 2026

Abstract

We optimize the probability that at least k of n independent Gamma requirements finish by a prescribed deadline, subject to a fixed total allocation rate. For a common shape greater than one, every maximizing allocation is uniform on its positive support. We completely classify the resulting support phases for $k \geq 2$: two support curves either are strictly ordered at every positive budget or have a unique transverse crossing in favor of the larger support. A continuous interpolation is strictly log-concave in the logarithm of support size. Consequently the optimal supports form a consecutive staircase from the small-budget winner through full support, with no re-entry or skipped transition and exactly two adjacent supports at each phase boundary. The allocation proof combines pivotal probabilities, log-concavity of Bernoulli convolutions, and a mixed Gamma tail-ratio bound; the support proof uses cumulative-hazard geometry, exponential order statistics, and the Prékopa–Leindler inequality. Shapes below one have a unique full-uniform optimum. At shape one, the same holds for $k \geq 2$, while every allocation ties for $k = 1$. For shapes greater than one and $k = 1$, one active coordinate is optimal. Integer shapes give Poissonized MDS recovery. We also derive exact local stability thresholds and sufficient conditions for other common recovery responses.

1 Introduction

A collection of types shares a fixed resource budget. Each type is recovered once its random requirement has been met, and success means that at least k types have recovered. Is it best to distribute the resource evenly, or to sacrifice some types and protect the rest more intensively? We determine both the form of the optimum and its complete support phases. For independent Gamma requirements with a common shape greater than one, every optimum has equal positive allocations and zero allocations elsewhere. We call this form *active-uniform*.

For Poisson read counts, an integer Gamma shape t is equivalent to requiring at least t reads of each recovered type. The cardinality condition is the recovery rule of an outer $[n, k]$ MDS code. The theorem thus applies to the nonuniform coverage-depth problem in DNA storage, but its continuous-shape formulation is independent of that application. Theorem 2.1 gives the full optimizer classification, including the different behavior at shapes at or below one.

The support phases have the following complete description. Here F_τ denotes the distribution function of a unit-rate Gamma variable of shape τ.

Theorem 1.1 (Complete Gamma MDS support staircase)**.** *Fix a real $\tau > 1$ and integers $n \geq k \geq 2$. Put*

$$V_q(M) = \mathbb{P}(\mathrm{Bin}(q, F_\tau(M/q)) \geq k),$$
$$c_{k,\tau}(q) = \frac{\binom{q}{k}}{q^{\tau k}}, \qquad q_0 = \max \arg\max_{k \leq q \leq n} c_{k,\tau}(q).$$

For each $q_0 \leq q < n$, the curves V_q, V_{q+1} have a unique crossing $\beta_q > 0$. If $q_0 < n$, these satisfy

$$0 < \beta_{q_0} < \beta_{q_0+1} < \cdots < \beta_{n-1} < \infty.$$

With $\beta_{q_0-1} = 0$ and $\beta_n = \infty$, for every $M > 0$,

$$\arg\max_{k \leq r \leq n} V_r(M) = \begin{cases} \{q\}, & \beta_{q-1} < M < \beta_q, \quad q_0 \leq q \leq n, \\ \{q, q+1\}, & M = \beta_q, \quad q_0 \leq q < n. \end{cases}$$

For independent $B_i \sim \mathrm{Bernoulli}(F_\tau(\lambda_i))$, all allocations maximizing $\mathbb{P}(\sum_{i=1}^n B_i \geq k)$ subject to $\lambda_i \geq 0$ and $\sum_{i=1}^n \lambda_i = M$ are exactly the permutations of

$$(\underbrace{M/q, \ldots, M/q}_{q \text{ entries}}, 0, \ldots, 0)$$

for the support sizes in this set. Every support from q_0 through n is uniquely optimal on a nonempty open budget interval; supports below q_0 are never optimal. If $q_0 = n$, the crossing list is empty and full uniformity is uniquely optimal for every $M > 0$.

The proof is in Section 4. Thus no optimal support can re-enter after losing optimality, and no transition after q_0 skips a support. The initial support itself can exceed k: for $k = 10, \tau = 2, n \geq 12$, one has $q_0 = 12$ (Example 3.3).

Proof and consequences. Conditioning on all but two Bernoulli recovery indicators produces a pair objective determined by two adjacent probabilities of the remaining sum. Its stationary equations and log-concavity restrict a nonuniform local maximum to one small coordinate and equal larger coordinates. A mixed Gamma tail-ratio inequality then gives a strictly positive second derivative in a feasible direction, excluding that last pattern. The argument is analytic for every real shape greater than one; it is not an extrapolation from computations at integer shapes.

Section 3 gives the endpoint support laws and the local stability criterion. The global support proof in Section 4 uses a different consequence of the Gamma tail ratios: geometric concavity of the cumulative hazard. Its inverse transfers a strict quantile comparison from exponential order statistics, proving uniqueness and orientation of every pairwise crossing. Exponential spacings and the Prékopa–Leindler inequality give strict concavity in logarithmic support, which orders the adjacent crossings and excludes skipped phases. Section 5 isolates sufficient hypotheses on a common recovery curve, and Section 6 gives a perturbation bound for finite comparisons.

Related work and scope. Nonuniform coupon collection and its links to caching were developed in [6]. Equal non-null probabilities are stochastically optimal for collecting a prescribed number of distinct types [3]; this supplies the first-copy benchmark. The DNA coverage study

[4] states its uniform-allocation theorem for the first-copy case. Computational methods for collecting multiple copies of any prescribed number of types are given in [2]. Our companion paper [8] treats the separable expected-defect objective and proves its exact support staircase. That is a different objective, and neither its optimizer nor its phase diagram automatically optimizes deadline success.

Equal-positive-coordinate extremizers also arise for weighted-exponential sum tails in the MISO outage theorem [1]; our objective is instead a Gamma order-statistic tail. Gamma order-statistic comparisons [11], Mills-ratio inequalities [5], and incomplete-Gamma bounds [10] provide additional context. We use log-concavity of Bernoulli convolutions [7], and prove the particular mixed tail-ratio estimate required here directly. The theorem concerns a common recovery response and an unweighted budget; heterogeneous requirements, fixed-read multinomial optimization beyond the first-copy case, and optimization of expected completion time require separate arguments.

2 MDS success-probability optimization

MDS success depends on the entire distribution of the number of recovered types. Its maximum is nevertheless attained by an active-uniform allocation. We prove this assertion together with a complete description of the maximizing allocations, including the constant-objective exception. The argument applies to a common Gamma recovery curve of any positive shape; the Poisson threshold model is its integer-shape specialization.

For $\tau > 0$, define

$$g_\tau(x) = \frac{x^{\tau-1}e^{-x}}{\Gamma(\tau)}, \qquad F_\tau(x) = \frac{1}{\Gamma(\tau)} \int_0^x s^{\tau-1}e^{-s}\, ds \quad (x > 0),$$

and set $F_\tau(0) = 0$. Thus F_τ is the distribution function of a unit-rate Gamma random variable with shape τ. For every integer $t \geq 1$, integration by parts gives

$$F_t(x) = 1 - e^{-x} \sum_{j=0}^{t-1} \frac{x^j}{j!} = \phi_t(x).$$

For a q-coordinate allocation, write

$$\mathcal{S}_{q,k,\tau}(\lambda_1, \ldots, \lambda_q) = \mathbb{P}\left(\sum_{i=1}^{q} B_i \geq k \right), \qquad B_i \sim \text{Bernoulli}(F_\tau(\lambda_i))$$

with independent Bernoulli variables. The constraint is $\lambda_i \geq 0$, $\sum_i \lambda_i = M$. In particular, when $\tau = t$ is an integer, this notation agrees with the Poissonized MDS success objective.

Theorem 2.1 (Gamma active-uniform theorem)**.** *Fix $n \geq 1$, $1 \leq k \leq n$, $\tau > 0$, and $M > 0$. Then*

$$\max_{\lambda_i \geq 0,\ \sum_{i=1}^n \lambda_i = M} \mathcal{S}_{n,k,\tau}(\lambda) = \max_{q=k,\ldots,n} \mathbb{P}\left(\text{Bin}(q, F_\tau(M/q)) \geq k\right).$$

The maximizing allocations are classified as follows.

1. *If $\tau > 1$, every maximizer is uniform on its positive support. More precisely, the complete set of maximizers consists of the permutations of*

$$(\underbrace{M/q, \ldots, M/q}_{q \text{ coordinates}}, 0, \ldots, 0)$$

for support sizes q attaining the finite maximum above.

2. *If $0 < \tau < 1$, the unique maximizing vector is $\lambda_1 = \cdots = \lambda_n = M/n$, for every k.*

3. *If $\tau = 1$ and $k \geq 2$, the unique maximizing vector is again full uniformity.*

4. *If $\tau = 1$ and $k = 1$, every allocation has success probability $1 - e^{-M}$.*

For $\tau > 1$ and $k = 1$, the maximizing support has size one. For $k = n$, full uniformity is the unique maximizer for every $\tau > 0$.

Theorem 2.2 (Poissonized active-uniform MDS success theorem)**.** *Fix integers $n \geq 1$, $1 \leq k \leq n$, and $t \geq 1$, and let $M > 0$. For independent $B_i \sim \mathrm{Bernoulli}(\phi_t(\lambda_i))$,*

$$\boxed{\max_{\lambda_i \geq 0,\ \sum_{i=1}^n \lambda_i = M} \mathbb{P}\left(\sum_{i=1}^n B_i \geq k\right) = \max_{q=k,\ldots,n} \mathbb{P}\left(\mathrm{Bin}(q, \phi_t(M/q)) \geq k\right).}$$

If $t \geq 2$, every maximizing allocation is active-uniform, and its support size attains the maximum on the right. Conversely, every active-uniform allocation with such a support size is a maximizer. If $t = 1$ and $k \geq 2$, full uniformity is the unique maximizer. If $t = k = 1$, every allocation is a maximizer.

The Poissonized theorem follows from Theorem 2.1 by taking $\tau = t$. We prove the Gamma theorem below. The proof uses pivotal probabilities to control the possible active levels and then excludes asymmetric local maxima by two feasible second variations.

Remark 2.3 (Gamma completion times). Let $Y_1, \ldots, Y_n$ be independent unit-rate Gamma variables of common shape τ, and let $p_i \geq 0$, $\sum_i p_i = 1$. Define $T_i = Y_i/p_i$ when $p_i > 0$, and $T_i = \infty$ otherwise. For the k-th order statistic,

$$\mathbb{P}(T_{(k)} \leq M) = \mathcal{S}_{n,k,\tau}(Mp).$$

Thus Theorem 2.1 also optimizes the probability of completing at least k independent Gamma requirements by a fixed deadline. Integer shapes recover the Poisson threshold model. Noninteger shapes describe a Gamma requirement rather than a fractional number of required reads.

2.1 Gamma tail-to-density ratios

Fix $\tau > 0$, put $m = \tau - 1$, and temporarily write $F = F_\tau$, $g = g_\tau$. Define

$$P(z) = \frac{1 - F(z)}{g(z)}, \qquad Q(z) = \frac{F(z)}{g(z)}.$$

Changes of variables in the two Gamma integrals give, for every $m > -1$,

$$P(z) = \int_0^\infty e^{-u} \left(1 + \frac{u}{z}\right)^m du, \tag{1}$$

$$Q(z) = z \int_0^1 e^{zv} (1 - v)^m \, dv. \tag{2}$$

In particular, $P, Q > 0$ and $Q', Q'' > 0$. When $m > 0$, also $P' < 0$ and $P'' > 0$, since

$$\frac{d^2}{dz^2}\left(1 + \frac{u}{z}\right)^m = \frac{mu}{z^3}\left(1 + \frac{u}{z}\right)^{m-2}\left(2 + (m+1)\frac{u}{z}\right) > 0 \qquad (u > 0).$$

Differentiation under these integrals is justified locally on $z > 0$ by an integrable dominating function.

For $m > 0$, set

$$\eta(z) = -\frac{zP'(z)}{P(z)}, \qquad \zeta(z) = \frac{zQ'(z)}{Q(z)}.$$

Both functions are positive. Since $g'/g = m/z - 1$,

$$zP' = (z-m)P - z, \qquad zQ' = (z-m)Q + z.$$

Writing $h = z - m + 1$, differentiation yields

$$z\eta' = \eta^2 + h\eta - m, \qquad z\zeta' = -\zeta^2 + h\zeta + m. \tag{3}$$

Lemma 2.4 (Mixed Gamma tail-ratio bound)**.** *For every real $m > 0$ and $z > 0$,*

$$\frac{3m}{5} < \eta(z)\zeta(z) < m.$$

Proof. Put $W = \eta\zeta$ and $\Delta = \zeta - \eta$. From (3),

$$zW' = 2hW - (W+m)\Delta, \qquad z\Delta' = -\Delta^2 - 2W + h\Delta + 2m.$$

At a critical point of W, the first identity gives $\Delta = 2hW/(W+m)$. Differentiating the first identity and substituting this expression into the second gives

$$\frac{z^2W''}{2W} = z - \left(\frac{m^2}{W} - W\right) - \frac{m-W}{m+W}(z-m+1)^2. \tag{4}$$

If $0 < W \le 3m/5$, then

$$\frac{m^2}{W} - W \ge \frac{16m}{15}, \qquad \frac{m-W}{m+W} \ge \frac{1}{4}.$$

Consequently, at any critical point with $W \le 3m/5$,

$$\frac{z^2W''}{2W} \le z - \frac{16m}{15} - \frac{(z-m+1)^2}{4} = -\frac{(z-m-1)^2}{4} - \frac{m}{15} < 0.$$

On the other hand, the endpoint expansions of the Gamma integrals give

$$\eta(z) \longrightarrow m, \qquad \zeta(z) \longrightarrow 1 \quad (z \downarrow 0),$$

and

$$\eta(z) = \frac{m}{z} + O(z^{-2}), \qquad \zeta(z) = z - m + o(1) \quad (z \to \infty).$$

For example, at zero one has $P(z) \sim \Gamma(m+1)z^{-m}$ and $Q(z) \sim z/(m+1)$; at infinity, (1) gives $P(z) = 1 + m/z + O(z^{-2})$. Thus $W(z) \to m$ at both ends. If W reached $3m/5$ or a smaller value, it would attain an interior minimum at or below that level. This contradicts (4), proving the lower bound.

For the upper bound, use the probability density on $(0,\infty)$ proportional to $e^{-u}(1+u/z)^m$, and let $V = u/(z+u)$. Then

$$\eta = m\mathbb{E}[V], \qquad z\eta' = -m\mathbb{E}[V(1-V)] - m^2\,\mathrm{Var}(V) < 0. \tag{5}$$

Under the probability density on $(0,1)$ proportional to $e^{zv}(1-v)^m$, one similarly has

$$\zeta = 1 + z\mathbb{E}[v], \qquad \zeta' = \mathbb{E}[v] + z\,\mathrm{Var}(v) > 0.$$

Equation (3) therefore implies

$$h < \frac{m}{\eta} - \eta, \qquad h > \zeta - \frac{m}{\zeta}.$$

Combining these inequalities gives $\eta + \zeta < m(1/\eta + 1/\zeta)$, which is equivalent to $\eta\zeta < m$. □

Lemma 2.5 (Two slope monotonicities)**.** *For $\tau > 1$, the functions*

$$L_1(z) = \frac{-P'(z)}{Q'(z)}, \qquad L_2(z) = \frac{(-P'(z))Q(z)^2}{Q'(z)P(z)^2}$$

are respectively strictly decreasing and strictly increasing on $(0,\infty)$.

Proof. The signs $P' < 0, P'' > 0, Q' > 0, Q'' > 0$ show that L_1 is strictly decreasing. Also

$$L_2 = \frac{\eta}{\zeta}\frac{Q}{P}.$$

Using (3) and Lemma 2.4,

$$z\frac{d}{dz}\log L_2 = (\eta+\zeta)\left(2 - \frac{m}{\eta\zeta}\right) > \frac{\eta+\zeta}{3} > 0.$$

This proves the second assertion. □

2.2 Endpoint faces and pivotal probabilities

Lemma 2.6 (Endpoint cases)**.** *On the closed q-coordinate simplex with total mass $M > 0$, the following statements hold.*

1. *For $\tau > 1$ and $k = 1$, the maximizers are exactly the one-point allocations $(M, 0, \ldots, 0)$ and their permutations.*
2. *For every $\tau > 0$ and $k = q$, the unique maximizer is the barycenter $\lambda_i = M/q$.*

Proof. If $\tau > 1$, P is strictly decreasing, so the hazard $g/(1-F) = 1/P$ is strictly increasing. Since $F(0) = 0$, integration of the hazard gives, for $x, y > 0$,

$$1 - F(x+y) < (1 - F(x))(1 - F(y)).$$

Indeed, the integral of the hazard on $[x, x+y]$ is strictly larger than its integral on $[0, y]$. Iterating the inequality shows that

$$\mathcal{S}_{q,1,\tau}(\lambda) = 1 - \prod_i (1 - F(\lambda_i)) \le F(M),$$

with equality exactly when only one coordinate is positive.

For $k = q$, the objective is $\prod_i F(\lambda_i)$. Formula (2) gives $Q' > 0$ for every $\tau > 0$, so

$$(\log F)' = \frac{g}{F} = \frac{1}{Q}$$

is strictly decreasing. Strict Jensen's inequality therefore gives

$$\prod_{i=1}^{q} F(\lambda_i) \le F(M/q)^q,$$

with equality only at the barycenter. Boundary allocations have zero product, whereas the barycenter has positive product. □

Lemma 2.7 (Support-shadow KKT equations)**.** *Let $\tau > 0$, $1 \le k \le q$, and $\lambda_i > 0$. Set*

$$R_i(\lambda) = \mathbb{P}\left(\sum_{j \ne i} B_j = k - 1\right).$$

Then

$$\frac{\partial \mathcal{S}_{q,k,\tau}}{\partial \lambda_i} = g_\tau(\lambda_i) R_i(\lambda) > 0.$$

At an interior stationary point under $\sum_i \lambda_i = M$, these derivatives are equal for all active coordinates.

Proof. Write $\theta_i = F_\tau(\lambda_i)$ and $Z_{-i} = \sum_{j \ne i} B_j$. Conditioning on B_i,

$$\mathcal{S}_{q,k,\tau} = \theta_i \mathbb{P}(Z_{-i} \ge k - 1) + (1 - \theta_i)\mathbb{P}(Z_{-i} \ge k).$$

Hence $\partial \mathcal{S}/\partial \theta_i = \mathbb{P}(Z_{-i} = k - 1)$. The chain rule gives the asserted derivative. Every active Bernoulli parameter lies in $(0, 1)$, so the pivotal probability is positive. The equality at a constrained stationary point is the Lagrange multiplier equation. □

Lemma 2.8 (Pair-shadow reduction)**.** *Let $q \ge 3$, $2 \le k \le q - 1$, and $\tau > 0$, and work on a positive q-coordinate face. For distinct i, j, define*

$$Z_{ij} = \sum_{\ell \ne i,j} B_\ell, \quad \beta_{ij} = \mathbb{P}(Z_{ij} = k - 1), \quad \gamma_{ij} = \mathbb{P}(Z_{ij} = k - 2), \quad \alpha_{ij} = 1 - \frac{\gamma_{ij}}{\beta_{ij}}.$$

Then $\beta_{ij}, \gamma_{ij} > 0$ and $\alpha_{ij} < 1$. With all other coordinates fixed, the pair contribution to the objective, up to an additive constant, is

$$\beta_{ij}\big(F_\tau(\lambda_i) + F_\tau(\lambda_j) - \alpha_{ij} F_\tau(\lambda_i) F_\tau(\lambda_j)\big).$$

At an interior stationary point,

$$L_{\alpha_{ij}}(\lambda_i) = L_{\alpha_{ij}}(\lambda_j), \qquad L_\alpha(x) = \frac{g_\tau(x)}{1 - \alpha F_\tau(x)}.$$

Proof. The shadow Z_{ij} assigns positive probability to every integer from 0 through $q - 2$, proving the positivity assertions. If $Z_{ij} \ge k$, the pair does not affect success; if $Z_{ij} \le k - 3$, the pair cannot produce success. When $Z_{ij} = k - 1$, at least one of the pair must succeed, and when $Z_{ij} = k - 2$, both must succeed. Writing $\theta_h = F_\tau(\lambda_h)$, the nonconstant contribution is therefore

$$\beta_{ij}(\theta_i + \theta_j - \theta_i \theta_j) + \gamma_{ij} \theta_i \theta_j,$$

which is the claimed expression. The two active partial derivatives are

$$\beta_{ij} g_\tau(\lambda_i)(1 - \alpha_{ij}\theta_j), \qquad \beta_{ij} g_\tau(\lambda_j)(1 - \alpha_{ij}\theta_i).$$

Their equality is equivalent to the marginal-ratio equation, since $1 - \alpha_{ij}\theta_h > 0$. □

2.3 Marginal-ratio and shadow monotonicity

Lemma 2.9 (Marginal-ratio shape)**.** *Let $\tau > 1$, put $m = \tau - 1$, and fix $\alpha < 1$. Then*

$$L_\alpha(x) = \frac{g_\tau(x)}{1 - \alpha F_\tau(x)}$$

is strictly unimodal on $(0, \infty)$. Its unique maximizer $x_*(\alpha)$ is strictly increasing in α, and

$$x_*(\alpha) < m \quad (\alpha < 0), \qquad x_*(0) = m, \qquad x_*(\alpha) > m \quad (0 < \alpha < 1).$$

Proof. Write $g = g_\tau$, $F = F_\tau$, $u = g'/g = m/x - 1$, and $D = 1 - \alpha F > 0$. The derivative of L_α has the sign of

$$N_\alpha = uD + \alpha g.$$

The terms involving g' cancel when this expression is differentiated:

$$N'_\alpha = u'D = -\frac{m}{x^2} D < 0.$$

Moreover, $N_\alpha(0+) = +\infty$ and $N_\alpha(\infty) = -(1 - \alpha) < 0$. Thus there is exactly one zero, with L_α strictly increasing before it and strictly decreasing after it.

At the zero of N_α,

$$\partial_\alpha N_\alpha = g - uF = \frac{g}{D} > 0.$$

Since $N'_\alpha < 0$, implicit differentiation shows that $x_*(\alpha)$ is strictly increasing. Finally, $N_\alpha(m) = \alpha g(m)$, giving the stated comparison with m. □

Lemma 2.10 (Pair-shadow monotonicity)**.** *Let W be a Poisson-binomial random variable with masses p_j, extended by zero outside its support. For $0 < \theta < 1$, put*

$$\beta(\theta) = (1 - \theta)p_{k-1} + \theta p_{k-2}, \qquad \gamma(\theta) = (1 - \theta)p_{k-2} + \theta p_{k-3}.$$

Wherever $\beta(\theta) > 0$, the function $\alpha(\theta) = 1 - \gamma(\theta)/\beta(\theta)$ is nondecreasing.

Proof. Direct differentiation gives

$$\alpha'(\theta) = \frac{p_{k-2}^2 - p_{k-1}p_{k-3}}{\beta(\theta)^2} \geq 0.$$

The last inequality is log-concavity of the Poisson-binomial mass sequence. Its probability generating polynomial is a product of linear factors with nonpositive real zeros, so Newton's inequalities give the required coefficient inequality; see also [7]. □

2.4 Exclusion of asymmetric critical patterns

Throughout this subsection, assume $\tau > 1$ and $2 \leq k \leq q - 1$.

Proposition 2.11 (No three active levels)**.** *At an interior stationary point of $\mathcal{S}_{q,k,\tau}$, the active coordinates take at most two distinct values.*

Proof. Suppose that three coordinates satisfy $x_1 < x_2 < x_3$. For the pairs 12 and 23, Lemmas 2.8 and 2.9 imply

$$x_1 < x_*(\alpha_{12}) < x_2 < x_*(\alpha_{23}) < x_3.$$

After removing these three coordinates, let the remaining Bernoulli sum be W. The 12-shadow is $W + B_3$, whereas the 23-shadow is $W + B_1$. Since $F_\tau(x_3) > F_\tau(x_1)$, Lemma 2.10 gives $\alpha_{12} \geq \alpha_{23}$. The increasing peak location then gives $x_*(\alpha_{12}) \geq x_*(\alpha_{23})$, a contradiction. □

Proposition 2.12 (Two-level local maxima are one-small)**.** *If an interior local maximum has two distinct active values $x < y$, then the smaller value occurs exactly once.*

Proof. For a pair consisting of one small and one large coordinate, the stationary equation gives $L_{\alpha_{xy}}(x) = L_{\alpha_{xy}}(y)$, so

$$x < x_*(\alpha_{xy}) < y.$$

Suppose that two small coordinates exist. The shadow of an xx-pair is obtained from the shadow of an xy-pair by replacing one small Bernoulli parameter by a large one. Hence

$$\alpha_{xx} \geq \alpha_{xy}, \qquad x_*(\alpha_{xx}) \geq x_*(\alpha_{xy}) > x,$$

and therefore $L'_{\alpha_{xx}}(x) > 0$.

Split the two small coordinates into $x + \varepsilon, x - \varepsilon$, holding all others fixed. This is a feasible variation for $|\varepsilon| < x$. By the pair-shadow formula, its second derivative at zero is

$$2\beta_{xx}\big[g'_\tau(x)(1 - \alpha_{xx}F_\tau(x)) + \alpha_{xx}g_\tau(x)^2\big].$$

The bracket equals $(1-\alpha_{xx}F_\tau(x))^2 L'_{\alpha_{xx}}(x) > 0$. The feasible variation thus increases the objective to second order, contradicting local maximality. □

Lemma 2.13 (One-small critical equation)**.** *At a point $(x, y, \ldots, y)$ with $0 < x < y$, set*

$$c = q - 1, \qquad a = \frac{q-k}{q-1}, \qquad b = \frac{k-1}{q-1},$$

and let D_x, D_y be the partial derivatives with respect to the small coordinate and one large coordinate. Then

$$\frac{D_y}{D_x} = a\frac{P(x)}{P(y)} + b\frac{Q(x)}{Q(y)}.$$

In particular, the stationary equation is $K(x, y) = 0$, where

$$K(x,y) = a\frac{P(x)}{P(y)} + b\frac{Q(x)}{Q(y)} - 1.$$

Proof. Write $\theta_x = F_\tau(x)$, $\theta_y = F_\tau(y)$. The two pivotal probabilities are

$$R_x = \mathbb{P}(\mathrm{Bin}(c, \theta_y) = k - 1)$$

and

$$R_y = (1 - \theta_x)\mathbb{P}(\mathrm{Bin}(c-1, \theta_y) = k - 1) + \theta_x\mathbb{P}(\mathrm{Bin}(c-1, \theta_y) = k - 2).$$

The binomial coefficient ratios give

$$\frac{R_y}{R_x} = a\frac{1-\theta_x}{1-\theta_y} + b\frac{\theta_x}{\theta_y}.$$

Since $D_x = g_\tau(x)R_x$, $D_y = g_\tau(y)R_y$, and $P = (1 - F_\tau)/g_\tau$, $Q = F_\tau/g_\tau$, multiplication by $g_\tau(y)/g_\tau(x)$ gives the asserted identity. □

Proposition 2.14 (No asymmetric one-small local maximum)**.** *An asymmetric one-small stationary point is a strict local minimum along the feasible path on which its large coordinates remain equal. In particular, it cannot be an interior local maximum of* $\mathcal{S}_{q,k,\tau}$*.*

Proof. Use the notation of Lemma 2.13. At the stationary point, put

$$U = \frac{P(x)}{P(y)} > 1, \qquad V = \frac{Q(x)}{Q(y)} < 1, \qquad S = \frac{P(x) - P(y)}{Q(y) - Q(x)}.$$

The equation $aU + bV = 1$, together with $a + b = 1$, is equivalent to

$$\frac{b}{a} = \frac{U - 1}{1 - V} = S\frac{Q(y)}{P(y)}.$$

Cauchy's mean-value theorem and the strict decrease of L_1 in Lemma 2.5 give

$$S = L_1(\xi) < L_1(x) \qquad \text{for some } \xi \in (x, y).$$

Applying the same theorem to $1/P$ and $-1/Q$, and using the strict increase of L_2, gives

$$S\frac{Q(x)Q(y)}{P(x)P(y)} = L_2(\widetilde{\xi}) < L_2(y) \qquad \text{for some } \widetilde{\xi} \in (x, y).$$

After substitution of b/a, these inequalities become

$$aU\eta(x) > bV\zeta(x), \qquad aU\eta(y) > bV\zeta(y).$$

Thus the partial derivatives of K, with x, y regarded as independent variables, satisfy

$$K_x = \frac{-aU\eta(x) + bV\zeta(x)}{x} < 0, \qquad K_y = \frac{aU\eta(y) - bV\zeta(y)}{y} > 0.$$

Along the fixed-mass path $y = (M - x)/c$, define $G(x) = \mathcal{S}_{q,k,\tau}(x, y, \dots, y)$. There are c large coordinates, so

$$G' = D_x - D_y = -D_x K.$$

At the stationary point $K = 0$, and hence

$$G'' = -D_x\left(K_x - \frac{1}{c}K_y\right) > 0.$$

This proves the asserted strict local minimum along the path and excludes an interior local maximum on the face. □

2.5 Completion of the proof and shapes at or below one

Proof of Theorem 2.1. The objective is continuous on the compact allocation simplex, so it attains its maximum. A maximizing allocation has at least k positive coordinates: a smaller support has success probability zero, whereas the uniform allocation on k coordinates has positive success probability.

First suppose $\tau > 1$, and let $q \geq k$ be the positive support size of a maximizer. The point is an interior local maximum on that face. The endpoint cases $k = 1$ and $k = q$ are covered by Lemma 2.6. For $2 \leq k \leq q - 1$, Proposition 2.11 allows at most two active values. If two

values occur, Proposition 2.12 forces a one-small configuration, which Proposition 2.14 excludes. Therefore all positive coordinates are equal to M/q.

Every active-uniform allocation on q coordinates has value

$$\mathbb{P}\left(\mathrm{Bin}(q, F_\tau(M/q)) \geq k\right),$$

because its zero coordinates never recover. Taking the maximum over $q = k, \ldots, n$ proves the formula and the complete equality classification for $\tau > 1$. The endpoint lemma also gives the asserted one-point optimum for $k = 1$.

Now suppose $0 < \tau < 1$, so $-1 < m < 0$. Formula (1) gives $P' > 0$, while $g' < 0$. Consequently,

$$g'(1 - F) + g^2 < 0,$$

since the hazard $g/(1 - F) = 1/P$ is strictly decreasing. For $\alpha < 1$,

$$L'_\alpha = \frac{g'(1 - \alpha F) + \alpha g^2}{(1 - \alpha F)^2} < 0.$$

For $\alpha \leq 0$, both terms in the numerator have the required sign. For $0 < \alpha < 1$, write that numerator as

$$(1 - \alpha)g' + \alpha\big(g'(1 - F) + g^2\big) < 0.$$

The derivative is also strictly negative at $\alpha = 1$.

Thus, when $2 \leq k \leq q - 1$, the pair-shadow stationary equations force all active coordinates to be equal. For $k = 1$, the objective is $1 - \prod_i (1 - F(\lambda_i))$; equality of its active derivatives is equality of the strictly decreasing hazards $g(\lambda_i)/(1 - F(\lambda_i))$, which again forces equal coordinates. The case $k = q$ is already covered by Lemma 2.6.

No boundary allocation can maximize in this range of shapes. Indeed, a boundary allocation with positive success has a positive support of size $q \geq k$. Activate a zero coordinate with mass ε and subtract this mass from one positive coordinate. Its new recovery probability is

$$F_\tau(\varepsilon) = \frac{\varepsilon^\tau}{\Gamma(\tau + 1)}(1 + O(\varepsilon)).$$

The pivotal probability for the new coordinate tends to the strictly positive probability that exactly $k - 1$ of the original active coordinates recover. The loss caused by the donor perturbation is $O(\varepsilon)$, since that coordinate remains bounded away from zero. Hence the gain is a positive multiple of ε^τ, up to $o(\varepsilon^\tau)$, and it dominates the donor loss. Every maximizer therefore has full support, and the preceding stationary-point argument proves that full uniformity is the unique maximizer.

Finally, let $\tau = 1$. For $k = 1$,

$$\mathcal{S}_{n,1,1}(\lambda) = 1 - \prod_i e^{-\lambda_i} = 1 - e^{-M},$$

so every allocation is optimal. Suppose $k \geq 2$. For every $\alpha < 1$,

$$L_\alpha(x) = \frac{e^{-x}}{1 - \alpha + \alpha e^{-x}}, \qquad L'_\alpha(x) = -\frac{(1 - \alpha)e^{-x}}{(1 - \alpha + \alpha e^{-x})^2} < 0.$$

The pair-shadow equations, together with the $k = q$ endpoint, show that any maximizing positive face must be uniform. If such a face has $q < n$ coordinates, put $p = 1 - e^{-M/q}$. On activating one

zero coordinate and taking the mass from one positive coordinate, the new-coordinate derivative and donor derivative are respectively

$$\mathbb{P}(\mathrm{Bin}(q,p)=k-1), \qquad (1-p)\mathbb{P}(\mathrm{Bin}(q-1,p)=k-1).$$

Their ratio is

$$\frac{\binom{q}{k-1}}{\binom{q-1}{k-1}} = \frac{q}{q-k+1} > 1.$$

The perturbation strictly improves the objective. Thus a maximizer has full support and is uniquely uniform.

In the cases $0 < \tau \le 1$, the established full-uniform optimum is one of the active-uniform candidates, so it also proves the finite-maximum formula. For $k = n$, uniqueness for every $\tau > 0$ follows directly from Lemma 2.6. This completes the proof. □

Proof of Theorem 2.2. Apply Theorem 2.1 with $\tau = t$, using $F_t = \phi_t$. □

Theorem 2.15 (Facewise active-uniform maximum)**.** *For $M > 0$, $\tau > 0$, and $1 \le k \le q$,*

$$\max_{\lambda_i \ge 0,\ \sum_{i=1}^{q} \lambda_i = M} \mathcal{S}_{q,k,\tau}(\lambda) = \max_{r=k,\dots,q} \mathbb{P}\left(\mathrm{Bin}(r, F_\tau(M/r)) \ge k\right).$$

The maximizing allocations have exactly the classification in Theorem 2.1, with $n = q$.

Proof. Apply Theorem 2.1 with $n = q$, deleting any coordinates fixed to zero outside the face. □

Remark 2.16 (Comparison with terminal-cloud minimization). The expected-cloud theorem in [8] and Theorem 2.2 have the same active-uniform form but optimize different objectives. For the expected terminal cloud at an integer threshold $t \ge 1$, the support size minimizes

$$(n-q) + q(1 - F_t(M/q)).$$

For MDS success, it maximizes

$$\mathbb{P}\left(\mathrm{Bin}(q, \phi_t(M/q)) \ge k\right).$$

These support sizes need not agree. The expected-cloud support size has an exact staircase [8]; Theorem 1.1 proves the MDS staircase by a separate comparison of binomial-tail probabilities. Neither staircase follows from the other objective's optimizer classification.

3 Support selection and local stability

Theorem 2.1 reduces the allocation problem to the finite family

$$V_q(M) = \mathbb{P}(\mathrm{Bin}(q, F_\tau(M/q)) \ge k), \qquad k \le q \le n. \tag{6}$$

The support selected by this binomial tail can differ from the support that minimizes the expected terminal cloud. We first give its small- and large-depth limits and a local test that eliminates unstable candidates. Section 4 then proves all global crossings and the complete support staircase. Throughout the asymptotic statements, n, k, τ are fixed.

3.1 Small-depth support selection

Proposition 3.1 (Small-depth expansion and tie-breaking)**.** *Let $n \geq k \geq 1$ and $\tau > 1$. Uniformly over the finite set $q \in \{k, \dots, n\}$, as $M \downarrow 0$,*

$$V_q(M) = \frac{\binom{q}{k}}{\Gamma(\tau+1)^k q^{\tau k}} M^{\tau k} \left[1 - \frac{\tau k}{(\tau+1)q} M + O(M^{\min\{2,\tau\}})\right]. \tag{7}$$

Put

$$c_{k,\tau}(q) = \frac{\binom{q}{k}}{q^{\tau k}}, \qquad q_0 = \max \arg\max_{k \leq q \leq n} c_{k,\tau}(q). \tag{8}$$

There exists $M_0 > 0$ such that q_0 is the unique maximizing support size in (6) whenever $0 < M < M_0$. Thus a tie between leading coefficients is resolved in favor of the larger support.

Proof. Expanding the exponential in the defining Gamma integral gives

$$F_\tau(z) = \frac{z^\tau}{\Gamma(\tau+1)} \left[1 - \frac{\tau}{\tau+1} z + O(z^2)\right].$$

For fixed $q \geq k$, the binomial tail at a small success probability p satisfies

$$\mathbb{P}(\mathrm{Bin}(q,p) \geq k) = \binom{q}{k} p^k (1 + O(p)).$$

Substitute $p = F_\tau(M/q)$. The linear correction comes from $F_\tau(M/q)^k$; the remaining relative errors have orders M^2 and M^τ. This proves (7), and finiteness of the set of supports makes the error uniform in q.

Any support with a smaller leading coefficient is eventually inferior to every support attaining the largest coefficient. If two such coefficients tie and $q_1 < q_2$, then the linear terms in (7) give

$$V_{q_2}(M) - V_{q_1}(M) = \frac{c_{k,\tau}(q_1)}{\Gamma(\tau+1)^k} \frac{\tau k}{\tau+1} \left(\frac{1}{q_1} - \frac{1}{q_2}\right) M^{\tau k+1} + o(M^{\tau k+1}) > 0$$

for sufficiently small $M > 0$. Here $\min\{2, \tau\} > 1$ is essential. A final minimum over the finitely many comparison intervals proves the assertion. □

Proposition 3.2 (Continuous localization of the leading support)**.** *Let $\tau > 1$. If $k = 1$, the coefficient $c_{1,\tau}(q) = q^{1-\tau}$ is strictly decreasing, so $q_0 = 1$. For $k \geq 2$, write*

$$H_k = \sum_{j=1}^{k} \frac{1}{j}, \qquad \binom{x}{k} = \frac{1}{k!} \prod_{j=0}^{k-1} (x - j), \quad x \geq k.$$

If $H_k \leq \tau$, the unique leading-coefficient maximizer is $q_0 = k$. If $H_k > \tau$, there is a unique $x_{k,\tau} > k$ satisfying

$$\sum_{j=0}^{k-1} \frac{x_{k,\tau}}{x_{k,\tau} - j} = \tau k. \tag{9}$$

Every integer maximizer of $c_{k,\tau}$ lies in

$$\{\kappa_{k,n}(\lfloor x_{k,\tau} \rfloor), \kappa_{k,n}(\lceil x_{k,\tau} \rceil)\}, \qquad \kappa_{k,n}(r) = \min\{n, \max\{k, r\}\}. \tag{10}$$

Proof. For real $x \geq k$,

$$\frac{d}{d\log x}\log c_{k,\tau}(x) = D_{k,\tau}(x) := \sum_{j=0}^{k-1}\frac{x}{x-j} - \tau k. \tag{11}$$

For $k \geq 2$,

$$D'_{k,\tau}(x) = -\sum_{j=1}^{k-1}\frac{j}{(x-j)^2} < 0, \qquad D_{k,\tau}(k) = k(H_k - \tau), \qquad \lim_{x\to\infty} D_{k,\tau}(x) = k(1-\tau) < 0.$$

These facts give the asserted monotonicity or the unique continuous maximum. A function that increases up to one point and decreases afterward attains its maximum on an integer interval at one of the two integers adjacent to that point, with the indicated endpoint clipping. □

Example 3.3 (The small-depth support can exceed the decoding threshold)**.** Take $k = 10$, $\tau = 2$, and any fixed $n \geq 12$. Exact evaluation of (11) gives

$$D_{10,2}(11) = \frac{5591}{2520} > 0, \qquad D_{10,2}(12) = -\frac{1759}{2310} < 0.$$

The only possible integer leading-coefficient maximizers are therefore 11 and 12. Their exact ratio is

$$\frac{c_{10,2}(12)}{c_{10,2}(11)} = 6\left(\frac{11}{12}\right)^{20} > 1,$$

where the last inequality is the integer comparison $6 \cdot 11^{20} > 12^{20}$. Consequently the unique optimal support is $q = 12$ for all sufficiently small positive M. Relative to using only $k = 10$ coordinates, its leading coefficient is larger by the factor

$$\frac{c_{10,2}(12)}{c_{10,2}(10)} = 66\left(\frac{5}{6}\right)^{20} > 1.$$

Thus the possibility of sacrificing coordinates does not imply that the smallest feasible support is optimal at small depth.

3.2 Large-depth support selection

Proposition 3.4 (Eventual full support)**.** *Let $\tau > 0$, $n \geq k \geq 1$, and $k \leq q \leq n$. Put $r = q-k+1$. As $M \to \infty$,*

$$1 - V_q(M) \sim \binom{q}{k-1}\frac{(M/q)^{(\tau-1)r}}{\Gamma(\tau)^r}e^{-rM/q}. \tag{12}$$

If $k \geq 2$, there is $M_1 < \infty$ such that $q = n$ is the unique maximizing support size for every $M > M_1$. In that range the unique maximizing allocation is $\lambda_i = M/n$.

Proof. The upper Gamma tail satisfies

$$1 - F_\tau(z) \sim \frac{z^{\tau-1}e^{-z}}{\Gamma(\tau)}.$$

For completeness, this follows by applying l'Hôpital's rule to $\int_z^\infty s^{\tau-1}e^{-s}\,ds$ divided by $z^{\tau-1}e^{-z}$; the ratio of their derivatives tends to one. Let $u = 1 - F_\tau(M/q)$. Failure on a q-point support means at least r unsuccessful coordinates, and hence

$$1 - V_q(M) = \mathbb{P}(\mathrm{Bin}(q,u) \geq r) \sim \binom{q}{r} u^r.$$

This proves (12) because $\binom{q}{r} = \binom{q}{k-1}$.

For $k \geq 2$, the exponential decay rate of failure,

$$\frac{r}{q} = 1 - \frac{k-1}{q},$$

strictly increases with q. The failure probability at $q = n$ is therefore smaller than that at every $q < n$ for sufficiently large M, regardless of the fixed polynomial factors in (12). There are only finitely many supports. The allocation statement now follows from Theorem 2.1. □

The restriction $k \geq 2$ in the support conclusion matters. For $k = 1$, Theorem 2.1 gives one active coordinate when $\tau > 1$, full uniformity when $0 < \tau < 1$, and the allocation-independent value $1 - e^{-M}$ when $\tau = 1$.

3.3 The constrained Hessian at an active barycenter

For $\tau > 1$, set $m = \tau - 1$ and define

$$\eta_\tau(z) = m - z + \frac{z g_\tau(z)}{1 - F_\tau(z)}, \qquad \zeta_\tau(z) = z - m + \frac{z g_\tau(z)}{F_\tau(z)}. \tag{13}$$

These are the logarithmic derivatives $-zP'(z)/P(z)$ and $zQ'(z)/Q(z)$ of the Gamma tail-to-density ratios in Section 2. The proof there gives

$$\eta_\tau \text{ strictly decreasing from } m \text{ to } 0, \qquad \zeta_\tau \text{ strictly increasing from } 1 \text{ to } \infty.$$

Proposition 3.5 (Barycenter stability and its threshold)**.** *Let $\tau > 1$ and $2 \leq k \leq q-1$. At the barycenter of a positive q-face, write*

$$z = M/q, \qquad p = F_\tau(z), \qquad R = \mathbb{P}(\mathrm{Bin}(q-1,p) = k-1), \qquad a = \frac{q-k}{q-1}, \quad b = \frac{k-1}{q-1}.$$

For every vector $v \in \mathbb{R}^q$ with $\sum_i v_i = 0$,

$$D^2\mathcal{S}_{q,k,\tau}(z,\dots,z)[v,v] = \frac{g_\tau(z)R}{z}\,[a\eta_\tau(z) - b\zeta_\tau(z)] \sum_{i=1}^q v_i^2. \tag{14}$$

Consequently:

(i) If $(\tau-1)(q-k) \leq k-1$, this barycenter is a strict facewise local maximum for every $M > 0$.

(ii) If $(\tau-1)(q-k) > k-1$, there is a unique $z_* > 0$ satisfying

$$(q-k)\eta_\tau(z_*) = (k-1)\zeta_\tau(z_*). \tag{15}$$

The barycenter is a strict facewise local minimum for $M < qz_*$ and a strict facewise local maximum for $M > qz_*$.

In particular, a candidate with $(q-k)\eta_\tau(M/q) > (k-1)\zeta_\tau(M/q)$ cannot maximize the full allocation problem.

Proof. Write

$$\beta = \mathbb{P}(\mathrm{Bin}(q-2,p) = k-1), \qquad \gamma = \mathbb{P}(\mathrm{Bin}(q-2,p) = k-2).$$

The pivotal derivative identity gives the diagonal entries of the Hessian as $g'_\tau(z)R$. Conditioning on a pair gives every off-diagonal entry as $g_\tau(z)^2(\gamma-\beta)$. Since $\sum_i v_i = 0$,

$$D^2\mathcal{S}_{q,k,\tau}[v,v] = [g'_\tau(z)R - g_\tau(z)^2(\gamma-\beta)]\sum_i v_i^2.$$

The elementary binomial ratios are

$$\frac{\beta}{R} = \frac{a}{1-p}, \qquad \frac{\gamma}{R} = \frac{b}{p}.$$

Using $g'_\tau/g_\tau = m/z - 1$ and $a+b=1$, the coefficient of $\sum_i v_i^2$ becomes

$$g_\tau(z)R\left[\frac{m}{z} - 1 + \frac{ag_\tau(z)}{1-p} - \frac{bg_\tau(z)}{p}\right] = \frac{g_\tau(z)R}{z}\big[a\eta_\tau(z) - b\zeta_\tau(z)\big],$$

which proves (14).

The barycenter is stationary by permutation symmetry. The function

$$(q-k)\eta_\tau(z) - (k-1)\zeta_\tau(z)$$

is strictly decreasing, has limit $m(q-k)-(k-1)$ at zero, and tends to $-\infty$ at infinity. Its sign gives the alternatives above. A definite constrained Hessian yields a strict local extremum on the positive face by the second-order Taylor expansion. A strict facewise local minimum cannot be a global maximizer on the simplex. □

At $M = qz_*$, the constrained Hessian vanishes and this test makes no local classification. Nor does facewise local maximality by itself compare different support sizes. The global comparisons among the $V_q(M)$ in Section 4 determine optimality, while the Hessian also describes suboptimal barycenters.

3.4 An obstruction to frozen-pair equalization

The full stationarity conditions in the active-uniform proof cannot be replaced by a rule that equalizes or collapses every unequal pair while holding its shadow fixed.

Example 3.6 (An unequal pair beats both equalization and collapse)**.** Let $F_2(x) = 1-(1+x)e^{-x}$, and put

$$\alpha = \frac{9999}{10000}, \qquad J(x) = F_2(x) + F_2(13-x) - \alpha F_2(x)F_2(13-x).$$

Then

$$J(3) - J(0) > \frac{12}{10^6}, \qquad J(3) - J(13/2) > \frac{9}{10^6}. \tag{16}$$

Thus replacing $(3,10)$ by either $(0,13)$, $(13,0)$, or $(13/2,13/2)$ decreases this frozen-pair objective. The shadow is realizable in an MDS problem: with $q=3, k=2$, let the third coordinate have recovery probability $10000/10001$. Its pair-shadow parameter is exactly the displayed α.

Proof. Put $E = e^{-1/2}$. Direct substitution gives

$$J(3) - J(0) = \frac{1 - 4E^6 - 11E^{20}}{10000} + (14 - 44\alpha)E^{26},$$
$$J(3) - J(13/2) = \frac{-4E^6 - 11E^{20} + 15E^{13}}{10000} + \frac{49\alpha}{4}E^{26}.$$

Here is a rational certificate of both signs. Set

$$L = \frac{6065306597126334}{10^{16}}, \qquad U = \frac{6065306597126335}{10^{16}}, \qquad S_j = \sum_{r=0}^{j} \frac{(-1/2)^r}{r!}.$$

The alternating-series bounds and rational comparisons give $L < S_{21} < E < S_{20} < U$. Since $14 - 44\alpha < 0$, substitution of the appropriate endpoints in each signed term yields

$$J(3) - J(0) > \frac{1 - 4U^6 - 11U^{20}}{10000} + (14 - 44\alpha)U^{26} > \frac{12}{10^6},$$
$$J(3) - J(13/2) > \frac{-4U^6 - 11U^{20} + 15L^{13}}{10000} + \frac{49\alpha}{4}L^{26} > \frac{9}{10^6}.$$

All quantities in the last comparisons are rational. Symmetry gives $J(13) = J(0)$.

For the asserted shadow, its two relevant masses are $\beta = 10000/10001$ and $\gamma = 1/10001$, so $1 - \gamma/\beta = \alpha$. Because F_2 is continuous and strictly increasing from zero to one, this third-coordinate probability is realized by a finite positive intensity. □

This example concerns a single conditional pair comparison. It does not require the fixed third coordinate to satisfy its own stationarity equation. The structured variations in Section 2 use those additional equations to exclude asymmetric local maxima of the full objective.

4 Global crossings and support phases

Throughout this section, $\tau > 1$ and $n \geq k \geq 2$ are fixed. For support-curve comparisons, we use the same formula for V_q at every integer $q \geq k$; feasible allocations still require $q \leq n$. Write $c(q) = c_{k,\tau}(q)$, with q_0 as in (8). We first classify every pair of support curves and prove strict concavity in logarithmic support. Together these yield Theorem 1.1. The support comparisons themselves do not use the active-uniform theorem; that theorem supplies the final allocation classification.

Theorem 4.1 (Complete pairwise crossing classification)**.** *Let $\tau > 1$ and let k, q, r be integers with $2 \leq k \leq q < r$. Exactly one of the following holds.*

(i) If $c_{k,\tau}(r) \geq c_{k,\tau}(q)$, then $V_r(M) > V_q(M)$ for every $M > 0$. This includes equality of the leading coefficients.

(ii) If $c_{k,\tau}(r) < c_{k,\tau}(q)$, there is a unique $\beta_{q,r} > 0$ at which $V_q = V_r$, with

$$V_q(M) > V_r(M) \quad (0 < M < \beta_{q,r}), \qquad V_q(M) < V_r(M) \quad (M > \beta_{q,r}).$$

The crossing is transverse: $V_r'(\beta_{q,r}) > V_q'(\beta_{q,r}) > 0$.

Theorem 4.2 (Strict concavity in logarithmic support)**.** *Let $\tau > 1$, $k \geq 2$ be an integer, and $M > 0$. There is a positive continuous interpolation $\widetilde{V}(q, M)$ for real $q \geq k$, agreeing with $V_q(M)$ at integers, such that*

$$s \longmapsto \log \widetilde{V}(e^s, M), \qquad s \geq \log k,$$

is strictly concave. In particular, for integers $k \leq q < r < t$,

$$V_r(M) > V_q(M)^{1-\theta} V_t(M)^{\theta}, \qquad \theta = \frac{\log r - \log q}{\log t - \log q}. \tag{17}$$

Among any finite set of consecutive feasible support sizes, at most two maximize at a given budget, and if two do, they are adjacent.

4.1 Cumulative-hazard geometry

Put

$$S(x) = 1 - F_\tau(x), \qquad h(x) = \frac{g_\tau(x)}{S(x)}, \qquad A(x) = -\log S(x) = \int_0^x h(t)\, dt.$$

The cumulative hazard A increases strictly from zero to infinity. Moreover $h = 1/P$, where P is the upper Gamma tail-to-density ratio from Section 2. Thus $xh'(x)/h(x) = \eta_\tau(x)$, and (5) proves that this elasticity is positive and strictly decreasing.

Lemma 4.3 (Cumulative-hazard geometry)**.** *For $\tau > 1$, the function $s \mapsto \log A(e^s)$ is strictly concave. Its inverse $s \mapsto \log A^{-1}(e^s)$ is strictly convex. If $T = A^{-1}$, then $e_T(y) = yT'(y)/T(y)$ is positive and strictly increasing.*

Proof. For $x > 0$,

$$\frac{A(x)}{xh(x)} = \int_0^1 \frac{h(xv)}{h(x)}\, dv. \tag{18}$$

For each $0 < v < 1$,

$$\frac{d}{d \log x} \log \frac{h(xv)}{h(x)} = \eta_\tau(xv) - \eta_\tau(x) > 0.$$

Hence the integral in (18) is strictly increasing. Its reciprocal

$$\alpha(x) = \frac{xh(x)}{A(x)} = \frac{d}{d \log x} \log A(x)$$

is positive and strictly decreasing, which proves strict concavity. The increasing inverse is strictly convex, and more explicitly $e_T(y) = 1/\alpha(T(y))$ is positive and strictly increasing. □

4.2 An exponential comparison and its Gamma transfer

Lemma 4.4 (Density-ratio crossing lemma)**.** *Let f, g be positive continuous probability densities on $(0, \infty)$. If $\log(g/f)$ is strictly concave, then*

$$D(x) = \int_0^x (g(t) - f(t))\, dt$$

has at most one zero in $(0, \infty)$. At such a zero, $D'(x) > 0$, and the sign changes from negative to positive.

Proof. The set $I = \{x : g(x) > f(x)\}$ is a nonempty interval and cannot be the entire half-line, since the densities have equal integral. Strict concavity excludes an interval on which $g/f = 1$. To the left of I, when nonempty, $D < 0$. To its right, when nonempty, total mass zero gives

$$D(x) = -\int_x^\infty (g(t) - f(t))\, dt > 0.$$

On I, $D' = g - f > 0$. Thus every zero lies in I, is unique, and has positive derivative. The same argument covers intervals I with zero or infinity as an endpoint. □

Lemma 4.5 (Exponential quantile-ratio monotonicity)**.** *For integers $2 \le k \le q < r$, let X_q, X_r be the k-th order statistics from q, r independent unit exponentials, with quantiles Q_q, Q_r. For $0 < u < 1$,*

$$Q_q(u) > Q_r(u), \qquad \frac{d}{du} \log \frac{Q_q(u)}{Q_r(u)} > 0.$$

Proof. Adding $r - q$ independent observations strictly increases the probability of at least k completions at every positive deadline, which gives the first inequality. The density for sample size q is

$$f_q(x) = k\binom{q}{k}(1 - e^{-x})^{k-1} e^{-(q-k+1)x}.$$

For $0 < c < 1$, set $R_c(x) = c f_r(cx)/f_q(x)$. Then

$$(\log R_c)'(x) = (k-1)\left(\frac{c}{e^{cx} - 1} - \frac{1}{e^x - 1}\right) + (q - k + 1) - c(r - k + 1),$$

and therefore

$$(\log R_c)''(x) = \frac{k-1}{x^2}[\Phi(x) - \Phi(cx)] < 0, \qquad \Phi(x) = \left(\frac{x}{2\sinh(x/2)}\right)^2.$$

Indeed, Φ is strictly decreasing: for $z > 0$, $\sinh z - z\cosh z < 0$, since its derivative is $-z\sinh z < 0$ and its limit at zero is zero.

Fix u, and put $a = Q_q(u)$, $b = Q_r(u)$, $c = b/a \in (0, 1)$. At $x = a$, the distribution functions corresponding to $c f_r(cx)$ and $f_q(x)$ agree. Lemma 4.4 gives $c f_r(b) > f_q(a)$, or $b f_r(b) > a f_q(a)$. Thus

$$\frac{d}{du} \log \frac{a}{b} = \frac{1}{a f_q(a)} - \frac{1}{b f_r(b)} > 0.$$

□

Lemma 4.6 (Transfer through a geometrically convex function)**.** *Suppose $a(u) > b(u) > 0$ are differentiable with positive derivatives and $(\log a - \log b)' > 0$. Let $T : (0, \infty) \to (0, \infty)$ be increasing and differentiable, with positive nondecreasing elasticity $e_T(y) = yT'(y)/T(y)$. Then*

$$\frac{d}{du} \log \frac{T(a(u))}{T(b(u))} > 0.$$

Proof. Write $A_1 = (\log a)'$, $B_1 = (\log b)'$, so $A_1 > B_1 > 0$. The derivative in question equals

$$e_T(a)A_1 - e_T(b)B_1 = e_T(a)(A_1 - B_1) + (e_T(a) - e_T(b))B_1 > 0.$$

□

Let $Y_{(k:q)}$ be the k-th order statistic of q unit-rate Gamma variables of shape τ, and let K_q denote the quantile function of $qY_{(k:q)}$, whose distribution function is V_q. These variables have positive continuous densities on $(0, \infty)$.

Proposition 4.7 (Strict scaled Gamma quantile order)**.** *For $\tau > 1$ and integers $2 \le k \le q < r$,*

$$\frac{d}{du} \log \frac{K_q(u)}{K_r(u)} > 0 \qquad (0 < u < 1). \tag{19}$$

Proof. The cumulative-hazard transform $A(Y)$ is unit exponential. Since A is increasing, $Y_{(k:q)}$ has the distribution of $T(X_q)$, where $T = A^{-1}$. Hence $K_q(u) = qT(Q_q(u))$. Apply Lemmas 4.3, 4.5, and 4.6; the factor q/r does not affect the logarithmic derivative. □

Proof of Theorem 4.1. For a fixed pair q, r, apply the endpoint estimates with a finite support cap at least r. Proposition 3.1 gives $V_q(M) \sim c(q)M^{\tau k}/\Gamma(\tau+1)^k$ at zero, so

$$\lim_{u \downarrow 0} \frac{K_q(u)}{K_r(u)} = \left(\frac{c(r)}{c(q)}\right)^{1/(\tau k)}. \tag{20}$$

At infinity, Proposition 3.4 yields

$$-\log(1 - V_q(M)) = \frac{q-k+1}{q} M + O(\log M), \qquad K_q(u) \sim \frac{q}{q-k+1} \log \frac{1}{1-u}.$$

Consequently

$$\lim_{u \uparrow 1} \frac{K_q(u)}{K_r(u)} = \frac{q(r-k+1)}{r(q-k+1)} > 1, \tag{21}$$

where the last inequality follows from $(r-q)(k-1) > 0$. All parameters in these asymptotics are fixed.

If $c(r) \ge c(q)$, the limit in (20) is at least one. Strict increase of the quantile ratio implies $K_q(u) > K_r(u)$ for every $0 < u < 1$, including when the limit equals one. Equivalently, $V_r(M) > V_q(M)$ for every $M > 0$.

If $c(r) < c(q)$, the two endpoint limits lie on opposite sides of one. The continuous strictly increasing ratio equals one at a unique u_*. Put $\beta_{q,r} = K_q(u_*) = K_r(u_*)$. Below u_*, one has $K_q < K_r$, and above it $K_q > K_r$. For a budget M, set $u = V_q(M)$, so $K_q(u) = M$. These quantile inequalities give the asserted signs of $V_r(M) - V_q(M)$. At equality,

$$0 < \frac{d}{du} \log \frac{K_q(u)}{K_r(u)}\bigg|_{u=u_*} = \frac{1}{\beta_{q,r} V_q'(\beta_{q,r})} - \frac{1}{\beta_{q,r} V_r'(\beta_{q,r})}.$$

Positivity of the densities proves transversality. □

4.3 Strict concavity in logarithmic support

Proof of Theorem 4.2. Let $E_0, \dots, E_{k-1}$ be independent unit exponentials. By memorylessness, the successive waiting times until the k-th of q exponential clocks rings are independent with rates $q, q-1, \dots, q-k+1$. Thus, for integer $q \ge k$,

$$X_q \stackrel{d}{=} \sum_{j=0}^{k-1} \frac{E_j}{q-j}.$$

The cumulative-hazard transform therefore gives

$$V_q(M) = \mathbb{P}\left(\sum_{j=0}^{k-1} \frac{E_j}{1-j/q} \le qA(M/q)\right). \tag{22}$$

Define $\widetilde{V}(q, M)$ for real $q \ge k$ by this same right-hand side. It is positive and continuous: the coefficients depend continuously on q, every weighted exponential sum has a continuous distribution, and dominated convergence applies.

Put $s = \log q$ and $U_j = \log E_j$. Their joint density is

$$\rho(u) = \exp\left(\sum_{j=0}^{k-1}(u_j - e^{u_j})\right), \qquad u \in \mathbb{R}^k, \tag{23}$$

which is positive and log-concave. Set

$$L(u, s) = \log\left(\sum_{j=0}^{k-1} \exp\{u_j - \log(1 - je^{-s})\}\right), \tag{24}$$

$$b_M(s) = s + \log A(Me^{-s}). \tag{25}$$

Then $\widetilde{V}(e^s, M) = \mu(K_s)$, where μ has density ρ and $K_s = \{u : L(u, s) \le b_M(s)\}$. The function L is jointly convex: log-sum-exp is convex and increasing in each argument, and

$$\frac{d^2}{ds^2}[-\log(1 - je^{-s})] = \frac{je^{-s}}{(1 - je^{-s})^2} \ge 0.$$

Lemma 4.3 makes b_M strictly concave.

Take $s_0 \ne s_1$, $0 < \theta < 1$, and $s_\theta = (1-\theta)s_0 + \theta s_1$. Put

$$\delta = b_M(s_\theta) - (1-\theta)b_M(s_0) - \theta b_M(s_1) > 0.$$

Joint convexity of L gives

$$(1-\theta)K_{s_0} + \theta K_{s_1} \subseteq K_\theta^- := \{u : L(u, s_\theta) \le b_M(s_\theta) - \delta\}.$$

The Prékopa–Leindler inequality for the log-concave measure μ [9, Theorem 2] yields

$$\mu(K_\theta^-) \ge \mu(K_{s_0})^{1-\theta}\mu(K_{s_1})^\theta.$$

The shell

$$\{u : b_M(s_\theta) - \delta < L(u, s_\theta) < b_M(s_\theta)\}$$

is nonempty and open, since $L(u + t(1, \dots, 1), s) = L(u, s) + t$. Its measure is positive because $\rho > 0$ everywhere. Hence

$$\widetilde{V}(e^{s_\theta}, M) > \widetilde{V}(e^{s_0}, M)^{1-\theta}\widetilde{V}(e^{s_1}, M)^\theta.$$

No equality classification for Prékopa–Leindler is needed. Restricting to integers proves (17). Two nonadjacent maximizers would force an intermediate integer support to have strictly larger value, which is impossible. □

Remark 4.8 (Beta form of the interpolation). The interpolation is $\widetilde{V}(q,M) = I_{F_\tau(M/q)}(k, q-k+1)$, where $I_p(a,b)$ is the distribution function of a Beta(a,b) variable at p. Indeed, for $Z = \sum_{j=0}^{k-1} E_j/(q-j)$,

$$\mathbb{E}e^{-tZ} = \prod_{j=0}^{k-1} \frac{q-j}{q-j+t} = \frac{\Gamma(q+1)\Gamma(q-k+1+t)}{\Gamma(q-k+1)\Gamma(q+1+t)}.$$

For nonnegative integer t, these are the moments of a Beta$(q-k+1,k)$ variable. Uniqueness of moments on $[0,1]$ identifies e^{-Z} with that law. Applying the symmetry of the beta distribution to $\mathbb{P}(Z \le A(M/q))$ gives the formula. This closed form is not required for the concavity proof.

4.4 Ordered transitions and the allocation phases

Lemma 4.9 (An adjacent tie is a global tie)**.** *Let $\tau > 1$, $n \ge k \ge 2$, and $k \le q < n$. If $V_q(M) = V_{q+1}(M)$ at $M > 0$, then*

$$V_r(M) < V_q(M) \qquad (r \in \{k, \dots, n\} \setminus \{q, q+1\}).$$

Proof. Let $f(s) = \log \widetilde{V}(e^s, M)$. Its secant slope between $\log q$ and $\log(q+1)$ is zero. Strict concavity makes every secant continuing to the right have negative slope, and every secant ending at $\log q$ from the left have positive slope. Thus every value outside the adjacent pair is strictly smaller. □

Proof of Theorem 1.1. The leading coefficients have a strictly concave logarithmic interpolation: using $\binom{x}{k} = k!^{-1} \prod_{j=0}^{k-1}(x-j)$,

$$\frac{d^2}{ds^2} \log c(e^s) = -\sum_{j=1}^{k-1} \frac{je^s}{(e^s-j)^2} < 0 \qquad (s \ge \log k).$$

Their integer values are strictly unimodal, apart from a possible tie at two adjacent maximizing integers. Since q_0 is the larger maximizer, $c(q) > c(q+1)$ for $q_0 \le q < n$. Theorem 4.1 supplies each unique adjacent crossing $\beta_q = \beta_{q,q+1} > 0$.

At $M = \beta_q$, Lemma 4.9 gives $V_{q+2}(M) < V_{q+1}(M)$ whenever $q+2 \le n$. The crossing of $q+1, q+2$ must occur later, so $\beta_q < \beta_{q+1}$. For $k \le r < q_0$, one has $c(q_0) \ge c(r)$, and Theorem 4.1 gives $V_{q_0}(M) > V_r(M)$ for every positive budget. Such supports are permanently dominated.

If $\beta_{q-1} < M < \beta_q$, each adjacent crossing below q has occurred and each crossing at or above q has yet to occur. Chaining their strict comparisons makes q the unique maximizing support. At $M = \beta_q$, Lemma 4.9 gives exactly the two maximizing supports $q, q+1$. If $q_0 = n$, all smaller supports are permanently dominated and there are no crossings to list. Finally, Theorem 2.1 gives exactly the stated maximizing allocations, including all equality cases at a boundary. □

Corollary 4.10 (Nonadjacent crossing locations)**.** *Let $\tau > 1$, $n \ge k \ge 2$, and $q_0 \le q < r-1 \le n-1$. Then*

$$\beta_q < \beta_{q,r} < \beta_{r-1}.$$

At $V_q(\beta_{q,r}) = V_r(\beta_{q,r})$, every integer support strictly between q and r has larger value. This equality is therefore not a global tie.

Proof. At β_q, support r is below q; at β_{r-1}, support q is below r. The unique crossing lies between these budgets. Theorem 4.2 gives the last assertion. □

For $k = 10, \tau = 2, n \geq 12$, Example 3.3 and the staircase theorem give the sequence $12, 13, \ldots, n$, with every listed support optimal on a nonempty open interval. Supports $10, 11$ never become optimal. This illustrates why absence of skipped transitions refers to the phases beginning at q_0, rather than asserting that $q_0 = k$. For $n = 16$, the four transition budgets are approximately

Transition	Budget
$12 \longrightarrow 13$	3.751806808726
$13 \longrightarrow 14$	8.851172168188
$14 \longrightarrow 15$	14.333018261348
$15 \longrightarrow 16$	20.197931039196

These diagnostic values were computed with 80-decimal working precision; the displayed decimals are approximations. The repository support-phase diagnostic records finite numerical checks. Existence, uniqueness, and ordering of the transitions follow from the analytic proof above. The cases $k = 1$, $\tau \leq 1$, and $k = n$ retain the classifications in Theorem 2.1.

Remark 4.11 (Support comparisons for other common responses). Let F be a continuously differentiable distribution function on $(0, \infty)$, with positive density, $F(0) = 0$, and $\lim_{x\to\infty} F(x) = 1$. Suppose that its cumulative hazard $A = -\log(1 - F)$ satisfies strict concavity of $s \mapsto \log A(e^s)$. The spacing and Prékopa argument then gives strict concavity in logarithmic support for the curves $\mathbb{P}(\mathrm{Bin}(q, F(M/q)) \geq k)$. For $k \geq 2$, the inverse-hazard comparison also gives strict quantile-ratio ordering. Any pair thus has at most one crossing, with the same transverse orientation in favor of the larger support. The Gamma endpoint asymptotics above are what decide crossing existence and identify q_0 and the terminal support in the present theorem. This remark concerns active-uniform candidates only: a global allocation theorem for a different response requires a separate argument, such as the criterion in Section 5.

5 A sufficient criterion for a common recovery response

The active-uniform proof uses a specific set of properties of the common recovery curve. We now state these properties as a sufficient criterion. The cardinality recovery rule and the unweighted budget are retained.

Proposition 5.1 (A common-response tail-ratio criterion). *Let F be an absolutely continuous distribution function on $[0, \infty)$, with $F(0) = 0$ and $\lim_{x\to\infty} F(x) = 1$. Suppose its density g is positive and C^2 on $(0, \infty)$, and satisfies*

$$g(0+) = 0, \qquad \lim_{x\to\infty} g(x) = 0, \qquad (\log g)''(x) < 0 \quad (x > 0). \tag{26}$$

Define

$$P(x) = \frac{1 - F(x)}{g(x)}, \qquad Q(x) = \frac{F(x)}{g(x)}.$$

Assume that on $(0, \infty)$,

$$L_1(x) = \frac{-P'(x)}{Q'(x)} \quad \textit{is strictly decreasing,} \qquad L_2(x) = \frac{(-P'(x))Q(x)^2}{Q'(x)P(x)^2} \quad \textit{is strictly increasing.} \tag{27}$$

For independent $B_i \sim \mathrm{Bernoulli}(F(\lambda_i))$*, every global maximizer of*

$$\mathcal{S}_F(\lambda) = \mathbb{P}\left(\sum_{i=1}^{n} B_i \geq k\right), \qquad \lambda_i \geq 0, \quad \sum_i \lambda_i = M,$$

is uniform on its positive support, for every $n \geq 1$*,* $1 \leq k \leq n$*, and* $M > 0$*. In particular,*

$$\max_{\lambda_i \geq 0,\ \sum_i \lambda_i = M} \mathcal{S}_F(\lambda) = \max_{q=k,\ldots,n} \mathbb{P}(\mathrm{Bin}(q, F(M/q)) \geq k). \tag{28}$$

The full set of maximizers consists of the permutations of the active-uniform allocations whose support sizes attain this finite maximum. If $k = 1$*, their support size is one; if* $k = n$*, the unique maximizer is full uniformity.*

Proof. We establish the properties of the response that are needed by the pair-shadow argument, and then exclude the possible asymmetric stationary patterns.

Tail ratios and endpoint faces. Put $u = (\log g)'$, so $u' < 0$. Differentiation gives

$$P' = -1 - uP, \qquad Q' = 1 - uQ. \tag{29}$$

In fact $P' < 0$ and $Q' > 0$. To see the first assertion, it is immediate from (29) when $u(x) \geq 0$. If $u(x) < 0$, strict log-concavity gives

$$\frac{g(x+s)}{g(x)} < e^{u(x)s} \quad (s > 0), \qquad P(x) = \int_0^\infty \frac{g(x+s)}{g(x)}\, ds < \frac{-1}{u(x)}.$$

Again $P'(x) < 0$. For Q', the case $u(x) \leq 0$ is immediate. If $u(x) > 0$, the corresponding tangent-line bound to the left gives

$$Q(x) = \int_0^x \frac{g(s)}{g(x)}\, ds < \int_0^x e^{u(x)(s-x)}\, ds < \frac{1}{u(x)},$$

so $Q'(x) > 0$. Thus the quotients in (27) are well defined.

The hazard $g/(1-F) = 1/P$ is strictly increasing, while $(\log F)' = g/F = 1/Q$ is strictly decreasing. The latter property and strict Jensen show that on a positive q-face the all-must-succeed objective $\prod_i F(\lambda_i)$ is uniquely maximized at its barycenter. For $k = 1$, let $A(x) = -\log(1 - F(x))$. Since $A(0) = 0$ and A' is strictly increasing,

$$A(x+y) > A(x) + A(y) \quad (x, y > 0).$$

Merging any two positive coordinates therefore strictly increases the at-least-one-success probability. This proves the stated $k = 1$ conclusion.

Marginal-ratio shape. For $\alpha < 1$, define

$$D_\alpha = 1 - \alpha F > 0, \qquad L_\alpha = \frac{g}{D_\alpha}, \qquad N_\alpha = uD_\alpha + \alpha g.$$

Then

$$L'_\alpha = \frac{g}{D_\alpha^2} N_\alpha, \qquad N'_\alpha = u' D_\alpha < 0.$$

The density assumptions imply $L_\alpha(0+) = L_\alpha(\infty) = 0$. Because L_α is positive in the interior, it has an interior maximum. The strict decrease of N_α makes this maximum unique; denote its location by $x_*(\alpha)$. At the critical point,

$$\partial_\alpha N_\alpha = g - uF = \frac{g}{D_\alpha} > 0.$$

The implicit-function theorem consequently shows that $x_*(\alpha)$ is strictly increasing in α.

Stationary-pattern exclusion. The differentiation and conditioning in Lemmas 2.7 and 2.8 use only $F' = g > 0$, $0 < F < 1$, and independence. They therefore hold for the present response without modification. Lemma 2.10 concerns only Bernoulli convolutions. Together with the marginal-ratio shape just proved, these statements supply precisely the hypotheses used in Propositions 2.11 and 2.12: three distinct active levels are impossible, and splitting two equal small coordinates gives positive second variation. Thus a nonuniform interior local maximum could only have the form $(x, y, \dots, y)$, where $0 < x < y$.

The binomial coefficient calculation in Lemma 2.13 also uses no Gamma identity. Writing $a = (q-k)/(q-1)$, $b = (k-1)/(q-1)$, it gives

$$\frac{D_y}{D_x} = 1 + K(x,y), \qquad K(x,y) = a\frac{P(x)}{P(y)} + b\frac{Q(x)}{Q(y)} - 1.$$

At stationarity, put $S = (P(x) - P(y))/(Q(y) - Q(x))$. Then $b/a = SQ(y)/P(y)$. Cauchy's mean-value theorem and the two slope assumptions give

$$S < L_1(x), \qquad S\frac{Q(x)Q(y)}{P(x)P(y)} < L_2(y).$$

Consequently

$$K_x = \frac{aQ'(x)}{P(y)}[S - L_1(x)] < 0, \qquad K_y = a\frac{P(x)(-P'(y))}{P(y)^2} - b\frac{Q(x)Q'(y)}{Q(y)^2} > 0.$$

Along $y = (M-x)/(q-1)$, the objective G satisfies $G' = -D_x K$. At a stationary point,

$$G'' = -D_x\left(K_x - \frac{K_y}{q-1}\right) > 0.$$

This is the same feasible-direction exclusion as in Proposition 2.14; it rules out the final asymmetric pattern.

Finally, compactness gives a global maximizer. Its positive support has size $q \geq k$, because smaller supports have zero success probability while a k-point uniform allocation has positive success probability. It is an interior local maximum on its positive face. The endpoint cases and the preceding exclusions force that face's barycenter. Evaluation on each possible support gives (28) and the complete equality description. □

The Gamma responses F_τ, $\tau > 1$, satisfy all these hypotheses by the tail-ratio inequalities proved in Section 2. Strict log-concavity of the density supplies the marginal-ratio and level reductions; the two slope conditions in (27) supply the final exclusion. The criterion applies to a common response under an unweighted total budget. It makes no optimizer claim for unequal response curves, unequal allocation costs, or a general upward-closed recovery family.

6 Perturbation bounds and finite support comparisons

A coordinatewise uniform coupling gives a two-sided perturbation estimate. It controls the effect of an approximate allocation or an approximate recovery curve without a new optimization argument.

Proposition 6.1 (Perturbation bound)**.** *Let* $B_i \sim \mathrm{Bernoulli}(a_i)$ *and* $\widetilde{B}_i \sim \mathrm{Bernoulli}(\widetilde{a}_i)$ *be independent within each family, with* $a_i, \widetilde{a}_i \in [0,1]$*. For every event* $\mathcal{A} \subseteq \{0,1\}^n$*,*

$$\left|\mathbb{P}(B \in \mathcal{A}) - \mathbb{P}(\widetilde{B} \in \mathcal{A})\right| \le \min\left\{1, \sum_{i=1}^{n} |a_i - \widetilde{a}_i|\right\}.$$

Consequently, for any common recovery curve F*,*

$$|\mathcal{S}_F(\lambda) - \mathcal{S}_F(\widetilde{\lambda})| \le \sum_i |F(\lambda_i) - F(\widetilde{\lambda}_i)|, \qquad \mathcal{S}_F(\lambda) := \mathbb{P}\left(\sum_i \mathrm{Bernoulli}(F(\lambda_i)) \ge k\right),$$

where the Bernoulli variables are independent. If F *is* L*-Lipschitz, the right-hand side is at most* $L\|\lambda - \widetilde{\lambda}\|_1$*. For the Gamma curve* F_τ *with* $\tau \ge 1$*, one can take*

$$L_\tau = \begin{cases} 1, & \tau = 1, \\ (\tau-1)^{\tau-1} e^{-(\tau-1)}/\Gamma(\tau), & \tau > 1. \end{cases}$$

Proof. Take independent uniform U_i and set $B_i = \mathbf{1}_{\{U_i \le a_i\}}$, $\widetilde{B}_i = \mathbf{1}_{\{U_i \le \widetilde{a}_i\}}$. The two event indicators agree unless at least one coordinate differs. The union bound gives the first inequality, and the remaining estimates follow by substitution and the Lipschitz property. For $\tau > 1$, differentiating the Gamma density shows that its maximum occurs at $\tau - 1$; at $\tau = 1$, the density is e^{-x}. □

Remark 6.2 (A certified finite comparison). Suppose one of the active-uniform candidates has value exceeding every other candidate by a positive gap δ. If every candidate allocation is perturbed by at most ε in ℓ^1, the winning candidate remains strictly best among these perturbed candidates whenever $2L_\tau\varepsilon < \delta$. The same conclusion holds for any candidate values known within a common error bound smaller than $\delta/2$. This is a gap between finitely many support comparisons; continuity prevents a positive objective gap from all nonuniform allocations arbitrarily close to a maximizer. For $0 < \tau < 1$, the density is unbounded at zero, but the CDF perturbation bound in Proposition 6.1 still applies.

7 Further questions and verification scope

Theorems 4.1, 4.2, and 1.1 settle the global support comparisons: crossings are unique whenever they exist, support re-entry is impossible, and the optimizing support visits every integer from q_0 through n. Quantitative questions remain, including estimates for the transition budgets β_q, their dependence on k, τ, and asymptotic descriptions when the parameters grow. The fixed-parameter endpoint expansions in Section 3 do not assert uniformity in such regimes.

Further directions are to identify common responses satisfying Proposition 5.1, sharpen the mixed Gamma bound, and determine what replaces equal-positive-coordinate extremizers for heterogeneous responses or unequal costs. For integer thresholds $t \ge 2$, exact fixed-read

optimization concerns coupled multinomial counts; a Poissonization sandwich transfers probability bounds, not coefficientwise optimizer classifications. Likewise, optimizing at each deadline does not justify interchanging optimization with integration to minimize expected completion time.

The repository includes exact finite checks, symbolic identity checks, and explicitly labeled numerical diagnostics. They supplement the analytic proofs above and do not certify their infinite parameter ranges. The global support proof uses decreasing Gamma hazard elasticity, an elementary exponential quantile comparison, and the classical Prékopa–Leindler inequality; its allocation conclusion additionally uses the active-uniform theorem. It does not require unimodality of a ratio of Gamma completion densities, the local barycenter Hessian, or the expected-terminal-cloud staircase. No proof-assistant formalization is claimed. Stable statement labels and a dependency ledger are supplied to support subsequent independent auditing and formalization.

References

[1] E. Abbe, S.-L. Huang, and E. Telatar, Proof of the outage probability conjecture for MISO channels, *IEEE Trans. Inf. Theory* 59 (2013), 2596–2602; arXiv:1103.5478.

[2] H. Abraham, I. Feldman, and E. Yaakobi, Analyzing Collection Strategies: A Computational Perspective on the Coupon Collector Problem, arXiv:2601.12351 (2026).

[3] E. Anceaume, Y. Busnel, E. Schulte-Geers, and B. Sericola, Optimization results for a generalized coupon collector problem, *J. Appl. Probab.* 53 (2016), 622–629; arXiv:1504.03878.

[4] D. Bar-Lev, O. Sabary, R. Gabrys, and E. Yaakobi, "Cover your bases: How to minimize the sequencing coverage in DNA storage systems," *IEEE Transactions on Information Theory*, vol. 71, no. 1, pp. 192–218, 2025. doi:10.1109/TIT.2024.3496587; arXiv:2305.05656.

[5] Á. Baricz, Mills' ratio: reciprocal convexity and functional inequalities, *Acta Universitatis Sapientiae Mathematica* 4 (2012), 26–35; arXiv:1010.3267.

[6] P. Flajolet, D. Gardy, and L. Thimonier, Birthday paradox, coupon collectors, caching algorithms and self-organizing search, *Discrete Applied Mathematics* 39 (1992), 207–229.

[7] T. M. Liggett, "Ultra logconcave sequences and negative dependence," *Journal of Combinatorial Theory, Series A*, vol. 79, no. 2, pp. 315–325, 1997.

[8] C. D. Long, Nonuniform Coverage and the Terminal-Cloud Staircase, companion manuscript (2026), repository source.

[9] A. Prékopa, On logarithmic concave measures and functions, *Acta Scientiarum Mathematicarum (Szeged)* 34 (1973), 335–343. Original paper. Theorem 2 gives log-concavity of the generated measure; Theorem 6 gives preservation under marginalization.

[10] J. Segura, Monotonicity properties and bounds for the chi-square and gamma distributions, arXiv:1406.6106 (2014).

[11] L. Sun and X. Zhang, Stochastic Comparisons of Order Statistics from Gamma Distributions, *Chinese Journal of Applied Probability and Statistics* 20 (2004), 404–408.