

# Agoh–Giuga conjecture

In [number theory](#) the **Agoh–Giuga conjecture** on the Bernoulli numbers  $B_k$  postulates that  $p$  is a [prime number](#) if and only if

$$pB_{p-1} \equiv -1 \pmod{p}.$$

It is named after [Takashi Agoh](#) and [Giuseppe Giuga](#).

## Equivalent formulation

---

The conjecture as stated above is due to [Takashi Agoh](#) (1990); an equivalent formulation is due to [Giuseppe Giuga](#), from 1950, to the effect that  $p$  is prime if and only if

$$1^{p-1} + 2^{p-1} + \cdots + (p-1)^{p-1} \equiv -1 \pmod{p}$$

which may also be written as

$$\sum_{i=1}^{p-1} i^{p-1} \equiv -1 \pmod{p}.$$

It is trivial to show that  $p$  being prime is sufficient for the second equivalence to hold, since if  $p$  is prime, [Fermat's little theorem](#) states that

$$a^{p-1} \equiv 1 \pmod{p}$$

for  $a = 1, 2, \dots, p-1$ , and the equivalence follows, since  $p-1 \equiv -1 \pmod{p}$ .

## Status

---

The statement is still a conjecture since it has not yet been proven that if a number  $n$  is not prime (that is,  $n$  is [composite](#)), then the formula does not hold. It has been shown that a composite number  $n$  satisfies the formula if and only if it is both a [Carmichael number](#) and a [Giuga number](#), and that if such a number exists, it has at least 13,800 digits (Borwein, Borwein, Borwein, Girgensohn 1996). Laerte Sorini, finally, in a work of 2001 showed that a possible counterexample should be a number  $n$  greater than  $10^{36067}$  which represents the limit suggested by Bedocchi for the demonstration technique specified by Giuga to his own conjecture.

## Relation to Wilson's theorem

---

The Agoh–Giuga conjecture bears a similarity to [Wilson's theorem](#), which has been proven to be true. Wilson's theorem states that a number  $p$  is prime if and only if

$$(p-1)! \equiv -1 \pmod{p},$$

which may also be written as

$$\prod_{i=1}^{p-1} i \equiv -1 \pmod{p}.$$

For an odd prime  $p$  we have

$$\prod_{i=1}^{p-1} i^{p-1} \equiv (-1)^{p-1} \equiv 1 \pmod{p},$$

and for  $p=2$  we have

$$\prod_{i=1}^{p-1} i^{p-1} \equiv (-1)^{p-1} \equiv 1 \pmod{p}.$$

So, the truth of the Agoh–Giuga conjecture combined with Wilson's theorem would give: a number  $p$  is prime if and only if

$$\sum_{i=1}^{p-1} i^{p-1} \equiv -1 \pmod{p}$$

and

$$\prod_{i=1}^{p-1} i^{p-1} \equiv 1 \pmod{p}.$$

## See also

- Bernoulli number § Arithmetical properties of the Bernoulli numbers

## References

- Giuga, Giuseppe (1951). "Su una presumibile proprietà caratteristica dei numeri primi". *Ist.Lombardo Sci. Lett., Rend., Cl. Sci. Mat. Natur.* (in Italian). **83**: 511–518. ISSN 0375-9164 (<https://search.worldcat.org/issn/0375-9164>) . Zbl 0045.01801 (<https://zbmath.org/?format=complete&q=an:0045.01801>) .
- Agoh, Takashi (1995). "On Giuga's conjecture". *Manuscripta Mathematica*. **87** (4): 501–510. doi:10.1007/bf02570490 (<https://doi.org/10.1007%2Fbf02570490>) . Zbl 0845.11004 (<https://zbmath.org/?format=complete&q=an:0845.11004>) .
- Borwein, D.; Borwein, J. M.; Borwein, P. B.; Girgensohn, R. (1996). "Giuga's Conjecture on Primality" (<https://web.archive.org/web/20050531164907/http://www.math.uwo.ca/~dborwein/cv/giuga.pdf>) (PDF). *American Mathematical Monthly*. **103** (1): 40–50. CiteSeerX 10.1.1.586.1424 (<https://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.586.14>

24) . doi:10.2307/2975213 (<https://doi.org/10.2307%2F2975213>) . JSTOR 2975213 (<https://www.jstor.org/stable/2975213>) . Zbl 0860.11003 (<https://zbmath.org/?format=complete&q=a+n:0860.11003>) . Archived from [the original](http://www.math.uwo.ca/~dborwein/cv/giuga.pdf) (<http://www.math.uwo.ca/~dborwein/cv/giuga.pdf>) (PDF) on 2005-05-31. Retrieved 2005-05-29.

- Sorini, Laerte (2001). "Un Metodo Euristico per la Soluzione della Congettura di Giuga". *Quaderni di Economia, Matematica e Statistica, DESP, Università di Urbino Carlo Bo* (in Italian). **68**. ISSN 1720-9668 (<https://search.worldcat.org/issn/1720-9668>) .