Random Number Generator

Long He

Design

Method I have tried:

www.random.org
Random.org is a website that produces “true” random numbers based on atmospheric noise.

Collect the noise in operating system: mouse movement and click, keyboard input, microphone.



Design

Why don't I choose these method?

www.random.org

I have to pay for true random number.



Noise in operating system

Low speed and easy to be exhausted.

Comparative low precision of CPU timer based on Java.

Mersenne Twister

The state succession algorithm of the Mersenne Twister can be depicted in an illustration as a kind of linear-feedback shift register.

http://www.quadibloc.com/crypto/co4814.htm

http://www.cs.gmu.edu/~sean/research/



Why I choose Mersenne Twister:

One of the best pseudorandom number generator

High speed (more details later)

Low bias

Long period period: 2^19937–1(Mersenne Prime)

Design Implement

Multithreading

Progress Bar

Java.math.BigInteger

System.nanoTime()



Multithreading
Show the number of “0”, number of “1”, data bias in the process dynamically.

Progress Bar
Show the progress rate.

Java.math.BigInteger
For extreme long input.

System.nanoTime()
Extreme high precision for seed generating.



Design Implement Usage



Trial	bias (%)	run time (s)
10	20	0.001
100	8	0.002
1k	3.2	0.002
10k	0.32	0.009
100k	0.4	0.056
1k^2	0.11	0.585
10k^2	0.018	5.459
100k^2	0.0085	51.366
1k^3	0.0022	455.416





My Future Idea:
Utilize laser or optical mouse to generate random binary numbers.

Advantages:
Real random number
High speed

IOP Publishing Laser Physics Letters
Laser Phys. Lett. **10** (2013) 045001 (5pp) doi:10.1088/1612-2011/10/4/045001

LETTER

Implementation of 1.6 Tb s^{-1} truly random number generation based on a super-luminescent emitting diode

Y Liu[1], M-Y Zhu[1], B Luo[1], J-W Zhang[2] and H Guo[1]

[1] State Key Laboratory of Advanced Optical Communication Systems and Networks, School of Electronics Engineering and Computer Science, Peking University, Beijing 100871, People's Republic of China
[2] School of Physics, Peking University, Beijing 100871, People's Republic of China

E-mail: hongguo@pku.edu.cn

Received 31 May 2012, in final form 27 September 2012
Accepted for publication 5 October 2012
Published 14 February 2013
Online at stacks.iop.org/LPL/10/045001

Paper on *Laser Physics Letters*:
http://iopscience.iop.org/1612-202X/10/4/045001/pdf/lpl13_4_045001.pdf

Thank you.