

SEM Growth Models

Elizabeth Hawkey

11/5/2017

```
## This is lavaan 0.5-23.1097
## lavaan is BETA software! Please report any bugs.
## Loading required package: Matrix
## -- Attaching packages ----- tidyverse 1.2.0 --
## √ ggplot2 2.2.1      √ purrr  0.2.3
## √ tibble  1.3.4      √ dplyr  0.7.4
## √ tidyr   0.7.2      √ stringr 1.2.0
## √ readr   1.1.1      √ forcats 0.2.0
## -- Conflicts ----- tidyverse_conflicts() --
## x tidyr::expand() masks Matrix::expand()
## x dplyr::filter() masks stats::filter()
## x dplyr::lag()     masks stats::lag()
##
## Attaching package: 'psych'
## The following objects are masked from 'package:ggplot2':
##
##   %+%, alpha
## The following object is masked from 'package:lavaan':
##
##   cor2cov
## Loading required package: arm
## Loading required package: MASS
##
## Attaching package: 'MASS'
## The following object is masked from 'package:dplyr':
##
##   select
##
## arm (Version 1.9-3, built: 2016-11-21)
## Working directory is /Users/elizabethhawkey/ejhawkey
##
## Attaching package: 'arm'
## The following objects are masked from 'package:psych':
##
##   logit, rescale, sim
##
## Attaching package: 'merTools'
```

```
## The following object is masked from 'package:psych':
##
##      ICC
##
## #####
## This is semTools 0.4-14
## All users of R (or SEM) are invited to submit functions or ideas for functions.
## #####
##
## Attaching package: 'semTools'
## The following object is masked from 'package:psych':
##
##      skew
```

1a. Start with a Univariate Growth Model

```
# Global Executive Composite raw scores
# with intercept only
Intercept.only= ' i=~ 1*T3gecrs_combined_conv + 1*T12gecrs_conv + 1*T14gecrs_conv'
Intercept.only.fit= growth(Intercept.only, data = growth_stats, missing= "ML")
```

```
## Warning in lav_data_full(data = data, group = group, cluster = cluster, : lavaan WARNING: some cases
## 6 19 21 28 29 34 36 49 52 64 72 81 83 84 86 91 94 103 113 115 141 169 180 187 194 198 199 205 217 :
```

```
summary (Intercept.only.fit)
```

```
## lavaan (0.5-23.1097) converged normally after 59 iterations
```

```
##
##
##              Used      Total
## Number of observations      302      348
##
## Number of missing patterns      7
##
## Estimator      ML
## Minimum Function Test Statistic      4.592
## Degrees of freedom      4
## P-value (Chi-square)      0.332
##
```

```
## Parameter Estimates:
```

```
##
##      Information      Observed
##      Standard Errors      Standard
##
```

```
## Latent Variables:
```


```
##      Estimate Std.Err z-value P(>|z|)
##      i =~
##      T3gecrs_cmbnd_c      1.000
##      T12gecrs_conv      1.000
##      T14gecrs_conv      1.000
##
```

```
## Intercepts:
```

```
##      Estimate Std.Err z-value P(>|z|)
```

```
##      .T3gecrs_cmbnd_c      0.000
##      .T12gecrs_conv       0.000
##      .T14gecrs_conv       0.000
##      i                     1.048      0.015      69.868      0.000
##
## Variances:
##              Estimate Std.Err z-value P(>|z|)
##      .T3gecrs_cmbnd_c   0.056   0.007   7.891   0.000
##      .T12gecrs_conv     0.018   0.004   4.589   0.000
##      .T14gecrs_conv     0.015   0.004   3.709   0.000
##      i                   0.044   0.005   8.056   0.000
```

```
semPaths(Intercept.only.fit, what = "paths", whatLabels= "est", layout = "tree")
```



```
# with a fixed slope
fixed.slope= ' i =~ 1*T3gecrs_combined_conv + 1*T12gecrs_conv + 1*T14gecrs_conv
s =~ 0*T3gecrs_combined_conv + 1*T12gecrs_conv + 2*T14gecrs_conv
s =~ 0*s' #fixes slope
fixed.slope.fit= growth(fixed.slope, data = growth_stats, missing= "ML")
```

```
## Warning in lav_data_full(data = data, group = group, cluster = cluster, : lavaan WARNING: some cases
##      6 19 21 28 29 34 36 49 52 64 72 81 83 84 86 91 94 103 113 115 141 169 180 187 194 198 199 205 217
##
## Warning in lav_object_post_check(object): lavaan WARNING: covariance matrix of latent variables
##      is not positive definite;
##      use inspect(fit,"cov.lv") to investigate.
```

```
inspect(fixed.slope.fit, "cov.lv")
```

```
##      i      s
## i 0.034
## s 0.005 0.000
```

```
summary (fixed.slope.fit)
```


```
## lavaan (0.5-23.1097) converged normally after 44 iterations
```

```

##
##
##      Number of observations      Used      Total
##      Number of missing patterns      7
##
##      Estimator      ML
##      Minimum Function Test Statistic      1.473
##      Degrees of freedom      2
##      P-value (Chi-square)      0.479
##
## Parameter Estimates:
##
##      Information      Observed
##      Standard Errors      Standard
##
## Latent Variables:
##      Estimate Std.Err z-value P(>|z|)
##      i =~
##      T3gcrs_cmbnd_c      1.000
##      T12gecrs_conv      1.000
##      T14gecrs_conv      1.000
##      s =~
##      T3gcrs_cmbnd_c      0.000
##      T12gecrs_conv      1.000
##      T14gecrs_conv      2.000
##
## Covariances:
##      Estimate Std.Err z-value P(>|z|)
##      i ~~
##      s      0.005      0.003      1.476      0.140
##
## Intercepts:
##      Estimate Std.Err z-value P(>|z|)
##      .T3gcrs_cmbnd_c      0.000
##      .T12gecrs_conv      0.000
##      .T14gecrs_conv      0.000
##      i      1.057      0.017      61.047      0.000
##      s      -0.011      0.012      -0.941      0.347
##
## Variances:
##      Estimate Std.Err z-value P(>|z|)
##      s      0.000
##      .T3gcrs_cmbnd_c      0.061      0.008      7.460      0.000
##      .T12gecrs_conv      0.020      0.004      4.632      0.000
##      .T14gecrs_conv      0.011      0.004      2.531      0.011
##      i      0.034      0.008      4.053      0.000

```

```
semPaths(fixed.slope.fit, what = "paths", whatLabels= "est", layout = "tree")
```



with a random slope

```
random.intercept= ' i=~ 1*T3gecrs_combined_conv + 1*T12gecrs_conv + 1*T14gecrs_conv
s=~ -1*T3gecrs_combined_conv + 0*T12gecrs_conv + 1*T14gecrs_conv'
random.intercept.fit= growth(random.intercept, data = growth_stats, missing= "ML")
```

```
## Warning in lav_data_full(data = data, group = group, cluster = cluster, : lavaan WARNING: some cases
## 6 19 21 28 29 34 36 49 52 64 72 81 83 84 86 91 94 103 113 115 141 169 180 187 194 198 199 205 217 2
```

```
summary (random.intercept.fit)
```

```
## lavaan (0.5-23.1097) converged normally after 67 iterations
```

```
##
```

```
##                               Used      Total
## Number of observations          302        348
```

```
##
```

```
## Number of missing patterns           7
```

```
##
```

```
## Estimator                        ML
```

```
## Minimum Function Test Statistic    0.477
```

```
## Degrees of freedom                 1
```

```
## P-value (Chi-square)               0.490
```

```
##
```

```
## Parameter Estimates:
```

```
##
```

```
## Information                      Observed
```

```
## Standard Errors                  Standard
```

```
##
```

```
## Latent Variables:
```

```
##           Estimate Std.Err z-value P(>|z|)
```


```
## i =~
```

```
## T3gecrs_cmbnd_c    1.000
```

```
## T12gecrs_conv      1.000
```

```
##      T14gecrs_conv      1.000
##      s =~
##      T3gecrs_cmbnd_c    -1.000
##      T12gecrs_conv      0.000
##      T14gecrs_conv      1.000
##
## Covariances:
##              Estimate Std.Err z-value P(>|z|)
##      i =~
##      s              0.005   0.003   1.471   0.141
##
## Intercepts:
##              Estimate Std.Err z-value P(>|z|)
##      .T3gecrs_cmbnd_c  0.000
##      .T12gecrs_conv    0.000
##      .T14gecrs_conv    0.000
##      i                 1.045   0.015  68.656   0.000
##      s                -0.013   0.012  -1.084   0.278
##
## Variances:
##              Estimate Std.Err z-value P(>|z|)
##      .T3gecrs_cmbnd_c  0.053   0.011   4.791   0.000
##      .T12gecrs_conv    0.021   0.005   4.688   0.000
##      .T14gecrs_conv    0.003   0.010   0.264   0.792
##      i                 0.044   0.006   7.739   0.000
##      s                 0.006   0.006   1.000   0.317
```

```
semPaths(random.intercept.fit, what = "paths", whatLabels= "est", layout = "tree")
```



```
# compare models
anova(Intercept.only.fit, fixed.slope)
```

```
## Chi Square Test Statistic (unscaled)
```

```
##
##           Df      AIC      BIC  Chisq Chisq diff Df diff Pr(>Chisq)
## Saturated  0              0.0000
## Model      4 39.036 57.589 4.5922      4.5922      4      0.3318
```

```
anova(fixed.slope.fit, random.intercept.fit)
```

```
## Chi Square Difference Test
```

```
##
##           Df      AIC      BIC  Chisq Chisq diff Df diff Pr(>Chisq)
## random.intercept.fit  1 40.921 70.604 0.4766
## fixed.slope.fit      2 39.917 65.890 1.4731      0.99658      1      0.3181
```

1b. Multivariate growth curves - start with this first (just using indicators - no latent variables) #As a rule of thumb you need at least three indicators for each latent variable.

```
## lavaan (0.5-23.1097) converged normally after 129 iterations
```

```
##
##                                     Used      Total
## Number of observations              310        348
##
## Number of missing patterns          44
##
## Estimator                          ML
## Minimum Function Test Statistic      6.070
## Degrees of freedom                   7
## P-value (Chi-square)                 0.532
##
```

```
## Parameter Estimates:
```

```
##
## Information                        Observed
## Standard Errors                    Standard
##
```

```
## Latent Variables:
```

```
##           Estimate Std.Err z-value P(>|z|)
## i.p =~
##   T3gcrs_cmbnd_c    1.000
##   T12gcrs_conv      1.000
##   T14gcrs_conv      1.000
## s.p =~
##   T3gcrs_cmbnd_c    0.000
##   T12gcrs_conv      1.000
##   T14gcrs_conv      2.000
## i.n =~
##   S1_FPNGEK1to5     1.000
##   S2_FPNGEK1to5     1.000
##   S3_FPNGEK1to5     1.000
## s.n =~
##   S1_FPNGEK1to5     0.000
##   S2_FPNGEK1to5     1.000
##   S3_FPNGEK1to5     2.000
##
```

```
## Covariances:
```

```
##           Estimate Std.Err z-value P(>|z|)
## i.p ~~
##   s.p          -0.002   0.007  -0.272   0.786
```

```

##      i.n          -0.000    0.003   -0.129    0.897
##      s.n          -0.001    0.002   -0.493    0.622
##    s.p ~~
##      i.n          0.000    0.002    0.205    0.838
##      s.n          0.000    0.001    0.013    0.990
##    i.n ~~
##      s.n          -0.000    0.001   -0.096    0.923
##
## Intercepts:
##              Estimate Std.Err  z-value  P(>|z|)
##    .T3gecrs_cmbnd_c    0.000
##    .T12gecrs_conv     0.000
##    .T14gecrs_conv     0.000
##    .S1_FPNGEK1to5     0.000
##    .S2_FPNGEK1to5     0.000
##    .S3_FPNGEK1to5     0.000
##      i.p             1.058    0.017   60.666    0.000
##      s.p            -0.013    0.012   -1.098    0.272
##      i.n             0.225    0.008   28.585    0.000
##      s.n             0.015    0.005    3.014    0.003
##
## Variances:
##              Estimate Std.Err  z-value  P(>|z|)
##    .T3gecrs_cmbnd_c    0.052    0.011    4.697    0.000
##    .T12gecrs_conv     0.021    0.005    4.702    0.000
##    .T14gecrs_conv     0.002    0.010    0.158    0.874
##    .S1_FPNGEK1to5     0.006    0.002    2.662    0.008
##    .S2_FPNGEK1to5     0.008    0.001    6.485    0.000
##    .S3_FPNGEK1to5     0.005    0.002    2.641    0.008
##      i.p             0.040    0.011    3.824    0.000
##      s.p             0.007    0.006    1.081    0.280
##      i.n             0.003    0.002    1.649    0.099
##      s.n             0.000    0.001    0.212    0.832
##
##              name idx nobs   type exo user  mean  var nlev lnam
##  1 T3gecrs_combined_conv 409  247 numeric  0  0 1.057 0.094  0
##  2      T12gecrs_conv 410  162 numeric  0  0 1.052 0.066  0
##  3      T14gecrs_conv 411   97 numeric  0  0 1.024 0.062  0
##  4      S1_FPNGEK1to5 339  124 numeric  0  0 0.231 0.009  0
##  5      S2_FPNGEK1to5 343  142 numeric  0  0 0.236 0.011  0
##  6      S3_FPNGEK1to5 401  132 numeric  0  0 0.258 0.009  0
##
## $lambda
##              i.p s.p i.n s.n
## T3gecrs_combined_conv  0  0  0  0
## T12gecrs_conv          0  0  0  0
## T14gecrs_conv          0  0  0  0
## S1_FPNGEK1to5         0  0  0  0
## S2_FPNGEK1to5         0  0  0  0
## S3_FPNGEK1to5         0  0  0  0
##
## $theta
##              T3gc_ T12gc_ T14gc_ S1_FPN S2_FPN S3_FPN
## T3gecrs_combined_conv 1
## T12gecrs_conv          0      2

```



```

## T14gecrs_conv      0      0      3
## S1_FPNGEK1to5      0      0      0      4
## S2_FPNGEK1to5      0      0      0      0      5
## S3_FPNGEK1to5      0      0      0      0      0      6
##
## $psi
##      i.p s.p i.n s.n
## i.p  7
## s.p 11  8
## i.n 12 14  9
## s.n 13 15 16 10
##
## $nu
##                               intrcp
## T3gecrs_combined_conv      0
## T12gecrs_conv              0
## T14gecrs_conv              0
## S1_FPNGEK1to5              0
## S2_FPNGEK1to5              0
## S3_FPNGEK1to5              0
##
## $alpha
##      intrcp
## i.p      17
## s.p      18
## i.n      19
## s.n      20

```

2a. Second order growth models - on BRIEF (using BRIEF composite scores (GEC = BRI + MI)

Begin with a simple CFA to determine if latent variable is appropriate

```

BRI.model <- ' BRI.T3 =~ T3_inhibrs + T3_shftrs + T3_emcnrs '
fit= cfa(BRI.model, data=growth_stats, missing= "ML")
summary(fit, fit.measures=TRUE)

```

```

## lavaan (0.5-23.1097) converged normally after  42 iterations
##
##                               Used      Total
##   Number of observations              67      348
##
##   Number of missing patterns              1
##
##   Estimator                          ML
##   Minimum Function Test Statistic      0.000
##   Degrees of freedom                   0
##
## Model test baseline model:

```

```

##
## Minimum Function Test Statistic          99.389
## Degrees of freedom                      3
## P-value                                0.000
##
## User model versus baseline model:
##
## Comparative Fit Index (CFI)              1.000
## Tucker-Lewis Index (TLI)                1.000
##
## Loglikelihood and Information Criteria:
##
## Loglikelihood user model (H0)            -563.818
## Loglikelihood unrestricted model (H1)    -563.818
##
## Number of free parameters                9
## Akaike (AIC)                            1145.636
## Bayesian (BIC)                          1165.479
## Sample-size adjusted Bayesian (BIC)     1137.141
##
## Root Mean Square Error of Approximation:
##
## RMSEA                                  0.000
## 90 Percent Confidence Interval          0.000  0.000
## P-value RMSEA <= 0.05                  NA
##
## Standardized Root Mean Square Residual:
##
## SRMR                                  0.000
##
## Parameter Estimates:
##
## Information                            Observed
## Standard Errors                      Standard
##
## Latent Variables:
##      Estimate  Std.Err  z-value  P(>|z|)
## BRI.T3 =~
##   T3_inhibrs      1.000
##   T3_shfters      0.691    0.112    6.167    0.000
##   T3_emcnrs       1.409    0.227    6.210    0.000
##
## Intercepts:
##      Estimate  Std.Err  z-value  P(>|z|)
##   .T3_inhibrs    18.731    0.717   26.128    0.000
##   .T3_shfters    14.194    0.464   30.590    0.000
##   .T3_emcnrs     19.045    0.736   25.870    0.000
##   BRI.T3         0.000
##
## Variances:
##      Estimate  Std.Err  z-value  P(>|z|)
##   .T3_inhibrs    16.907    3.474    4.867    0.000
##   .T3_shfters     6.056    1.379    4.392    0.000
##   .T3_emcnrs     1.510    3.743    0.404    0.687

```

```
##      BRI.T3      17.529      5.514      3.179      0.001
```

2b. Build second order growth models

```
sec.order <- '
###define latent variables
BRI_T3 =~ NA*T3_inhibrs + L1*T3_inhibrs + L2*T3_shftrs + L3*T3_emcnrs
BRI_T12 =~ NA*T12_inhibrs + L1*T12_inhibrs + L2*T12_shftrs + L3*T12_emcnrs
BRI_T14 =~ NA*T14_inhibrs + L1*T14_inhibrs + L2*T14_shftrs + L3*T14_emcnrs

### intercepts
T3_inhibrs ~ t1*1
T3_shftrs ~ t2*1
T3_emcnrs ~ t3*1

T12_inhibrs ~ t1*1
T12_shftrs ~ t2*1
T12_emcnrs ~ t3*1

T14_inhibrs ~ t1*1
T14_shftrs ~ t2*1
T14_emcnrs ~ t3*1

#this is setting the means equal across waves

## correlated residuals across time
T3_inhibrs ~~ T12_inhibrs + T14_inhibrs
T12_inhibrs ~~ T14_inhibrs
T3_shftrs ~~ T12_shftrs + T14_shftrs
T12_shftrs ~~ T14_shftrs
T3_emcnrs ~~ T12_emcnrs + T14_emcnrs
T12_emcnrs ~~ T14_emcnrs

## latent variable intercepts
BRI_T3 ~ 0*1
BRI_T12 ~ 0*1
BRI_T14 ~ 0*1

#model constraints for effect coding
## loadings must average to 1 (the three here changes to how many indicators you have; so change this b
L1 == 2 - L2
## means of indicators must average to 0 (in terms of the indicator means; )
t1 == 0 - t2

#the intercept and slope done with effect coding will give you the actual metric from your indicator va

#final step is the normal growth model
i =~ 1*BRI_T3 + 1*BRI_T12 + 1*BRI_T14
s =~ 0*BRI_T3 + 1*BRI_T12 + 2*BRI_T14 '

fit.sec.order <- growth(sec.order, data=growth_stats, missing = "ML")
```

```

## Warning in lav_data_full(data = data, group = group, cluster = cluster, : lavaan WARNING: some cases
##   1 4 6 8 9 11 13 15 19 21 22 25 27 28 29 31 33 34 36 37 38 39 40 41 45 48 49 51 52 53 54 55 56 57 5
## Warning in lav_data_full(data = data, group = group, cluster = cluster, :
## lavaan WARNING: due to missing values, some pairwise combinations have less
## than 10% coverage

## Warning in lav_object_post_check(object): lavaan WARNING: some estimated lv
## variances are negative

## Warning in lav_object_post_check(object): lavaan WARNING: the covariance matrix of the residuals of
##           variables (theta) is not positive definite;
##           use inspect(fit,"theta") to investigate.
summary(fit.sec.order, fit.measures=TRUE)

## lavaan (0.5-23.1097) converged normally after 173 iterations
##
##                                     Used      Total
##   Number of observations                205      348
##
##   Number of missing patterns              7
##
##   Estimator                          ML
##   Minimum Function Test Statistic      77.404
##   Degrees of freedom                   24
##   P-value (Chi-square)                 0.000
##
## Model test baseline model:
##
##   Minimum Function Test Statistic      680.471
##   Degrees of freedom                   36
##   P-value                             0.000
##
## User model versus baseline model:
##
##   Comparative Fit Index (CFI)          0.917
##   Tucker-Lewis Index (TLI)            0.876
##
## Loglikelihood and Information Criteria:
##
##   Loglikelihood user model (H0)        -2380.070
##   Loglikelihood unrestricted model (H1) -2341.368
##
##   Number of free parameters            30
##   Akaike (AIC)                        4820.139
##   Bayesian (BIC)                      4919.830
##   Sample-size adjusted Bayesian (BIC)  4824.779
##
## Root Mean Square Error of Approximation:
##
##   RMSEA                                0.104
##   90 Percent Confidence Interval        0.079  0.131
##   P-value RMSEA <= 0.05                0.000
##
## Standardized Root Mean Square Residual:

```

```

##
##      SRMR                                0.189
##
## Parameter Estimates:
##
##      Information                                Observed
##      Standard Errors                                Standard
##
## Latent Variables:
##      Estimate   Std.Err   z-value   P(>|z|)
##      BRI_T3 =~
##      T3_inhbrs (L1)    1.137    0.033    34.328    0.000
##      T3_shftrs (L2)    0.863    0.033    26.059    0.000
##      T3_emcnrs (L3)    1.341    0.067    19.986    0.000
##      BRI_T12 =~
##      T12_nhbrs (L1)    1.137    0.033    34.328    0.000
##      T12_shftr (L2)    0.863    0.033    26.059    0.000
##      T12_mcnrs (L3)    1.341    0.067    19.986    0.000
##      BRI_T14 =~
##      T14_nhbrs (L1)    1.137    0.033    34.328    0.000
##      T14_shftr (L2)    0.863    0.033    26.059    0.000
##      T14_mcnrs (L3)    1.341    0.067    19.986    0.000
##      i =~
##      BRI_T3            1.000
##      BRI_T12            1.000
##      BRI_T14            1.000
##      s =~
##      BRI_T3            0.000
##      BRI_T12            1.000
##      BRI_T14            2.000
##
## Covariances:
##      Estimate   Std.Err   z-value   P(>|z|)
##      .T3_inhibrs ~~
##      .T12_inhibrs      4.908    1.571    3.124    0.002
##      .T14_inhibrs      0.953    2.627    0.363    0.717
##      .T12_inhibrs ~~
##      .T14_inhibrs      4.287    0.837    5.124    0.000
##      .T3_shftrs ~~
##      .T12_shftrs      2.014    0.752    2.680    0.007
##      .T14_shftrs     -1.146    0.995   -1.152    0.249
##      .T12_shftrs ~~
##      .T14_shftrs      1.648    0.527    3.124    0.002
##      .T3_emcnrs ~~
##      .T12_emcnrs      0.949    1.346    0.705    0.481
##      .T14_emcnrs     -1.424    1.283   -1.110    0.267
##      .T12_emcnrs ~~
##      .T14_emcnrs      1.162    0.620    1.876    0.061
##      i ~~
##      s                2.367    1.671    1.417    0.157
##
## Intercepts:
##      Estimate   Std.Err   z-value   P(>|z|)
##      .T3_inhbrs (t1)  -0.491    0.420   -1.168    0.243

```

```



##      .T3_shftrs (t2)      0.491      0.420      1.168      0.243
##      .T3_emcnrs (t3)     -2.289      0.841     -2.720      0.007
##      .T12_nhbrs (t1)     -0.491      0.420     -1.168      0.243
##      .T12_shftr (t2)      0.491      0.420      1.168      0.243
##      .T12_mcnrs (t3)     -2.289      0.841     -2.720      0.007
##      .T14_nhbrs (t1)     -0.491      0.420     -1.168      0.243
##      .T14_shftr (t2)      0.491      0.420      1.168      0.243
##      .T14_mcnrs (t3)     -2.289      0.841     -2.720      0.007
##      BRI_T3              0.000
##      BRI_T12             0.000
##      BRI_T14             0.000
##      i                   13.277      0.356     37.302      0.000
##      s                   -0.915      0.196     -4.681      0.000
##
## Variances:
##      Estimate Std.Err z-value P(>|z|)
##      .T3_inhibrs      16.007      3.332      4.805      0.000
##      .T3_shftrs        5.298      1.282      4.131      0.000
##      .T3_emcnrs        5.594      2.388      2.342      0.019
##      .T12_inhibrs      6.317      0.904      6.989      0.000
##      .T12_shftrs       3.527      0.513      6.876      0.000
##      .T12_emcnrs       2.881      0.791      3.643      0.000
##      .T14_inhibrs      5.870      1.075      5.462      0.000
##      .T14_shftrs       4.034      0.677      5.960      0.000
##      .T14_emcnrs       0.947      0.725      1.306      0.192
##      BRI_T3           18.434      4.566      4.038      0.000
##      BRI_T12           1.314      1.073      1.225      0.221
##      BRI_T14           5.078      2.210      2.298      0.022
##      i                 3.950      2.990      1.321      0.186
##      s                -3.088      1.567     -1.971      0.049
##
## Constraints:
##                                     |Slack|
##      L1 - (2-L2)                  0.000
##      t1 - (0-t2)                  0.000

```

```

#josh's code
#fit.sec.order <- growth(sec.order, data=long, missing = "ML")
#summary(fit.sec.order, fit.measures=TRUE)

```



For longitudinal models, occasion specific variance can lead to biased estimates. We want to separate the time specific variance from the overall construct variance. Or, we want to make sure that the time specific variance doesn't make it appear that a construct is changing when really it is not.