

# Functions of miRNAs in Rice

Qian-Hao Zhu, Julien Curaba, Júlio César de Lima, and Chris Helliwell

**Abstract** As rice is both an important crop and a well-developed model for monocotyledonous plants, the study of miRNAs and their functions in this species is an active area of research. In this chapter, we summarize the progress that has been made in identifying miRNAs and their targets in rice. The identified miRNAs and their target mRNAs suggest that miRNAs function in a wide range of biological processes. The expression patterns of the miRNAs support their roles not only in development but also in response to environmental stresses. Demonstrations of biological functions for rice miRNAs are beginning to emerge and are detailed in this chapter.

## 1 Introduction

Rice is an important food crop. Having a relatively small genome (approximately 420 MB), it was among the earliest genomes sequenced (Goff et al. 2002; Yu et al. 2002). Coupled with extensive genomic resources such as EST collections and insertion mutant populations, this has led to rice becoming an important research species. The extensive synteny between rice and less tractable cereal crops such as wheat has meant that rice has also become a model system for cereal research. The availability of the rice genome sequence allowed identification of rice miRNA genes soon after the discovery of miRNAs in *Arabidopsis*. Numerous studies of rice

---

Q.-H. Zhu • J. Curaba • C. Helliwell (✉)  
CSIRO Plant Industry, GPO Box 1600, Canberra, ACT 2601, Australia  
e-mail: [chris.helliwell@csiro.au](mailto:chris.helliwell@csiro.au)

J.C. de Lima  
CSIRO Plant Industry, GPO Box 1600, Canberra, ACT 2601, Australia  
Department of Genetics, UFRGS, 99070-160 Porto Alegre, Brazil

miRNA populations and their biological roles have shown that they play important roles through the life cycle of the plant and in its response to environmental challenges.

## 2 Rice miRNAs and Their Target Genes

In the current miRBase (release 17), there are 491 rice miRNAs of 205 families (<http://www.mirbase.org/>; Table 1). These miRNAs were identified by sequencing of small RNA populations or by computational prediction using miRNAs first identified in other plant species. The first group of rice miRNAs to be identified (miR156–167 and miR169–172) was predicted computationally based on sequenced *Arabidopsis* miRNAs (Park et al. 2002; Reinhart et al. 2002) that are well conserved between dicotyledonous and monocotyledonous plants (Bonnet et al. 2004; Wang et al. 2004a). All of these bioinformatically predicted rice miRNAs were subsequently confirmed by small RNA cloning (Sunkar et al. 2005, 2008). Small RNA cloning followed by traditional Sanger sequencing played an important role in discovery of rice miRNAs, but this approach is biased toward miRNAs that are broadly and relatively highly expressed. miRNAs that are expressed at low abundance or only in certain cell types are not easily captured by this method (Jones-Rhoades et al. 2006). The development and application of next-generation high-throughput sequencing technology has greatly accelerated the discovery of miRNAs. It is now feasible to identify miRNAs with low expression, from specific tissues or induced by altered environmental conditions (Heisel et al. 2008; Li et al. 2010a; Lu et al. 2008; Morin et al. 2008; Sunkar et al. 2008; Xue et al. 2009; Zhu et al. 2008), and as a result, many more rice miRNAs have been identified (Table 1). There is often limited overlap between the newly identified miRNAs in different studies. This is a reflection of the differing tissue specificities or induction by environmental stimuli of the miRNAs, and it also suggests that new studies will identify further miRNAs.

### 2.1 Categories of Rice miRNAs

Rice miRNAs can be grouped into four categories based on their conservation in other plant species. Group I miRNAs (27 families) are conserved in at least one nonmonocotyledonous plant, Group II miRNAs (24 families) are conserved in at least one monocotyledonous plant, Group III miRNAs (145 families) are rice specific, and Group IV miRNAs (nine families) appear to be nonauthentic because they have not been found by high-throughput small RNA sequencing in any plant (Table 1).

**Table 1** Rice miRNAs, target genes, and their functions

miRNA family	No. of family members	Identification method <sup>a</sup>	Conservation <sup>b</sup>	Target genes confirmed by degradome sequencing or 5' RACE <sup>c</sup>	Functional investigation of miRNAs and their targets
<i>MIR156</i>	12	I, II <sup>[2,18]</sup>	I	<b><i>Os01g69830</i></b> (I) <sup>[33]</sup> <i>Os02g04680</i> (I) (II) <i>Os06g45310</i> (I) (II) <i>Os07g32170</i> (III) (II) <i>Os11g30370</i> (I) (II) <i>Os02g07780</i> (II) (II) <b><i>Os08g39890</i></b> (III) <sup>[6,33]</sup> <b>(<i>OsSPL14</i>)</b> <i>Os06g49010</i> (III) (II) <i>Os09g32944</i> (II) <i>Os09g31438</i> (III) <i>Os05g48800</i> (III) <i>Os07g36500</i> (III)	Plants overexpressing of <i>MIR156b</i> or <i>MIR156h</i> were semidwarf and had increased number of tillers (Xie et al. 2006). Interruption of miR156-mediated <i>OsSPL14</i> repression leads to higher level of <i>OsSPL14</i> , less tillers, and highly branched panicles with more grains (Jiao et al. 2010; Miura et al. 2010)
<i>MIR159</i>	6	I, II <sup>[17,22]</sup>	I	<b><i>Os01g59660</i></b> (I) (II) <sup>[15,33]</sup> <b>(<i>OsGAMYB</i>)</b> <i>Os06g40330</i> (II) (II) <b>(<i>OsGAMYBL1</i>)</b> <i>Os01g11430</i> (III) <i>Os10g05230</i> (III)	Overexpression of <i>MIR159a</i> causes shortened top internode, and shrunken, whitened anthers because of downregulation of <i>OsGAMYB</i> and <i>OsGAMYBL1</i> (Tsuji et al. 2006)
<i>MIR160</i>	6	I, II <sup>[2,18]</sup>	I	<i>Os06g47150</i> (I) (I) <i>Os10g33940</i> (I) (I) <i>Os02g41800</i> (II) (II) <i>Os04g43910</i> (II) (II) <i>Os04g59430</i> (I)	
<i>MIR162</i>	2	I, II <sup>[18,23]</sup>	I	<i>Os03g02970</i> (I)	

(continued)

**Table 1** (continued)

miRNA family	No. of family members	Identification method <sup>a</sup>	Conservation <sup>b</sup>	Target genes confirmed by degradome sequencing or 5' RACE <sup>c</sup>	Functional investigation of miRNAs and their targets
<i>MIR164</i>	6	I, II <sup>[2,18]</sup>	I	<i>Os06g46270</i> (I) (I) <i>Os08g10080</i> (I) (I) <i>Os12g41680</i> (I) (I) <b><i>Os02g36880</i></b> (III) (I) <sup>[33]</sup> <i>Os04g38720</i> (III) (II) <i>Os03g50040</i> (I) <i>Os05g39650</i> (I) <i>Os12g05260</i> (III)	Overexpression of <i>MIR164b</i> causes semidwarfism and low fertility (Zhu et al. unpublished)
<i>MIR166</i>	14	I, II <sup>[2,18]</sup>	I	<i>Os03g01890</i> (I) (III) ( <i>OSHB1</i> ) <i>Os10g33960</i> (I) (II) ( <i>OSHB2</i> ) <i>Os12g41860</i> (I) ( <i>OSHB3</i> ) <b><i>Os03g43930</i></b> (I) <sup>[13]</sup> ( <i>OSHB4</i> ) <i>Os04g48290</i> (III) <i>Os04g57610</i> (II)	Overexpressing miR166-resistant versions of <i>OSHB1</i> , <i>OSHB3</i> , and <i>OSHB5</i> in Taichung 65 exhibited rolled and/or filamentous leaves (Itoh et al. 2008). Overexpressing miR166-resistant version of <i>OSHB1</i> in <i>shoot organization1</i> ( <i>sho1</i> ) partially restored the morphological defects of <i>sho1</i> seedlings (Nagasaki et al. 2007)
<i>MIR167</i>	10	I, II <sup>[2,18]</sup>	I	<b><i>Os04g57610</i></b> (II) <sup>[13,29]</sup> <i>Os02g06910</i> (I) <i>Os06g46410</i> (I) <i>Os12g41950</i> (I) <b><i>Os06g03830</i></b> (I) <sup>[10]</sup> <i>Os07g33790</i> (I) <i>Os07g29820</i> (II) <b><i>Os09g39420</i></b> <sup>[13]</sup> <b><i>Os04g47870</i></b> (II) <sup>[33]</sup> <b><i>Os02g45070</i></b> (III) <sup>[33]</sup> <i>Os06g51310</i> (III) <b><i>Os02g58490</i></b> (III) <sup>[13,33]</sup>	Cultured rice cells delivered with artificial miR167 showed decreased mRNAs of <i>ARF8</i> and <i>OsGH3-2</i> , a rice IAA-conjugating enzyme (Yang et al. 2006a)
<i>MIR168</i>	2	I, II <sup>[8,22]</sup>	I		

<i>MIR169</i>	17	I, II <sup>[18,22]</sup>	I	<i>Os02g53620</i> (I) (II) <sup>[13]</sup> <i>Os03g07880</i> (I) (I) <sup>[10]</sup> <i>Os03g29760</i> (I) (II) <sup>[30]</sup> <i>Os03g44540</i> (I) <sup>[10]</sup> <i>Os03g48970</i> (I) <i>Os07g06470</i> (I) (III) <i>Os07g41720</i> (I) (II) <i>Os12g42400</i> (I) (I) <sup>[13,33]</sup>	
<i>MIR171</i>	9	I, II <sup>[17,25]</sup>	I	<i>Os02g44360</i> (I) (II) <i>Os04g46860</i> (I) (II) <i>Os02g44370</i> (III) (II) <i>Os04g39864</i> (I) <sup>[10]</sup>	
<i>MIR172</i>	4	I, II <sup>[17,22]</sup>	I	<i>Os05g03040</i> (I) (II) <sup>[13]</sup> <i>Os07g13170</i> (III) (II) <sup>[32]</sup> <i>Os04g55560</i> (I) (II) <sup>[32]</sup> <i>Os03g60430</i> (III) <i>Os08g39630</i> (I) <i>Os01g04550</i> (I) <i>Os06g43220</i> <sup>[32]</sup>	Overexpression of <i>MIR172b</i> but not <i>MIR172a</i> causes deformed spikelets (Zhu et al. 2009)
<i>MIR319</i>	2	I, II <sup>[8,23]</sup>	I	<i>Os07g05720</i> (III) (III) <i>Os03g57190</i> (III)	
<i>MIR390</i>	1	II <sup>[22]</sup>	I	<i>EU293144</i> (I) <i>AU100890</i> (I) <i>CA765877</i> (I)	
<i>MIR393</i>	2	I, II <sup>[8,22]</sup>	I	<i>Os05g05800</i> (I) (I) <i>Os04g32460</i> (I) (II) <i>Os03g36080</i> (II) <i>Os04g58734</i> (III) <sup>[10]</sup>	Overexpression of <i>MIR393</i> decreases salt and alkali stress tolerance (Gao et al. 2011)
<i>MIR394</i>	1	I, II <sup>[8,23]</sup>	I	<i>Os01g69940</i> (I) (II)	
<i>MIR395</i>	25	I, II <sup>[8,23]</sup>	I	<i>Os06g05160</i> (II)	

(continued)

**Table 1** (continued)

miRNA family	No. of family members	Identification method <sup>a</sup>	Conservation <sup>b</sup>	Target genes confirmed by degradome sequencing or 5' RACE <sup>c</sup>	Functional investigation of miRNAs and their targets
<i>MIR396</i>	9	I, II <sup>[8,23]</sup>	I	<i>Os02g47280</i> (I) (I) <i>Os03g47140</i> (I) <i>Os12g29980</i> (I) <i>Os04g51190</i> (III) <i>Os11g35030</i> (I) (I) <i>Os06g02560</i> (I) (II) <i>Os02g45570</i> (I) <i>Os02g53690</i> (III) <i>Os12g29980</i> (I) <i>Os04g48510</i> (III) <i>Os03g51970</i> (III) <i>Os01g32750</i> (III) <b><i>Os11g35030</i></b> (I) <sup>[10]</sup> <i>Os01g44230</i> (III) <i>Os06g29430</i> (I)	Overexpression of <i>MIR396c</i> causes decreased salt and alkali stress tolerance (Gao et al. 2010)
<i>MIR397</i>	2	I, II <sup>[8,23]</sup>	I	<i>Os11g48060</i> (III)	
<i>MIR398</i>	2	I, II <sup>[8,23]</sup>	I	<i>Os07g46990</i> (I) (II) <i>Os03g22810</i> (III) <b><i>Os04g48410</i></b> (II) <sup>[10]</sup>	
<i>MIR399</i>	11	I, II <sup>[8,23]</sup>	I	<i>Os05g48390</i> (III)	
<i>MIR408</i>	1	I, II <sup>[21,22]</sup>	I	<i>Os02g52180</i> (I) <i>Os04g46130</i> (I) <i>Os06g11490</i> (I) <i>Os08g37670</i> (I)	
<i>MIR413–MIR426</i>	1 for each	I <sup>[24]</sup>	IV		
<i>MIR435</i>	1	II <sup>[22]</sup>	III		

<i>MIR437</i>	1	II <sup>[22]</sup>	II	
<i>MIR438, MIR440,</i> <i>MIR442, MIR443</i>	1 for each	II <sup>[22]</sup>	III	
<i>MIR439</i>	10	II <sup>[22]</sup>	III	
<i>MIR441</i>	3	II <sup>[22]</sup>	III	
<i>MIR444</i>	6	II <sup>[22]</sup>	II	<i>Os08g33488</i> (I) <b><i>Os02g36924</i></b> (I) (I) <sup>[14]</sup> <b><i>Os04g38780</i></b> (I) (II) <sup>[14]</sup> <b><i>Os02g49840</i></b> (I) (II) <sup>[14]</sup> <i>Os02g49090</i> (I) <b><i>Os03g63750</i></b> (I) <sup>[10]</sup> <i>Os05g08410</i> (I) <i>Os08g33479</i> (III) <b><i>Os02g13420</i></b> (I) <sup>[10]</sup> <i>Os03g54084</i> (II) <i>Os05g47560</i> (II) <i>Os08g06510</i> (I)
<i>MIR445</i>	9	II <sup>[22]</sup>	III	
<i>MIR446</i>	1	II <sup>[22]</sup>	III	<i>Os02g29140</i> (III) <i>Os04g38450</i> (III) <i>Os06g19990</i> (III) <i>Os10g26720</i> (III) <b><i>Os06g06050</i></b> (I) (III) <sup>[10,33]</sup> <i>Os06g11310</i> (I) <b><i>Os06g37150</i></b> (I) (I) <sup>[33]</sup> <i>Os06g47780</i> (I) <i>Os06g04270</i> (II) <i>Os02g14990</i> (I)
<i>MIR528</i>	1	II <sup>[12]</sup>	II	
<i>MIR529</i>	2	II <sup>[12]</sup>	I	
<i>MIR530</i>	1	II <sup>[12]</sup>	I	
<i>MIR531</i>	2	II <sup>[12]</sup>	III	
<i>MIR535</i>	1	I, II <sup>[1,23]</sup>	I	
<i>MIR806</i>	8	II <sup>[15]</sup>	III	

(continued)

**Table 1** (continued)

miRNA family	No. of family members	Identification method <sup>a</sup>	Conservation <sup>b</sup>	Target genes confirmed by degradome sequencing or 5' RACE <sup>c</sup>	Functional investigation of miRNAs and their targets
<i>MIR807, MIR815</i>	3 for each	II <sup>[15]</sup>	III		
<i>MIR808</i>	1	II <sup>[15]</sup>	III	<i>Os10g39970</i> (I) <i>Os12g40920</i> (I) <i>Os12g44250</i> (II) <i>Os04g08415</i> (I) <i>Os06g33210</i> (I) <i>Os10g40540</i> (I) <i>Os02g55480</i> (II) <b><i>Os10g39520</i></b> <sup>[15]</sup>	
<i>MIR809</i>	8	II <sup>[15]</sup>	III		
<i>MIR810</i>	2	II <sup>[15]</sup>	III		
<i>MIR811</i>	4	II <sup>[15]</sup>	III		
<i>MIR812</i>	15	II <sup>[15]</sup>	III	<i>Os03g22050</i> (II)	
<i>MIR813, MIR816, MIR817</i>	1 for each	II <sup>[15]</sup>	III		
<i>MIR814</i>	3	II <sup>[15]</sup>	III	<i>Os12g39380</i> (I)	
<i>MIR818</i>	5	II <sup>[15]</sup>	III	<b><i>Os09g10710</i></b> <sup>[15]</sup>	
<i>MIR819</i>	11	II <sup>[15]</sup>	III		
<i>MIR820</i>	3	II <sup>[15]</sup>	III	<b><i>Os03g02010</i></b> (II) <sup>[14,15,27]</sup>	
<i>MIR821</i>	3	II <sup>[15]</sup>	II ( <i>S. bicolor</i> )		
<i>MIR827</i>	3	II <sup>[3]</sup>	I	<b><i>Os04g48390</i></b> (II) <sup>[3,9]</sup> <b><i>Os02g45520</i></b> <sup>[3]</sup>	
<i>MIR1317, MIR1319</i>	1 for each	II <sup>[16]</sup>	III		
<i>MIR1318/1432</i>	1	II <sup>[16]</sup>	II		
<i>MIR1320</i>	1	II <sup>[16]</sup>	III	<i>Os05g47550</i> (II)	
<i>MIR1423</i>	2	II <sup>[14]</sup>	III		



<i>MIR1424, MIR1427,</i> <i>MIR1429, MIR1430,</i> <i>MIR1425</i>	1 for each  1	II <sup>[14]</sup>  II <sup>[14]</sup>	III  III	<i>Os08g01640</i> (I) <i>Os08g01650</i> (I) <i>Os08g01870</i> (I) <i>Os10g35230</i> (I) <b><i>Os10g35240</i></b> (I) (I) <sup>[9,14,33]</sup> <b><i>Os10g35436</i></b> (I) <sup>[9,14]</sup> <b><i>Os10g35640</i></b> (I) <sup>[14,33]</sup> <i>Os01g49580</i> (III) <i>Os01g49614</i> (III) <i>Os05g40700</i> (III) <i>Os05g30350</i> (I) <b><i>Os03g17980</i></b> (III) (II) <sup>[31]</sup> <b><i>Os08g37800</i></b> <sup>[31]</sup> <i>Os02g16670</i> (I)
<i>MIR1426</i>	1	II <sup>[14]</sup>	III	
<i>MIR1428</i>	7	II <sup>[14]</sup>	III	
<i>MIR1431</i>	1	II <sup>[14]</sup>	III	
<i>MIR1433</i>	1	II <sup>[14]</sup>	II ( <i>T. aestivum</i> )	
<i>MIR1435</i>	1	II <sup>[23]</sup>	II ( <i>S. bicolor</i> )	
<i>MIR1436</i>	1	II <sup>[23]</sup>	II ( <i>H. vulgare</i> )	<i>Os03g48010</i> (II)
<i>MIR1437, MIR1438,</i> <i>MIR1440, MIR1441</i>	1	II <sup>[23]</sup>	III	
<i>MIR1439</i>	1	II <sup>[23]</sup>	III	<i>Os09g33690</i> (I)
<i>MIR1442</i>	1	II <sup>[23]</sup>	III	<i>Os02g58670</i> (II)
<i>MIR1846</i>	5	II <sup>[31]</sup>	III	<i>Os06g14060</i> (I) <i>Os09g07510</i> (I)
<i>MIR1847</i>	1	II <sup>[31]</sup>	III	<i>Os02g34990</i> (I) <i>Os04g39170</i> (I)
<i>MIR1848</i>	1	II <sup>[31]</sup>	III	<i>Os01g49720</i> (I)
<i>MIR1849</i>	1	II <sup>[31]</sup>	III	

(continued)

**Table 1** (continued)

miRNA family	No. of family members	Identification method <sup>a</sup>	Conservation <sup>b</sup>	Target genes confirmed by degradome sequencing or 5' RACE <sup>c</sup>	Functional investigation of miRNAs and their targets
<i>MIR1850</i>	1	II <sup>[31]</sup>	III	<i>Os01g02360</i> (I) <i>Os09g32250</i> (I) <i>Os05g05140</i> (I) <i>Os01g05790</i> (I) <i>Os04g47410</i> (II) <b><i>AK068321</i></b> <sup>[31]</sup>	
<i>MIR1851–MIR1856</i>	1 for each	II <sup>[31]</sup>	III		
<i>MIR1857</i>	1	II <sup>[31]</sup>	III	<i>Os11g45590</i> (III)	
<i>MIR1858</i>	2	II <sup>[31]</sup>	III		
<i>MIR1859–MIR1860</i> , <i>MIR1864–MIR1872</i> , <i>MIR1874–MIR1875</i> , <i>MIR1877</i> , <i>MIR1879–MIR1881</i>	1 for each	II <sup>[31]</sup>	III		
<i>MIR1861</i>	14	II <sup>[31]</sup>	III	<b><i>Os05g51790</i></b> <sup>[31]</sup>	
<i>MIR1862</i>	7	II <sup>[31]</sup>	III		
<i>MIR1863</i>	3	II <sup>[31]</sup>	I ( <i>P. abies</i> )	<i>^Os06g38480</i> <sup>[27]</sup>	
<i>MIR1873</i>	1	II <sup>[31]</sup>	III	<i>^Os05g01790</i> <sup>[27]</sup>	
<i>MIR1876</i>	2	II <sup>[31]</sup>	III	<i>^Os07g41090</i> <sup>[27]</sup> <i>^Os02g05890</i> <sup>[27]</sup>	
<i>MIR1878</i>	1	II <sup>[31]</sup>	II ( <i>B. distachyon</i> )		
<i>MIR1882</i>	8	II <sup>[31]</sup>	III		
<i>MIR1883</i>	2	II <sup>[31]</sup>	III		

<i>MIR1884</i>	2	II <sup>[31]</sup>	III	<i>Os01g59720</i> (II) <i>Os01g64520</i> (III) <i>Os02g22610</i> (I) <i>Os03g10250</i> (I) <i>Os03g19380</i> (II) <i>Os11g34910</i> (II)
<i>MIR2055</i>	1	II <sup>[9]</sup>	III	
<i>MIR2090–MIR2097</i> , <i>MIR2099</i> , <i>MIR2100</i> , <i>MIR2104</i> , <i>MIR2106</i>	1 for each	II <sup>[28]</sup>	III	
<i>MIR2098</i>	1	II <sup>[28]</sup>	III	<i>Os03g23050</i> (I) (for 3p) <i>Os07g03279</i> (II) <i>Os07g03368</i> (II) <i>Os07g03458</i> (II) (for 5p)
<i>MIR2101</i>	1	II <sup>[28]</sup>	III	<i>Os12g08210</i> (I)
<i>MIR2102</i>	1	II <sup>[28]</sup>	III	<i>Os05g37460</i> (I)
<i>MIR2103</i>	1	II <sup>[28]</sup>	III	<i>Os03g52880</i> (I)
<i>MIR2105</i>	1	II <sup>[28]</sup>	III	<i>Os01g14040</i> (I) <i>Os01g25484</i> (I)
<i>MIR2118</i>	18	II <sup>[7]</sup>	I	
<i>MIR2120</i> , <i>MIR2122</i> , <i>MIR2125</i>	1 for each	II <sup>[4]</sup>	III	
<i>MIR2121</i>	2	II <sup>[4]</sup>	III	
<i>MIR2123</i>	3	II <sup>[4]</sup>	III	
<i>MIR2124</i>	9	II <sup>[4]</sup>	III	
<i>MIR2275</i>	2	II <sup>[7]</sup>	II ( <i>Z. mays</i> )	
<i>MIR2862</i>	1	II <sup>[26]</sup>	III	
<i>MIR2863</i>	2	II <sup>[26]</sup>	III	

(continued)

**Table 1** (continued)

miRNA family	No. of family members	Identification method <sup>a</sup>	Conservation <sup>b</sup>	Target genes confirmed by degradome sequencing or 5' RACE <sup>c</sup>	Functional investigation of miRNAs and their targets
<i>MIR2864–MIR2867</i> , <i>MIR2869</i> , <i>MIR2870</i> , <i>MIR2872</i> , <i>MIR2874–MIR2880</i> , <i>MIR2905</i>	1	II <sup>[26]</sup>	III		
<i>MIR2868</i>	1	II <sup>[26]</sup>	III	<i>Os11g14140</i> (II)	
<i>MIR2871</i> , <i>MIR2873</i>	2 for each	II <sup>[26]</sup>	III		
<i>MIR2906</i>	2	II <sup>[5]</sup>	III		
<i>MIR2907</i>	4	II <sup>[5]</sup>	III		
<i>MIR2918–MIR2932</i>	1	II <sup>[19]</sup>	III		
<i>MIR3979</i> , <i>MIR3981</i> , <i>MIR3982</i>	1 for each	II <sup>[10]</sup>	III		
<i>MIR3980</i>	2	II <sup>[10]</sup>	III		
<i>MIR5071–MIR5083</i>	1 for each	I <sup>[20]</sup>	II		

<sup>a</sup>I—homology searching using miRNAs identified in other species; II—small RNA cloning. The paper that first reported the miRNA is shown. For the miRNAs that were first identified by homology searching, the paper that first confirmed their presence in rice is also shown

<sup>b</sup>I—conserved in rice and at least one nonmonocotyledonous plant; II—conserved in rice and at least one other monocotyledonous plant; III—rice specific; IV—supposed to be conserved in *Arabidopsis* and rice, but they have not been discovered by high-throughput sequencing in both species. For the rice miRNA families that are conserved only in one other plant species, the name of the species is given (based on the information available in miRBase)

<sup>c</sup>Target genes that were identified by degradome sequencing, confirmed by 5' RACE or shown to be methylated, are shown. I, II, and III in parentheses indicate the classes of target as shown in the original papers [11] and [26] (the first and second pair of parentheses, respectively). Targets that were confirmed by 5' RACE are bolded, and the reference that performed the confirmation is also shown. Targets starting with ^ are not cleaved but methylated by miRNA as shown in ref. [27]

[1] Arazi et al. (2005), [2] Bonnet et al. (2004), [3] Heisel et al. (2008), [4] Huang et al. (2009), [5] Jian et al. (2010), [6] Jiao et al. (2010), [7] Johnson et al. (2009), [8] Jones-Rhoades and Bartel (2004), [9] Lacombe et al. (2008), [10] Li et al. (2010a), [11] Li et al. (2010b), [12] Liu et al. (2005), [13] Liu et al. (2009), [14] Lu et al. (2008), [15] Luo et al. (2006), [16] Morin et al. (2008), [17] Park et al. (2002), [18] Reinhart et al. (2002), [19] Sanan-Mishra et al. (2009), [20] Schreiber et al. (2011), [21] Sunkar and Zhu (2004), [22] Sunkar et al. (2005), [23] Sunkar et al. (2008), [24] Wang et al. (2004b), [25] Wang et al. (2004a), [26] Wu et al. (2009), [27] Wu et al. (2010), [28] Xue et al. (2009), [29] Yang et al. (2006a), [30] Zhao et al. (2009), [31] Zhu et al. (2008), [32] Zhu et al. (2009), [33] Zhu et al. (unpublished)

### 2.1.1 Group I miRNAs and Their Targets

Of the 27 Group I miRNA families, 21 families (miR156, 159, 160, 162, 164, 166–169, 171, 172, 319, 390, 393–399, and 408) are well conserved among diverse angiosperms. At least three families (miR156, 160, and 166) are also conserved in gymnosperms, lycopods, and bryophytes, indicating an ancient origin of these miRNAs. Target genes of 11 conserved miRNA families (miR156, 159, 160, 164, 166, 167, 169, 171, 172, 319, and 396) encode transcription factors that have a conserved role in plant development. For example, miR172 functions in regulating the transitions between developmental stages and in specifying floral organ identity by regulating expression of AP2-domain-containing transcription factors in monocots and dicots (Zhu and Helliwell 2011). The Class III *HD-ZIP* transcription factors are targets of miR166. The rice genome contains five Class III *HD-ZIP* genes: *OSHB1* to *OSHB5*. All these five genes have the miR166 binding site, and four of them (*OSHB1* to *OSHB4*) have been confirmed to be regulated by miR166 (Table 1). Overexpression of miR166-resistant versions of *OSHB1*, *OSHB3*, and *OSHB5* exhibited rolled and/or filamentous leaves caused by loss-of-leaf polarity (Itoh et al. 2008). Mutations in Class III *HD-ZIP* genes, which reduce miR166 complementarity in *Arabidopsis* and maize, confer similar phenotypes (Juarez et al. 2004; McConnell et al. 2001). Further studies have shown that proper leaf polarity in rice and maize is established by a coordinated expression of miR166 and *tasiARF* in opposing domains along the adaxial–abaxial axis (Nagasaki et al. 2007; Nogueira et al. 2007). These results suggest that conserved miRNAs play a similar role by targeting conserved genes in different plants, although nonconserved target genes have been identified (Li et al. 2010b).

Of the remaining six Group I miRNAs (miR529, 530, 535, 827, 1863, and 2118), only miR827 is conserved in *Arabidopsis* (Heisel et al. 2008; Lacombe et al. 2008), which targets *SPX*-domain-containing genes in rice and *Arabidopsis*. The *SPX*-domain is associated with proteins that have been implicated in transport or sensing of phosphate or nitrogen in *Arabidopsis* (Peng et al. 2007; Wang et al. 2004c). It is expected the two confirmed miR827 targets (*Os02g45520* and *Os02g48390*) could play a similar role in rice development. Homologs of miR529 and miR535 are found in moss (Arazi et al. 2005), and a homolog of miR1863 has been found in gymnosperms (*Picea abies*) (Yakovlev et al. 2010), suggesting that these miRNAs have an ancient origin but could have been lost in eudicots (miR529) or *Arabidopsis* (miR535) after their separation from the monocotyledonous plant lineage. No homolog of miR530 has been found in *Arabidopsis*, but it is conserved in several dicotyledonous and monocotyledonous plants.

### 2.1.2 Group II miRNAs and Their Targets

Group II, the monocot-conserved miRNAs, contains 24 miRNA families, of which miR444 and miR528 are well conserved in all monocotyledons investigated.

The miR444 family is encoded by four loci in the rice genome. A few features of this miRNA family make it unique. First, in each locus, the pri-miR444 is overlapping and antisense to its target MADS box gene. Second, the pre-miR444 is processed from a spliced pri-miR444 that possesses large introns. Third, more than one functional miRNA (overlapping or in phase) is processed from each pre-miR444 (Lu et al. 2008). Due to the natural antisense characteristics of pre-miR444 and its MADS targets, different miR444 variants from the same pre-miR444 are expected to cleave MADS target mRNAs. This was confirmed by recent degradome sequencing (Li et al. 2010b; Wu et al. 2009), indicating that different miR444 variants are functional, and the MADS genes are collectively regulated by miR444 variants at multiple positions with different cleavage frequencies. In both of the degradome sequencing studies (Li et al. 2010b; Wu et al. 2009), non-MADS targets were also found, and two of them were further confirmed by 5' RACE (Li et al. 2010b), demonstrating that miR444, like some other deeply conserved miRNAs, can regulate targets unrelated to the conserved ones.

The miR528 family of rice contains a single miRNA (Liu et al. 2005), which is predicted to target several genes with unrelated functions (Li et al. 2010b; Wu et al. 2009; Xue et al. 2009), three of them (*Os06g06050*, *Os06g37150*, and *Os06g11310*) were confirmed by degradome sequencing and/or 5' RACE (Table 1). *Os06g37150* encodes L-ascorbate oxidase, which catalyzes the oxidation of ascorbic acid to yield dehydroascorbate to regulate the apoplastic redox state and modulate plant growth and defense responses by regulating signal transduction cascades (Pignocchi et al. 2006). miR528 is highly expressed in the embryo compared to vegetative tissues (Xue et al. 2009; consistent with our *P<sub>miR528</sub>::GUS* transgene results, Zhu et al. unpublished). miR528-mediated downregulation of *Os06g37150* has been observed in the embryo (Xue et al. 2009). This may be required to maintain a higher level of cell division in developing rice grains, because it has been found that accumulation of dehydroascorbate oxidized by L-ascorbate oxidase can trigger the arrest of cell division (Potters et al. 2000). miR528-mediated regulation of *OsCBP1* ( $\text{Cu}^{2+}$ -binding domain-containing protein 1, *Os07g38290*), one of the predicted miR528 targets (Wu et al. 2009), has not been confirmed in rice, but its sugarcane ortholog *SsCBP1* has been shown to be cleaved by miR528 (Zanca et al. 2010). Regulation of plant *CBPs* by miR528 seems to be conserved in monocotyledons and has evolved exclusively within the monocotyledons after their divergence from eudicots (Zanca et al. 2010). As for miR528, a few other miRNAs that are highly expressed in embryo also have target genes encoding copper-binding proteins that are downregulated in embryo, suggesting a role for these miRNAs in maintaining copper and reactive oxygen species homeostasis in the rice embryo (Xue et al. 2009).

Of the other Group II miRNAs, miR821, miR1433, miR1435, miR1436, miR1878, and miR2275 are only conserved in one other monocotyledonous plant (Table 1), whereas miR1318/1432 and miR437 are conserved across multiple monocotyledonous plants. With the exception of miR1436, all these miRNAs do not have targets confirmed by degradome sequencing or 5' RACE (Table 1). The remaining 13 Group II miRNAs (miR5071–5083) are computational predictions

based on cloned barley miRNAs (Schreiber et al. 2011). Their presence in rice has not been confirmed by small RNA sequencing.

### 2.1.3 Group III miRNAs and Their Targets

The majority of annotated rice miRNAs appear to be species specific. They were identified based on small RNA cloning followed by traditional or high-throughput sequencing using various tissues, such as seedlings (Morin et al. 2008; Sunkar et al. 2005; 2008), seeds (Xue et al. 2009; Zhu et al. 2008) and callus (Luo et al. 2006), or using stress-treated samples (Huang et al. 2009; Li et al. 2010a; Lu et al. 2008). miRNAs were also identified from small RNA populations associated with AGO1 (Wu et al. 2009). As most of these miRNAs were identified from specific tissues or plants growing under altered growth conditions, it is expected that some of them will have a specific expression profile. For example, miR1428e is highly expressed in developing grains but not detected in roots (Zhu et al. 2008). However, for the majority of rice-specific miRNAs, their tissue specificity is yet to be explored.

The target genes of rice-specific miRNAs encode a wide range of proteins; some of these mRNAs have been shown to be cleaved by miRNAs (Table 1). Generally, the low expression level and/or tissue specificity of these miRNAs means that confirmation of miRNA-mediated regulation of putative target mRNAs is not as easy as for highly expressed conserved miRNAs. In addition to miRNA-mediated cleavage and translational repression, recent investigation in rice showed that 24-nt miRNAs were absent from the AGO1 complex but were loaded into AGO4 clade proteins to induce DNA methylation in *cis* at loci from which they are generated or in *trans* at their targets, suggesting that these miRNAs may transcriptionally regulate expression of themselves and their target genes (Wu et al. 2009, 2010). It is worthy of attention that *Os03g02010*, a target of miR820a, was subjected to both miRNA-mediated cleavage (Lu et al. 2008; Luo et al. 2006) and DNA methylation (Wu et al. 2010), presumably induced by 21-nt and 24-nt miRNA variants, respectively. This is consistent with our previous small-RNA-sequencing result for the *MIR820a* locus (Zhu et al. 2008), from which 94 reads of 21-nt and 2,208 reads of 24-nt miRNAs (the most abundant miRNA variant) were generated. This phenomenon may apply to many other *MIRNA* genes because different lengths of miRNA variants are generated from the same region of many pre-miRNAs.

miRNAs derived from exonic regions are rare but are found in RNA populations from human cancer cells (Lui et al. 2007). It was recently demonstrated in rice plants that miR3981 is processed from an exon region from a gene that encodes for a glyoxylase. More interestingly, by 5' RACE, it was demonstrated that miR3981 targets its own precursor gene (Li et al. 2010a). Similarly, degradome sequencing has shown that miRNAs regulate their primary precursors (German et al. 2008).

### 3 miRNAs Are Required for Correct Growth and Development in Rice

miRNA-mediated gene regulation is implicated in most if not all major developmental steps of the plant life cycle, and the function of the well-conserved miRNA families seems to have been preserved across species (Chen 2009). For example, miR156, which is involved in shoot apical meristem (SAM) development, controls vegetative to floral phase transition in *Arabidopsis* and maize, and shoot branching in rice (Chuck et al. 2007; Jiao et al. 2010; Miura et al. 2010; Yang et al. 2010). Although specific functions of each conserved miRNA may differ across species, the key role of miRNAs in regulating plant development is undeniable, and plants affected in miRNA biogenesis or action show severe pleiotropic phenotypes.

In *Arabidopsis*, biogenesis of most miRNAs requires DCL1 for the two initial processing steps of the precursor (Bouche et al. 2006; Kurihara and Watanabe 2004). Loss-of-function *AtDCL1* mutants dramatically affect the accumulation of miRNAs and lead to a wide range of phenotypes, ranging from altered organ morphology to developmental arrest (McElver et al. 2001; Park et al. 2002; Reinhart et al. 2002; Schauer et al. 2002). Consistently, mutants of *AtAGO1*, an essential enzyme of the miRNA effector complex, display similar pleiotropic developmental defects associated with impaired miRNA function (Vaucheret et al. 2004).

In rice, RNA-interference-mediated silencing of *OsDCL1a* (*Os03g02970*), the homolog of *AtDCL1*, leads to abnormal shoot and root development with eventual growth arrest for the strongest RNAi lines (Liu et al. 2005). The aerial part of the plant shows severe dwarfism and rolled dark green leaves. The root system is dramatically affected, having reduced elongation and fewer adventitious roots together with the presence of ectopic chloroplasts. At a molecular level, RNAi-*OsDCL1a* plants show impaired accumulation of all seven mature miRNAs tested but none of the tested siRNAs, suggesting that *OsDCL1a*, like *AtDCL1*, is required for miRNA biogenesis. The phenotypes of RNAi-*OsDCL1a* can be therefore considered as direct consequences of a loss-of-miRNA function.

miRNA action in rice seems to be driven by at least four partially redundant AGO1-like proteins. Plants in which the *OsAGO1s* are silenced using an inverted repeat RNAi construct targeting all four AGO1 homologs (*OsAGO1a* (*Os02g45070*), *OsAGO1b* (*Os04g47870*), *OsAGO1c* (*Os02g58490*), and *OsAGO1d* (*Os06g51310*)) show pleiotropic phenotypes similar to those observed in RNAi-*OsDCL1a*, including dwarfism with narrow, rolled leaves, and developmental arrest for the strongest line (Wu et al. 2009). Accordingly, five previously confirmed targets of known miRNAs were shown to be upregulated in both RNAi-*OsAGO1s* and RNAi-*OsDCL1a* plants. The severe phenotypes associated with loss-of-miRNA function highlight the crucial role of miRNAs in the growth and development of the rice plant.



## 4 Expression of miRNAs and miRNA Biogenesis Machinery

### 4.1 Regulation of the miRNA Biogenesis Machinery

The initial step of plant miRNA biogenesis is the transcription of a *MIR* gene by RNA polymerase II. The primary transcript (pri-miRNA) forms an imperfect hairpin that is processed into a small RNA duplex in two or three steps by an enzyme complex containing at least three proteins: DICER-LIKE1 (DCL1), HYPONASTIC LEAVES1 (HYL1), and SERRATE (SE). The small RNA duplex, composed of the miRNA and its opposite strand, the miRNA\*, is protected from exonuclease degradation through the methylation of its 3' ends by HUA ENHANCER1 (HEN1). Exported outside the nucleus by HASTY (HST), the miRNA is then incorporated into an ARGONAUTE (AGO) protein to form the miRNA-induced silencing complex (miRISC) and trigger the posttranscriptional regulation of its target gene(s). A genome-wide analysis of the rice genome revealed the presence of homologs for all the currently known components of the miRNA biogenesis machinery, including eight *OsDCLs*, 19 *OsAGOs*, three *SE* (*Os06g48530*, *Os08g40560*, *Os02g05610*), four *HYL1* (*Os12g01916*, *Os05g24160*, *Os11g01869*, *Os01g56520*), and interestingly, only one copy of *HST* (*Os01g26160*) and *HEN1* (*Os07g06970*) homologs (data collected from Kapoor et al. 2008 and BLAST searches using the *Arabidopsis* gene sequences against the rice genome available at <http://rice.plantbiology.msu.edu/index.shtml>).

miRNA biogenesis in *Arabidopsis* requires *AtDCL1*, a gene with three homologs in rice: *OsDCL1a* (*Os03g02970*), *OsDCL1b* (*Os06g25250*), and *OsDCL1c* (*Os05g18850*) (Kapoor et al. 2008; Margis et al. 2006). Based on their sequence similarity and chromosome location none appear to be the result of tandem duplication. *OsDCL1a* is most closely related to *AtDCL1*, possessing all the six domains characteristic of the Dicer family; silencing this gene abolishes the biogenesis of miRNAs (Liu et al. 2005). On the other hand, *OsDCL1b* and *OsDCL1c* encode for smaller proteins containing only the RNaseIII and dsRB domains, matching the C-terminal region of *OsDCL1a*. Although these two domains are crucial for the recognition and spatial cleavage of dsRNAs (Zhang et al. 2004), both *OsDCL1b* and *OsDCL1c* are unlikely to be functionally redundant with *OsDCL1a*. Consistent with this, microarray analysis shows that *OsDCL1a* is ubiquitously expressed at all stages of the plant development with a peak of expression in the vegetative tissue, whereas *OsDCL1b* and *OsDCL1c* are only weakly expressed (Kapoor et al. 2008). However, *OsDCL1c* has a peak of expression in the panicle during the anthesis stage, suggesting a function distinct from *OsDCL1a*. It is also worth noticing that the silencing of *OsDCL4*, the closest homolog of *OsDCL1a* after *OsDCL1b* and *OsDCL1c*, does not affect the accumulation of any tested miRNAs (Liu et al. 2005). Taken together, these data strongly suggest that *OsDCL1a* is the primary dicer responsible for miRNA processing in rice.

Once processed, most of the mature miRNAs interact with *AtAGO1* in *Arabidopsis*. Rice has five *AtAGO1* homologs: *OsAGO1a*, *b*, *c*, and *d* (with *OsAGO1a*

and *OsAGO1b* being segmentally duplicated) and *OsAGO10* (also known as *PIN-HEAD*, *Os06g39640*) (Kapoor et al. 2008). All of them possess the conserved metal chelating residues in the PIWI domain that are critical for endonuclease activity, suggesting that all encode for a functional enzyme. The four *OsAGO1*s probably have some functional redundancy since they mostly interact with the same known miRNAs (Wu et al. 2009). On the other hand, their spatiotemporal expression does not necessarily overlap (Fujita et al. 2010; Jain et al. 2007; Kapoor et al. 2008, <http://www.plexdb.org/index.php>) and may explain why a subset of miRNAs were found to differentially interact with *OsAGO1a*, *b*, and *c*. Here are a few striking examples of the differential expression among the *OsAGO1* family. During vegetative development, *OsAGO1b* and *c* are preferentially expressed in the shoot rather than the root. Later on in the developing anther, the expression constantly drops for all but *OsAGO1a* which is significantly upregulated in mature pollen. *OsAGO1b* and *OsAGO1d* are preferentially expressed in the top part of the ovary, but dramatically decrease just after fertilization. Finally, during seed development, the expression of *OsAGO1a* and *OsAGO1c* drops, whereas *OsAGO1b* and *OsAGO1d* gradually increase. These data clearly suggest that despite their similarity, the four *OsAGO1*s are probably involved, at least partially, in different biological processes. It has also been found that some miRNA\*s accumulate in certain tissue more than their counterpart miRNA. miR390\* is present at higher level in the embryo than miR390, suggesting that miRNAs and miRNA\*s could function by association with different AGO1 family members (Xue et al. 2009). AGO10, which also belongs to the AGO1 family, has a miRNA-related function and is partially redundant with AGO1 in *Arabidopsis* (Mallory et al. 2009). Mutation of *AtAGO10* causes a critical alteration of the maintenance of the SAM (Lynn et al. 1999). Similarly in rice, *OsAGO10* is highly expressed in the SAM and is required for the maintenance of the indeterminate state of the SAM, suggesting a conserved function with *Arabidopsis* (Jain et al. 2007; Nishimura et al. 2002; <http://www.plexdb.org/index.php>). On the other hand, the proper development of the adaxial side of the leaves, which requires *AtAGO1* in *Arabidopsis*, seems to be dependent on *OsAGO10* in rice, suggesting that the specialization of AGO1 and AGO10 occurred after the split between monocot and dicot lineages (Kidner and Martienssen 2004; Nishimura et al. 2002; Yang et al. 2006b). Overall miRNA-mediated regulation in rice involves at least five AGO1 homologs, with potentially different tissue-specific functions.

Across species, the function of the miRNA pathway seems to be highly dependent on the *DCL1* and *AGO1* gene families. In the absence of AGO1, newly synthesized miRNAs are subject to degradation, suggesting that miRNA biogenesis and action need to be coupled in order for the miRNA to be functional. Interestingly, in tricellular mature pollen, *OsDCL1a* is not expressed, whereas *OsAGO1a* mRNA and two known miRNAs (miR166 and miR167) can be detected, suggesting that miRNAs previously processed and incorporated into miRISC can maintain their regulatory function in later developmental stages (Fujioka et al. 2008; Fujita et al. 2010; <http://www.plexdb.org/index.php>).

## 4.2 Regulation of miRNA Expression

While the overall control of the miRNA pathway depends on common enzymes, the specific function of a miRNA relies on its transcriptional regulation. Several mature miRNAs have been shown to accumulate preferentially in one or two tissues with a pattern that correlates with their associated function (Xue et al. 2009). The expression profile of a *MIR* gene provides valuable information in determining its role in plants. There is currently relatively little expression profile data for the rice miRNAs. However, the reduced cost and increased accessibility of next-generation high-throughput sequencing technology now allows generation of miRNA expression data at high sensitivity and resolution.

The *MIR* genes in rice are predominantly found in intergenic regions suggesting an autonomous control of their transcription, as opposed to pri-miRNA located in intronic regions that could be processed from the host gene transcript. Most of the *MIR* promoters possess *cis*-elements characteristic of RNA polymerase II activity (TATA-box, CAAT-box, and transcription start site), mainly located within 400 bp upstream of the pre-miRNA (Cui et al. 2009; Meng et al. 2009). The conserved *MIR* genes tend to contain more core promoter motifs than the nonconserved ones, suggesting that new pri-miRNAs gain their functionality by acquiring *cis*-regulatory elements in order to be specifically transcribed. Indeed some transcription factors, such as Auxin Response Factors (ARFs), may require the combination of a specific *cis*-element and the TATA-box in order to efficiently interact with the promoter (Inukai et al. 2005). In contrast, some *MIR* promoters are TATA-less; these can still be transcribed by RNA polymerase II but may belong to a different subclass of *MIR* genes expressed in more specific conditions (Bernard et al. 2010).

Regulation of rice miRNAs under abiotic stress conditions has been investigated by several research groups using different methods, including stem-loop RT-qPCR, mature-miRNA-specific RT-qPCR, miRNA microarray, Affymetrix GeneChip, and deep sequencing of small RNA libraries (Chen et al. 2009; Huang et al. 2009; Lv et al. 2010; Meng et al. 2009; Shen et al. 2010; Zhou et al. 2010). A large number of miRNAs, including most of the conserved miRNAs, have been shown to respond to abiotic stress. Six rice-specific *MIR* gene families have been identified in response to toxic exposure to the heavy metal cadmium ( $\text{Cd}^{2+}$ ) (Huang et al. 2009).

Under drought stress during vegetative and reproductive stages, Zhou et al. (2010) showed that after a few days of treatment, nine miRNA families found in the rice genome were downregulated and four were upregulated. Two miRNA families, miR172 and miR396, were found to be consistently repressed at both developmental stages, suggesting an important role in drought response. Interestingly, one member of the miR393 family showed an opposite expression at the reproductive stage between 5 and 6 days of treatment.

Using a rice-specific miRNA microarray chip, Lv et al. found 15 miRNA families affected by cold stress (Lv et al. 2010). Five of those miRNA families (miR156, 168, 169, 171, and 319) were also deregulated under drought stress, with

miR169 showing an opposite response between both stresses. In comparison, two other groups found additional miRNA families showing a slight deregulation within a few hours after drought, salt, cold, and abscisic acid (ABA) treatments (Shen et al. 2010; Zhao et al. 2007). A majority of those miRNAs were deregulated in at least two stresses and were sensitive to ABA treatment, suggesting that a common regulatory mechanism exists in response to different abiotic stresses in rice (Shen et al. 2010). Additionally, 5% of the miRNAs in rice (miRBase release 10) have been shown to be sensitive to auxin, and expression of seven miRNA families was affected in *osaxr*, an auxin-resistant mutant (Meng et al. 2009). Interestingly, miR528, which was the only *MIR* gene sensitive to both auxin treatment and the *osaxr* mutation, has not been linked to any other abiotic stress.

Analysis of the promoter region of stress-sensitive *MIR* genes revealed the presence of multiple stress or phytohormone regulatory *cis*-elements (Lv et al. 2010; Meng et al. 2009; Shen et al. 2010; Zhou et al. 2010). Among the more represented are *cis*-element linked with ABA response (“ABRE”ACGTG, CAAACACC, CACCTCCA, CACGTGGC), early response to dehydration (ACGT), MYB transcription factor (CTAACCA, CNGTTR), salt response (CANNTG), gibberellin response (“P-box” CCTTTT, TATC-box), salicylic acid response, auxin response (“AuxRE” TGTCTC), and endosperm expression (“Skn-1” TRTCAT, “CGN4” TGAGTCA). So far, none of these regulatory elements has been functionally validated, and the diversity and elevated number of sequences found in the promoters make it difficult for any confident interpretation. Still, we can observe that almost all *MIR* genes that are drought-sensitive have the Skn-1 *cis*-element in their promoter (Zhou et al. 2010). Also, AuxREs are found more frequently in the promoters of auxin-related *MIR* genes (Meng et al. 2009).

Interestingly, most stress-sensitive miRNAs show dynamic expression patterns (as opposed to a constitutive response) within the first hours following the treatment (Lv et al. 2010; Shen et al. 2010; Zhao et al. 2007). So depending on the experimental protocol, some stress-sensitive miRNAs may have escaped detection or shown different responses. For example, Shen et al. (2010) observed an upregulation of miR408 at 20 min after treatment, whereas Zhou et al. (2010) observed a downregulation at 14 days. Additionally, the ability to detect different miRNA homologs is important since miRNAs that belong to the same family can show different stress–response patterns, an example of this is the miR169 family (Zhao et al. 2007). Indeed, precursors that encode for the same mature miRNA can also be expressed differently under the same stress. A good example is the response of miR171 family members to cold stress (Lv et al. 2010). In rice, there are nine *MIR171* genes (a–i), which encode for five different mature miRNAs. Under cold treatment, pri-miR171a/e/f/i are downregulated, whereas pri-miR171c/d/h are upregulated. Pri-miR171c/d and e/f encode for the same miRNA and so potentially target the same genes, suggesting that those precursors have different tissue specificity. This example illustrates the complexity of miRNA-mediated regulation during abiotic stress responses and highlights the need to investigate them in different tissues and at multiple time points.

An efficient way of coordinating the regulation of specific molecular processes is through the simultaneous expression of clustered miRNAs, a dominant

characteristic of animal *MIR* genes. Interestingly, the rice genome contains 18–24 miRNA clusters (reported in two different studies), which represent roughly 20% of the total miRNAs (Cui et al. 2009; Merchan et al. 2009). This suggests that coordinated expression of miRNAs via a polycistronic transcript may be an important biosynthesis pathway in rice. Indeed, two miRNA clusters have already been validated, the first located in an intergenic region encodes for two miRNA homologs (miR159a.1 and a.2) and the second encodes for three intronic miRNAs (miR1876, miR1884b, and miR1862d) that are cotranscribed with the host gene (Lacombe et al. 2008; Lv et al. 2010; Zhu et al. 2008). Additionally, some miRNA clusters originating from a gene duplication event may have conserved regulatory expression (Meng et al. 2009). Further study of the *MIR* clusters, particularly those containing nonhomolog miRNAs (*MIR172c-820d*, *MIR172b-806a*, *MIR399d-439h*, and *MIR171f-443*), will certainly provide new insights to investigate the connections between miRNA-regulated networks in rice.

Overall, the dynamic expression patterns of miRNAs in response to abiotic stress, the presence of multiple phytohormone-related *cis*-elements in their promoters, and the evidence of direct regulation of phytohormone-related transcription factors by miRNAs (such as miR160/miR167-targeting ARF and miR159-targeting GAMyb) suggest complex interconnections between miRNA and phytohormone regulatory networks. In rice, the miRNAs are likely to be early components in the cascade of regulation that spreads and specifies hormone responses during a stress or a developmental event.

## 5 Biological Functions of Rice miRNAs

As detailed above, many miRNAs (both conserved and nonconserved) have target mRNAs that have been confirmed cleaved at the predicted target site. However, how the regulation of these target mRNAs by miRNAs contributes to biological processes has only been defined in a few cases. This is particularly the case for members of the Group II and III miRNAs where no information is available from *Arabidopsis* studies.

It is well established from *Arabidopsis* studies that miR156 regulates the *SPL* genes. Rice plants overexpressing miR156 are semidwarf, have altered numbers of leaves, and have longer vegetative phase (Xie et al. 2006). The downregulation of miR156 expression as plants progress from the juvenile to the adult phase agrees with observations across multiple species that this miRNA has a conserved role in regulating phase changes. In contrast, miR172 is upregulated during rice development (Zhu et al. 2009) also consistent with a role in regulating developmental phase changes. miR172 targets a small group of genes encoding *AP2-like* transcription factors. Overexpression of miR172b leads to a delay in the transition from spikelet to floral meristem, changing the numbers and identities of floral organs (Zhu et al. 2009). The miR172b overexpression lines also had enlarged lodicules which prevent closing of the spikelets after flowering. Loss of the miR172 target site in

a barley *AP2-like* gene, *Cly*, leads to cleistogamous (closed) flowering; these results suggest that a balance of miR172-mediated repression is important for controlling the normal noncleistogamous phenotype (Zhu and Helliwell 2011).

A major target for plant breeders is improvement of plant architecture to optimize grain yields. The map-based cloning and characterization of a semidominant QTL (*IPA1*) revealed that it contains the miR156-targeted *SPL14* gene (Jiao et al. 2010). The basis of the QTL appears to be a mutation of the miR156 target site in the *SPL14* gene. This leads to increased abundance of the SPL14 protein. As a consequence, the number of tillers is reduced, and grain yield increased compared to wild type. This result suggests that identifying alleles with altered miRNA target sites may be a fruitful method of identifying useful phenotypic variation.

Recently, it was demonstrated that miR396 controls cell proliferation in *Arabidopsis* (Rodriguez et al. 2009). Rice plants overexpressing miR396c have reduced tolerance to salt and alkali stress (Gao et al. 2010). miR396 targets GRFs (growth regulating factors) that are important proteins in plant growth and development. Overexpression of rice miR396c caused the downregulation of different GRFs, which may confer the increased sensitivity of the plants to salt and alkali stress. It has also been reported that members of the highly conserved miR393 family have important roles in response to development and stress (Arenas-Huertero et al. 2009; Liu et al. 2008; Sunkar and Zhu 2004). In rice, miR393 expression shows a trend to decreased expression in response to either high salt or alkali treatments, while expression of the related miR393b is relatively constant (Gao et al. 2011). Rice plants overexpressing miR393 showed severe symptoms than wild type when treated with 150 mM of NaCl. Bioinformatic predictions suggest that miR393 targets rice genes that code for stress-related proteins (Gao et al. 2011), and thus, the reduction of miR393 under salt stress conditions may allow expression of stress-response proteins to increase.

## 6 Other miRNA-Like siRNAs in Rice

In addition to the miRNAs discussed above, a number of other siRNAs have been identified with miRNA-like properties but which differ in aspects of their biogenesis. One example is a putative mirtron identified from deep sequencing of grain small RNAs (Zhu et al. 2008). Mirtrons are miRNAs processed from spliced and debranched introns; they were originally defined in animal systems where intron splicing essentially replaces the need for processing of the pri-miRNA hairpin structure by Droscha (Ruby et al. 2007). In the case of the rice mirtron (miR1429.2), small RNA reads align exactly to each end of the intron of a putative homeobox gene which is predicted to form a miRNA-like stem-loop once debranched. miR1429.2 is predominantly expressed in developing grains, but does not have a demonstrated target gene, so whether this miRNA has a biological role is unclear.

A large number of phased siRNAs have been identified in rice; most of these appear to be generated by a similar mechanism to ta-siRNAs and are expressed specifically in the developing inflorescence (Johnson et al. 2009). Two examples of long hairpin structures that are transcribed and processed to phased 21-nt siRNAs have been identified in rice (Zhu et al. 2008). Among the small RNAs produced from these loci two have been shown to cleave target mRNA transcripts by 5' RACE, indicating that these siRNAs can act to cleave target transcripts in a miRNA-like manner. At present, the biological significance of these miRNA-like small RNAs is unclear.

## 7 Conclusion

In conclusion, it is clear that as for other species, the rice genome encodes many miRNAs that potentially target a wide range of biological processes. To date, our understanding of the function of these miRNAs is largely limited to those that are conserved across species. The many rice-specific miRNAs tend to be present at low abundance, so many have very subtle or cell-specific roles that will require careful investigation.

## References

- Arazi T, Talmor-Neiman M, Stav R, Riese M, Huijser P, Baulcombe DC (2005) Cloning and characterization of micro-RNAs from moss. *Plant J* 43:837–848
- Arenas-Huertero C, Perez B, Rabanal F, Blanco-Melo D, De la Rosa C, Estrada-Navarrete G, Sanchez F, Covarrubias AA, Reyes JL (2009) Conserved and novel miRNAs in the legume *Phaseolus vulgaris* in response to stress. *Plant Mol Biol* 70:385–401
- Bernard V, Brunaud V, Lecharny A (2010) TC-motifs at the TATA-box expected position in plant genes: a novel class of motifs involved in the transcription regulation. *BMC Genomics* 11:166
- Bonnet E, Wuyts J, Rouze P, Van de Peer Y (2004) Detection of 91 potential in plant conserved plant microRNAs in *Arabidopsis thaliana* and *Oryza sativa* identifies important target genes. *Proc Natl Acad Sci USA* 101:11511–11516
- Bouche N, Lauressergues D, Gascolli V, Vaucheret H (2006) An antagonistic function for Arabidopsis DCL2 in development and a new function for DCL4 in generating viral siRNAs. *EMBO J* 25:3347–3356
- Chen SY, Li XQ, Zhao AG, Wang LJ, Li XF, Shi QY, Chen M, Guo J, Zhang JC, Qi DM, Liu GS (2009) Genes and pathways induced in early response to defoliation in rice seedlings. *Curr Issues Mol Biol* 11:81–100
- Chen X (2009) Small RNAs and their roles in plant development. *Annu Rev Cell Dev Biol* 25:21–44
- Chuck G, Cigan AM, Saeteurn K, Hake S (2007) The heterochronic maize mutant Corngrass1 results from overexpression of a tandem microRNA. *Nat Genet* 39:544–549
- Cui X, Xu SM, Mu DS, Yang ZM (2009) Genomic analysis of rice microRNA promoters and clusters. *Gene* 431:61–66



- Fujioka T, Kaneko F, Kazarna T, Suwabe K, Suzuki G, Makino A, Mae T, Endo M, Kawagishi-Kobayashi M, Watanabe M (2008) Identification of small RNAs in late developmental stage of rice anthers. *Genes Genet Syst* 83:281–284
- Fujita M, Horiuchi Y, Ueda Y, Mizuta Y, Kubo T, Yano K, Yamaki S, Tsuda K, Nagata T, Niihama M, Kato H, Kikuchi S, Hamada K, Mochizuki T, Ishimizu T, Iwai H, Tsutsumi N, Kurata N (2010) Rice expression atlas in reproductive development. *Plant Cell Physiol* 51:2060–2081
- Gao P, Bai X, Yang L, Lv DK, Li Y, Cai H, Ji W, Guo DJ, Zhu YM (2010) Over-expression of osa-MIR396c decreases salt and alkali stress tolerance. *Planta* 231:991–1001
- Gao P, Bai X, Yang LA, Lv DK, Pan X, Li Y, Cai H, Ji W, Chen Q, Zhu YM (2011) osa-MIR393: a salinity- and alkaline stress-related microRNA gene. *Mol Biol Rep* 38:237–242
- German MA, Pillay M, Jeong DH, Hetawal A, Luo S, Janardhanan P, Kannan V, Rymarquis LA, Nobuta K, German R, De Paoli E, Lu C, Schroth G, Meyers BC, Green PJ (2008) Global identification of microRNA-target RNA pairs by parallel analysis of RNA ends. *Nat Biotechnol* 26:941–946
- Goff SA, Ricke D, Lan TH, Presting G, Wang R, Dunn M, Glazebrook J, Sessions A, Oeller P, Varma H, Hadley D, Hutchison D, Martin C, Katagiri F, Lange BM, Moughamer T, Xia Y, Budworth P, Zhong J, Miguel T, Paszkowski U, Zhang S, Colbert M, Sun WL, Chen L, Cooper B, Park S, Wood TC, Mao L, Quail P, Wing R, Dean R, Yu Y, Zharkikh A, Shen R, Sahasrabudhe S, Thomas A, Cannings R, Gutin A, Pruss D, Reid J, Tavtigian S, Mitchell J, Eldredge G, Scholl T, Miller RM, Bhatnagar S, Adey N, Rubano T, Tusneem N, Robinson R, Feldhaus J, Macalma T, Oliphant A, Briggs S (2002) A draft sequence of the rice genome (*Oryza sativa* L. ssp. japonica). *Science* 296:92–100
- Heisel SE, Zhang YJ, Allen E, Guo L, Reynolds TL, Yang X, Kovalic D, Roberts JK (2008) Characterization of unique small RNA populations from rice grain. *PLoS One* 3:e2871
- Huang S, Peng J, Qiu C, Yang Z (2009) Heavy metal-regulated new microRNAs from rice. *J Inorg Biochem* 103:282–287
- Inukai Y, Sakamoto T, Ueguchi-Tanaka M, Shibata Y, Gomi K, Umemura I, Hasegawa Y, Ashikari M, Kitano H, Matsuoka M (2005) Crown rootless1, which is essential for crown root formation in rice, is a target of an AUXIN RESPONSE FACTOR in auxin signaling. *Plant Cell* 17:1387–1396
- Itoh J, Hibara K, Sato Y, Nagato Y (2008) Developmental role and auxin responsiveness of Class III homeodomain leucine zipper gene family members in rice. *Plant Physiol* 147:1960–1975
- Jain M, Nijhawan A, Arora R, Agarwal P, Ray S, Sharma P, Kapoor S, Tyagi AK, Khurana JP (2007) F-box proteins in rice. Genome-wide analysis, classification, temporal and spatial gene expression during panicle and seed development, and regulation by light and abiotic stress. *Plant Physiol* 143:1467–1483
- Jian X, Zhang L, Li G, Wang X, Cao X, Fang X, Chen F (2010) Identification of novel stress-regulated microRNAs from *Oryza sativa* L. *Genomics* 95:47–55
- Jiao YQ, Wang YH, Xue DW, Wang J, Yan MX, Liu GF, Dong GJ, Zeng DL, Lu ZF, Zhu XD, Qian Q, Li JY (2010) Regulation of OsSPL14 by OsmiR156 defines ideal plant architecture in rice. *Nat Genet* 42:541–536
- Johnson C, Kasprzewska A, Tennessen K, Fernandes J, Nan GL, Walbot V, Sundaresan V, Vance V, Bowman LH (2009) Clusters and superclusters of phased small RNAs in the developing inflorescence of rice. *Genome Res* 19:1429–1440
- Jones-Rhoades MW, Bartel DP (2004) Computational identification of plant MicroRNAs and their targets, including a stress-induced miRNA. *Mol Cell* 14:787–799
- Jones-Rhoades MW, Bartel DP, Bartel B (2006) MicroRNAs and their regulatory roles in plants. *Annu Rev Plant Biol* 57:19–53
- Juarez MT, Kui JS, Thomas J, Heller BA, Timmermans MC (2004) microRNA-mediated repression of rolled leaf1 specifies maize leaf polarity. *Nature* 428:84–88
- Kapoor M, Arora R, Lama T, Nijhawan A, Khurana JP, Tyagi AK, Kapoor S (2008) Genome-wide identification, organization and phylogenetic analysis of dicer-like, Argonaute and RNA-



- dependent RNA polymerase gene families and their expression analysis during reproductive development and stress in rice. *BMC Genomics* 9:451
- Kidner CA, Martienssen RA (2004) Spatially restricted microRNA directs leaf polarity through ARGONAUTE1. *Nature* 428:81–84
- Kurihara Y, Watanabe Y (2004) Arabidopsis micro-RNA biogenesis through Dicer-like 1 protein functions. *Proc Natl Acad Sci USA* 101:12753–12758
- Lacombe S, Nagasaki H, Santi C, Duval D, Piegu B, Bangratz M, Breitler JC, Guiderdoni E, Brugidou C, Hirsch J, Cao X, Brice C, Panaud O, Karlowski WM, Sato Y, Echeverria M (2008) Identification of precursor transcripts for 6 novel miRNAs expands the diversity on the genomic organisation and expression of miRNA genes in rice. *BMC Plant Biol* 8:123
- Li T, Li H, Zhang YX, Liu JY (2010a) Identification and analysis of seven H<sub>2</sub>O<sub>2</sub>-responsive miRNAs and 32 new miRNAs in the seedlings of rice (*Oryza sativa* L. ssp. indica). *Nucleic Acids Res* 39:2821–2833
- Li YF, Zheng Y, Addo-Quaye C, Zhang L, Saini A, Jagadeeswaran G, Axtell MJ, Zhang WX, Sunkar R (2010b) Transcriptome-wide identification of microRNA targets in rice. *Plant J* 62:742–759
- Liu B, Li PC, Li X, Liu CY, Cao SY, Chu CC, Cao XF (2005) Loss of function of OsDCL1 affects microRNA accumulation and causes developmental defects in rice. *Plant Physiol* 139:296–305
- Liu HH, Tian X, Li YJ, Wu CA, Zheng CC (2008) Microarray-based analysis of stress-regulated microRNAs in *Arabidopsis thaliana*. *RNA* 14:836–843
- Liu Q, Zhang CY, Wang CY, Luo YC, Huang QJ, Chen SY, Zhou H, Qu LH, Chen YQ (2009) Expression analysis of phytohormone-regulated microRNAs in rice, implying their regulation roles in plant hormone signaling. *FEBS Lett* 583:723–728
- Lui WO, Pourmand N, Patterson BK, Fire A (2007) Patterns of known and novel small RNAs in human cervical cancer. *Cancer Res* 67:6031–6043
- Lu C, Jeong DH, Kulkarni K, Pillay M, Nobuta K, German R, Thatcher SR, Maher C, Zhang L, Ware D, Liu B, Cao X, Meyers BC, Green PJ (2008) Genome-wide analysis for discovery of rice microRNAs reveals natural antisense microRNAs (nat-miRNAs). *Proc Natl Acad Sci USA* 105:4951–4956
- Luo YC, Zhou H, Li Y, Chen JY, Yang JH, Chen YQ, Qu LH (2006) Rice embryogenic calli express a unique set of microRNAs, suggesting regulatory roles of microRNAs in plant post-embryogenic development. *FEBS Lett* 580:5111–5116
- Lv DK, Bai X, Li Y, Ding XD, Ge Y, Cai H, Ji W, Wu NLH, Zhu YM (2010) Profiling of cold-stress-responsive miRNAs in rice by microarrays. *Gene* 459:39–47
- Lynn K, Fernandez A, Aida M, Sedbrook J, Tasaka M, Masson P, Barton MK (1999) The PINHEAD/ZWILLE gene acts pleiotropically in Arabidopsis development and has overlapping functions with the ARGONAUTE1 gene. *Development* 126:469–481
- Mallory AC, Hinze A, Tucker MR, Bouche N, Gascioli V, Elmayan T, Lauressergues D, Jauvion V, Vaucheret H, Laux T (2009) Redundant and specific roles of the ARGONAUTE proteins AGO1 and ZLL in development and small RNA-directed gene silencing. *PLoS Genet* 5:e1000646
- Margis R, Fusaro AF, Smith NA, Curtin SJ, Watson JM, Finnegan EJ, Waterhouse PM (2006) The evolution and diversification of Dicers in plants. *FEBS Lett* 580:2442–2450
- McConnell JR, Emery J, Eshed Y, Bao N, Bowman J, Barton MK (2001) Role of PHABULOSA and PHAVOLUTA in determining radial patterning in shoots. *Nature* 411:709–713
- McElver J, Tzafrir I, Aux G, Rogers R, Ashby C, Smith K, Thomas C, Schetter A, Zhou Q, Cushman MA, Tossberg J, Nickle T, Levin JZ, Law M, Meinke D, Patton D (2001) Insertional mutagenesis of genes required for seed development in *Arabidopsis thaliana*. *Genetics* 159:1751–1763
- Meng Y, Huang F, Shi Q, Cao J, Chen D, Zhang J, Ni J, Wu P, Chen M (2009) Genome-wide survey of rice microRNAs and microRNA-target pairs in the root of a novel auxin-resistant mutant. *Planta* 230:883–898

- Merchan F, Boualem A, Crespi M, Frugier F (2009) Plant polycistronic precursors containing non-homologous microRNAs target transcripts encoding functionally related proteins. *Genome Biol* 10:R136
- Miura K, Ikeda M, Matsubara A, Song XJ, Ito M, Asano K, Matsuoka M, Kitano H, Ashikari M (2010) OsSPL14 promotes panicle branching and higher grain productivity in rice. *Nat Genet* 42:545–549
- Morin RD, Aksay G, Dolgosheina E, Ebhardt HA, Magrini V, Mardis ER, Sahinalp SC, Unrau PJ (2008) Comparative analysis of the small RNA transcriptomes of *Pinus contorta* and *Oryza sativa*. *Genome Res* 18:571–584
- Nagasaki H, Itoh J, Hayashi K, Hibara K, Satoh-Nagasawa N, Nosaka M, Mukouhata M, Ashikari M, Kitano H, Matsuoka M, Nagato Y, Sato Y (2007) The small interfering RNA production pathway is required for shoot meristem initiation in rice. *Proc Natl Acad Sci USA* 104:14867–14871
- Nishimura A, Ito M, Kamiya N, Sato Y, Matsuoka M (2002) OsPNH1 regulates leaf development and maintenance of the shoot apical meristem in rice. *Plant J* 30:189–201
- Nogueira FT, Madi S, Chitwood DH, Juarez MT, Timmermans MC (2007) Two small regulatory RNAs establish opposing fates of a developmental axis. *Genes Dev* 21:750–755
- Park W, Li JJ, Song RT, Messing J, Chen XM (2002) CARPEL FACTORY, a Dicer homolog, and HEN1, a novel protein, act in microRNA metabolism in *Arabidopsis thaliana*. *Curr Biol* 12:1484–1495
- Peng M, Hannam C, Gu H, Bi YM, Rothstein SJ (2007) A mutation in NLA, which encodes a RING-type ubiquitin ligase, disrupts the adaptability of Arabidopsis to nitrogen limitation. *Plant J* 50:320–337
- Pignocchi C, Kiddle G, Hernandez I, Foster SJ, Asensi A, Taybi T, Barnes J, Foyer CH (2006) Ascorbate oxidase-dependent changes in the redox state of the apoplast modulate gene transcript accumulation leading to modified hormone signaling and orchestration of defense processes in tobacco. *Plant Physiol* 141:423–435
- Potters G, Horemans N, Caubergs RJ, Asard H (2000) Ascorbate and dehydroascorbate influence cell cycle progression in a tobacco cell suspension. *Plant Physiol* 124:17–20
- Reinhart BJ, Weinstein EG, Rhoades MW, Bartel B, Bartel DP (2002) MicroRNAs in plants. *Genes Dev* 16:1616–1626
- Rodriguez RE, Mecchia MA, Debernardi JM, Schommer C, Weigel D, Palatnik JF (2009) Control of cell proliferation in *Arabidopsis thaliana* by microRNA miR396. *Development* 137:103–112
- Ruby JG, Jan CH, Bartel DP (2007) Intronic microRNA precursors that bypass Drosha processing. *Nature* 448:83–86
- Sanan-Mishra N, Kumar V, Sopory SK, Mukherjee SK (2009) Cloning and validation of novel miRNA from basmati rice indicates cross talk between abiotic and biotic stresses. *Mol Genet Genomics* 282:463–474
- Schauer SE, Jacobsen SE, Meinke DW, Ray A (2002) DICER-LIKE1: blind men and elephants in Arabidopsis development. *Trends Plant Sci* 7:487–491
- Schreiber AW, Shi BJ, Huang CY, Langridge P, Baumann U (2011) Discovery of barley miRNAs through deep sequencing of short reads. *BMC Genomics* 12:129
- Shen JQ, Xie KB, Xiong LZ (2010) Global expression profiling of rice microRNAs by one-tube stem-loop reverse transcription quantitative PCR revealed important roles of microRNAs in abiotic stress responses. *Mol Genet Genomics* 284:477–488
- Sunkar R, Girke T, Jain PK, Zhu JK (2005) Cloning and characterization of microRNAs from rice. *Plant Cell* 17:1397–1411
- Sunkar R, Zhou XF, Zheng Y, Zhang WX, Zhu JK (2008) Identification of novel and candidate miRNAs in rice by high throughput sequencing. *BMC Plant Biol* 8:25
- Sunkar R, Zhu JK (2004) Novel and stress-regulated microRNAs and other small RNAs from Arabidopsis. *Plant Cell* 16:2001–2019

- Tsuji H, Aya K, Ueguchi-Tanaka M, Shimada Y, Nakazono M, Watanabe R, Nishizawa NK, Gomi K, Shimada A, Kitano H, Ashikari M, Matsuoka M (2006) GAMYB controls different sets of genes and is differentially regulated by microRNA in aleurone cells and anthers. *Plant J* 47:427–444
- Vaucheret H, Vazquez F, Crete P, Bartel DP (2004) The action of ARGONAUTE1 in the miRNA pathway and its regulation by the miRNA pathway are crucial for plant development. *Genes Dev* 18:1187–1197
- Wang JF, Zhou H, Chen YQ, Luo QJ, Qu LH (2004a) Identification of 20 microRNAs from *Oryza sativa*. *Nucleic Acids Res* 32:1688–1695
- Wang XJ, Reyes JL, Chua NH, Gaasterland T (2004b) Prediction and identification of *Arabidopsis thaliana* microRNAs and their mRNA targets. *Genome Biol* 5:R65
- Wang Y, Ribot C, Rezzonico E, Poirier Y (2004c) Structure and expression profile of the *Arabidopsis* PHO1 gene family indicates a broad role in inorganic phosphate homeostasis. *Plant Physiol* 135:400–411
- Wu L, Zhang QQ, Zhou HY, Ni FR, Wu XY, Qi YJ (2009) Rice microRNA effector complexes and targets. *Plant Cell* 21:3421–3435
- Wu L, Zhou H, Zhang Q, Zhang J, Ni F, Liu C, Qi Y (2010) DNA methylation mediated by a microRNA pathway. *Mol Cell* 38:465–475
- Xie K, Wu C, Xiong L (2006) Genomic organization, differential expression, and interaction of SQUAMOSA promoter-binding-like transcription factors and microRNA156 in rice. *Plant Physiol* 142:280–293
- Xue LJ, Zhang JJ, Xue HW (2009) Characterization and expression profiles of miRNAs in rice seeds. *Nucleic Acids Res* 37:916–930
- Yakovlev IA, Fossdal CG, Johnsen O (2010) MicroRNAs, the epigenetic memory and climatic adaptation in Norway spruce. *New Phytol* 187:1154–1169
- Yang JH, Han SJ, Yoon EK, Lee WS (2006a) Evidence of an auxin signal pathway, microRNA167-ARF8-GH3, and its response to exogenous auxin in cultured rice cells. *Nucleic Acids Res* 34:1892–1899
- Yang L, Conway SR, Poethig RS (2010) Vegetative phase change is mediated by a leaf-derived signal that represses the transcription of miR156. *Development* 138:245–249
- Yang L, Huang W, Wang H, Cai R, Xu Y, Huang H (2006b) Characterizations of a hypomorphic argonaute1 mutant reveal novel AGO1 functions in *Arabidopsis* lateral organ development. *Plant Mol Biol* 61:63–78
- Yu J, Hu S, Wang J, Wong GK, Li S, Liu B, Deng Y, Dai L, Zhou Y, Zhang X, Cao M, Liu J, Sun J, Tang J, Chen Y, Huang X, Lin W, Ye C, Tong W, Cong L, Geng J, Han Y, Li L, Li W, Hu G, Li J, Liu Z, Qi Q, Li T, Wang X, Lu H, Wu T, Zhu M, Ni P, Han H, Dong W, Ren X, Feng X, Cui P, Li X, Wang H, Xu X, Zhai W, Xu Z, Zhang J, He S, Xu J, Zhang K, Zheng X, Dong J, Zeng W, Tao L, Ye J, Tan J, Chen X, He J, Liu D, Tian W, Tian C, Xia H, Bao Q, Li G, Gao H, Cao T, Zhao W, Li P, Chen W, Zhang Y, Hu J, Liu S, Yang J, Zhang G, Xiong Y, Li Z, Mao L, Zhou C, Zhu Z, Chen R, Hao B, Zheng W, Chen S, Guo W, Tao M, Zhu L, Yuan L, Yang H (2002) A draft sequence of the rice genome (*Oryza sativa* L. ssp. indica). *Science* 296:79–92
- Zanca AS, Vicentini R, Ortiz-Moreira FA, Del Bem LE, da Silva MJ, Vincenz M, Nogueira FT (2010) Identification and expression analysis of microRNAs and targets in the biofuel crop sugarcane. *BMC Plant Biol* 10:260
- Zhang H, Kolb FA, Jaskiewicz L, Westhof E, Filipowicz W (2004) Single processing center models for human Dicer and bacterial RNase III. *Cell* 118:57–68
- Zhao B, Ge L, Liang R, Li W, Ruan K, Lin H, Jin Y (2009) Members of miR-169 family are induced by high salinity and transiently inhibit the NF-YA transcription factor. *BMC Mol Biol* 10:8
- Zhao BT, Liang RQ, Ge LF, Li W, Xiao HS, Lin HX, Ruan KC, Jin YX (2007) Identification of drought-induced microRNAs in rice. *Biochem Biophys Res Commun* 354:585–590
- Zhou LG, Liu YH, Liu ZC, Kong DY, Duan M, Luo LJ (2010) Genome-wide identification and analysis of drought-responsive microRNAs in *Oryza sativa*. *J Exp Bot* 61:4157–4168

- Zhu QH, Spriggs A, Matthew L, Fan L, Kennedy G, Gubler F, Helliwell CA (2008) A diverse set of microRNAs and microRNA-like small RNAs in developing rice grains. *Genome Res* 18:1456–1465
- Zhu QH, Upadhyaya NM, Gubler F, Helliwell CA (2009) Over-expression of miR172 causes loss of spikelet determinacy and floral organ abnormalities in rice (*Oryza sativa*). *BMC Plant Biol* 9:17
- Zhu QH, Helliwell CA (2011) Regulation of flowering time and floral patterning by miR172. *J Exp Bot* 62:487–495