

Expression variations of miRNAs and mRNAs in rice (*Oryza sativa*)

Ming Wen^{1§}, Munan Xie¹, Lian He^{1§}, Yushuai Wang¹, Suhua Shi¹, Tian Tang^{1*}

¹ State Key Laboratory of Biocontrol, Guangdong Key Laboratory of plant Resources and Key Laboratory of Biodiversity Dynamics and Conservation of Guangdong Higher Education Institutes, School of Life Sciences, Sun Yat-sen University, Guangzhou 510275, Guangdong, China

*Corresponding author: lsstt@mail.sysu.edu.cn, +86-020-84110750

Ming Wen: wenming126@gmail.com

Munan Xie: meximus@126.com

Lian He: lhe@ibt.tamhsc.edu

Yushuai Wang: buyejue2012@163.com

Suhua Shi: lssssh@mail.sysu.edu.cn

§Present Address: Ming Wen, Room 029, 6th floors, The Academia, 20 College Road, Singapore, 169856. Lian He, Institute of Biosciences and Technology, Texas Agriculture and Mechanics University, Houston, Texas 77030, U.S.A.

Abstract

Differences in expression levels are an important source of phenotypic variation within and between populations. MicroRNAs (miRNAs) are key players in post-transcriptional gene regulation that are important for plant development and stress responses. We surveyed expression variation of miRNAs and mRNAs of six accessions from two rice subspecies *Oryza sativa* L. ssp. *indica* and *Oryza sativa* L. ssp. *japonica* using deep sequencing. While more than half (53.7%) of the mature miRNAs exhibit differential expression between grains and seedlings of rice, only 11.0% show expression differences between subspecies, with an additional 2.2% differentiated for the development-by-subspecies interaction. Expression variation is greater for lowly conserved miRNAs than highly conserved miRNAs, whereas the latter show stronger negative correlation with their targets in expression changes between subspecies. Using a permutation test, we identified 51 miRNA-mRNA pairs that correlate negatively or positively in expression level among cultivated rice. Genes involved in various metabolic processes and stress responses are enriched in the differentially expressed genes between rice *indica* and *japonica* subspecies. Our results indicate that stabilizing selection is the major force governing miRNA expression in cultivated rice, albeit positive selection may be responsible for much of the between-subspecies expression divergence.

Keywords: Expression variation; microRNA; mRNA; Rice

Introduction

MicroRNAs (miRNAs) are short (~ 21 nt) non-coding RNAs that regulate target transcripts post-transcriptionally via translational repression or mRNA degradation (Jones-Rhoades et al. 2006, Filipowicz et al. 2008, Bartel 2009). Mature miRNAs are incorporated into RNA-induced silencing complexes (RISCs) and guide RISCs to specific targets through Watson-Crick base-pairing (Jones-Rhoades et al. 2006, Filipowicz et al. 2008, Bartel 2009). A large proportion of miRNAs that were initially discovered in plants regulate transcription factors (Zhang et al. 2006); some of them are highly conserved miRNAs and play important roles in plant development (Carrington et al. 2003, Chuck et al. 2009, Carlsbecker et al. 2010), stress adaptation (Lv et al. 2010, Ding et al. 2011) and hormone signaling (Liu et al. 2009). Sequence evolution of miRNAs and their target sites have been extensively explored between closely related plant species (Fahlgren et al. 2010, Ma et al. 2010). For example, rapid sequence evolution of the miR482/2118 gene family has promoted the evolution of resistance genes in the Solanaceae (de Vries et al. 2015). However, little is known about how expression variation in miRNAs may affect variation in gene expression among individuals, which is an important source of phenotypic variation within species (Britten et al. 1971, King et al. 1975).

Rice is one of the most important crops in the world. The two rice subspecies, *Oryza sativa* L. ssp. *indica* (*indica*) and *Oryza sativa* L. ssp. *japonica* (*japonica*), show significant phenotypic and genetic differentiations (He et al. 2011). To date, 592 precursor miRNAs, representing 334 miRNA families, have been identified in rice (miRBase, v20.0). MiRNAs play a significant role in regulating rice development and stress responses (Campo et al. 2013). For example, regulation of *OsSPL14* by OsmiR156 defines ideal plant architecture in rice (Jiao et al. 2010, Luo et al. 2012) and OsmiR397

substantially enhance grain yield in rice through the repression of *OsLAC* (Zhang et al. 2013). Moreover, differential expression of miRNAs was reported among *japonica* cv. Nipponbare and *indica* cv. 93-11, probably playing roles in heterosis (Chen et al. 2010). However, variation in miRNA expression among rice accessions and its effects on target and global gene expression remain largely unknown.

Here, we surveyed the expression variation in miRNAs and mRNAs among six rice accessions (three from each of *indica* and *japonica* subspecies) using RNA-seq. In contrast to extensive differential miRNA expression between developmental stages, only a few miRNAs showed expression differences between *indica* and *japonica* subspecies of rice. Highly conserved miRNAs exhibited less expression variation but repressed target gene expression more strongly than lowly conserved miRNAs. Our analysis also identified significantly correlated miRNA-mRNA pairs that differed in expression levels among *indica* and *japonica* cultivars, which may be candidates for adaptive regulatory evolution underlying *indica-japonica* differentiation in cultivated rice.

Results

Small RNA data processing and expression analysis of rice miRNAs

Small RNA (sRNA) libraries of rice grains and seedlings were constructed and sequenced for three *indica* cultivars Khal Dawk Mali 105, Guangluai 4 and Rathuwee, and three *japonica* cultivars Taipei 309, Heukgyeong and Nipponbare individually. About 9-14 million short reads were obtained for each library, representing an average of 14,849,295 unique reads for *indica* grain library, 12,978,406 for *japonica* grain library, 9,924,065 for *indica* seedling library and 9,395,033 for *japonica* seedling library (Table S1). About 94% of the sRNAs were 20-24 nucleotides (nts) in length, with 21 and 24nt as the two major size classes (Figure S1). In plants, most miRNAs are 21-nt in length while 24-nt sRNAs consist mainly of sRNAs that are associated with repeats and transposable

elements (TEs) (Axtell 2013). These results indicate that rice has a highly complex sRNA population to which repetitive sequences are the major contributors (Zhu et al. 2008). Small reads matching rice plastid DNA, structural non-coding RNAs (ncRNAs) and repetitive sequences were removed before miRNA annotation. On average, only a small portion of the total reads were mapped to the chloroplast (6.64%) or mitochondria (1.71%) genome, whereas short reads matching ncRNAs and repetitive sequences accounted for 49.3% and 14% of the total reads, respectively. While the sRNAs belonging to the above categories showed an uniform distribution among all 12 libraries, the proportion of signatures that matched miRNA precursors were consistently higher in seedlings (31.3%) than in grain (3.8%), consistent with a previous report (Xue et al. 2009).

To date, 592 rice miRNAs representing 334 families have been registered in the miRBase database (v20.0). Among them, 22 miRNA families were found conserved in both monocots and eudicots previously (Cuperus et al. 2011). We denoted these miRNA families as highly conserved miRNAs and the others as lowly conserved miRNAs in this study. MiRNAs were considered to be expressed by requiring at least ten raw reads in all three accessions from the same developmental stage and the same subspecies. As a result, a total of 272 mature miRNAs were expressed among the 12 libraries, comprising 111 highly conserved miRNAs and 161 lowly conserved miRNAs (Table 2). The expression level of each mature miRNA was measured as Reads Per Million (RPM). A pairwise comparison of $\log_2(\text{RPM})$ of all miRNAs across the 12 libraries demonstrated the measurement of miRNA expression is highly reproducible between biological replicates in our data (Pearson's correlation, $r = 0.86$ to 0.97 ; Figure S2). In addition, we also predicted 20 novel miRNAs using the criteria for plant miRNA annotation (Meyers et al. 2008). Most of these novel miRNAs exhibited developmental stage- or subspecies-specific expression and were predicted to regulate 12 target genes (Table S3).

Highly conserved miRNAs exhibit more stable expression than the lowly conserved miRNAs

We first explored the miRNA expression variation across the 12 samples by hierarchical clustering (Figure 1A). Dendrogram classification grouped the samples first by subspecies (*indica-japonica*) and then by developmental stage (grain-seedling), indicating that rice miRNA expression, which is largely remodeled between the different developmental stages, exhibits a relatively small difference between the two subspecies. The mature miRNAs were clustered into several clades (Figure 1A). MiRNAs from the basal clades showed high expression levels with similar expression profiles across rice cultivars from different developmental stages and subspecies; most of them are highly conserved miRNAs, including miR156, miR164, miR166, miR167 and miR168, and two are lowly conserved miR535 and miR5794. In contrast, the majority of miRNAs showed grain- or seedling-biased expression between developmental stages. A number of miRNA families, including miR1861, miR1868, miR319, as well as miR444, exhibited higher expression in grains, whereas other miRNAs, such as members of the miR160, miR169, miR172 and miR529 families, mainly showed increased expression in seedlings. These stage-biased miRNAs are promising candidates for developmental regulation in rice (Figure 1A and Table S2).

Using a Log-linear Poisson Regression model (see Methods), 146 (53.7%), 30 (11.0%) and 6 (2.2%) of the 272 expressed mature miRNAs exhibited significantly differential expression for the factors developmental stage, subspecies and development-by-subspecies interaction (fold change ≥ 2 , FDR ≤ 0.05), respectively (Table 2). These miRNAs were considered to play important roles in regulating rice development or subspecies differentiation. For example, miR159a-f showed a

grain-biased expression (Table S2) and is involved in ABA-mediated responses in the hormonal and abiotic stress signaling networks (Reyes et al. 2007); members of miR156, miR166, miR169, miR171, and miR444 families showed a leaf-biased expression (Table S2) and are known to play defensive roles against stress by targeting transcription factors (Macovei et al. 2012).

When grouping the mature miRNAs according to their conservation, expression variation was less prominent in highly conserved miRNAs (Figure 1B) than lowly conserved miRNAs (Figure 1C). Only 1.8% (2/111) of highly conserved miRNAs were differentially expressed between subspecies while the proportion was 17.4% (28/161) for lowly conserved miRNAs. An additional 50.9% (82/161) and 3.7% (6/161) of lowly conserved miRNAs were also differentially expressed between developmental stages or development-by-subspecies interaction (Table 2 and Figure 2A). Furthermore, magnitudes of the expression fold changes are much higher for lowly conserved miRNAs than highly conserved miRNAs (Figure 1B vs. C and Figure 2B). These results strongly suggest that variation in expression is higher for the less conserved miRNAs than for the more conserved miRNAs found in both monocots and dicots. A possible explanation for this result is that conserved miRNAs are processed better from their foldback structures (Shen et al. 2011). Another explanation could also be that conserved miRNAs are under strong functional constraints (Cuperus et al. 2011). Using qRT-PCR, we validated the differential expression between subspecies for eight out of nine lowly conserved miRNAs in rice seedlings but not for the highly conserved miR166j-5p due to its low expression (Figure S3A and C).

Differential expression of transcriptome between *indica* and *japonica* cultivars

To understand how miRNA expression may affect mRNA expression, we conducted RNA-seq for each of the seedling samples with 30 X coverage (Table S1). We report the results of the whole transcriptome analysis here and analyzed the expression correlation between miRNAs and targets with the transcriptome as a control in the next two sections. All reads were mapped to the *japonica* cv. Nipponbare genome (Kawahara et al. 2013). Approximately 81% and 86% of the unique reads could be mapped to the reference genome for *indica* and *japonica* cultivars, respectively. To allow differential expression analysis between rice subspecies, 32,044 homologous genes were identified between the genomes of *japonica* cv. Nipponbare and *indica* cv. 93-11 (see Methods), of which 18,288 were considered to be expressed, i.e. having more than ten raw reads across all three accessions from the same developmental stage and the same subspecies (Table S4). A regression comparison was then performed based on the expression level of these 18,288 expressed homologous genes. Among them, 2,530 genes (13.8%) were differentially expressed (fold-change ≥ 2 and FDR $\leq 5\%$) between the two rice subspecies, with 1,452 up-regulated in *indica* and 1,078 in *japonica*, respectively (Figure S4). To confirm the results of RNA-seq, we conducted qRT-PCR of ten experimentally validated or predicted miRNA targets and confirmed the differential expression patterns between rice subspecies in all the comparisons (Figure S3B and C).

The differentially expressed genes (DEGs) are significantly enriched in Gene Ontology categories, including “plasma membrane”, “cell wall”, “endoplasmic reticulum” and “extracellular region” for “Cellular Component”, categories of molecular binding (“Hydrocarbon binding”, “Oxygen binding”, “protein binding” etc.) and enzyme activities (“hydrolase activity”, “catalytic activity”, “kinase activity” etc.) for “Molecular function”, and “cellular process”, “flower development” and many categories of metabolic process (“Secondary metabolic process”, “lipid metabolic process”, etc.) or response to stimuli

and stress (“endogenous stimulus”, “abiotic stimulus” and “biotic stimulus”) for “Biological Process” (Figure 3A). Consistent with the GO analysis, DEGs are significantly enriched in the KEGG pathways related to secondary metabolites, such as “Glutathione metabolism” (Figure S5) and “starch and sucrose metabolism” (Figure S6 and Figure 3B). The former is known to play important roles in plant stress tolerance (Noctor et al. 1998), while the latter is presumably affecting the varietal difference of soluble sugar in different rice varieties (Yang et al. 2014). The overrepresentative KEGG pathways in DEGs also included “Plant-pathogen interaction” (Figure S7) and “Plant hormonal signal transduction” (Figure S8 and Figure 3B).

MicroRNAs, particularly highly conserved miRNAs, show a negatively-correlated expression pattern with their direct targets

As miRNAs negatively regulated target gene expression, one may expect to see negative correlation between expression of miRNAs and target mRNAs. To test this speculation, we performed correlation analyses either for all the miRNAs and their targets globally using the log2 fold changes of expression between *indica* and *japonica* subspecies, or for individual miRNA-target pairs using their expression levels across six rice cultivars. We shall focus on the global analysis in this section and present the results of the individual analysis in the next. Such analyses were based on four target sets, including one predicted target set for all the expressed miRNAs (target set I; see Methods for rice miRNA target prediction) and three experimentally-verified target sets by high-throughput degradome sequencing for *japonica* cv. Nipponbare (target sets II, III) or *indica* cv. 93-11 (target set IV) (Wu et al. 2009, Li et al. 2010, Zhou et al. 2010).

As for the global analysis, no correlation (Pearson’s correlation hereafter, P -value = 0.420, $r = -0.033$, $n = 609$) was observed for all the expressed miRNAs and their

predicted targets (target set I) (Figure S9A). However, when separating miRNAs according to their conservation, a weak but significant negative correlation was observed for the highly conserved miRNAs and their predicted targets (P -value = 0.019, r = -0.119, $n=390$ Figure 4A), but not for the lowly conserved miRNAs and targets (P -value = 0.611, r = 0.035, n = 219, Figure 4B). The negative correlation in expression fold changes was highly significant for the co-expressed miRNAs and degradome-verified target set IV (Zhou et al. 2010) (P -value = 0.002, r = -0.364, n = 68; Figure 4C), and nearly significant for target set II (Wu et al. 2009) (P -value = 0.121, r = -0.224, n = 49; Figure 4D) and target set III (P -value = 0.119, r = -0.161, $n=95$; Figure S9B) (Li et al. 2010). The extent of negative correlation seemed to be associated with the proportion of targets of highly conserved miRNAs in the analyzed target sets, which is 86%, 81% and 91% for target set II, III and IV, respectively. Taken together, the overall expression of miRNAs correlates negatively with the expression of their targets in rice, which is consistent with miRNA functions in guiding mRNA cleavage in plants (Bartel 2004). Highly conserved miRNAs exhibit less variation in expression but repress targets more strongly than lowly conserved miRNAs.

We then compared the *indica-japonica* expression differentiation of miRNAs relative to that of target genes using transcriptome as a control. Interestingly, the extent of differential expression between rice subspecies was much greater for miRNAs (median absolute fold change = 0.630) than for the whole transcriptome (median absolute fold change = 0.263). Kolmogorov-Smirnov test (KS test) on the cumulative plot shows that the differences are highly significant (P -value \ll 0.001; Figure S10A). In contrast, the extent of differential expression was comparable between miRNA targets (median absolute fold change = 0.279) and the transcriptome (median absolute fold change = 0.263, Figure S10B). Although lowly conserved miRNAs (median absolute fold

change= 0.677) were more variable in expression than highly conserved miRNAs (median absolute fold change= 0.457, KS test P -value $\ll 0.001$, Figure S10A), no such expression difference was observed for their targets (KS test, P -value = 0.926, Figure S10B). It is thus evident that miRNA targets exhibit less variation in expression than miRNAs.

Identification of individual miRNA-target pairs with significant expression correlations

We further examined the individual correlations of the expression of miRNA-mRNA pairs across the 6 accessions. The average correlation coefficient for the 476 miRNA-target pairs (in target set I) was -0.11. The highly conserved miRNAs tend to be more negatively correlated with their corresponding target mRNAs than the lowly conserved miRNAs in expression variation among accession (KS test, P -value = 0.0008, Figure 5A) but no pair passed the 0.05 FDR level after the multiple-testing correction.

We then asked whether the mean expression correlation among all miRNA-target pairs shifted to the negative end in comparison with the random miRNA-gene pairs. Using a permutation test as previously described (Nunez-Iglesias et al. 2010), we identified 51 significantly correlated miRNA-target pairs, including 29 negatively-correlated and 22 positively-correlated pairs (Table 3). Indeed, the mean correlation of the real miRNA-mRNA pairs is more negative than the permuted pairs (empirical P -value $\ll 0.001$, Figure 5B), both for the highly conserved (empirical P -value $\ll 0.001$, Figure 5C) and lowly conserved miRNAs (empirical P -value = 0.019, Figure 5D). Therefore, the overall tendency in the individual miRNA-target correlations is towards the negative correlation, which is in contrast to a similar test performed in human brain samples (Nunez-Iglesias et al. 2010). Also, the degradome verified targets

are significantly enriched in the correlated pairs (27 verified targets, Chi-squared test P -value $\ll 0.001$, Table 3), suggesting that those genes are indeed under the control of miRNA regulation.

Significantly-correlated miRNA-target pairs, both negatively and positively, participate in key biological processes such as reproduction, development, pigmentation, and stress responses (Figure S11). The negatively-correlated pairs included the well-documented miRNA-target pairs, such as miR156:*SQUAMOSA promoter binding protein-like (SPL)* (Miura et al. 2010); miR169:*Nuclear transcription factor Y (NF-Y)* (Zhao et al. 2009), miR172-*Apetala2 (AP2)* (Zhu et al. 2011), and miR396:*Growth-regulating factor (GRF)* (Debernardi et al. 2012) (Table 3). The involvement of the positively-correlated miRNA-target pairs in cell cycle, cell death, pollination and the transport process suggest that these pairs also play significant roles in rice development. Positive correlations between the miRNA and target expression levels across rice cultivars may be due to the miRNA-mediated regulatory circuits such as negative feedback loops or incoherent feedforward loops (Wu et al. 2009).

Discussion

We present here the first survey of miRNA expression variation in rice cultivars. Substantial miRNA expression changes were detected between rice grains and seedlings, consistent with the regulatory role of miRNAs in rice seed development (Zhu et al. 2008). In contrast, only a small fraction of miRNAs, mainly lowly conserved miRNAs, differed in expression level between rice *indica* and *japonica* subspecies. While miRNA genes are under strong purifying selection (Ehrenreich et al. 2008, Wang et al. 2010), the single nucleotide polymorphisms (SNP) densities of pre-miRNAs and mature miRNAs are significantly higher than their flanking regions in rice (Liu et al. 2013), implying that

cis-regulatory mutations affecting miRNA expression level are mostly deleterious and are thus quickly purged from the rice population. The low level of miRNA expression variation coupled with low level of *cis*-regulatory sequence polymorphism is consistent with the scenario that stabilizing selection commonly uses purifying selection to select against extreme values of characters, i.e. miRNA expression profiles in this study. Indeed, it has been demonstrated both empirically and theoretically that stabilizing selection is the major evolutionary force governing the evolution of gene expression (Denver et al. 2005, Rifkin et al. 2005, Hutter et al. 2008, Bedford et al. 2009, Hodgins-Davis et al. 2015). A recent study revealed that a vast majority of miRNAs are under stabilizing selection at the onset of *Drosophila* metamorphosis (Yeh et al. 2014). Using the same methodology, we estimated that the counts for rice miRNAs with expression variation compatible with particular evolutionary modes are 196, 29, 1, and 46 for stabilizing selection, directional selection, genetic drift and complex scenarios, respectively (see Supplementary text). About 72% of miRNAs in this study are not significantly differentiated within or between rice subspecies, compatible with the evolutionary mode of stabilizing selection. These miRNAs under stabilizing selection are mostly conserved between eudicots and monocots, whereas more than half of the miRNAs under directional selection are species-specific to rice.

It is remarkable that lowly and highly conserved miRNAs showed sharply contrasting patterns in expression variation and regulation strength. The great expression variation of the lowly conserved miRNAs is largely coupled with their high level of sequence polymorphism among cultivated rice (Liu et al. 2013), suggesting they are under weak selection pressures. Such a coupling of expression variation and sequence polymorphism of rice miRNAs is compatible with the correlated divergences between gene sequences and expression patterns during organ evolution in angiosperms (Yang et

al. 2013). The negligible overall correlation between expression changes of the lowly conserved miRNAs, also the lowly expressed miRNAs, and their targets further indicates that these miRNAs exert very modest, if any, repression on target genes. Therefore, the lowly conserved miRNAs are more like young miRNAs, which are expressed lowly or in specialized tissues, evolve rapidly, and tend to be lack of targets (Rajagopalan et al. 2006, Fahlgren et al. 2010, Ma et al. 2010), rather than the old, deeply conserved miRNAs.

While most young miRNAs are evolutionarily transient (Fahlgren et al. 2010), they can occasionally be selectively favored. A good case in point is a new miRNA specific in *japonica* rice, osa-miR7695, which negatively regulates an alternatively spliced transcript of OsNramp6 (Natural resistance-associated macrophage protein 6), conferring pathogen resistance (Campo et al. 2013). The differentially-expressed miRNAs between *indica* and *japonica* subspecies, which are resulted from low expression polymorphism within subspecies and high expression divergence between subspecies, may represent a class of miRNAs that are favored by artificial selection in rice domestication and/or improvement. Previous evolutionary analyses have identified rice miRNA genes that are putative candidates of positive selection, including highly conserved MIR164e, MIR395a/b, and MIR399d (Wang et al. 2010), and lowly conserved or rice-specific osa-miR5513, osa-miR818e, osa-miR1847, osa-miR1865, osa-miR160f, osa-miR5143, and osa-miR2118h-l (Liu et al. 2013). Among them, osa-miR395a-v,y and miR399a-d,i-j (unadjusted P -value < 0.05, FDR \approx 0.2, Table S2) showed significant differential expression between *indica* and *japonica* subspecies and the former correlated negatively with a putative target LOC_Os03g53230 in this study. In the Solanaceae, differential expression of the miR482/2188 gene family members that are under different evolutionary constraints also suggested miRNA subfunctionalization between closely related plant species (de Vries et al. 2015).

Another pattern that we observed is the prevalent positive correlation between individual miRNA-target pairs in the cultivated rice, of which many target genes have been experimentally verified by degradome data. Given miRNA negatively regulated target expression, this observation may reflect the composite effect of miRNA-mediated circuits, such as negative feedback loops (FBLs) and incoherent feedforward loops (FFLs) (Alon 2007). miRNA-mediated FBLs and FFLs are recurrent network motifs in animals (Tsang et al. 2007, Wu et al. 2009), and also play significant roles in plants as evidenced by both experiments (Xie et al. 2003, Bari et al. 2006, Vaucheret et al. 2006) and computational analyses (Meng et al. 2009, Wu et al. 2009). From the network perspective, miRNAs can thus play a role in expression buffering and biological robustness, as well recognized in animal systems (Wu et al. 2009, Herranz et al. 2010, Ebert et al. 2012, Pelaez et al. 2012). We found that targets of rice miRNAs overall have less expression variation in comparison with the transcriptome, albeit miRNAs themselves are more variable in expression. These results are consistent with the notion that miRNAs act as dampers in buffering target expression variation (Fei et al. 2013) or target sequence diversity (de Vries et al. 2015). In contrast, miRNA targets compared with non-targets overall have higher level of expression variation in human, despite a small number of highly expressed targets with decreased expression variation, suggesting the dual function of miRNA regulation (Lu et al. 2012).

Functional enrichment tests of the DEGs highlight that expression changes of genes involved in various metabolism processes and stress responses are important for the *indica-japonica* differentiation. Genes involved in plant-pathogen pathways are the most prominent examples, since the interaction between rice-host cultivar (genotype) and parasite population is shown to be critical in determining parasite affecting (Huang et al.

2012). Interestingly, we identified a target (LOC_Os02g30900, *PBS1*) of a lowly conserved miRNA, miR1857, which is involved in this pathway (Figure S7). The orthologous *PBS1* gene in *Arabidopsis* encodes a putative serine-threonine kinase, which is required for specific recognition of the bacterial protein *AvrPphB* (Swiderski et al. 2001). Another related example has also been recently reported in *Arabidopsis*, where a peptide derived from the plant pathogen *Pseudomonas syringae* induces the expression of the host miR393 that targets auxin receptor (*TIR1*, *AFB2*, and *AFB3*), and consequently inhibits the growth of the bacteria (Navarro et al. 2006). It would be interesting to explore the role of miRNAs in mediating biotic stress response in the context of differentiation between *indica* and *japonica* subspecies.

Our permutation analysis provides a promising approach to further classify the important miRNA regulation pairs in rice. For example, besides the well documented miRNA-target pairs mentioned previously, recent study revealed that osa-miR171c controls the floral transition and maintenance of shoot apical meristem (SAM) indeterminacy in rice by targeting *GRAS* (*GAI-RGA-SCR*), a plant-specific transcription factors (LOC_Os02g44370, also for *OsHAM2*, Table 3) (Fan et al. 2015); In addition, miR166-mediated posttranscriptional gene silencing of rice Class III HD-Zip genes (LOC_Os03g43930, also for *OsHB5*) is reported to be responsible for the auxin signals to regulate leaf and shoot development (Itoh et al. 2008, Toriba et al. 2010). There are also many miRNAs with un-verified targets in our list of correlated miRNA-target pairs (Table 3). Further studies are necessary to confirm the functional contribution of these miRNAs to the process of rice development and/or differentiation. The joint miRNA-mRNA data and permutation analyses used in this study provide a novel idea to study miRNA regulatory relationships in rice. Our results may provide a valuable

resource for further investigation of miRNA functions in rice developmental regulation, stress responses and biomass yields under domestication.

Methods

Plant materials

Seeds of *O. sativa* L. ssp. *indica* and ssp. *japonica* accessions as listed in Table 1 were obtained from the International Rice Research Institute (IRRI, Manila, Philippines). For grains, the husks of the seeds were removed before RNA extraction. For seedlings, rice seeds were sterilized and germinated in Petri dishes containing distilled water at 37°C under dark conditions for 2 days. The uniformly germinated seeds were transferred into Yoshida nutrient solution and grown under a 16-hour light (28°C)/8-hour dark (25°C) photoperiod for one week. The samples were collected and rinsed with double distilled water 3 times, and then immediately frozen in liquid nitrogen until use.

RNA isolation and preparation of sequencing libraries

Total RNA were extracted from rice grains and seedlings using TRIzol Reagent (Invitrogen), and evaluated using an Agilent 21100 Bioanalyzer (Agilent Technologies). Small RNA and transcriptome libraries were prepared using standard protocols of the Illumina Small RNA Sample Prep Kit or the Illumina TruSeq RNA Sample Prep kit, and sequenced using an Illumina Genome Analyzer (Illumina, San Diego, CA, USA) at BGI (Shenzhen, China). As we did not obtain enough quality RNA in grain, only the seedling samples were used for RNA-seq.

Expression analysis of miRNAs

For all small RNA libraries, after trimming prime adaptors and filtering low quality or adaptor contaminated reads, clean reads within the length range of 19 to 30 nt were

retained for further analysis. These reads were searched against the Rfam database (Griffiths-Jones et al. 2003) and the RepBase database (Jurka et al. 2005) using the SOAP software (Li et al. 2008) with two mismatches, in order to remove reads matching structural RNAs, including rRNA, tRNA, small nuclear RNA (snRNA) and small nucleolar RNA (snoRNA), and repeats/transposons. The remaining reads were mapped to the precursors of all known rice miRNAs registered in miRBase (Release 20, <http://www.mirbase.org/index.shtml>) using SOAP (Li et al. 2008). We grouped all the variants of rice mature miRNAs into 553 distinct mature miRNAs since some miRNA precursors produced more than one different mature sequences and some identical mature sequences were generated from distinct precursors. Read counts corresponding to individual mature miRNA sequences were normalized and rescaled as RPM (Reads Per Million), which divided the raw read count of each mature miRNA by the total mapped read count in each library and multiplied it by one million. Subsequent heatmap clustering and differential expression analysis were performed based on the average RPM of each distinct mature miRNA. An unsupervised, two-dimensional, hierarchical clustering was conducted using R package gplots (Warnes et al. 2009). For differential expression analysis, a generalized Poisson-regression linear model was fitted in the R package edgeR (Nikolayeva et al. 2014) to identify the differentially expressed miRNAs for the factors of development, subspecies, and development-by-subspecies. MiRNAs with fold change ≥ 2 and FDR ≤ 0.05 are denoted as significantly mi-expressed.

Expression analysis of mRNAs

For the transcriptome libraries, after removing adaptors and low-quality reads from raw reads, clean reads were mapped to the reference genome of rice (*japonica* cv. Nipponbare) using the Bowtie software package (Langmead et al. 2009). Reads that could be mapped equally well to multiple locations without mismatch or with up to 2 mismatches were

randomly assigned to one position and were retained for further analyses as previously described (Wang et al. 2009). According to the TIGR 6.0 gff3 file (Kawahara et al. 2013), reads matching gene or genomic region were recovered and expression level of each transcript was measured as numbers of reads per kilobase of exon region in a gene per million mapped reads (RPKM) as previously described (Mortazavi et al. 2008).

To identify homologous genes between rice subspecies, the longest isoform of each gene model of *japonica* cv. Nipponbare was aligned to the genome of *indica* cv. 93-11 by GMAP (Wu et al. 2005). We considered a Nipponbare gene and its counterpart in the 93-11 genome as an homologous gene pair if they share at least 95% sequence identity in at least 95% of their longest isoforms (He et al. 2010). The resulting 32,044 homologous gene pairs were further filtered by requiring at least 10 raw reads in all the three accessions sequenced for at least one developmental stage of a subspecies. These expressed homologous genes were then used for the subsequent differential expression analysis. The generalized Poisson-regression linear model was fitted and the likelihood ratio test was performed with lmtest (Zeileis et al. 2002) to identify the mRNAs with differential expression between rice subspecies. Genes with fold-change ≥ 2 and FDR $\leq 5\%$ were denoted as significantly differentially expressed.

Gene functional annotation

The gene ontology classification developed in the TIGR Rice Genome Annotation Project (<http://rice.plantbiology.msu.edu/>) was used to assign genes to a hierarchical biological process using the criteria of the Gene Ontology (GO) Consortium databases (Ashburner et al. 2000, Kawahara et al. 2013), and KEGG pathway annotation was identified according to KEGG database (<http://www.genome.jp/kegg/>) (Kanehisa et al. 2000, Kanehisa et al. 2016). The GO and KEGG enrichment test (FDR ≤ 0.05) was performed

by using the CORNA and clusterProfiler package, respectively (Wu et al. 2009, Yu et al. 2012). KEGG pathway rendering of the expressed genes was conducted by the R package Pathview (Luo et al. 2013). The DEGs with a fold change larger than 2 were used for the GO enrichment test and 1.5 for the KEGG pathway enrichment test, with a $FDR \leq 0.05$ was used as the significance threshold.

Identification of novel miRNAs

After removing the reads mapped to the structural RNAs, the known miRNAs from miRBase and repeats/transposons, the remaining sequences were mapped on the rice genome and analyzed by an adjusted miRDeep script for novel plant miRNA prediction (Wen et al. 2012). The following criteria for miRNA annotation (Wang et al. 2009) were further applied to filter the novel miRNAs, including: (1) a secondary structure must have a hairpin with at least 18 paired nucleotides in its stem region; (2) the hairpin must have free energy less than or equal to -18 kCal/mol and no more than two central loops; (3) the miRNA and miRNA* form a duplex with a two-nucleotide overhang; (4) fewer than four mismatches exist in the miRNA/miRNA* duplex; and (5) $\geq 75\%$ of the sRNAs mapped onto a miRNA precursor are derived from the miRNA or miRNA* region. The candidate sequences were extracted and folded with Vienna RNAfold (Lorenz et al. 2011). The miRNAs with more than 10 raw reads matching the mature sequences in at least two replicate sRNA libraries were considered to be expressed.

MiRNA target prediction and integration

A search for the miRNA target genes was performed for all known miRNAs and newly identified miRNA sequences on the *japonica* cv. Nipponbare cDNA dataset (TIGR 6.0) using psRNAtarget (Dai et al. 2011) with a maximum expectation score of 2.5 to reduce the false discovery rate (Klevebring et al. 2009). The predicted target set was referred to

as target set I in this study. Three additional sets of miRNA targets that are verified by degradome analysis were also used, which include target set II (Wu et al. 2009) and target set III (Li et al. 2010) for rice *japonica* cv. Nipponbare, and target set IV for rice *indica* cv. 93-11 (Zhou et al. 2010).

miRNA target permutation analysis

The permutation test and weighted shift for the miRNA-target correlations were processed as described previously (Nunez-Iglesias et al. 2010). In short, the miRNA-target pairs can be considered a bipartite graph, with nodes representing the miRNAs on one side and the targets on the other, and the edges represent the target prediction relationships. The nodes have associated expression measurements. Correlation statistics were then computed for each miRNA-target pair across the 6 rice lineages. Finally, the network was permuted by shuffling the edges, re-computing the statistics, and then repeating this processes 1000 times. From these, we can obtain an empirical P-value and a “weighted correlation shift”, W , which was defined as the difference between the true value and the mean permuted value divided by the standard deviation of the permuted values: $W=(r-r_0)/S_0$.

Quantitative miRNA and mRNA analysis by qRT-PCR

For miRNA quantification, total RNA from one week old seedlings was subjected to stem-loop reverse transcription (RT) (Chen et al. 2005) followed by Taqman PCR (Applied Biosystems) using the miRNA UPL (Roche Diagnostics) probe assay as described previously (Varkonyi-Gasic et al. 2007, He et al. 2016). 5.8S rRNA was used as an internal control. Three biological replicates were examined. For mRNA quantification, the same total RNA was first treated with TURBO DNA-free kit (Ambion)

to remove potential genomic DNA contamination and then used for reverse transcription with SuperScript III First-Strand Synthesis System (Invitrogen). qRT-PCR was performed with Platinum SYBR Green qPCR SuperMix (Invitrogen) according to the manufacturer's instructions. *Actin* was used as an internal control. Three biological replicates with two technique repeats each were examined to ensure reproducibility. The relative levels of miRNA or mRNA were calculated using the $2^{-\Delta\Delta CT}$ method (Livak et al. 2001). The sequences of primers used are listed in Supplementary Table S5.

Data availability

The expression data generated by this study are available in the NCBI Gene Expression Omnibus (GEO) under accession GSE71925.

References

- Alon, U. (2007). "Network motifs: theory and experimental approaches." *Nat Rev Genet* **8**(6): 450-461.
- Ashburner, M., C. A. Ball, J. A. Blake, D. Botstein, H. Butler, et al. (2000). "Gene Ontology: tool for the unification of biology." *Nature Genetics* **25**(1): 25-29.
- Axtell, M. J. (2013). "Classification and comparison of small RNAs from plants." *Annu Rev Plant Biol* **64**: 137-159.
- Bari, R., B. Datt Pant, M. Stitt and W. R. Scheible (2006). "PHO2, microRNA399, and PHR1 define a phosphate-signaling pathway in plants." *Plant Physiol* **141**(3): 988-999.
- Bartel, D. (2004). "MicroRNAs: Genomics, biogenesis, mechanism, and function." *Cell* **116**(2): 281-297.
- Bartel, D. P. (2009). "MicroRNAs: target recognition and regulatory functions." *Cell* **136**(2): 215-233.
- Bedford, T. and D. L. Hartl (2009). "Optimization of gene expression by natural selection." *Proc Natl Acad Sci U S A* **106**(4): 1133-1138.
- Britten, R. J. and E. H. Davidson (1971). "Repetitive and non-repetitive DNA sequences and a speculation on the origins of evolutionary novelty." *Q Rev Biol* **46**(2): 111-138.

- Campo, S., C. Peris-Peris, C. Sire, A. B. Moreno, L. Donaire, et al. (2013). "Identification of a novel microRNA (miRNA) from rice that targets an alternatively spliced transcript of the Nramp6 (Natural resistance-associated macrophage protein 6) gene involved in pathogen resistance." New Phytol **199**(1): 212-227.
- Carlsbecker, A., J.-Y. Lee, C. J. Roberts, J. Dettmer, S. Lehesranta, et al. (2010). "Cell signalling by microRNA165/6 directs gene dose-dependent root cell fate." Nature **465**(7296): 316-321.
- Carrington, J. C. and V. Ambros (2003). "Role of microRNAs in plant and animal development." Science **301**(5631): 336-338.
- Chen, C., D. A. Ridzon, A. J. Broomer, Z. Zhou, D. H. Lee, et al. (2005). "Real-time quantification of microRNAs by stem-loop RT-PCR." Nucleic Acids Res **33**(20): e179.
- Chen, F. F., G. M. He, H. He, W. Chen, X. P. Zhu, et al. (2010). "Expression Analysis of miRNAs and Highly-expressed Small RNAs in Two Rice Subspecies and their Reciprocal Hybrids." Journal of Integrative Plant Biology **52**(11): 971-980.
- Chuck, G., H. Candela and S. Hake (2009). "Big impacts by small RNAs in plant development." Current Opinion in Plant Biology **12**(1): 81-86.
- Cuperus, J. T., N. Fahlgren and J. C. Carrington (2011). "Evolution and functional diversification of MIRNA genes." Plant Cell **23**(2): 431-442.
- Dai, X. B. and P. X. Zhao (2011). "psRNATarget: a plant small RNA target analysis server." Nucleic Acids Research **39**: W155-W159.
- de Vries, S., T. Kloesges and L. E. Rose (2015). "Evolutionarily Dynamic, but Robust, Targeting of Resistance Genes by the miR482/2118 Gene Family in the Solanaceae." Genome Biol Evol **7**(12): 3307-3321.
- Debernardi, J. M., R. E. Rodriguez, M. A. Mecchia and J. F. Palatnik (2012). "Functional Specialization of the Plant miR396 Regulatory Network through Distinct MicroRNA-Target Interactions." Plos Genetics **8**(1).
- Denver, D. R., K. Morris, J. T. Streelman, S. K. Kim, M. Lynch, et al. (2005). "The transcriptional consequences of mutation and natural selection in *Caenorhabditis elegans*." Nat Genet **37**(5): 544-548.
- Ding, Y. F., Z. Chen and C. Zhu (2011). "Microarray-based analysis of cadmium-responsive microRNAs in rice (*Oryza sativa*)." Journal of Experimental Botany **62**(10): 3563-3573.
- Ebert, M. S. and P. A. Sharp (2012). "Roles for microRNAs in conferring robustness to biological processes." Cell **149**(3): 515-524.

Ehrenreich, I. M. and M. D. Purugganan (2008). "Sequence variation of MicroRNAs and their binding sites in Arabidopsis." *Plant Physiol* **146**(4): 1974-1982.

Fahlgren, N. and J. C. Carrington (2010). "miRNA Target Prediction in Plants." *Methods Mol Biol* **592**: 51-57.

Fahlgren, N., S. Jogdeo, K. D. Kasschau, C. M. Sullivan, E. J. Chapman, et al. (2010). "MicroRNA gene evolution in Arabidopsis lyrata and Arabidopsis thaliana." *Plant Cell* **22**(4): 1074-1089.

Fan, T., X. Li, W. Yang, K. Xia, J. Ouyang, et al. (2015). "Rice osa-miR171c Mediates Phase Change from Vegetative to Reproductive Development and Shoot Apical Meristem Maintenance by Repressing Four OsHAM Transcription Factors." *PLoS One* **10**(5): e0125833.

Fei, Q., R. Xia and B. C. Meyers (2013). "Phased, secondary, small interfering RNAs in posttranscriptional regulatory networks." *Plant Cell* **25**(7): 2400-2415.

Filipowicz, W., S. N. Bhattacharyya and N. Sonenberg (2008). "Mechanisms of post-transcriptional regulation by microRNAs: are the answers in sight?" *Nature Reviews Genetics* **9**(2): 102-114.

Griffiths-Jones, S., A. Bateman, M. Marshall, A. Khanna and S. R. Eddy (2003). "Rfam: an RNA family database." *Nucleic Acids Res* **31**(1): 439-441.

He, G., X. Zhu, A. A. Elling, L. Chen, X. Wang, et al. (2010). "Global Epigenetic and Transcriptional Trends among Two Rice Subspecies and Their Reciprocal Hybrids." *Plant Cell* **22**(1): 17-33.

He, L., M. Xie, J. Huang, T. Zhang, S. Shi, et al. (2016). "Efficient and specific inhibition of plant microRNA function by anti-microRNA oligonucleotides (AMOs) in vitro and in vivo." *Plant Cell Rep* **35**(4): 933-945.

He, Z. W., W. W. Zhai, H. J. Wen, T. Tang, Y. Wang, et al. (2011). "Two Evolutionary Histories in the Genome of Rice: the Roles of Domestication Genes." *Plos Genetics* **7**(6).

Herranz, H. and S. M. Cohen (2010). "MicroRNAs and gene regulatory networks: managing the impact of noise in biological systems." *Genes Dev* **24**(13): 1339-1344.

Hodgins-Davis, A., D. P. Rice and J. P. Townsend (2015). "Gene Expression Evolves under a House-of-Cards Model of Stabilizing Selection." *Mol Biol Evol* **32**(8): 2130-2140.

Huang, K., R. Whitlock, M. C. Press and J. D. Scholes (2012). "Variation for host range within and among populations of the parasitic plant *Striga hermonthica*." *Heredity* **108**(2): 96-104.

Hutter, S., S. S. Saminadin-Peter, W. Stephan and J. Parsch (2008). "Gene expression variation in African

and European populations of *Drosophila melanogaster*." Genome Biol **9**(1): R12.

Itoh, J., K. Hibara, Y. Sato and Y. Nagato (2008). "Developmental role and auxin responsiveness of Class III homeodomain leucine zipper gene family members in rice." Plant Physiol **147**(4): 1960-1975.

Jiao, Y. Q., Y. H. Wang, D. W. Xue, J. Wang, M. X. Yan, et al. (2010). "Regulation of OsSPL14 by OsmiR156 defines ideal plant architecture in rice." Nature Genetics **42**(6): 541-U536.

Jones-Rhoades, M. W., D. P. Bartel and B. Bartel (2006). "MicroRNAs and their regulatory roles in plants." Annual Review of Plant Biology **57**: 19-53.

Jurka, J., V. V. Kapitonov, A. Pavlicek, P. Klonowski, O. Kohany, et al. (2005). "Repbase Update, a database of eukaryotic repetitive elements." Cytogenet Genome Res **110**(1-4): 462-467.

Kanehisa, M. and S. Goto (2000). "KEGG: kyoto encyclopedia of genes and genomes." Nucleic Acids Res **28**(1): 27-30.

Kanehisa, M., Y. Sato, M. Kawashima, M. Furumichi and M. Tanabe (2016). "KEGG as a reference resource for gene and protein annotation." Nucleic Acids Res **44**(D1): D457-462.

Kawahara, Y., M. de la Bastide, J. P. Hamilton, H. Kanamori, W. R. McCombie, et al. (2013). "Improvement of the *Oryza sativa* Nipponbare reference genome using next generation sequence and optical map data." Rice (N Y) **6**(1): 4.

King, M. C. and A. C. Wilson (1975). "Evolution at two levels in humans and chimpanzees." Science **188**(1126): 107-116.

Klevebring, D., N. R. Street, N. Fahlgren, K. D. Kasschau, J. C. Carrington, et al. (2009). "Genome-wide profiling of populus small RNAs." BMC Genomics **10**: 620.

Langmead, B., C. Trapnell, M. Pop and S. L. Salzberg (2009). "Ultrafast and memory-efficient alignment of short DNA sequences to the human genome." Genome Biol **10**(3): R25.

Li, R., Y. Li, K. Kristiansen and J. Wang (2008). "SOAP: short oligonucleotide alignment program." Bioinformatics **24**(5): 713-714.

Li, Y. F., Y. Zheng, C. Addo-Quaye, L. Zhang, A. Saini, et al. (2010). "Transcriptome-wide identification of microRNA targets in rice." Plant J **62**(5): 742-759.

Liu, Q., H. Wang, L. Zhu, H. Hu and Y. Sun (2013). "Genome-wide identification and analysis of miRNA-related single nucleotide polymorphisms (SNPs) in rice." Rice (N Y) **6**(1): 10.

- Liu, Q., Y. C. Zhang, C. Y. Wang, Y. C. Luo, Q. J. Huang, et al. (2009). "Expression analysis of phytohormone-regulated microRNAs in rice, implying their regulation roles in plant hormone signaling." *Febs Letters* **583**(4): 723-728.
- Livak, K. J. and T. D. Schmittgen (2001). "Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2- $\Delta\Delta$ CT Method." *Methods* **25**(4): 402-408.
- Lorenz, R., S. H. Bernhart, C. Honer Zu Siederdissen, H. Tafer, C. Flamm, et al. (2011). "ViennaRNA Package 2.0." *Algorithms Mol Biol* **6**: 26.
- Lu, J. and A. G. Clark (2012). "Impact of microRNA regulation on variation in human gene expression." *Genome Res* **22**(7): 1243-1254.
- Luo, L., W. Q. Li, K. Miura, M. Ashikari and J. Kozuka (2012). "Control of Tiller Growth of Rice by OsSPL14 and Strigolactones, Which Work in Two Independent Pathways." *Plant and Cell Physiology* **53**(10): 1793-1801.
- Luo, W. and C. Brouwer (2013). "Pathview: an R/Bioconductor package for pathway-based data integration and visualization." *Bioinformatics* **29**(14): 1830-1831.
- Lv, D. K., X. Bai, Y. Li, X. D. Ding, Y. Ge, et al. (2010). "Profiling of cold-stress-responsive miRNAs in rice by microarrays." *Gene* **459**(1-2): 39-47.
- Ma, Z., C. Coruh and M. J. Axtell (2010). "Arabidopsis lyrata small RNAs: transient MIRNA and small interfering RNA loci within the Arabidopsis genus." *Plant Cell* **22**(4): 1090-1103.
- Macovei, A., S. S. Gill and N. Tuteja (2012). "microRNAs as promising tools for improving stress tolerance in rice." *Plant Signaling & Behavior* **7**(10): 1296-1301.
- Meng, Y., F. Huang, Q. Shi, J. Cao, D. Chen, et al. (2009). "Genome-wide survey of rice microRNAs and microRNA-target pairs in the root of a novel auxin-resistant mutant." *Planta* **230**(5): 883-898.
- Meyers, B. C., M. J. Axtell, B. Bartel, D. P. Bartel, D. Baulcombe, et al. (2008). "Criteria for Annotation of Plant MicroRNAs." *Plant Cell* **20**(12): 3186-3190.
- Miura, K., M. Ikeda, A. Matsubara, X. J. Song, M. Ito, et al. (2010). "OsSPL14 promotes panicle branching and higher grain productivity in rice." *Nature Genetics* **42**(6): 545-U102.
- Mortazavi, A., B. A. Williams, K. McCue, L. Schaeffer and B. Wold (2008). "Mapping and quantifying mammalian transcriptomes by RNA-Seq." *Nature Methods* **5**: 621-628.

- Navarro, L., P. Dunoyer, F. Jay, B. Arnold, N. Dharmasiri, et al. (2006). "A plant miRNA contributes to antibacterial resistance by repressing auxin signaling." *Science* **312**(5772): 436-439.
- Nikolayeva, O. and M. D. Robinson (2014). "edgeR for differential RNA-seq and ChIP-seq analysis: an application to stem cell biology." *Methods Mol Biol* **1150**: 45-79.
- Noctor, G., A. C. M. Arisi, L. Jouanin, K. J. Kunert, H. Rennenberg, et al. (1998). "Glutathione: biosynthesis, metabolism and relationship to stress tolerance explored in transformed plants." *Journal of Experimental Botany* **49**(321): 623-647.
- Nunez-Iglesias, J., C. C. Liu, T. E. Morgan, C. E. Finch and X. J. Zhou (2010). "Joint genome-wide profiling of miRNA and mRNA expression in Alzheimer's disease cortex reveals altered miRNA regulation." *PLoS One* **5**(2): e8898.
- Pelaez, N. and R. W. Carthew (2012). "Biological robustness and the role of microRNAs: a network perspective." *Curr Top Dev Biol* **99**: 237-255.
- Rajagopalan, R., H. Vaucheret, J. Trejo and D. P. Bartel (2006). "A diverse and evolutionarily fluid set of microRNAs in Arabidopsis thaliana." *Genes Dev* **20**(24): 3407-3425.
- Reyes, J. L. and N. H. Chua (2007). "ABA induction of miR159 controls transcript levels of two MYB factors during Arabidopsis seed germination." *Plant J* **49**(4): 592-606.
- Rifkin, S. A., D. Houle, J. Kim and K. P. White (2005). "A mutation accumulation assay reveals a broad capacity for rapid evolution of gene expression." *Nature* **438**(7065): 220-223.
- Shen, Y., Y. Lv, L. Huang, W. Liu, M. Wen, et al. (2011). "Testing hypotheses on the rate of molecular evolution in relation to gene expression using microRNAs." *Proc Natl Acad Sci U S A* **108**(38): 15942-15947.
- Swiderski, M. R. and R. W. Innes (2001). "The Arabidopsis PBS1 resistance gene encodes a member of a novel protein kinase subfamily." *Plant Journal* **26**(1): 101-112.
- Toriba, T., T. Suzuki, T. Yamaguchi, Y. Ohmori, H. Tsukaya, et al. (2010). "Distinct regulation of adaxial-abaxial polarity in anther patterning in rice." *Plant Cell* **22**(5): 1452-1462.
- Tsang, J., J. Zhu and A. van Oudenaarden (2007). "MicroRNA-mediated feedback and feedforward loops are recurrent network motifs in mammals." *Mol Cell* **26**(5): 753-767.
- Varkonyi-Gasic, E., R. Wu, M. Wood, E. F. Walton and R. P. Hellens (2007). "Protocol: a highly sensitive RT-PCR method for detection and quantification of microRNAs." *Plant Methods* **3**: 12.

- Vaucheret, H., A. C. Mallory and D. P. Bartel (2006). "AGO1 homeostasis entails coexpression of MIR168 and AGO1 and preferential stabilization of miR168 by AGO1." Mol Cell **22**(1): 129-136.
- Wang, X. F., A. A. Elling, X. Y. Li, N. Li, Z. Y. Peng, et al. (2009). "Genome-Wide and Organ-Specific Landscapes of Epigenetic Modifications and Their Relationships to mRNA and Small RNA Transcriptomes in Maize." Plant Cell **21**(4): 1053-1069.
- Wang, Y., D. Shen, S. Bo, H. Chen, J. Zheng, et al. (2010). "Sequence variation and selection of small RNAs in domesticated rice." BMC Evol Biol **10**: 119.
- Warnes, G. R., B. Bolker, L. Bonebakker, R. Gentleman, W. Huber, et al. (2009). "gplots: Various R programming tools for plotting data." R package version **2**(4).
- Wen, M., Y. Shen, S. Shi and T. Tang (2012). "miREvo: an integrative microRNA evolutionary analysis platform for next-generation sequencing experiments." BMC Bioinformatics **13**: 140.
- Wu, C. I., Y. Shen and T. Tang (2009). "Evolution under canalization and the dual roles of microRNAs: a hypothesis." Genome Res **19**(5): 734-743.
- Wu, L., Q. Zhang, H. Zhou, F. Ni, X. Wu, et al. (2009). "Rice MicroRNA effector complexes and targets." Plant Cell **21**(11): 3421-3435.
- Wu, T. D. and C. K. Watanabe (2005). "GMAP: a genomic mapping and alignment program for mRNA and EST sequences." Bioinformatics **21**(9): 1859-1875.
- Wu, X. and M. Watson (2009). "CORNA: testing gene lists for regulation by microRNAs." Bioinformatics **25**(6): 832-833.
- Xie, Z., K. D. Kasschau and J. C. Carrington (2003). "Negative feedback regulation of Dicer-Like1 in Arabidopsis by microRNA-guided mRNA degradation." Curr Biol **13**(9): 784-789.
- Xue, L. J., J. J. Zhang and H. W. Xue (2009). "Characterization and expression profiles of miRNAs in rice seeds." Nucleic Acids Res **37**(3): 916-930.
- Yang, R. and X. Wang (2013). "Organ evolution in angiosperms driven by correlated divergences of gene sequences and expression patterns." Plant Cell **25**(1): 71-82.
- Yang, Y., Y. Rao, J. Xu, G. Shao, Y. Leng, et al. (2014). "Genetic analysis of sugar-related traits in rice grain." South African Journal of Botany **93**: 137-141.
- Yeh, S. D., M. von Grotthuss, K. A. Gandasetiawan, S. Jayasekera, X. Q. Xia, et al. (2014). "Functional

divergence of the miRNA transcriptome at the onset of *Drosophila* metamorphosis." *Mol Biol Evol* **31**(10): 2557-2572.

Yu, G., L. G. Wang, Y. Han and Q. Y. He (2012). "clusterProfiler: an R package for comparing biological themes among gene clusters." *OMICS* **16**(5): 284-287.

Zeileis, A. and T. Hothorn (2002). "Diagnostic Checking in Regression Relationships." *R News* **2**(3): 7-10.

Zhang, B. H., X. P. Pan, G. P. Cobb and T. A. Anderson (2006). "Plant microRNA: A small regulatory molecule with big impact." *Developmental Biology* **289**(1): 3-16.

Zhang, Y. C., Y. Yu, C. Y. Wang, Z. Y. Li, Q. Liu, et al. (2013). "Overexpression of microRNA OsmiR397 improves rice yield by increasing grain size and promoting panicle branching." *Nature Biotechnology* **31**(9): 848-+.

Zhao, B. T., L. F. Ge, R. Q. Liang, W. Li, K. C. Ruan, et al. (2009). "Members of miR-169 family are induced by high salinity and transiently inhibit the NF-YA transcription factor." *Bmc Molecular Biology* **10**.

Zhou, M., L. Gu, P. Li, X. Song, L. Wei, et al. (2010). "Degradome sequencing reveals endogenous small RNA targets in rice (*Oryza sativa* L. ssp. indica)." *Frontiers in Biology* **5**(1): 67-90.

Zhu, Q. H. and C. A. Helliwell (2011). "Regulation of flowering time and floral patterning by miR172." *J Exp Bot* **62**(2): 487-495.

Zhu, Q. H., A. Spriggs, L. Matthew, L. Fan, G. Kennedy, et al. (2008). "A diverse set of microRNAs and microRNA-like small RNAs in developing rice grains." *Genome Res* **18**(9): 1456-1465.

Acknowledgements

We thank International Rice Research Institute for providing rice germplasm. We thank Han Liang for reading the early version of this manuscripts and providing instructive comments. This work was supported by National Science Foundation of China (31170308 and 91231117 to T.T., and 41130208 to S.S.), the Science Foundation for Outstanding Young Teachers in Higher Education of Guangdong (Yq2013005) and the Fundamental Research Funds for the Central Universities (16lgjc75) to T. T., and the General Financial Grant from the China Postdoctoral Science Foundation (2013M531981) to M.W.

Table 1. Summary of the reads mapping of small RNAs and transcriptomes by RNA sequencing.

Rice variety				Small RNAs			Transcriptome		
Tissue	Subspecies	Accession Name	Accession No. ^a	No. of sequences generated	No. of sequences matching the rice genome	No. of unique sequences	No. of sequences generated ^b	No. of sequences matching the rice genome ^c	Unique mapped reads ^c
Seedling	<i>Indica</i>	Khao Dawk Mali 105	IRGC 27748	9 985 373	9 758 028(97.7%)	2 786 744(27.9%)	20 326 850	34 063 783(83.79%)	33 569 936(82.6%)
		Guangluai 4	IRGC 114900	9 788 880	9 558 792(97.7%)	3 048 520(31.1%)	20 216 686	32 972 884(81.54%)	32 494 623(80.4%)
		Rathuwee	IRGC 8952	9 997 943	9 701 803(97.0%)	3 403 862(34.0%)	19 876 912	32 789 681(82.48%)	32 328 952(81.3%)
	<i>Japonica</i>	Taipei 309	IRGC 42576	8 221 032	8 051 214(97.9%)	2 282 686(27.8%)	19 094 551	33 286 973(87.16%)	32 855 807(86.0%)
		Heukgyeong	IRGC 55530	9 371 607	8 988 554(95.9%)	2 203 943(23.5%)	19 889 198	34 863 682(87.64%)	34 372 869(86.4%)
		Nipponbare	IRGC 12731	10 592 461	10 416 128(98.3%)	3 010 461(28.4%)	18 818 192	32 995 013(87.67%)	325 922 62(86.6%)
Grain	<i>Indica</i>	Khao Dawk Mali 105	IRGC 27748	14 396 308	13 994 396(97.2%)	5 408 929(37.6%)	NA	NA	NA
		Guangluai 4	IRGC 114900	14 433 295	14 137 205(98.0%)	4 056 622(28.1%)	NA	NA	NA
		Rathuwee	IRGC 8952	15 718 282	15 185 127(96.6%)	5 740 388(36.5%)	NA	NA	NA
	<i>Japonica</i>	Taipei 309	IRGC 42576	14 846 530	14 293 090(96.3%)	2 786 744(18.8%)	NA	NA	NA
		Heukgyeong	IRGC 55530	10 225 740	9 512 313(93.0%)	3 081 796(30.1%)	NA	NA	NA
		Nipponbare	IRGC 12731	13 862 947	13 356 435(96.4%)	5 014 395(36.2%)	NA	NA	NA
	Total			141 440 398	136 953 085(96.8%)	42 825 090(30.3%)	NA	NA	NA

^a International Rice Germplasm Collection at IRRI in the Philippines (<http://archive.irri.org/GRC/requests/requests.htm>)

^b Number of pairs of 75-nt paired-ends sequencing reads.

^c The proportion was calculated as the number of mapped reads versus the number of total reads.

NA, Not applicable

Table 2. Summary of differentially expressed miRNA families. MiRNAs with fold change ≥ 2 and $FDR \leq 0.05$ are defined as differentially expressed.

Effect	Highly conserved (n=111)	Lowly conserved (n=161)	Total (n=272)
Developmental stage	64 (57.7%)	82 (50.9%)	146 (53.7%)
Subspecies	2 (1.8%)	28 (17.4%)	30 (11.0%)
Development×Subspecies	0 (0%)	6 (3.7%)	6 (2.2%)

Table 3. Significantly correlated miRNA-mRNA pairs in the permutation test.

miRNA	miRNA FC ^a	mRNA FC ^b	Correla tion ^c	W	Target	TIGR annotation ^d	Target verification ^e
miR156k	0.080	0.142	0.953	2.082	LOC_Os01g 69830	OsSPL2-SBP-box gene family member	Y ^{3,4}
miR160a-d	-0.048	-0.040	-0.815	-2.096	LOC_Os06g 47150	Auxin response factor 18	Y ^{2,3,4}
miR160e	0.537	-0.040	-0.817	-1.923	LOC_Os06g 47150	Auxin response factor 18	Y ^{2,3,4}
		0.222	0.873	1.573	LOC_Os04g 43910	Auxin response factor	Y ^{2,3,4}
miR160f	0.379	0.222	0.702	1.593	LOC_Os04g 43910	Auxin response factor	Y ^{2,3,4}
miR166a-d,f,n	0.073	2.722	0.888	1.845	LOC_Os04g 48290	MATE efflux family protein	Y ²
miR166g,h	0.203	0.383	0.791	1.427	LOC_Os03g 43930	START domain containing protein	Y ^{3,4}
miR166m	-0.252	-0.223	0.864	2.339	LOC_Os08g 34740	SGT1 protein	N
miR168a	0.624	-1.165	-0.902	-1.639	LOC_Os11g 44860	Cysteine-rich receptor-like protein kinase 28 precursor	N
miR169f-g	0.661	-0.930	-0.940	-1.922	LOC_Os03g 29760	Nuclear transcription factor Y subunit	Y ^{2,3,4}
miR169h-m	0.846	-1.145	-0.948	-1.747	LOC_Os07g 41720	Nuclear transcription factor Y subunit	Y ^{2,3,4}
miR171b-f	0.242	-0.126	-0.774	-1.880	LOC_Os05g 34460	OsDegp7 - Putative Deg protease homologue	N
		0.098	0.668	1.649	LOC_Os02g 44370	Myosin	Y ^{2,3,4}
miR171i	-0.426	-0.375	0.882	1.933	LOC_Os03g 04300	Targeting protein-related	N
miR172b	-0.175	0.105	-0.867	-1.747	LOC_Os05g 03040	AP2 domain containing protein	Y ^{2,3,4}
		0.321	-0.772	-1.593	LOC_Os03g 44420	Tubulin/FtsZ domain containing protein	N
miR172c	0.072	-0.044	0.855	1.921	LOC_Os07g 13170	AP2 domain containing protein	Y ^{2,3,4}
miR319a-b	0.171	-0.065	0.930	2.056	LOC_Os08g 16660	Aspartic proteinase nepenthesin precursor	N
miR393	0.766	-1.128	-0.695	-1.645	LOC_Os03g 36080	Expressed protein	Y ³
miR395b,d-e, g-n,p-s,y	1.449	-0.505	-0.972	-1.944	LOC_Os03g 53230	Bifunctional 3-phosphoadenosine 5-phosphosulfate synthetase	N
miR396d-e	-0.035	0.053	-0.991	-2.419	LOC_Os06g	Growth-regulating	Y ^{2,3,4}

		0.129	-0.855	-1.999	02560	factor	
					LOC_Os03g	Growth regulating	Y ^{3,4}
		-0.325	-0.812	-1.887	47140	factor protein	
					LOC_Os02g	Ankyrin repeat domain	Y ^{3,4}
		0.241	-0.749	-1.690	53690	containing protein	
					LOC_Os04g	Growth-regulating	Y ^{3,4}
		-0.008	-0.693	-1.654	51190	factor	
					LOC_Os11g	Growth regulating	Y ^{2,3,4}
					35030	factor protein	
miR396f	-0.054	0.053	-0.987	-2.399	LOC_Os06g	Growth-regulating	Y ^{2,3,4}
					02560	factor	
		0.129	-0.870	-2.030	LOC_Os03g	Growth regulating	Y ^{3,4}
					47140	factor protein	
		-0.325	-0.795	-1.843	LOC_Os02g	Growth regulating	Y ^{3,4}
					53690	factor protein	
		0.241	-0.744	-1.671	LOC_Os04g	Growth-regulating	Y ^{3,4}
					51190	factor	
miR397a	-0.076	0.413	-0.812	-1.833	LOC_Os11g		Y ³
					48060	Laccase-22 precursor	
miR397b	-0.108	0.413	-0.828	-1.875	LOC_Os11g		Y ³
					48060	Laccase-22 precursor	
		1.263	0.731	1.745	LOC_Os09g		N
					27950	Galactosyltransferase	
miR444b-c	-0.466	0.006	0.678	1.797	LOC_Os02g		Y ^{2,3}
					49840	Scarecrow	
miR444f	1.947	-0.646	-0.941	-1.789	LOC_Os02g	Targeting protein for	N
					34080	Xklp2	
miR528	-0.581	0.136	-0.897	-1.583	LOC_Os08g	Copper/zinc superoxide	N
					44770	dismutase	
miR529b	-1.633	0.219	-0.876	-1.923	LOC_Os04g	Peptidyl-prolyl	N
					28420	isomerase	
		-0.038	-0.687	-1.529	LOC_Os12g	Carboxyvinyl-carboxyp	N
					08760	hosphonate	
						phosphorylmutase	
miR530	-0.900	-0.678	0.794	2.288	LOC_Os05g	Ubiquinone	N
					09650	biosynthesis protein	
						COQ4	
miR1847	-0.376	-0.150	0.912	1.924	LOC_Os01g	Laccase precursor	N
					63190	protein	
miR1856	-6.129	-1.509	0.827	1.785	LOC_Os09g		N
					10274	Expressed protein	
miR1860	0.174	-0.069	0.920	2.125	LOC_Os01g		N
					01030	Monocopper oxidase	
miR1861a	1.984	0.021	0.796	1.590	LOC_Os03g	PPR repeat containing	N
					40020	protein	
miR1864	-1.514	0.414	-0.863	-1.374	LOC_Os01g		N
					14020	Expressed protein	
miR1873	0.003	0.223	-0.717	-2.359	LOC_Os05g		N
					01790	Expressed protein	

miR1884a	0.575	-1.074	-0.685	-1.786	LOC_Os11g34910	Expressed protein	Y ²
		0.435	0.952	1.782	LOC_Os06g14780	Expressed protein	N
miR1884b	0.083	-0.039	-0.677	-2.294	LOC_Os12g01680	Macrophage migration inhibitory factor	N
		0.036	-0.573	-1.865	LOC_Os02g49870	Expressed protein	N
		-0.231	0.575	1.640	LOC_Os01g64520	Uricase	Y ²
		0.435	0.636	1.786	LOC_Os06g14780	Expressed protein	N
miR2097	0.543	0.787	0.779	1.585	LOC_Os08g43920	Carrier	N

^a Fold change in the log2 ratio of miRNA expression between *indica* and *japonica* subspecies in seedlings. The mean expression levels averaged from three accessions were used for the calculation. MiRNAs with significant differentially expressions (unadjusted *P*-value ≤ 0.05 and Fold change ≥ 2) between rice subspecies are marked in bold.

^b Fold change in the log2 ratio of mRNA expression between *indica* and *japonica* subspecies in seedlings. The mean expression levels averaged from three accessions were used for the calculation. Genes with significantly differential expression are marked in bold.

^c Pearson's correlation coefficient.

^d Gene annotations from TIGR (Version 6).

^e The Y^{2,3,4} Target is verified in the corresponding degradome target set II, III, or IV, respectively; N: not yet verified.

Figure legends

Figure 1. Expression variation of known miRNAs in rice. (A) Heat map and unsupervised hierarchical clustering of known miRNA expression. The color key represented the scale of the relative expression levels of the miRNAs (log2 RPM). KDM: *indica* cv. Khal Dawk Mali 105; GLA4: *indica* cv. Guangluai 4; PATH: *indica* cv. Rathuwee; TP309: *japonica* cv. Taipei 309; HEUK: *japonica* cv. Heukgyeong; NIPP: *japonica* cv. Nipponbare. (B, C) Scatter plot of differentially expressed highly conserved (B) and lowly conserved (C) miRNAs. A generalized Poisson-regression linear model was used to identify the differentially expressed miRNAs for the factors of development, subspecies, and development-by-subspecies interaction. MiRNAs with fold change ≥ 2 and FDR ≤ 0.05 are denoted as significantly differentially expressed. Mature miRNAs that show no differential expression (black) or show significant differential expression between subspecies (green), developmental stage (blue) and both (red) are indicated by circles in different colors, while those with differential expression for the additional factor of development-by-subspecies interaction are indicated by crosses with the same color setting.

Figure 2. Expression variation of the highly conserved and lowly conserved miRNAs between subspecies or developmental stages. (A) The proportions of differentially expressed miRNAs in both sets of the highly conserved and lowly conserved miRNAs. A generalized Poisson-regression linear model was used to identify the differentially expressed miRNAs for the factors of development, subspecies, and development-by-subspecies interaction. MiRNAs with fold change ≥ 2 and FDR ≤ 0.05 are denoted as significantly differentially expressed. MiRNAs that show significantly differential expression between subspecies or interactions are enriched in the lowly conserved miRNAs (Fisher's Exact Test, P -value $\ll 0.01$). (B) Fold changes in expression of the highly conserved and lowly conserved miRNAs between subspecies or developmental stages. The lowly conserved miRNAs exhibit significantly more variation in expression than the highly conserved miRNAs for both comparisons (Kolmogorov-Smirnov test, P -value $\ll 0.01$).

Figure 3. Gene ontology (GO) and KEGG pathway enrichment analyses of DEGs. The DEGs (FDR ≤ 0.05) with a fold change larger than 2 or 1.5 were used for the enrichment analyses of GO terms and KEGG pathways, respectively. The significantly over-represented and under-represented GO terms (A) and KEGG pathways (B) with a FDR ≤ 0.05 were presented. Grey and black bars indicate the percentages of DEGs and the whole transcriptome that were classified into different functional annotations, respectively.

Figure 4. Correlation between the coexpressed miRNAs and their targets in seedlings. (A) highly conserved miRNAs and their predicted targets (390 pairs); (B) lowly conserved miRNAs and their predicted targets (219 pairs); (C) miRNAs and the degradome-verified targets in target set IV (Zhou et al. 2010) (68 pairs) and (D) miRNAs and the degradome-verified targets in target set II (Wu et al. 2009) (49 pairs). The log₂ fold changes of miRNA or mRNA expression between rice *indica* and *japonica* subspecies in seedlings were used for Pearson's correlation analysis.

Figure 5. Permutation of miRNA-mRNA target relationships at the lineage level. (A) The empirical distribution of the Pearson's correlation coefficient values for 436 miRNA-mRNA pairs between expression levels of 96 miRNAs and those of their target mRNAs across 6 lineages. (B) The histogram plot represents the distribution of the global mean correlation values for the expression levels of all miRNA-mRNA pairs for 1000 permutations, (C) for the highly conserved miRNAs and (D) for the lowly conserved miRNAs. The black arrowhead indicates the true value.




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5