

# **Transfer reactions with Be-Li isotopes near the drip-line**

*LISE Workshop 2024*

**M. Lozano-González, A. Matta, B. Fernández-Domínguez,  
J. Lois-Fuentes, F. Delaunay**  
on behalf on the E748 collaboration


IGFAE-USC and LPC-Caen

# Overview of the exotic Be-Li region

Be and Li isotopes close to the neutron drip line have been extensively studied due to their exotic properties.

Two prime examples can be showcased:

**$^{11}\text{Li}$**  is a neutron-rich nuclei displaying a 2n **halo** structure.




$^{10}\text{B}$	$^{11}\text{B}$	$^{12}\text{B}$	$^{13}\text{B}$	$^{14}\text{B}$
$^9\text{Be}$	$^{10}\text{Be}$	$^{11}\text{Be}$	$^{12}\text{Be}$	$^{13}\text{Be}$
$^8\text{Li}$	$^9\text{Li}$	$^{10}\text{Li}$	$^{11}\text{Li}$	$^{12}\text{Li}$
$^7\text{He}$	$^8\text{He}$	$^9\text{He}$	$^{10}\text{He}$	

# Overview of the exotic Be-Li region

Be and Li isotopes close to the neutron drip line have been extensively studied due to their exotic properties.

Two prime examples can be showcased:

<sup>11</sup>Be presents parity inversion: g.s. has **positive** parity when negative expected.




I. Talmi and I. Unna, PRL 4 (1960).

<sup>10</sup> B	<sup>11</sup> B	<sup>12</sup> B	<sup>13</sup> B	<sup>14</sup> B
<sup>9</sup> Be	<sup>10</sup> Be	<sup>11</sup> Be	<sup>12</sup> Be	<sup>13</sup> Be
<sup>8</sup> Li	<sup>9</sup> Li	<sup>10</sup> Li	<sup>11</sup> Li	<sup>12</sup> Li
<sup>7</sup> He	<sup>8</sup> He	<sup>9</sup> He	<sup>10</sup> He	

## Recently gathered information

During the MUST2 @ RIKEN campaign, an unexpected **reduction** of the cross-section was observed in  $^9\text{Li}(d, ^3\text{He})^8\text{He}$  reactions.




A. Matta et al., PRC 92 (2015).

Possible **explanations**:

- Role of the many-body interactions.

## Recently gathered information

During the MUST2 @ RIKEN campaign, an unexpected **reduction** of the cross-section was observed in  ${}^{9,11}\text{Li}(d, {}^3\text{He}){}^{8,10}\text{He}$  reactions.



GMF of neutron w.f for  ${}^{12}\text{Be}(d, {}^3\text{He}){}^{11}\text{Li}$


Possible **explanations**:

- Role of the many-body interactions.
- Overestimation of the nuclear overlap  $\langle {}^{9,11}\text{Li} | {}^{8,10}\text{He} \rangle$ .

**Collect** more  
 $d\sigma/d\Omega$  data!


# Reactions to be studied

**E748** at GANIL during the MUST2@LISE campaign. Neutron and proton removal reactions from  $^{10,12}\text{Be}$  beams have been performed to probe key nuclei:



# Experimental setup for E748


Traditional **solid target** experiment @ D6. Below a sketch of the setup:



# Analysis at a glance

A **common** procedure is employed in all the reactions. Different gates are applied in a sequential manner, as follows:

1. **Heavy ID:** Only distinction in  $Z$ : separation of Be from Li residuals, but not along isotopic chain.






# Analysis at a glance

A **common** procedure is employed in all the reactions. Different gates are applied in a sequential manner, as follows:

2. **Light ID:** Using only stopped particles in Si layer, but low TOF resolution. Separation of **t-<sup>3</sup>He** attained with kinematics!



Missing mass technique:

$$E_{\text{beam}} + (E, \theta)_{\text{Lab}} \rightarrow \mathbf{E}_{\mathbf{x}}$$



# Elastic: $^{10}\text{Be}(d, d)^{10}\text{Be}$

Serves as a test of the analysis, allowing us to ascertain the **normalization** factors  $N_t$  and  $N_b$ .





- Modern models (Haixia and DA1p) adjust better the minimum.
- Overall agreement in magnitude.

Error in **efficiency**

Proton **contamination** at  
low  $E$

# Elastic: $^{10}\text{Be}(d, d)^{10}\text{Be}$

Cross-section for the 1st excited state is also achievable.





- DWBA: potential deformed in both Coulomb and nuclear parts
- Using  $B(E2)$  from other experiments
- **Same** error as before?

To be further investigated.  
 $\Rightarrow C^2S = 0.270(22)$   
with Daehnick

# Neutron removal: $^{10}\text{Be}(d, t)^9\text{Be}$

Only the **ground state** is accessible. Angular distributions are determined in the interval  $\theta_{\text{CM}} \in [5, 20]^\circ$ .




- **DWBA** with DAEHNICK (d) and PANG (t) OMPs
- **Best fit** is  $\ell = 1$  with  $j = 3/2$

$$\Rightarrow C^2S = 1.522(44)$$

## Neutron removal: $^{10}\text{Be}(d, t)^9\text{Be}$

Another measurement is available in D.L Auton Nucl. Phys. A (1970). A reanalysis with **our model** is executed:




- No errors could be extracted from paper
- Poor quality at large  $\theta_{CM}$
- $C^2S = 1.951(54)$

Agreement with our results


# Proton removal: $^{10}\text{Be}(d, ^3\text{He})^9\text{Li}$

**E748** can be compared with a recent experiment carried out at the Acculinn facility. For the  $E_x$ :

*Our experiment @ 30 A MeV*



*E. Y. Nikolskii et al. @ 42 A MeV*




*Recently published: NIMPR B 541 (2023)*

# Proton removal: $^{10}\text{Be}(d, ^3\text{He})^9\text{Li}$


Angular distributions for the **ground state** are extracted:

*Our experiment,  $\theta_{\text{CM}} \in [6, 14]^\circ$*



Again  $\ell = 1 \implies 3/2^-$ .  
 $C^2S = 1.80(11)$

*Acculinnane one,  $\theta_{\text{CM}} \in [3, 13]^\circ$*




Original publication:  
 $C^2S = 1.74$


# Proton removal: $^{10}\text{Be}(d, ^3\text{He})^9\text{Li}$

Angular distributions for the **ground state** are extracted:

*Our experiment,  $\theta_{\text{CM}} \in [6, 14]^\circ$*



*Reanalysis of Acculinnä's data*





Again  $\ell = 1 \implies 3/2^-$ .  
 $C^2S = 1.80(11)$

$\ell = 1 \implies C^2S = 2.679(45)$   
Investigate **input parameters**  
in the models!



# Proton removal: $^{10}\text{Be}(d, ^3\text{He})^9\text{Li}$



A first excited state is also accesible.



- **Best fit** is  $\ell = 1$
- Assuming  $j = 1/2$
- Spectroscopic factor: 0.185(36)

# Elastic: $^{12}\text{Be}(d, d)^{12}\text{Be}$

Yet another validation method of the normalization.





- Same behaviour as for  $^{10}\text{Be}$
- Normalization fine for this beam also

Clearly the same  
*systematic* error

# Elastic: $^{12}\text{Be}(d, d)^{12}\text{Be}$


Cross-section for the 1st excited state is also achievable.



- Same procedure as before



To be further investigated.  
 $\Rightarrow C^2S = 0.519(30)$  with Daehnick

# Challenging channels: $^{12}\text{Be}(d, t)^{11}\text{Be}$



- Strong **inhibition** of the ground state
- Why? Related to parity inversion in  $^{11}\text{Be}$ ?

# Challenging channels: $^{12}\text{Be}(d, ^3\text{He})^{11}\text{Li}$



- Low cross-section
- Subject to **contamination**: hard to disentangle  $A = 3$

$\Rightarrow C^2S = 0.510(85)$  with Daehnick

## Conclusions and outlook

We investigated several proton and neutron pick-up reactions on  $^{10,12}\text{Be}$ :

	Channel	Status	Pending
$^{10}\text{Be}$	(d,d)	Normalization OK	Requires study
	(d,t)	Completed	$C^2S$ matches other measures.
	(d, $^3\text{He}$ )	Completed	Two new $C^2S$
$^{12}\text{Be}$	(d,d)	Normalization OK	Same as $^{10}\text{Be}$
	(d,t)	Puzzled	?
	(d, $^3\text{He}$ )	Needs clean	New $C^2S$

# Conclusions and outlook

## Future prospects

Solve discrepancies in **elastic** channels

Reduce contamination in  $^3\text{He}$  PID

Compare with **shell model** calculations

**Thanks for your attention!**


And special thanks to the **E748** collaboration.




**Not part of the talk!**

## CsI on or off?

So far, studied excited states are *compressed* in the DSSD layer:




For  $^{10}\text{Be}(d, t)^9\text{Be}$



For  $^{10}\text{Be}(d, ^3\text{He})^9\text{Li}$


# Beam ID

Using Caviar to CATS1 TOF and energy loss in IC




*For  $^{10}\text{Be}$*

# Kinematic lines



For  $^{10}\text{Be}$  beam



For  $^{12}\text{Be}$  beam