



IGFAE

INSTITUTO GALEGO
DE FÍSICA
DE ALTAS ENERXÍAS

25 → 1999
2024

$\nu 0p_{1/2} - \nu 0p_{3/2}$ spin-orbit splitting in ^{20}O


M. Lozano-González, B. Fernández-Domínguez, J. Lois-Fuentes,
T. Roger, F. Delaunay

IGFAE-USC, GANIL and LPC-Caen

EuNPC 2025 - Caen

A recap on the SO splitting

Introduced by M. Goeppert-Mayer, reproduces magic numbers for stable nuclei.



SO splitting is mainly a surface effect:

$$V_{SO} = -\frac{1}{\hbar^2} V_{so} (\vec{l} \cdot \vec{s}) \left(\frac{1}{r} \frac{dV}{dr} \right)$$


which yields a ℓ -depending gap:

$$\Delta_{SO} = \frac{\hbar^2}{2} (2\ell + 1) \xi$$

⇒ Expected to evolve towards more exotic nuclei, where surface blurs

A recap on the SO splitting


G .Mairle et al. (PLB 304 (1993)) found systematic trends easily parametrizable.



Proton-neutron
interactions drive **shell**
evolution

Deviations from the trend are found due to:

- Loosely bound orbitals
- Nuclear matter depletion (^{35}Si)
- Role of **tensor force**





T.Otsuka and Y. Tsunoda, JPG 43 (2016)

Physics case

E796 to measure **transfer** reactions probing single-particle occupancies in ^{20}O .

1. Proton removal $^{20}\text{O}(\text{d}, ^3\text{He})^{19}\text{N}$ to investigate persistence of **Z = 6**





Tensor V_{pn} reduces $Z = 6$ gap as neutrons are added to $v0d_{5/2}$

Physics case

E796 to measure **transfer** reactions probing single-particle occupancies in ^{20}O .

2. Neutron removal $^{20}\text{O}(\text{d},\text{t})^{19}\text{O}$ to extract **N = 6 SO gap**




Would the gap decrease in ^{20}O ?



Can we extract magnitude of tensor force?

Experimental setup

E796 @ LISE in 2022. First transfer experiment with ACTAR TPC!





Silicon sizes:
 $80 \times 50 \times 0.5 \text{ mm}^3$



Gas mixture:
 $90\% \text{ D}_2 + 10\% \text{ iC}_4\text{H}_{10}$
at 952 mbar

A window to the analysis

Intricate analysis to extract reactions of interest out of noisy data.



Unique advantages from the TPC:

- Precise **vertex** determination
- Improved ΔE corrections


- Factor 10 in target number
- Implicit PID with ΔE_{gas}

A window to the analysis

Two steps are needed after a binary reaction has been identified.


1. PID of tritons by plotting

$$\Delta E_{\text{gas}} \text{ vs } \Delta E_{\text{Sil}}$$




Masked punch-through to 2nd front layer

2. E_x reconstructed by the **missing-mass** technique




Results: E_x spectrum



- 11 observed states
- At $E_x > 15$ MeV (**p,d**) contamination appears

- 1n phase space considered:
 $^{19}\text{O} \Rightarrow ^{18}\text{O} + n$
- 2n phase space is negligible



Results: E_x spectrum



T = 3/2 states @ $E_x < 10$ MeV

T = 5/2 at $E_x > 10$ MeV, based on comparison with $^{20}\text{O}(\text{d}, \text{He})^{19}\text{N}$

Results: cross-sections





DBWA with Fresco

- OMP:
 - ²⁰O + d: Daehnick
W. W. Daehnick et al. PRC 21 (1980)
 - ¹⁹O + t: Pang
D.Y. Pang et al. PRC 79 (2009)
- ⟨ d | t ⟩ from ab-initio GFMC
I. Brida et al., PRC 84 (2011)
- ⟨ ²⁰O | ¹⁹O ⟩ from standard WS

- g.s: 5/2⁺, taking up 71% of the occupancy
- 1st: 1/2⁺, with 8% of 1s_{1/2} occupancy

Results: cross-sections



Based on shell-model calculations (see next slide):

- $E_x = 3.1$ and 4.6 MeV $\Rightarrow 0p_{1/2}$
- $E_x = 6.7, \dots, 8.9$ MeV $\Rightarrow 0p_{3/2}$

T = 3/2 states:


- $0p_{1/2}$: 61 % of strength
- $0p_{3/2}$: just 14 % of occupancy!

Results: comparison with models

SFO-tls interaction

T. Suzuki, T. Otsuka PRC 78 (2008)

- For p -sd neutron-rich nuclei
- **Modified**: reduced tensor $\nu\nu$ and $\nu\pi$ monopole matrix el.
- C²S reduced wrt SFO-tls
- Great reproduction of low-lying states
- 0p_{3/2} less fragmented than predicted



Results: comparison with models


SFO-tls interaction

T. Suzuki, T. Otsuka PRC 78 (2008)

- For p -sd neutron-rich nuclei
- **Modified**: reduced tensor $\nu\nu$ and $\nu\pi$ monopole matrix el.

- C²S reduced wrt SFO-tls
- Great reproduction of low-lying states

- 0p_{3/2} less fragmented than predicted
- T = 3/2 0p_{3/2} predicted at much higher E_x



Results: centroids


Modified SFO-tls:

- Excellent agreement for $0d_{5/2}$, $1s_{1/2}$ and $0p_{1/2}$
- $0p_{3/2}$ shifted towards high E_x

$0p_{1/2}$ – $0p_{3/2}$ SO gap:

- Exp: 3.79(9) MeV
 - Theo: 5.64 MeV
- ⇒ Gap is reduced by ~ 1.8 MeV!

Systematic error of 20% from OMPs variety has been included in error bars



No $0p$ vacancies were observed through $^{20}\text{O}(\text{d},\text{p})^{21}\text{O}$
B. Fernández-Domínguez et al. PRC 84 (2011)

Conclusions

$^{20}\text{O}(\text{d},\text{t})^{19}\text{O}$ reaction as a means to measure SO gap
in exotic O isotopes


DWBA analysis to extract spectroscopic factors and
 E_x centroids for $T=3/2$ states

Comparison with SFO-tls interaction overestimates
SO gap due to $0p_{3/2}$ states

Theoretical efforts needed to reconcile those states
with experimental data!

Results: gap evolution


- $^{16}\text{O}(\text{d},\text{t})$: *K. H. Purser et al. NPA 132 (1969)*. No need to reanalyze xs bc there is only one state per nlj; just take the Ex
- $^{16}\text{O}(\text{d},\text{p})$: Alleged state $3/2^-$ is not *single-particle* but j-forbidden stripping. See *K. Hosono JPSP 25 (1968) Table II*. This state is neutron $0\text{d}_{5/2}$ + proton $0\text{p}_{1/2}^{-1}0\text{d}_{5/2}^1$ as I understood from it
- $^{18}\text{O}(\text{d},\text{t})$: *G. Mairle et al. NPA 280 (1977)*. Reanalysis of xs with our OMPs and Fresco. Major discrepancies with their paper
- $^{20}\text{O}(\text{d},\text{t})$: this experiment



We are lacking theo calculations for ^{16}O and ^{18}O

Conclusions

- To be determined



Acknowledgements



T. Roger
J. Pancin
M. Fisichella
C. Nicole
F. Saillant
G. Wittwer
V. Morel
A. Cassisa
J.C. Thomas
O. Sorlin
L. Cáceres
C. Stodel
F. de Oliveira



J. Lois-Fuentes
B. Fernández-Domínguez
M. Caamaño
D. Fernández
D. Regueira
C. Cabo
H. Álvarez-Pol
Y. Ayyad
G. Mantovani



F. Delaunay
L. Achouri



J. Giovannazzo
A. Ortega-Moral
S. Grevy
Q. Delignac
T. Kurtikian



R. Raabe
O. Poleschuk
A. Ceuleman
S. Fracassetti
M. Latif



A. M. Martínez
J. Dueñas



D. Suzuki
B. Mauss



Istituto Nazionale di Fisica Nucleare

M. Pellegretti
T. Marchi

Extra slides

A window to the analysis

