



INSTITUTO GALEGO
DE FÍSICA
DE ALTAS ENERXÍAS

25 → * 1999
2024

Quenching of spectroscopic factors in $^{10,12}\text{Be}(d, ^3\text{He})$ reactions

M. Lozano-González, A. Matta, B. Fernández-Domínguez

IGFAE and LPC-Caen

Zakopane 2024 Conference



Cofinanciado por
la Unión Europea



Fondos Europeos


A recap on spectroscopic factors

Spectroscopic factors arise from the breakdown of the single-particle scheme to describe nuclear reactions:

$$\sigma = C^2 S \cdot \sigma_{s.p}$$

- **Long-range** correlations: vibrations, giant resonances,...
- **Short-range:** tensor forces,...


Reduction of
~65 %!



T. Aumann et al. Prog. Part. Nucl. Phys. 118 (2021)

A long-standing puzzle

A trend with asymmetry $\Delta S \equiv S_n - S_p$ is found depending on the experimental **probe**!




T. Aumann et al. Prog. Part. Nucl. Phys. 118 (2021)


⇒ measure towards more exotic nuclei:
 $^{10,12}\text{Be}$!

Importance of the overlaps

In a previous experiment by A. Matta and colleagues, for the $^{11}\text{Li}(d, ^3\text{He})^{10}\text{Be}$ reaction they found that the **geometrical mismatch factor** (GMF) plays an essential role.



A.Matta et al.



A.Matta et al.