



INSTITUTO GALEGO  
DE FÍSICA  
DE ALTAS ENERXÍAS


25 → \* 1999  
2024

# Quenching of spectroscopic factors in $^{10,12}\text{Be}(d, ^3\text{He})$ reactions

M. Lozano-González, A. Matta, B. Fernández-Domínguez,  
F. Delaunay, J. Lois-Fuentes

USC-IGFAE, LPC-Caen and FRIB

Zakopane 2024 Conference




# A recap on spectroscopic factors

**Spectroscopic factors** shed light on the occupancy of single-particle states:

$$\left. \frac{d\sigma}{d\Omega} \right|_{exp} = C^2 S \cdot \left. \frac{d\sigma}{d\Omega} \right|_{s.p}, \quad C^2 S = \begin{cases} (2j+1) & \text{removing} \\ 1 & \text{adding} \end{cases} \quad \text{in IPSM}$$

**Experimentally:**  
Reduction of  $\sim 65\%$ !


- **Short-range:** tensor forces,...
- **Long-range** correlations: vibrations, giant resonances,...



*L. Lapikás, Nuclear Phys. A 553 (1993)*

# A long-standing puzzle

A trend with asymmetry energy  $\Delta S \equiv S_n - S_p$  is found depending on the experimental **probe**!





*T. Aumann et al. Prog. Part. Nucl. Phys. 118 (2021)*

⇒ measure towards more exotic nuclei:  $|\Delta S| \uparrow$

# Status with light isotopes

Several experiments allowed for the extraction of  $C^2S$  with Li-induced (d,  $^3\text{He}$ ) reactions:



A. Matta et al., Phys. Rev. C 92 (2015)


Several challenges in this region:

1 Dealing with **unbound** nuclei ( $^{10}\text{He}$ )


2 Impact of core excitations (completar algo +)

# Importance of GMF

Towards exotic nuclei (loosely bound or halo), a **geometrical mismatch factor** emerges from the very different w.f. in the overlap:



A.Matta et al., Phys. Rev. C 92 (2015)



N. K. Timofeyuk, private communication (in E748 proposal)


⇒ Need to establish more systematics for this parameter

# Physics case of E748

E748 @ GANIL back in 2017. Using  $^{10,12}\text{Be}(d, ^3\text{He})$  reactions to:


Overlaps:

- $\langle ^{10}\text{Be} | ^9\text{Li} \rangle, \Delta S = -12.8 \text{ MeV}$
- $\langle ^{12}\text{Be} | ^{11}\text{Li} \rangle, \Delta S = -19.8 \text{ MeV}$




Explore effects of GMF:

- $\langle ^{10}\text{Be} | ^9\text{Li} \rangle, \text{GMF} \sim 1$
- $\langle ^{12}\text{Be} | ^{11}\text{Li} \rangle, \text{GMF} \sim 0.5?$




# Experimental setup

## Traditional solid target experiment @ LISE




# A glance at the analysis

## 1 Heavy ID at 0°



## 2 Light PID in DSSD





## 3 $E_x$ from missing mass technique

$$E_{beam} + (E, \theta)_{Lab} \rightarrow E_x$$




# Results: $^{10}\text{Be}(d, ^3\text{He})^9\text{Li}$




# Results: $^{10}\text{Be}(d, ^3\text{He})^9\text{Li}$

Recent experiment @ Acculina. Different beam energy of 40 A MeV





Their analysis:  
 $C^2S = 1.74$



Our  $C^2S$ :

- Pang: 2.679(48)
- HT1p: 1.848(33)

# Results: $^{12}\text{Be}(d, ^3\text{He})^{11}\text{Li}$



# Conclusions

Ola  
A ver  
Que  
Poñemos  
Aqui

# Acknowledgments

## The E748 collaboration:

- Santiago:  
B. Fernández
- LPC-Caen:  
A. Matta  
F. Delaunay  
N. L. Achouri  
F. Flavigny  
J. Gibelin  
M. Marques  
N. Orr
- IJCLab:  
D. Beaumel  
M. Assié  
Y. Blumenfeld  
S. Franchoo  
A. Georgiadou  
V. Girard-Alcindor  
F. Hammache  
N. de Séreville  
A. Meyer  
I. Stefan
- GANIL:  
B. Jacquot  
O. Kamalou  
A. Lemasson  
M. Rejmund  
T. Roger  
O. Sorlin  
J.C. Thomas  
M. Vandebrout  
B. Bastin  
F. de Oliveira  
C. Stodel
- RIKEN:  
S. Koyama  
D. Suzuki
- Surrey:  
N. Timofeyuk

