Beyond the static Einstein–Euler–Heisenberg description of an extreme electric field: a formally verified validity map for discharge and trapping

Álvaro López Almeida
(Dated: November 2, 2026)

We ask what lies beyond the point where the static, quartic-truncated Einstein–Euler–Heisenberg (EEH) description of a point-like electric charge breaks down ($\varepsilon \equiv eE/m_e^2 \gtrsim 1$, $r \lesssim \lambda_C$), and whether QED can act as a "universal quantum censor" that prevents the gravitational collapse of the field. Every equation used is derived or checked with a computer algebra system (112 gates, 24 of them negative controls that must fail on altered models). We find: (i) the exact one-loop (proper-time) real part does not saturate the field; it reproduces Reissner–Nordström with a running charge, whereas the quartic truncation and a matched Born–Infeld model yield a spurious single Schwarzschild-like horizon; the dominant-energy condition is violated for $\varepsilon >$1.88218 with relative margin $\to -\alpha/3\pi$ (the one-loop trace anomaly). (ii) The condition $\theta_+ = 0 \Leftrightarrow 2GM_{\rm MS} = R$, the decomposition $M_{\rm ADM} = M_{\rm MS}(R) + U_C(R)$ and the exact cancellation of the electrostatic energy are derived inside the pipeline for time-symmetric spherical data, together with the exact one-loop correction to U_C. (iii) A Pauli-blocked Markov kinetic model keeps $0 \le f \le \frac{1}{2}$ and gives $\omega_p =$0.186 $e\sqrt{eE_0}$; replacing the Markov ansatz by the exact Bloch–Bogoliubov equations coupled to Maxwell's equation (the normalization of the coupling is fixed by a Landau–Zener test) lets the occupation reach $f_{\max} =$1.0000, advances the first field zero by 19 per cent, raises the plasma frequency to $\omega_p =$0.194 $e\sqrt{eE_0}$, and shows that the pairs form a collinear gas with $\langle w_\parallel \rangle =$0.99 rather than $\frac{1}{3}$, with energy conserved to 6.2×10^{-12}. (iv) The ratio of the discharge time to the free-fall time obeys $\tau_{\rm dis}/\tau_{\rm ff} = (r_0^*/r_0)^{1/2}$ with r_0^* independent of the number of charges N, and a quantitative (N, r_0) phase map is given. (v) Without time symmetry, the Misner–Sharp balance $\partial_t M_{\rm MS} = -4\pi r^2 F_{\rm flux}$ is derived and the internal field-to-plasma conversion is shown to leave $M_{\rm MS}$ unchanged; a 1+1D solver of the discharge of a charged ball in general relativity shows that the plasma leaving through the surface removes only a fraction 0.50 of the interior field energy while the ingoing half of the plasma converted outside adds 2.5 times more, so the net effect is a gain; the dynamics shifts the critical mass by less than 0.8 per cent, and the horizon boundary is set instead by the static interior field energy, which places the shell curve $r_s(N)$ inside an initial apparent horizon for $N >4.6 \times 10^4$. Several ingredients are known results that we re-derive and verify; Sec. VI separates them from what is specific to this work, and Sec. VII states the regimes where the model is not controlled.

I. INTRODUCTION

The Euler–Heisenberg (EH) effective action [1–3] coupled to Einstein gravity has been used to study QED corrections to charged black holes in its quartic truncation [4, 5]. Its validity requires $\varepsilon \ll 1$, and for a point charge $Q = Ne$ the static field reaches $\varepsilon = 1$ at $r_\varepsilon = \sqrt{N\alpha}\,\lambda_C$, long before any horizon unless $N \gtrsim N_{\rm crit} = M_{\rm Pl}^2/m_e^2 \simeq 5.71 \times 10^{44}$. Beyond that point two qualitatively different effects matter: nonperturbative vacuum pair creation, which discharges the field on plasma time scales [6–10], and the gravitational response, which depends on whether a trapped surface can exist at all. The loss of charge by black holes through a Schwinger-type mechanism goes back to Gibbons [11], and the structure of charged thin-shell collapse to Israel, Boulware and Bekenstein [12, 13].

Aim and scope. We do not claim a new law of nature. We build a reproducible, formally gated computation that (a) quantifies where the quartic EEH extrapolation is qualitatively wrong, (b) re-derives the trapping criterion without invoking external theorems, (c) extends the pair-plasma back-reaction through several field reversals with Pauli blocking, and (d) assembles the resulting scales in one phase map. The pipeline (Appendix A) executes symbolic derivations (SymPy), adaptive numerical integrations with independent cross-checks, and negative controls; it fails if any gate fails. All numbers quoted below are read automatically from the pipeline output.

II. STATIC NONLINEAR ELECTRODYNAMICS COUPLED TO GRAVITY

A. Reduction for an arbitrary $\mathcal{L}(E)$

For the static spherical ansatz $ds^2 = -f\,dt^2 + dr^2/g + r^2 d\Omega^2$, $A = \Phi(r)\,dt$, a purely electric field $E = \sqrt{g/f}\,\Phi'$ and any Lagrangian $\mathcal{L}(X)$, $X = F_{\mu\nu}F^{\mu\nu} = -2E^2$, the stress tensor derived from the full tensor expression obeys

$$T^t{}_t = T^r{}_r = -\rho, \quad \rho = E\mathcal{L}' - \mathcal{L}, \quad T^\theta{}_\theta = \mathcal{L}, \tag{1}$$

$$r^2\mathcal{L}'(E) = Q/4\pi, \qquad f = g = 1 - \frac{2Gm(r)}{r}, \quad m' = 4\pi r^2\rho, \tag{2}$$

and $\rho' = -(2/r)(\rho + p_\perp)$ holds identically (gates S1–S4; the control C1 with $p_\perp = \mathcal{L}/2$ is detected). For Maxwell, $\rho = E^2/2$; for the quartic EH term, $\mathcal{L} = E^2/2 + 4cE^4$ with $c = \alpha^2/90\,m_e^4$.

B. One-loop proper-time Lagrangian and strong-field behaviour

For purely electric fields the spinor one-loop correction is

$$\mathcal{L}_1(x) = -\frac{1}{8\pi^2}\int_0^\infty \frac{ds}{s^3} e^{-s}\Big[xs\cot(xs) - 1 + \tfrac{(xs)^2}{3}\Big], \quad x = eE, \tag{3}$$

evaluated on the ray $s = \rho e^{-i\pi/4}$ because the poles of cot lie on the real axis. The real part is the effective Lagrangian; the imaginary part reproduces the Schwinger series $\mathrm{Im}\,\mathcal{L}_1 = \frac{x^2}{8\pi^3}\sum_n n^{-2}e^{-n\pi/x}$ to a relative error of 8.5×10^{-15} (gate N1), and for $\varepsilon \ll 1$ it matches $4cE^4$ with $\mathcal{L}_1/(4cE^4) - 1 \propto \varepsilon^{2.00}$ (N2; the Taylor coefficient is checked symbolically, S6).

At strong field the expansion is logarithmic, in agreement with the known renormalization-group structure [14, 15]:

$$\mathcal{L} = \frac{E^2}{2}\Big[1 - \frac{\alpha}{3\pi}\big(\ln\varepsilon + 3K\big)\Big], \quad K = -0.7640, \tag{4}$$

where the coefficient is verified symbolically (S7) and the slope $d(\mathcal{L}_1/x^2)/d\ln x = -4.22171597\times10^{-3}$ agrees with $-1/24\pi^2 = -4.22171599\times10^{-3}$ (N3). The pole of Eq. (4) sits at $\log_{10}\varepsilon_L = 561.9$, i.e. $\mu_L = \sqrt{eE} \approx 10^{277.7}$ GeV, consistent with the usual Landau scale of QED, whereas the Planck density corresponds to $\log_{10}\varepsilon_{\rm Pl} = 44.4$. This holds *at one loop*: the Ritus–Narozhny conjecture, that each loop in a strong field costs $\alpha\chi^{2/3}$ and perturbation theory fails for $\chi \gtrsim \alpha^{-3/2} \simeq 1604$ [16, 17], would invalidate all one-loop statements far below ε_L; its applicability to a pure electric field is not established here.

C. Energy conditions and the trace anomaly

WEC and NEC hold for the one-loop Lagrangian up to $\varepsilon = 10^{40}$ (the minima of ρ/E^2 and ED/E^2 are 0.465 and 0.930, gate N7), and $\mathcal{L} > 0$ (SEC) until the pole (N8). The dominant energy condition $\rho \geq |p_\perp|$ requires $ED - 2\mathcal{L} \geq 0$, and SymPy establishes the identity (gate S10)

$$ED - 2\mathcal{L} = x\mathcal{L}_1' - 2\mathcal{L}_1 = x^3\frac{d}{dx}\Big(\frac{\mathcal{L}_1}{x^2}\Big). \tag{5}$$

Hence the DEC is violated exactly where $\mathcal{L}_1/x^2$ decreases, i.e. for $\varepsilon > 1.88218$ (root found with the exact integral, not the interpolation). The relative margin $(\rho - |\mathcal{L}|)/\rho$ tends to $-\alpha/3\pi = -7.743\times10^{-4}$ and keeps growing slowly with $\ln\varepsilon$ (-7.875×10^{-4} at $\varepsilon = 10^{10}$) up to the pole: it is *not* bounded. This is the one-loop trace anomaly $T^\mu{}_\mu \propto \beta F^2$ [18, 19] written in terms of ρ and $p_\perp$, not a new quantum violation. For nonlinear electrodynamics of Plebański class, causality implies the DEC [20]; the corresponding causality conditions for the full $\mathcal{L}_1(S,P)$ have not been evaluated here, so we do not draw a conclusion on (a)causality.

D. Interior profiles

Figure 1 compares the metric function and the field for $N = 10^6$ and $M = 1.5M_{\rm ext}$. The quartic extrapolation gives $E \propto r^{-2/3}$ and a finite interior mass (gate S9), and a Born–Infeld model matched to the quartic term ($b^2 = 1/32c$, S5; field bounded by $\varepsilon = 69.6$) yields the same single horizon at $r = 2GM$, $r_h = 1.07263\times10^{-17}\,\lambda_C$. The exact one-loop result retains two horizons, $r_- = 1.48259\times10^{-18}$ and $r_+ = 9.24904\times10^{-18}$ (Reissner–Nordström: 1.36569×10^{-18} and 9.36055×10^{-18}), consistent with an effective charge $Q^2 \to \lambda Q^2$, $\lambda = E/E_{\rm Coul}(r_+) = 1.0710$: the measured shift $\delta r_+/r_+ = -1.191\times10^{-2}$ agrees with the running-charge prediction -1.230×10^{-2} (N6). The static nonlinearity therefore does not censor the horizon. These profiles are *formal solutions of the locally-constant-field model*: for $r \sim r_Q$ the field varies on a scale $\ell \simeq r_Q/2 \sim 1.8\times10^{-18}\,\lambda_C \ll \lambda_C$, outside the derivative expansion (Sec. VII).

III. TRAPPING CRITERION DERIVED INSIDE THE PIPELINE

The statement that the electrostatic energy does not help to trap a charged shell is standard in terms of the Misner–Sharp mass [12, 21, 22]; the charged Penrose inequality is a known theorem under stated hypotheses [23, 24]. To remove any dependence on citations we derive what we use (gate `P10_misner_sharp_symbolic`, 16 sub-checks).

(a) Geometry. In double-null coordinates, $ds^2 = -2e^{2\sigma}du\,dv + R^2d\Omega^2$, with $M_{\rm MS} = (R/2G)(1 - g^{ab}\partial_a R\,\partial_b R)$ and expansions $\theta_\ell = 2\partial_u R/R$, $\theta_n = 2\partial_v R/R$, SymPy proves

$$1 - \frac{2GM_{\rm MS}}{R} = -\frac{R^2e^{-2\sigma}}{2}\,\theta_\ell\theta_n. \tag{6}$$

Hence $\theta_+ = 0 \Leftrightarrow 2GM_{\rm MS} = R$ (for $\theta_- \neq 0$), and a trapped surface is $2GM_{\rm MS} > R$.

(b) Constraint. For time-symmetric data ($K_{ij} = 0$) the Hamiltonian constraint follows from computing $G_{ab}n^an^b = 2Gm'/r^2$ for a metric with arbitrary second time derivatives, which drop out; thus $m' = 4\pi r^2\rho$. The radial Maxwell field obeys Gauss' law $r^2E_{\rm orth} = $ const independently of the metric and $T_{ab}n^an^b = E_{\rm orth}^2/2$.

(c) ADM decomposition. Integrating from R to infinity,

$$M_{\rm ADM} = M_{\rm MS}(R) + U_C(R), \quad U_C(R) = \frac{Q^2}{8\pi R}, \tag{7}$$

so the exterior metric *is* Reissner–Nordström, $g = 1 - 2GM_{\rm ADM}/r + GQ^2/4\pi r^2$ (derived, not assumed).

(d) Cancellation. At $\theta_+ = 0$, $GM_{\rm ADM} = (R + r_Q^2/R)/2$ with $r_Q^2 = GQ^2/4\pi$ (the equality case of the charged Penrose inequality). With $M_{\rm MS}(R) = M_n$ the electrostatic




FIG. 1. Metric function $g(r) = 1 - 2Gm(r)/r$ (left) and electric field relative to Coulomb (right) for $N = 10^6$ charges and $M = 1.5M_{\rm ext}$, in the exact one-loop model, Maxwell (Reissner–Nordström), the quartic Euler–Heisenberg truncation extrapolated beyond its validity, and a Born–Infeld model matched to the quartic term.

term cancels identically: $G(M_n + U_C) - \frac{1}{2}(R + r_Q^2/R) = GM_n - R/2$. Non-charged exterior matter with $\rho \geq 0$ only adds mass, giving the inequality.

Scope. Proven in this subsection: spherical time-symmetric data, $\rho \geq 0$, no charged matter outside the surface. The dynamical case ($K_{ij} \neq 0$) is treated in Sec. III A; the persistence of trapped surfaces under NEC is cited, not proven. The control C4 verifies that wrong factors in $M_{\rm MS}$, in the constraint ($m' = 2\pi r^2\rho$) or in U_C are detected.

One-loop correction to U_C. For a general $\mathcal{L}$ with $H(D) = ED - \mathcal{L}$ and $dH/dD = E$ (proved for arbitrary $\mathcal{L}$), $\rho = H(D)$, $D = a/r^2$, $a = Q/4\pi$, and

$$U(R) = 2\pi a^{3/2} \int_0^{D_R} H(D)\, D^{-5/2} dD, \quad D_R = a/R^2, \tag{8}$$

which gives $Q^2/8\pi R$ for Maxwell and $-16\pi c\, a^4/5R^5$ for the quartic term (symbolic). With the exact one-loop H and $N = 10^6$, $U_{\rm QED}/U_C - 1 = 7.047 \times 10^{-2}$ at $R = r_Q$, in agreement with a direct radial quadrature to 8.9×10^{-14} (gate P11). The slope with respect to $\ln \varepsilon$ is 7.8527×10^{-4} ($\ln \varepsilon \in [9, 23]$) and 8.1870×10^{-4} ($[18, 60]$), approaching $\alpha/3\pi = 7.7427 \times 10^{-4}$ with the $O(\alpha \ln \varepsilon)$ sub-leading logarithm.

A. Dynamical Misner–Sharp balance without time symmetry

For the general metric $ds^2 = -e^{2\phi(t,r)}dt^2 + e^{2\lambda(t,r)}dr^2 + r^2 d\Omega^2$ and $M_{\rm MS} = (r/2G)(1 - e^{-2\lambda})$, SymPy computes the full Einstein tensor and proves $G^t{}_t = -2GM_r/r^2$ and $G^r{}_t = 2GM_t/r^2$, i.e.

$$\partial_r M_{\rm MS} = 4\pi r^2 \rho, \qquad \partial_t M_{\rm MS} = 4\pi r^2 T^r{}_t = -4\pi r^2 F_{\rm flux}, \tag{9}$$

together with $K^r{}_r = -e^{-\phi}\partial_t \lambda = -4\pi G r e^{2\lambda - \phi} T^r{}_t$ and $K^\theta{}_\theta = 0$ (D_MS1): the slices have $K_{ij} \neq 0$ if and only if there is an energy flux, so the data of gate P10 are exactly the zero-flux case. For a radial electric field (the Maxwell tensor is built from $F_{tr} = -e^{\phi+\lambda}E$, with $T^t{}_t = T^r{}_r = -E^2/2$, $T^\theta{}_\theta = E^2/2$ and $T^r{}_t = 0$) SymPy verifies $\nabla_\mu T^\mu{}_\nu{}^{(\rm EM)} = F_{\nu\lambda} J^\lambda$ in the dynamical metric, with $J^\nu \equiv \nabla_\mu F^{\mu\nu}$, and that its t-component is $-\partial_t(E^2/2)$. If the plasma obeys $\nabla_\mu T^\mu{}_\nu{}^{(\rm pl)} = -F_{\nu\lambda} J^\lambda$, the two Einstein constraints $G^r{}_t$ and $G^r{}_r - G^t{}_t$ give

$$\partial_t \rho_{\rm tot} = \frac{1}{r^2}\partial_r\big(r^2 T^r{}_t\big) \quad \text{independently of } F_{t\lambda} J^\lambda, \tag{10}$$

so the internal conversion $E^2/2 \to \rho_{\rm pl}$ in $r < R$ conserves $M_{\rm MS}(t, R)$ exactly and $\partial_t M_{\rm MS}(t, R) = 4\pi R^2 T^r{}_t(R)$ depends only on the plasma flux through $r = R$ (D_MS2); if the transfer law is altered to $(1+\delta)F_{t\lambda}J^\lambda$ a spurious volume source $4\pi\delta \int_0^R r^2 F_{t\lambda} J^\lambda dr$ appears (control C_MS). For a boundary $r = R(t)$, $dM_{\rm MS}/dt = 4\pi R^2(T^r{}_t + \dot{R}\rho)$ vanishes for comoving matter, so only the flux *relative* to the surface matters (D_MS3). Radial null streams $T^{\mu\nu} = \rho_\pm \ell_\pm^\mu \ell_\pm^\nu$ propagate with $v = \alpha/a = e^{\phi-\lambda}$ and, with $W_\pm = r^2\rho_\pm$,

$$\partial_t W_+ + \partial_r(vW_+) = -W_+\big[v(\phi_r + \lambda_r) + 2\lambda_t\big], \tag{11}$$
$$\partial_t W_- - \partial_r(vW_-) = +W_-\big[v(\phi_r + \lambda_r) - 2\lambda_t\big], \tag{12}$$

with $\lambda_t = -\frac{1}{2}a^2 v(W_+ - W_-)/r$ and $\phi_r = \frac{1}{2}a^2 r p_r + (a^2 - 1)/2r$ ($8\pi G = 1$ in these two expressions), where both the t and r components of $\nabla_\mu T^{\mu\nu} = 0$ give the same equation (D_MS4). Together with the conversion (field loses what the plasma gains) these equations preserve the momentum constraint, $r^2\partial_t\rho_{\rm tot} - \partial_r(r^2 T^r{}_t) = 0$ identically (D_MS5); a statically "held" mass $\rho_{\rm mat}$ with $p = 0$ violates it by the exact amount $4\pi G r^4 \rho_{\rm mat} v(\rho_- - \rho_+)/(r - 2GM)$, which is why M_n is treated below as a test mass.

Initial data of a charged ball. For a uniformly charged ball of radius r_0 with non-electromagnetic mass M_n, with $q \equiv r_Q/r_0$ and $C_n \equiv 2GM_n/r_0$, the compactness profile outside the ball is $C(x) = [C_n + \frac{6}{5}q^2 - q^2/x]/x$, $x = r/r_0$ (interior field energy $q^2/5$, exterior q^2), whose maximum is $(C_n + \frac{6}{5}q^2)^2/4q^2$. SymPy gives an apparent horizon already in the data when $C_n = 0$ for $q \geq 1.6667 = 5/3$, i.e. $r_0 \leq 0.60\, r_Q$ (P12): this is the extremality bound $M_{\rm ADM} \geq Q$, and the discharge cannot undo it (NEC). The interior field energy of a ball, which does gravitate at the surface, is absent from the shell criterion $C_n = 1$ used in Sec. V.

IV. PAIR-PLASMA BACK-REACTION

A. Markov source up to the first zero

For a homogeneous field $x(t) = eE(t)$ with local pair-creation rate $\Gamma(x) = x^2 e^{-\pi/x}/4\pi^3$ (the $n = 1$ Schwinger term, which is the *mean pair number*; the vacuum-decay rate $2\,\mathrm{Im}\mathcal{L}_1$ contains the full series), conduction and polarization currents give

$$\dot{x} = -2e^2\Big[\int_0^t dt'\,\Gamma(x')\langle v\rangle + \Gamma(x)\frac{\langle m_\perp\rangle}{x}\Big], \qquad (13)$$

with $p(t,t') = \int_{t'}^t x$, $v = p/\sqrt{p^2+m_\perp^2}$ and $m_\perp^2 = 1 + xs/\pi$ averaged over $s \sim \mathrm{Exp}(1)$. This is the Markov limit of the quantum Vlasov equation [8, 25]. The first zero of the field occurs at $\tau_{\rm dis} = \kappa/(e\sqrt{eE_0})$ with $\kappa = 13.5394$ and exponent -0.50001 (the exponent follows from scale invariance for $m \to 0$, so only κ carries information; it is specific to this model). The scheme is second order (observed order 1.98) and an independent implicit-trapezoid solver agrees ($\tau_0 = 1.41658$ vs 1.41658). As an order-of-magnitude check against a secondary report of Ref. [9], our first zero is 17.1 at $E_0 = 10E_c$ and 8.8×10^6 at $0.15E_c$ (in units of $\hbar/m_e c^2$), against ~ 10 and $\sim 10^7$ quoted there for an oscillation length; the definitions are not identical, so this is consistency, not validation.

B. Pauli-blocked kinetics through repeated field reversals

Removing the restriction to the first zero requires tracking occupation. We evolve $f(P, p_\perp, t)$ on a canonical-momentum grid with kinetic momentum $p = P + \Phi(t)$, $\dot{\Phi} = x$:

$$\partial_t f = (1-2f)\,S, \quad S = |x|e^{-\pi m_\perp^2/|x|}\,G_\sigma(p), \qquad (14)$$

$$\dot{x} = -2e^2\Big[\sum W f\frac{p}{\omega} + \sum W\omega\,\frac{(1-2f)S}{x}\Big], \qquad (15)$$

with $\rho_{\rm pl} = 2\sum W f\omega$ and a Gaussian of width $\sigma = 0.25\sqrt{\varepsilon_0}$. SymPy proves that

$$\frac{d}{dt}\Big(\frac{x^2}{2e^2} + \rho_{\rm pl}\Big) = 0 \qquad (16)$$

holds identically *only if* the polarization current carries the factor $(1-2f)$ (K1; the control C5 yields the residual $-4\omega S f \neq 0$ otherwise), and that the blocked equation has the exact solution $f = \frac{1}{2}(1 - e^{-2\int S}) \in [0, \frac{1}{2})$ (K2).

Results (Fig. 2). Using an adaptive DOP853 integrator with zero-crossing detection over $\varepsilon_0 \in \{30, 10^2, 10^3, 10^4\}$ we obtain at least 10 zero crossings (five full oscillations) with $\max|\Delta E|/E_0 = 2.9 \times 10^{-6} < 10^{-4}$ (K5); $0 \leq f \leq 0.500000$ everywhere (K4) whereas without blocking f reaches $6.53 > 1$; without the blocking factor the model reproduces the cohort model of the previous subsection (K3, $\tau_0 = 4.55758$ vs 4.55709), while blocking lengthens the first zero to 5.5684 at $\varepsilon_0 = 100$. The plasma frequency scales as $\omega_p \propto \varepsilon_0^{0.501}$ with $\omega_p = 0.1865\, e\sqrt{eE_0}$ (K7), and the amplitude decays slowly, $\gamma/\omega_p = 0.0052$ (the amplitude falls from 0.73 to 0.62 of E_0 in nine half-cycles). Changing the tolerance, grid and transverse quadrature alters τ_0 by at most 1.5×10^{-5} (K6); varying the source width within $[0.15, 0.40]\sqrt{\varepsilon_0}$ changes it by 5.0×10^{-4}. The negative controls (no $(1-2f)$ in $j_{\rm pol}$: $|\Delta E|/E_0 = 5.4 \times 10^{-2}$ against 2.3×10^{-8} for the consistent model; $(1+2f)$: $f = 135 \gg \frac{1}{2}$; wrong conduction sign) are all detected.

Limitation. A Markov equation with $S \geq 0$ cannot raise f above $\frac{1}{2}$ from the vacuum, whereas the exact constant-field occupation $e^{-\pi m_\perp^2/\varepsilon}$ exceeds $\frac{1}{2}$ for $\varepsilon > 4.5$ and small $p_\perp$; for a single crossing the blocked rule gives $f = \frac{1}{2}(1 - e^{-2f_0})$, which is only 0.46 of the exact value at $f_0 = 0.9$. The quantitative values of ω_p and γ of this subsection are therefore properties of the Markov model; Sec. IV C replaces the ansatz by the exact kinetics [25, 26] and quantifies the difference.

C. Exact non-Markovian quantum kinetics

The Markov limit is removed by evolving, for every mode $(P, s{=}p_\perp^2)$ with kinetic momentum $p = P + \Phi(t)$, $\dot{\Phi} = x$ and $m_\perp^2 = 1 + s$, the Bloch vector of the exact two-level Dirac problem $H = p\,\sigma_z + m_\perp\sigma_x$, with $f = (1-z)/2$, u and v the components of the Bloch vector along the adiabatic axes (Bloch–Bogoliubov form of the quantum Vlasov equation [8, 25]):

$$\dot{f} = \tfrac{1}{2}\lambda\,u, \quad \dot{u} = \lambda(1-2f) - 2\omega v, \quad \dot{v} = 2\omega u, \qquad (17)$$

$$\dot{x} = -2e^2\sum W\Big[\frac{p}{\omega}f + \frac{m_\perp}{2\omega}u\Big], \quad \lambda = \frac{x\,m_\perp}{\omega^2}, \qquad (18)$$

with $\omega = \sqrt{p^2+m_\perp^2}$, $\rho_{\rm pl} = 2\sum W\omega f$ and the parallel pressure $P_\parallel = \sum W p\,[2(p/\omega)f + (m_\perp/\omega)u]$ (Hellmann–Feynman). There is no source term, no locally-constant-field approximation and no source width: phase coherence between successive crossings and Pauli blocking are contained exactly. SymPy derives Eq. (17) from $\dot{\mathbf{n}} = 2\mathbf{h}\times\mathbf{n}$ together with the velocity and energy identifications (Q0), proves that $\frac{d}{dt}[(1-2f)^2 + u^2 + v^2] = 0$ identically, so that the vacuum $(0,0,0)$ implies $0 \leq f \leq 1$ without any truncation at $\frac{1}{2}$ (Q1), proves $\frac{d}{dt}(x^2/2e^2 + \rho_{\rm pl}) = 0$




FIG. 2. Left: electric field $\varepsilon(t)/\varepsilon_0$ in units of $1/(e\sqrt{\varepsilon_0})$ for the Pauli-blocked Markov kinetic equation (solid) and without blocking (dashed), exhibiting the collapse of different ε_0 onto one curve (scale invariance). Right: $\omega_p/(e\sqrt{\varepsilon_0})$ and γ/ω_p as functions of ε_0.

with the polarization current of Eq. (18) (Q2; omitting it leaves the residual $Wx\,m_\perp u/\omega \neq 0$, control C10), and the trace identity $\rho_{\rm pl} - P_\parallel - 2P_\perp = \sum W m^2(2f - pu/m_\perp)/\omega$ (Q4).

Normalization of the coupling. The form $\dot f = \lambda u$, $\dot u = 2\lambda(1-2f) - 2\omega v$ (the coupling of Eq. (17) doubled) is not excluded by the algebraic gates: it also conserves purity and energy (with a correspondingly doubled polarization current), so these gates cannot discriminate it, but it fails the Landau–Zener test: for $eE = 10$, $m_\perp = 1$ it gives $f_\infty = 0.2942$ instead of $e^{-\pi m_\perp^2/eE} = 0.7304$ (control C11). The coefficient of Eq. (17) is the one derived from the Dirac equation (Q0) and reproduces $f_\infty = e^{-\pi m_\perp^2/eE}$ to 4.1×10^{-7} or better in all 8 test cases (Q3; the C kernel gives 2.1×10^{-6}, Q3b), with f_∞ up to 0.969, above the Markov ceiling $\frac12$ for $\varepsilon > \pi/\ln 2 = 4.5324$ (Q3a).

Numerics. A fixed-step RK4 scheme in C/OpenMP (time step $0.2/\omega_{\max}$, P-spacing $0.04\sqrt{\varepsilon_0}$, 30 Gauss–Legendre nodes in $p_\perp^2$) integrates 39450 modes at $\varepsilon_0 = 100$. The discrete system conserves $x^2/2e^2 + \rho_{\rm pl}$ exactly, and the integrator preserves it to 6.2×10^{-12} over at least five oscillations for $\varepsilon_0 \geq 10$ (Q2b). Halving ΔP or Δt, doubling the transverse nodes or widening the P range changes τ_0 by $\leq 1.7\times10^{-8}$ and the first four half-periods by $\leq 3.4\times10^{-3}$ (K6); a narrow P range is detected by the coverage gate (C12). The initial state is the zeroth-order adiabatic vacuum at $t = 0$, as in the Markov model; replacing it with the first-order superadiabatic vacuum changes the first half-periods by 1.5×10^{-2} (K8).

Results (Table I, Fig. 3). The exact system is not capped at $\frac12$: $f_{\max} = 0.999996$ at $\varepsilon_0 = 100$ and ≥ 0.99997 for every $\varepsilon_0 \geq 10$ (Q1c), against 0.5000 for the blocked Markov model; even at $\varepsilon_0 = 4 < \pi/\ln 2$ interference between successive crossings reaches $f_{\max} = 0.9990$. For a single crossing the blocked Markov rule undershoots by the fraction 0.536 at $f_0 = 0.9$ (limit 0.568 for $f_0 \to 1$). The first zero of the field occurs earlier than in the Markov model: $\tau_0 e\sqrt{\varepsilon_0} = 13.44$ at $\varepsilon_0 = 10^3$ against 16.67, a ratio 0.806 (0.851 at $\varepsilon_0 = 10$, 0.899 at $\varepsilon_0 = 4$). The exact value is within 0.8 per cent of the constant $\kappa = 13.54$ of the unblocked source of Sec. IV, so the lengthening attributed above to Pauli blocking is an artefact of the ceiling at $\frac12$. The plasma frequency (mean of the last three half-periods) scales as $\omega_p \propto \varepsilon_0^{0.505}$ with $\omega_p/(e\sqrt{\varepsilon_0}) = 0.1944$ for $\varepsilon_0 = 10^3$, i.e. 4.3 per cent above the Markov value 0.1864 (K7). The amplitude decays slowly and non-exponentially (it falls from 0.78 to 0.67 of E_0 in six half-cycles), and the log-linear fit gives $\gamma/\omega_p = 0.0048$ against 0.0052; with ten amplitudes this number is a rough descriptor and the difference is not significant. *Equation of state.* The total parallel $w_{\rm tot} = (P_\parallel - E^2/2)/(\rho_{\rm pl} + E^2/2)$ starts at -1 (pure field) and first reaches zero at $t\,e\sqrt{\varepsilon_0} = 6.23$ (6.57 at $\varepsilon_0 = 30$), before the first field zero at 13.4. The pairs themselves have $w_\parallel^{\rm pl} = P_\parallel/\rho_{\rm pl} > \frac13$ from the first instant ($t\,e\sqrt{\varepsilon_0} \simeq 0.020$) and its average over the last period is 0.988 (at most 1.054 instantaneously, the excess over unity being the coherent term): it tends to the collinear value 1, not to the isotropic $\frac13$, because the pairs are created with $p_\perp^2 \sim \varepsilon/\pi \ll p_\parallel^2$ and the homogeneous model has no collisions or transverse spreading that could isotropize them. *Controls and sensitivity.* Without the polarization current the energy drifts by 4.1×10^{-2} (C13); with the doubled coupling (and a consistent doubled current) energy is conserved but $\tau_0 e\sqrt{\varepsilon_0}$ changes to 15.18 (C14), so only Landau–Zener fixes the coefficient. The initial state is the zeroth-order adiabatic vacuum; the first-order superadiabatic vacuum changes τ_0 by 0.015 of its value (K8).

TABLE I. Exact (Bloch–Bogoliubov) versus Markov kinetics at the same ε_0: maximum occupation, first zero $\tau_0 e\sqrt{\varepsilon_0}$, plasma frequency $\omega_p/(e\sqrt{\varepsilon_0})$ (mean of the last three half-periods), γ/ω_p (log-linear fit of the half-cycle amplitudes), and the time-averaged parallel equation of state of the pairs over the last period. For $\varepsilon_0 = 4$ only four zero crossings were computed.

ε_0	$f_{\max}^{\mathrm{ex}}$	$f_{\max}^{\mathrm{M}}$	τ_0^{ex}	τ_0^{M}	ω_p^{ex}	ω_p^{M}	$\gamma/\omega_p^{\mathrm{ex}}$	$\gamma/\omega_p^{\mathrm{M}}$	$\langle w_\parallel^{\mathrm{pl}}\rangle$
4	0.99902	0.47109	21.29	23.69	0.1236	0.1169	0.0140	0.0167	0.991
10	1.00000	0.50000	16.17	19.01	0.1790	0.1740	0.0047	0.0057	0.988
30	1.00000	0.50000	14.28	17.38	0.1907	0.1824	0.0044	0.0053	0.988
100	1.00000	0.50000	13.67	16.86	0.1947	0.1853	0.0047	0.0052	0.988
300	0.99999	0.50000	13.50	16.72	0.1945	0.1861	0.0045	0.0052	0.988
1000	0.99997	0.50000	13.44	16.67	0.1944	0.1864	0.0048	0.0052	0.988




FIG. 3. Exact non-Markovian quantum Vlasov (solid) versus the Pauli-blocked Markov model (dashed). Top left: field oscillations at $\varepsilon_0 = 10$ and 100. Top right: maximum occupation f; the exact system exceeds the Markov ceiling $\frac{1}{2}$ above $\varepsilon = \pi/\ln 2$. Bottom left: plasma frequency. Bottom right: parallel equation of state $w_{\mathrm{tot}} = (P_\parallel - E^2/2)/(\rho_{\mathrm{pl}} + E^2/2)$ (solid) and $w_\parallel^{\mathrm{pl}} = P_\parallel/\rho_{\mathrm{pl}}$ (dotted).

V. SCALES AND PHASE MAP

For N electrons of charge Ne confined in radius r_0 (lengths in λ_C): the field reaches $\varepsilon = 1$ at $r_\varepsilon = \sqrt{N\alpha}$; the Reissner–Nordström radius is $r_Q = N\sqrt{\alpha}/M_{\mathrm{Pl}}$; the Coulomb density equals M_{Pl}^4 at r_P (with a floor of $10\,l_P$); and the trapping radius $r_s(N)$ solves $r = 2G\,[N + \frac{3}{4}(9\pi/4)^{1/3}N^{4/3}/r]$ (rest mass plus ultra-relativistic Fermi energy, gate P4). Because of the cancellation of Sec. III, r_s is independent of the charge. The ratio of the electromagnetic to the gravitational force of the electron is $z^2 = \alpha M_{\mathrm{Pl}}^2 = 4.17 \times 10^{42}$.

Time scales. With $E_0 = Q/4\pi r_0^2$, $\varepsilon_0 = N\alpha/r_0^2$ and the result of Sec. IV, $\tau_{\mathrm{dis}} = \kappa\, r_0/(\alpha\sqrt{4\pi N})$, equal to the light-crossing time at $N^* = \kappa^2/(4\pi\alpha^2) = 2.74 \times 10^5$. For the free fall of a dust ball of mass Nm_e, $\tau_{\mathrm{ff}} = \frac{\pi}{2}\sqrt{r_0^3/2GM}$ (symbolic, P8). Both times scale as $N^{-1/2}$,

so

$$\frac{\tau_{\rm dis}}{\tau_{\rm ff}} = \sqrt{\frac{r_0^*}{r_0}}, \qquad r_0^* = \frac{2\kappa^2}{\pi^3\alpha^2 M_{\rm Pl}^2} = 9.29\times10^{-18}\, l_P, \tag{19}$$

independent of N (gate P9). We stress that this closed form is the product of two power laws and of the electron's z^2; it only expresses that discharge is faster than free fall for every $r_0 > l_P$ (since $r_0^* \ll l_P$ the value of r_0^* itself lies outside the validity of either model). On the trapping boundary with the Fermi energy included, the discharge wins only for $N \gtrsim 1.11\times10^5$.

Phase map (Fig. 4). Four regions: static EFT ($\varepsilon < 1$); Schwinger discharge ($r_P < r_0 < r_\varepsilon$ and untrapped); trapping ($r_0 \le r_s$); and trans-Planckian. The trapped region always lies at $\varepsilon \gg 1$ and above the Planck floor for $N \gtrsim 15$; a classical horizon enters the static-EFT region only for $N \gtrsim \alpha M_{\rm Pl}^4/4 = 5.94\times10^{86}$. The Coulomb density at r_Q is $\rho/M_{\rm Pl}^4 = 1/(8\pi\alpha N^2)$ (P6), sub-Planckian for $N \gtrsim 2.3$. The "trapping" region $r_0 \le r_s$ is the shell criterion $2GM_n \ge r_0$ (matter surface with no interior field energy); Sec. V A shows that for $N > N_A \simeq 4.6\times10^4$ this curve lies inside the region $r_0 \le 0.6\, r_Q$ in which the initial data of a uniformly charged ball already contain an apparent horizon. Table II lists the scales. The electrons in region (ii) are not bound (the Coulomb energy exceeds the Fermi energy for $N \gtrsim 6.0\times10^3$); r_s is a condition on initial data, not a formation scenario.

A. Spatially resolved discharge and trapping

We evolve the system of Sec. III A on a radial grid in polar–areal gauge (script `06b_dynamical_trapping_1d`; units $r_0 = c = 1$, $q = r_Q/r_0$, $s = \sqrt{N/N_*}$). The field of a uniformly charged ball, $\tilde\rho_{\rm EM} \propto q^2 r^2$ ($r < 1$) and q^2/r^4 ($r > 1$), is converted locally with the clock $\dot s_c = \alpha\,\kappa s\sqrt{E/E_{\rm surf}}$ and the universal first-oscillation curve of the exact kinetics (Sec. IV C): the field decays as $E = E_{\rm geom}\, g(s_c)$ until its first zero, the energy removed is shared equally between an outgoing and an ingoing null stream (zero net momentum; the collinear plasma of Sec. IV C has $w_\parallel \simeq 1$), and the metric follows from the constraints at every step. At the surface the discharge time is r_0/s (so $s = 1$ is $N = N_*$). The held mass M_n enters as a static test mass. The flat limit with instantaneous discharge has a closed form, including the passage of ingoing rays through the centre: the solver reproduces $E_{\rm out}$ and $E_{\rm in}$ to 1.3×10^{-5} and 1.7×10^{-6} with observed order 2.00 (N1; omitting the centre reflection gives an error of 3%, control C_N1). In the full theory the balance $m(r_0,t) - m(r_0,0) = \int(F_{\rm in} - F_{\rm out})dt$ of Eq. (9) holds to 1.0×10^{-3} relative (2.4×10^{-5} absolute) in 10 cases with q up to 1.3 (N2); a plasma that receives 1.2× what the field loses breaks it by 4.0×10^{-2} (C_N2). Grid convergence with Δr, $\Delta r/2$, $\Delta r/4$ is 5.5×10^{-4} at most (N3).

How much mass-energy leaves, and how much arrives. Take weak gravity ($q = 0.2$) and instantaneous discharge, evaluated at $t = r_0/c$ (perfectly radial streams form a central caustic after that time; beyond it, up to the free-fall time $\frac{\pi}{2}r_0$ of the trapping boundary, a reflecting core of radius $r_{\rm core} = 0.1\, r_0$ stands in for the angular momentum of the rays, $\theta_\perp \simeq \sqrt{1-w_\parallel} \simeq 0.11$ from Sec. IV C; it conserves energy exactly, $|\Delta m - \int F| \le 2.4\times10^{-5}$, changes ΔC by 0.049 for $r_{\rm core} = 0.05 \to 0.2$ at $q = 0.5$ and by 3.4×10^{-8} at $q = 1$, N10). A neutral matter element released from rest follows the radial geodesic of the evolving metric; in a point-mass test its proper fall time agrees with the exact cycloid to 3.5×10^{-5} (N9), which validates α, a, ϕ_r and λ_t used for the dynamics. The interior channel (field energy inside the ball converted into plasma that crosses r_0 outwards) removes $\Delta C = -0.0997\, q^2$ from $C = 2GM_{\rm MS}(r_0)/r_0$, i.e. a fraction 0.497 of the interior field energy (0.422 for $s = 1$) and 0.083 of the total field energy U_C. The exterior channel (field energy outside the ball converted in place, whose ingoing half enters r_0) adds $0.2481\, q^2$, 2.49 times larger, so the net effect is a *gain*, $\Delta C = +0.1493\, q^2$: the quasilocal mass at the surface is increased, not decreased, by the discharge. The net gain depends on the discharge speed as $\Delta C \simeq q^2 F_\infty s^p/(s^p + s_h^p)$ with $F_\infty = 0.1493$, $p = 1.61$, $s_h = 1.62$ (N4, N8): it is the instantaneous-discharge value for $N \gg N_*$ and falls as s^p for $N \ll N_*$, where the plasma escapes ballistically before the field is converted ($0.0065\, q^2$ at $s = 0.3$). Gravity suppresses the inflow: at $q = 1$, $\Delta C/q^2 = 0.096$ (instantaneous) and at $q = 1.5$ it is 0.0004, and a held mass $C_n = 0 \to 0.6$ lowers ΔC at $q = 1$, $s = 3$ from 0.0626 to 0.0011, because the coordinate speed $v = \alpha/a$ vanishes as $\theta_+ \to 0$ (N5).

Displacement of the trapping threshold. Just below the static horizon threshold of P12, at $C_n = C_n^{\rm AH}(q) - 0.05$ with $C_n^{\rm AH} = 1 - q^2/5$ for $q < 1$ and $2q - \frac{6}{5}q^2$ for $q \ge 1$, the maximum of the compactness profile grows by at most $\delta C_n = 0.0033$ until $t = r_0/c$ and 0.0080 until the free-fall time $\frac{\pi}{2}r_0$, over $q \in [0.3, 1.5]$ and $s \in [0.1, \infty)$ (0.0035 at $s = 1$); no case forms a new horizon, and the numerical initial compactness reproduces the closed form to 4.3×10^{-3} (N6). Therefore the dynamics shifts the critical non-electromagnetic mass, $\Delta M_{\rm crit}/M_{\rm crit}$, by at most 0.0080 (0.8%) within the free-fall time, and in no regime does the escaping plasma delay trapping: the effect that dominates the retained mass is the *static* interior field energy $q^2/5$, which the shell criterion of Sec. V omits. Figure 5 shows the resulting curves in the (N, r_0) plane: the horizon boundary $r_{\rm AH}(N)$ of the ball (P12 plus dynamics) lies above the shell curve $r_s(N)$ by a factor up to 1402 for $N \le 10^{14}$, the two cross at $N_A \simeq 4.62\times10^4$ (where $r_s = 0.6 r_Q$), and the dynamic correction is $r^{\rm dyn}/r_{\rm AH} - 1 \le 1.1\times10^{-3}$. For $N > N_A$ the curve $r_s(N)$ of the previous sections lies *inside* the region where the initial data already contain an apparent horizon; there the race between discharge and free fall (Eq. (19)) is moot, because discharge cannot remove a trapped surface that already exists.




FIG. 4. Phase map in the (N, r_0) plane. Solid lines: $r_\varepsilon = \sqrt{N\alpha}$ and $r_s(N)$; dashed: r_Q; dotted: Planck-density radius. Vertical lines mark $N_{\rm crit} = M_{\rm Pl}^2/m_e^2$ and N^* ($\tau_{\rm dis} = r_0$).

TABLE II. Characteristic radii (in λ_C) and ratios for selected N (computed in `06_phase_diagram.py`).

N	r_ε	r_Q	r_s	$\varepsilon(r_s)$	$\tau_{\rm dis}/r_0$
1.00×10^{0}	8.54×10^{-2}	3.58×10^{-24}	7.10×10^{-23}	1.45×10^{42}	5.23×10^{2}
1.00×10^{3}	2.70×10^{0}	3.58×10^{-21}	7.10×10^{-21}	1.45×10^{41}	1.66×10^{1}
1.00×10^{6}	8.54×10^{1}	3.58×10^{-18}	7.10×10^{-19}	1.45×10^{40}	5.23×10^{-1}
1.00×10^{12}	8.54×10^{4}	3.58×10^{-12}	7.10×10^{-15}	1.45×10^{38}	5.23×10^{-4}
1.00×10^{20}	8.54×10^{8}	3.58×10^{-4}	1.53×10^{-9}	3.12×10^{35}	5.23×10^{-8}
1.00×10^{30}	8.54×10^{13}	3.58×10^{6}	7.10×10^{-3}	1.45×10^{32}	5.23×10^{-13}
5.71×10^{44}	2.04×10^{21}	2.04×10^{21}	4.89×10^{7}	1.74×10^{27}	2.19×10^{-20}
1.00×10^{48}	8.54×10^{22}	3.58×10^{24}	7.10×10^{9}	1.45×10^{26}	5.23×10^{-22}
1.00×10^{60}	8.54×10^{28}	3.58×10^{36}	7.12×10^{17}	1.44×10^{22}	5.23×10^{-28}

VI. WHAT IS KNOWN AND WHAT IS SPECIFIC TO THIS WORK

VII. REGIMES WHERE THE MODEL IS NOT CONTROLLED

(1) *Higher loops.* All strong-field statements are one-loop; Ritus–Narozhny scaling may invalidate them for $\varepsilon \gtrsim 10^3$–10^6. The sweep of the kinetic model reaches $\varepsilon_0 = 10^{10}$, 6.8 orders above that threshold. (2) *Derivative expansion and curvature.* The profiles of Sec. II descend to $r \sim 10^{-18}\lambda_C$; the curvature couplings of the one-loop action [27] are of relative order $\alpha(\lambda_C/L)^2$ and are neglected, and the local-field Lagrangian is not controlled there. (3) *Imaginary part.* For $\varepsilon > 1$ the vacuum-decay rate is $O(1)$ and the real part alone is not a stable effective action. (4) *Kinetics.* The exact Bloch system of Sec. IV C has no LCFA and no $f \leq \frac{1}{2}$ cap, but it is homogeneous, has no collisions or spatial transport (the pairs do not isotropize: $\langle w_\parallel \rangle \simeq 1$), starts from the zeroth-order adiabatic vacuum (a 1.5×10^{-2} effect on the half-periods), uses a finite $(P, p_\perp)$ quadrature (converged to 3.4×10^{-3}), and only reaches $\varepsilon_0 = 10^3$; the Markov constants of Sec. IV are model-specific, and the first-zero scale κ of the cohort model (LCFA, Markov, $t_{\rm form}/\tau_0 \simeq e/\kappa = 0.022$) agrees with the exact kinetics only to the 0.8 per cent quoted in Sec. IV C. (5) *Dynamics of trapping.* The dynamical Misner–Sharp balance is proven (Sec. III A) and the 1+1D solver is validated (flat limit, balance, convergence, geodesic test), but the




FIG. 5. Spatially resolved discharge. Top left: retained compactness at r_0 for the interior (leak), exterior (inflow) and combined channels ($q = 0.2$, instantaneous discharge). Top right: dependence on $s = \sqrt{N/N_*}$ and fit. Bottom left: shift δC_n of the critical $2GM_n/r_0$ produced by the dynamics (solid: $t = r_0/c$; dashed: $t = \pi r_0/2$ with a reflecting core). Bottom right: boundaries in the (N, r_0) plane: shell criterion r_s, r_Q and $0.6\,r_Q$, apparent-horizon boundary of the static ball $r_{\rm AH}$ and the corrected curve $r^{\rm dyn}$ (indistinguishable from $r_{\rm AH}$ at this scale).

collapse race is not evolved self-consistently: the time available for trapping is the *prescribed* free-fall time $\frac{\pi}{2}r_0$ of the trapping boundary, and the matter M_n is held at r_0 as a static test mass (its constraint violation is 0.17–0.31 of ΔC in the cases tested); a moving charged shell would also need its radial stress (the lapse jumps across it) and the Coulomb force on its carriers, neither of which is modelled. The plasma is a set of radial null streams whose central caustic is regularized by a reflecting core ($r_{\rm core} = 0.1r_0$); the local discharge law is the homogeneous kinetics of Sec. IV C applied cell by cell, Gauss's law is not imposed during the conversion, and collisions, transverse spreading and the Coulomb explosion of the charge carriers are not included. The polar–areal gauge fails once an apparent horizon forms, so no evolution beyond that point is claimed. (6) *Landau and Planck.* The Landau pole and the Planck density are never crossed in any calculation; the Planck floor defines region (iii) and no statement is made there.

VIII. CONCLUSIONS

Beyond the point where the quartic static EEH description fails, the exact one-loop static field does not saturate and does not censor the horizon; the electrostatic energy *outside* the surface plays no role in the trapping of that surface, which for a shell requires non-electromagnetic mass, $GM_n \geq R/2$ (the interior field energy of a ball does gravitate, Sec. III A); and pair discharge converts the field into plasma on the time $\tau_{\rm dis} = \kappa/(e\sqrt{eE_0})$ while conserving energy-momentum, so it cannot remove mass or undo a trapped surface. In the exact homogeneous kinetics the resulting collinear pair plasma oscillates for many periods with $\omega_p \simeq 0.194\,e\sqrt{eE_0}$ (the Pauli-blocked Markov model gave 0.186), reaches occupations up to unity, and has $\langle w_\parallel \rangle \simeq 1$. What discharge does remove is the net charge whose repulsion exceeds gravity by $z^2 \sim 10^{42}$, and it does so faster than free fall. These conclusions are consistent with the classical picture of Misner–Sharp, Boulware and Gibbons; the added value of this work is a verified quanti-

TABLE III. Attribution of the results. "Known" means that the mathematical structure was published before; "here" lists what the pipeline adds.

Result	Prior work	Contribution of this work
Coulomb energy cancels in trapping	Misner–Sharp mass with charge, thin-shell energy equation, charged Penrose inequality [12, 21–24]	Derivation inside a CAS under explicit hypotheses (time-symmetric data, $\rho \geq 0$, no exterior charge); one-loop-exact $U_C(R)$
Plasma oscillations, Markov source, $\omega_p \propto e\sqrt{eE}$	[6–10, 25]	Exact energy identity with $(1-2f)$ in $j_{\rm pol}$, $f \leq \frac{1}{2}$, constants ω_p, γ for this source; convergence and negative controls
Bloch–Bogoliubov quantum Vlasov with Maxwell back-reaction	[8, 25, 26]	Same equations (a rediscovery); here: symbolic Dirac→Bloch derivation, a Landau–Zener test that fixes the normalization of the coupling, a like-for-like Markov–exact comparison (τ_0, ω_p, γ, $f_{\max}$) and the parallel equation of state; not compared with published curves
Strong-field logarithm, Landau scale	[14, 15]	Exact evaluation of the electric proper-time integral and the constant K
DEC violation	One-loop trace anomaly [18, 19]	Exact onset ε =1.88218 and identity $x^3(\mathcal{L}_1/x^2)'$
Dynamical Misner–Sharp equations, quasilocal flux balance, radial null-dust transport in polar–areal gauge	[21, 22] (standard)	Derivation in a CAS for the general $\phi(t,r), \lambda(t,r)$ metric with a Maxwell field and an exchange term, negative control for the transfer law; the conclusion that field→plasma conversion inside R conserves $M_{\rm MS}$ and only the surface flux counts is classical
Channel decomposition (leak from inside, inflow from outside) and its dependence on q and N/N_*	none found; radiation collapse of null shells is classical	Quantitative numbers for this specific discharge closure; not a prediction (closure ansatz, test-mass matter, radial beams)
Discharge of charged black holes, $N_{\rm crit}$	Gibbons [11]	Reformulation of the scale in the (N, r_0) plane
EH black holes (quartic)	[4, 5]	Quantitative comparison with the exact one-loop interior in the coupled system, with explicit limits of validity
$\tau_{\rm dis}/\tau_{\rm ff} = (r_0^*/r_0)^{1/2}$, phase map	Elementary combination of known scalings	Closed forms with κ, verified symbolically; organization in one plane

tative map of where each description holds, together with an open, reproducible pipeline. Collisional isotropization, a comparison with published non-Markovian curves and the evolution of the matter and of the plasma beyond the first light-crossing time (central caustic, formation of the horizon) remain open. For the spatial and gravitational gap, the complete Misner–Sharp balance with $K_{ij} \neq 0$ is derived, the discharge of a charged ball is simulated, and the result is that the dynamics moves the horizon threshold by less than 0.8 per cent up to the prescribed free-fall time while the static interior field energy, absent from the shell criterion, relocates the threshold to $r_0 \simeq 0.6\, r_Q$ for large N.

ACKNOWLEDGMENTS

The symbolic derivations, code and first draft of the text were produced with an AI assistant (Claude, Anthropic) operating the open pipeline described in Appendix A; the author is responsible for the content. Bibliographic details marked in `references.bib` must be verified against the original publications before submission.

Appendix A: Verification pipeline and reproducibility

The repository (https://github.com/lopezalmeidaalvaro/experimento) contains numbered scripts executed by `run_all.sh`, which stops at the first failing gate. Symbolic gates assert simplify(LHS − RHS) = 0; numerical gates include convergence under refinement and an independent method; negative controls assert that altered signs, factors or conservation laws are detected. The totals are 112 gates, of which 24 are negative controls. Table IV lists the gates by script.

[1] W. Heisenberg and H. Euler, Folgerungen aus der diracschen theorie des positrons, Z. Phys. **98**, 714 (1936).

[2] V. Weisskopf, Über die elektrodynamik des vakuums auf grund der quantentheorie des elektrons, Kgl. Danske Vi-




FIG. 6. Phase map of Fig. 4 with the horizon boundary of the charged ball: the shaded region has an apparent horizon in the initial data for arbitrarily small M_n; the curve $r^{\rm dyn}(N)$ includes the discharge up to $t = \pi r_0/2$ and coincides with $r_{\rm AH}$ to 10^{-3}. For $N > N_A$ the shell curve r_s lies inside the shaded region.

TABLE IV. Verification gates by script (symbolic, numerical and negative controls).

Script	gates	neg. controls
`04_nonperturbative_qed`	22	3
`05_dynamical_backreaction`	13	4
`05b_plasma_oscillations`	12	4
`05c_quantum_vlasov_exact`	24	6
`06_phase_diagram`	20	4
`06b_dynamical_trapping_1d`	21	3
01–03 (baseline)		re-run at each build

densk. Selsk. Mat.-Fys. Medd. **14** (1936).
[3] J. Schwinger, On gauge invariance and vacuum polarization, Phys. Rev. **82**, 664 (1951).
[4] H. Yajima and T. Tamaki, Black hole solutions in euler-heisenberg theory, Phys. Rev. D **63**, 064007 (2001), gr-qc/0005016.
[5] R. Ruffini, Y.-B. Wu, and S.-S. Xue, Einstein-euler-heisenberg theory and charged black holes, Phys. Rev. D **88**, 085004 (2013), 1307.4951.
[6] Y. Kluger, J. M. Eisenberg, B. Svetitsky, F. Cooper, and E. Mottola, Pair production in a strong electric field, Phys. Rev. Lett. **67**, 2427 (1991).
[7] Y. Kluger, J. M. Eisenberg, B. Svetitsky, F. Cooper, and E. Mottola, Fermion pair production in a strong electric field, Phys. Rev. D **45**, 4659 (1992).
[8] Y. Kluger, E. Mottola, and J. M. Eisenberg, The quantum vlasov equation and its markov limit, Phys. Rev. D **58**, 125015 (1998), hep-ph/9803372.
[9] R. Ruffini, G. Vereshchagin, and S.-S. Xue, Electron-positron pairs in physics and astrophysics: from heavy nuclei to black holes, Phys. Rep. **487**, 1 (2010), 0910.0974.
[10] F. Gelis and N. Tanji, Schwinger mechanism revisited, Prog. Part. Nucl. Phys. **87**, 1 (2016), 1510.05451.
[11] G. W. Gibbons, Vacuum polarization and the spontaneous loss of charge by black holes, Commun. Math. Phys. **44**, 245 (1975).
[12] D. G. Boulware, Naked singularities, thin shells, and the reissner-nordström metric, Phys. Rev. D **8**, 2363 (1973).
[13] J. D. Bekenstein, Hydrostatic equilibrium and gravitational collapse of relativistic charged fluid balls, Phys. Rev. D **4**, 2185 (1971).
[14] V. I. Ritus, Lagrangian of an intense electromagnetic field and quantum electrodynamics at short distances, Zh. Eksp. Teor. Fiz. **69**, 1517 (1975).
[15] G. V. Dunne, Heisenberg–euler effective lagrangians: basics and extensions, in *From Fields to Strings: Circumnavigating Theoretical Physics* (World Scientific, 2005) hep-th/0406216.
[16] A. M. Fedotov, Conjecture of perturbative qed breakdown at $\alpha\chi^{2/3} \gtrsim 1$, J. Phys.: Conf. Ser. **826**, 012027 (2017), 1608.02261.
[17] A. Fedotov, A. Ilderton, F. Karbstein, B. King, D. Seipt, H. Taya, and G. Torgrimsson, Advances in qed with intense background fields, Phys. Rep. **1010**, 1 (2023), 2203.00019.
[18] S. L. Adler, J. C. Collins, and A. Duncan, Energy-momentum-tensor trace anomaly in spin-1/2 quantum electrodynamics, Phys. Rev. D **15**, 1712 (1977).
[19] J. C. Collins, A. Duncan, and S. D. Joglekar, Trace and dilatation anomalies in gauge theories, Phys. Rev. D **16**, 438 (1977).
[20] J. G. Russo and P. K. Townsend, Causality and energy conditions in nonlinear electrodynamics, J. High Energy Phys. **2024** (06), 191, 2404.09994.
[21] C. W. Misner and D. H. Sharp, Relativistic equations for adiabatic, spherically symmetric gravitational collapse, Phys. Rev. **136**, B571 (1964).
[22] S. A. Hayward, General laws of black-hole dynamics, Phys. Rev. D **49**, 6467 (1994).
[23] E. Malec and N. Ó Murchadha, Trapped surfaces and the penrose inequality in spherically symmetric geometries, Phys. Rev. D **49**, 6931 (1994), gr-qc/9401014.
[24] M. A. Khuri, G. Weinstein, and S. Yamada, Proof of the riemannian penrose inequality with charge for multiple black holes, J. Differential Geom. **106**, 451 (2017), 1409.3271.
[25] S. M. Schmidt, D. Blaschke, G. Röpke, S. A. Smolyansky, A. V. Prozorkevich, and V. D. Toneev, A quantum kinetic equation for particle production in the schwinger mechanism, Int. J. Mod. Phys. E **7**, 709 (1998), hep-ph/9809227.
[26] M. Prakapenia and G. Vereshchagin, Pauli blocking effects on pair creation in strong electric field (2023), arXiv:2306.12956.
[27] I. T. Drummond and S. J. Hathrell, Qed vacuum polarization in a background gravitational field and its effect on the velocity of photons, Phys. Rev. D **22**, 343 (1980).