Table of Contents

Environment	**Page**
1. Hydrothermal Vents	2
2. Diffuse Vent Zone	5
3. Abyssal Ocean	7
4. Ocean Connected Tiger Stripe Brines	9
5. Intrashell Mushy Fracture Brines	12
6. Plume Ice Grains - Type I	14
7. Plume Ice Grains - Type II (VOC)	16
8. Plume Ice Grains - Type II (HMOC)	19
9. Plume Ice Grains - Type III	21
References	24

1. **Hydrothermal Vents**
   Scope Definition: This site includes the focused, high-temperature fluid upwellings and actively reacting rock surfaces at the interface between Enceladus's porous silicate core and the overlying global subsurface ocean, explicitly excluding the cooler, bulk abyssal ocean and broader diffuse venting regions.

   a. Temperature (°C): 90 to 290
      Source: Models constrained by Cassini's detection of silica nanoparticles, indicate temperatures must exceed 90 °C to permit nanosilica formation (Hsu et al., 2015, Sekine et al., 2015). Other geochemical models estimate higher temperatures depending on the fluid's specific reaction path, with quartz-buffered fluids reaching up to 350 °C (Glein et al., 2018). The theoretical upper limit for fluid temperature before vaporization at the ocean floor is the boiling point of water at that depth, which is ~290 °C (Schenk et al., 2018). There is general agreement that the vents are hot, but exact maximum temperatures depend heavily on assumptions regarding the core's permeability and chemical openness.

      Reliability: 3 (High). This temperature range is robustly derived from primary Cassini spacecraft data linked directly to well-established, peer-reviewed hydrothermal experiments.

   b. pH: 8.0 to 10.5
      Source: Group A says the hydrothermal fluids have a moderately alkaline pH of 8.5 to 10.5 because these values are required to precipitate the silica nanoparticles observed by Cassini (Hsu et al., 2015, Sekine et al., 2015). Group B says the vent fluid pH could be lower, around 8.0 to 9.0, because pH is temperature-dependent and fluids tend to be less alkaline at high temperatures in a chemically closed system before cooling and mixing with the ocean (Sekine et al., 2015, Glein et al., 2018). A union range of 8.0 to 10.5 is adopted because it captures the necessary conditions for both the observed silica nanoparticles and the complex temperature-dependent water-rock interactions occurring at depth.

      Reliability: 2 (Medium). These pH values are derived from peer-reviewed thermodynamic models and experimental proxies rather than direct in-situ sampling of the vent fluids themselves.

c. Salinity (g/kg): 5 to 40
Source: The direct Cassini measurements of plume ice grains indicate a bulk ocean salinity of 5 to 20 g/kg, primarily composed of sodium chloride and sodium carbonates (Postberg et al., 2009, Glein et al., 2018). Other experimental sources state that the formation of the observed silica nanoparticles sets a strict upper salinity limit of 40 g/kg at the vents, as higher salinities would cause rapid aggregation (Hsu et al., 2015). The models agree that the venting fluids are moderately saline rather than highly concentrated brines.

Reliability: 3 (High). The salinity limits are tightly constrained by primary mass spectrometry of plume grains and corresponding peer-reviewed laboratory silica coagulation experiments.

d. Pressure (MPa): 4 to 10
Source: Current work has provided critical constraints on the seafloor pressure at ~40–100 bar (Postberg et al. 2009, 2011; Hsu et al. 2015; Glein et al. 2018). This converts to 4–10 MPa. The best interior structure models, based on Cassini gravity and topography data, dictate that the hydrostatic pressure at the ocean floor is approximately 7.4 MPa (Glein et al., 2018). Other sources estimate that fluids percolating deeper into the highly porous core could experience pressures up to 37.7 MPa at the moon's center before ascending (Glein et al., 2018, Neveu et al., 2014). There is no major disagreement, as these values are mathematically derived from Enceladus's known mass, radius, and internal density profile.

Reliability: 3 (High). Pressure profiles are mathematically calculated directly from highly precise, peer-reviewed Cassini gravity and libration measurements of the moon's interior structure.

e. Isolation score (1-5): 5
(i) Physical Confinement (1-5): 5
Source: As an extraterrestrial environment, Enceladus is separated from Earth by vast interplanetary distances, completely preventing any direct physical transfer of terrestrial biological contaminants or matter (Worth et al., 2013).

(ii) Hydrologic Connectivity (1-5): 5
Source: The subsurface ocean and hydrothermal systems of Enceladus are entirely self-contained within the moon, meaning there is an absolute lack of hydrological connectivity with Earth's ecosystems (Worth et al., 2013).

(iii) Isolation Time (1-5): 5
Source: Enceladus has been completely isolated from Earth since the formation of the solar system, ensuring any potential biosphere has evolved independently for billions of years without terrestrial contact (Worth et al., 2013).

Reliability: 3 (High). The absolute physical separation and lack of biological connectivity between Earth and the Saturnian system is a fundamental and undisputed astronomical reality.

f. Redox Score (1 -5): 5
(i) Redox Species (1-5): 5
Source: Cassini's Ion and Neutral Mass Spectrometer provided confirmed, direct measurements of both molecular hydrogen and carbon dioxide in the plume, which are directly sourced from the hydrothermal vents (Waite et al., 2017).

(ii) Equilibrium (1-5): 5
Source: The coexistence of abundant molecular hydrogen and carbon dioxide demonstrates a well-documented, strong chemical disequilibrium that represents a constant flux of available free energy (Waite et al., 2017, Ray et al., 2021).

(iii) Chemolithotrophic Metabolism (1-5): 5
Source: Strong evidence confirms that the thermodynamic drive for hydrogenotrophic methanogenesis is highly favorable in these fluids, closely matching the energy availability in alkaline hydrothermal ecosystems on Earth (Taubner et al., 2018, Waite et al., 2017).

Reliability: 3 (High). Redox conditions are robustly validated by primary Cassini mass spectrometry data and peer-reviewed bioenergetic modeling of methanogenic pathways.

2. **Diffuse Venting Zone**
   Scope Definition: This site includes the broad regions of Enceladus's highly porous, unconsolidated rocky core where ocean water percolates and undergoes lower-temperature fluid exchanges, explicitly excluding the focused, high-temperature hydrothermal vents and the cold, unmixed abyssal ocean.

   a. Temperature (°C): 10 to 90
   Source: The best peer-reviewed models, such as those by Choblet et al. (2017) and Barge and White (2017), indicate that diffuse fluid circulation within the highly porous core encompasses lower-temperature gradients leading up to hot upwellings. Other sources establish that silica nanoparticle formation requires temperatures above 90 °C, setting a strict upper limit for diffuse venting before it transitions into focused white smokers (Hsu et al., 2015, Sekine et al., 2015). Disagreements largely center on whether these lower-temperature regimes can independently sustain large-scale geochemical alterations without localized high-temperature spikes.

   Reliability: 2 (Medium). This temperature range is derived from thermodynamic models of silica precipitation and large-scale tidal dissipation rather than direct thermal measurements.

   b. pH: 8.0 to 10.5
   Source: Group A, supported by Glein and Waite (2020) and Fifer et al. (2022), says the pore fluids within the diffuse venting zone have a pH of 8.0 to 9.0 because fluids exchanging efficiently with the overlying ocean would be buffered by the bulk ocean's moderately alkaline composition. Group B says the pH of fluids interacting with the rocky core could range from 8.5 to 10.5 because serpentinization and water-rock interactions in chemically closed pore spaces naturally drive the pH higher as the fluids cool (Sekine et al., 2015, Glein et al., 2015). A union range of 8.0 to 10.5 is adopted because diffuse venting encompasses both chemically open regions mixing with the bulk ocean and more restricted pore spaces within the core.

   Reliability: 2 (Medium). The pH values rely on complex geochemical modelling of carbonate speciation and laboratory serpentinization experiments instead of direct sampling.

   c. Salinity (g/kg): 5 to 40
   Source: Cassini measurements of plume ice grains indicate a baseline ocean salinity of 5 to 20 g/kg, establishing the minimum salinity for percolating fluids (Postberg et al., 2009, Glein et al., 2018). Other experimental sources state that ongoing water-rock interactions and the prevention of silica coagulation set a strict upper salinity limit of 40 g/kg within the core (Hsu et al., 2015). There is general agreement on this moderate salinity, though exact concentrations vary based on localized freezing or fluid pathways.

   Reliability: 3 (High). The salinity limits are tightly constrained by primary mass spectrometry of plume grains and corresponding peer-reviewed laboratory silica coagulation experiments.

e. Pressure (MPa): 4 to 10
Source: Current work has provided critical constraints on the seafloor pressure at ~40–100 bar (Postberg et al. 2009, 2011; Hsu et al. 2015; Glein et al. 2018). This converts to 4–10 MPa. Interior structure models based on Cassini gravity data dictate that the hydrostatic pressure at the ocean floor is approximately 7.4 MPa (Glein et al., 2018). Other sources estimate that fluids percolating deeper into the highly porous core could experience pressures up to 37.7 MPa near the moon's center (Glein et al., 2018, Choblet et al., 2017). There is no major disagreement, as these pressure gradients are mathematically derived from Enceladus's known internal density profile and porous core structure.

Reliability: 3 (High). Pressure profiles are mathematically calculated directly from highly precise, peer-reviewed Cassini gravity and libration measurements of the moon's interior structure.

e. Isolation Score (1-5): 5
(i) Physical Confinement (1-5): 5
Source: As an extraterrestrial environment, Enceladus is separated from Earth by vast interplanetary distances, completely preventing any direct physical transfer of terrestrial biological contaminants or matter (McKay et al., 2014).

(ii) Hydrologic Connectivity (1-5): 5
Source: The subsurface ocean and hydrothermal systems of Enceladus are entirely self-contained within the moon, meaning there is an absolute lack of hydrological connectivity with Earth's ecosystems (McKay et al., 2014).
(iii) Isolation Time (years) (1-5): 5

Source: Enceladus has been completely isolated from Earth since the formation of the solar system, ensuring any potential biosphere has evolved independently for billions of years without terrestrial contact (McKay et al., 2014).

Reliability: 3 (High). The absolute physical separation and lack of biological connectivity between Earth and the Saturnian system is a fundamental and undisputed astronomical reality.

f. Redox Score (1-5): 5
(i) Redox Species (1-5): 5
Source: Cassini's Ion and Neutral Mass Spectrometer provided confirmed measurements of molecular hydrogen and carbon dioxide in the plume, which are generated by widespread water-rock interactions in the porous core (Waite et al., 2017).

(ii) Equilibrium (1-5): 5
Source: The coexistence of molecular hydrogen and oxidized carbon species in the percolating core fluids demonstrates a strong chemical disequilibrium that provides a constant flux of free energy (Waite et al., 2017, Ray et al., 2021).

(iii) Chemolithotrophic Metabolism (1-5): 5
Source: Peer-reviewed bioenergetic models confirm that the thermodynamic drive for hydrogenotrophic methanogenesis is highly favorable in these diffuse, lower-temperature core fluids, providing ample energy for potential microbial life (Taubner et al., 2018, Affholder et al., 2021).

Reliability: Score 3 (High). Redox conditions are robustly validated by primary Cassini mass spectrometry data and peer-reviewed bioenergetic modeling of methanogenic pathways.

3. **Abyssal Ocean**
Scope Definition: This site encompasses the bulk global subsurface liquid water reservoir of Enceladus, situated between the overlying icy crust and the rocky core, explicitly excluding the highly localized high-temperature hydrothermal vents and the dynamic, rapidly fractionating conduit environments near the surface.

   a. Temperature (°C): -1.0 to 1
   Source: Models based on Cassini data state that the ocean's temperature must be near the freezing point of water, depressed by the presence of dissolved salts to approximately -1.0 °C. Other sources land on similar values around 0 °C to 0.4 °C, as the bulk ocean acts as a massive thermal reservoir insulated by the ice shell. There is virtually no disagreement, as thermodynamics strictly dictate that a liquid ocean in contact with an ice shell must sit near the pure water freezing point (Glein et al., 2018, Fifer et al., 2022).

   Reliability: 3 (High). This temperature is a robust thermodynamic certainty derived from peer-reviewed interpretations of Cassini's confirmation of a global liquid water ocean interfacing with an ice shell.

   b. pH: 8.5 to 11.6
   Source: Group A says the abyssal ocean has a moderately alkaline pH of 8.5 to 9.0 because these values best fit thermodynamic models of carbonate speciation when correcting for CO2 exsolution and condensation during plume eruption. Group B says the ocean pH is significantly higher, between 10.1 and 11.6, because the newly analysed distribution of phosphate species found in Cassini's ice grain data strongly acts as an alkaline pH indicator. A union range of 8.5 to 11.6 is adopted because both estimates rely on rigorous, peer-reviewed chemical fractionation models from primary Cassini mass spectrometry data (Glein & Waite, 2020, Fifer et al., 2022, Glein & Truong, 2025).

   Reliability: 3 (High). These pH values are robustly derived from primary Cassini mass spectrometry data of plume ice grains, modelled through peer-reviewed thermodynamic speciation frameworks.

c. Salinity (g/kg): 5 to 40
Source: Cassini measurements of salt-rich E-ring ice grains indicate a baseline bulk ocean salinity of 5 to 20 g/kg, dominated by sodium chloride and sodium carbonates. Other recent peer-reviewed models extending thermodynamic speciation to include newly detected phosphates land on a broader possible range of 5.1 to 40 g/kg. The nature of this minor disagreement stems from varying assumptions about the exact extent of salt fractionation and cryoconcentration that occurs when ocean water freezes to form the sampled plume ice grains (Postberg et al., 2009, Glein & Truong, 2025).

Reliability: 3 (High). Salinity ranges are tightly constrained by direct primary mass spectrometry from Cassini's Cosmic Dust Analyzer and subsequent peer-reviewed geochemical modelling.

d. Pressure (MPa): 0.2 to 7.4
Source: The best peer-reviewed interior structure models dictate that hydrostatic pressures at the ocean floor reach approximately 7.4 MPa due to Enceladus's weak gravity. Other sources establish that at the very top of the abyssal ocean, where the ice shell is thinnest at the south pole, the pressure lands around 0.2 MPa. There is no major disagreement, as these pressure gradients are strictly mathematically derived from Enceladus’s known internal density profile and variations in the ice shell thickness (Glein et al., 2018).

Reliability: 3 (High). Pressure profiles are mathematically calculated directly from highly precise, peer-reviewed Cassini gravity and libration measurements of the moon's interior structure.

e. Isolation score (1-5): 5
(i) Physical Confinement (1-5): 5
Source: As an extraterrestrial environment, Enceladus is separated from Earth by vast interplanetary distances, completely preventing any direct physical transfer of terrestrial biological contaminants or matter (Worth et al., 2013).

(ii) Hydrologic Connectivity (1-5): 5
Source: The subsurface ocean and hydrothermal systems of Enceladus are entirely self-contained within the moon, meaning there is an absolute lack of hydrological connectivity with Earth's ecosystems (Worth et al., 2013).

(iii) Isolation Time (years) (1-5): 5
Source: Enceladus has been completely isolated from Earth since the formation of the solar system, ensuring any potential biosphere has evolved independently for billions of years without terrestrial contact (Worth et al., 2013).

Reliability: 3 (High). The absolute physical separation and lack of biological connectivity between Earth and the Saturnian system is a fundamental and undisputed astronomical reality.

f. Redox Score (1 -5): 5
(i) Redox Species (1-5): 5
Source: Cassini's Ion and Neutral Mass Spectrometer detected substantial levels of molecular hydrogen, carbon dioxide, and methane that have accumulated and mixed throughout the bulk ocean. This creates a bulk fluid exceptionally rich in biologically viable redox pairs (Waite et al., 2017).

(ii) Equilibrium (1-5): 5
Source: The coexistence of abundant reduced hydrogen and oxidized carbon dioxide in the bulk ocean demonstrates a chemical disequilibrium. This persistent thermodynamic imbalance confirms that these species have not fully reacted abiotically, leaving a constant flux of available free energy (Waite et al., 2017, Ray et al., 2021).

(iii) Chemolithotrophic Metabolism (1-5): 5
Source: Peer-reviewed bioenergetic models confirm that the thermodynamic drive for hydrogenotrophic methanogenesis is highly favourable within the bulk ocean's conditions. The available energy from the H2-CO2 redox pair easily meets the maintenance and growth requirements for terrestrial-type methanogenic archaea (Affholder et al., 2021, Fifer et al., 2022).

Reliability: 3 (High). Redox conditions and bioenergetic potentials are robustly validated by primary Cassini mass spectrometry data and subsequent peer-reviewed ecological thermodynamic modelling.

4. **Ocean Connected Tiger Stripe Brines**
Scope Definition: This site encompasses the deep-fissure plumbing system where bulk ocean water ascends directly through the ice shell to the surface, including the liquid water table where boiling, gas exsolution, and evaporative cooling occur, but explicitly excluding the abyssal ocean below and the ejected plume ice grains in space.

a. Temperature (°C): -1.0 to 1
Source: Peer-reviewed models establish that liquid water ascending through the tiger stripe fractures must remain near the freezing point of water, approximately 1 °C to -1.0 °C, at the boiling liquid-vapor interface. Other sources investigating evaporative cooling confirm that as the water boils into the vacuum, it loses latent heat and stays at or just below 0 °C before flash-freezing. There is general agreement that the liquid phase within these conduits is thermodynamically bound to temperatures just below the freezing point of pure water due to its salt content (Ingersoll & Pankine, 2010, Kite & Rubin, 2016, Fifer et al., 2022).

Reliability: 3 (High). The temperatures are tightly constrained by robust thermodynamic principles dictating phase equilibria between liquid brine, water vapor, and the surrounding ice shell.

b. pH: 8.5 to 11.6
Source: Group A says the ascending brines maintain a pH of 8.5 to 9.0 because models of plume eruption indicate significant water vapor condensation on fracture walls alongside rapid ejection, preventing massive pH shifts before eruption (Glein & Waite, 2020, Fifer et al., 2022). Group B says the pH could increase significantly up to 11.6 because as the brines ascend and experience pressure drops, carbon dioxide vigorously exsolves and degasses from the liquid, naturally driving the remaining fluid to become more alkaline (Glein & Truong, 2025, Glein et al., 2015). A union range of 8.5 to 11.6 is adopted because it reflects both the baseline oceanic pH entering the cracks and the potential alkalinisation resulting from extensive volatile exsolution during ascent.

Reliability: 3 (High). These pH values are rigorously derived from primary mass spectrometry data combined with peer-reviewed thermodynamic modelling of volatile exsolution and carbonate speciation.

c. Salinity (g/kg): 5 to 40
Source: Sources utilizing Cassini data report that the baseline salinity of ascending fluid ranges from 5 to 40 g/kg, consisting primarily of sodium salts. Other geochemical models suggest that preferential evaporation at the water table within the fractures causes salts to become progressively more concentrated in the residual liquid. There is no major disagreement, though the exact concentration depends on the dynamic balance between active boiling, vapour escape, and the continuous flushing of fresh ocean water into the conduits (Postberg et al., 2009, Glein & Truong, 2025, Kite & Rubin, 2016).

Reliability: 3 (High). Salinity estimates are robustly supported by direct in-situ sampling of ejected salt-rich ice grains analyzed by Cassini's Cosmic Dust Analyzer.

d. Pressure (MPa): 0
Source: Thermodynamic models state that the pressure at the liquid-vapor interface within the shallow fractures is determined by the saturation vapor pressure of water at 0 °C, approximately 0.0006 MPa. Other sources confirm that deeper within the conduits, the hydrostatic pressure increases linearly with depth, reaching a maximum of roughly 7.4 MPa where the fractures connect to the bulk abyssal ocean. There is universal agreement on these limits, as they are mathematically dictated by the phase diagram of water and Enceladus's internal gravity profile (Ingersoll & Pankine, 2010, Glein et al., 2018).

Reliability: 3 (High). Pressure profiles are strictly derived from fundamental thermodynamic laws governing the triple point of water and Cassini's peer-reviewed gravity measurements.

e. Isolation Score (1-5): 5
(i) Physical Confinement (1-5): 5
Source: As an extraterrestrial environment, Enceladus is separated from Earth by vast interplanetary distances, completely preventing any direct physical transfer of terrestrial biological contaminants or matter (Worth et al., 2013).

(ii) Hydrologic Connectivity (1 -5): 5
Source: The subsurface ocean and hydrothermal systems of Enceladus are entirely self-contained within the moon, meaning there is an absolute lack of hydrological connectivity with Earth's ecosystems (Worth et al., 2013).

(iii) Isolation Time (1 -5): 5
Source: Enceladus has been completely isolated from Earth since the formation of the solar system, ensuring any potential biosphere has evolved independently for billions of years without terrestrial contact (Worth et al., 2013).

Reliability: 3 (High). The absolute physical separation and lack of biological connectivity between Earth and the Saturnian system is a fundamental and undisputed astronomical reality.

f. Redox Score (1 -5): 5
(i) Redox Species (1-5): 5
Source: Cassini's Ion and Neutral Mass Spectrometer detected substantial concentrations of molecular hydrogen, carbon dioxide, and methane exsolving from the ascending brines within the tiger stripe conduits (Waite et al., 2017).

(ii) Equilibrium (1-5): 5
Source: The rapid boiling and volatile exsolution occurring within these conduits actively preserves the strong chemical disequilibrium between reduced hydrogen and oxidized carbon species ascending from the ocean (Waite et al., 2017, Fifer et al., 2022).

(iii) Chemolithotrophic Metabolism (1-5): 5
Source: Peer-reviewed bioenergetic models demonstrate that the abundant molecular hydrogen and carbon dioxide streaming through these fractures provide highly favourable thermodynamic drives and abundant free energy for terrestrial-type methanogenic archaea (Affholder et al., 2022, Waite et al., 2017).

Reliability: 3 (High). Redox potentials are strongly supported by primary Cassini mass spectrometry of the outgassed species and validated by rigorous bioenergetic thermodynamic models.

5. **Intrashell Mushy Fracture Brines**
   Scope Definition: This site encompasses the two-phase coexistence region (mushy zone) of solid ice and interstitial saline liquid generated by localized shear heating along tiger stripe fractures in Enceladus's ice shell, explicitly excluding the underlying bulk ocean and the high-velocity erupting jet conduits.

   a. Temperature (°C): -21.15 to 0
      Source: Peer-reviewed models, such as Meyer et al. (2025), define the temperature of the mushy zone as existing between the eutectic temperature of the local salt mixture and the pure ice melting point. For a sodium chloride-dominated system, this corresponds to temperatures ranging from approximately -21.15 °C up to 0 °C depending on the distance from the fracture's active shear zone. Other sources assessing pure ice shells do not account for this freezing point depression. Disagreement is minimal as the phase diagram of salty water strictly dictates these thermal boundaries (Meyer et al., 2025).

      Reliability: 3 (High). These temperature boundaries are robustly determined by fundamental thermodynamic phase diagrams for binary eutectic systems combined with newly published geophysical shear heating models.

   b. pH: 7.95 to 11.6
      Source: Group A says the pH of the mushy brine is around 7.95 to 9.05, closely reflecting the moderately alkaline bulk ocean that was originally frozen into the ice shell before being remelted by shear heating (Fifer et al., 2022). Group B says the pH could range significantly higher, from 10.1 to 11.6, because recent phosphate speciation models demand a highly alkaline source and fractional melting within the mushy zone inherently alters the residual liquid's alkalinity (Glein & Truong, 2025, Klenner et al., 2025). A union range of 7.95 to 11.6 is adopted because the interstitial brine's pH is primarily inherited from the highly contested bulk ocean composition, further complicated by localized fractionation during the partial melting process (Fifer et al., 2022, Glein & Truong, 2025, Klenner et al., 2025).

      Reliability: 2 (Medium). The pH is uncertain because it relies on contested ocean pH models compounded by the complex, unmeasured chemical fractionation occurring during partial melting within the ice shell.

   c. Salinity (g/kg): 35 to 230
      Source: The thermomechanical models by Meyer et al. (2025) indicate that partial melting concentrates salts into the interstitial fluid, leading to salinities ranging from the baseline ice shell content of roughly 2 g/kg up to the eutectic concentration of sodium chloride near 230 g/kg. Other sources studying ice-ocean exchange confirm that freezing and subsequent partial melting strongly fractionates the fluid, creating localized pockets of highly concentrated brine. Disagreements are negligible, as this wide range reflects the dynamic, depth-dependent transition from solid ice to fully mobilized eutectic liquid (Meyer et al., 2025, Klenner et al., 2025).

Reliability: 2 (Medium). Salinity limits are theoretically robust based on phase diagrams but rely on idealized geophysical models rather than direct sampling of the interstitial fluid itself.

d. Pressure (MPa): 0 to 0.6
Source: The best geophysical models estimate an active south polar ice shell thickness of approximately 6 km, dictating interstitial hydrostatic pressures from near-vacuum triple-point pressures (0.0006 MPa) at shallow melt zones up to 0.6 MPa at the shell's base. Other models considering thicker regional shells suggest pressures could reach 7.4 MPa if the mushy zone extends deeper. There is general agreement on the pressure gradient, with variance reflecting uncertainty in the exact thickness of the active ice shell (Meyer et al., 2025, Glein et al., 2018).

Reliability: 3 (High). Pressures are strictly constrained by the well-established hydrostatic gradient derived from Cassini gravity data and the moon's known mass and density profile.

e. Isolation Score (1-5): 5
(i) Physical Confinement (1-5): 5
Source: As an extraterrestrial environment, Enceladus is separated from Earth by vast interplanetary distances, completely preventing any direct physical transfer of terrestrial biological contaminants or matter to its subsurface environments (Worth et al., 2013).

(ii) Hydrologic Connectivity (1-5): 5
Source: The subsurface ocean, ice shell, and hydrothermal systems of Enceladus are entirely self-contained within the moon, meaning there is an absolute lack of hydrological connectivity with Earth's diverse ecosystems (Worth et al., 2013).

(iii) Isolation Time (1-5): 5
Source: Enceladus has been completely isolated from Earth since the formation of the solar system, ensuring any potential biosphere has evolved independently for billions of years without terrestrial contact (Worth et al., 2013).

Reliability: 3 (High). The absolute physical separation and lack of biological connectivity between Earth and the Saturnian system is a fundamental and undisputed astronomical reality.

f. Redox Score (1-5): 5
(i) Redox Species (1-5): 5
Source: Cassini mass spectrometry detected species like molecular hydrogen, carbon dioxide, and methane that would be incorporated into the ice shell from the ocean and released into the interstitial brine during melting (Meyer et al., 2025, Waite et al., 2017).

(ii) Equilibrium (1-5): 5
Source: The periodic phase changes, fluid mobilization, and volatile exsolution occurring within the mushy zone actively maintain a state of strong chemical disequilibrium between reduced and oxidized species trapped within the ice (Meyer et al., 2025, Fifer et al., 2022).

(iii) Chemolithotrophic Metabolism (1-5): 5
Source: Peer-reviewed bioenergetic models demonstrate that if trapped oceanic volatiles are remobilized within the mushy zone brine, the available molecular hydrogen and carbon dioxide would provide highly favorable free energy for hydrogenotrophic methanogenesis (Affholder et al., 2022, Meyer et al., 2025).

Reliability: 2 (Medium). Redox potential is inferred by extrapolating ocean composition into the ice shell matrix, lacking direct empirical validation of the exact chemical state within the mushy zone.

6. **Plume Ice Grains - Type I**
   Scope Definition: This site includes the small, almost pure water ice grains that form primarily via the homogenous condensation of supersaturated water vapor within the cold, upper regions of Enceladus's tiger stripe fissures, excluding salt-rich or complex-organic-rich grains.

   a. Temperature (°C): -76 to -43
      Source: Thermal mapping of the active tiger stripe fissures establishes a surface outlet temperature of approximately -76 °C (197 K), meaning condensing grains must form around or below this temperature (Goguen et al., 2013). Other thermomechanical models assume gas cools from 0 °C (273 K) at the ocean interface down to these cryogenic temperatures during ascent, predicting condensation primarily occurs in the narrowest, coldest upper vent regions (Schmidt et al., 2008). Disagreements are minor and generally relate to the exact depth and width of the condensation zone rather than the exit temperature (Ingersoll & Pankine, 2010).

      Reliability: 3 (High). This temperature range is highly reliable because it is derived directly from Cassini's infrared thermal mapping of the active fissures (Goguen et al., 2013).

   b. pH: ~7.0
      Source: Mass spectrometry data establishes that Type I grains are almost pure water ice, entirely lacking the alkaline sodium carbonates that buffer the subsurface ocean (Postberg et al., 2009). Other analyses agree that because these grains form from condensed water vapor rather than frozen liquid spray, they do not inherit the ocean's high pH of 9 to 11 (Glein et al., 2015). There is no significant disagreement regarding this, as the absence of alkaline buffering salts in vapor-condensed ice naturally implies a neutral pH for the pure water matrix (Postberg et al., 2011).

Reliability: 3 (High). This is highly reliable as Cassini Cosmic Dust Analyzer spectra definitively show a lack of pH-altering salts in this specific grain population (Postberg et al., 2009).

c. Salinity (g/kg): ~0
Source: In-situ measurements from Cassini's Cosmic Dust Analyzer demonstrate that Type I grains are almost pure water ice, with a sodium-to-water ratio on the order of 10^{-7} (Postberg et al., 2009). Other plume models consistently land on this same conclusion, noting that salts cannot easily condense from the gaseous phase, thereby leaving vapor-condensed grains practically salt-free (Ershova et al., 2024). There is no disagreement in the literature regarding this parameter, as the compositional dichotomy between salt-poor condensed grains and salt-rich frozen ocean spray is universally accepted (Postberg et al., 2011).

Reliability: 3 (High). This parameter is exceptionally reliable because it is based on the direct, in-situ compositional analysis of thousands of individual ice grains by Cassini (Postberg et al., 2009).

d. Pressure (MPa): 0
Source: Thermodynamic models of the plume dictate that pressures cannot exceed the triple point of water (0.0006 MPa) at the liquid-vapor interface and rapidly drop toward zero as the gas expands upward (Nakajima & Ingersoll, 2016). Other hydrodynamic simulations agree with this bounding, showing that the vapor must expand into the vacuum of space to achieve the observed supersonic velocities (Schmidt et al., 2008). There is no substantial disagreement regarding this pressure range, as it is a fundamental thermodynamic requirement for a system venting directly into space (Kite & Rubin, 2016).

Reliability: 3 (High). This pressure range is highly reliable as it is dictated by fundamental thermodynamic laws governing liquid water exposed to the vacuum of space (Nakajima & Ingersoll, 2016).

e. Isolation Score (1-5): 5
(i) Physical Confinement (1-5): 5
Source: As an extraterrestrial environment, Enceladus is separated from Earth by vast interplanetary distances, completely preventing any direct physical transfer of terrestrial biological contaminants or matter (Worth et al., 2013).

(ii) Hydrologic Connectivity (1-5): 5
Source: The subsurface ocean and hydrothermal systems of Enceladus are entirely self-contained within the moon, meaning there is an absolute lack of hydrological connectivity with Earth's ecosystems (Worth et al., 2013).

(iii) Isolation Time (1-5): 5
Source: Enceladus has been completely isolated from Earth since the formation of the solar system, ensuring any potential biosphere has evolved independently for billions of years without terrestrial contact (Worth et al., 2013).

Reliability: 3 (High). The absolute physical separation and lack of biological connectivity between Earth and the Saturnian system is a fundamental and undisputed astronomical reality.

f. Redox Score (1-5): 1
(i) Redox Species (1-5): 1
Source: Type I grains are formed from pure water vapor, inherently lacking the dissolved reductants and oxidants present in the bulk ocean (Postberg et al., 2009). Condensation physically separates the solid ice from these dissolved chemical energy sources entirely (Schmidt et al., 2008).

(ii) Equilibrium (1-5): 1
Source: The rapid flash-freezing and extreme cryogenic temperatures of the vapor condensation process prevent any meaningful chemical equilibrium from being established within the solid ice grain itself (Schmidt et al., 2008). The system is a kinetically frozen solid.

(iii) Chemolithotrophic Metabolism (1-5): 1
Source: Pure, salt-free water ice grains in a vacuum provide absolutely no solvent, nutrients, or chemical energy gradients to support chemolithotrophic metabolism (Postberg et al., 2009). They are completely biologically inert physical structures.

Reliability: 3 (High). This is highly reliable because the completely frozen, pure-water composition of these grains inherently precludes any complex redox chemistry or biological metabolism (Postberg et al., 2009).

7. **Plume Ice Grains - Type II (VOC)**
Scope Definition: This site includes salt-poor water ice grains enriched in low-mass (under 100 atomic mass units) volatile organic compounds (VOCs), such as nitrogen-bearing, oxygen-bearing, and simple aromatic species. It specifically excludes salt-rich frozen ocean droplets and grains bearing high-mass, refractory macromolecular organics (HMOC).

a. Temperature (°C): -76 to -37
Source: The thermal and adsorption models dictate that temperatures range from approximately -37 °C deep in the upper fractures to -76 °C at the surface vents, which provides the necessary conditions for volatile organics to adsorb onto condensing ice grains (Bouquet et al., 2019, Goguen et al., 2013). Other physical models agree that the grains undergo severe and continuous cooling during their supersonic ascent, eventually reaching ambient space temperatures (Yeoh et al., 2015). There is no significant disagreement, as this temperature gradient directly drives the observed sequential condensation of the VOCs (Khawaja et al., 2019).

Reliability: 3 (High). This parameter is highly reliable because it combines Cassini's direct infrared thermal mapping with robust thermodynamic adsorption models for organic species (Goguen et al., 2013).

b. pH: ~7.0
Source The in-situ mass spectrometry by Cassini demonstrates that Type II (VOC) grains are intrinsically salt-poor, containing negligible amounts of the alkaline sodium carbonates that buffer the subsurface ocean (Postberg et al., 2009). Other studies independently confirm that because these grains form via the condensation of water vapor followed by the adsorption of volatile organic gases, they cannot inherit the highly alkaline pH of the bulk ocean (Khawaja et al., 2019). There is no disagreement regarding this neutral pH, as the absence of buffering salts in vapor-condensed grains is firmly established (Ershova et al., 2024).

Reliability: 3 (High). This is highly reliable because Cassini Cosmic Dust Analyzer spectra explicitly confirm the absence of pH-altering salts within this specific sub-population of ice grains (Postberg et al., 2009).

c. Salinity (g/kg): ~0
Source: Cassini measurements definitively classify Type II (VOC) grains as salt-poor, confirming they contain only trace amounts of sodium and potassium salts (Postberg et al., 2009). Other plume formation models align perfectly with this finding, explaining that these grains grow from water vapor and organic gases which cannot easily transport non-volatile salts from the liquid ocean phase (Khawaja et al., 2019). There is no disagreement in the literature, as the compositional dichotomy between salt-poor vapor-condensed grains and salt-rich frozen ocean spray is universally accepted (Postberg et al., 2011).

Reliability: 3 (High). This is exceptionally reliable as it relies on direct, in-situ compositional analyses of thousands of individual ice grains by Cassini (Postberg et al., 2009).

d. Pressure (MPa): 0
Source: Hydrodynamic models establish that the vapor pressure inside the fractures cannot exceed the triple point of water (~0.0006 MPa) and rapidly approaches zero as the plume expands (Nakajima & Ingersoll, 2016). Other physical simulations agree, showing that this continuous pressure drop into a vacuum drives the supersonic acceleration of the gas and its entrained ice grains (Schmidt et al., 2008). There is no disagreement regarding this pressure bounding, as it is a strict thermodynamic requirement for vapor venting into space from a subsurface liquid water table (Kite & Rubin, 2016).

Reliability: 3 (High). This pressure limit is highly reliable because it is derived from the fundamental thermodynamic laws governing the liquid-vapor interface of water venting to a vacuum (Nakajima & Ingersoll, 2016).

e. Isolation Score (1-5): 5
(i) Physical Confinement (1-5): 5
Source: As an extraterrestrial environment, Enceladus is separated from Earth by vast interplanetary distances, completely preventing any direct physical transfer of terrestrial biological contaminants or matter (Worth et al., 2013).

(ii) Hydrologic Connectivity (1-5): 5
Source: The subsurface ocean and hydrothermal systems of Enceladus are entirely self-contained within the moon, meaning there is an absolute lack of hydrological connectivity with Earth's ecosystems (Worth et al., 2013).

(iii) Isolation Time (1-5): 5
Source: Enceladus has been completely isolated from Earth since the formation of the solar system, ensuring any potential biosphere has evolved independently for billions of years without terrestrial contact (Worth et al., 2013).

Reliability: 3 (High). The absolute physical separation and lack of biological connectivity between Earth and the Saturnian system is a fundamental and undisputed astronomical reality.

f. Redox Score (1-5): 1.33
(i) Redox Species (1-5): 2
Source: Although these grains are enriched in reactive volatile organic compounds like amines and carbonyls, the species are physically immobilized within a solid, frozen water ice matrix in a vacuum (Khawaja et al., 2019, Bouquet et al., 2019).

(ii) Equilibrium (1-5): 1
Source: The extreme cryogenic temperatures and rapid cooling processes entirely halt chemical kinetics, preventing any active reactions or chemical equilibrium from being reached among the adsorbed organic species (Bouquet et al., 2019).

(iii) Chemolithotrophic Metabolism (1-5): 1
Source: Despite carrying potentially biogenic organic precursors, the completely frozen, solid-state nature of these grains in the vacuum of space offers zero liquid solvent to support any chemolithotrophic metabolism (Khawaja et al., 2019).

Reliability: 3 (High). This is highly reliable because the cryogenic, solid-state environment of these grains inherently precludes active biological metabolism or ongoing aqueous redox chemistry (Khawaja et al., 2019).

8. **Plume Ice Grains - Type II (HMOC)**
   Scope Definition: This site encompasses salt-poor water ice grains that are highly enriched in complex, refractory high-mass organic cations (HMOC) exceeding 200 atomic mass units, formed via the aerosolization of an insoluble organic-rich film at the liquid water table by bursting gas bubbles, specifically excluding volatile-only organics and salt-rich frozen ocean spray grains.

   a. Temperature (°C): ~0 to -76
      Source: Models for HMOC grain formation indicate they originate from aerosolized droplets of an organic-rich film at the liquid water table, where temperatures are near the freezing point of water at approximately 0 °C (Postberg et al., 2018). Other thermal models of the tiger stripe fissures agree that these aerosolized cores rapidly freeze and cool to approximately -76 °C as they are carried upward to the surface vents (Goguen et al., 2013, Schmidt et al., 2008). There is no significant disagreement regarding this temperature gradient driving the initial freezing of these organic cores.

      Reliability Score: 3 (High). This parameter is highly reliable as it combines in-situ mass spectrometry with robust infrared thermal mapping of the active surface fissures (Postberg et al., 2018).

   b. pH: ~7.0
      Source: In-situ measurements from Cassini's Cosmic Dust Analyzer demonstrate that Type II HMOC grains are intrinsically salt-poor, lacking the alkaline sodium carbonates that buffer the subsurface ocean (Postberg et al., 2018). Other studies independently confirm that these grains form from organic nucleation cores condensing pure water vapor, preventing them from inheriting the ocean's high pH of 9 to 11 (Khawaja et al., 2019, Glein & Waite, 2020). There is no disagreement regarding this neutral pH, as the compositional separation between salt-poor organic grains and alkaline ocean spray is universally accepted.

      Reliability: 3 (High). This is highly reliable because Cassini spectra explicitly confirm the absence of pH-altering salts within this specific sub-population of complex organic grains (Postberg et al., 2018).

   c. Salinity (g/kg): ~0
      Source: Cassini measurements definitively classify HMOC-bearing grains as salt-poor, containing only trace amounts of sodium and potassium compared to Type III grains (Postberg et al., 2009, Postberg et al., 2018). Other plume formation models align with this, explaining that these grains grow around insoluble organic cores from water vapor, which cannot easily transport non-volatile salts from the liquid phase (Khawaja et al., 2019). There is no disagreement in the literature, as the compositional dichotomy between salt-poor HMOC grains and salt-rich frozen ocean spray is universally accepted.

      Reliability: 3 (High). This parameter is exceptionally reliable as it relies on direct, in-situ compositional analyses of thousands of individual ice grains by the Cassini spacecraft (Postberg et al., 2009).

d. Pressure (MPa): ~0
Source: Hydrodynamic models establish that bubble-bursting aerosolization at the subsurface water table occurs at the triple point of water, corresponding to approximately 0.0006 MPa (Nakajima & Ingersoll, 2016). Other physical simulations agree, showing that the pressure drops continuously toward the vacuum of space, driving the supersonic acceleration of the gas and entrained organic grains (Schmidt et al., 2008). There is no disagreement regarding this pressure bounding, as it is a strict thermodynamic requirement for liquid venting into space (Kite & Rubin, 2016).

Reliability: 3 (High). This pressure limit is highly reliable because it is derived from fundamental thermodynamic laws governing the liquid-vapor interface of water venting to vacuum (Nakajima & Ingersoll, 2016).

e. Isolation Score (1-5): 5
(i) Physical Confinement (1-5): 5
Source: As an extraterrestrial environment, Enceladus is separated from Earth by vast interplanetary distances, completely preventing any direct physical transfer of terrestrial biological contaminants or matter (Worth et al., 2013).

(ii) Hydrologic Connectivity (1-5): 5
Source: The subsurface ocean and hydrothermal systems of Enceladus are entirely self-contained within the moon, meaning there is an absolute lack of hydrological connectivity with Earth's ecosystems (Worth et al., 2013).

(iii) Isolation Time (1-5): 5
Source: Enceladus has been completely isolated from Earth since the formation of the solar system, ensuring any potential biosphere has evolved independently for billions of years without terrestrial contact (Worth et al., 2013).

Reliability: 3 (High). The absolute physical separation and lack of biological connectivity between Earth and the Saturnian system is a fundamental and undisputed astronomical reality.

f. Redox Score (1-5): 1
(i) Redox Species (1-5): 2
Source: Although highly enriched in complex, macromolecular organic compounds which could act as potential redox reactants, these species are physically immobilized within a solid water ice matrix in a vacuum (Postberg et al., 2018).

(ii) Equilibrium (1-5): 1
Source: The rapid flash-freezing of the aerosolized organic cores and extreme cryogenic temperatures entirely halt chemical kinetics, preventing any active reactions or chemical equilibrium from being reached (Schmidt et al., 2008).

(iii) Chemolithotrophic Metabolism (1-5): 1
Source: Despite carrying complex organic material, the completely frozen, solid-state nature of these grains in the vacuum of space offers zero liquid solvent to support any chemolithotrophic metabolism (Postberg et al., 2018).

Reliability: 3 (High). This is highly reliable because the cryogenic, solid-state environment of these grains inherently precludes active biological metabolism or ongoing aqueous redox chemistry (Postberg et al., 2018).

9. **Plume Ice Grains - Type III**
Scope Definition: This site includes the relatively large, salt-rich water ice grains formed via the direct flash-freezing of bulk ocean water spray at the subsurface liquid-vapor interface, specifically excluding salt-poor vapor-condensed grains (Type I) and grains dominated by complex macromolecular organics (Type II).

   a. Temperature (°C): -1 to -76
   Source: Thermomechanical models dictate that Type III grains originate from aerosolized liquid droplets at the subsurface water table, where temperatures are near the freezing point of moderately saline water, around 0 to -1 °C (Glein et al., 2018). Other studies agree that these droplets subsequently undergo rapid evaporative cooling and flash-freezing as they ascend through the icy conduits, reaching approximately -76 °C at the surface vents (Goguen et al., 2013). There is no disagreement regarding this gradient, as it is a thermodynamic requirement for liquid boiling into a vacuum.

   Reliability: 3 (High). This is highly reliable because it combines fundamental thermodynamic properties of saline water with Cassini's direct infrared thermal mapping of the fissures (Goguen et al., 2013).

b. pH: 8.5 to 11.6
Source: Group A says the pH is between 8.5 and 9.0 because this range best fits the sodium carbonate speciation directly measured in the ice grains by Cassini (Postberg et al., 2009, Glein & Waite, 2020). Group B says the pH is significantly more alkaline, between 10.1 and 11.6, because recent evaluations of the exact phosphate speciation found in these grains require a highly alkaline environment to remain chemically stable (Glein & Truong, 2025). The union range of 8.5 to 11.6 is adopted here to encompass both the established carbonate-based models and the most recent phosphate-based models.

Reliability: 2 (Medium). Although based on in-situ mass spectrometry, conflicting theoretical interpretations of carbonate versus phosphate speciation introduce uncertainty into the exact pH value.

c. Salinity (g/kg): 5 to 26
Source: In-situ measurements from Cassini identify Type III grains as salt-rich, containing 0.5 to 2 percent by mass of sodium and potassium salts, which equates to 5 to 20 g/kg (Postberg et al., 2009, Postberg et al., 2011). Other recent compositional models of Enceladus's ocean agree, predicting a bulk salinity between 8 and 26 g/kg based on the expected presence of ammonium and phosphates (Klenner et al., 2025). There is no significant disagreement, as all studies concur these grains represent frozen spray from a moderately saline ocean.

Reliability: 3 (High). This parameter is exceptionally reliable as it is derived from direct, in-situ mass spectrometry of thousands of individual salt-rich ice grains by Cassini (Postberg et al., 2009).

d. Pressure (MPa): ~0
Source: Hydrodynamic models of the Enceladus plume dictate that liquid water droplets form and flash-freeze at the liquid-vapor interface where the pressure corresponds to the triple point of water, approximately 0.0006 MPa (Nakajima & Ingersoll, 2016). Other physical simulations universally agree that these grains then experience a continuous pressure drop as they are propelled upward into the vacuum of space (Schmidt et al., 2008). There is no disagreement on this parameter, as it is a strict thermodynamic requirement for liquid water exposed to a vacuum.

Reliability: 3 (High). This pressure bounding is highly reliable because it is derived from fundamental thermodynamic laws governing the liquid-vapor interface of water venting to space (Nakajima & Ingersoll, 2016).

e. Isolation Score (1-5): 5
(i) Physical Confinement (1-5): 5
Source: As an extraterrestrial environment, Enceladus is separated from Earth by vast interplanetary distances, completely preventing any direct physical transfer of terrestrial biological contaminants or matter (Worth et al., 2013).

(ii) Hydrologic Connectivity (1-5): 5
Source: The subsurface ocean and hydrothermal systems of Enceladus are entirely self-contained within the moon, meaning there is an absolute lack of hydrological connectivity with Earth's ecosystems (Worth et al., 2013).

(iii) Isolation Time (1-5): 5
Source: Enceladus has been completely isolated from Earth since the formation of the solar system, ensuring any potential biosphere has evolved independently for billions of years without terrestrial contact (Worth et al., 2013).

Reliability: Score 3 (High). The absolute physical separation and lack of biological connectivity between Earth and the Saturnian system is a fundamental and undisputed astronomical reality.

f. Redox Score (1-5): 1.33
(i) Redox Species (1-5): 2
Source: Because Type III grains freeze directly from bulk ocean water, they capture oceanic salts, phosphates, and potentially trace dissolved reductants and oxidants, but these remain locked in a solid ice matrix (Postberg et al., 2009, Glein & Truong, 2025).

(ii) Equilibrium (1-5): 1
Source: The extreme cryogenic temperatures and rapid flash-freezing of the ocean spray completely halt chemical kinetics, preventing any active aqueous reactions or chemical equilibrium from being reached within the grain (Schmidt et al., 2008).

(iii) Chemolithotrophic Metabolism (1-5): 1
Source: Despite representing the nutrient-rich ocean, the completely frozen, solid-state nature of these grains in the vacuum of space provides zero liquid solvent to support any chemolithotrophic metabolism (Postberg et al., 2011).

Reliability: 3 (High). This is highly reliable because the cryogenic, solid-state environment of these grains inherently precludes active biological metabolism or ongoing aqueous redox chemistry in space.

References:

1. Affholder, A., Guyot, F., Sauterey, B., Ferrière, R. & Mazevet, S. (2021) Bayesian analysis of Enceladus's plume data to assess methanogenesis. Nature Astronomy. 5 (8), 805–814. doi:10.1038/s41550-021-01372-6.
2. Affholder, A., Guyot, F., Sauterey, B., Ferrière, R. & Mazevet, S. (2022) Putative Methanogenic Biosphere in Enceladus's Deep Ocean: Biomass, Productivity, and Implications for Detection. The Planetary Science Journal. 3 (12), 270. doi:10.3847/PSJ/aca275.
3. Barge, L.M. & White, L.M. (2017) Experimentally Testing Hydrothermal Vent Origin of Life on Enceladus and Other Icy/Ocean Worlds. Astrobiology. 17 (9), 820–833. doi:10.1089/ast.2016.1633.
4. Bouquet, A., Glein, C.R. & Waite, J.H. (2019) How Adsorption Affects the Gas–Ice Partitioning of Organics Erupted from Enceladus. The Astrophysical Journal. 873 (1), 28. doi:10.3847/1538-4357/ab0100.
5. Choblet, G., Tobie, G., Sotin, C., Běhounková, M., Čadek, O., Postberg, F. & Souček, O. (2017) Powering prolonged hydrothermal activity inside Enceladus. Nature Astronomy. 1 (12), 841–847. doi:10.1038/s41550-017-0289-8.
6. Ershova, A., Schmidt, J., Postberg, F., Khawaja, N., Nölle, L., Srama, R., Kempf, S. & Southworth, B. (2024) Modeling the Enceladus dust plume based on in situ measurements performed with the Cassini Cosmic Dust Analyzer. Astronomy & Astrophysics. 689, A114. doi:10.1051/0004-6361/202450429.
7. Fifer, L.M., Catling, D.C. & Toner, J.D. (2022) Chemical Fractionation Modeling of Plumes Indicates a Gas-rich, Moderately Alkaline Enceladus Ocean. The Planetary Science Journal. 3 (8), 191. doi:10.3847/PSJ/ac7a9f.
8. Glein, C.R., Baross, J.A. & Waite, J.H. (2015) The pH of Enceladus' ocean. Geochimica et Cosmochimica Acta. 162, 202–219. doi:10.1016/j.gca.2015.04.017.
9. Glein, C.R., Postberg, F. & Vance, S.D. (2018) The Geochemistry of Enceladus: Composition and Controls. In: Enceladus and the Icy Moons of Saturn. The University of Arizona Press. p. doi:10.2458/azu_uapress_9780816537075-ch003.
10. Glein, C.R. & Truong, N. (2025) Phosphates reveal high pH ocean water on Enceladus. Icarus. 441, 116717. doi:10.1016/j.icarus.2025.116717.
11. Glein, C.R. & Waite, J.H. (2020) The Carbonate Geochemistry of Enceladus' Ocean. Geophysical Research Letters. 47 (3), e2019GL085885. doi:10.1029/2019GL085885.
12. Goguen, J.D., Buratti, B.J., Brown, R.H., Clark, R.N., Nicholson, P.D., Hedman, M.M., Howell, R.R., Sotin, C., Cruikshank, D.P., Baines, K.H., Lawrence, K.J., Spencer, J.R. & Blackburn, D.G. (2013) The temperature and width of an active fissure on Enceladus measured with Cassini VIMS during the 14 April 2012 South Pole flyover. Icarus. 226 (1), 1128–1137. doi:10.1016/j.icarus.2013.07.012.
13. Hsu, H.-W., Postberg, F., Sekine, Y., Shibuya, T., Kempf, S., Horányi, M., Juhász, A., Altobelli, N., Suzuki, K., Masaki, Y., Kuwatani, T., Tachibana, S., Sirono, S., Moragas-Klostermeyer, G. & Srama, R. (2015) Ongoing hydrothermal activities within Enceladus. Nature. 519 (7542), 207–210. doi:10.1038/nature14262.
14. Ingersoll, A.P. & Pankine, A.A. (2010) Subsurface heat transfer on Enceladus: Conditions under which melting occurs. Icarus. 206 (2), 594–607. doi:10.1016/j.icarus.2009.09.015.

15. Khawaja, N., Postberg, F., Hillier, J., Klenner, F., Kempf, S., Nölle, L., Reviol, R., Zou, Z. & Srama, R. (2019a) Low-mass nitrogen-, oxygen-bearing, and aromatic compounds in Enceladean ice grains. Monthly Notices of the Royal Astronomical Society. 489 (4), 5231–5243. doi:10.1093/mnras/stz2280.
16. Khawaja, N., Postberg, F., Hillier, J., Klenner, F., Nölle, L. & Reviol, R. (2019b) Astrobiologically relevant precursor organic compounds from the subsurface ocean of Enceladus. In: 1 September 2019 p. EPSC-DPS2019-474. https://ui.adsabs.harvard.edu/abs/2019EPSC...13..474K.
17. Kite, E.S. & Rubin, A.M. (2016) Sustained eruptions on Enceladus explained by turbulent dissipation in tiger stripes. Proceedings of the National Academy of Sciences. 113 (15), 3972–3975. doi:10.1073/pnas.1520507113.
18. Klenner, F., Bönigk, J., Napoleoni, M., Hillier, J., Khawaja, N., Olsson-Francis, K., Cable, M.L., Malaska, M.J., Kempf, S., Abel, B. & Postberg, F. (2024) How to identify cell material in a single ice grain emitted from Enceladus or Europa. Science Advances. 10 (12), eadl0849. doi:10.1126/sciadv.adl0849.
19. M. Schenk, P., N. Clark, R., J. A. Howett, C., J. Verbiscer, A. & Waite, J.H. (2018) Enceladus and the Icy Moons of Saturn. The University of Arizona Press. doi:10.2458/azu_uapress_9780816537075.
20. McKay, C.P., Porco, C.C., Altheide, T., Davis, W.L. & Kral, T.A. (2008) The Possible Origin and Persistence of Life on Enceladus and Detection of Biomarkers in the Plume. Astrobiology. 8 (5), 909–919. doi:10.1089/ast.2008.0265.
21. Meyer, C.R., Buffo, J.J., Nimmo, F., Wells, A.J., Boury, S., Fox-Powell, M., Tomlinson, T.C., Parkinson, J.R.G. & Vasil, G.M. (2025) A Potential Mushy Source for the Geysers of Enceladus and Other Icy Satellites. Geophysical Research Letters. 52 (3), e2024GL111929. doi:10.1029/2024GL111929.
22. Nakajima, M. & Ingersoll, A.P. (2016) Controlled boiling on Enceladus. 1. Model of the vapor-driven jets. Icarus. 272, 309–318. doi:10.1016/j.icarus.2016.02.027.
23. Neveu, M. & Glein, Christopher & Anbar, A. & McKay, C. & Desch, Steven & Castillo-Rogez, J. & Tsou, P.. (2014). Enceladus' Fully Cracked Core: Implications for Habitability.
24. Postberg, F., Clark, R.N., Hansen, C.J., Coates, A.J., Dalle Ore, C.M., Scipioni, F., Hedman, M.M. & Waite, J.H. (2018a) Plume and Surface Composition of Enceladus. In: Enceladus and the Icy Moons of Saturn. The University of Arizona Press. p. doi:10.2458/azu_uapress_9780816537075-ch007.
25. Postberg, F., Kempf, S., Schmidt, J., Brilliantov, N., Beinsen, A., Abel, B., Buck, U. & Srama, R. (2009) Sodium salts in E-ring ice grains from an ocean below the surface of Enceladus. Nature. 459 (7250), 1098–1101. doi:10.1038/nature08046.
26. Postberg, F., Khawaja, N., Abel, B., Choblet, G., Glein, C.R., et al. (2018b) Macromolecular organic compounds from the depths of Enceladus. Nature. 558 (7711), 564–568. doi:10.1038/s41586-018-0246-4.
27. Postberg, F., Schmidt, J., Hillier, J., Kempf, S. & Srama, R. (2011) A salt-water reservoir as the source of a compositionally stratified plume on Enceladus. Nature. 474 (7353), 620–622. doi:10.1038/nature10175.
28. Ray, C., Glein, C. & Waite, H. (2021) Can abiotic synthesis generate complex organic matter on Enceladus? In: 1 December 2021 pp. P32A-02. https://ui.adsabs.harvard.edu/abs/2021AGUFM.P32A..02R.
29. Schmidt, J., Brilliantov, N., Spahn, F. & Kempf, S. (2008) Slow dust in Enceladus' plume from condensation and wall collisions in tiger stripe fractures. Nature. 451 (7179), 685–688. doi:10.1038/nature06491.

30. Sekine, Y., Shibuya, T., Postberg, F., Hsu, H.-W., Suzuki, K., Masaki, Y., Kuwatani, T., Mori, M., Hong, P.K., Yoshizaki, M., Tachibana, S. & Sirono, S. (2015) High-temperature water–rock interactions and hydrothermal environments in the chondrite-like core of Enceladus. Nature Communications. 6 (1), 8604. doi:10.1038/ncomms9604.
31. Taubner, R.-S., Pappenreiter, P., Zwicker, J., Smrzka, D., Pruckner, C., Kolar, P., Bernacchi, S., Seifert, A.H., Krajete, A., Bach, W., Peckmann, J., Paulik, C., Firneis, M.G., Schleper, C. & Rittmann, S.K.-M.R. (2018a) Biological methane production under putative Enceladus-like conditions. Nature Communications. 9 (1), 748. doi:10.1038/s41467-018-02876-y.
32. Taubner, R.-S., Pappenreiter, P., Zwicker, J., Smrzka, D., Pruckner, C., Kolar, P., Bernacchi, S., Seifert, A.H., Krajete, A., Bach, W., Peckmann, J., Paulik, C., Firneis, M.G., Schleper, C. & Rittmann, S.K.-M.R. (2018b) Simulating putative Enceladus-like conditions: The possibility of biological methane production on Saturn's icy moon. Proceedings of the International Astronomical Union. 14 (S345), 219–221. doi:10.1017/S1743921319001789.
33. Waite, J.H., Glein, C.R., Perryman, R.S., Teolis, B.D., Magee, B.A., Miller, G., Grimes, J., Perry, M.E., Miller, K.E., Bouquet, A., Lunine, J.I., Brockwell, T. & Bolton, S.J. (2017) Cassini finds molecular hydrogen in the Enceladus plume: Evidence for hydrothermal processes. Science. 356 (6334), 155–159. doi:10.1126/science.aai8703.
34. Worth, R.J., Sigurdsson, S. & House, C.H. (2013) Seeding Life on the Moons of the Outer Planets via Lithopanspermia. Astrobiology. 13 (12), 1155–1165. doi:10.1089/ast.2013.1028.
35. Yeoh, S.K., Chapman, T.A., Goldstein, D.B., Varghese, P.L. & Trafton, L.M. (2015) On understanding the physics of the Enceladus south polar plume via numerical simulation. Icarus. 253, 205–222. doi:10.1016/j.icarus.2015.02.020.