

بسم الله الرحمن الرحيم

قناة روح مريم التعليمية

Zainab's Textbook of Common Pediatric Dosages

Written By :

Dr. Sidiqa Mohamad Osman

Dr. Israa mustafa

Pediatric specialists/ Rufaa hospital

إهداء

إلى من كان لهم الفضل بعد الله في كل بداية جميلة، إلى والدينا الاعزاء

نهديكم هذا العمل المتواضع

ليس وفاء لحقكم فحقكم اعظم

ولكن تعبيراً عن حب يسكن القلوب

وامتنان لا تصفه الكلمات

كنتم الداعمين حين ضعفنا

والنور حين أظلم الطريق

والأمان الذي نلجأ إليه دائماً

نسأل الله أن يحفظكم

ويبارك فيكم

ويجعل كل ما قدمتموه في ميزان حسناتكم

بكل الحب والتقدير، ابنتيكم صديقة وإسراء

لو الكتاب دا كان مفيد بالنسبة ليك ما تنسى تدعي معاك لي يمة مريم
وزينوبة وجميع اموات المسلمين



A

- Adrenaline (epinephrine) :1
 - Anaphylaxis **IM** = 0.01mg/kg/dose.
 - Cardiac arrest = 0.01mg/kg/dose, i.v.
 - Nebulized = 0.5ml/kg of 1:1000. Max 5ml/session.
(1:1000, 1:10.000)
- Artesunate injection :
 - Less than 20 kg = wt x 3 per dose. 1,2,3
 - 20 kg or more = wt x 2.4 per dose. 1,2,3
(injection 30mg, 60mg, 120mg).
- Albumin :

Hypoalbuminemia 0.5 - 1g/kg/dose, for 1 - 2days. Max 25g/dose iv
over 120minutes.

- Aciclovir :

- Injection :

- regular dose = 10 mg/kg per dose, 8 hourly. 1, 2

- Sepsis or encephalitis :

- Birth to 12 yrs = 20mg/kg/dose, 8 hourly. 3

- More than 12yrs = 30mg/kg/day, 8 hourly.

- Tabs : 15 - 20mg/kg/dose, 5 times/day, Max 800mg. 1,2

(vial 200mg, 250mg, 500mg, 1g, tabs 200mg, 400mg).

- Albendazole tabs :

- Less than 2 yrs = 200mg single dose. 1

- 2 yrs or more = 400mg single dose. 1

(can be repeated 2 - 3 wks later).

(tabs 200mg, 400mg).

- Amikacin :

15 - 22.5mg/kg/day, Od, BD, or TDS. 1,2,3

(injection 250mg/ml, 500mg/ml).

- Aminophylline in asthma exacerbation :
 - Loading dose = 5 - 6 mg/kg in 20 - 30 minutes. ^{1,2}
 - Maintenance :
 - 1month - 11yrs = 1 mg/kg/hour.
 - 12yrs - 17yrs = 0.5 - 0.7mg/kg/hr

(vial 250mg/10ml, 500mg/20ml).

- Amlodipine :
 - Less than 6 yrs = 0.1 - 0.3 mg/kg once, Max 5mg/day. ³
 - 6 yrs or more = 0.05 - 0.2 mg/kg once, Max 10mg/day. ¹

(tabs 5mg, 10mg).

- Amoxicillin :
 - Regular dose = 25 - 50 mg/kg/day, BD or TDS. ¹
 - Tonsillitis & pharyngitis in 5 - 15yrs = 40mg/kg/day, BD, for 10days.⁴
 - Sever infection = 80 - 100mg/kg/day, BD. ²

(syrup 125mg/5ml, 250mg/5ml, Caps 250mg, 500mg).

- Ampicillin :
 - Regular dose = 50 - 100mg/kg/day, 6 hourly. 1
 - Meningitic = 200 - 400 mg/kg/day, 4 - 6 hourly. 1Max 8 - 12 g/day.
(vial 250mg, 500mg, 1g).

- Artemether with lumefantrine (Co-artem) 20/120 mg :
 - 2 to 14 kg = 1 tab per dose. 2
 - 15 to 24 kg = 2 tabs per dose.
 - 25 to 34 kg = 3 tabs per dose.
 - 35 and above = 4 tabs per dose (1 tab per dose of 80/480 mg).
(tabs 20/120mg, 80/480mg).

- Asprin (salicylic acid) :
 - Rhumatic fever :
 - anti-inflammatory 50 - 100mg/kg/day. 1
 - In local protocol we give 60mg/kg/day, for 2 wks then taper.
No max dose.
 - Kawasaki disease = 80 - 100mg/kg/day, 6hourly, during febrile phase - up to 14days - for 48hrs - 72hrs, then decrease to 3 - 5mg/kg/day.
 - Anti-platelet 3 - 5 mg/kg/day, (Max 75 - 100mg/day) 1
(tabs 75mg, 80mg, 100mg, 300mg).

- Azithromycine :
 - Pharyngitis & tonsillitis = 12mg/kg/day, od for 5 days.
 - Whooping cough & community Acquired pneumonia :
 - o Less than 6 months 10mg/kg/day, od for 5 days. 1,2
 - o > 6 months 10mg/kg/day, od on day 1, then 5mg/kg/day on the 4 subsequent days. 1,
- (caps 250mg, 500mg).

B

- Baclofen :
 - Start with = 0.3mg/kg/day, TDS or 6hourly.¹
 - Maintenance = 0.75mg - 2mg/kg/day.¹ Max 40 - 60mg/day.
(to increase every 3 days).
 - (tabs 5mg, 10mg, 25mg).
- Benzyle penicillin (penicillin G) :
 - Moderate/severe infection = 100.000 - 250.000 u/kg/day. 4 - 6hrs.
¹
 - Meningitic dose = 250.000 - 400.000 u/kg/day. ¹
(injection 1 million units, 5 million units).
- Blood transfusion :^{1,2,3}
 - Packed RBCs = 10 - 15ml/kg.
 - Whole blood = 20ml/kg. ⁴
 - Platelets = 10 - 15ml/kg.

C

- Calcium carbonate :
45 - 65mg/kg/day, 6hourly.
- Calcium gluconate :
 - As bolus in convulsions = 0.5 - 2ml/kg in 20minutes in Dextrose 10% (1 : 4 concentration. 1
 - Maintenance = 0.5 - 1.5 ml/kg/hour. 3
- Captopril tabs :
 - In Hypertension :
 - starting dose is 0.3 - 0.5 mg/kg/dose, TDS. 1
 - Maintenance = 1 - 6 mg/kg/day, BD or TDS. Max 6mg/kg/day.1
 - In heart failure = starting dose is 0.1 to 0.3 mg/kg/dose BD, or TDS, Max 4mg/kg/day.4
(tabs 6.25mg, 12.5mg, 25mg, 50mg)

- Carbamazepine tabs :
 - starting = 5mg/kg/day, BD. ^{1,3}
 - (Increasing by 5mg/kg/day every 3 - 7days.)
 - Maintenance = 20 - 30mg/kg/day. Max 35mg/kg/day.¹
 - (tabs 100mg, 200mg, 400mg, syrurp 100mg/5ml)

- Cefaclor (2nd generation cephalosprin) :

20 - 40mg/kg/day, 8 hourly, Max 1g/day.¹

(caps 250mg, 500mg, syrup 125mg/5ml, 250mg/ml)

- Cefadroxil (1st generation cephalosprin) :

30 - 50mg/kg/day, Od or BD. ¹

(syrup 125mg/5ml, 250mg/5ml, Caps 250mg, 500mg)

- Cefalexin, 1st generation cephalosprin :
 - regular dose :
 - Streptococcal pharyngitis & skin infection = 25 - 50mg/kg/day, BD or 6hourly.^{1, 3}
 - UTI = 12.5mg/kg/dose, BD. Max 1g/dose.
 - Sever infection = 50 - 100mg/kg/day, TDS or 6 hourly. (Max 4g/day). ^{1,3}

(syrup 125mg/5ml, 250mg/ml, caps 250mg, 500mg)

- Cefepime injection 4th generation cephalosprin :
50mg/kg/dose, TDS. Max 2g/dose.^{1,3}
(500mg, 1g, 2g)
- Cefpodoxime proxetil (cefodox), 3rd generation cephalosporin :
10mg/kg/day, BD, for 5 - 10days. Max 200mg/day.
In AOM & maxillary sinusitis Max 400mg/day.
- Cefixime, 3rd generation cephalosprin :
8mg/kg per day. Od or BD. Max 400mg/day.^{1,3}
Syrup 100mg/5ml (0.4ml/kg/day).
(caps 200mg, 400mg).
- Cefotaxime injection , 3rd generation cephalosprin :
 - Regular dose = 150 - 200mg/kg/day 6 - 8hourly. ^{1,3}
 - Meningitc dose = 200mg/kg/day 6 - 8hourly. Max 12g/dady. ¹
(vial 500mg, 1g)
- Ceftazidim (fortum)injection , 3rd generation cephalosprin :
 - Regular dose = 100 - 150mg/kg/day TDS. ³
 - Meningitic dose = 200mg/kg/day 8hourly. Max 6g/day. ¹
(vial 500mg, 1g).

- Ceftriaxone injection (Samixone), 3rd generation cephalosprin :
 - Mild/moderate = 50 - 75mg/kg/day. Od or BD. Max 2g/day. 1
 - Septic dose (sever infections) = 80 - 100mg/kg/day.1
 - Meningitic = 100mg/kg/day. Max 4g/day. 1

(vial 500mg, 1g)

- Cefuroxime injection (maxil), 2nd generation cephalosprin :
 - Mild/moderate = 75 - 100 mg/kg/day.TDS. 3
 - Severe infection = 100 - 200mg/kg/day, 6 - 8 hourly . Max 3g/day i.v.1

(vial 750mg)

- Cetirizine hydrochloride (2nd generation antihistamine) : 1
 - 6months - 2yrs = 2.5mg,od.
 - 2yrs - 6yrs = 2.5mg, BD or Od. Max 5mg.
 - 6yrs - 12yrs = 10mg, Devide BD or given Od.
 - More than 12yrs = 10mg, Od.

(syrup 5mg/5ml, tabs 10mg).

- Ciprofloxacin :

20 - 30 mg/kg/day, BD or TDS. Max 400mg/dose.1

(syrup 250mg/5ml, tabs 250mg, 500mg, infusions 2mg/ml)

- Clarithromycin :

15mg/kg/day. BD. Max 1g/day.^{1,3}

(Caps 250mg, 500mg, syrup 125mg/5ml, 250mg/5ml).

- Clindamycin :

20 - 40mg/kg/day, 6hourly or TDS. Max 2.7g/day.¹

(ampoules 150mg/1ml).

- Co-amoxiclav (Amocaln) :

➤ Regular dose = 25 - 45mg/kg/day. BD. ^{1,3}

➤ Severe infections (CAP, orbital cellulitis, AOM) = 80 - 90mg/kg/day, BD or TDS. ^{1,3}

(syrup 156mg/5ml, 200mg/ml, 400mg/5ml, tabs 375mg, 625mg, 1g).

- Co-trimoxazole syrp (septrin) :

➤ Mild - moderate infections = 8mg/kg/day, BD. ^{1,3}

➤ In immunocompromised pts = 15 - 20mg/kg/day, 6hourly or TDS. Max 320mg/day.^{1,3}

D

- Dexamethazone :1,3
 - In Croup = 0.15 - 0.6mg/kg/single dose. Max 10-12mg.
 - In Meningitis = 0.15mg/kg/dose, 6hourly (2 - 4 days). Max 16mg/day.
(4mg/ml, 8mg/2ml)
- Diazepam :
 - Acute seizures :
 - 0.1 - 0.3mg/kg/dose, i.v.1
 - Rectal = 0.5mg/kg/dose.1,3
(10mg/2ml)
- Diclofinac sodium, suppositories :
1 - 3mg/kg/day, Od, or BD. 3
(12.5mg, 25mg, 50mg, 100mg)

- Domperidone :3

0.2 - 0.4mg/kg/dose, TDS (before meals).

(syrup mg/ml, 5mg/5ml, tabs 10mg).

- Doxycyclin :

8 yrs and older = 2 - 4mg/kg/day, BD. Max 200mg/day.1

(Contraindicated in below 8yrs)

(caps 50mg, 100mg).

- Dobutamine :

5 - 20mics/kg/minute, infusion in NS of D5%. Max 30mic/kg/minutes.

(vial 250mg/20ml).

- Dopamin :

➤ Medium dose (cardiac effect) = 3 - 10mic/kg/min, increases contractility & cardiac output.

➤ High dose (vasopressor effect) = 10 - 20 mic/kg/min, vasoconstriction & increase blood pressure.

➤ Usual clinical range = 2 - 20mic/kg/min, infusion in NS or D5%. Max 20mg/mic/min.

(vial 200mg/5ml, 400mg/5ml).

E

- Erythromycin :
 - Whooping cough = 40 - 50mg/kg/day, 6hourly, for 14days.
 - Chlamydial conjunctivitis & pneumonia = 50mg/kg/day, 6hourly, for 14days. Max 2g/day.^{1, 2,3}(syrup 125mg/5ml, 250mg/5ml, tabs 250mg, 500mg).
- Esmoprazole :

0.5 - 1mg/kg/day, Od. 1

(caps 20mg, 40mg, ampule 40mg).

F

- Ferrous sulfate, (saferon) :1,3
 - Prophylaxis = 1 - 2mg/kg/day, Od or BD,TDS.
 - Therapeutic = 3 - 6mg/kg/day. Max 150 - 200mg/day
(syrup 50mg/5ml, tabs 200mg).
- Ferrous sulfate with Folic acid, (hematin) :
Same dose as above.
(syrup 30mg/5ml).
- Fluconazol :1,2,3
 - mucocutaneous candidiasis = day1 = 6mg/kg/od. Then
3mg/kg/Od.
 - Systemic invasive = 6 - 12mg/kg/od. Max 400 - 800mg/day.
(syrup 50mg/5ml, 200mg/5ml, tabs 50mg, 100mg, 150mg, 200mg,
infusion 2mg/ml).

- Folic acid :1

- Folate deficiency :

- o Infant = 0.5mg/day, od for 4months.

- o 1yr - 17yrs = 5mg, od for 4months

- Haemolytic anemia = 2.5 - 5mg, od.

- Pregnant & lactating women = 0.4mg/day, od.

- Prophylaxis :

- o 0.1mg/day in infants

- o 0.2 - 0.5mg/day in children.

(tabs 0.5mg, 1mg, 5mg, drops 0.2mg/ml).

- Furosemide (Lasix) :1,3

0.5mg - 1mg/kg/dose, od, BD, TDS or 6 hourly. Max 3mg/kg/day.

(tabs 20mg, 40mg, injection 20mg/2ml).

G

- Gentamicin :1

In normal patient = 5 - 7.5mg/kg/day, od or 8 hourly.

In malnutrition = 7.5mg/kg/day, od.

(ampoules 10mg/ml, 40mg/ml).

- Griseofulvin :

12.5mg/kg/day. Max 500mg/day.

- Glucose :1,3

➤ All pediatrics & neonate as bolus = 2ml/kg of D10%.

➤ Malnutrition :

- o Conscious : 50ml of D10%, orally or NG tube.

- o Unconscious = 5ml/kg D10% iv bolus.

(D5W = 5g/100ml, D10W = 10g/100ml).

H

- Haloperidol :1,3

Agitation = 0.01 - 0.02mg/kg/day, od or BD. Max 5 - 10mg/day.

(tabs 0.5mg, 1.5mg, 5mg, ampoules 5mg/ml)

- Hepatitis B immunoglobulin :1,3

➤ Post-exposure Prophylaxis = 0.06ml/kg **I.M.**

➤ In neonate = 0.5ml **I.M.**

(100iu/ml, 220iu/ml).

- Hydralazine hydrochloride :1,3

➤ Hypertensive crises = 0.1 - 0.5mg/kg (100 - 500mic/kg) slowly i.v.
4hourly or infusion if required.

➤ Chronic HTN = 0.5 - 1mg/kg/day, BD or 6hourly. Max 6mg/kg/day.

(injection 20mg/ml, tabs 25mg, 50mg, 100mg).

- Hydrocortisone :1,3

- Acute adrenal insufficiency :

- o Less than 1yr = 25mg, im, then divided 6hourly iv.
 - o 1 - 5yrs = 50mg, im, then divided 6hourly iv.
 - o 6yrs & more = 100mg, im, then divided 6hourly iv.

- Status asthmaticus :

- o 4mg/kg/day, 6hourly. Max 100mg/dose.

- Anti-inflammatory :

- o 1 - 5mg/kg/day, od or BD.

(injection 100mg, tabs 10mg, 20mg).

- Hyoscine butylbromide :1,3

0.3 - 0.6mg/kg/dose, i.v or oral. Max 20mg/kg/dose.

(tabs 10mg, injection 20mg/ml).

- Ibuprofen :1,2,3
 - Analgesic/antipyretic = 5 - 10mg/kg/dose, 6 - 8 hourly. Max lesser of 40mg/kg/day. Preferred in 6months & above only.
 - JRA (6months - 12yrs) = 30 - 50mg/kg/day, 6hourly. Max 800mg/dose Or 2400mg/day.
(tabs 200mg, 400mg).

- Ipratropium bromide :1,3,4
 - nebulizer in Moderate/severe asthma exacerbations :
 - Less than 5yrs = 250mic/dose, 3 doses then 6hourly. Max 1mg/day.
 - 5yrs & more = 500mic/dose, 3 doses then 6hourly. Max 2mg/day. (GINA guideline)
 - (Nebulized solution 250mic, 500mic).

- Itraconazol :1,2,3
 - Regular = 3 - 5mg/kg/day, od or BD.
 - Severe Systemic = 10mg/kg/day. Max 200 - 400mg/day.
(oral 10mg/ml, caps 100mg, 200mg).

- Imipenem :

- Age of 4wks - 3months = 100mg/kg/day, 6hourly.
- More than 3months = 60 - 100mg/kg/day, 6hourly. Max 4g/day.

- Ivermectin :1,3

To be given to Children 15kg or more only

- Scabies = 200mic/kg (0.2mg/kg) as a single dose, repeat after 7 - 14 days.

(tabs 3mg, 6mg, 12mg).

J

- ?

K

- Ketotifen : 3

For children 6 months and older.

- 6months - 3yrs = 0.5mg, BD.
 - 3yrs and more = 1mg, BD. Max 2mg Bd.
- (syrup 1mg/5ml, tabs 1mg).

L

- Lactulose :1,3

- Constipation :

- o 1 - 3ml/kg/day, od or Bd.

Or

- o 1months - 11months = 2.5ml, BD.

- o 1yr - 4yrs = 2.5 - 10ml, BD.

- o 5yrs - 17yrs = 5 - 20ml, BD

- Hepatic encephalopathy = 1 - 2ml/kg/dose, 6 - 8 hourly. Target to reach 2 - 3 soft stools/day :

- o Infant = 2.5 - 10ml/day, 6 - 8hourly.

- o Children & adolescents = 40 - 90ml/day, 6 - 8 hourly.

(syrup 10g/15ml).

- Lansoprazole :1

0.5 - 1mg/kg/day, od.

(caps 15mg, 30mg).

- Levetiracetam (kepra) :1

- Starting dose :

- o Less than 12yrs (less than 50kg) = 10 - 20mg/kg/day, od or BD. Increased by 10mg/kg/day every 3 - 7days.
- o 12yrs - 18yrs = 250mg, BD. Increasing by 500mg, BD every 7 - 14days.

- Maintenance :

- o Less than 50kg = 40mg/kg/day, Bd. Max 3000mg/day.
- o More than 50kg = 60mg/kg/day, BD. Max 1.5g, BD.

(syrup 100mg/ml, tabs 250mg, 500mg, 750mg, 1g)

- Labetalol :

Hypertensive emergency = 500mic/kg/hr (0.5mg/kg/hr). Max 3mg/kg/hr. Dilute to 1mg in 1ml or 2mg in 1ml, in D5% or D5%/NaCl0.18%.

- Levofloxacin :

6months - 5yrs = 10mg/kg/dose, BD. Max 500mg/day.

5yrs or more = 10mg/kg/dose, Od. Max 750mg/day.

- Levocetirizine hydrochloride (2nd generation) :

- 0.1mg/kg/day, once. Max 5mg/day.¹

Or

- 6months - 5yrs = 1.25mg, od.

- 6yrs - 11yrs = 2.5mg - 5mg, od.

- 12yrs and more = 5mg, od. ³

(syrup 0.5mg/ml, tabs 5mg).

- Levothyroxine sodium :¹

- newborns = 10 - 15mic/kg/day, od. Max 50mic/day.

- 1months - 1yr = 5mic/kg/day. Max 50mic/day.

- 2yrs - 11yrs = initially 50mic/day, **increased by 25mic every 2 - 4wks, maintenance 75 - 100mic.**

- 12yrs - 17yrs = initially 50mic/day, **increased by 25 - 50mic every 3 - 4wks, maintenance 100 - 200mic.**

(tabs 25mg, 50mg, 75mg, 100mg, 125mg, 150mg)

- Loratadine (2nd generation) :

- 0.2mg/kg/day, once. 1

Or

- 2 - 5yrs = 5mg, od.

- 6 and more 10mg, od.3

(syrup 5mg/5ml, tabs 10mg).

M

- Magnesium sulfate :1,3

Asthma exacerbations (2yrs - 17yrs) = 40 - 50mg/kg/dose, infusion over 20 - 30minutes. Max 2g.

(injection 50% solution = 500mg/ml, 20% solution = 200mg/ml).

- Mannitol :1,3

Raised Intracranial pressure/cerebral edema = 0.25 - 1g/kg/dose, over 30 - 60minutes, may repeat every 4 - 8hourly according to response.

(infusion 20% mannitol = 200mg/ml, 10% solution =100mg/ml).

- Mebendazole :1,3

➤ Ascaris, Hookworm (roundworm) , Trichuris (whipworm) infection = 100mg, BD for 3 days, then repeat after 2 - 4wks.

➤ Enterobius (pinworm) = 100mg single dose, repeat after 2 wks.

(Contraindicated in less than 2 yr)

(tabs 100mg, syrup 100mg/5ml).

- Mefenamic acid :1,3

5 - 10mg/kg/dose, 6 - 8hourly. Max 1500mg/day.

(syrup 50mg/5ml, tabs 250mg, 500mg).

- Meropenem :1,3

➤ Mild/moderate, fever neutropenia, empirical therapy = 20mg/kg/dose, 8hourly. Max 1g/dose.

➤ Meningitis/severe infection & cystic fibrosis pulmonary exacerbations = 40mg/kg/dose, 8 hourly. Max 2g/dose.

(vial 500mg, 1g).

- Methylprednisolone :1,3

pulse therapy = 10 - 30mg/kg/day once for 3 - 5 days. Max 1g/day.

(vial 40mg, 125mg, 500mg, 1g).

- Metoclopramide hydrochloride :1,3

0.1 - 0.15mg/kg/dose, TDS. Max 10mg/dose.

(syrup 5mg/5ml, tabs 10mg, injection 10mg/2ml).

- Metronidazole, (flagyle) :1,3
 - Giardiasis = 15 - 30mg/kg/day, TDS. For 5 - 7days. Max 750mg/day.
 - H.pylori = 20mg/kg/day, BD, for 14days. Max 1g/day.
 - Amoebiasis = 35 - 50mg/kg/day, TDS, for 10days. Max 750mg/dose.
 - Anaerobic infection :
 - Orally = 30 - 50mg/kg/day, 8 hourly. Max 2250mg/day.
 - IV = 22.5 - 40mg/kg/day, 6 - 8 hourly. Max 4g/day.
 - Malnutrition = 7.5mg/kg/dose, TDS.^{1, 2, 3,4}. Max 400 - 500mg
(syrup 200mg/5ml, tabs 200mg, 400mg, 500mg, infusion 500mg/100ml).

- Midazolam :1,3
 - As bolus in status epilepticus (preferred in 2months & older) :
 - Loading = 0.2mg/kg, followed by :
 - 0.05 - 0.1mg/kg/min slow i.v.
 (injection 1mg/ml, 5mg/ml).

- Montelukast :3

- Asthma control :

- o 6months - 5yrs = 4mg, od.
 - o 6yrs - 14yrs = 5mg, od.
 - o 15yrs or more = 10mg, od.

(granules 4mg sachet, chewable tabs 4mg, 5mg, normal tabs 10mg).

- Morphine sulfate :

- Cyanotic spells = 0.05mg/kg/dose, 8 hourly.

- Sickler :

- o Starting dose = 0.1 - 0.15mg/kg/dose, 4hourly, slowly iv.
 - o Maintenance = 0.04 - 0.07mg/kg/hr.

- Mupriocin :

2months or more for impetigo = apply small amount, TDS, for 5 - 10days.

N

- Neomycin sulfate :1,3

Hepatic encephalopathy = 50 - 100mg/kg/day, oral, 6hourly. Max 4g/day.

(tabs 500mg).

- Nifedipine :1,3

➤ HTN in acute state (immediate-release IR) = 0.25 - 5mg/kg/dose, 6 - 8hourly.

➤ Chronic HTN (sustained-release SR) = 0.5 - 3mg/kg/day, of or BD.

(tabs of IR 10mg, 20mg, tabs of SR 30mg, 60mg).

- Nitrofurantoin :1,3

➤ Prophylaxis in VUR = 1 - 2mg/kg, od, at night.

➤ Treatment UTI = 5 - 7mg/kg/day, 6 - 12hourly. Max 400mg/day.

(tabs 50mg, 100mg, syrup 25mg/5ml).

- Nystatin :1,3

100,000 unite (1ml), 6hourly.

(syrup 100,000 u/ml, cream for skin use 100,000 u/g).

O

- Omeprazole :

- 0.5 - 1mg/kg/day. Od or bd.¹

Or

- Less than 10kg = 5 - 10 mg, od.
 - 10 - 20kg = 10 mg, od.
 - More than 20kg = 20mg, od. Max 40mg/day.³
- (tabs 10mg, 20mg, 40mg, syrup 2mg/ml, injections 40mg).

- Ondansetron :

- 0.1 - 0.15mg/kg/dose, 8 - 12hourly. Max 4mg in young children, 8mg in Older and adolescents.^{1,3}

(Contraindicated in less than 6months)

Or

- Less than 15kg = 2mg.
- 15 - 30kg = 4mg.
- More than 30kg = 8mg.³

(syrup 4mg/5ml, tabs 4mg, 8mg, injections 2mg/ml).

P

- Paracetamol (panadol) :

10 - 15mg/kg/dose, 4 - 6 hourly. Max 60mg/kg/day (up to 75mg/kg/day for SHORT duration if needed). 1

(syrup 120mg/5ml, 250mg/5ml, tabs 250mg, 500mg, 1g, suppositories 60mg, 125mg, 250mg, 500mg).

- Pethidine hydrochloride :1,3

0.5 - 1mg/kg/dose, 4 - 6hourly if needed. Max 100mg/dose.

(injection 50mg/ml, 100mg/ml).

- Phenobarbital :

➤ Loading = 15 - 20mg/kg.1,3

➤ Maintenance = 2.5 - 5mg/kg/day, (most of the time we use 5mg/kg/day) od at night or BD. 3

(tabs 15mg, 30mg, 60mg, 100mg).

- Phenytoin :

- Loading = 15 - 20mg/kg. Max 1500mg/day.^{1, 3}
- Maintenance = 4 - 8mg/kg/day, (most of the time we use 5mg/kg/day) BD.¹
(injection 50mg/ml).

- Potassium chloride :

- Maintenance :
 - o Orally = 0.5 - 1.5ml/kg/day (1 - 2 mmol/kg).
 - o IV = 1ml/kg/day.
- Rapid correction : according to it's level, if :
 - o $3.5 - 3 = (\text{wt} \times 0.25) / 2 = \text{ml} / \text{over 1 hr.}$
 - o $2.9 - 2.5 = (\text{wt} \times 0.5) / 2 = \text{ml} / \text{over 2 hrs.}$
 - o $2.4 \text{ and less} = (\text{wt} \times 0.75) / 2 = \text{ml} / \text{over 3 hrs.}$ ⁴

(kcl ampule 20mmol/10ml).

- Praziquantel :^{1,3}

- Schistosomiasis = 40 - 60mg/kg/day, 4 - 6 hourly for 1day.^{1,3} Or single dose.²
- Taeniasis = 5 - 19mg/kg/single dose.
(tabs 150mg, 600mg).

- Prednisolone :1,3

Mild/moderate asthma = 1 - 2mg/kg/day, od or BD. Max 20mg in less than 3yrs, 40mg in less than 12 yrs for 5 - 7 days. (Gina guideline).

(tabs 5mg, 10mg, 20mg, syrup 5mg/5ml, 15mg/5ml).

- Primaquine :1, 3

0.5mg/kg/day, od, for 14days. Max 30mg/day.

(contraindicated in less than 6months)

(tabs 7.5mg, 15mg).

- Propranolol hydrochloride :1,3

➤ In TOF = 0.2 - 0.4mg/kg/dose, BD or TDS. Increase gradually to 1 - 1.5mg/kg/dose (common side effect Hypotension & hypoglycemia).

➤ Thyrotoxicosis = 0.5 - 2mg/kg/day, 8hourly. Max 40mg/dose

➤ Infantile hemangioma :

o Start with 1mg/kg/day, TDS.

o Maintenance (24hrs later) = 2mg/kg/day, BD or TDS.

➤ Migraine Prophylaxis :

o 2yrs - 11yrs = 10 - 20mg, BD. Max 2mg/kg BD.

o 12yrs - 17yrs = 20 - 40mg, BD.

(tabs 10mg, 40mg, 80mg, syrup 5mg/5ml).

- Pyridoxine :
 - pyridoxin-dependent seizures :
 - o Loading = 50 - 10mg/dose, im or rapid iv.
 - o Maintenance = 50 - 100mg/day, orally.

Q

- Quinine :1,2,3
 - Loading dose = 20mg/kg over 4 hours. Deluted in D5%, infusion over 4hrs, amount of fluids same as quinine dose.
 - Maintenance = 10mg/kg, 8hourly.

(any more than 6 doses, you should reduce maintenance to 7mg/kg/dose).

(In local protocol we start maintenance direct)

(injection 600mg/2ml, tabs 300mg).

R

- Rantidine :
 - Oral = 2 - 4mg/kg/dose, BD. Max 300mg/day.^{1, 3}
 - Injectable = 1mg/kg/dose, 6 - 8hourly.(syrup 75mg/5ml, tabs 150mg, 300mg, injections 25mg/ml).

S

- Salbutamole :1,3

Nebulization :

- o less than 5 yrs = 2.5mg/dose.
- o 5yrs - 11yrs = 2.5 - 5mg/dose.
- o 12yrs & more = 5mg.

(nebules 2.5mg/2.5ml, 5mg/2.5ml).

- Sildenafil :1,3

Pulmonary HTN = 0.25 - 2mg/kg/dose, 6 - 8hourly. Max 60mg/day.

(tabs 25mg, 50mg, 100mg).

- Sodium nitroprusside :1,3

HTN emergency = start with = 0.3 - 0.5mic/kg/minute up to = 0.5 - 5mic/kg/minute. Max 8mic/kg/minutes.

(injection 50mg **diluted in D5% & protected from light**).

- Sodium valproate :1,3

- Epilepsy initially = 10 - 15mg/kg/day. Max 60mg/kg/day.

- Increment = 5 - 10mg/kg/day. Max 60mg/kg/day.

- Maintenance = 20 - 60mg/kg/day, BD.

(syrup 200mg/5ml, 250mg/5ml, tabs 200mg, 300mg, 500mg).

- Spironolactone (aldactone) :1,3

1 - 3mg/kg/day, od or Bd. Max 100mg/day.

(tabs 25mg, 50mg, 100mg).

T

- Tobramycin :
 - ophthalmic drop = 1 - 2 drops, BD for 6 - 8days.
 - Severe infection = 2 drops, 6hourly, **then taper.**
- Thiamine
Deficiency = 10 - 50mg/day.¹
(tabs 25mg, 50mg, 100mg).
- Tinidazole :^{1,3}
 - Giardiasis = 50mg/kg/, once.
 - Amoebiasis = 50mg/kg/day, od for 3 - 5days.
(Contraindicated in less than 3 yrs)(tabs 300mg, 500mg).
- Tranexamic acid :^{1,3}
10 - 15mg/kg/dose, 6 - 8hourly. Max 1g/dose.
(tabs 500mg, 650mg).

U

- Ursodexoycholic acid :1,3

10 - 15mg/kg/day, od or Bd or TDS. Max 20mg/kg/day.

(tabs 150mg, 300mg, 500mg).

V

- Vancomycin :1,3

40 - 60mg/kg/day, 6 - 8hourly.

(We give 45mg/kg/day due renal issues and facilities)

(vial 500mg, 1g).

- Vitamin A :

- Prophylaxis :

- o less than 6months = 50,000i.u, od.

- o 6months - 11months = 100,000i.u, od.

- o 12months and above = 200,000i.u, od. 4

- Therapeutic & measles = same doses with 3 intervals = day1, day2, and day14. 1,3

- Vitamin D3:1,3

- Prophylaxis :

- o 1yr of life = 400i.u, od.

- o In second yr of life and more = 600i.u, od.

- Therapeutic = 1,000 - 10,000i.u/day

- Or

- 50,000i.u weekly for 6 - 8wks.

- (caps 1000i.u, 2000i.u, 5000i.u, 50,000i.u, 60,000i.u).

- Vitamin E :1,3

- 5 - 10mg/kg/day.

- (caps 100i.u, 200i.u, 400i.u)

- Vitamin K :1,3

- Prophylaxis = 1mg, i.m, once. (wt less than 2500 give 400mic/kg).

- Therapeutic = 0.1 - 2mg/kg/day, i.v. Max 10mg/day.

W

• ?

X

• ?

Y

• ?

Z

- Zinc sulfate :1,2,3
 - Less than 6months = 10mg, od.
 - 6months or more = 20mg/day, od. For 10 - 14days.
- (syrup 12mg/5ml, 20mg/5ml, tabs 20mg).

Reference :

1. Nelson Textbook of Pediatrics.
2. World Health Organization WHO.
3. British National Formulary BNF.
4. Local Portocol.
5. Harriet Lane.
6. Oxford Nephrology.
7. Oxford Neurology.
8. NICE Guidelines.
9. Gina guideline

Neonatology drug section

(all doses are taken from NeoFax 2026)

- Acetaminophen (panadole) :

Route	Less than 32wks	32wks - 37	Term
Intravenous	10mg/kg/dose. Max 40mg/day/6hr	12.5mg/kg/dose. Max 50mg/day/6hr	Same
Oral	Loading : 20 - 25mg/kg Maintenance = 12 - 15mg/kg/dose/BD	L = 20 - 25mg/kg M = 12 - 15mg/kg/dose/TDS	L = 20 - 25mg/kg M = 12 - 15mg/kg/dose/6hr
Special situation	15mg/kg/dose, iv or orally, 6hrs for 3 days in active PDA in preterm		

- Acyclovir :

Dosage	Less than 34wks, postmenstrual	More than 34wks
	20mg/kg/BD	20mg/kg/TDS

- Aminophylline :
 - Loading = 5 - 8mg/kg.
 - Maintenance = 1 - 3mg/kg/dose, BD or TDS or 6hourly.

(start maintenance after 12hrs from loading dose).

- Ampicillin :

Gestational age	Postnatal age	Dosage
34wks or less	7days and less	50mg/kg/dose/BD
	8 days - 28days	75mg/kg/dose/BD
More than 34wks	28days or less	50mg/kg/dose/TDS

- Artesunate :
2.4 mg/kg.

- Azithromycine :
 - whooping cough = 10mg/kg/dose, od, for 5days.
 - Ophthalmia neonatorum = 20mg/kg/dose, od, for 3days.

- Benzyle penicillin :

Gestational age	Postnatal age	Dose	Regimen
Regular dose			
32 - 34wks	Less than 7days	100.000u/kg/dose	12hrs
	7 - 28days	100.000u/kg/dose	8hrs
34 wks & more	Less than 7 days	100.000u/kg/dose	8hrs
	7 - 28days	100.000u/kg/dose	6hrs
Meningitc dose			
All ages	7days & younger	150.000u/kg/dose	8hrs
	8days & more	125.000u/kg/dose	6hrs

- Cefotaxime :

Gestational age	Postnatal age	Dosage
All weeks	Less than 7days	50mg/kg/dose/BD
Less than 32wks	7days and more	50mg/kg/dose/BD
32wks or more	7days and more	50mg/kg/dose/6hourly

- Ciprofloxacin :

Gestational age	Postnatal age	Dosage
32wks - 34wks	Less than 7 days	7.5mg/kg/dose/BD
32wks - 34wks	8 - 14days	12.5mg/kg/dose/BD
34wks and more	All age	12.5mg/kg/dose/BD

- Erythromycin :

- Whooping cough = 10 - 12.5mg/kg/dose, 6hourly.
- Chlamydia conjunctivitis = 50mg/kg/day, 6hourly for 14days.

- Ferrous sulfate :

	Less than 37wks	37wks and more
Prophylaxis	2 - 3mg/kg/day	1mg/kg/day
Therapeutic	3 - 6mg/kg/day	

- Folic acid :

- Preterm = 25 - 50mic/kg/day. Max 65mic/day.
- Term = 65mic/day.

- Gentamicin :

Gestational age	Postnatal age	Dosage	Regmien in hours
29wks or less	0 - 7days	5	48hrs
	8 - 28days	4	36hrs
	More than 28days	4	24hrs
30 - 34wks	0 - 7days	4.5	36hrs
	8 and more	4	24 hrs
35wks or more	All age	4	24 hrs

- Meropenem :

Gestational age	Postnatal age	Usual dose	Meningitc dose
Less than 32wks	Less than 14days	20mg/kg/dose/BD	40mg/kg/dose/BD
	14 days or more	20mg/kg/dose/TDS	40mg/kg/dose/TDS
32wks and and more	Less than 14days	20mg/kg/dose/TDS	40mg/kg/dose/TDS for all ages
	14days and more	30mg/kg/dose/TDS	

- Metronidazole :

Gestational age	Loading	Maintenance	Regimen
24 - 25wks	15mg/kg	7.5mg/kg	24hrs
26 - 27wks	15mg/kg	10mg/kg	24hrs
28 - 33wks	15mg/kg	7.5mg/kg	12hrs
34 - 40wks	15mg/kg	7.5mg/kg	8hrs
More than 40wks	15mg/kg	7.5mg/kg	6hrs

- Omeprazole :

0.5 - 1.5mg/kg/od.

- Phenobarbital :

➤ Loading = 20mg/kg.

(Second loading dose if convulsion didn't stop)

➤ Maintenance = 3 - 5mg/kg/day, od or BD.

- Phenytoin :

➤ Loading = 15 - 20mg/kg.

➤ Maintenance = 4 - 8mg/kg/day, od or BD.

- Predinslon :

Weaning from oxygen = 2mg/kg/day, Bd for 5days, then 1mg/kg/day, od for 3days, then 1mg/kg/day every other day for 3 doses.

- Rantidine :

➤ Oral = 2mg/kg/dose, TDS

Or iv :

➤ Term = 1.5mg/kg/dose, TDS.

➤ Preterm = 0.5mg/kg/dose, BD.

- Vancomycin :

10 - 15mg/kg/dose.

Gestational age	Postnatal age	Regimen
29wks & less	0 - 14days	18hrs
	Older than 14days	12hrs
30 - 36wks	0 - 14days	12hrs
	Older than 14days	8hrs
37 - 44wks	0 - 7days	12hrs
	Older than 7days	8hrs
45wks & more	All	6hrs

- Vitamin K :

➤ Less than 1500g = 0.3 - 0.5mg/kg, within 6 hrs.

➤ More than 1500 = 1mg/day