


Stern-Gerlach Experiment

Luca Cicu - Lorenzo Liuzzo

27/11/2025

Experiment description

Stern-Gerlach Experiment Apparatus



Angular & Magnetic momenta





Figure 2: **Angular** and **magnetic** momenta generated by an electron orbiting circularly.

Angular & Magnetic momenta




Angular momentum $\mathbf{L} = \mathbf{r} \times \mathbf{p}$

Magnetic momentum $\boldsymbol{\mu} = I \Sigma \hat{\mathbf{n}}$

Figure 2: **Angular** and **magnetic** momenta generated by an electron orbiting circularly.

Angular & Magnetic momenta



Angular momentum $\mathbf{L} = \mathbf{x} \times \mathbf{p}$

Magnetic momentum $\boldsymbol{\mu} = I \Sigma \hat{\mathbf{n}}$

$$\boldsymbol{\mu} = \frac{qv}{2\pi r} \pi r^2 \hat{\mathbf{n}} = \frac{q}{2m} \mathbf{L} \Rightarrow \boldsymbol{\mu} \propto \mathbf{L}$$

Figure 2: **Angular** and **magnetic** momenta generated by an electron orbiting circularly.

Angular & Magnetic momenta




Figure 2: **Angular** and **magnetic** momenta generated by an electron orbiting circularly.

Angular momentum $\mathbf{L} = \mathbf{x} \times \mathbf{p}$

Magnetic momentum $\boldsymbol{\mu} = I \Sigma \hat{\mathbf{n}}$


$$\boldsymbol{\mu} = \frac{qv}{2\pi r} \pi r^2 \hat{\mathbf{n}} = \frac{q}{2m} \mathbf{L} \Rightarrow \boldsymbol{\mu} \propto \mathbf{L}$$

Magnetic Hamiltonian

$$H_{\text{magn}} = -\boldsymbol{\mu} \cdot \mathbf{B}$$

Stern-Gerlach Experiment

Magnetic Interaction




$$F_{\text{magn}} = -\nabla H_{\text{magn}} = \nabla(\mu \cdot B)$$

Figure 3: Microscopic origin of the magnetic force.

Stern-Gerlach Experiment

Magnetic Interaction



$$F_{\text{magn}} = -\nabla H_{\text{magn}} = \nabla(\mu \cdot B)$$

$$F_z = \mu_z \frac{\partial B_z}{\partial z}$$

Figure 3: Microscopic origin of the magnetic force.

Stern-Gerlach Experiment

Magnetic Interaction




Figure 3: Microscopic origin of the magnetic force.


$$F_{\text{magn}} = -\nabla H_{\text{magn}} = \nabla(\mu \cdot B)$$

$$F_z = \mu_z \frac{\partial B_z}{\partial z} = m\ddot{z}$$

$$z(t) = z_0 + \dot{z}_0 t + \frac{\mu_z}{2m} \frac{\partial B_z}{\partial z} t^2$$

Stern-Gerlach Experiment

Classical Results




classically predicted

Diocane

Stern-Gerlach Experiment

Observed Results




observed

Why two deflected spots?

Historical Context

Stern-Gerlach Experiment

Historical Context



Quantum Mechanics

QM — States

$|\psi\rangle \in \mathbb{H}$ **Hilbert space**

QM — States

$|\psi\rangle \in \mathbb{H}$ **Hilbert space**

- **Basis Properties**

$$\langle e_i | e_j \rangle = \delta_{ij} \qquad I = \sum_i |e_i\rangle \langle e_i|$$

QM — States

$|\psi\rangle \in \mathbb{H}$ **Hilbert space**

- **Basis Properties**

$$\langle e_i | e_j \rangle = \delta_{ij} \quad I = \sum_i |e_i\rangle \langle e_i|$$

- **Superposition Principle**

$$|\psi\rangle = \sum_i |e_i\rangle \langle e_i | \psi \rangle = \sum_i c_i^\psi |e_i\rangle$$

$c_i^\psi = \langle e_i | \psi \rangle$ are **probability amplitudes**

Stern-Gerlach Experiment

QM — States

$|\psi\rangle \in \mathbb{H}$ **Hilbert space**

- **Basis Properties**

$$\langle e_i | e_j \rangle = \delta_{ij} \quad I = \sum_i |e_i\rangle \langle e_i|$$

- **Superposition Principle**

$$|\psi\rangle = \sum_i |e_i\rangle \langle e_i | \psi \rangle = \sum_i c_i^\psi |e_i\rangle$$

$c_i^\psi = \langle e_i | \psi \rangle$ are **probability amplitudes**

- **Inner product**

$$\langle \varphi | \psi \rangle = \sum_i \langle \varphi | e_i \rangle \langle e_i | \psi \rangle = \sum_i (c_i^\varphi)^* c_i^\psi$$

Stern-Gerlach Experiment

QM — States

$|\psi\rangle \in \mathbb{H}$ **Hilbert space**

- **Basis Properties**

$$\langle e_i | e_j \rangle = \delta_{ij} \quad I = \sum_i |e_i\rangle \langle e_i|$$

- **Superposition Principle**

$$|\psi\rangle = \sum_i |e_i\rangle \langle e_i | \psi \rangle = \sum_i c_i^\psi |e_i\rangle$$

$c_i^\psi = \langle e_i | \psi \rangle$ are **probability amplitudes**

- **Inner product**

$$\langle \varphi | \psi \rangle = \sum_i \langle \varphi | e_i \rangle \langle e_i | \psi \rangle = \sum_i (c_i^\varphi)^* c_i^\psi$$

- **Born's rule**

$$\langle \psi | \psi \rangle = \sum_i (c_i^\psi)^* c_i^\psi = \sum_i |c_i^\psi|^2 = 1$$

$|c_i^\psi|^2$ is the **probability** to find $|\psi\rangle$ in $|e_i\rangle$

QM — Operators

A **linear operator** $\hat{O}$ is a transformation from an Hilbert space into itself, that acts like

$$\hat{O}(\alpha|\psi\rangle + \beta|\varphi\rangle) = \alpha\hat{O}|\psi\rangle + \beta\hat{O}|\varphi\rangle = \alpha|\psi'\rangle + \beta|\varphi'\rangle$$

Stern-Gerlach Experiment

QM — Operators

A **linear operator** $\hat{O}$ is a transformation from an Hilbert space into itself, that acts like

$$\hat{O}(\alpha|\psi\rangle + \beta|\varphi\rangle) = \alpha\hat{O}|\psi\rangle + \beta\hat{O}|\varphi\rangle = \alpha|\psi'\rangle + \beta|\varphi'\rangle$$

The action of $\hat{O}$ on a state can be entirely represented by its action on some **basis**:

$$|\psi'\rangle = \hat{O}|\psi\rangle = \sum_i \hat{O}|e_i\rangle\langle e_i|\psi\rangle = \sum_{ij} |e_j\rangle\langle e_j|\hat{O}|e_i\rangle\langle e_i|\psi\rangle = \sum_{ij} |e_j\rangle O_{ij} c_i^\psi$$

Stern-Gerlach Experiment

QM — Operators

A **linear operator** $\hat{O}$ is a transformation from an Hilbert space into itself, that acts like

$$\hat{O}(\alpha|\psi\rangle + \beta|\varphi\rangle) = \alpha\hat{O}|\psi\rangle + \beta\hat{O}|\varphi\rangle = \alpha|\psi'\rangle + \beta|\varphi'\rangle$$

The action of $\hat{O}$ on a state can be entirely represented by its action on some **basis**:

$$|\psi'\rangle = \hat{O}|\psi\rangle = \sum_i \hat{O}|e_i\rangle\langle e_i|\psi\rangle = \sum_{ij} |e_j\rangle\langle e_j|\hat{O}|e_i\rangle\langle e_i|\psi\rangle = \sum_{ij} |e_j\rangle O_{ij} c_i^\psi$$

We can also define the **adjoint** operator $\hat{O}^\dagger$: $\langle\psi'| = \langle\psi|\hat{O}^\dagger$ $\hat{O}_{ij}^\dagger = (\hat{O}_{ji})^*$

QM — Operators & Observables

We can associate every **observable** of a system to an operator

$$\hat{O} = \sum_i \lambda_i |e_i\rangle \langle e_i| \quad \hat{O}|e_j\rangle = \sum_i \lambda_i |e_i\rangle \langle e_i|e_j\rangle = \lambda_j |e_j\rangle$$

QM — Operators & Observables

We can associate every **observable** of a system to an operator

$$\hat{O} = \sum_i \lambda_i |e_i\rangle \langle e_i| \quad \hat{O}|e_j\rangle = \sum_i \lambda_i |e_i\rangle \langle e_i|e_j\rangle = \lambda_j |e_j\rangle$$

where λ_i are the **possible results** and $|e_i\rangle$ are the **possible states** in which the system is founded **after** the measurement.

The requirements are that, **independently by the choice of the basis**,

$$\langle e_i|e_j\rangle = \delta_{ij} \quad \lambda_i^* = \lambda_i$$

QM — Operators & Observables

We can associate every **observable** of a system to an operator

$$\hat{O} = \sum_i \lambda_i |e_i\rangle \langle e_i| \quad \hat{O}|e_j\rangle = \sum_i \lambda_i |e_i\rangle \langle e_i|e_j\rangle = \lambda_j |e_j\rangle$$

where λ_i are the **possible results** and $|e_i\rangle$ are the **possible states** in which the system is founded **after** the measurement.

The requirements are that, **independently by the choice of the basis**,

$$\langle e_i|e_j\rangle = \delta_{ij} \quad \lambda_i^* = \lambda_i$$

Such operators are called **hermitian** and satisfy $\hat{O}_{ij}^\dagger = \lambda_i^* \delta_{ij} = \lambda_i \delta_{ij} = \hat{O}_{ij}$

QM — Commutator of Operators

We define the **commutator** between two operators $\hat{A}$ and $\hat{B}$ as

$$[\hat{A}, \hat{B}] = \hat{A}\hat{B} - \hat{B}\hat{A} = -[\hat{B}, \hat{A}]$$

QM — Commutator of Operators

We define the **commutator** between two operators $\hat{A}$ and $\hat{B}$ as

$$[\hat{A}, \hat{B}] = \hat{A}\hat{B} - \hat{B}\hat{A} = -[\hat{B}, \hat{A}]$$

$\hat{A}, \hat{B}$ are said to be **compatible iff they commute**, so that $[\hat{A}, \hat{B}] = 0$.

QM — Commutator of Operators

We define the **commutator** between two operators $\hat{A}$ and $\hat{B}$ as

$$[\hat{A}, \hat{B}] = \hat{A}\hat{B} - \hat{B}\hat{A} = -[\hat{B}, \hat{A}]$$

$\hat{A}, \hat{B}$ are said to be **compatible iff they commute**, so that $[\hat{A}, \hat{B}] = 0$.

Two observables can be **simultaneously** determined iff their operators are compatible, that is if they share a **common basis** of eigenstates:

$$\hat{A}|e_i\rangle = \lambda_i|e_i\rangle \quad \hat{B}|e_i\rangle = \mu_i|e_i\rangle \quad \Leftrightarrow \quad \langle e_i | [\hat{A}, \hat{B}] | e_j \rangle = (\lambda_i \mu_i - \lambda_j \mu_j) \langle e_i | e_j \rangle = 0$$

Position and impulse
don't commute!

$$[\hat{p}, \hat{x}] = -i\hbar$$

QM — Uncertainty Principle

We define the **uncertainty** of an operator $\hat{A}$ in a state $|\psi\rangle$ as its **standard deviation**:

$$\Delta^2 \hat{A}_\psi = \langle \hat{A}^2 \rangle_\psi - (\langle \hat{A} \rangle_\psi)^2 = \langle (\hat{A} - \langle \hat{A} \rangle_\psi)^2 \rangle_\psi$$

QM — Uncertainty Principle

We define the **uncertainty** of an operator $\hat{A}$ in a state $|\psi\rangle$ as its **standard deviation**:

$$\Delta^2 \hat{A}_\psi = \langle \hat{A}^2 \rangle_\psi - (\langle \hat{A} \rangle_\psi)^2 = \langle (\hat{A} - \langle \hat{A} \rangle_\psi)^2 \rangle_\psi$$

The uncertainty of a pair of operators must satisfy the **uncertainty principle**:

$$\frac{1}{4} |\langle [\hat{A}, \hat{B}] \rangle_\psi|^2 \leq |\langle \hat{A}\hat{B} \rangle_\psi|^2 \leq \langle \hat{A}^2 \rangle_\psi \langle \hat{B}^2 \rangle_\psi$$

Heisenberg's Principle

$$\Delta^2 \hat{x} \Delta^2 \hat{p} \geq \frac{\hbar^2}{4}$$

QM — Angular Momentum

$$\hat{L}_k = (\hat{\mathbf{x}} \times \hat{\mathbf{p}})_k = \varepsilon_{kij} \hat{x}_i \hat{p}_j$$

QM — Angular Momentum

$$\hat{L}_k = (\hat{\mathbf{x}} \times \hat{\mathbf{p}})_k = \varepsilon_{kij} \hat{x}_i \hat{p}_j$$

Commutation rules

$$[\hat{L}_i, \hat{L}_j] = i\hbar \varepsilon_{ijk} \hat{L}_k \quad [\hat{L}_i, \hat{L}^2] = 0$$

QM — Angular Momentum

$$\hat{L}_k = (\hat{\mathbf{x}} \times \hat{\mathbf{p}})_k = \varepsilon_{kij} \hat{x}_i \hat{p}_j$$

Commutation rules

$$[\hat{L}_i, \hat{L}_j] = i\hbar \varepsilon_{ijk} \hat{L}_k \quad [\hat{L}_i, \hat{L}^2] = 0$$

Spectrum

$$\hat{L}^2 |lm\rangle = \hbar^2 l(l+1) |lm\rangle \quad l \in \mathbb{N}$$

$$\hat{L}_z |lm\rangle = \hbar m |lm\rangle \quad -l \leq m \leq l$$

QM — Angular Momentum

$$\hat{L}_k = (\hat{\mathbf{x}} \times \hat{\mathbf{p}})_k = \varepsilon_{kij} \hat{x}_i \hat{p}_j$$

Commutation rules

$$[\hat{L}_i, \hat{L}_j] = i\hbar \varepsilon_{ijk} \hat{L}_k \quad [\hat{L}_i, \hat{L}^2] = 0$$


Spectrum

$$\hat{L}^2 |lm\rangle = \hbar^2 l(l+1) |lm\rangle \quad l \in \mathbb{N}$$

$$\hat{L}_z |lm\rangle = \hbar m |lm\rangle \quad -l \leq m \leq l$$

$$\text{Degeneracy} \quad d_l = 2l + 1$$

$$|\psi\rangle = |lm_l\rangle$$




observed


$$|\psi\rangle = |lm_l\rangle$$

We can rewrite the magnetic moment of e^-

$$\hat{\mu} = \frac{\hbar q_e}{2m_e} \frac{\hat{\mathbf{L}}}{\hbar} = -g_l \mu_B \frac{\hat{\mathbf{L}}}{\hbar}$$



observed




$$|\psi\rangle = |lm_l\rangle$$

We can rewrite the magnetic moment of e^-

$$\hat{\mu} = \frac{\hbar q_e}{2m_e} \frac{\hat{\mathbf{L}}}{\hbar} = -g_l \mu_B \frac{\hat{\mathbf{L}}}{\hbar}$$

$$\langle \hat{H} \rangle_\psi = -\langle \hat{\mu}_z B_z \rangle_\psi = g_l \mu_B B_z \frac{1}{\hbar} \langle \hat{L}_z \rangle_\psi$$



$$|\psi\rangle = |lm_l\rangle$$

We can rewrite the magnetic moment of e^-

$$\hat{\mu} = \frac{\hbar q_e}{2m_e} \frac{\hat{\mathbf{L}}}{\hbar} = -g_l \mu_B \frac{\hat{\mathbf{L}}}{\hbar}$$

$$\langle \hat{H} \rangle_\psi = -\langle \hat{\mu}_z B_z \rangle_\psi = g_l \mu_B B_z \frac{1}{\hbar} \langle \hat{L}_z \rangle_\psi$$

But $\hat{L}_z$ exhibits an odd number of discrete eigenvalues $(2l + 1)$, so...

Why two deflected spots?

Spin

QM — Intrinsic Angular Momentum

Commutation rules

$$[\hat{S}_i, \hat{S}_j] = i\hbar\epsilon_{ijk}\hat{S}_k \quad [\hat{S}_i, \hat{S}^2] = 0$$

QM — Intrinsic Angular Momentum

Commutation rules

$$[\hat{S}_i, \hat{S}_j] = i\hbar\epsilon_{ijk}\hat{S}_k \quad [\hat{S}_i, \hat{S}^2] = 0$$

Spectrum

$$\hat{S}^2|sm_s\rangle = \hbar^2 s(s+1)|sm_s\rangle$$

$$\hat{S}_z|sm_s\rangle = \hbar m_s|sm_s\rangle \quad -s \leq m_s \leq s$$

QM — Intrinsic Angular Momentum

Commutation rules

$$[\hat{S}_i, \hat{S}_j] = i\hbar\epsilon_{ijk}\hat{S}_k \quad [\hat{S}_i, \hat{S}^2] = 0$$

Spectrum

$$\hat{S}^2|sm_s\rangle = \hbar^2 s(s+1)|sm_s\rangle$$

$$\hat{S}_z|sm_s\rangle = \hbar m_s|sm_s\rangle \quad -s \leq m_s \leq s$$

$$\text{Degeneracy} \quad d_s = 2s + 1$$

QM — Theoretical Review

$$|\psi\rangle = |lm_l\rangle \otimes |sm_s\rangle \quad \leftrightarrow \quad \{\hat{\mathbf{L}}^2, \hat{L}_z, \hat{\mathbf{S}}^2, \hat{S}_z\}$$

QM — Theoretical Review

$$|\psi\rangle = |lm_l\rangle \otimes |sm_s\rangle \quad \leftrightarrow \quad \{\hat{\mathbf{L}}^2, \hat{L}_z, \hat{\mathbf{S}}^2, \hat{S}_z\}$$

$$\hat{\mu} = \hat{\mu}_l + \hat{\mu}_s = -\frac{\mu_B}{\hbar} (g_l \hat{\mathbf{L}} + g_s \hat{\mathbf{S}})$$

$$|\psi\rangle = |lm_l\rangle \otimes |sm_s\rangle \quad \leftrightarrow \quad \{\hat{\mathbf{L}}^2, \hat{L}_z, \hat{\mathbf{S}}^2, \hat{S}_z\}$$

$$\hat{\mu} = \hat{\mu}_l + \hat{\mu}_s = -\frac{\mu_B}{\hbar} (g_l \hat{\mathbf{L}} + g_s \hat{\mathbf{S}})$$

$$E_{m_l m_s} = \frac{\mu_B B_z}{\hbar} (g_l \langle \hat{L}_z \rangle_\psi + g_s \langle \hat{S}_z \rangle_\psi)$$

$$|\psi\rangle = |lm_l\rangle \otimes |sm_s\rangle \quad \leftrightarrow \quad \{\hat{\mathbf{L}}^2, \hat{L}_z, \hat{\mathbf{S}}^2, \hat{S}_z\}$$

$$\hat{\mu} = \hat{\mu}_l + \hat{\mu}_s = -\frac{\mu_B}{\hbar} (g_l \hat{\mathbf{L}} + g_s \hat{\mathbf{S}})$$

$$\begin{aligned} E_{m_l m_s} &= \frac{\mu_B B_z}{\hbar} (g_l \langle \hat{L}_z \rangle_\psi + g_s \langle \hat{S}_z \rangle_\psi) \\ &= \mu_B B_z (g_l m_l + g_s m_s) \end{aligned}$$

$$|\psi\rangle = |lm_l\rangle \otimes |sm_s\rangle \quad \leftrightarrow \quad \{\hat{\mathbf{L}}^2, \hat{L}_z, \hat{\mathbf{S}}^2, \hat{S}_z\}$$


$$\hat{\mu} = \hat{\mu}_l + \hat{\mu}_s = -\frac{\mu_B}{\hbar} (g_l \hat{\mathbf{L}} + g_s \hat{\mathbf{S}})$$

$$\begin{aligned} E_{m_l m_s} &= \frac{\mu_B B_z}{\hbar} (g_l \langle \hat{L}_z \rangle_\psi + g_s \langle \hat{S}_z \rangle_\psi) \\ &= \mu_B B_z (g_l m_l + g_s m_s) \end{aligned}$$

$$d = (2l + 1)(2s + 1)$$


QM — Experimental Outcome $l = 0$

The energy degeneracy is $d = d_s = 2$



observed

QM — Experimental Outcome $l = 0$




observed

The energy degeneracy is $d = d_s = 2$

$$s = \frac{1}{2} \Rightarrow m_s = \pm \frac{1}{2}$$

we are measuring the spin!

QM — Experimental Outcome $l = 0$



observed

The energy degeneracy is $d = d_s = 2$

$$s = \frac{1}{2} \Rightarrow m_s = \pm \frac{1}{2}$$

we are measuring the spin!

Measuring the spacing

$$\Delta E_{0, \pm \frac{1}{2}} = g_s (\mu_B B_z)$$

We also conclude that $g_s = 2$

Conclusion

In comparison with L , spin can assume both **integer** and **semi-integer** eigenvalues. Particles with integer and half-integer spin are called respectively **bosons** and **fermions**. This distinction governs their quantum behavior.

Thank you for the attention!

Total Angular Momentum

Consider a state $|\psi\rangle = |lm_l\rangle \otimes |sm_s\rangle$

Definition

$$\hat{\mathbf{J}} = \hat{\mathbf{L}} + \hat{\mathbf{S}}$$

$$\hat{\mathbf{J}}^2 = \hat{\mathbf{L}}^2 + \hat{\mathbf{S}}^2 + 2\hat{\mathbf{S}} \cdot \hat{\mathbf{L}}$$

Commutation rules

$$[\hat{L}_i, \hat{S}_j] = 0 \quad [\hat{J}_i, \hat{\mathbf{J}}^2] = 0$$

Spectrum

$$\hat{\mathbf{J}}^2 |jm_j\rangle = \hbar^2 j(j+1) |jm_j\rangle \quad |l-s| \leq j \leq l+s$$

$$\hat{J}_z |jm_j\rangle = \hbar m_j |jm_j\rangle \quad -j \leq m_j \leq j$$

Degeneracy

$$d_j = 2j + 1 = (2l + 1)(2s + 1)$$

Coupling of Angular Momenta

Consider the magnetic field experienced by an e^- orbiting around a nucleus with $Z p^+$

$$\mathbf{B}(\mathbf{r}) = \frac{Zq_e}{4\pi\epsilon_0 c^2 m_e} \frac{\hat{\mathbf{L}}}{|\mathbf{r}|^3}$$

Spin-Orbit coupling $\hat{H}_{\text{s-o}} = -\frac{1}{2}\hat{\boldsymbol{\mu}}_s \cdot \mathbf{B}$

$$\begin{aligned}\hat{H}_{\text{s-o}} &\propto \frac{1}{r^3} \hat{\mathbf{S}} \cdot \hat{\mathbf{L}} \\ &\propto \frac{1}{r^3} (\hat{\mathbf{J}}^2 - \hat{\mathbf{L}}^2 - \hat{\mathbf{S}}^2)\end{aligned}$$

We can represent the e^- state using the basis

$$\{\hat{\mathbf{L}}^2, \hat{L}_z, \hat{\mathbf{S}}^2, \hat{S}_z\}$$

Another option is the **coupled** basis

$$\{\hat{\mathbf{J}}^2, \hat{J}_z, \hat{\mathbf{L}}^2, \hat{\mathbf{S}}^2\}$$

since

$$[\hat{J}_z, \hat{\mathbf{L}}^2] = [\hat{J}_z, \hat{\mathbf{S}}^2] = 0$$

$$[\hat{\mathbf{J}}^2, \hat{\mathbf{L}}^2] = [\hat{\mathbf{J}}^2, \hat{\mathbf{S}}^2] = 0$$

Energy Spectra in Magnetic Field

$$\hat{H} = \hat{H}_0 + \hat{H}_{\text{s-o}} + \frac{\mu_B B_z}{\hbar} (\hat{L}_z + 2\hat{S}_z)$$

Energy Spectra in Magnetic Field

Paschen-Back Effect $\mu_B B \gg \xi$

$$\begin{aligned}\xi_{m_l m_s} &= \frac{\mu_B B_z}{\hbar} \left(\langle \hat{L}_z \rangle_\psi + 2 \langle \hat{S}_z \rangle_\psi \right) \\ &= \frac{\mu_B B_z}{\hbar} (m_l + 2m_s)\end{aligned}$$

Features

- Spin-orbit coupling negligible
- L and S precess independently around B
- Good quantum numbers: m_l, m_s
- Linear splitting in B

Zeeman Effect $\mu_B B \ll \xi$

Anomalous Zeeman Effect

- Spin-orbit coupling dominant
- L and S coupled to J
- Good quantum numbers: j, m_j
- Energy shift: $\Delta E = \mu_B B g_j m_j$

Landé g-factor

$$g_j = 1 + \frac{j(j+1) + s(s+1) - l(l+1)}{2j(j+1)}$$

Stern-Gerlach Experiment

Boson & Fermions

Hello world