



Analyzing Gravitational Waves through Numerical Simulations

Lorenzo Speri

Department of Physics
University of Trento
17th July 2018



Contents

1. Introduction
2. Einstein's Field Equation
3. Effects of Gravitational Waves
4. Production of Gravitational Waves
5. Numerical Evolution of Compact Binaries
6. Conclusions



Why Gravitational Waves?

The speed of light is the speed of causality.

Gravity must be causal.

A change in a gravitating source is communicated to a distant observer not faster than light.

Communication of a change in spacetime → Gravitational Waves



Einstein's Field Equation

Einstein's Field Equation is a rather complicated tensor equation

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = \frac{8\pi G}{c^4}T_{\mu\nu}$$

where

$$\Gamma_{\beta\gamma}^\alpha = \frac{1}{2}g^{\alpha\rho}(\partial_\beta g_{\gamma\rho} + \partial_\gamma g_{\rho\beta} - \partial_\rho g_{\beta\gamma})$$

$$R^\alpha_{\beta\gamma\sigma} = \Gamma^\alpha_{\gamma\lambda}\Gamma^\lambda_{\sigma\beta} - \Gamma^\alpha_{\sigma\lambda}\Gamma^\lambda_{\gamma\beta} + \partial_\gamma\Gamma^\alpha_{\sigma\beta} - \partial_\sigma\Gamma^\alpha_{\gamma\beta}$$

$$R_{\mu\nu} = R^\alpha_{\mu\alpha\nu} \quad R = \eta^{\mu\nu}R_{\mu\nu} = R^\mu_{\mu}$$



Linearized Einstein's Field Equation

- Weak field linearized approximation

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu} \quad |h_{\mu\nu}| \ll 1$$

- Trace-reversed perturbation metric

$$\bar{h}_{\mu\nu} = h_{\mu\nu} - \frac{1}{2}h\eta_{\mu\nu} \quad \bar{h} = \bar{h}^{\mu\nu}\eta_{\mu\nu} = -h$$

- Lorenz gauge $\partial_\mu h^{\mu\nu} = 0$

leads to the linearized Einstein's field equation

$$\left(\frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \nabla^2 \right) \bar{h}_{\mu\nu} = \frac{16\pi G}{c^4} T_{\mu\nu}$$



Linearized Einstein's Field Equation $T_{\mu\nu} = 0$

In vacuum $T_{\mu\nu} = 0$

$$\left(\frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \nabla^2 \right) \bar{h}_{\mu\nu} = 0$$

Plane wave solution

$$\bar{h}_{\mu\nu} = A_{\mu\nu} \cos [i\omega(t - z/c)]$$



Radiative degrees of freedom

The conditions imposed by the Transverse Traceless gauge

$$h_{0\nu}^{\text{TT}} = 0 \quad \eta^{\mu\nu} h_{\mu\nu}^{\text{TT}} = 0 \quad \partial^\mu h_{\mu\nu}^{\text{TT}} = 0$$

allow us to obtain the only two radiative degrees of freedom

$$\bar{h}_{\mu\nu}^{\text{TT}} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & h_+ & h_\times & 0 \\ 0 & h_\times & -h_+ & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

where

$$h_+ = A_{11} \cos(\omega(t - z/c))$$

$$h_\times = A_{12} \cos(\omega(t - z/c))$$



Geodesic Deviation Equation

For slowly-moving free-falling particles the geodesic deviation equations become

$$\frac{\partial^2}{\partial t^2} S^1 = \frac{1}{2} S^1 \frac{\partial^2}{\partial t^2} h_+ + \frac{1}{2} S^2 \frac{\partial^2}{\partial t^2} h_\times$$

$$\frac{\partial^2}{\partial t^2} S^2 = \frac{1}{2} S^1 \frac{\partial^2}{\partial t^2} h_\times - \frac{1}{2} S^2 \frac{\partial^2}{\partial t^2} h_+$$

first order solution

$$S^1 = S^1(t=0) \left(1 + \frac{1}{2} h_+ \right) + \frac{1}{2} h_\times S^2(t=0)$$

$$S^2 = S^2(t=0) \left(1 - \frac{1}{2} h_+ \right) + \frac{1}{2} h_\times S^1(t=0)$$



+ polarization $h_x = 0$



× polarization $h_+ = 0$



Left-handed polarization



Right-handed polarization



Linearized Einstein's Field Equation $T_{\mu\nu} \neq 0$

$$\left(\frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \nabla^2 \right) \bar{h}_{\mu\nu} = \frac{16\pi G}{c^4} T_{\mu\nu}$$

- far field approximation
- slowly moving source
- isolated system



Quadrupole Formula

A solution of the Linearized Einstein's Field equation is the Quadrupole Formula

$$h_{ij}^{\text{TT}} = \frac{2G}{c^4 r} \frac{d^2 \mathcal{I}_{kl}(t_r)}{dt^2} \left(P_i^k P_j^l - \frac{1}{2} P_{ij} P^{kl} \right)$$

where

$$\mathcal{I}_{kj} = \int \rho(\mathbf{y}) \left(y_k y_j - \frac{1}{3} \delta_{kj} y^l y_l \right) d^3 y$$

and

$$P_{ij} = \delta_{ij} - n_i n_j$$



slowly-moving equal-mass binary system

For an equal-mass binary system rotating at angular frequency ω and orbital radius R , the quadrupole formula becomes

$$h_{ij}^{\text{TT}} = \frac{8 GM R^2 \omega^2}{c^4 r} \begin{bmatrix} -\cos 2\omega t_r & -\sin 2\omega t_r & 0 \\ -\sin 2\omega t_r & \cos 2\omega t_r & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

estimate of the amplitude

$$\frac{8 GM R^2 \omega^2}{c^4 r} = 1.6 \times 10^{-22}$$

setting $M = M_\odot = 2 \times 10^{30}$ kg, $\omega = \sqrt{GM/(4R^3)}$,
 $R = 3r_s = 8.86$ km and $r = 100$ Mpc.



slowly-moving equal-mass binary system



Numerical Evolution of Compact Binaries

Quasi-equilibrium initial conditions of compact objects

- **Binary Black Holes (BBH)**
- **Binary Neutron Stars (BNS)**

Gravitational Radiation \Leftrightarrow Dynamical Evolution



BBH-b3




BBH-b4




BBH-b5



BBH-b6, BBH-b7, BBH-b10



BNS



Rest Mass Density Evolution



Conclusions

*"Gravitational waves will show us details of the bulk motion
of dense concentrations of energy"*

Kip S. Thorne

