

Class09

Loretta Cheng

##PDB Statistics

Today we will be using the Data Export Summary from PDB to figure out the following questions

```
db <- read.csv("Data Export Summary.csv")
```

Q1. What percentage of structures in the PDB are solved by X-Ray and Electron Microscopy.

```
xray.total <- sum(as.numeric(gsub(",", "", db$X.ray)))  
em.total <- sum(as.numeric(gsub(",", "", db$EM)))
```

```
# use `x` has input  
sum.comma <- function(x) {  
  #substitute comma and convert to numeric  
  sum(as.numeric(gsub(",", "", x)))  
}
```

For x-ray:

```
sum.comma(db$X.ray) / sum.comma(db$Total)
```

```
[1] 0.8553721
```

For EM:

```
round(sum.comma(db$EM) / sum.comma(db$Total), 2)
```

```
[1] 0.07
```

Q2. What proportion of structures in the PDB are protein?

```
round(sum.comma(db$Total[1])/sum.comma(db$Total),2)
```

[1] 0.87

Q3. Type HIV in the PDB website search box on the home page and determine how many HIV-1 protease structures are in the current PDB?




Figure 1: HIV-PR Structure from MERK with a bound drug

##Visualizing the HIV-1 protease structure

Q4: Water molecules normally have 3 atoms. Why do we see just one atom per water molecule in this structure?

The structure is too low to see a resolution of H atoms. We need a sub 1 Angstrom resolution to see Hydrogen.

Q5: There is a critical “conserved” water molecule in the binding site. Can you identify this water molecule? What residue number does this water molecule have?

HOH308

Working with Structures in R

We can use the `bio3d` package to read and perform bioinformatics about calculations on PDB structures.

```
library(bio3d)

pdb <- read.pdb("1hsg")
```

Note: Accessing on-line PDB file

```
pdb
```

```
Call: read.pdb(file = "1hsg")
```

```
Total Models#: 1
Total Atoms#: 1686, XYZs#: 5058 Chains#: 2 (values: A B)

Protein Atoms#: 1514 (residues/Calpha atoms#: 198)
Nucleic acid Atoms#: 0 (residues/phosphate atoms#: 0)

Non-protein/nucleic Atoms#: 172 (residues: 128)
Non-protein/nucleic resid values: [ HOH (127), MK1 (1) ]
```

```
Protein sequence:
```

```
PQITLWQRPLVTIKIGGQLKEALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYD
QILIEICGHKAIGTVLVGPTPVNIIGRNLLTQIGCTLNFPQITLWQRPLVTIKIGGQLKE
ALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYDQILIEICGHKAIGTVLVGPTP
VNIIGRNLLTQIGCTLNF
```

```
+ attr: atom, xyz, seqres, helix, sheet,
      calpha, remark, call
```

```
attributes(pdb)
```

```
$names
```

```
[1] "atom" "xyz" "seqres" "helix" "sheet" "calpha" "remark" "call"
```

```
$class
```

```
[1] "pdb" "sse"
```

```
head(pdb$atom)
```

	type	eleno	elety	alt	resid	chain	resno	insert	x	y	z	o	b
1	ATOM	1	N	<NA>	PRO	A	1	<NA>	29.361	39.686	5.862	1	38.10
2	ATOM	2	CA	<NA>	PRO	A	1	<NA>	30.307	38.663	5.319	1	40.62
3	ATOM	3	C	<NA>	PRO	A	1	<NA>	29.760	38.071	4.022	1	42.64
4	ATOM	4	O	<NA>	PRO	A	1	<NA>	28.600	38.302	3.676	1	43.40
5	ATOM	5	CB	<NA>	PRO	A	1	<NA>	30.508	37.541	6.342	1	37.87
6	ATOM	6	CG	<NA>	PRO	A	1	<NA>	29.296	37.591	7.162	1	38.40

	segid	elesy	charge
1	<NA>	N	<NA>
2	<NA>	C	<NA>
3	<NA>	C	<NA>
4	<NA>	O	<NA>
5	<NA>	C	<NA>
6	<NA>	C	<NA>

Read an ADK structure

```
adk <- read.pdb("6s36")
```

Note: Accessing on-line PDB file

PDB has ALT records, taking A only, rm.alt=TRUE

```
adk
```

Call: read.pdb(file = "6s36")

Total Models#: 1

Total Atoms#: 1898, XYZs#: 5694 Chains#: 1 (values: A)

Protein Atoms#: 1654 (residues/Calpha atoms#: 214)

Nucleic acid Atoms#: 0 (residues/phosphate atoms#: 0)

Non-protein/nucleic Atoms#: 244 (residues: 244)

Non-protein/nucleic resid values: [CL (3), HOH (238), MG (2), NA (1)]

Protein sequence:

```
MRILLGAPGAGKGTQAQFIMEKYGIPQISTGDMRLRAAVKSGSELGKQAKDIMDAGKLV  
DELVIALVKERIAQEDCRNGFLDGFPRTPQADAMKEAGINVDYVLEFDVPDELIVDKI  
VGRRVHAPSGRVYHVKFNPPKVEGKDDVTGEELTTRKDDQEETVRKRLVEYHQMTAPLIG  
YYSKEAEAGNTKYAKVDGTPVAEVRADLEKILG
```


+ attr: atom, xyz, seqres, helix, sheet,
calpha, remark, call

Perform a prediction of flexibility with a technique called NMA(normal mode analysis)

```
# predict flexibility  
m <- nma(adk)
```

```
Building Hessian...      Done in 0.013 seconds.  
Diagonalizing Hessian... Done in 0.263 seconds.
```

```
plot(m)
```



Write out a “movie” (a.k.a trajectory) of the motion for viewing in Molstar

```
mktrj(m, file="adk_m7.pdb")
```

Q7: How many amino acid residues are there in this pdb object?

198

Q8: Name one of the two non-protein residues?

MK1

Q9: How many protein chains are in this structure?

2