

Sex-biased drug-sensitivity hits by class and Egr1 dependence

Drug class	WT-only (n = 27) lost after Egr1 KD	Shared (n = 22) Egr1-independent	KD-only (n = 3) gained after KD
Mevalonate-sterol & glucocorticoid	Lovastatin Simvastatin Dexamethasone Methylprednisolone Flumethasone Fluocinolone	Atorvastatin	–
Estrogen receptor (SERM)	Tamoxifen	Clomiphene Bazedoxifene	Toremifene
Other steroid / adrenergic	Allylestrenol	–	Prasterone (DHEA) Guanabenz
Antimetabolite / nucleoside	Thioguanine (run 2)	Fluorouracil (5-FU) Carmofur Floxuridine Azacitidine Thioguanine (run 1) Mycophenolic acid	–
DNA-damaging (crosslink / intercalate)	Amsacrine Aminacrine	Oxaliplatin Mitomycin	–
Microtubule	Vinblastine	Docetaxel	–
Cationic-amphiphilic neuroactive	Pimozide Fluphenazine Metitepine Suloctidil	Ethopropazine	–
Oxidative / thiol-reactive	Phenylmercuric acetate Thimerosal Bronopol Menadione	Artesunate Furazolidone	–
Mito / ionophore / antiparasitic	Atovaquone Monensin Pyrvinium	–	–
Antimicrobial (azole / macrolide / quinoline)	Benzoxiquine Oxyquinoline Dirithromycin	Oxiconazole Itraconazole	–
Anti-inflammatory (NSAID / COX)	Oxyphenbutazone	Celecoxib	–
Other (protein synth / CNS / saponin)	–	Cycloheximide Ketanserin Riluzole Escin	–

Compartments = $q \leq 0.05$ Male-vs-Female sensitivity hits (Stouffer combined Z, BH-FDR). All WT & shared hits are male-biased; the 2 female-biased hits (Prasterone, Guanabenz) are KD-only.

Thioguanine was screened twice: run 1 is shared, run 2 is WT-only.