

Sex-biased drug-sensitivity hits by class and Egr1 dependence

Drug class	WT-only (n = 27) lost after Egr1 KD	Shared (n = 22) Egr1-independent	KD-only (n = 3) gained after KD
Sterol / steroid	Lovastatin Allylestrenol Flumethazone Tamoxifen Simvastatin Dexamethasone Fluocinolone Methylprednisolone	Atorvastatin Bazedoxifene Clomiphene	Toremifene Prasterone (DHEA)
Oxidative / anti-inflammatory	Menadione Oxyphenbutazone Phenylmercuric acetate Thimerosal Bronopol	Celecoxib Cycloheximide Escin	—
Antimicrobial / antiparasitic	Atovaquone Benzoxiquine Pyrvinium Dirithromycin Oxyquinoline Monensin	Oxiconazole Itraconazole Furazolidone Artesunate	—
Neuroactive	Metitepine Pimozide Suloctidil Fluphenazine	Ethopropazine Ketanserin Riluzole	Guanabenz
Antiproliferative	Thioguanine (run 2) Amsacrine Aminacrine Vinblastine	Azacitidine Carmofur Oxaliplatin Floxuridine Fluorouracil (5-FU) Thioguanine (run 1) Mitomycin Mycophenolic Acid Docetaxel	—

Compartments = $q \leq 0.05$ Male-vs-Female sensitivity hits (BH-FDR over 85 runs), grouped into the 5 panel-D classes.

All hits are male-biased except the 2 KD-only female-biased hits (Guanabenz, Prasterone). Thioguanine was screened twice.