

Coalgebra, Categories, Automata, and MDPs

lorisj

May 2025

Contents

1	Introduction	3
2	Notation	4
3	Category Theory	5
3.1	Categories	5
3.2	Basic Categorical Structures	7
3.2.1	Span and Cospan	7
3.2.2	Monoid	7
3.2.3	Isomorphism and Isomorphic objects	8
3.2.4	Initial and Terminal/Final Objects	8
3.2.5	Product and Coproduct	9
3.3	Duality	10
3.4	Functors	11
3.5	Coalgebra	12
4	Slice and product category	15
5	Final Coalgebras, Behavioral Equivalence, Bisimulation	16
5.1	Final Coalgebras	16
5.2	Behavioral Equivalence and Bisimulation	16
6	Couplings (new)	18
6.0.1	New definitions of couplings (new)	18
7	Coalgebra of MDPs (new)	21
7.1	Bisimulation	21
8	LDBA/LDBW Coalgebraically (new)	26
8.1	Automata review	26
9	Coalgebraic Behavioral Metrics	31
9.1	Metric review and notation	31
9.2	Categorical Prerequisites	31
9.3	Evaluations	33
9.4	Kantorovic Metric	33
10	More complicated results	36
10.1	Limits/Colimits, Pullbacks	36
10.2	Comparing Bisimulation	36
10.3	Existence of final coalgebra	37

1 Introduction

The main goal of these notes is to explore new ideas in the intersection of:

- Category theory/coalgebra
- MDPs (Markov Decision Processes)
- LDBA (Limit Deterministic Buchi Automata),
- Bisimulation and bisimulation metrics

and also serve as a reference for some results/definitions in these areas.

2 Notation

Definition: Notation Used


These are default notations that are used unless otherwise noted:

- $\text{TFDC} \stackrel{\text{def}}{=} \text{The following diagram(s) commute(s).}$
- $\text{s.t.} \stackrel{\text{def}}{=} \text{such that}$
- $A^B \stackrel{\text{def}}{=} \{f : B \rightarrow A\}$ for sets A, B .
- $2^A = \mathcal{P}[A]$ (powerset of A) for set A , note that this means the powerset can be interpreted as the set of functions from $A \rightarrow 2$.
- $\omega \stackrel{\text{def}}{=} \{\text{natural numbers including } 0\} = \{0, 1, 2, \dots\}$.
- $\forall_v^V, \exists_v^V \stackrel{\text{def}}{=} \forall_{v \in V}, \exists_{v \in V}$, where the superscript refers to the set being quantified over (domain) and the subscript denotes the variable bound by the quantifier.
- $f[X], f^{-1}[X] \stackrel{\text{def}}{=} \text{image, preimage for a set } X \text{ under function } f \text{ (in contrast } f(x), f^{-1}(x) \text{ denote element application)}$
- $(\cdot)$ denotes a unique free variable (i.e. $(\cdot) + 2$ is the function that takes in a variable and adds 2)
- ΔX DEF distribution-functor

3 Category Theory

3.1 Categories

Intuitively, a category can be thought of as a more structured graph, where you can compose edges. Specifically, a category is a directed multigraph, and also there is a composition operation, as well as an identity. The nodes are called Objects, and the edges are called arrows (or morphisms). Essentially a category will look something like this:



The important notes are that:

1. You can have multiple arrows between two nodes that are not equal
2. If I have two arrows: $f : A \rightarrow B, g : B \rightarrow C$, then there must exist some arrow $g \circ f : A \rightarrow C$.
3. There exists at least one self loop for each object, called the identity, $\text{id}_C : C \rightarrow C$ such that its composition with any other object does not change it.

More formally:

Definition: Category

category

A category C consists of:

- **Obj**(C): A class of objects
- **Hom**(C): A class of morphisms (sometimes called arrows)
- $\circ$: A class function : **Hom**(C) $\times$ **Hom**(C) $\rightarrow$ **Hom**(C)

Such that:

1. Each morphism has a *domain* and *codomain*:

For every $f \in \mathbf{Hom}(C)$ there must be two objects, a *domain* and *codomain*, both in **Obj**(C).

As notation, $f : X \rightarrow Y$ denotes a morphism with domain X , codomain Y , and **Hom**(X, Y) for objects X, Y denotes the class of all morphisms with domain X , codomain Y .

2. Composition is *type-respecting*:

If $f : X \rightarrow Y$ and $g : Y \rightarrow Z$, then the composite $g \circ f$ must exist in **Hom** C , and must be of type: $g \circ f : X \rightarrow Z$.

3. Composition is *associative*:

For morphisms $f : W \rightarrow X$, $g : X \rightarrow Y$, and $h : Y \rightarrow Z$:

$$(h \circ g) \circ f = h \circ (g \circ f) \quad (1)$$

4. *Identity morphisms* exist:

Every $X \in \mathbf{Obj}(C)$ has a *identity* (or *unit*) morphism $\text{id}_X : X \rightarrow X$ such that for every morphism $f : X \rightarrow Y$ and $g : Z \rightarrow X$:

$$\text{id}_Y \circ f = f \quad (2)$$

$$f \circ \text{id}_X = f \quad (3)$$

This definition is very abstract, so the following examples should make the idea more concrete.

Example: Common categories

Here are some common categories: (See DEF category)

Set, the category of sets :

catset

- **Obj**: All sets.
- **Hom**: All functions between sets.
- $\circ$: Normal function composition.
- id_X : The identity function on a set X .

Rel, the category of relations :

catrel

- **Obj**: All sets.
- **Hom**: A morphism $R : X \rightarrow Y$ is a binary relation $R \subseteq X \times Y$.
- $\circ$: For $R : X \rightarrow Y$ and $S : Y \rightarrow Z$, their composite $S \circ R : X \rightarrow Z$ is given by:
$$(x, z) \in (S \circ R) \quad \text{iff} \quad \exists y \in Y \text{ such that } (x, y) \in R \text{ and } (y, z) \in S.$$
- id_X : For each set X , $\text{id}_X = \{(x, x) : x \in X\}$, i.e. the diagonal relation.

Typ, the category of types :

cattypes

- **Obj** : Types in a given type theory or programming language.
(think `string`, `int`, `double`, `string[]`)
- **Hom** : Pure functions (i.e. programs with no side effects) from the input type to the output type (think `length : string → number`). Note only pure functions are allowed, so something that has side effects would not work.
- $\circ$: Composition of functions.
- id_X : Identity function on type X .

Grp, the category of groups :

catgroup

- **Obj**: All groups.
- **Hom**: Group homomorphisms. Formally, a morphism between groups $f : (G, \cdot^G) \rightarrow (H, \cdot^H)$ respects the group operation: $f(g_1 \cdot^G g_2) = f(g_1) \cdot^H f(g_2)$.
- $\circ$: Composition is the usual function composition of homomorphisms.
- $\text{id}_{(G, \cdot)}$: The identity on a group $(G, \cdot)$ is the identity homomorphism id_G which sends each $g \in G$ to itself.

The following example shows that morphisms on a category need not reduce to relations on the class of objects:

Category of Dimensions of Linear maps :

- **Obj**: The natural numbers ω .
- **Hom**: A morphism $R : n \rightarrow m$ is a $n \times m$ matrix $\in \mathbb{R}^{n \times m}$.
- $\circ$: matrix multiplication.
- id_n : For $n \in \omega$, id_n is simply the $n \times n$ identity matrix.

3.2 Basic Categorical Structures

Here are some basic categorical structures that we will use, and these should give some intuition as to what a categorical definition looks like.

3.2.1 Span and Cospan

Maybe the easiest objects to define.

Definition: Span and cospan

span-cospan

For a category C :

Span :

span

A span consists of a diagram of the form:

$$A \xleftarrow{f} B \xrightarrow{g} C$$

Cospan :


cospan

A cospan consists of a diagram of the form:

$$A \xrightarrow{f} B \xleftarrow{g} C$$

3.2.2 Monoid

If we look at a category, sometimes we can see an isolated object, i.e. something that looks like this:



where there are no arrows going from A outside of A . This is called a monoid. More formally:

Definition: Monoid

monoid

A monoid in a category C is an object M such that every morphism f with domain M must also have codomain M .

Recall the traditional definition of a monoid consists of:

Set theoretic monoid: A monoid $(M, +, e)$ is a set M , function $+: M \times M \rightarrow M$, element $e \in M$ s.t.:

1. $+$ is associative
2. For all $m \in M$: $e + m = m + e = m$

Category Theory Monoid is Set Monoid: Let $*$ be a monoid. Then $(\mathbf{Hom}(*, *), \circ)$ is a set theoretic monoid. (recall $\circ$ is the composition operator of C)

Proof. Associativity and the existence of identity follow immediately from the category theory axioms. $\square$

3.2.3 Isomorphism and Isomorphic objects

An isomorphism generalizes the idea of a bijection. Namely:

Definition: Isomorphism

isomorphism

For a category C , an isomorphism is any morphism $f: X \rightarrow Y$ such that there exists another morphism $f^{-1}: Y \rightarrow X$ such that $f^{-1} \circ f = \text{id}_X$, and $f \circ f^{-1} = \text{id}_Y$. As a diagram:

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \xleftarrow{f^{-1}} & \end{array}$$

We can transfer this definition from morphisms to objects:

Definition: Isomorphic objects

isomorphic

In a category C , objects $X, Y \in C$ are isomorphic (denoted $X \cong Y$) if there exists some isomorphism $f: X \rightarrow Y$.

Isomorphisms/Isomorphic objects:

- In **Set**, isomorphisms are exactly bijective functions, and two sets are isomorphic iff they have the same cardinality.
- In **Grp**, isomorphisms are group homomorphisms that are invertible, i.e. group isomorphisms.

Many categorical definitions are only unique up to isomorphism, meaning that any two objects that satisfy the condition must be isomorphic. For example see [terminal-unique](#).

3.2.4 Initial and Terminal/Final Objects

Definition: Initial Object

initial-obj

An object $*$ in $\mathbf{Obj}(C)$ for some category C is called initial if for any other object $X \in \mathbf{Obj}(C)$, there is a unique morphism $! : * \rightarrow X$.

Definition: Terminal/Final Object

terminal-obj

An object $*$ in $\mathbf{Obj}(C)$ for some category C is called terminal (aka final) if for any other object $X \in \mathbf{Obj}(C)$, there is a unique morphism $! : X \rightarrow *$.

Example: Initial/Terminal Objects

Initial for **Set** : Note that if we look at **Set**, the set with a single element (usually denoted $1 = \{0\} = \{\emptyset\}$) is terminal. (For any other set, there is a unique map that just sends everything to 0)

Terminal for **Set** : By convention, we assume that there is a unique function from any set to the emptyset. This means that the empty set (aka 0) is the terminal object in **Set**.

Terminal for **Grp** : Note that the trivial group $(\{0\}, +, 0)$ is terminal in **Grp**. Namely for any other group $(G, +, 0)$ there is only one homomorphism to the trivial group, namely the one that sends everything to the identity.

Result: Terminal Objects are Unique up to Isomorphism

terminal-unique

Let $*_0, *_1$ both be terminal objects of a category C . Then $*_0 \cong *_1$.

Proof. Because $*_0$ is terminal, there must be some (unique) $!_0 : *_1 \rightarrow *_0$. Likewise because $*_1$ is terminal, there must be some (unique) $!_1 : *_0 \rightarrow *_1$. So notice that $!_0 \circ !_1 : *_0 \rightarrow *_0$, so by finality of $*_0$ the identity map is the unique morphism $!_0 \circ !_1 = \text{id}_{*_0}$. The same argument holds for $!_1 \circ !_0 = \text{id}_{*_1}$. Thus by **DEF isomorphism** $!_0$ is an isomorphism, and so $*_0 \cong *_1$. $\square$

Note that the same is true for initial objects. This means that we can say **the** [initial/terminal] object instead of **an/a** [initial/terminal] object.

3.2.5 Product and Coproduct

Definition: Product

product

Let $A, B \in \mathbf{Obj}(C)$ for some category C . A product consists of:

- An object $A \times B \in \mathbf{Obj}(C)$
- A morphism $\pi_A : A \times B \rightarrow A$
- A morphism $\pi_B : A \times B \rightarrow B$

s.t. for any other span between A and B : $A \xleftarrow{f} Q \xrightarrow{g} B$ (**DEF span**) we have that there exists a unique map $r_Q : Q \rightarrow A \times B$ s.t. TFDC:

$$\begin{array}{ccccc} & & A \times B & & \\ \pi_A \swarrow & & \uparrow \exists ! r_Q & & \searrow \pi_B \\ A & \xleftarrow{f} & Q & \xrightarrow{g} & B \end{array}$$

Intuitively, this means that if I know a product exists, and I have some span $A \xleftarrow{f} Q \xrightarrow{g} B$, we have that $f = \pi_A \circ r_Q$, and $g = \pi_B \circ r_Q$. This means that we can factor/reduce/simplify f, g by going through the product.

IMPORTANT NOTE: Not all categories have a product for every two objects, however many of the ones we will see do.

Products of some categories : The product for objects $A, B \in \mathbf{Obj}(C)$ in the categories C below is given by:

Set : The product of sets A, B consists of the cartesian product $\{(a, b) | a \in A, b \in B\}$, where π_A, π_B are the projection functions from $A \times B$ to A and B respectively. Note that given a span $A \xleftarrow{f} Q \xrightarrow{g} B$, the unique r_Q (see **DEF product**) is given by $r_Q = (q : Q) \mapsto (f(q), g(q))$.

Grp : The product of groups $(G_0, +_0, e_0), (G_1, +_1, e_1)$ is a group $(G_0 \times G_1, (+_0, +_1), (e_0, e_1))$, where $G_0 \times G_1$ is the cartesian product on sets, and $(+_0, +_1)$ acts elementwise.

Definition: Coproduct

coproduct

Let $A, B \in \mathbf{Obj}(C)$ for some category C . A coproduct consists of:

- An object $A + B \in \mathbf{Obj}(C)$
- A morphism $i_A : A \rightarrow A + B$
- A morphism $i_B : B \rightarrow A + B$

s.t. for any other cospan between A and B : $A \xrightarrow{f} Q \xleftarrow{g} B$ (**DEF cospan**) we have that there exists a unique map $r_Q : A + B \rightarrow Q$ s.t. TFDC:

$$\begin{array}{ccccc} & & A + B & & \\ & i_A \nearrow & & \nwarrow i_B & \\ A & \xrightarrow{f} & Q & \xleftarrow{g} & B \\ & & \downarrow \exists! r_Q & & \end{array}$$

Coproduct :

Set : The coproduct of sets A, B consists of the disjoint union $A \sqcup B$, where the $i_A : A \rightarrow A \sqcup B$ is the inclusion map of A , and $i_B : B \rightarrow A \sqcup B$ is the inclusion map of B . Note that for any cospan $A \xrightarrow{f} Q \xleftarrow{g} B$, the unique r_Q (see **DEF coproduct**) is given by $r_Q = (q : Q) \mapsto \begin{cases} f(q) & q \in A \\ g(q) & q \in B \end{cases}$

3.3 Duality

You may have noticed many of these definitions look the same, but are sort of opposite to each other. For instance the definition of a coproduct is the same as a product, but where all of the arrows are flipped. Likewise a cospan is a span but with all arrows are flipped. This notion is called duality, many times in category theory there is some definition that is of interest, and it turns out that usually, the dual definition (consisting of reversing all of the arrows) usually describes something interesting as well. When we have a name for the base thing, and we want a name for the dual, we just add the word co- in front. (coproduct, cospan, colimit, coalgebra, comonad, ...)

3.4 Functors

A functor is sort of like a homomorphism between categories. A functor should take an object to another object, and an arrow should take an arrow to another arrow. It should also preserve composition, as well as identity morphisms. More formally:

Definition: Functor

functor

Let C, D be two categories. A (covariant) *functor* $F : C \rightarrow D$ consists of:

- A class function on objects:
For each object $X \in \mathbf{Obj}(C)$, there must exist an object $FX \in \mathbf{Obj}(D)$
- A class function on morphisms:
For each morphism $f : X \rightarrow Y$ in $\mathbf{Hom}(C)$, there must exist a morphism $Ff : FX \rightarrow FY$.

These assignments must satisfy the following conditions:


- *Preservation of id*:

$$F(\text{id}_X^C) = \text{id}_{FX}^D \quad \text{for all } X \in \mathbf{Obj}(C) \quad (4)$$


- *Preservation of $\circ$* :

$$F(g \circ^C f) = F(g) \circ^D F(f) \quad (5)$$

Intuitively, a functor basically just adds F in front of everything. In particular, if you have any diagram that commutes:



then after lifting the diagram (applying F everywhere) it should still commute:



Definition: Endofunctor

endofunctor

An endofunctor is simply a functor that has the same codomain, namely an endofunctor on a category C is a functor $F : C \rightarrow C$.

Example: Functor Examples

On **Set** : , these are functors, which take a set X and a morphism $f : X \rightarrow Y$ to:

- $FX = X$, and $F(f) = f$. This is called the identity functor.
- $FX = X^2$, and $F(f) = (f, f) : X^2 \rightarrow Y^2$.
- $FX = X + A$ for some fixed set A , where a morphism $f : X \rightarrow Y$ maps to:

$$F(f) : X + A \rightarrow Y + A \quad (6)$$

$$F(f) : (u : X + A) \mapsto \begin{cases} f(u) & u \in X \\ u & u \in A \end{cases} \quad (7)$$

- $FX = 2^X$, and $F(f) = f[\cdot]$. (also called the covariant powerset functor)
- The distribution functor:

Definition: Distribution Functor

distribution-functor

The distribution functor on **Set**:

- On objects: maps a set X to the set

$$\Delta X = \{P : X \rightarrow [0, 1]_{\mathbb{R}} \mid \sum_{x \in X} P(x) = 1, P \text{ has finite support}\} \quad (8)$$

- On morphisms: maps a function $f : X \rightarrow Y$ to the function $\Delta f : \Delta X \rightarrow \Delta Y$, taking $P \mapsto (y : Y) \mapsto \sum_{x \in f^{-1}[\{y\}]} P(x)$.

Functors for programmers : If you come from a programming background, the easiest way of thinking about a functor is of a generic type. Recall **Typ** from **DEF cattytypes**. A functor on **Typ** consists of a generic type/way to lift types (a function $\text{Type} \rightarrow \text{Type}$), as well as a way to lift functions. The typical example is the list functor:

$$: X \mapsto X[]$$

On morphisms, given a function/program $f : X \rightarrow Y$, you get the function $\text{map}(f) : X[] \rightarrow Y[]$, which just applies f elementwise. For example take the function: $\text{len} : \text{string} \rightarrow \text{Nat}$. This function takes in a string, and returns its length as a natural number. The lifted function $\text{map}(\text{len})$ will take in a string array, and output each strings length:

$$\text{map}(\text{len}) : \text{string}[] \rightarrow \text{Nat}[] \quad (9)$$

$$: ["abc", "a", ""] \mapsto [3, 1, 0] \quad (10)$$

If you are familiar with generic types, then a functor is exactly a generic type (usually written as $T<X>$) as well as a way to lift maps to the generic type.

3.5 Coalgebra

As it turns out, many systems can be modeled in terms of category theory, using coalgebras (which we will define shortly). In fact the standard notion of bisimulation and trace equivalence can be modeled canonically using fairly simple definitions, and because of how general the language of category theory is, it allows us to come up with canonical notions of bisimulation for many different systems. For now we have to define what a system is.

Definition: Coalgebra**coalgebra**

A coalgebra for an endofunctor $F : C \rightarrow C$ is a morphism

$$c_X : X \rightarrow FX \quad (11)$$

Note that there are no additional constraints. Note that if we think of F adding structure to X , then a coalgebra can model many different forms of systems:

Coalgebras on **Set** : Consider **Set**. Here are some standard functors, along with the systems that they represent:

- $F(X) = X$
Deterministic system (each state transitions to exactly one next state).
- $F(X) = A \times X$
Labeled deterministic system (each state emits a label in A and transitions to a unique next state).
- $F(X) = 1 + X$
System that can either terminate (landing in the singleton 1) or transition to a next state.
- $F(X) = 1 + X + X^2$ (+ is disjoint union/coproduct of sets)
System where each state can either terminate, transition deterministically, or offer two next states, equivalently a potentially infinite tree.
- $F(X) = X^A$
System where each action $a \in A$ leads to a next state (an “action-indexed” deterministic system).
- $F(X) = 2^X$ (also called powerset, $\mathcal{P}A$)
System where each state has a set of next states, or equivalently a non deterministic transition system.
- $F(X) = \Delta X$ (where Δ is the distribution functor, and so ΔX is the set of distributions on X which acts on morphisms via the pushforward measure)
System with probabilistic transitions.
- $F(X) = \mathbb{R} \times (\Delta X)^A$
Markov Decision Process (MDP) with state valued reward. See here [DEF mdp-functor](#) for more.

We will also need this structure:

Definition: Coalgebra Homomorphism**coalgebra-hom**

Let $c_X : X \rightarrow FX$, $c_Y : Y \rightarrow FY$ be two coalgebras.

A coalgebra homomorphism is a function $\phi : X \rightarrow Y$ such that TFDC:

$$\begin{array}{ccc} X & \xrightarrow{\phi} & Y \\ c_X \downarrow & & \downarrow c_Y \\ FX & \xrightarrow{F\phi} & FY \end{array}$$

Result: Coalgebras form a category

For a given functor F , the set of F -coalgebras form a category. Namely this category has:

- Objects: F -coalgebras

- Morphisms: F -coalgebra morphisms
- Composition: Function composition
- id_X : Identity functions

Proof. The only things to check is that the composition of coalgebra homomorphisms is a coalgebra homomorphism, and that there are identities (which are just the identity functions). Assume you have 3 coalgebras c_X, c_Y, c_Z . Let $\phi : c_X \rightarrow c_Y$, and $\psi : c_Y \rightarrow c_Z$.

$$\begin{array}{ccccc} X & \xrightarrow{\phi} & Y & \xrightarrow{\psi} & Z \\ c_X \downarrow & & \downarrow c_Y & & \downarrow c_Z \\ FX & \xrightarrow{F\phi} & FY & \xrightarrow{F\psi} & FZ \end{array}$$

The diagram commutes because both ϕ and ψ are coalgebras.

Now because F is a functor, $F\phi \circ F\psi = F(\phi \circ \psi)$, and so you can make the homomorphism $: c_X \rightarrow c_Z$:

$$\begin{array}{ccc} X & \xrightarrow{\psi \circ \phi} & Z \\ c_X \downarrow & & \downarrow c_Z \\ FX & \xrightarrow{F(\psi \circ \phi)} & FZ \end{array}$$

The identity part can be shown as well by just assuming $\phi = \text{id}^X, Y = X$ or $\psi = \text{id}^Y, Z = Y$. Associativity can be shown as well because coalgebra homomorphism composition is the same as function composition. $\square$

4 Slice and product category

Later we will use the following constructions:

Definition: Slice Category

slice-cat

Let C be some category, and let $A \in \mathbf{Obj}(C)$. The slice category C/A is the category with:

- $\mathbf{Obj}(C/A)$: V_X , where $V_X : X \rightarrow A$ is a morphism in C .
- $\mathbf{Hom}(C/A)$: $f : V_X \rightarrow V_Y$ reduces to $f : X \rightarrow Y$ s.t. TFDC:

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \searrow V_X & \swarrow V_Y \\ & A & \end{array}$$

Set/2: $\mathbf{Set}/2$ is the category with:

- $\mathbf{Obj}(\mathbf{Set}/2) \stackrel{\text{def}}{=} \{X \mid X \in \mathbf{Obj}(\mathbf{Set})\}$
- $\mathbf{Hom}(\mathbf{Set}/2) \stackrel{\text{def}}{=} \{m_X : X \rightarrow 2 \mid X \in \mathbf{Obj}(\mathbf{Set})\}$
- $\text{id}_X, \circ$ just like in $\mathbf{Set}$.

Note that objects in this category are just binary partitions of $\mathbf{Obj}(\mathbf{Set})$, and morphisms are just functions that preserve the partition.

In particular if you have an object $m_X : X \rightarrow 2$, this induces a partition $X = m_X^{-1}[\{0\}] + m_X^{-1}[\{1\}]$.

Furthermore any morphism $f : m_X \rightarrow m_Y$ must preserve the partition, i.e. $m_Y(f(x)) = m_X(x)$ for all $x \in X$.

Definition: Product Category

product-cat

Let C and D be two categories. Define the product category $C \times D$ as the category with:

- $\mathbf{Obj}(C \times D) \stackrel{\text{def}}{=} \{(X, Y) \mid X \in \mathbf{Obj}(C), Y \in \mathbf{Obj}(D)\}$ (as a class, not necessarily a set)
- $\mathbf{Hom}(C \times D) = \{(f, g) \mid f \in \mathbf{Hom}(C), g \in \mathbf{Hom}(D)\}$

Set $\times$ Set: $\mathbf{Set} \times \mathbf{Set}$ is the category with:

- $\mathbf{Obj}(\mathbf{Set} \times \mathbf{Set}) \stackrel{\text{def}}{=} \text{all } (X, Y) \text{ where } X, Y \in \mathbf{Obj}(\mathbf{Set}).$
- $\mathbf{Hom}(\mathbf{Set} \times \mathbf{Set}) \stackrel{\text{def}}{=} \text{all } (f, g) \text{ where } f, g \in \mathbf{Hom}(\mathbf{Set}).$
- $\circ: (f, g) \circ (h, k) \stackrel{\text{def}}{=} (f \circ h, g \circ k)$
- $\text{id}_X: \text{id}_{(X, Y)} \stackrel{\text{def}}{=} (\text{id}_X, \text{id}_Y)$

5 Final Coalgebras, Behavioral Equivalence, Bisimulation

5.1 Final Coalgebras

One natural question to answer is given two systems, are they behaviorally equivalent. This question can be formalized by looking at terminal objects in the category of coalgebras.

Terminal/Final Coalgebra : Recall DEF terminal-obj.

Fix a functor F . The terminal/final coalgebra for F is the terminal object in the category of F -coalgebras.

Terminal Coalgebra : As it turns out, most categories have a notion of a terminal object. Here we give some examples:

$FX = A \times X$: The final coalgebra consists of $* : A^\omega \rightarrow A \times A^\omega$, which maps $(a_0, a_1, \dots) \mapsto (a_0, (a_1, \dots))$.

Any coalgebra $c : X \rightarrow A \times X$ has a unique coalgebra morphism $! : c \rightarrow *$, namely the map which sends x to the sequence $(a_0, a_1, \dots) : A^\omega$ that happens after applying c infinitely often:

$c(x) = (a_0, x_1), c(x_1) = (a_1, x_2), \dots$

$FX = 2 \times X^A$: The final coalgebra consists of 2^{A^*} , the set of languages over A .

Result: Lambek's theorem (dual)

If F has a final coalgebra $\alpha : A \rightarrow FA$, then $A \cong FA$.

Proof. By finality of α , there must be a unique coalgebra hom $f : F\alpha \rightarrow \alpha$. Thus TFDC:

$$\begin{array}{ccc} A & \xleftarrow{\exists! f} & FA \\ \alpha \downarrow & & \downarrow F\alpha \\ FA & \xleftarrow{Ff} & FFA \end{array}$$

Now note that α is also a coalgebra hom : $\alpha \rightarrow F\alpha$ (trivially TFDC):

$$\begin{array}{ccc} A & \xrightarrow{\alpha} & FA \\ \alpha \downarrow & & \downarrow F\alpha \\ FA & \xrightarrow{F\alpha} & FFA \end{array}$$

So there are two coalgebra homomorphisms : $\alpha \rightarrow \alpha$, namely id_A and $f \circ \alpha$. By finality of α , we have $f \circ \alpha = \text{id}_A$.

So because $f \circ \alpha = \text{id}_A$, it must be that $Ff \circ F\alpha = \text{id}_{FA}$, by functoriality of F .

Thus by the first diagram, we have $\text{id}_{FA} = Ff \circ F\alpha = \alpha \circ f$. □

Note that Lambek's Theorem is for F algebras, but the dual statement is what we proved above.

5.2 Behavioral Equivalence and Bisimulation

There are two notions of coalgebraic bisimulation that we cover:

Definition: Behavioral Equivalence Bisimilarity (Set)

beq-bisimulation

Define the bisimilarity relation between two F -coalgebras:

$$c_X : X \rightarrow FX$$

$$c_Y : Y \rightarrow FY$$

as a relation on $X \times Y$ given by:

$$x \sim y \iff [x]_{\text{final}(c_X)} = [y]_{\text{final}(c_Y)} \quad (12)$$

where $[\cdot]_{\text{final}(c_X)}$ and $[\cdot]_{\text{final}(c_Y)}$ are the unique homomorphisms to the final F -coalgebra from c_X and c_Y .

Behavioral Equivalence on a set is an equivalence relation: If $X = Y$ and $c_X = c_Y$, then this will be an equivalence relation.

Definition: Aczel-Mendler/Span Bisimulation

AM-bisimulation

An Aczel Mendler bisimulation (AM-bisimulation) is a DEF span in the category of F -coalgebras.

I.e. for two coalgebras $c_X : X \rightarrow FX$, and $c_Y : Y \rightarrow FY$, an AM bisimulation is a span:

$$c_X \xleftarrow{\phi_X} c_R \xrightarrow{\phi_Y} c_Y$$

Note that the two are related. In fact for most common functors (any functor that preserves weak pullbacks) AM Bisimulation is weaker than Behavioral equivalence.

The result is a bit more complicated, but can be found here

RES AM-vs-beh-equiv.

6 Couplings (new)

Set Element Coupling :

Let

- $F : \mathbf{Set} \rightarrow \mathbf{Set}$,
- $n \in \omega$,
- $X \in \mathbf{Obj}(\mathbf{Set})$,
- $t_i \in FX$ for all $i \in n$.

Then $t \in F(X^n)$ is a *set element coupling* iff for all i we have $F\pi_i t = t_i$. Or equivalently if TFDC:

$$\begin{array}{ccc} * & \xrightarrow{t} & FX^n \\ & \searrow t_i & \downarrow F\pi_i \\ & & FX \end{array}$$

set-element-coupling

6.0.1 New definitions of couplings (new)

We can generalize this definition so that it does not need to operate on set, also so that it acts on objects instead of elements, and not limit it to one object X but instead a collection of different objects:

Definition: Coupling

coupling

A *coupling* for $\prod_{i \in I} F(X_i)$ is a morphism $c : \prod_{i \in I} F(X_i) \rightarrow F(\prod_{i \in I} X_i)$ such that TFDC:

$$\begin{array}{c} \begin{array}{ccc} & \cdots & \\ & \swarrow & \searrow \\ & F_{X_{i_1}} & \\ \pi_{i_1} \nearrow & & \nwarrow F(\pi_{i_1}) \\ \pi_{i_0} \nearrow & F_{X_{i_0}} & \nwarrow F(\pi_{i_0}) \\ \prod_{i \in I} F(X_i) & \xrightarrow{c} & F(\prod_{i \in I} X_i) \end{array} \end{array}$$

i.e. $\forall_i^I \pi_i = F\pi_i \circ c$

this relates to the earlier definition above:

Coupling $\implies$ Coupling (original) for every object : Let c be a coupling DEF coupling on **Set** between $X_i = X$ for all $i \in I$ for a finite I (so $|I| \in \omega$). Then for any $(t_i)_{i \in I} \in \prod_{i \in I} F(X)$ is a set element coupling

DEF set-element-coupling.

Proof. Take $p \stackrel{\text{def}}{=} (t_i)_{i \in I} \in \prod_{i \in I} F(X)$. From the definition of coupling, we have that $\pi_i(p) = (F(\pi_i) \circ c)p = (F(\pi_i)(c(p)))$. Letting $t \stackrel{\text{def}}{=} c(p)$, we get $F(\pi_i)t = F\pi_i t$. Letting $n = |I| \in \omega$ we get exactly the definition of a set element coupling. $\square$

(note this is a simpler version of a distributive law for a monad, however we do not assume F is a monad, and also specify that the other functor is the product functor)

Suppose now that $c_R : R \rightarrow FR$ is a bisimulation between $c_X : X \rightarrow FX$, and $c_Y : Y \rightarrow FY$. Then note because of **DEF coalgebra-hom**, there must be some span on sets $X \xleftarrow{\phi_X} R \xrightarrow{\phi_Y} Y$, and because **Set** has products, by **DEF product** these must factor out into product maps. The question is, is there any way to given a bisimulation $c_R : R \rightarrow FR$, reduce it to a coalgebra of type $X \times Y \rightarrow F(X \times Y)$? this would allow for an easier way to interpret what the bisimulation is doing.

Result: Products of Coalgebras vs Products on Original Category

Let C be a category with products.

If a coupling $\gamma : FX \times FY \rightarrow F(X \times Y)$ exists, then for any two coalgebras $c_x : X \rightarrow FX$, $c_y : Y \rightarrow FY$ we have:

- The product of c_X, c_Y exists in the category of F -coalgebras.
- This product can be of type $c_{X \times Y} : X \times Y \rightarrow F(X \times Y)$

Proof. Let $c_X : X \rightarrow FX$, and $c_Y : Y \rightarrow FY$, and let $c_R : R \rightarrow FR$ be a bisimulation, with homomorphisms $\phi_X : R \rightarrow X$, $\phi_Y : R \rightarrow Y$. The goal is to show that a product exists, i.e. $\exists c_{X \times Y} : X \times Y \rightarrow F(X \times Y)$ such that for any bisimulation c_R , TFDC (in the category of F -coalgebras):

$$\begin{array}{ccccc}
 & & c_{X \times Y} & & \\
 & \swarrow \pi_X & \uparrow \exists! & \searrow \pi_Y & \\
 c_X & \xleftarrow{\phi_X} & c_R & \xrightarrow{\phi_Y} & c_Y
 \end{array}$$

Now going back to C , notice that since C has products, there exists an $f : R \rightarrow X \times Y$ such that TFDC (in C):

$$\begin{array}{ccccc}
 & & X \times Y & & \\
 & \swarrow \pi_X & \uparrow \exists! f & \searrow \pi_Y & \\
 X & \xleftarrow{\phi_X} & R & \xrightarrow{\phi_Y} & Y
 \end{array}$$

□

Now note that we assume c_R is a bisimulation, so we have TFDC (in C)

$$\begin{array}{ccccc}
 X & \xleftarrow{\phi_X} & R & \xrightarrow{\phi_Y} & Y \\
 c_X \downarrow & & c_R \downarrow & & \downarrow c_Y \\
 FX & \xleftarrow{F\phi_X} & FR & \xrightarrow{F\phi_Y} & FY
 \end{array}$$

Notice, that if we then consider the object $FX \times FY$, this is a product object in C . So it must have projection maps π_{FX}, π_{FY} . Furthermore, since $X \times Y \xrightarrow{\pi_X} X \xrightarrow{c_X} FX$ and $X \times Y \xrightarrow{\pi_Y} Y \xrightarrow{c_Y} FY$, this forms a span of FX and FY , so there must be a unique map $\exists! : X \times Y \rightarrow FX \times FY$ such that TFDC:

$$\begin{array}{ccccc}
 X & & & & \\
 \downarrow c_X & \swarrow \phi_X & & \searrow \pi_X & \\
 & R & \xrightarrow{\phi_Y} & Y & \xrightarrow{\pi_Y} & X \times Y \\
 & \downarrow c_R & & \downarrow c_Y & & \downarrow \exists! = \langle c_X, c_Y \rangle \\
 FX & \xleftarrow{F\phi_X} & FR & \xrightarrow{F\phi_Y} & FY & \xrightarrow{\pi_{FY}} & FX \times FY
 \end{array}$$

Now because F is a functor, the following diagram must commute as well (it is just applying the functor everywhere on the product diagram for $X \times Y$, and functors preserve commutative diagrams)

$$\begin{array}{ccccc}
 & & F(X \times Y) & & \\
 & \swarrow F(\pi_X) & \uparrow F(f) & \searrow F(\pi_Y) & \\
 FX & \xleftarrow{F(\phi_X)} & FR & \xrightarrow{F(\phi_Y)} & FY
 \end{array}$$

So we add it to our diagram:

$$\begin{array}{ccccccc}
 X & & & & & & \\
 \downarrow c_X & \swarrow \phi_X & & \searrow \pi_X & & & \\
 & R & \xrightarrow{\exists! f} & X \times Y & & & \\
 & \downarrow c_R & \searrow \phi_Y & \swarrow \pi_Y & & & \\
 FX & & Y & & & & \\
 \downarrow F\phi_X & \downarrow c_Y & \downarrow \pi_{FX} & & & & \\
 & FR & & FX \times FY & & & \\
 \downarrow F\pi_X & \downarrow F\phi_Y & \downarrow \pi_{FY} & & & & \\
 & Ff & & FY & & & \\
 & \downarrow Ff & & & & & \\
 & F(X \times Y) & & & & &
 \end{array}$$

Now note that since we have a coupling $\left(\begin{array}{c} \text{DEF} \\ \text{coupling} \end{array} \right) \gamma : FX \times FY \rightarrow F(X \times Y)$, it must commute with $\pi_X, \pi_Y, F\pi_X, F\pi_Y$ in the diagram above, so we get TFDC:

$$\begin{array}{ccccccc}
 X & & & & & & \\
 \downarrow c_X & \swarrow \phi_X & & \searrow \pi_X & & & \\
 & R & \xrightarrow{\exists! f} & X \times Y & & & \\
 & \downarrow c_R & \searrow \phi_Y & \swarrow \pi_Y & & & \\
 FX & & Y & & & & \\
 \downarrow F\phi_X & \downarrow c_Y & \downarrow \pi_{FX} & & & & \\
 & FR & & FX \times FY & & & \\
 \downarrow F\pi_X & \downarrow F\phi_Y & \downarrow \pi_{FY} & & & & \\
 & Ff & & FY & & & \\
 & \downarrow Ff & & & & & \\
 & F(X \times Y) & & & & & \\
 & & \nearrow \gamma & & & &
 \end{array}$$

Thus finally we see that there is a coalgebra $c_{X \times Y} \stackrel{\text{def}}{=} \gamma \circ \langle c_x, c_y \rangle : X \times Y \rightarrow F(X \times Y)$, and there are homomorphisms $f : c_R \rightarrow c_{X \times Y}$, $\pi_X : c_{X \times Y} \rightarrow c_X$, and $\pi_Y : c_{X \times Y} \rightarrow c_Y$. Now to show that the coalgebra homomorphism f (not f on **Set**, but f as a coalgebra homomorphism) is unique, simply note that any coalgebra homomorphism $p : c_R \rightarrow c_{X \times Y}$ must be a map $f : R \rightarrow X \times Y$, and since on **Set** this is unique, it must also be unique.

7 Coalgebra of MDPs (new)

Recall the MDP functor:

Definition: MDP functor

mdp-functor

Define the MDP functor on **Set** as:

$$F : S \rightarrow \mathbb{R} \times (\Delta S)^A \quad (13)$$

Note this will consider MDPs with state valued reward. Alternatively you could consider $F : S \rightarrow (\mathbb{R} \times \Delta S)^A$, which assigns a reward to a transition (s, a, s') instead of a state s . Much of the work we do should transfer up to some minor changes.

Result: MDP homomorphism

MDP-hom

Consider MDP coalgebras:

$$\mathcal{M}_X : X \rightarrow FX \quad (14)$$

$$: x \mapsto (r_x^X : \mathbb{R}, P_x^X : (\Delta X)^A) \quad (15)$$

$$\mathcal{M}_Y : Y \rightarrow FY \quad (16)$$

$$: y \mapsto (r_y^Y : \mathbb{R}, P_y^Y : (\Delta Y)^A) \quad (17)$$

Note that ϕ is a MDP-coalgebra homomorphism iff:

$$\begin{array}{ccc} X & \xrightarrow{\phi} & Y \\ \mathcal{M}_X \downarrow & & \downarrow \mathcal{M}_Y \\ FX & \xrightarrow{F\phi} & FY \end{array}$$

$$\dots \iff (F\phi) \circ \mathcal{M}_X = \mathcal{M}_Y \circ \phi \quad (18)$$

$$\iff \forall_x^X (F\phi) \circ \mathcal{M}_X(x) = \mathcal{M}_Y \circ \phi(x) \quad (19)$$

$$\iff \forall_x^X (F\phi)(r_x^X, P_x^X) = (r_{\phi(x)}^Y, P_{\phi(x)}^Y) \quad (20)$$

$$\iff \forall_x^X (r_x^X, (\Delta\phi)P_x^X) = (r_{\phi(x)}^Y, P_{\phi(x)}^Y) \quad (21)$$

$$\iff \forall_x^X \left(r_x^X, (y : Y, a : A) \mapsto \sum_{\substack{x' \in X \\ \phi(x')=y}} P_x^X(a, x') \right) = (r_{\phi(x)}^Y, P_{\phi(x)}^Y) \quad (22)$$

$$\iff \forall_x^X (r_x^X = r_{\phi(x)}^Y) \wedge \forall_y^Y \forall_a^A \left(\sum_{\substack{x' \in X \\ \phi(x')=y}} P_x^X(a, x') = P_{\phi(x)}^Y(a, y) \right) \quad (23)$$

$$\iff \forall_x^X (r_x^X = r_{\phi(x)}^Y) \wedge \forall_y^Y \forall_a^A \left(P^Y(\phi(x) \xrightarrow{a} y) = P^X(x \xrightarrow{a} \phi^{-1}[\{y\}]) \right) \quad (24)$$

7.1 Bisimulation

Recall the transition system version of bisimulation:

Definition: Standard MDP Bisimulation

standard-mdp-bisim

For an MDP $\mathcal{M} = (S, A, P, R, \gamma)$, a bisimulation is an equivalence relation $B \subseteq S^2$ such that:

$$sBs' \iff R(s) = R(s') \wedge \forall_a^A \forall_C^{S/B} P[s \xrightarrow{a} C] = P[s' \xrightarrow{a} C] \quad (25)$$

And bisimilarity is the union of all such B .

Now conversly, when you look at the coalgebraic version, some natural questions arise:

1. How can you get a relation out of a span?
2. If you take the relation, also assume $c_y = c_x$, does this definition correspond to the standard one?

For question 1, the natural choice for a relation is $\{(\phi_x(b), \phi_y(b)) | b \in B\}$. Or equivalently using categories, simply take the unique map f to the product, and the image of that map:

$$\begin{array}{ccccc} & & X \times Y & & \\ & \swarrow & \uparrow & \searrow & \\ X & \xleftarrow{\phi_X} & B & \xrightarrow{\phi_Y} & Y \end{array}$$

Definition: Induced Relation for **Set** bisimulation

induced-relation-bisimulation

Let F be an endofunctor on **Set**. For a bisimulation c_B of type given below:

$$\begin{array}{ccccc} X & \xleftarrow{\phi_0} & B & \xrightarrow{\phi_1} & X \\ c_X \downarrow & & \downarrow c_B & & \downarrow c_x \\ FX & \xleftarrow{F\phi_0} & FB & \xrightarrow{F\phi_1} & FX \end{array}$$

Let B_I denote the induced relation on X^2 , namely $\{\phi_X(b), \phi_X(b) | b \in B\}$.

For 2, it is a bit more tricky. I think it can be done if you assume that the resulting relation is an equivalence relation.

We give the formal proof first, then give some intuition.

Result: Coalgebraic Bisimulation is MDP bisimulation

Suppose c_B is an MDP-coalgebraic bisimulation:

$$c_X \xleftarrow{\phi_{x_0}} c_B \xrightarrow{\phi_{x_1}} c_X$$

Such that the induced relation B_I is an equivalence relation. Then B_I is a traditional MDP bisimulation.

Proof.

Forward direction: WTS $\forall_{x,x'} xB_I x' \implies r^X(x) = r^X(x') \wedge \forall_a^A \forall_C^{X/B_I} P^X[x \xrightarrow{a} C] = P^X[x' \xrightarrow{a} C]$
 Assume $xB_I x'$.

From DEF induced-relation-bisimulation, we have $b \in B$ such that $\phi_0(b) = x$, and $\phi_1(b) = x'$.

$$\implies r^B(b) = r^X(\phi_0(b)) = r^X(\phi_1(b)) \quad (26)$$

$$= r^X(x) = r^X(x') \quad (27)$$

Take arbitrary $a \in A$, $C \in X/B_I$. Note $\phi_0^{-1}[C] = \phi_1^{-1}[C]$, because:

$\phi_0^{-1}[C] = \phi_1^{-1}[C]$: Recall $B_I = \{\phi_0(b), \phi_1(b) | b \in B\}$.

So $\phi_0(b) \in C \iff \phi_1(b) \in C$. Thus:

$$\phi_0^{-1}[C] = \{b \in B | \phi_0(b) \in C\} \quad (28)$$

$$= \{b \in B | \phi_1(b) \in C\} \quad (29)$$

$$= \phi_1^{-1}[C] \quad (30)$$

Using the above, and RES MDP-hom:

$$\forall_c^C P^B[b \xrightarrow{a} \phi_0^{-1}[\{c\}]] = P^X[x \xrightarrow{a} c] \quad (31)$$

$$\implies P^B[b \xrightarrow{a} \phi_0^{-1}[C]] = P^X[x \xrightarrow{a} C] \quad (32)$$

$$= P^B[b \xrightarrow{a} \phi_1^{-1}[C]] = P^X[x' \xrightarrow{a} C] \quad (33)$$

Thus $P^X[x \xrightarrow{a} C] = P^X[x' \xrightarrow{a} C]$, and $r^X(x) = r^X(x')$.

Backward direction: WTS $\forall_{x,x'} r^X(x) = r^X(x') \wedge \forall_a^A \forall_C^{X/B_I} P^X[x' \xrightarrow{a} C] = P^X[x \xrightarrow{a} C] \implies xB_I x'$

Prove. May not be true, but you may be able to add more conditions until you get it. This in some sense says that the bisimulation is larger,

□

Intuition for Bisimulation: Now, notice that since B_I is an equivalence relation, you can view it as a matrix of size $|X| \times |X|$, where each entry corresponds to elements of $B_I \subseteq X^2$. Furthermore, you can reorder the indices so that the matrix is block diagonal (using transitivity, reflexivity, symmetry of B). In the following figure, the red dots are elements of B , the green squares correspond to the actual matrix entries (so each green square corresponds to a pair $(x_0, x_1) \in B_I \subseteq X^2$, and the blue squares represent equivalence classes of B . Everything outside of a blue square will be empty.




Figure 1: Bisimulation B grouped by equivalence classes

Notice that $P^B[b \xrightarrow{a} \phi_{X_0}^{-1}[\{x\}]] = P^X[\phi_{X_0}(b) \xrightarrow{a} x]$ and $P^B[b \xrightarrow{a} \phi_{X_1}^{-1}[\{x'\}]] = P^X[\phi_{X_1}(b) \xrightarrow{a} x']$ imply:




Figure 2: The Transitions from any element $b \in \phi_{X_0}^{-1}[\{x\}]$, to the entirety of $\phi_{X_0}^{-1}[\{x'\}]$

Let b be one of the red dots pictured
(i.e. some $b \in B$ s.t. $\phi_{X_0}(b) = x$)
Each transition above will happen with probability:

$$P^B[b \xrightarrow{a} \phi_{X_0}[\{x'\}]] = P^X[x \xrightarrow{a} x']$$




Figure 3: The transitions from any element $b \in \phi_{X_1}^{-1}[\{x\}]$, to the entirety of $\phi_{X_1}^{-1}[\{x''\}]$

Let b be one of the red dots pictured
(i.e. some $b \in B$ s.t. $\phi_{X_1}(b) = x$)
Each transition above will happen with probability:

$$P^B[b \xrightarrow{a} \phi_{X_1}[\{x'\}]] = P^X[x \xrightarrow{a} x']$$

Now let $C = \{q \mid qB_I x'\}$ be an equivalence class of B_I that includes x' .
So we can see that by summing up instances of the last picture, we get:




Figure 4: Transitions from any element $b \in \phi_{X_1}^{-1}[\{x\}]$, to $\phi_{X_1}^{-1}[C]$, i.e. anything related to x'

In the figure above, the probability of each of these transitions is

$$P^B \left[b \xrightarrow{a} \phi_{X_1}^{-1}[C] \right] = P^X[x \xrightarrow{a} C]$$

Likewise the same argument holds when you sum up columns instead, and you get

$$P^B \left[b \xrightarrow{a} \phi_{X_0}^{-1}[C] \right] = P^X[x \xrightarrow{a} C]$$

Note that because B_I is an equivalence relation, it must be that $\phi_{X_0}^{-1}[C] = \phi_{X_1}^{-1}[C]$.

8 LDBA/LDBW Coalgebraically (new)

8.1 Automata review

Recall:

Definition: Generalized Buchi Automaton (GBA)

GBA

A Generalized Buchi Automaton (GBA) is a tuple

$$\mathcal{B} = (Q, \Sigma, \delta, \alpha) \quad (34)$$

such that:

- Q is a set (states)
- Σ is a set (alphabet)
- $\delta : Q \rightarrow \Sigma \rightarrow 2^Q$ is a transition function (NDTM)
- $\alpha = \{F_1, \dots, F_n\}$ with $F_i : 2^Q$ is a set of accepting conditions

Coalgebraically, we can view a GBA as $F : Q \rightarrow (2^Q)^\Sigma \times 2^n$, where the left part yields the next possible states given a letter, and the right part gives which (if any) of the acceptance conditions the current state is a part of.

Definition: Pointed GBA

A pointed GBA is a GBA $\mathcal{B} = (Q, \Sigma, \delta, \alpha)$ along with an initial state $q_0 \in Q$.

For now we will consider non pointed GBA, but there exist pointed coalgebras, so perhaps pointed GBAs reduce to pointed coalgebras.

Definition: $\inf \tau$ for trajectory τ

For a run/trajectory $\tau : (Q \times \Sigma \times Q)$, let:

$$\inf \tau \stackrel{\text{def}}{=} \{q | (q, \sigma, q') \in \tau \text{ occurs infinitely often in } \tau\} \quad (35)$$

Accepting Run: A run $\tau : (Q \times \Sigma \times Q)$ is accepted if $\inf \tau \circ F_i \neq 0$ for all F_i .

Definition: Limit Deterministic Buchi Automata (LDBA)

A GBA is a LDBA iff Q can be partitioned $Q = Q_I \sqcup Q_D$ such that:

- Transitions from Q_D must remain in Q_D , and must be deterministic:
For any $q_d : Q_D, \sigma : \Sigma, \delta(q, \sigma) \subseteq Q_D$ and $|\delta(q, \sigma)| = 1$.
- Accepting states are in Q_D :
For any $F_i : \alpha, F_i \subseteq Q_D$.

So the definition of a GBA can be refined for a LDBA to be:

$$(\Sigma, Q_I \sqcup Q_D, \delta_I, \delta_D, G) \quad (36)$$

where:

- $\delta_I : Q_I \rightarrow \Sigma \rightarrow 2^{Q_I} \times 2^{Q_D}$

- $\delta_D : Q_D \rightarrow \Sigma \rightarrow Q_D$
- $G : Q_D \rightarrow 2^k$ (tells you which of the acceptance criteria are met)

Coalgebraically this is more tricky. Our goal is to find a functor F such that every F -coalgebra can be modeled by an LDBA, and likewise that every LDBA can be modeled by a F -coalgebra.

The naïve approach would be something like: $F : Q \rightarrow (2^Q)^\Sigma + (Q^\Sigma \times 2^k)$. The problem is enforcing that transitions from Q_D remain in Q_D . (otherwise you can construct a coalgebra where the output of one of the deterministic transitions is something not in Q_D). So we need to enforce that if the coalgebra “knows” that the input was in Q_D , it should only output elements in Q_D .

We can model this however by going to **Set/2**, i.e. **slice-cat**. This will consist of objects of the form $m_X : X \rightarrow 2$ for any set X , and morphisms of the form $f : m_X \rightarrow m_Y$ which make TFDC:

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \searrow m_x & \swarrow m_y \\ & 2 & \end{array}$$

Note that any object $m_X : X \rightarrow 2$ in **Set/2** can be thought of as a partition $X = m_X^{-1}[\{0\}] + m_X^{-1}[\{1\}]$.

Furthermore any morphism $m_X \rightarrow m_Y$ must preserve the partition, i.e. $m_X^{-1}[\{0\}] = m_Y^{-1}[\{0\}]$ and $m_X^{-1}[\{1\}] = m_Y^{-1}[\{1\}]$.

Equivalently, we can think of **Set/2** as **Set** $\times$ **Set**. In particular, notice that **Set** $\times$ **Set** is exactly **Set/2**, just with objects relabeled.

Namely there is a bijection:

Set $\times$ **Set** is **Set/2**: There is a bijective functor $F : \mathbf{Set} \times \mathbf{Set} \rightarrow \mathbf{Set/2}$.

Namely for any $(X, Y) : \mathbf{Obj}(\mathbf{Set} \times \mathbf{Set})$, define F :

- On objects:

$$F(X) : X + Y \rightarrow 2 \tag{37}$$

$$F(X) \stackrel{\text{def}}{=} (x : X + Y) \mapsto \begin{cases} 0 & x : X \\ 1 & x : Y \end{cases} \tag{38}$$

- On morphisms (given $(f, g) : (X_0, X_1) \rightarrow (Y_0, Y_1)$):

$$F((f, g)) : (X_0 + X_1) \rightarrow (Y_0 + Y_1) \tag{39}$$

$$F((f, g)) \stackrel{\text{def}}{=} (x : X_0 + X_1) \mapsto \begin{cases} f(x) & x : X_0 \\ g(x) & x : X_1 \end{cases} \tag{40}$$

Proof. To show this is a bijection, define the inverse $G : \mathbf{Set/2} \rightarrow \mathbf{Set} \times \mathbf{Set}$:

- On objects: $G(m_X) \stackrel{\text{def}}{=} (m_X^{-1}[\{0\}], m_X^{-1}[\{1\}])$.
- On morphisms: $G(f : m_X \rightarrow m_Y) \stackrel{\text{def}}{=} (f|_{m_X^{-1}[\{0\}]}, f|_{m_X^{-1}[\{1\}]})$, i.e. the restriction of f to everything that maps to 0, and to 1.

Note that $GF = 1_{\mathbf{Set} \times \mathbf{Set}}$, and likewise $FG = 1_{\mathbf{Set/2}}$ for both objects and morphisms. □

(This is called an isomorphism of categories, it essentially means one is just relabeling the other)

This gives a more convenient way to package **Set/2**, and also gives an easier way to define how functors act on it.

Definition: LDBA Functor

Consider the category $\mathbf{Set} \times \mathbf{Set}$. Define the LDBA functor such that:

- On objects:

$$F(Q_I, Q_D) \stackrel{\text{def}}{=} ((2^{Q_I} \times 2^{Q_D})^\Sigma, Q_D^\Sigma \times 2^k) \quad (41)$$

Here we can see that an element of $F(X_I, X_D)$ is of the form:

$$\left[\left((\sigma : \Sigma) \mapsto (A_\sigma : 2^{Q_I}, B_\sigma : 2^{Q_D}) \right), \left((\sigma : \Sigma) \mapsto (\delta : Q_D^\Sigma, G_\sigma : 2^k) \right) \right] \quad (42)$$

- **On morphisms:** Given a morphism $(f : X_I \rightarrow Y_I, g : X_D \rightarrow Y_D)$, define

$$F(f, g) \stackrel{\text{def}}{=} \left[\left(\sigma \mapsto (A_\sigma, B_\sigma) \right), \left(\sigma \mapsto (\delta, G_\sigma) \right) \right] : F(X_I, X_D) \quad (43)$$

$$\mapsto \left[\left(\sigma \mapsto (f[A_\sigma], g[B_\sigma]) \right), \left(\sigma \mapsto (g \circ \delta, G_\sigma) \right) \right] : F(Y_I, Y_D) \quad (44)$$

or equivalently:

$$F(f, g) \stackrel{\text{def}}{=} \left[\left((f[\cdot], g[\cdot]) \circ (\cdot) \right), \left(g \circ (\cdot), \text{id}_{2^k} \right) \right] \quad (45)$$

Notice that because we work on $\mathbf{Set} \times \mathbf{Set}$, and due to the product nature of the functor, we can decompose any LDBA coalgebra:

Equivalent LDBA coalgebra : Any LDBA coalgebra $c_Q : Q_I \times Q_D \rightarrow ((2^{Q_I} \times 2^{Q_D})^\Sigma, Q_D^\Sigma \times 2^k)$ must consist of two maps:
(from definition of morphisms on $\mathbf{Set} \times \mathbf{Set}$)

$$c_{Q_I} : Q_I \rightarrow (2^{Q_I} \times 2^{Q_D})^\Sigma \quad (46)$$

$$c_{Q_D} : Q_D \rightarrow (Q_D^\Sigma \times 2^k) \quad (47)$$

So a LDBA coalgebra can also be defined by (c_{Q_I}, c_{Q_D}) with types given above.

Coalgebraic LDBA model LDBA : There is a (class) bijection between the class of all LDBA and the objects of the category of LDBA-coalgebras, **up to an initial state**.

Proof. The following maps are (class) inverses of each other:

- **(LDBA $\rightarrow$ Coalgebra)** Given a LDBA (without initial state) $\mathcal{B} = (\Sigma, Q_I \sqcup Q_D, \delta_I, \delta_D, G)$, you can define a LDBA coalgebra with:

$$- c_{Q_I} = \delta_I$$

$$- c_{Q_D} = q \mapsto (\delta_D(q), G(q))$$

- **(Coalgebra $\rightarrow$ LDBA)** Given a coalgebra $(c_{Q_I}, c_{Q_D}) : (Q_I, Q_D) \rightarrow F(Q_I, Q_D)$, define a LDBA as:

$$(\Sigma, Q_I \sqcup Q_D, c_{Q_I}, \pi_0 \circ c_{Q_D}, \pi_1 \circ c_{Q_D}) \quad (48)$$

□

Result: LDBA Homomorphisms

Let:

- $c_X = (c_{X_I}, c_{X_D})$ be a LDBA coalgebra : $(X_I, X_D) \rightarrow F(X_I, X_D)$.

Let $(X_I \sqcup X_D, \delta_{X_I}, \delta_{X_D}, G_X)$ be the corresponding LDBA.

- $c_Y = (c_{Y_I}, c_{Y_D})$ be a LDBA coalgebra : $(Y_I, Y_D) \rightarrow F(Y_I, Y_D)$

Let $(Y_I \sqcup Y_D, \delta_{Y_I}, \delta_{Y_D}, G_Y)$ be the corresponding LDBA.

Then a coalgebra homomorphism $\phi = (\phi_I, \phi_D) : c_X \rightarrow c_Y$ is:

ϕ will make TFDC:

$$\begin{array}{ccc} (X_I, X_D) & \xrightarrow{(\phi_I, \phi_D)} & (Y_I, Y_D) \\ \downarrow (c_{X_I}, c_{X_D}) & & \downarrow (c_{Y_I}, c_{Y_D}) \\ F(X_I, X_D) & \xrightarrow{F(\phi_I, \phi_D)} & F(Y_I, Y_D) \end{array}$$

which commutes iff both of the following commute:

$$\begin{array}{ccc} X_I & \xrightarrow{\phi_I} & Y_I \\ \downarrow c_{X_I} & & \downarrow c_{Y_I} \\ (2^{X_I} \times 2^{X_D})^\Sigma & \xrightarrow{(\phi_I[\cdot], \phi_D[\cdot]) \circ (\cdot)} & (2^{Y_I} \times 2^{Y_D})^\Sigma \end{array} \quad \begin{array}{ccc} X_D & \xrightarrow{\phi_D} & Y_D \\ \downarrow c_{X_D} & & \downarrow c_{Y_D} \\ (X_D^\Sigma \times 2^k) & \xrightarrow{(\phi_D \circ (\cdot), \text{id}_{2^k})} & (Y_D^\Sigma \times 2^k) \end{array}$$

$$\begin{aligned} \text{LHS Commutes} &\iff (\phi_I[\cdot], \phi_D[\cdot]) \circ c_{X_I} = c_{Y_I} \circ \phi_I \\ &\iff \forall_{x_I}^{X_I} (\phi_I[\cdot], \phi_D[\cdot]) \circ c_{X_I}(x_I) = c_{Y_I} \circ \phi_I(x_I) \end{aligned}$$

$$\begin{aligned} \text{RHS commutes} &\iff (\phi_D \circ (\cdot), \text{id}_{2^k}) \circ c_{X_D} = c_{Y_D} \circ \phi_D \\ &\iff \forall_{x_D}^{X_D} (\phi_D \circ (\cdot), \text{id}_{2^k}) \circ c_{X_D}(x_D) = c_{Y_D} \circ \phi_D(x_D) \end{aligned}$$

So to summarize, the initial part looks like (for all $\sigma : \Sigma$):




Figure 5: Initial part of the homomorphism

For the final part:




Figure 6: Deterministic part of the homomorphism

9 Coalgebraic Behavioral Metrics

This section summarizes

Reference coalgebraic behavioral metrics paper

The goal of this section is to define a quantitative notion of bisimulation, namely a bisimulation (pseudo) metric for coalgebras.

The main idea: If you have some pseudometric d^X on X , and also a way to lift d^X to a pseudometric d^{FX} on FX , then

Essentially the goal is to lift a metric on X^2 to a metric on $F(X^2)$.

First some review some definitions on pseudometrics, and some notation/assumptions.

Then we cover some category theory definitions that we will use.

9.1 Metric review and notation

Definition: Pseudometric Space

A pseudometric space is a pair (X, d^X) , for a set X and function $d^X : X \times X \rightarrow [0, T]_{\mathbb{R}}$ for some $T \in \mathbb{R}$, subject to: $(\forall_{x,y,z}^X)$

- (Symmetry) $d^X(x, y) = d^X(y, x)$
- (Triangle Inequality) $d^X(x, z) \leq d^X(x, y) + d^X(y, z)$
- (Reflexivity) $d^X(x, x) = 0$

(Note we are assuming a pseudometric is bounded by some fixed $T \in \mathbb{R}$)

Definition: Metric

A pseudometric space (X, d^X) is a metric space iff it satisfies Definiteness, i.e.

$$\forall_{x,y}^X d^X(x, y) \neq 0 \implies d(x, y) > 0 \quad (49)$$

i.e. equivalently (under the pseudometric axioms):

$$d^X(x, y) = 0 \iff x = y \quad (50)$$

Notation T, F : For the rest of this section, let $T \in \mathbb{R}$ be some fixed real number, and $F : \mathbf{Set} \rightarrow \mathbf{Set}$ be some endofunctor on $\mathbf{Set}$.

9.2 Categorical Prerequisites

Definition: PMet**PMet**

Define **PMet** as the category with:

- $\mathbf{Obj}(\mathbf{PMet}) \stackrel{\text{def}}{=} \text{all pseudo-metric spaces } (X, d^X).$
- $\mathbf{Hom}(\mathbf{PMet}) \stackrel{\text{def}}{=} \text{all nonexpansive functions, i.e. } f : (X, d^X) \rightarrow (Y, d^Y) \text{ where } d^Y \circ (f, f) \leq d^X.$
(here $\leq$ between functions is the pointwise order, i.e. $f \leq g$ iff $f(x) \leq g(x)$ for all $x \in X$)
- $\circ$ is the composition of functions.
- id_X is the identity function on X .

Note that this is indeed a category, as nonexpansive functions are closed under composition, and the identity function is nonexpansive.

Definition: Lifting of endofunctor from Set to PMet**lift-functor-pmet**

Let $F : \mathbf{Set} \rightarrow \mathbf{Set}$. A lifting of F to **PMet** is a functor $\bar{F}$ such that TFDC:

$$\begin{array}{ccc} \mathbf{PMet} & \xrightarrow{\bar{F}} & \mathbf{PMet} \\ \text{Forget} \downarrow & & \downarrow \text{Forget} \\ \mathbf{Set} & \xrightarrow{F} & \mathbf{Set} \end{array}$$

(in the category of small categories)

Given a pseudometric space (X, d) , define the metric d^F as the output $(FX, d^F) = \bar{F}(X, d)$.

The general idea is this:

General idea for Coalgebraic Behavioral Metrics :

Given an F -coalgebra $c : X \rightarrow FX$, we want to turn this into a $\bar{F}$ -coalgebra $((X, d_C), c)$ such that $c : (X, d_C) \rightarrow (FX, d_C^F)$ is nonexpansive (c is a **PMet** morphism). Note that $d \mapsto d^F \circ (c, c)$ is monotone. By Knaster Tarski, this has a fixed point. Let d_C be this fixed point:

$$d_C \stackrel{\text{def}}{=} \inf \{d : \text{pseudometric}(: X^2 \rightarrow [0, T]_{\mathbb{R}}) | d_F \circ (c, c) \leq d\} \quad (51)$$

(here inf is taken over the poset of pseudometrics)

Since $d_C = d_C^F \circ (c, c)$, we have that c is an isometry

An important result is that $\bar{F}$ on objects is monotone:

Result: $\bar{F}$ is monotone

1. Let $F : \mathbf{Set} \rightarrow \mathbf{Set}$.
2. Let $\bar{F} : \mathbf{PMet} \rightarrow \mathbf{PMet}$ be as in **DEF lift-functor-pmet**.
3. Let $(X, d_0), (X, d_1)$ be pseudometrics.

Then:

$$d_0^F \leq d_1^F$$

Proof. Recall first that from **DEF PMet**, morphisms only consist of non expansive functions.

Thus $d_0 \leq d_1 \iff$ there is some $f : \mathbf{Hom}^{\mathbf{PMet}}((X, d_0), (X, d_1))$ such that $Uf = \text{id}_X$.

Now assume $d_0 \leq d_1$. Then take f like above.

Then $\bar{F}f : (FX, d_0^F) \rightarrow (FX, d_1^F)$ must exist by functoriality.

Furthermore, because $\bar{F}$ is a lifting, $U\bar{F}f = \text{id}_X$. Thus $f_0^F \leq f_1^F$. □

9.3 Evaluations

Now consider coalgebras on **Set**.

We can think of $\mathbf{Set}/[0, T]_{\mathbb{R}}$ (recall DEF slice-cat) as having objects that have a sort of value function $V_X : X \rightarrow [0, T]_{\mathbb{R}}$, which assigns a value to each state.

Note that also any pseudometric space (X, d^X) can be thought of as an object in $\mathbf{Set}/[0, T]_{\mathbb{R}}$, namely $d^X : X \times X \rightarrow [0, T]_{\mathbb{R}}$.

Definition: Evaluation Function, Evaluation Functor

Given a fixed **Set** endofunctor F . We want to find a way to lift F to $\mathbf{Set}/[0, T]_{\mathbb{R}}$:


Evaluation Function **eval**: An evaluation function is any F algebra with domain $[0, T]_{\mathbb{R}}$, i.e. any map:

$$\text{eval} : F([0, T]_{\mathbb{R}}) \rightarrow [0, T]_{\mathbb{R}} \quad (52)$$

Evaluation Functor $\tilde{F}$: Given a fixed **eval**, $\tilde{F}$ is an endofunctor on $\mathbf{Set}/[0, T]_{\mathbb{R}}$ that takes:

- On objects: $V_X \mapsto \text{eval} \circ FV_X$
- On morphisms: $f \mapsto Ff$

This means you can lift any diagram in the slice category. Pictorially, this means that you can apply $\tilde{F}$ everywhere to a diagram:



9.4 Kantorovic Metric

Definition: Kantorovic Metric

Given F , **eval**, then for any pseudometric (X, d) define d_K :

$$d_{K(X, d^X)}(t_1, t_2) \stackrel{\text{def}}{=} \sup \left\{ d^{[0, T]_{\mathbb{R}}}(\tilde{F}f(t_1), \tilde{F}f(t_2)) \mid f : \mathbf{Hom}^{\mathbf{PMet}}((X, d), ([0, T]_{\mathbb{R}}, d^{[0, T]_{\mathbb{R}}})) \right\} \quad (53)$$

$$= \sup \left\{ \left| \tilde{F}V_X(t_1) - \tilde{F}V_X(t_2) \right| \mid V_X : X \rightarrow [0, T]_{\mathbb{R}}, V_X \text{ non expansive} \right\} \quad (54)$$

Pictorially, it is sup over all non expansive functions $V_X : (X, d^X) \rightarrow ([0, T]_{\mathbb{R}}, d^{[0, T]_{\mathbb{R}}})$ such that TFDC:

$$\begin{array}{ccc}
 FX \times FX & \xrightarrow{\sup_{V_X} FV_X \times FV_X} & F[0, T]_{\mathbb{R}} \times F[0, T]_{\mathbb{R}} \\
 \downarrow d_K & & \downarrow \text{eval} \times \text{eval} \\
 [0, T]_{\mathbb{R}} & \xleftarrow{d^{[0, T]_{\mathbb{R}}}} & [0, T]_{\mathbb{R}} \times [0, T]_{\mathbb{R}}
 \end{array}$$

where $d^{[0, T]_{\mathbb{R}}}$ is L1 distance on $\mathbb{R}$.

Definition: Kantorovich Lifting/Functor

The Kantorovich lifting (functor) of $F : \mathbf{Set} \rightarrow \mathbf{Set}$ is $F_K : \mathbf{PMet} \rightarrow \mathbf{PMet}$ such that $F_K(X, d^X) = (X, d_{K(X, d^X)})$, and $F_K f = Ff$.

Show more examples, in particular on Δ

Useful results :

1. If (X, d) is a pseudometric, then so is d_K .
2. F_K for a functor F preserves isometries
- 3.

Kantorovich metric for MDP (new) : Recall the MDP functor DEF mdp-functor.

Below represents applying $\text{eval} \circ FV_X$ for some arbitrary $V_X : X \rightarrow \mathbb{R}$.
Suppose you have an element

$$(r_0, P_{X_0}) : FX = \mathbb{R} \times (\Delta(X))^A$$

After applying FV_X , we get:

$$(r_0, P^{\mathcal{M}}[x_0 \xrightarrow{a} V_X^{-1}[\{\cdot\}]]) : F[0, T]_{\mathbb{R}}$$

Now let $\text{eval} : F[0, T]_{\mathbb{R}} \rightarrow \mathbb{R}$ be the evaluation function. This can be arbitrary, but a natural choice is to take the current reward, and sum it with the discounted expected next value. To get rid of a we can apply max. Applying this to the above:

$$r_0 + \gamma \max_{a:A} \sum_{t \in [0, T]_{\mathbb{R}}} t P^{\mathcal{M}}[x_0 \xrightarrow{a} V_X^{-1}[\{t\}]]$$

In particular, if (r_0, P_{X_0}) is the output from state x_0 , then we get that the above is:

$$= r^{\mathcal{M}}(x_0) + \gamma \max_{a:A} \mathbb{E}_{x_0 \xrightarrow{a} x'} [V_X(x')]$$

Note that this is exactly the DEF bellman-operator $\mathcal{T}$, applied to V_X .

you also need to bound reward suitably, like instead of $\mathbb{R}$ do $[0, (1 - \gamma)T]_{\mathbb{R}}$

Thus the Kantorovich metric for MDPs is:

Not nice metric for functor: Consider the functor $F : X \rightarrow A \times X$. Suppose you wanted to find the wasserstein metric.

Note that from DEF coupling, suppose $(x_0, b_0), (x_1, b_1)$ are two elements of FX . Then (x_0, b_0) can only be coupled iff $b_0 = b_1$.

This is not what you would expect if B had some sort of metric structure already, say $B = \mathbb{R}$. In fact the wasserstein metric removes all metric information (namely it assumes the discrete topology on B by default).

To fix this issue, we will introduce multifunctors, and lifting pseudometrics of mutlifunctors.

10 More complicated results

10.1 Limits/Colimits, Pullbacks

Definition: Limit

limit

Fill in

Weak limit :

weak-limit

A weak limit is just like a limit, but the map does not need to be unique.

Definition: Colimit


colimit

Fill in

Definition: Pullback


pullback

Let $X \xrightarrow{\phi_X} Z \xleftarrow{\phi_Y} Y$ be a cospan. A pullback of this cospan is the limit of the diagram. More concretely, it is an object P with maps $p_X : P \rightarrow X$ and $p_Y : P \rightarrow Y$ such that for any other object Q with maps $q_X : Q \rightarrow X$, $q_Y : Q \rightarrow Y$, there is a unique map $! : Q \rightarrow P$ TFDC:



Definition: Weak Pullback

A weak pullback is a weak limit of a cospan $X \xrightarrow{\phi_X} Z \xleftarrow{\phi_Y} Y$. I.e. see **DEF pullback**, but the map $! : Q \rightarrow P$ does not have to be unique:



10.2 Comparing Bisimulation

First we will need the following definitions/results:

Weak pullback preserving functor : Let $F : C \rightarrow D$ be a functor.

F preserves weak pullbacks iff for any pullback

$$\begin{array}{ccc} P & \xrightarrow{p_Y} & Y \\ p_X \downarrow & & \downarrow \phi_Y \\ X & \xrightarrow{\phi_X} & Z \end{array}$$

we have that the following commutative diagram is also a pullback of cospan $FX \xrightarrow{F\phi_X} FZ \xleftarrow{F\phi_Y} FY$:

$$\begin{array}{ccc} FP & \xrightarrow{Fp_Y} & FY \\ Fp_X \downarrow & & \downarrow F\phi_Y \\ FX & \xrightarrow{F\phi_X} & FZ \end{array}$$

(meaning if there is a $Q \in \mathbf{Obj}(D)$ with morphisms to FX and FY , it must factor through FP)

Result: AM Bisimulation is weaker than Behavioral Equivalence

AM-vs-beh-equiv

Suppose $F : C \rightarrow C$ preserves weak pullbacks, and C has weak pullbacks. Then for any AM bisimulation $c_X \xleftarrow{\phi_X} c_R \xrightarrow{\phi_Y} c_Y$ there is a behavioral equivalence.

We will use the following:

$$\begin{array}{ccc} \exists c_R & \longrightarrow & c_Y \\ \downarrow & & \downarrow \\ c_X & \longrightarrow & * \end{array}$$

So basically we want to show that given a

10.3 Existence of final coalgebra

(A slightly more involved result shows how to build a final coalgebra)

Definition: (Ordinal Indexed) Chain, Cochain (new?)

Given an ordinal α , an α -chain is a diagram of the form:

$$A_0 \longrightarrow A_1 \longrightarrow A_2 \longrightarrow \cdots \longrightarrow A_\omega \cong \text{colim}(A_0 \rightarrow \cdots) \longrightarrow A_{\omega+1} \longrightarrow \cdots \longrightarrow A_\alpha$$

Likewise a α -cochain is the dual, i.e. a diagram of the form:

$$A_0 \longleftarrow A_1 \longleftarrow A_2 \longleftarrow \cdots \longleftarrow A_\omega \cong \text{lim}(A_0 \leftarrow \cdots) \longleftarrow A_{\omega+1} \longleftarrow \cdots \longleftarrow A_\alpha$$

Result: Final coalgebra exists (Pohlova '73, Adamek '74)

Let C be a category with terminal object 1.

Consider the following ω -cochain:

$$\begin{array}{c} L_F = \text{lim}(1 \leftarrow F(1) \leftarrow \cdots) \\ \swarrow \exists ! l_0 \quad \downarrow l_1 \quad \downarrow l_2 \quad \searrow \\ 1 \xleftarrow{\exists ! f} F(1) \xleftarrow{F(f)} F^2(1) \xleftarrow{\quad} \cdots \end{array}$$

Suppose that L_F exists, and that F preserves this limit. Then:

- There exists a coalgebra $: L_F \rightarrow F(L_F)$
- This coalgebra is final

Proof. Now consider shifted chain:

$$\begin{array}{ccccc}
 & L_F^{\text{SH}} = \lim(F(1) \leftarrow \dots) & & & \\
 & \swarrow l_1 \quad \downarrow l_2 \quad \searrow & & & \\
 F(1) & \xleftarrow{F(f)} & F^2(1) & \xleftarrow{\quad} & \dots
 \end{array}$$

Since F preserves this limit, we have that $F(L_F)$ is a limit of the shifted chain and $l_1, \dots$. Thus we have two limits so they must be isomorphic: $\text{pres}_F : F(L_F) \cong L_F^{\text{SH}}$.

Note L_F^{SH} is also a cone over $1 \xleftarrow{f} F(1) \leftarrow \dots$ (take $! : L_F^{\text{SH}} \rightarrow 1$)
 $(L_F \text{ is a limit}) \implies \exists ! : L_F^{\text{SH}} \rightarrow L_F$.

Note L_F is a cone over $F(1) \xleftarrow{F(f)} F^2(1) \leftarrow \dots$
 $(L_F^{\text{SH}} \text{ is a limit}) \implies \exists ! : L_F \rightarrow L_F^{\text{SH}}$.

Thus $L_F \cong L_F^{\text{SH}}$.

Now take the coalgebra $L_F \rightarrow L_F^{\text{SH}} \xrightarrow{\text{pres}_F} F(L_F)$.

□