

Rational Polynomials: Graphing and Asymptotes

Find the intercepts, if there are any.

Step 1: Set the numerator to 0 to solve for horizontal intercepts.

Step 2: Set the x to 0 to solve for vertical intercept.

Step 3: Set the denominator to 0 to solve for vertical asymptotes.

Step 4: Perform a long division to find the quotient which specifies the oblique asymptote.


Note: Blue curve the actual Rational function.

Red and Gold asymptotes.

Example: Horizontal Asymptote

$$\frac{x-3}{2x-5}$$


$$\begin{array}{r} + \left(\frac{1}{2} \right) \\ \hline \boxed{2x-5} \quad (1)x + (-3) \\ \quad \quad \quad \left(\frac{x}{2} \right) + \left(-\frac{5}{2} \right) \\ \quad \quad \quad + \left(-\frac{1}{2} \right) \end{array}$$



Example: Oblique Linear Asymptote

$$\frac{(x-1)(x+2)}{x+1}$$

$$\begin{array}{r} + \left(x \right) \\ \hline \boxed{x+1} \quad (1)x^2 + (1)x + (-2) \\ \quad \quad \quad \left(x^2 \right) + \left(x \right) \\ \quad \quad \quad + \left(-2 \right) \end{array}$$



Example: Multiple Vertical Asymptotes

$$\frac{x-2}{(x+1)(x+2)}$$

$$\begin{array}{r} + \left(0 \right) \\ \hline \left(x \right) + \left(-2 \right) \end{array}$$

