

Intercepts of the Quadratic

Given a quadratic $k(m) = am^2 + bm + c$ compute its discriminant Δ :

$$\Delta = \sqrt{b^2 - 4ac}$$

Case1: $\Delta > 0$

$m_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ computes the m-intercepts of multiplicity 1.
 $k(0) = c$ computes the single k-intercept.


Example 1.

$k(m) = 3m^2 - 16m - 64$ compute its discriminant Δ :

$$\Delta = 1024 > 0$$

$$m_{1,2} = -\frac{8}{3}, 8$$

$k(0) = -64$ k-intercept.



Case2: $\Delta = 0$

$m_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-b \pm 0}{2a} = \frac{-b}{2a}$ single m-intercept of multiplicity 2.


Example 2.

$k(m) = -3m^2 - 6m - 3$ compute its discriminant Δ :

$$\Delta = 0$$

$$m_{1,2} = -1, -1$$

$k(0) = -3$ k-intercept.



Case3: $\Delta < 0$

$\sqrt{b^2 - 4ac}$ has no value in Real Numbers. Therefore there are no m-intercepts.

However there is a k-intercept.

Example 3.

$k(m) = 4m^2 + 80m + 500$ compute its discriminant Δ :

$$\Delta = -1600 < 0$$

$k(0) = 500$ k-intercept.

