

Rational Polynomials: Graphing and Asymptotes

Find the intercepts, if there are any.

Step 1: Set the numerator to 0 to solve for horizontal intercepts.

Step 2: Set the x to 0 to solve for vertical intercept.

Step 3: Set the denominator to 0 to solve for vertical asymptotes.


Step 4: Perform a long division to find the quotient which specifies the oblique asymptote.

Note: Blue curve the actual Rational function.
Red and Gold asymptotes.

Example: Horizontal Asymptote

$$\frac{x-2}{5x-3}$$


$$\begin{array}{r} + \left(\frac{1}{5}\right) \\ \hline 5x-3 \quad (1)x + (-2) \\ \left(\frac{x}{1}\right) + \left(-\frac{3}{5}\right) \\ + \left(-\frac{7}{5}\right) \end{array}$$



Example: Oblique Linear Asymptote

$$\frac{(x-2)(x+3)}{x-1}$$

$$\begin{array}{r} + \left(x\right) + \left(2\right) \\ \hline x-1 \quad (1)x^2 + (1)x + (-6) \\ \left(x^2\right) + \left(-x\right) \\ + (2)x + (-6) \\ + \left(2x\right) + \left(-2\right) \\ + \left(-4\right) \end{array}$$



Example: Multiple Vertical Asymptotes

$$\frac{x+3}{(x-2)(x-1)}$$

$$\begin{array}{r} + \left(0\right) \\ \hline \left(x\right) + \left(3\right) \end{array}$$

