

Intercepts of the Quadratic

Given a quadratic $g(k) = a k^2 + b k + c$ compute its discriminant Δ :

$$\Delta = \sqrt{b^2 - 4ac}$$

Case1: $\Delta > 0$

$k_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ computes the k-intercepts of multiplicity 1.

$g(0) = c$ computes the single g-intercept.


Example 1.

$g(k) = -2k^2 + 16k + 18$ compute its discriminant Δ :

$$\Delta = 400 > 0$$

$$k_{1,2} = -1, 9$$

$g(0) = 18$ g-intercept.



Case2: $\Delta = 0$

$k_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-b \pm 0}{2a} = \frac{-b}{2a}$ single k-intercept of multiplicity 2.


Example 2.

$g(k) = 3k^2 - 36k + 108$ compute its discriminant Δ :

$$\Delta = 0$$

$$k_{1,2} = 6, 6$$

$g(0) = 108$ g-intercept.



Case3: $\Delta < 0$

$\sqrt{b^2 - 4ac}$ has no value in Real Numbers. Therefore there are no k-intercepts.

However there is a g-intercept.

Example 3.

$g(k) = -9k^2 - 162k - 810$ compute its discriminant Δ :

$$\Delta = -2916 < 0$$

$g(0) = -810$ g-intercept.

