

Rational Polynomials: Graphing and Asymptotes

Find the intercepts, if there are any.

Step 1: Set the numerator to 0 to solve for horizontal intercepts.

Step 2: Set the x to 0 to solve for vertical intercept.

Step 3: Set the denominator to 0 to solve for vertical asymptotes.

Step 4: Perform a long division to find the quotient which specifies the oblique asymptote.


Note: Blue curve the actual Rational function.

Red and Gold asymptotes.

Example: Horizontal Asymptote

$$\frac{4x-2}{5x-1}$$


$$\begin{array}{r} + \left(\frac{4}{5} \right) \\ \hline 5x - 1 \quad (4)x \quad + (-2) \\ \quad (4x) \quad + \left(-\frac{4}{5} \right) \\ \quad \quad + \left(-\frac{6}{5} \right) \end{array}$$



Example: Oblique Linear Asymptote

$$\frac{(x-3)(x+2)}{x-2}$$

$$\begin{array}{r} + \left(x \right) \quad + \left(1 \right) \\ \hline x - 2 \quad (1)x^2 \quad + (-1)x \quad + (-6) \\ \quad (x^2) \quad + \left(-2x \right) \\ \quad \quad + (1)x \quad + (-6) \\ \quad \quad + (x) \quad + \left(-2 \right) \\ \quad \quad \quad + \left(-4 \right) \end{array}$$



Example: Multiple Vertical Asymptotes

$$\frac{x+2}{(x-1)(x+1)}$$

$$\begin{array}{r} + \left(0 \right) \\ \hline (x) \quad + \left(2 \right) \end{array}$$

