

Rational Polynomials: Graphing and Asymptotes

Find the intercepts, if there are any.

Step 1: Set the numerator to 0 to solve for horizontal intercepts.

Step 2: Set the x to 0 to solve for vertical intercept.

Step 3: Set the denominator to 0 to solve for vertical asymptotes.


Step 4: Perform a long division to find the quotient which specifies the oblique asymptote.

Note: Blue curve the actual Rational function.
Red and Gold asymptotes.

Example: Horizontal Asymptote

$$\frac{3x-5}{2x-4}$$


$$\begin{array}{r} + \left(\frac{3}{2}\right) \\ \hline 2x-4 \quad (3)x \quad + (-5) \\ \quad (3x) \quad + (-6) \\ \quad + (1) \end{array}$$



Example: Oblique Linear Asymptote

$$\frac{(x-1)(x+3)}{x-2}$$

$$\begin{array}{r} + (x) \quad + (4) \\ \hline x-2 \quad (1)x^2 + (2)x \quad + (-3) \\ \quad (x^2) \quad + (-2x) \\ \quad + (4)x \quad + (-3) \\ \quad + (4x) \quad + (-8) \\ \quad + (5) \end{array}$$



Example: Multiple Vertical Asymptotes

$$\frac{x+2}{(x-1)(x+3)}$$

$$\begin{array}{r} + (0) \\ \hline (x) \quad + (2) \end{array}$$

