

# Intercepts of the Quadratic

Given a quadratic  $y(g) = a g^2 + b g + c$  compute its discriminant  $\Delta$ :

$$\Delta = \sqrt{b^2 - 4ac}$$

**Case1:  $\Delta > 0$**

$g_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$  computes the g-intercepts of multiplicity 1.  
 $y(0) = c$  computes the single y-intercept.


**Example 1.**

$y(g) = 25 - g^2$  compute its discriminant  $\Delta$ :

$$\Delta = 100 > 0$$

$$g_{1,2} = 5, -5$$

$y(0) = 25$  y-intercept.



**Case2:  $\Delta = 0$**

$g_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-b \pm 0}{2a} = \frac{-b}{2a}$  single g-intercept of multiplicity 2.


**Example 2.**

$y(g) = -2g^2 + 24g - 72$  compute its discriminant  $\Delta$ :

$$\Delta = 0$$

$$g_{1,2} = 6, 6$$

$y(0) = -72$  y-intercept.



**Case3:  $\Delta < 0$**

$\sqrt{b^2 - 4ac}$  has no value in Real Numbers. Therefore there are no g-intercepts.

However there is a y-intercept.

**Example 3.**

$y(g) = -4g^2 + 80g - 500$  compute its discriminant  $\Delta$ :

$$\Delta = -1600 < 0$$

$y(0) = -500$  y-intercept.

