



Antelope Valley Medical Center

Antelope Valley Medical Center

1600 West Avenue J

Lancaster, CA 93534

661-949-5115

Discharge Instructions (Patient)

Name: HERNANDEZ, NICOLAS **Current Date:** 07/12/26 03:04:18

DOB: [REDACTED] **MRN:** [REDACTED] 80 **FIN:** [REDACTED] 04

Diagnosis: 1:Anxiety; 2:Benzodiazepine dependence; 3:Opioid dependence

Visit Date: 07/12/26 01:32:08

Address: [REDACTED] CA 9[REDACTED]

Phone:

Primary Care Provider: Bartz-Altadonna, Community Health Ctr-Palmdale AVH **Phone:**
(661)874-4050

Emergency Department Providers:

Provider Name	Role
Meredith Graham	ED APP

Antelope Valley Hospital would like to thank you for allowing us to assist you with your healthcare needs. The following instructions include patient education materials and information regarding your injury/illness.

The examination and treatment you have received today has been on an emergency basis only and has not been intended to be a substitute for complete medical care. Since it is impossible to recognize and treat all elements of injury or illness in a single visit, for your protection in preventing possible complications, if you are not clearly improving in the next 12 hours, you should be seen for a re-check at this emergency room or at your own doctor's office immediately.

IF YOU FEEL YOUR CONDITION HAS AGAIN BECOME AN EMERGENCY, RETURN TO THIS DEPARTMENT AT ONCE, DAY OR NIGHT.

IF YOU EXPERIENCE SHORTNESS OF BREATH OR CHEST PAIN. PLEASE CALL 9-1-1.

Continue your current medications as you have been, unless instructed otherwise by your Provider today. If any new medication were prescribed today, take them as directed.

If you have been prescribed medication(s) as part of your care plan, read the medication package insert carefully and completely AND ask your pharmacist for complete information about medication use, side effects, precautions, and interactions with other medications that you currently take.

Make sure to bring any attached imaging reports and lab results to your doctor. As we discussed, these must be reviewed by your doctor to decide what additional testing or treatment may be needed to address any abnormal results.

Patient Instructions:

Please return to the ER in 24-48 hour for a recheck of your symptoms.

Please return sooner for any changes or concerns.

If you can follow up with your PMD instead that is OK.

Please consider these lifestyle changes:

Please eliminate processed foods and focus on eating vegetables and meat (such as red meat, chicken, turkey and fish- red meat has the most vitamins and nutrients)

**DO NOT EAT ANY PROCESSED CARBOHYDRATES OR SUGARS
DO NOT EAT ANYTHING YOU CANNOT PRONOUNCE (CHEMICALS)
DO NOT EAT ANYTHING YOU DO NOT MAKE YOURSELF
DO NOT EAT ANYTHING WITH AN INGREDIENT LIST**

Please increase your activity: example walking 30 minutes a day

Please eliminate monosodium glutamate (MSG)

Please do not eat any type of Hot Chip or Hot Cheetos or Takis.

Please eliminate gluten and any other inflaming food products (please do not eat regular chips or hot chips of any kind)

Please do not drink soda or juice or prepackaged drinks, Please aim to drink WATER- 60-90oz a day

Please do not use cigarettes, vapes, marijuana, methamphetamines, cocaine, heroin or other illicit drugs.

Please do not drink Alcohol

Please avoid seed oils (corn, soy, rapeseed, canola, sunflower, high oleic oils)

Taking Opioids Safely: Key Information for Patients

Before You Start Opioid Treatment

As your healthcare provider, I want to ensure you understand the important information about taking opioid medications. Here are the key points:

1. Risk of Addiction and Overdose

- o Opioid medications carry a risk of addiction, even when taken as prescribed.
- o Overdose can happen, especially if opioids are misused or combined with other substances.

2. Increased Risks for Certain Individuals

- o If you have mental health conditions or a history of substance use, your risk of opioid addiction may be higher. Share your medical history to help us manage your care safely.

3. Avoid Dangerous Interactions

- o Combining opioids with alcohol, benzodiazepines, or other sedatives can cause severe complications, including overdose or death.
- o Always inform us about all medications or substances you are taking.

Additional Safety Tips

- Take your medication exactly as prescribed. Do not take extra doses or stop suddenly without consulting your doctor.
- Keep your medications in a secure place and never share them with others.
- Avoid driving or operating machinery if you feel drowsy or impaired.

For More Information

Visit the FDA webpage for detailed resources:

<https://www.fda.gov/drugs/information-drug-class/opioid-medications>

Your safety is our top priority. Please don't hesitate to ask questions or share any concerns during your visit and with your healthcare providers after your visit.
???????

Follow-up Instructions:

With:

Address:

When:

Follow-up in Emergency
Department

In 1 day, only if
needed

With:

Address:

When:

Follow up with Primary Care
Provider

In 1 day

Allergies: No Known Medication Allergies

Medication Information:

If you have been instructed to stop taking a medication, please ensure you also follow up with this information to your Primary Care Physician. Unless otherwise noted, please continue to take medications as prescribed prior to your Emergency Room visit. Any specific questions regarding your chronic medications and dosages should be discussed with your physician(s) and pharmacist.

New Medications and Prescriptions

 **CVS/pharmacy #9785, 2006 W Avenue J Lancaster, CA 935365913, (661) 945 - 2729**

diazepam (diazepam 10 mg oral tablet) 1 tab Oral (given by mouth) 2 times a day as needed as needed for anxiety for 3 Days. Refills: 0.

Last Dose: _____ Next Dose: _____

Medications That Were Changed - Changes Shown Below

CVS/pharmacy #9785, 2006 W Avenue J Lancaster, CA 935365913, (661) 945 - 2729

Current **buprenorphine-naloxone (Suboxone 4 mg-1 mg sublingual film)** 1 Each Sublingual (dissolve under the tongue) 2 times a day for 10 Days. Refills: 0., XDEA XG5094444

Last Dose: _____ Next Dose: _____

Other Medications

Current **buprenorphine-naloxone (Suboxone 4 mg-1 mg sublingual film)** 1 Each Sublingual (dissolve under the tongue) 2 times a day for 10 Days. Refills: 0., XDEA XG5094444

Last Dose: _____ Next Dose: _____

Patient Education:

Managing Anxiety, Adult

After being diagnosed with anxiety, you may be relieved to know why you have felt or behaved a certain way. You may also feel overwhelmed about the treatment ahead and what it will mean for your life. With care and support, you can manage your anxiety.

How to manage lifestyle changes