

Worksheet on

Toric Geometry

Written by Louan Mol

Last update on June 27, 2022.

Contents

1	What are toric varieties ?	1
1.1	Cones	1
1.2	Toric varieties	2
2	Constructing toric affine varieties	2
2.1	Toric variety of a cones	2
2.2	Toric variety of a fan	4
2.3	Fan from a toric variety	6
2.4	More general constructions	6
3	Relation between fan varieties, cone varieties and local coordinates	6
4	Calabi-Yau toric varieties	7
5	Toric diagrams and p-q webs	7
6	Singularities	8
7	Important examples	8
7.1	The conifold	8
7.2	The suspended pinch point	8
7.3	del Pezzo surfaces	8

1 | What are toric varieties ?

The interest of toric varieties lies in the fact that all its defining data can be encoded in a simple auxiliary object called a fan. This data is purely combinatoric (discrete quantities) so complicated geometric problems are often reduced to simpler combinatorics problems.

1.1 | Cones

Let us consider a lattice $N = \mathbb{Z}^m$ and $N_{\mathbb{R}} = N \otimes \mathbb{R}$ the real vector space that we get by allowing real coefficients. Given v_1, \dots, v_n of the lattice N , a *cone*¹ $\sigma \subset N_{\mathbb{R}}$ is a the subset of the vector space $N_{\mathbb{R}}$ containing all points that are positive linear combination of the vectors v_1, \dots, v_n , that is

$$\sigma = \left\{ \sum_{i=1}^n a_i v_i \mid a_i \geq 0, a_i \in \mathbb{R} \right\} \quad (1.1)$$

¹More precisely, a strongly convex rational polyhedral cone.

and such that $\sigma \cap (-\sigma) = \{0\}$. The last condition is the strong convexity condition and it imposes that our cone has to be “acute”. The dimension of a cone is the dimension of $\text{Span}(\sigma)$.

The dual lattice N^* of N is the set of linear functionals on L which take integer values on each point of N :

$$N^* = \{f \in (\text{Span}(L))^* \mid \forall x \in N, f(x) \in \mathbb{Z}\}. \quad (1.2)$$

We denote by M the dual lattice of N and by $M_{\mathbb{R}}$ the vector space we get from it by allowing real coefficients. For a given cone σ in $N_{\mathbb{R}}$, we can define the *dual cone* σ^\vee in $M_{\mathbb{R}}$ as

$$\sigma^\vee = \{m \in M_{\mathbb{R}} \mid \langle u, m \rangle \geq 0, \forall u \in \sigma\}. \quad (1.3)$$

Let us denote by H_m the hyperplane given by the dual lattice point $m \in \sigma^\vee$: $H_m = \{v \in N_{\mathbb{R}} \mid \langle v, m \rangle = 0\}$ and the closed half-space $H_m^+ = \{v \in N_{\mathbb{R}} \mid \langle v, m \rangle \geq 0\}$. We then say that a hyperplane *supports* a cone if the closed half-space of the hyperplane completely contains the cone. A *face* of a cone is the intersection of the cone with a supporting hyperplane. Put differently, a convex subset τ is a face of σ if and only whenever $u, v \in \sigma$ satisfy $u + v \in \tau$ then $u \in \tau$ and $v \in \tau$. Note that a hyper plane H_m supports σ if and only if $m \in \sigma^\vee$.

1.2 | Toric varieties

An *algebraic group* is a group that is also an algebraic variety and such the product and inversion are regular maps on the variety. Any product of \mathbb{C}^* is an algebraic group that we call *algebraic tori*. An affine variety $X \subset \mathbb{C}^n$ is a *affine toric variety* if it contains the algebraic torus $\mathbb{T} = (\mathbb{C}^*)^n$ as a dense open subset such that the action of \mathbb{T} on itself extends to an action $\mathbb{T} \times X \rightarrow X$ on X .

Example. Let us enumerate some examples of affine toric varieties.

- $(\mathbb{C}^*)^n$ and \mathbb{C}^n are naturally provided with an embedding and an action of the torus and are toric varieties
- $V = Z(x^3 - y^2)$ with torus embedding

$$\left(\begin{array}{ccc} \mathbb{C}^* & \hookrightarrow & V \\ t & \longmapsto & (t^2, t^3) \end{array} \right). \quad (1.4)$$

and the action $t \cdot (u, v) \mapsto (t^2 u, t^3 v)$.

2 | Constructing toric affine varieties

2.1 | Toric variety of a cones

A *semigroup* S is a set with an internal associative $+$ operation and a neutral element 0. It differs from a group in that elements need not have an inverse. A semigroup is *affine* if it can be embedded as a subsemigroup in a lattice \mathbb{Z}^m (*integral*) and if there exists a finite set \mathcal{A} such that $S = \mathbb{N}\mathcal{A}$ (*finitely generated*).

What is interesting is that affine semigroups allows us to construct affine varieties. For this, we need to introduce the notion of *semigroup algebra* $\mathbb{C}[S]$ associated to any semi group S . It is the algebra generated by elements χ^u indexed by elements $u \in S$. The semigroup operation $+$ induces a multiplication operation for the χ^u in $\mathbb{C}[S]$ as $\chi^u \cdot \chi^v = \chi^{u+v}$. For instance, the semigroup algebra of $S = \mathbb{N}^n$ is simply $\mathbb{C}[x_1, \dots, x_n]$ and the semigroup algebra of $S = \{2, 3, \dots\} \subset \mathbb{N}$ is $\mathbb{C}[x, y]/I(x^3 - y^2)$. The important result is now that if S is an affine semigroup then $\text{Spec}(\mathbb{C}[S])$ is an affine variety.

Given a cone σ and its dual cone σ^\vee , $S_\sigma = \sigma^\vee \cap M$ is finitely generated and hence an affine semigroup. In this way, we may obtain an affine variety from a cone as $U_\sigma = \text{Spec}(\mathbb{C}[S_\sigma])$, this the *variety if the cone* σ . An important result is that if $K[V]$ is the coordinate ring of an affine variety V , then $V = \text{Spec}(K[V])$. From this, we see that $\mathbb{C}[S_\sigma]$ is exactly the coordinate ring of the variety U_σ we are looking for.

We have seen how to get an affine variety U_σ from a cone σ . This variety is in fact toric. If n denotes the rank of the lattice $\mathbb{Z}S_\sigma$, then there is torus action $\mathbb{T} = (\mathbb{C}^*)^n$ acting on U_σ . The torus that it contained is called the *torus corresponding to a lattice* $N = \mathbb{Z}^m$ and is $\mathbb{Z}_N = (\mathbb{C}^*)^n$, so that the rank of the lattice equals the dimension of the torus:

$$T_N = N \otimes_{\mathbb{Z}} \mathbb{C}^* = \text{Hom}_{\mathbb{Z}}(M, \mathbb{C}^*). \quad (2.1)$$

To torus acts on U_σ as $t \cdot \gamma : S_\sigma \rightarrow \mathbb{C} : m \mapsto \chi^m(t)\gamma(m)$, for all $t \in T_N$ and $\gamma \in \sigma$ (so $\gamma : S_\sigma \rightarrow \mathbb{C}$). To summarize, the steps to extract the m -complex dimensional toric variety from a given cone σ in $N_{\mathbb{R}}$ are the following:

1. find the dual cone σ^\vee
2. find the intersection $S_\sigma = \sigma^\vee \cap \mathbb{Z}^m$, it is a finitely generated semigroup. Find a minimal set of generating vectors $\{w_1, \dots, w_r\}$
3. find the polynomial ring $\mathbb{C}[S_\sigma]$ by exponentiating the coordinates of S_σ
4. find the relations between the elements of $\mathbb{C}[S_\sigma]$ corresponding to the generating vectors w_i . Such a relation looks like

$$\sum_{i \in I} m_i w_i = \sum_{j \in J} m_j w_j, \quad m_i, m_j \in \mathbb{N} \Rightarrow p(x) = \prod_{i \in I} x^{m_i} x_i - \prod_{j \in J} x^{m_j} x_j \quad (2.2)$$

where $I \cup J = \{1, \dots, r\}$ and $I \cap J = \emptyset$. If $\{p_1, p_2, \dots\}$ is the set of all those relations, then $I(\{p_1, p_2, \dots\})$ is a prime ideal and $\mathbb{C}[S_\sigma] = \mathbb{C}[x_1, \dots, x_r]/I(p_1, p_2, \dots)$

5. $\mathbb{C}[S_\sigma]$ is the coordinate ring of desired the toric variety. We can recover the variety explicitly as $\text{Spec}(\mathbb{C}[S_\sigma])$. It might be easier to use the fact that $\text{Spec}[K[V]] = V$, so $\mathbb{C}[S_\sigma]$ is actually the coordinate ring of U_σ and $U_\sigma = Z(\{p_1, p_2, \dots\})$.

The vtoric variety U_σ has the same dimension than σ .

Example. If we consider the cone σ generated by e_2 and $2e_1 - e_2$ in \mathbb{Z}^2 (where $\{e_1, e_2\}$ is the canonical basis of \mathbb{Z}^2), then the dual cone σ^\vee is generated by e_1 and $e_1 + 2e_2$. We therefore have $S_\sigma = \sigma^\vee \cap \mathbb{Z}^2$. This subset of \mathbb{Z}^2 is spanned by $e_1, e_1 + e_2$ and $e_1 + 2e_2$:

$$S_\sigma = \text{Span}(\{(1, 0), (1, 1), (1, 2)\}). \quad (2.3)$$

Note that, even though it is a 2-dimensional cone, i.e. the intersection of this cone and the lattice actually need three vectors to be completely generated. This is often the case when dealing with lattices. By exponentiating we get the semigroup algebra $\mathbb{C}[S_\sigma] = \mathbb{C}[x, xy, xy^2]$. If we denote $u = x, v = xy$ and $w = xy^2$, relation between those three variables is $uw = v^2$, so $\mathbb{C}[S_\sigma] = \mathbb{C}[u, v, w]/I(v^2 - uw)$. It only remains to compute the spectrum, and we find that

$$\text{Spec}(\mathbb{C}[S_\sigma]) = \mathbb{C}^2/\mathbb{Z}^2. \quad (2.4)$$

In conclusion the toric variety associated to the cone σ is the abelian orbifold $\mathbb{C}^2/\mathbb{Z}^2$.




Figure 1: Representations of the cone and dual cone in the lattice \mathbb{Z}^2 .

The previous example illustrate a more genral result:

Proposition. All abelian orbifolds are toric varities.

Note that this result is not that surprising considering the general expression (2.6).

Example. Let us start with $\sigma = \text{Cone}(\{e_1, 2e_1 + e_2\})$. We then get $\sigma^\vee = \text{Cone}(\{e_2, e_1 - 2e_2\})$ so $S_\sigma = \text{Span}_{\mathbb{Z}^+}(\{e_2, e_1 - 2e_2\})$ and $\mathbb{C}[S_\sigma] = \mathbb{C}[y, xy^{-2}] = \mathbb{C}[u, v]$.

Now what happens with faces? An inclusion of face $\tau \subset \sigma$ gives an inclusions of the semigroups $S_\sigma \subset S_\tau$ and $\mathbb{C}[S_\sigma] \subset \mathbb{C}[S_\tau]$. This induces a morphism $U_\tau \rightarrow U_\sigma$ of affine toric varieties and it happens the this morphism is an open embedding.

Example. If $\sigma = \text{Cone}(e_1, e_2)$ then we find $U_\sigma = \mathbb{C}^2$. There are three faces: $\rho_1 = \text{Cone}(e_1)$, $\rho_2 = \text{Cone}(e_2)$ and the origin 0. These faces can be described by hyperplane H_{e_2} , H_{e_1} and $H_{e_1+e_2}$ respectively, so we get

$$\mathbb{C}[S_{\rho_1}] = \mathbb{C}[x, y, y^{-1}], \mathbb{C}[S_{\rho_2}] = \mathbb{C}[x, x^{-1}, y], \mathbb{C}[S_0] = \mathbb{C}[x, y, x^{-1}, y^{-1}]. \quad (2.5)$$

They define the varieties $U_{\rho_1} = \mathbb{C}^* \times \mathbb{C}$, $U_{\rho_2} = \mathbb{C} \times \mathbb{C}^*$ and $(\mathbb{C}^*)^2$, which are all open subvarieties of \mathbb{C}^2 .

2.2 | Toric variety of a fan

A *fan* is a collection Δ of cones in $N_{\mathbb{R}}$ such that each face of a cone is also a cone and the intersection of two cones is a face of each. Those two requirement imply in particular that the intersection of two cones in a fan is again a cone in the fan. Note that to any cone corresponds a fan containing the cone itself and all its faces.

For a cone σ in $N_{\mathbb{R}}$ the variety U_σ contains the torus T_N so fan in $N_{\mathbb{R}}$ produces a collection of varieties that all contain th same torus T_N . Gluing together the affine varities we obtain X_Δ , it is clear that T_N is an open subset of X_Δ and that T_N acts on X_Δ . This is called the *toric variety of a fan*. For the toric varitey of a fan, each cone in the fan turns out to be determines an orbit $O(\sigma)$ of the torus.

Generally speaking, an m -dimensional algebraic variety of a fan can be obtained as as particular holomorphic quotient of \mathbb{C}^n . If $\mathbb{Z}_\Delta \subset \mathbb{C}^n$ a set of points and $G \cong (\mathbb{C}^*)^{n-m} \times \Gamma$ is the group formed by the algebraic torus and an abelian discrete group Γ , then

$$X_\Delta = \frac{\{\mathbb{C}^n \setminus \mathbb{Z}_\Delta\}}{G}. \quad (2.6)$$

Let us explain this construction.

If Δ is an m -dimensional fan, we denote by $\Delta(j)$ ($j \leq n$) the collection of j -dimensional cones in Δ . The set $\Delta(1) \subset \Delta$ is then the collection of all one-dimensional cones in Δ , i.e. the set of edges. To each cone $\sigma \in \Delta(1)$ is associated a unique vector $v_\sigma \in N$ that generates the sublattice $\sigma \cap N$. This vector is called the *primitive generator* of σ . Let v_1, \dots, v_n be the primitive generators of of each $\sigma \in \Delta(1)$. Note that we always have $n \geq m$ since Δ is m -dimensional. Each vector v_i has component v_i^k ($i = 1, \dots, n, k = 1, \dots, m$) so we can construct an $m \times n$ matrix by putting the vectors in columns:

$$V = \begin{bmatrix} v_1^1 & \cdots & v_n^1 \\ \vdots & & \vdots \\ v_1^m & \cdots & v_n^m \end{bmatrix} \quad (2.7)$$

This defines a linear map from \mathbb{C}^n to \mathbb{C}^m whose kernel is \mathbb{C}^{n-m} . Let Q^a ($a = 1, \dots, n-m$) be vectors that generate this kernel, then we must have

$$VQ^a = 0 \Leftrightarrow \sum_{i=1}^n v_i^k Q_i^a = 0, \quad k = 1, \dots, m \quad (2.8)$$

for all $a = 1, \dots, n - m$. We now put that on the side for a moment. To each edge $\sigma \in \Delta(1)$ with associate a complex coordinate x_σ and consider the new map

$$\phi : \begin{pmatrix} \mathbb{C}^n & \longrightarrow & \mathbb{C}^m \\ (z_1, \dots, z_n) & \longmapsto & (\prod_{i=1}^n z_i^{v_i^1}, \dots, \prod_{i=1}^n z_i^{v_i^m}) \end{pmatrix}. \quad (2.9)$$

It is clear that the kernel of ϕ is $\tilde{G} \equiv \text{Ker}(\phi) = (\mathbb{C}^*)^{n-m}$. One can see that this kernel naturally acts on \mathbb{C}^n as

$$\begin{pmatrix} (\mathbb{C}^*)^a \times \mathbb{C}^n & \longrightarrow & \mathbb{C}^n \\ (\lambda, z_1, \dots, z_n) & \longmapsto & (\lambda^{Q_1^a} z_1, \dots, \lambda^{Q_n^a} z_n) \end{pmatrix} \quad (2.10)$$

where Q_i^a are the component of the vectors Q^a defined above. We defined the action of $\tilde{G} = (\mathbb{C}^*)^{n-m}$ on \mathbb{C}^n for each factor separately for simplicity.

The n vectors v_i generate over \mathbb{Z} a sublattice of N that we denote N' . The discrete group obtained by taking the quotient of N by this sublattice

$$\Gamma = N/N' \quad (2.11)$$

is a subgroup of G and taking the quotient by Γ gives rise to the orbifold singularities.

The last piece of data in the construction is the zero set Z_Δ . Let us denote by \mathcal{S} any subset of $\Delta(1)$ that does not generate a cone in Δ and $V(\mathcal{S}) \subset \mathbb{C}^n$ the linear subspace defined by setting $x_\sigma = 0$ for all $\sigma \in \mathcal{S}$, i.e. the hyperplane generated by the edges that does not generate a cone in Δ . We denote Z_Δ the union of all those hyperplanes, i.e. of all the $V(\mathcal{S})$. Our toric variety is defined as a quotient of $\mathbb{C}^n \setminus Z_\Delta$.

To summarize, we can construct affine toric variety X_Δ from the m -fan Δ by following these steps:

- find the primitive generators v_1, \dots, v_n of the 1-cones of Δ
- find the $n - m$ vectors Q^a such that $\sum_i v_i^k Q_i^a = 0$
- compute $\Gamma = N/N'$
- find the subsets of $\{v_1, \dots, v_n\}$ that don't generate in Δ and compute Z_Δ
- the toric affine variety is $X_\Delta = (\mathbb{C}^n \setminus Z_\Delta) / ((\mathbb{C}^*)^{n-m} \times \Gamma)$.

Example. If we consider the 2-dimensional fan $\Delta = \text{Fan}(e_1, e_2, -e_1 - e_2)$ in $N = \mathbb{Z}^3$ (so $m = 2$), we have three one-dimensional cones generated by $v_1 = e_1, v_2 = e_2$ and $v_3 = -e_1 - e_2$ (so $n = 3$) and the component matrix is

$$\begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & -1 \end{bmatrix}. \quad (2.12)$$

We only have one vector Q^a to find and since it must satisfy the relation (2.8), i.e. in this case $Q_1 v_1 + Q_2 v_2 + Q_3 v_3 = 0$, it implies that $Q = (1, 1, 1)$. The group action of $\tilde{G} = \mathbb{C}^*$ on the \mathbb{C}^3 is then

$$(z_1, z_2, z_3) \mapsto (\lambda z_1, \lambda z_2, \lambda z_3). \quad (2.13)$$

Since v_1, v_2 and v_3 generate all of \mathbb{Z}^2 , $\Gamma = \{0\}$ is the trivial group and $G = \tilde{G} \times \Gamma \cong \tilde{G} = \mathbb{C}^*$. There is no subset of $\{v_1, v_2, v_3\}$ that generates a cone which is not in Δ (because we defined Δ as the fan generated by those vectors in the first place) therefore $Z_\Delta = \{(0, 0, 0)\}$. Finally, the affine toric variety of Δ is

$$X_\Delta = \frac{\mathbb{C}^3 \setminus Z_\Delta}{G} = \frac{(\mathbb{C}^*)^3}{\mathbb{C}^*} = \mathbb{CP}^2. \quad (2.14)$$

Proposition. A toric variety X_Δ is compact if and only if its fan Δ span the $N_{\mathbb{R}}$.

2.3 | Fan from a toric variety

2.4 | More general constructions

In general, an affine variety can be defined by an ideal in $\mathbb{C}[x_1, \dots, x_n]$. In the same way, a toric affine variety can be defined by a *toric ideal* (or *binomial ideal*), i.e. an ideal in $\mathbb{C}[x_1, \dots, x_n]$ generated by binomials (polynomials with precisely two non-zero coefficients).

Theorem. Let V be an affine variety. The following are equivalent:

- V is toric,
- $V = \text{Spec}(\mathbb{C}[S])$ for an affine semigroup S ,
- $V = V(I)$ for a toric ideal.

This is a very powerful theorem since it states that an affine variety is toric if and only if the generating polynomials of its ideal are binomials.

3 | Relation between fan varieties, cone varieties and local coordinates

Given an m -dimensional cone spanned by n vectors v_1, \dots, v_n , we want to find the local coordinates associated to it. Since the toric variety can be expressed as the quotient by G , the local coordinates should be G -invariant polynomials, that is polynomials $x = z_1^{n_1} \dots z_n^{n_n}$ such that

$$x \mapsto \lambda^{\sum_i Q_i^n n_i} x = x \quad (3.1)$$

under G . But remember that (2.8) so we can take $n_i = \langle w, v_i \rangle$ for any $w \in M$. We found that the local coordinates are in one-to-one correspondence with the dual lattice M .

Given a fan and its fan toric variety, we can associate a toric affine variety to each top-dimensional cone. These cone toric varieties will be affine patches of the fan variety. The transition functions between these patches are also encoded in the initial fan. Let us explain those statements in greater details. If $\sigma_i \in \Delta$ are the top-dimensional cones of a given fan Δ , we can construct their toric variety U_{σ_i} by following the procedure that we presented above. The toric varieties U_{σ_i} act as affine patches and can be patched together to form X_Δ . Now suppose that τ is a face of both σ_i and σ_j , one can show that

$$\sigma_{i,j}^\vee \subset \tau^\vee \Rightarrow \mathbb{C}[\sigma_{i,j}^\vee \cap M] \subset \mathbb{C}[\tau^\vee \cap M] \Rightarrow U_\tau \subset U_{\sigma_i} \cap U_{\sigma_j} \quad (3.2)$$

so the affine set associated to the face τ is in the intersection of the affine sets of the cones. The relation between local coordinates $x^{(i)}$ of U_{σ_i} and $x^{(j)}$ of U_{σ_j} can then be read from the relations between the generators of $\sigma_{i,j}^\vee \cap M$ and $\sigma_{i,j}^\vee \cap M$:

$$\sum_{l=1}^{r_i} q_l w_l^{(i)} = \sum_{k=1}^{r_j} q_k w_k^{(j)}, \quad q_l, q_k \in \mathbb{Z} \Rightarrow \prod_{l=1}^{r_i} (x_l^{(i)})^{q_l} = \prod_{k=1}^{r_j} (x_k^{(j)})^{q_k}. \quad (3.3)$$

We see that those transition functions are always rational functions. This is a crucial property of toric varieties.

Example. Going back to the fan of \mathbb{CP}^2 , we see that there are three top-dimensional cones σ_1, σ_2 and σ_3 . For each we have $U_{\sigma_i} = \mathbb{C}^2$. Applying (3.3) we find that the transition functions between the coordinates (x_1, x_2) of U_{σ_1} and (y_1, y_2) of U_{σ_2} are

$$x_1 = \frac{y_1}{y_2}, \quad x_2 = \frac{1}{y_2}. \quad (3.4)$$

4 | Calabi-Yau toric varieties

As motivated in the beginning, we are mainly interested in Calabi-Yau affine varieties, so Calabi-Yau affine toric varieties in this case. A very convenient property of the toric varieties is that the CY condition is translated into a simple condition on the combinatoric data of the variety.

Recall that a divisor of an affine variety is a linear combination of codimension-one irreducible subvarieties. A *toric divisor* is a divisor invariant under the action of G . Using the homogeneous coordinates (z_i) , we can easily construct G -invariant subvarieties. The simple algebraic subsets

$$\{(z_1, \dots, z_n) | z_i = 0 \forall i \in I \subset \{1, \dots, n\}\} \quad (4.1)$$

are G -invariant so the subvarieties

$$D_i = \{z_i = 0\} \cap X_\Delta \quad (4.2)$$

are toric divisors. Even stronger, one can actually show that they generate the full group of divisors of X_Δ and that if X_Δ is smooth with canonical bundle K_X , we have

$$K_X = \mathcal{O}\left(-\sum_{i=1}^n D_i\right). \quad (4.3)$$

This important result allows us to state the Calabi-Yau condition (triviality of the canonical bundle) in a very simple way. The only thing left is to see that any G -invariant function is a section of the trivial bundle and that $K_X = \mathcal{O}(-\sum_{i=1}^n D_i)$ is trivial if and only if

$$G : z_1 \dots z_n \mapsto \lambda^{\sum_{i=1}^n Q_i^a} z_1 \dots z_n \Leftrightarrow \sum_{i=1}^n Q_i^a = 0. \quad (4.4)$$

This last condition is equivalent to the existence of a vector $w \in M$ such that $\langle w, v_i \rangle = 1$ for all v_i in the fan. Finally, we get the following criteria:

Proposition. The toric variety X_Δ is CY if and only if all the vectors v_i in Δ end on the same hyperplane in N , which happens if and only if $\sum_{i=1}^n Q_i^a = 0$ for $a = 1, \dots, n - m$.

Following from the proposition on compactness of toric varieties that we gave before, this implies that toric CY varieties are necessarily non-compact.

X_Δ	$\dim_{\mathbb{C}}$	generating polynomial	compact	CY	Δ
\mathbb{C}^2	2		no	yes	$\text{Fan}(e_1, e_2)$
$\mathbb{C}^2/\mathbb{Z}_k$	2	$uv - w^2$	no	yes	$\text{Fan}(e_2, ke_1 - e_2)$
$\mathbb{C}^3/\mathbb{Z}_k$	3		no	yes	$\text{Fan}(e_2, ke_1 - e_2, e_3)$
\mathcal{C}_0	3	$x_1x_2 - x_3x_4$	no	yes	$\text{Fan}(e_3, e_1 + e_3, e_1 + e_2 + e_3, e_2 + e_3)$
SSP					

Table 1: Useful list of toric varieties.

5 | Toric diagrams and p-q webs

For CY toric varieties, the combinatoric information encoded in the fan can be expressed in terms of a reduced lattice of one less dimension. This is especially convenient for us since we are mainly concerned about toric CY threefolds, which can therefore be fully encoded in the 2-dimensional lattice. Indeed, instead of drawing a 3-dimensional fan, we can simply project it on the special plane defined by $\langle w, v_i \rangle = 1$, and we get the *toric diagram*.

We can also write the dual of the toric diagram by replacing each line with an orthogonal line. This is called the *peq-web*. They have a nice physical interpretation as webs intersecting 5-banes.

6 | Singularities

A cone is said to be *smooth* if it is generated by part of a lattice basis.

Theorem. A cone σ is smooth if and only if U_σ is smooth.

Example. The affine toric variety $Z(x^2 - y^3)$ discussed above is not smooth.

Proposition. The variety X_Δ is non-singular if and only if Δ is a smooth fan.

A toric variety is nonsingular if its cones of maximal dimension are generated by a basis of the lattice. This implies that every toric variety has a resolution of singularities given by another toric variety, which can be constructed by subdividing the maximal cones into cones of non-singular toric varieties.

7 | Important examples

7.1 | The conifold

The singular conifold \mathcal{C}_0 can be viewed as the toric variety of the fan $\Delta = \text{Fan}(e_3, e_1 + e_3, e_1 + e_2 + e_3, e_2 + e_3)$ in \mathbb{Z}^3 . This fan contains 10 cones on total. Denoting the four generating vectors as v_1, v_2, v_3 and v_4 , we find that the only charge vector Q must satisfy $Q_1 v_1 + Q_2 v_2 + Q_3 v_3 + Q_4 v_4 = 0$ which gives $Q = (1, -1, 1, -1)$ so $\tilde{G} = \mathbb{C}^*$ acts as

$$(z_1, z_2, z_3, z_4) \mapsto (\lambda z_1, \lambda^{-1} z_2 \lambda z_3, \lambda^{-1} z_4). \quad (7.1)$$

We also find that $\Gamma = \{0\}$ is the trivial group and that $Z_\Delta = Z(z_1, z_3) \cup Z(z_2, z_4)$. Finally,

$$\mathcal{C}_0 = \frac{\mathbb{C}^3 \setminus Z_\Delta}{\mathbb{C}^*}. \quad (7.2)$$

Note that, by definition we also have $\mathcal{C}_0 = Z(x_1 x_4 - x_2 x_3)$ so it is generated by a binomial and is therefore expected to be toric.



Figure 2: Toric diagram and pq-web of the conifold.

7.2 | The suspended pinch point

7.3 | del Pezzo surfaces




Figure 3: Toric diagram of the SPP.