

Time-frequency representation of autoionization dynamics in helium

D. Busto¹, L. Barreau², M. Isinger¹, M. Turconi², C. Alexandridi², A. Harth¹, S. Zhong¹, R. J. Squibb³, D. Kroon¹, S. Plogmaker¹, M. Miranda¹, Á. Jiménez-Galán⁴, L. Argenti⁵, C. L. Arnold¹, R. Feifel³, F. Martín^{4,6,7}, M. Gisselbrecht¹, A. L'Huillier¹, P. Salieres²

¹Department of Physics, Lund University, P. O. Box 118, SE-22100 Lund, Sweden

²LIDYL, CEA, CNRS, Université Paris-Saclay, CEA Saclay, 91191 Gif-Sur-Yvette, France

³Department of Physics, University of Gothenburg, Origovägen 6B, SE-41296 Gothenburg, Sweden

⁴Departamento de Química, Módulo 13, Universidad Autónoma de Madrid, 28049 Madrid, Spain

⁵Department of Physics & CREOL, University of Central Florida, FL32816, USA

⁶Instituto Madrileño de Estudios Avanzados en Nanociencia (IMDEA-Nanociencia), Cantoblanco, 28049 Madrid, Spain

⁷Condensed Matter Physics Center (IFIMAC), Universidad Autónoma de Madrid, 28049 Madrid, Spain

E-mail: david.busto@fysik.lth.se

Abstract. Autoionization, which results from the interference between direct photoionization and photoexcitation to a discrete state decaying to the continuum by configuration interaction, is a well known example of the important role of electron correlation in light-matter interaction. Information on this process can be obtained by studying the spectral, or equivalently, temporal complex amplitude of the ionized electron wavepacket. Using an energy-resolved interferometric technique, we measure the spectral amplitude and phase of autoionized wavepackets emitted via the $sp2^+$ and $sp3^+$ resonances in helium. These measurements allow us to reconstruct the corresponding temporal profiles by Fourier transform. In addition, applying various time-frequency representations, we observe the build up of the wavepackets in the continuum, monitor the instantaneous frequencies emitted at any time and disentangle the dynamics of the direct and resonant ionization channels.