Supplement to "CITIES: Clinical Trials With Intercurrent Events Simulator"

Ahmad Hakeem Abdul Wahab*,¹, Yongming Qu ², Hege Michiels ³, Junxiang Luo ⁴, Run Zhuang ⁵, Dominique McDaniel ⁶, Dong Xi ⁷, Elena Polverejan ¹, Steven Gilbert ⁸, Stephen Ruberg ⁹, and Arman Sabbaghi ⁵

Key words: Causal Inference; Estimands; Intercurrent Events; Repeated Measures.

1 Supplementary Material

To access the Shiny application, simply use the following link (https://hakeemwahabapp.shinyapps.io/cities/) where the default settings are set to the ones used in Section 2. The file app.R contains all the script used for the R shiny app. Although all CITIES outputs are in color, they appear as greyscale in this manuscript for the purposes of publication. The CITIES packages is available on CRAN (https://cran.r-project.org/web/packages/cities/index.html). A vignette (CITIES_demo.Rmd) is provided to replicate all output in this manuscript, and an additional example for custom covariates and covariances. Further, all code is publicly available on github (https://github.com/hakeemwahab/cities).

For the custom covariates and covariances example in the Rshiny app, users can use the attached csv file "custom_covariate_3arm.csv" for the custom covariates and "custom_covariance.csv" for the custom covariances (for all arms). The former is a csv containing 2 continuous covariates simulated from a standard normal and one binary covariate sampled from a bernoulli with probability 0.5 per subject with column names specified, while the latter contains a 5×5 covariance matrix without any column and row names. Note that users can have different custom covariances for each arm as well. For this example, you need to specify the following:

- 1. a vector of length 5 for the "Time Indices", e.g. 0, 12, 24, 48, 55.
- 2. a total of 300 subjects across all arms for the "Arms" input, e.g. 150, 50, 100 for arms 1, 2, and 3 respectively.
- 3. the "Baseline Covariate Coefficients" should all be of length 3 since we have 3 columns or covariates in teh supplied covariance csv file, e.g. 1.25, 1.25, 1 for all arms.

¹ Janssen Pharmaceuticals (Statistics and Decision Sciences, 08560)

² Eli Lilly and Company (Department of Statistics, Data and Analytics, 46285)

³ Ghent University (Department of Applied Mathematics, Computer Science and Statistics, 9000)

⁴ Moderna (Department of Biostatistics and Programming, 02139)

⁵ Purdue University (*Department of Statistics*, 47907)

⁶ Drexel University (Department of Epidemiology and Biostatistics, 19121)

⁷ Gilead Sciences (Department of Biostatistics, 94404)

⁸ Pfizer (Global Product Development, 02139)

⁹ Analytix Thinking LLC (Indianapolis, 46236)

^{*}Corresponding author: aabdulw1@its.jnj.com

4. the "Means" should all be of length 5, and all have the same initial baseline value. For a 3-arm example, you could supply "0, 0, 0, 0, 0" for arm 1, and "0, 0.2, 0.4, 0.6, 0.8" for arms 2 and 3 respectively.

1.1 Diabetes Clinical Trial Figures




Figure S1 The Mean Treatment Response Settings tab in CITIES for the Diabetes case study.




Figure S2 The Mean Treatment Response Settings Profile in CITIES for the Diabetes case study.



Figure S3 The LoE & EE tab in CITIES for the Diabetes case study.



Figure S4 The Admin & AE tab in CITIES for the Diabetes case study.




Figure S5 Treatment effect profiles for the three estimands in CITIES for the Diabetes case study. Note that the "Trt Policy Data" is greyed out, since the Treatment Policy options were not elected in Figure S1.

Cumulative Proportion Discontinued




Figure S6 Percentages of treatment discontinuations, and the corresponding ICEs in CITIES for the Diabetes case study.

1.2 Alzheimer's Clinical Trial Figures




Figure S7 The Mean Treatment Response Settings tab in CITIES for the Alzheimer's case study.




Figure S8 The Mean Treatment Response Settings Profile in CITIES for the Alzheimer's case study.




Figure S9 The LoE & EE tab in CITIES for the Alzheimer's case study.




Figure S10 The Admin & AE tab in CITIES for the Alzheimer's case study.




Figure S11 Treatment effect profiles for the three estimands in CITIES for the Alzheimer's case study. Note that the "Trt Policy Data" is greyed out, since the Treatment Policy options were not elected in Figure S7.

Cumulative Proportion Discontinued




Figure S12 Percentages of treatment discontinuations, and the corresponding ICEs in CITIES for the Alzheimer's case study.