

MATH574 - Dynamical Systems

Based on lectures from Winter 2025 by Prof. Antony Humphries.

Notes by Louis Meunier

Contents

1 Examples of Dynamical Systems	2
2 Existence-Uniqueness Theory	2
3 Limit Sets and the Evolution Operator	2

§1 EXAMPLES OF DYNAMICAL SYSTEMS

Roughly speaking, a dynamical system is a system that evolves in time, with common examples being a differential equation, in the continuous case, or a map, in the discrete case.

⊗ **Example 1.1** (The Logistic Map):

§2 EXISTENCE-UNIQUENESS THEORY

↪ **Definition 2.1** (Lipschitz): We say a function $f : \mathbb{R}^p \rightarrow \mathbb{R}^p$ is Lipschitz on $B \subseteq \mathbb{R}^p$ if there is a constant $L > 0$ such that $\|f(x) - f(y)\| \leq L \|x - y\|$ for every $x, y \in B$. We call L a “Lipschitz” constant. It is certainly not unique in general.

We say f *globally Lipschitz* if it is Lipschitz on $B = \mathbb{R}^p$, and f *locally Lipschitz* if f is Lipschitz on every bounded domain $B \subseteq \mathbb{R}^p$ (note: the L will in general depend on the domain).

↪ **Theorem 2.1**: Let $f : \mathbb{R}^p \rightarrow \mathbb{R}^p$ be a locally Lipschitz function. Then, there exists a unique solution to the initial value problem $\dot{u} = f(u)$, $u(0) = u_0$ on some interval $t \in (-T_1(u_0), T_2(u_0))$, where $-T_1(u_0) < 0 < T_2(u_0)$ and

- either $T_2(u_0) = +\infty$ or $\|u(t)\| \rightarrow \infty$ as $t \rightarrow T_2(u_0)$, and
- either $T_1(u_0) = -\infty$ or $\|u(t)\| \rightarrow -\infty$ as $t \rightarrow -T_1(u_0)$.

Heuristically, this first condition states that either our solution exists for all (forward) time after $-T_1(u_0)$, or it blows up in finite time, with a similar interpretation for the second, going backwards.

↪ **Proposition 2.1**: Let $\dot{u} = f(u)$ where f is locally Lipschitz. Let $B \subseteq \mathbb{R}^p$ be a bounded subset such that initial conditions $u_0, v_0 \in B$ define solutions $u(t), v(t)$ with $u(t), v(t) \in B$ for all $t \in [0, T]$. Let L be a Lipschitz constant for f on B . Then,

$$e^{-Lt} \|u_0 - v_0\| \leq \|u(t) - v(t)\| \leq e^{Lt} \|u_0 - v_0\| \quad \forall t \in [0, T].$$

This provides a bound on how quickly solutions grow, decay in B .

↪ **Corollary 2.1**: Let f be locally Lipschitz and $u_0 \neq v_0$. Then, $u(t) \neq v(t)$ for all time such that the solutions both exist.

§3 LIMIT SETS AND THE EVOLUTION OPERATOR

We state definitions in this section first for ODEs, but they generalize.

↪ **Definition 3.1** (Evolution Operator): Given $\dot{u} = f(u)$, the *evolution operator* is the map

$$S(t) : \mathbb{R}^p \rightarrow \mathbb{R}^p, \quad t \geq 0$$

such that $u(t) = S(t)u_0$.

Such a map also defines a *semi-group* $\{S(t) : t \geq 0\}$ under composition, namely it is closed under repeated composition and this operator is associative.

For $B \subseteq \mathbb{R}^p$ define

$$S(t)B := \bigcup_{u \in B} S(t)u = \{u(t) = S(t)u_0 : u_0 \in B\}.$$

↪ **Definition 3.2** (Forward/Positive Orbit): We define the *forward orbit* of a point u_0 as

$$\Gamma^+(u_0) := \bigcup_{t \geq 0} S(t)u_0,$$

i.e. the set of all points u_0 may “visit” as time increases.

↪ **Definition 3.3** (Backwards/Negative Orbit): Similarly, define a *backwards orbit* (if one exists)

$$\Gamma^-(u_0) := \{u(t) : t \leq 0\},$$

s.t. $\forall t \leq s \leq 0, S(-t)u(t) = u_0$ and $S(s-t)u(t) = u(s)$.

Note that a negative orbit won't be unique in general, eg in maps, periodic points may multiple preimages.

↪ **Definition 3.4** (Complete Orbit): If a negative orbit for u_0 exists, define the *complete orbit* through u_0 as

$$\Gamma(u_0) := \Gamma^+(u_0) \cup \Gamma^-(u_0).$$

Notice that if $v \in \Gamma(u_0)$, then $\Gamma(v) = \Gamma(u_0)$; namely a complete orbit through v exists.

↪ **Definition 3.5** (Invariance): The set B is said to be *positively invariant* if $S(t)B \subseteq B$ for all $t \geq 0$. Similarly, B is said to be *negatively invariant* if $B \subseteq S(t)B$ for all $t \geq 0$.

↪ **Definition 3.6** (ω -limit sets): A point $x \in \mathbb{R}^p$ is called an ω -limit point of u_0 if there exists a sequence $\{t_n\}$ with $t_n \rightarrow \infty$ such that $S(t_n)u_0 \rightarrow x$ as $n \rightarrow \infty$. The set of all such points for an initial condition u_0 is denoted $\omega(u_0)$, and called the ω -limit set of u_0 .

Given a bounded set B , the ω -limit set of B is defined as

$$\omega(B) := \{x \in \mathbb{R}^p : \exists t_n \rightarrow \infty, y_n \in B \text{ s.t. } S(t_n)y_n \rightarrow x\}.$$

Remark 3.1: In general, $\omega(B)$ is *not* the union of ω -limit sets of points in B .

↪ **Theorem 3.1:** For any $u_0 \in \mathbb{R}^p$,

$$\omega(u_0) = \bigcap_{s \geq 0} \overline{\bigcup_{t \geq s} \{S(t)u_0\}},$$

and similarly for any bounded $B \subseteq \mathbb{R}^p$,

$$\omega(B) = \bigcap_{s \geq 0} \overline{\bigcup_{t \geq s} S(t)B}.$$

↪ **Definition 3.7** (α -limit set): A point $x \in \mathbb{R}^p$ is called an α -limit point for $u_0 \in \mathbb{R}^p$ if there exists a negative orbit through u_0 and a sequence $\{t_n\}$ with $t_n \rightarrow -\infty$ such that $u(t_n) \rightarrow x$. The set of all such points for u_0 is denoted $\alpha(u_0)$.

↪ **Theorem 3.2:** If $\Gamma^+(u_0)$ bounded, then $\omega(u_0)$ is a non-empty, compact, invariant, connected set.

↪ **Definition 3.8** (Attraction): We say a set A attracts B if for every $\varepsilon > 0$, there is a $t^* = t^*(\varepsilon, A, B)$ such that $S(t)B \subseteq N(A, \varepsilon)$ for every $t \geq t^*$, where $N(A, \varepsilon)$ denotes the ε -neighborhood of A .

A compact, invariant set A is called an *attractor* if it attracts an open neighborhood of itself, i.e. $\exists \varepsilon > 0$ such that A attracts $N(A, \varepsilon)$.

A *global attractor* is an attractor that attracts every bounded subset of \mathbb{R}^p .

↪ **Theorem 3.3** (Continuous Gronwall Lemma): Let $z(t)$ be such that $\dot{z} \leq az + b$ for some $a \neq 0, b \in \mathbb{R}$ and $z(t) \in \mathbb{R}$. Then, $\forall t \geq 0$,

$$z(t) \leq e^{at}z(0) + \frac{b}{a}(e^{at} - 1).$$

↪ **Theorem 3.4** (ω -limit sets as attractors): Assume $B \subseteq \mathbb{R}^p$ is a bounded, open set such that $S(t)B \subseteq \bar{B} \forall t > 0$. Then, $\omega(B) \subseteq B$, and $\omega(B)$ is an attractor, which attracts B . Furthermore,

$$\omega(B) = \bigcap_{t \geq 0} S(t)B.$$

↪ **Definition 3.9** (Dissipative): A dynamical system is called *dissipative* if there exists a bounded set B such $\forall A$ bounded, there exists a $t^* = t^*(A) > 0$ such that $S(t)A \subseteq B \forall t \geq t^*$. We then call such a B an *absorbing set*.

Remark 3.2: B absorbing $\Rightarrow \omega(A) \subseteq \omega(B)$. Moreover, $\omega(B)$ attracts A for every bounded set A . I.e., $\omega(B)$ is a global attractor.