

Course Outline:

Based on Lectures from Winter, 2024 by Prof. Anush Tserunyan.

Contents

1	Introduction	2
1.1	Vector Spaces	2
1.2	Creating Spaces from Other Spaces	4
1.3	Linear Combinations and Space	6

1 Introduction

Remark 1.1. *This course is about vector spaces and linear transformations between them; a vector space involves multiplication by scalars, where the scalars come from some field. We recall first examples of fields, then vector spaces, as a motivation, before presenting a formal definition.*

1.1 Vector Spaces

Remark 1.2. *Much of this is recall from [Algebra 1](#).*

⊗ Example 1.1: Examples of Fields

1. \mathbb{Q} ; the field of rational numbers.
2. \mathbb{R} ; the field of real numbers; $\mathbb{Q} \subseteq \mathbb{R}$.
3. \mathbb{C} ; the field of complex numbers; $\mathbb{Q} \subseteq \mathbb{R} \subseteq \mathbb{C}$.
4. $\mathbb{F}_p \equiv \mathbb{Z}/p\mathbb{Z} \equiv \{0, 1, \dots, p-1\}$; the (unique) field of p elements, where p prime.^a
 - (a) $p = 2$; $\mathbb{F}_2 \equiv \{0, 1\}$.
 - (b) $p = 3$; $\mathbb{F}_3 \equiv \{0, 1, 2\}$.
 - (c) \dots

^awhere $a +_p b := \text{remainder of } \frac{a+b}{p}$, $a \cdot_p b := \text{remainder of } \frac{a \cdot b}{p}$.

Remark 1.3. *Throughout the course, we will denote an abstract field as \mathbb{F} .*

⊗ Example 1.2: Examples of Vector Spaces

1. $\mathbb{R}^3 := \{(x, y, z) : x, y, z \in \mathbb{R}\}$. We can add elements in \mathbb{R}^3 , and multiply them by real scalars.
2. $\mathbb{F}^n := \underbrace{\mathbb{F} \times \mathbb{F} \times \dots \times \mathbb{F}}_{n \text{ times}} := \{(a_1, a_2, \dots, a_n) : a_i \in \mathbb{F}\}$, where $n \in \mathbb{N}^+$; this is a generalization of the previous example, where we took $n = 3$, $\mathbb{F} = \mathbb{R}$. Operations follow identically; addition:

$$(a_1, a_2, \dots, a_n) + (b_1, b_2, \dots, b_n) := (a_1 + b_1, a_2 + b_2, \dots, a_n + b_n)$$

and, taking a scalar $\lambda \in \mathbb{F}$, multiplication:

$$\lambda \cdot (a_1, a_2, \dots, a_n) := (\lambda \cdot a_1, \lambda \cdot a_2, \dots, \lambda \cdot a_n).$$

We refer to these elements (a_1, \dots, a_n) as *vectors* in \mathbb{F}^n ; the vector for which

$a_i = 0 \forall i$ is the 0 *vector*, and is the additive identity, making \mathbb{F}^n an abelian group under addition, that admits multiplication by scalars from \mathbb{F} .

3. $C(\mathbb{R}) := \{f : \mathbb{R} \rightarrow \mathbb{R} : f \text{ continuous}\}$. Here, we have the constant zero function as our additive identity ($x \mapsto 0 \forall x$), and addition/scalar multiplication of two continuous real functions are continuous.

4. $\mathbb{F}[t] := \{a_0 + a_1t + a_2t^2 + \cdots + a_nt^n : a_i \in \mathbb{F} \forall i, n \in \mathbb{N}\}$, ie, the set of all polynomials in t with coefficients from \mathbb{F} . Here, we can add two polynomials;

$$(a_0 + a_1t + \cdots + a_nt^n) + (b_0 + b_1t + \cdots + b_mt^m) := \sum_{i=0}^{\max\{n,m\}} (a_i + b_i)t^i,$$

(where we “take” undefined a_i/b_i ’s as 0; that is, if $m > n$, then $a_{m-n}, a_{m-n+1}, \dots, a_m$ are taken to be 0). Scalar multiplication is defined

$$\lambda \cdot (a_0 + a_1t + a_2t^2 + \cdots + a_nt^n) := \lambda a_0 + \lambda a_1t + \lambda a_2t^2 + \cdots + \lambda a_nt^n.$$

Here, the zero polynomial is simply 0 (that is, $a_i = 0 \forall i$).

↪ **Definition 1.1: Vector Space**

A *vector space* V over a field \mathbb{F} is an *abelian group* with an operation denoted $+$ (or $+_V$) and identity element² denoted 0_V , equipped with *scalar multiplication* for each scalar $\lambda \in \mathbb{F}$ satisfying the following axioms:

1. $1 \cdot v = v$ for $1 \in \mathbb{F}, \forall v \in V$.
2. $\alpha \cdot (\beta \cdot v) = (\alpha \cdot \beta)v, \forall \alpha, \beta \in \mathbb{F}, v \in V$.
3. $(\alpha + \beta) \cdot v = \alpha \cdot v + \beta \cdot v, \forall \alpha, \beta \in \mathbb{F}, v \in V$.
4. $\alpha \cdot (u + v) = \alpha \cdot u + \alpha \cdot v, \forall \alpha \in \mathbb{F}, u, v \in V$.

We refer to elements $v \in V$ as *vectors*.

↪ **Proposition 1.1**

For a vector space V over a field \mathbb{F} , the following holds:

1. $0 \cdot v = 0_V, \forall v \in V$ (where $0 := 0_{\mathbb{F}}$)
2. $-1 \cdot v = -v, \forall v \in V$ (where $1 := 1_{\mathbb{F}}$)³

¹Where we take $0 \in \mathbb{N}$, for sake of consistency. Moreover, by convention, we define \mathbb{F}^0 (that is, when $n = 0$) to be $\{0\}$; the trivial vector space.

²The “zero vector”.

$$3. \alpha \cdot 0_V = 0_V, \forall \alpha \in \mathbb{F}$$

³NB: “additive inverse”

Proof. 1. $0 \cdot v = (0 + 0) \cdot v = 0 \cdot v + 0 \cdot v \implies 0 \cdot v = 0_V$ (by “cancelling” one of the $0 \cdot v$ terms on each side).

$$2. v + (-1 \cdot v) = (1 \cdot v + (-1) \cdot v) = (1 - 1) \cdot v = 0 \cdot v = 0_V \implies (-1 \cdot v) = -v.$$

$$3. \alpha \cdot 0_V = \alpha \cdot (0_V + 0_V) = \alpha \cdot 0_V + \alpha \cdot 0_V \implies \alpha \cdot 0_V = 0_V \text{ (by, again, cancelling a term on each side).}$$

■

↪ Mon Jan 8 13:50:37 EST 2024

1.2 Creating Spaces from Other Spaces

↪ [Definition 1.2: Product/Direct Sum of Vector Spaces](#)

For vector spaces U, V over the same field \mathbb{F} , we define their *product* (or *direct sum*) as the set

$$U \times V = \{(u, v) : u \in U, v \in V\},$$

with the operations:

$$(u_1, v_1) + (u_2, v_2) := (u_1 + u_2, v_1 + v_2)$$

$$\lambda \cdot (u, v) := (\lambda \cdot u, \lambda \cdot v)$$

⊗ [Example 1.3: \$\mathbb{F}\$](#)

$\mathbb{F}^2 = \mathbb{F} \times \mathbb{F}$, where \mathbb{F} is considered as the vector space over \mathbb{F} (itself).

↪ Definition 1.3: Subspace

For a vector space V over a field \mathbb{F} , a *subspace* of V is a subset $W \subseteq V$ s.t.

1. $0_V \in W$ ⁴
2. $u + v \in W \forall u, v \in W$ (closed under addition)
3. $\alpha \cdot u \in W \forall u \in W, \alpha \in \mathbb{F}$ ⁵

Then, W is a vector space in its own right.

⊗ Example 1.4: Examples of Subspaces

1. Let $V := \mathbb{F}^n$.

- $W := \{(x_1, x_2, \dots, x_n) \in \mathbb{F}^n : x_1 = 0\} = \{(0, x_2, x_3, \dots, x_n) : x_i \in \mathbb{F}\}$.
- $W := \{(x_1, x_2, \dots, x_n) \in \mathbb{F}^n : x_1 + 2 \cdot x_2 = 0\}$

Proof. Let $x = (x_1, \dots, x_n), y = (y_1, \dots, y_n) \in W$. Then, $x + y = (x_1 + y_1, \dots, x_n + y_n)$, and $x_1 + y_1 + 2 \cdot (x_2 + y_2) = x_1 + 2 \cdot x_2 + y_1 + 2 \cdot y_2 = 0 + 0 = 0 \implies x + y \in W$. Similar logic follows for axioms 2., 3. ■

- (More generally)

$$W := \{(x_1, \dots, x_n) \in \mathbb{F}^n : \begin{array}{l} a_{11}x_1 + \dots + a_{1n}x_n = 0 \\ a_{21}x_1 + \dots + a_{2n}x_n = 0 \\ \vdots \\ a_{k1}x_1 + \dots + a_{kn}x_n = 0 \end{array} \},$$

that is, a linear combination of homogenous “conditions” on each term.

- $W^* := \{(x_1, \dots, x_n) : x_1 + x_2 = 1\}$ is *not* a subspace; it is not closed under addition, nor under scalar multiplication.

2. Let $\mathbb{F}[t]_n := \{a_0 + a_1t + \dots + a_nt^n : a_i \in \mathbb{F}\}$. Then, $\mathbb{F}[t]_n$ is a subspace of $\mathbb{F}[t]$, the more general polynomial space. However, the set of all polynomials of degree *exactly* n (all axioms fail, in fact) is not a subspace of $\mathbb{F}[t]_n$.

- $W := \{p(t) \in \mathbb{F}[t]_n : p(1) = 0\}$.
- $W := \{p(t) \in \mathbb{F}[t]_n : p''(t) + p'(t) + 2p(t) = 0\}$.

3. Let $V := C(\mathbb{R})$ be the space of continuous function $\mathbb{R} \rightarrow \mathbb{R}$.

⁴This is equivalent to requiring that $W \neq \emptyset$; stated this way, axiom 3. would necessitate that $0 \cdot w = 0_V \in W$.

⁵Note that these axioms are equivalent to saying that W is a subgroup of V with respect to vector addition; 2. ensures closed under addition, and 3. ensures the existence of additive inverses (as per $-1 \cdot v = -v$).

- $W := \{f \in C(\mathbb{R}) : f(\pi) + 7f(\sqrt{2}) = 0\}$.
- $W := C^1(\mathbb{R}) := \text{everywhere differentiable functions}$.
- $W := \{f \in C(\mathbb{R}) : \int_0^1 f \, dx = 0\}$.

↪ **Proposition 1.2**

Let W_1, W_2 be subspaces of a vector space V over \mathbb{F} . Then, define the following:

1. $W_1 + W_2 := \{w_1 + w_2 : w_1 \in W_1, w_2 \in W_2\}$
2. $W_1 \cap W_2 := \{w \in V : w \in W_1 \wedge w \in W_2\}$

These are both subspaces of V .

- Proof.
1. (a) $0_V \in W_1$ and $0_V \in W_2 \implies 0_V = 0_V + 0_V \in W_1 + W_2$.
 (b) $(u_1 + u_2) + (v_1 + v_2) = (u_1 + v_1) + (u_2 + v_2) \in W_1 + W_2$.
 (c) $\alpha \cdot (u + v) = \alpha \cdot u + \alpha \cdot v \in W_1 + W_2$
 2. (a) $0_V \in W_1$ and $0_V \in W_2 \implies 0_V = 0_V + 0_V \in W_1 \cap W_2$.
 (b) $u, v \in W_1 \cap W_2 \implies u + v \in W_1 \wedge u + v \in W_2 \implies u + v \in W_1 \cap W_2$.
 (c) $\alpha \cdot u \in W_1 \wedge \alpha \cdot u \in W_2 \implies \alpha \cdot u \in W_1 \cap W_2$.

■

1.3 Linear Combinations and Space

↪ **Definition 1.4: Linear Combination**

Let V be a vector space over a field \mathbb{F} . For finitely many vectors v_1, v_2, \dots, v_n , their *linear combination* is a sum of the form

$$\sum_{i=1}^n a_i v_i = a_1 \cdot v_1 + \dots + a_n \cdot v_n,$$

where $a_i \in \mathbb{F} \forall i$.

A linear combination is called *trivial* if $a_i = 0 \forall i$, that is, all coefficients are 0.

↪ Mon Jan 8 18:19:28 EST 2024