

# MATH457 - Algebra 4

Representation Theory; Galois Theory

Based on lectures from Winter 2025 by Prof. Henri Darmon.

Notes by Louis Meunier

## Contents

1 Representation Theory .....	2
1.1 Introduction .....	2

## §1 REPRESENTATION THEORY

### §1.1 Introduction

↪ **Definition 1.1** (Linear Representation): A *linear representation* of a group  $G$  is a vector space  $V$  over a field  $\mathbb{F}$  equipped with a map  $G \times V \rightarrow V$  that makes  $V$  a  $G$ -set in such a way that for each  $g \in G$ , the map  $v \mapsto gv$  is a linear homomorphism of  $V$ .

This induces a homomorphism

$$\rho : G \rightarrow \text{Aut}_{\mathbb{F}}(V),$$

or, in particular, when  $n = \dim_{\mathbb{F}} V < \infty$ , a homomorphism

$$\rho : G \rightarrow \text{GL}_n(\mathbb{F}).$$

Alternatively, a linear representation  $V$  can be viewed as a module over the group ring  $\mathbb{F}[G] = \left\{ \sum_{g \in G} \lambda_g g : \lambda_g \in \mathbb{F} \right\}$  (where we require all but finitely many scalars  $\lambda_g$  to be zero).

↪ **Definition 1.2** (Irreducible Representation): A linear representation  $V$  of a group  $G$  is called *irreducible* if there exists no proper, nontrivial *subspace*  $W \subsetneq V$  such that  $W$  is  $G$ -stable.

#### ⊗ Example 1.1:

1. Consider  $G = \mathbb{Z}/2 = \{1, \tau\}$ . If  $V$  a linear representation of  $G$  and  $\rho : G \rightarrow \text{Aut}(V)$ . Then,  $V$  uniquely determined by  $\rho(\tau)$ . Let  $p(x)$  be the minimal polynomial of  $\rho(\tau)$ . Then,  $p(x) \mid x^2 - 1$ . Suppose  $\mathbb{F}$  is a field in which  $2 \neq 0$ . Then,  $p(x) \mid (x - 1)(x + 1)$  and so  $p(x)$  has either  $1, -1$ , or both as eigenvalues and thus we may write

$$V = V_+ \oplus V_-,$$

where  $V_{\pm} := \{v \mid \tau v = \pm v\}$ . Hence,  $V$  is irreducible only if one of  $V_+, V_-$  all of  $V$  and the other is trivial, or in other words  $\tau$  acts only as multiplication by  $1$  or  $-1$ .

2. Let  $G = \{g_1, \dots, g_N\}$  be a finite abelian group, and suppose  $\mathbb{F}$  an algebraically closed field of characteristic  $0$  (such as  $\mathbb{C}$ ). Let  $\rho : G \rightarrow \text{Aut}(V)$  and denote  $T_j := \rho(g_j)$  for  $j = 1, \dots, N$ . Then,  $\{T_1, \dots, T_N\}$  is a set of mutually commuting linear transformations. Then, there exists a simultaneous eigenvector, say  $v$ , for  $\{T_1, \dots, T_N\}$ , and so  $\text{span}(v)$  a  $G$ -stable subspace of  $V$ . Thus, if  $V$  irreducible, it must be that  $\dim_{\mathbb{F}} V = 1$ .

↪ **Theorem 1.1:** If  $G$  a finite abelian group and  $V$  an irreducible finite dimensional representation over an algebraically closed field of characteristic  $0$ , then  $\dim V = 1$ .

PROOF. Let  $\rho : G \rightarrow \text{Aut}(V)$ , label  $G = \{g_1, \dots, g_N\}$  and put  $T_j := \rho(g_j)$  for  $j = 1, \dots, N$ . Then,  $\{T_1, \dots, T_N\}$  a family of mutually commuting linear transformations on  $V$ . Then, there is a simultaneous eigenvector  $v$  for  $\{T_1, \dots, T_N\}$  and thus  $\text{span}(v)$  is  $T_1, \dots, T_N$ -stable and so  $V = \text{span}(v)$ . ■

↪ **Lemma 1.1:** Let  $V$  be a finite dimensional vector space over  $\mathbb{C}$  and let  $T_1, \dots, T_N : V \rightarrow V$  be a family of mutually commuting linear automorphisms on  $V$ . Then, there is a simultaneous eigenvector for  $T_1, \dots, T_N$ .

↪ **Proposition 1.1:** Let  $\mathbb{F}$  a field where  $2 \neq 0$  and  $V$  an irreducible representation of  $S_3$ . Then, there are three distinct (i.e., up to homomorphism) possibilities for  $V$ .

PROOF. Let  $\rho : G \rightarrow \text{Aut}(V)$  and let  $T = \rho((23))$ . Then, notice that  $p_T(x) \mid (x^2 - 1)$  so  $T$  has eigenvalues in  $\{-1, 1\}$ .

If the only eigenvalue of  $T$  is  $-1$ , we claim that  $V$  one-dimensional.

If  $T$  has 1 as an eigenvalue. ■

↪ **Proposition 1.2:**  $D_8$  has a unique faithful irreducible representation, of dimension 2 over a field  $F$  in which  $0 \neq 2$ .

PROOF. Write  $G = D_8 = \{1, r, r^2, r^3, v, h, d_1, d_2\}$  as standard. Let  $\rho$  be our irreducible, faithful representation and let  $T = \rho(r^2)$ . Then,  $p_T(x) \mid x^2 - 1 = (x - 1)(x + 1)$  and so  $V = V_+ \oplus V_-$ , the respective eigenspaces for  $\lambda = +1, -1$  respectively for  $T$ . Then, notice that since  $r^2$  in the center of  $G$ , both  $V_+$  and  $V_-$  are preserved by the action of  $G$ , hence one must be trivial and the other the entirety of  $V$ .  $V$  can't equal  $V_+$ , else  $T = I$  on all of  $V$  hence  $\rho$  not faithful so  $V = V_-$ .

Next, it must be that  $\rho(h)$  has both eigenvalues 1 and  $-1$ . Let  $v_1 \in V$  be such that  $hv_1 = v_1$  and  $v_2 = rv_1$ . We claim that  $W := \text{span}\{v_1, v_2\}$ , namely  $V = W$  2-dimensional.

We simply check each element.  $rv_1 = v_2$  and  $rv_2 = r^2v_1 = -v_1$  which are both in  $W$  hence  $r$  and thus  $\langle r \rangle$  fixes  $W$ . Next,  $hv_1 = v_1$  and  $vv_2 = vrv_1 = rhv_1 = rv_1 = v_2$  (since  $rhr^{-1} = v$ ) and so  $hv_2 = -v_2$  and  $vv_1 = -v_1$  and so  $W$   $G$ -stable. Finally,  $d_1$  and  $d_2$  are just products of these elements and so  $W$   $G$ -stable. ■